Temporal dynamics of collateral RNA cleavage by LbuCas13a in human cells

Jorik F. Bot[1,2]*, Zhihan Zhao[1,2,3]*, Mengyuan Li[1,2], Darnell Kammeron[1,2], Peng Shang[1,2,‡] and Niels Geijsen[1,2,‡]

[1] Dept. of Anatomy & Embryology, Leiden University Medical Center, Einthovenweg 20, 2300 RC Leiden, The Netherlands.

[2] The Novo Nordisk Foundation Center for Stem Cell Medicine (reNEW), Leiden node, The Netherlands

[3] Biomedical Pioneering Innovation Center, Beijing Advanced Innovation Center for Genomics, Peking-Tsinghua Center for Life Sciences, Peking University Genome Editing Research Center, State Key Laboratory of Protein and Plant Gene Research, School of Life Sciences, Peking University, Beijing, China.

* These authors contributed equally to this work.

‡ Leiden University Medical Center, Dept. of Anatomy & Embryology, Building 2, Einthovenweg 20, 2300 RC Leiden, The Netherlands. +31-(0)71-526 9302, p.shang@lumc.nl and n.geijsen@lumc.nl

Abstract

CRISPR-Cas13 exclusively targets RNA. In prokaryotic cells, Cas13 cleaves both target and
non-target RNA indiscriminately upon activation by a specific target RNA, but in eukaryotic cells
collateral cleavage activity has been limited. To investigate collateral cleavage by Cas13 in
eukaryotic cells, we first compared various Cas13 orthologs and found that specifically
LbuCas13a exhibits strong collateral RNA cleavage activity in human cells when delivered as
ribonucleoprotein, independent of cell line and targeting both exogenous and endogenous
transcripts. Collateral RNA cleavage started within 50 minutes of ribonucleoprotein delivery
resulting in major alterations to the total RNA profile. In response to the collateral RNA
cleavage, cells upregulated genes associated with stress and innate immune response,
ultimately leading to apoptotic cell death. This enabled us to use LbuCas13a as a flexible and
repeatable target-RNA-specific cell elimination tool. Finally, we used Nanopore sequencing to
explore the identity of collaterally cleaved RNAs, the nucleotide position at which they are
cleaved, and the temporal dynamics of collateral RNA cleavage. This revealed that LbuCas13a
activation leads to rapid and global cleavage of cytoplasmic RNAs at specific nucleotide
positions. In conclusion, we here report that LbuCas13a has high collateral activity in human
cells and describe the temporal dynamics of the collateral RNA cleavage, the cellular responses
ultimately leading to apoptosis, how this can be exploited as a cell elimination tool, and the
collateral cleavage preferences of LbuCas13a.

Introduction

CRISPR-Cas is a prokaryotic adaptive immune system that protects bacteria and archaea from invading viruses and plasmids (Marraffini, 2015; Mohanraju et al., 2016; Oost et al., 2014). Since its initial discovery, an impressive diversity of CRISPR-Cas systems has been revealed (Makarova et al., 2020). Especially the Class 2 DNA targeting effector proteins, Cas9 and Cas12, have been extensively studied and have found widespread adoption for biotechnological and clinical use (Liu et al., 2022; Sharma et al., 2021). A recently discovered Class 2 system, CRISPR-Cas13, targets and cleaves RNA. The signature element of CRISPR-Cas13 systems are the two Higher Eukaryotes and Prokaryotes Nucleotide-binding (HEPN) RNase domains of the effector protein (Koonin et al., 2017; Makarova et al., 2015). When a Cas13 protein is activated by a guide-matching target RNA, these domains cooperate to form a single active RNA nuclease site (Abudayyeh et al., 2016; Cox et al., 2017; East-Seletsky et al., 2016; Konermann et al., 2018; Smargon et al., 2017; Xu et al., 2021; Yan et al., 2018). This catalytic site is located on the outside of the protein, facing away from the guide RNA-target RNA complex (Knott et al., 2017; Liu et al., 2017b, 2017a; Slaymaker et al., 2019; Zhang et al., 2018). As a consequence, Cas13 cleaves both target and non-target RNA indiscriminately *in vitro* (Abudayyeh et al., 2016; East-Seletsky et al., 2017, 2016; Konermann et al., 2018; Smargon et al., 2017; Xu et al., 2021; Yan et al., 2018) and in bacteria (Abudayyeh et al., 2016; Meeske et al., 2019; Smargon et al., 2017; Yan et al., 2018) when activated by a specific target RNA. *In vitro,* this collateral RNA cleavage has been employed to develop portable, rapid, and highly sensitive nucleic acid detection methods (East-Seletsky et al., 2017; Gootenberg et al., 2018, 2017; Myhrvold et al., 2018), including for Coronavirus disease 2019 (COVID-19) diagnostics (Arizti-sanz et al., 2020; Fozouni et al., 2021; Patchsung et al., 2020). In bacteria, collateral RNA cleavage probably serves to protect bacterial communities by inducing dormancy after phage infection (Meeske et al., 2019; Mohanraju et al., 2022).

Initially, no evidence of collateral cleavage was found when Cas13 was first tested in eukaryotic cells (Abudayyeh et al., 2017; Cox et al., 2017; Konermann et al., 2018). Instead, Cas13 induced efficient and specific target RNA knockdown. Because Cas13-based knockdown was shown to have higher specificity and efficiency than RNAi (Abudayyeh et al., 2017; Cox et al., 2017; He et al., 2020; Huynh et al., 2020; Konermann et al., 2018; Li et al., 2021; Zhang et al.,

2021) and CRISPRi (Huynh et al., 2020; Konermann et al., 2018), it has seen enthusiastic

adoption as a specific RNA knockdown tool. For example, Cas13 has been adapted for use in

mice (He et al., 2020; Kushawah et al., 2020; Powell et al., 2022; Zhou et al., 2020), fish

(Kushawah et al., 2020; Wang et al., 2021), yeast (Jing et al., 2018), fly (Huynh et al., 2020),

plants (Aman et al., 2018; Mahas et al., 2019), and as an anti-viral tool (Blanchard et al., 2021;

Freije et al., 2019; Tng et al., 2020). However, recently several groups have reported that Cas13

is capable of collateral cleavage in eukaryotes in certain conditions, in some cases causing

cytotoxicity (Ai et al., 2022; Buchman et al., 2020; Y. Li et al., 2023; Özcan et al., 2021; Shi et

al., 2023; Wang et al., 2019; Xu et al., 2021; Zhang et al., 2019). To further advance CRISPR-

Cas13 as a specific RNA knockdown method, some groups have made progress to minimize

the Cas13 collateral cleavage effects (Ai et al., 2022; Tong et al., 2022). We took the opposite

approach and set out to exploit Cas13's collateral cleavage activity as a target-RNA-specific cell

elimination tool. A comparison of the collateral cleavage activity of different Cas13 orthologs

identified LbuCas13a as a particularly active Cas13 variant. In this set out to explore the on-

target and collateral RNA cleavage activity of LbuCas13a in cells. We observed that specifically

ribonucleoprotein complex (RNP) delivery of LbuCas13a induces robust collateral RNA

cleavage in human cells. The collateral activity is cell line and RNP delivery method

independent, and is induced by targeting both exogenous and endogenous transcripts. The

resulting collateral RNA cleavage leads to transcriptomic changes associated with the activation

of an innate immune response. We demonstrate that collateral RNA cleavage triggers target

RNA-specific apoptosis, which is strongly correlated with the expression level of the target RNA.

Consequently, LbuCas13a can be used as a target RNA-specific cell selection tool which

eliminates specific cell types from a heterogenous cell population. We demonstrate that this can

be used to enrich for cells that have undergone gene editing. Finally, using Nanopore

sequencing, we observed that upon activation by the target RNA, LbuCas13a indiscriminately

cleaves cytoplasmic RNAs. This cleavage occurs at very specific positions, often within the

loops of stem-loop structures. In conclusion, here we describe the highly active LbuCas13a, the

temporal dynamics of target and collateral RNA cleavage, the cellular response to collateral

RNA cleavage, the application of LbuCas13a as a negative cell selection tool, and the identity

and cleavage positions of the RNAs subject to collateral RNA cleavage by LbuCas13a.

Results

LbuCas13a exhibits strong collateral RNA cleavage activity in human cells

The enzymatic activity of different Cas13 orthologs *in vitro* is known to range over seven orders of magnitude (East-Seletsky et al., 2017). We hypothesized that the Cas13 orthologs with the strongest collateral cleavage activity *in vitro* are the most likely to show any collateral cleavage in human cells. Therefore, we compared the collateral RNA cleavage activity of six different CRISPR-Cas13 effectors (LbuCas13a, LwaCas13a, BzCas13b, PspCas13b, RspCas13d and RfxCas13d) *in vitro*, as well as in cells. We produced and purified all six recombinant proteins (Supplementary figure 1A). Next, the *in vitro* collateral RNA cleavage activity of these proteins was assessed using the RNaseAlert® assay (IDT), which quantitatively detects non-specific RNA cleavage. LbuCas13a exhibited the strongest *in vitro* collateral cleavage activity (Supplementary figure 1B). To test these orthologs in human cells, we transfected them as recombinant protein together with guide RNAs into HAP1 cells constitutively overexpressing a fluorescence-inactivated EGFP^{Y66S} (Zhao et al., 2022), referred to as HAP1-dEGFP from hereon. As transfection method we used 'induced Transduction by Osmocytosis and Propanebetaine' (iTOP), a previously reported method for the intracellular delivery of recombinant proteins (D'Astolfo et al., 2015). Next, the total RNA integrity was assessed using a Bioanalyzer. The total RNA profile of dEGFP expressing cells transfected with LbuCas13a and a dEGFP targeting guide RNA clearly revealed a degradation pattern mainly consisting of several novel fragments preceding the 18S ribosomal RNA (rRNA) peak (Figure 1A), while the total RNA of wildtype HAP1 cells transfected with LbuCas13a and a dEGFP targeting guide RNA was intact at all timepoints. This collateral RNA cleavage pattern was time-dependent and peaked at 100 to 200 minutes after transfection, with some cleavage fragments remaining at 24 hours after transfection. None of the other Cas13 orthologs were able to generate such a distinct change to the total RNA pattern for any of the target RNAs tested (Supplementary figures 2 and 3). Next we performed an LbuCas13a protein and guide RNA concentration gradient, ranging from 1 µM to 15 µM and using an equimolar ratio of protein and guide RNA (Supplementary figure 4). Collateral cleavage was evident in the total RNA profile from 1 µM and increased logarithmically with the concentration. Because 15 µM yielded the strongest

collateral effect and further increases in concentration probably would only give a minimal

improvement, we used 15 µM of protein and guide RNA in all further experiments.

To specifically assess the cleavage of the target transcript and a collateral transcript, we used a

modified RT-qPCR strategy designed to quantify transcript integrity (Auer and Lyianarachchi,

2003; Nolan et al., 2006; Swift et al., 2000), here named 5'-3' RT-qPCR. In the 5'-3' RT-qPCR

assay, cDNA is created by reverse transcription using oligo-dT primers, driving cDNA synthesis

from the 3' end of the mRNA transcript (Figure 1B). Thus, transcript degradation results in a 3'

bias of the cDNA. Consequently, a 5' located primer pair is expected to return higher Cq values

than a primer pair located at the 3' end of the transcript. To validate this assay, we first

transfected cells with RNase A protein. This indeed resulted in the expected Cq value increase

of the 5' primer pairs relative to the 3' primer pairs (Supplementary figure 5A). Using the 5'-3'

RT-qPCR assay, we observed that LbuCas13a was able to degrade both target RNA (dEGFP)

and collateral RNA (GAPDH) (Figure 1C). While target RNA cleavage seemingly started almost

immediately, the collateral GAPDH cleavage started later and peaked at around 100 minutes

after transfection. Both target and collateral RNA integrity returned to normal from about 500

minutes after transfection. dEGFP targeting with LwaCas13a, PspCas13b and BzCas13b also

resulted in a significant relative increase of the 5' Cq values of the dEGFP transcript, indicating

that these orthologs degraded the dEGFP target RNA in HAP1-dEGFP cells (Supplementary

figure 5B). While RfxCas13d did increase the relative 5' Cq values of the target transcript when

targeting dEGFP (Supplementary figure 5C), this increase was not significant (P = 0.076).

RspCas13d also did not show significant target RNA cleavage. Although LwaCas13a,

BzCas13b and PspCas13b showed a minimal increase in non-target RNA 5' Cq values

compared to the non-targeting control (Supplementary figure 5B), suggesting low levels of

collateral cleavage, these increases were not significant. RspCas13d and RfxCas13d did not

show an increase of collateral RNA 5' Cq values (Supplementary figure 5B and C).

Taken together, we observed that under these specific conditions (I.e. iTOP RNP delivery,

HAP1 cells, for these specific guide RNAs), LbuCas13a exhibited the highest collateral RNA

cleavage activity both *in vitro* and in cells. Therefore, we used LbuCas13a for all further

experiments. While collateral cleavage inside cells has also been reported for some of the other

orthologs compared here (Ai et al., 2022; Y. Li et al., 2023; Shi et al., 2023), we could not find

strong evidence for collateral cleavage with these orthologs. This further highlights the

situational nature of collateral cleavage by Cas13 in human cells (Bot et al., 2022), which we

further discuss in supplementary note 1.

Different RNP delivery methods support LbuCas13a collateral cleavage

We next examined whether collateral cleavage could also be induced using different delivery

methods, and whether it depended on targeting exogenous RNA. First, we electroporated HAP1

cells with LbuCas13a and a guide RNA targeting one of the highly abundant endogenous

transcripts RPS19, GAPDH or 18S rRNA. Delivery by electroporation resulted in strong total

RNA degradation for all three guide RNAs, demonstrating that collateral cleavage is not the

result of the iTOP delivery method per se, and does not require targeting an overexpressed

exogenous transcript (Supplementary figure 6A).

To explore if the collateral RNA cleavage also occurs when LbuCas13a is overexpressed in the

cell, we transiently transfected an LbuCas13a expression plasmid into HAP1-dEGFP cells. The

transiently transfected cells showed a clear LbuCas13a band on western blot (supplementary

figure 7A), despite an average transfection efficiency of only 9.2%. We enriched successfully

transfected cells by FACS two days after plasmid transfection. Both the unsorted and enriched

cells were transfected with a dEGFP-targeting guide RNA after allowing the enriched cells to

recover overnight from sorting. No collateral cleavage pattern was observed in the protein

expressing cells transfected with guide RNA (Supplementary figure 7B). Altogether, our data

demonstrate that LbuCas13a is capable of collateral cleavage, targeting both exogenous and

endogenous transcripts, and using two different RNP delivery methods. However, no collateral

cleavage was observed when the LbuCas13a protein was expressed.

Collateral cleavage occurs in all tested cell lines

Since previous reports have demonstrated that Cas13 activity can be cell-line dependent (Bot et

al., 2022), we next explored whether collateral RNA cleavage by LbuCas13a RNP delivery was

also cell-type dependent. We used iTOP to transfect the chronic myeloid leukemia derived line

HAP1, the rhabdomyosarcoma line RD, the osteosarcoma line U2OS, and the retinal pigment

epithelia line ARPE19 with LbuCas13a targeting the highly abundant endogenous transcripts

RPS19, GAPDH, or 18S rRNA. All four cell lines demonstrated robust collateral cleavage after

targeting any of these target transcripts (Figure 2A). Across all four cell lines, 18S rRNA

targeting resulted in the most robust collateral RNA cleavage activity, followed by GAPDH and

RPS19 respectively. Notably, the degradation pattern was exactly the same in all conditions and

does not depend on the target RNA or cell line. This consistent RNA degradation pattern

allowed us to quantify the total RNA degradation by taking the area under the curve of the

region where most of the cleaved RNA fragments occur (See Figure 2A). HAP1 cells

consistently demonstrated significantly more total RNA degradation than the other three cell

lines (Figure 2B). Next, we utilized the 5'-3' RT-qPCR assay to further quantify the collateral

cleavage effect in these samples. RNA degradation was presented by both an overall increase

in Cq values compared to untreated cells as well as an increased difference between the 5' and

3' primer pair (Figure 2C), with HAP1 again showing the biggest effect. HAP1 cells are known

as near haploid cells, however they diploidize quickly in just several passages (Yaguchi et al,

2018). The cells we used in this study are mostly diploid. A possible reason for the higher

collateral activity in HAP1 cells could be a higher RNP delivery efficiency, as iTOP was

developed using this cell type (D'Astolfo et al, 2015).

Several reports showing specific RNA knockdown with Cas13 used HEK293T cells (Abudayyeh

et al., 2017; Cox et al., 2017; Konermann et al., 2018), while others explicitly report low or even

the absence of collateral cleavage in this cell type (Ai et al., 2022; Wang et al., 2019),

suggesting that Cas13 may be less active in HEK293T cells. Using electroporation to deliver

LbuCas13a protein and guide RNA into HEK293T cells, we observed the characteristic RNA

degradation pattern upon targeting GAPDH and 18S rRNA (Supplemental figure 6B), indicating

that HEK293T cells are not completely void of LbuCas13a collateral cleavage activity. However,

in agreement with above mentioned publications, the extent of the collateral cleavage was

decreased compared to HAP1 when targeting GAPDH and even undetectable when targeting

the highly abundant RPS19 transcript.

RNA targeting with LbuCas13a induces cell death via apoptosis

Over the course of our experiments we noticed that a day after targeting dEGFP with LbuCas13a cells were rounding up and detaching from the bottom of the plate (Figure 3A), implying loss of viability. To examine this further, we performed live cell imaging with Annexin V as a marker for early apoptosis and CellTox Green to stain for dead cells (Wallberg et al., 2016). When cells die by necrosis they become positive for both these dyes simultaneously. However, if they go into apoptosis Annexin V staining precedes CellTox green staining. We transfected LbuCas13a and one of three different guide RNAs targeting dEGFP into either wildtype or dEGFP expressing cells HAP1. In wildtype cells the number of Annexin V and CellTox green positive cells was similar to the empty transfection control for all three dEGFP targeting guide RNAs (Figure 3B). However, in HAP1-dEGFP the number of Annexin V positive cells rapidly increased from approximately 7.5 hours after transfection, followed by an increase in CellTox Green positive cells approximately 15 hours after transfection (Figure 3B and Supplementary movies 1 and 2). This shows that LbuCas13a is capable of inducing cell death, and that this is specific to cell expressing the target RNA. Furthermore, it demonstrates that the targeted cells go into apoptosis, not necrosis.

Application of LbuCas13a as a cell selection tool

Given that target-activated collateral RNA cleavage by LbuCas13a can induce cell death, we wondered whether we could exploit this property to eliminate cells that express a specific target RNA from a heterogeneous cell population (Figure 4A). To test this, we mixed fluorescent EGFP expressing HAP1-EGFP cells with wildtype HAP1 cells and transfected this mixed population with LbuCas13a and a guide RNA targeting EGFP. It is important to note that this EGFP targeting guide does not cause collateral RNA cleavage or reduced viability in wildtype cells (Figure 1B and Figure 3B). We then measured the percentage of EGFP positive cells by FACS over time. The percentage EGFP positive cells rapidly decreased when targeting EGFP (Figure 4B), with 36.3% EGFP-positive cells remaining at day 6 compared to more than 60% EGFP positive cells when using a non-targeting guide. The decrease in GFP positive cells was maintained over the entire six-day assay, showing that it was not due to temporary knockdown of the GFP transcript itself, as the GFP transcript integrity is largely restored after 500 minutes

to 24 hours after RNP transfection (see Figure 1C). Next, we examined whether multiple

sequential selection events could further eliminate the EGFP positive cell population (Figure

4C). We performed several serial transfections at 2-day intervals, with up to 3 subsequent

selection rounds. Every additional selection round decreased the percentage of EGFP positive

cells in the population by approximately 20 percentage points. From the above data we

conclude that Cas13-based cell-selection is able to reduce the target cell population and is

repeatable.

Target RNA expression level theoretically limits the number of active LbuCas13a molecules and

therefore the amount of collateral cleavage, which could influence the efficiency of cell selection.

To test this, we FACS sorted a heterogeneous population of EGFP expressing cells into cell

populations with different levels of EGFP intensity (Figure 4D and Supplementary figure 8A).

RT-qPCR confirmed EGFP expression levels increased with EGFP intensity (Supplementary

figure 8B). We then tested the depletion efficiency of these cells by mixing them with wildtype

HAP1 cells and targeting EGFP with LbuCas13a. In parallel we also performed RT-qPCR on

these cell lines to quantify their EGFP expression at the time of the experiment. We found a

strong correlation between the expression level of the target RNA and the depletion of EGFP

positive cells (Figure 4E), suggesting that activating more LbuCas13a proteins results in more

collateral RNA cleavage and a high probability of inducing cell death. However, depletion

efficiency seems to plateau at the highest levels of target RNA expression. At this point the RNP

delivery efficiency may have become the limiting factor.

We next examined whether the same strategy could be used to positively select for cells that

underwent successful gene editing. For these experiments we used a HAP1-EGFP$^{\Delta fluor}$ cell line

containing a 9 nt deletion in EGFP, rendering it non-fluorescent (Supplementary figure 8C)

(Zhao et al., 2022). Using this system, we previously reported Cas12a-mediated correction of

EGFP fluorescence, with an efficiency of about 5% (Zhao et al., 2022). When we employed

LbuCas13a and a guide RNA that targets the unedited cells as a cell selection tool, the

percentage of EGFP expressing cells increased to around 15% (Figure 4F). Thus, a single

round of selection using LbuCas13a resulted in a 3-fold enrichment. Altogether, these proof-of-

principle studies show the potential of Cas13 as a potent cell selection tool, including the

enrichment of successfully edited cells.

Genes involved in the innate immune system are upregulated after collateral RNA cleavage

To gain further insight into the cellular response to the collateral RNA cleavage and the

mechanism by which it triggers apoptosis, we performed RNA sequencing at 16 hours after

transfection. Based on the 5'-3' RT-qPCR data (Figure 1C), at this timepoint most mRNA

cleavage has been resolved, allowing us to focus on the cellular response following the

collateral cleavage. STAR (Dobin et al., 2013) aligned sequences at similar rates across all

conditions, however, a higher percentage of the reads from the dEGFP targeting samples

mapped to introns compared to the controls, consistent with a relative increase in immature

nuclear transcripts generated as a result of cytoplasmic RNA depletion in the cell

(Supplementary figure 9A). First, we performed principal component analysis (PCA)

(Supplementary figure 9B). The PCA plot shows a clear separation between the dEGFP

targeting samples and the controls, suggesting that collateral cleavage by LbuCas13a leads to

substantial changes in the transcriptome. Indeed, we found 721 significantly upregulated and 36

significantly downregulated genes (adjusted P-value < 0.05, |Log2 FC| > 1) in response to

LbuCas13a activation (Figure 5A). The relatively low number of downregulated genes and high

number of upregulated genes is perhaps somewhat counterintuitive in the context of collateral

cleavage, but consistent with another report documenting the cellular response to Cas13

collateral cleavage (Y. Li et al., 2023). In agreement with Li et al., we observed that many of the

upregulated genes we find are immediate early response genes, such as JUN, FOS, EGR1,

EGR2, EGR3, EGR4 and ARC. GO enrichment analysis of the significantly upregulated genes

revealed an enrichment of many biological processes related to the innate cellular immune

response (Figure 5B), and molecular functions related to chemokine and cytokine activity

(Supplementary figure 9C). KEGG pathway analysis also showed enrichment for immune

related pathways, such as viral protein interaction with cytokine and cytokine receptor, IL17

signaling pathway, TNF signaling pathway and NF-kappa B signaling pathway (Supplementary

figure 9D). These are known factors of the cellular immune response that are triggered, for

example, by intracellular recognition of viral RNA and suggests that collateral RNA cleavage by

LbuCas13a and subsequent increase in 5' uncapped RNA and 3' ends without poly-A tail

triggers a similar immune response which ultimately leads to apoptosis.

Characterizing collateral RNA cleavage using nanopore sequencing

We decided to use Nanopore sequencing to explore the identity of the RNAs that are subject to

collateral cleavage, where RNAs get cleaved by LbuCas13a, and the dynamics of the collateral

cleavage process. HAP1-dEGFP cells were transfected with LbuCas13a protein and either a

non-targeting or a dEGFP targeting guide RNA. Nanopore read lengths of dEGFP targeting

samples were much shorter than control samples (Supplementary figure 10C), containing

markedly fewer reads longer than 1000 nt, suggesting widespread collateral RNA cleavage. We

found that when targeting dEGFP, the number of protein coding transcripts decreased relative

to all other transcripts biotypes (Supplementary figure 11A). On the other hand, the percentage

of reads mapping to lncRNAs, snoRNAs, and (in replicate 1) snRNAs increased, suggesting that

nuclear RNAs were relatively protected from collateral cleavage. The abundance and read

length of reads mapped to protein coding transcripts greatly decreased at 50 and 200 minutes

after transfection (Figure 6A and Supplementary figure 11B). At these timepoints, almost all

reads were much shorter than the reference they mapped to, whereas most reads in the empty

transfection and non-targeting controls were full length transcripts. This suggests unbiased

cleavage of all mRNAs after LbuCas13a activation by dEGFP.

Next we explored the dynamics of cleavage of individual transcripts by calculating their relative

transcript coverage. We found that at 50, 200 and 500 minutes after transfection, the overall

coverage of dEGFP-IRIS-PuroR is much lower for the dEGFP targeting samples than the

control conditions (Figure 6B). Additionally, the target RNA shows a decrease in coverage at the

exact location of the protospacer at all timepoints (Figure 6B, dotted lines), which fits with the 5'-

3' RT-qPCR data that showed target RNA cleavage immediately after transfection. This suggest

LbuCas13a cleaves in, or very near the protospacer, which is surprising since others have

shown that Cas13 cleavage sites do not depend on protospacer position (Abudayyeh et al.,

2016; Konermann et al., 2018; Smargon et al., 2017). Alternatively, the protospacer specific dip

in coverage may be a technical artifact, for example, due to LbuCas13a still being bound to the

target RNA after purification, preventing reverse transcription at this location. Besides the

coverage dip around the protospacer, additional dips in coverage were observed, which were

consistent between timepoints and replicates. We suspect that these dips are LbuCas13a
cleavage sites. Non-target mRNAs and cytoplasmic non-coding RNAs also show these dips in
coverage at specific positions, indicating collateral cleavage of these transcripts (Figure 6C,
Supplementary figures 12 and 13). We did not observe any differences in coverage between
targeting and control samples for small nuclear or mitochondrial RNAs (Supplementary figure
14). The dips in transcript coverage tend to be 10-20 nucleotides wide, so we cannot discern the
exact cleavage position (Figure 7A). *In vitro*, LbuCas13a was shown to prefer cleaving at uracil
residues in or near single stranded loop structures (East-Seletsky et al., 2017). In our data,
transcript coverage dips often map to predicted stem-loop structures, with uracil containing
single stranded loops on the 5' side of the coverage dip (Figure 7B), suggesting that the 5' side
of the dip is the actual cleavage location. These results indicate that LbuCas13a cleavage
preferences follow the same pattern in human cells as observed *in vitro*. Next, we manually
annotated 100 potential cleavage sites (Supplementary table 4), assuming that the Uracil
closest to the 5' side of the coverage dip was the cleavage position. This revealed a US (S = G
or C) di-nucleotide preference (Figure 7C). Taken together, the Nanopore sequencing provides
further evidence of collateral RNA cleavage in human cells by LbuCas13a and suggests that
both mRNAs and cytoplasmic ncRNAs are cleaved without bias. Additionally, LbuCas13a
cleaves RNA at specific positions, showing a preference for uracil-containing single stranded
loop structures.

Discussion

In this work, we revealed the temporal dynamics of both target and collateral RNA cleavage by
LbuCas13a, as well as the cellular responses to this collateral RNA cleavage (Figure 8). Target
and collateral RNA cleavage start within 50 minutes after RNP delivery, with target cleavage
preceding collateral RNA cleavage. All cytoplasmic RNAs seems to be cleaved in an unbiased
manner, with a cleavage preference for Uracil containing stem loop structures. Target and
collateral mRNA integrity mostly returns to normal at around 500 minutes after RNP delivery,
while ribosomal RNA takes longer to recover, with some cleavage fragments still being present
at 24 hours after delivery. The recovery of mRNA integrity coincides with the cells starting to
become positive for Annexin V, indicating early apoptosis, at about 6-8 hours after delivery.
Stress and innate immune response genes are upregulated, probably contributing to the

induction of apoptosis. From about 15 hours after delivery, cell death starts occurring. As such,

LbuCas13a can be used as a negative cell selection tool.

The collateral RNA cleavage activity of Cas13 is well established *in vitro* (Abudayyeh et al.,

2016; East-Seletsky et al., 2017, 2016; Konermann et al., 2018; Smargon et al., 2017; Xu et al.,

2021; Yan et al., 2018) and in bacteria (Abudayyeh et al., 2016; Meeske et al., 2019; Smargon

et al., 2017; Yan et al., 2018), however in eukaryotes the situation is less clear. While many

groups report that no collateral cleavage seems to occur (Abudayyeh et al., 2017; Cox et al.,

2017; Konermann et al., 2018; Tieu et al., 2024; Wessels et al., 2024), others have recently run

into unexpected cytotoxicity and/or apparent collateral RNA cleavage when using CRISPR-

Cas13 in eukaryotic cells (Ai et al., 2022; Buchman et al., 2020; Y. Li et al., 2023; Özcan et al.,

2021; Shi et al., 2021; Tong et al., 2021; Wang et al., 2019). From these reports and our work,

several factors have emerged that are likely important in determining if and how much collateral

cleavage occurs, namely the collateral activity of the ortholog, the intracellular RNP

concentration, the target RNA expression level and cell line specific differences (Bot et al.,

2022).

When comparing the collateral activity of several orthologs *in vitro*, in our hands LbuCas13a

was the most active ortholog. However, different orthologs may have different preferences for

factors such as the assay buffer, the protospacer sequence (Wessels et al., 2024, 2020) and

the cleavage substrate (East-Seletsky et al., 2017). When comparing the collateral RNA

cleavage activities of these Cas13 orthologs in cells, only LbuCas13a exhibited robust collateral

cleavage across a range of different guide RNAs. While other reports have documented

collateral cleavage or toxicity when using LwaCas13a (Özcan et al., 2021; Wang et al., 2019),

or RfxCas13d (Ai et al., 2022; Y. Li et al., 2023; Shi et al., 2023), we did not find collateral RNA

cleavage activity in human cells with these orthologs (see supplementary note 1).

We detected collateral cleavage when delivering LbuCas13a as RNP, but not when LbuCas13a

was expressed as a transgene. Cas13 expression and protein levels may be limited by guide

RNA independent RNA cleavage activity of Cas13 (Z. Li et al., 2023), potentially explaining why

only RNP delivery allowed us to detect collateral RNA cleavage. Here we showed that by

combining the highly active LbuCas13a with RNP delivery, apoptosis can be induced in target

RNA expressing cells. This allowed us to specifically deplete target cells from a mixed

population. With just one round of selection, we reach a similar efficiency as Shi and colleagues

(Shi et al., 2023). However, unlike Shi et al., we show that multiple selection rounds are

possible, achieving much greater negative selection efficiency, namely reducing target cells

from 80% to just 13% of the population. In addition, Shi and colleagues use cells stably

expressing RfxCas13d, which requires the generation of such an RfxCas13s expression cell line

before cell selection is possible. By transfecting Cas13 as RNP, our system can be immediately

applied to any cell type without the need for introducing any transgenes. Finally, to our

knowledge we are the first to show that Cas13 can be applied to enrich for cells that have

undergone a rare gene editing event, achieving a three times enrichment. Enriching for edited

cells with CRISPR-Cas13 has major advantages over other methods that select for genome

editing events, as it does not require the co-insertion of a selection marker (Shy et al., 2016) or

a second selectable edit (Ewen-Campen and Perrimon, 2018). Furthermore, it does not depend

on co-expression of gene editing components with a selection marker, which risks selection for

permanent genomic integration of the gene editing plasmid.

Although we were able to deplete cells from a mixed population, we were not able to completely

eliminate all target cells. In addition, high target RNA expression is required to enable the

induction of cell death, which limits the potential target RNAs that can be used. Nevertheless,

we are optimistic that future iterations of the CRISPR-Cas13 cell selection system will overcome

these limitations. There are several potential strategies to improve the CRISPR-Cas13 cell

selection systems. For example, the number of selection rounds could be increased, since we

observed no indication of reduced efficiency with subsequent selection rounds. Furthermore, the

sensitivity to collateral cleavage of cells might differ between different phases of the cell cycle,

as it does for many anti-cancer drugs (Johnson et al., 2021), so cell cycle synchronization may

be a factor to consider as well. Finally, orthologs with even higher collateral RNA cleavage

activity than LbuCas13a could be discovered or engineered through directed protein evolution

approaches.

In conclusion, here we demonstrated that LbuCas13a is capable of high collateral cleavage

activity in human cell lines. We described in detail the events occurring following activation of

LbuCas13a by a target RNA. Finally, we showed how CRISPR-Cas13 can be employed as a

specific and effective cell elimination tool. We envision that this new system will find broad

application in both basic and clinical research, enabling the selective elimination of unedited

cells, specific cell types to improve stem cell differentiation yields or perhaps the specific

elimination of cancer cells.

Materials and Methods

Contact for reagent and resource sharing

434 Recombinant LbuCas13a and AsCas12a proteins, the HAP1-dEGFPY66S and the HAP1-

435 dEGFPΔfluor cell lines are available through Divvly (https://divvly.com/geijsenlab).

Molecular cloning

For recombinant protein production, all Cas13 orthologs were cloned into a pET15b expression

vector (Addgene: 62731). LbuCas13a and LshCas13a were a gift from Jennifer Doudna

(Addgene: 83482 and 83487). LwaCas13a, BzCas13b and PspCas13b were a gift from Feng

Zhang (Addgene: 91924, 89898 and 103862). RspCas13d was a gift from Arbor

Biotechnologies (Addgene: 108304). pET-28b-RfxCas13d-His was a gift from Ariel Bazzini &

Miguel Angel Moreno-Mateos (Addgene: 141322). Cloning was done using standard molecular

biology techniques using In-Fusion cloning according to the manufacturer's instruction. For the

construction of the lentiviral LbuCas13a and RfxCas13d protein and guide expression vectors,

all fragments were ordered from IDT as gBlocks. The fragments were cloned into the pEF-

1a_dEGFP^{Y66S}-IRIS-PuroR-WPRE lentiviral plasmid backbone. Cloning was done using

standard molecular biology techniques using In-Fusion cloning according to the manufacturer's

instruction. All plasmid sequences are available in supplementary file 1. Plasmids will be made

available on Addgene.

Cell culture

HAP1 cells were a gift from Dr. Brummelkamp, NKI Amsterdam. RD cells were a gift from Dr.

Gerben Schaaf, Erasmus MC. HEK293T cells were ordered from Takara (#632180). U2OS and

ARPE19 cells were ordered from ATCC (HTB-96 and CRL-2302 respectively). All cell lines were

tested for mycoplasma monthly.

HAP1 cells were cultured in Iscove's Modified Dulbecco's Medium (IMDM) supplemented with

10% Fetal Bovine Serum (FBS). HEK293T cells were cultured in Dulbecco's Modified Eagle

Medium (DMEM) with 10% FBS. U2OS and RD cells were cultured in DMEM supplemented

with 10% FBS and 2 mM L-Glutamine. ARPE19 cells were cultured in DMEM/F12 with 10%

FBS. All culture medium was purchased from Gibco. All cells were grown at 37°C in a

humidified atmosphere containing 5% CO_2.

To generate the HAP1-dEGFP^{Y66S}, HAP1-dEGPF$^{\triangle fluor}$ and HAP1-EGPF lines (Zhao et al.,

2022), HAP1 cells were transduced with the pEF-1a_dEGFP^{Y66S}-IRIS-PuroR-WPRE, pEF-

1a_EGFP$^{\triangle fluor}$-IRIS-PuroR-WPRE or pEF-1a_EGFP-IRIS-PuroR-WPRE plasmids

(Supplementary file 1) using lentivirus. Two days after transduction, puromycin selection was

started at 0.5 µg/mL to select for integration of the plasmid. After resistant cells had established

themselves, the puromycin concentration was increased every 2 days to 1, 2, 5 and finally 10

µg/mL. HAP1-dEGFP^{Y66S}, HAP1-dEGFP$^{\triangle fluor}$ and HAP1-EGFP cells were maintained in HAP1

medium supplemented with 10 µg/mL puromycin.

To generate the LbuCas13a and RfxCas13d protein or guide RNA expressing cells, HAP1-

dEGFP cells were transduced with lentivirus (See Supplementary file 1 for plasmid sequences).

Two days after transduction, blasticidin selection was started at 10 µg/mL. Because the

percentage of BFP positive cells remained low, cells were sorted for being BFP or mScarlet

positive at 14 after infection on a CytoFLEX SRT benchtop cell sorter (Beckman Coulter). Cells

were maintained in HAP1 medium supplemented with 10 µg/mL blasticidin.

Lentivirus production

Lentivirus was produced in HEK293T cells. Briefly, 4 packaging plasmids pHDM-Hgpm2,

pHDM-Rev1b, pHDM-Tat1b, pHDM-G and the transfer plasmid (in the ratio of 2:1:1:2:4) were

transfected to HEK293T cells with FuGENE HD transfection reagent (Promega, E2311)

according to the manufacturer's protocol. The FuGENE to DNA ratio was 2.5:1. One day after

transfection, the cell culture medium was supplemented with medium containing caffeine

(Sigma C0750) and HEPES (Gibco 15630-080), to a final concentration of 3 mM and 25 mM

respectively. The supernatant containing the lentivirus was harvested 48 hours after

transfection, filtered with 0.22 μm filter (EMD Millipore, SCGP00525), and stored at 4°C.

Expression and purification of recombinant Cas13 proteins

The sequence of the expression vectors is available in supplementary file 1. Expression vectors

will be made available on Addgene. The different recombinant Cas13 orthologs were produced

as described previously for Cas12 (Zhao et al., 2022). In brief, the proteins were expressed in *E.*

coli BL21(DE3) and purified by Ni-NTA affinity purification followed by gel-filtration

chromatography. The purified Cas13 proteins were concentrated to 75 μM using the Amicon

ultra centrifugal filter MWCO 50 kDa (Merck), flash frozen in liquid nitrogen, and stored at -80°C.

Preparation of guide RNAs and target RNA by in vitro transcription

Oligonucleotides carrying the T7 promoter and appropriate downstream sequence were

synthesized by IDT and PCR-amplified to generate *in vitro* transcription (IVT) templates (see

supplementary table 1 for template and primer sequences). The EF-1a_dEGFP^{Y66S}-IRIS-PuroR-

WPRE plasmid (supplementary file 1) was used as PCR template to generate the IVT template

for the target RNA. Transcription reactions were performed at 37°C for 4 hours in a buffer

containing 40 mM Tris-HCl (pH 8), 20 mM $MgCl_2$, 5 mM DTT, 2 mM NTPs, 50 μg/mL T7 RNA

polymerase, and 250 ng/μL DNA template. Next, to remove the DNA template, 1 μL Turbo

DNase (Invitrogen: AM2238) per 100 μL IVT reaction was added and incubated at 37°C for 30

minutes. The RNA was then purified using the Zymo RNA Clean & Concentrator -25 kit

(R1017). Typically, four different guide RNAs were tested for each target RNA, assessing

collateral cleavage by comparing total RNA degradation using the Bioanalyzer (see below). The

guide RNA with the highest amount of total RNA degradation was selected for use in

experiments.

In vitro collateral cleavage activity assay

Collateral RNA cleavage detection assays were performed with IDT RNaseAlert (IDT, 11-04-02-

03) according to the manufacturer's protocol. Reactions were performed in different cleavage

buffers depending on the Cas13 ortholog. For LshCas13a, LwaCas13a, LbuCas13a and

RfxCas13d the buffer (Abudayyeh et al., 2016) consisted of 40 mM Tris-HCl pH 7.3, 60 mM

NaCl and 6 mM MgCl$_2$. For Cas13b (BzCas13b and PspCas13b), the cleavage buffer contained

10 mM Tris-HCl pH 7.5, 50 mM NaCl and 0.5 mM MgCl$_2$ (Smargon et al., 2017). For

RspCas13d, the cleavage buffer used was 20 mM HEPES pH 7.1, 50 mM KCl, 5 mM MgCl$_2$ and

5% glycerol (Yan et al., 2018). Before the RNaseAlert assay, 1 µM Cas13 and 500 nM guide

RNA were preassembled in 10X cleavage buffer for 10 min at 37°C. The RNaseAlert assays

were performed using 100 nM Cas13 protein and 50 nM guide RNA with the appropriate

amount of target RNA, using a 50 µL reaction volume. Reactions were prepared on ice, in

opaque 96-well plates (Greiner, 655076). Fluorescence was measured every 5 minutes using a

Tecan Spark plate reader, while incubating at 37°C. Reactions were allowed to proceed for at

least one hour. The collateral cleavage rate was calculated in Excel by applying the SLOPE

function to the initial linear increase in fluorescence.

Ribonucleoprotein transfection using iTOP

The Cas13 proteins and their guide RNAs were transfected into target cells using iTOP, using a

protocol adapted from D'Astolfo et al. (D'Astolfo et al., 2015). The iTOP protocol consists of four

steps: (1) seeding the cells, (2) iTOP transfection mixture preparation, (3) incubating the cells in

the iTOP mixture, and (4) recovery. First, cells were plated the day before transfection in

Matrigel coated pates (Corning: 356231, diluted 1:100 in PBS), such that the confluency was

~70% the next day. Second, to prepare the iTOP transfection mixture, 2/5 of transfection

supplement B (Opti-MEM media supplemented with 810 mM NaCl, 504 mM GABA, 1.875X

concentrated N2 (Gibco: 17502048), 1.875X concentrated B27 (Gibco: 17504044), 1.875X

concentrated MEM non-essential amino acids (Gibco: 11140050), 3.75 mM L-Glutamine (Gibco:

25030081), 2.5 µg/mL bFGF (Gibco: PHG0023), and 0.625 µg/mL EGF (Gibco: PHG0313)), 1/5

of Cas13 protein (75 µM) or empty protein storage buffer (Zhao et al., 2022), 1/5 of guide RNA

(75 µM) or nuclease-free water, and finally 1/5 nuclease-free water were combined. iTOP

mixture volumes used per well were: 50 µL for 96-well plates, 150 µL for 48-well plates, 200 µL

for µ-slide 8-well (ibidi, 80827) and 1000 µL for 6-well plates. Third, the culture medium was

aspirated and iTOP mixture was carefully added into the wells. Next the cells were returned to

the cell culture incubator for 45 minutes. Fourth, immediately after incubation the iTOP mixture

was aspirated and replaced by pre-warmed cell culture medium (96-well: 200 µL, 48-well: 600

µL, µ-slide 8-well: 300 µL, 6-well: 4000 µL). Finally cells were returned to the cell culture

incubator or immediately harvested for downstream analysis.

Ribonucleoprotein transfection using electroporation

A Lonza Nucleofector system including a Nucleofector 2b Device and the Cell Line Nucleofector

Kit V (Lonza) were used for electroporation of HAP1-dEGFP and HEK293T cells targeting

endogenous transcripts (Supplementary figure 6A and 6B), according to the manufactures

instructions. One million cells were used per electroporation, resuspended in 100 µL

Nucleofector solution V buffer containing 1 µM LbuCas13a and 1 µM guide RNA. The

electroporation was performed in the Nucleofector 2b device using the 'Cell-line T-020' program.

For the comparison of LbuCas13a with RfxCas13d (Supplementary figure 3 and 5C), HAP1-

dEGFP cells were electroporated using a Bio-Rad Gene Pulser Xcell. One million cells were

resuspended in Gene Pulser electroporation buffer (Bio-Rad #1652677) containing 1 µM Cas13

protein and 1 µM guide RNA in a total volume of 150 µl. Cells were electroporated in 4mm

cuvettes using a single 400V, 5ms square wave. After electroporation, the cells were incubated

at 37°C.

Plasmid transfection

Cells were plated in 6-well plates such that the confluence was ~80% at the time of transfection.

Five µg plasmid DNA was mixed with 15 µg PEI (MW 25000, Polysciences #23966) in a total

volume of 500 µl opti-MEM (Gibco #31985070). The mixture was incubated at room

temperature for twenty minutes. After incubation, the mixture was added dropwise to the culture

medium in the well. Next, the 6-well plate was spun at 200g for five minutes, after which the

cells were return to the incubator.

Total RNA isolation

Cells were collected in TRIzol reagent (Invitrogen: 15596018) and stored at -80°C until RNA

isolation. To isolate total RNA, chloroform was added to the TRIzol at a ratio of 1:5 and samples

were mixed by briefly vortexing. Next, samples were centrifuged for 15 minutes at 12000 g at

4°C. Total RNA was isolated from the colorless upper aqueous phase using the Zymo RNA

Clean & Concentrator -5 kit, including an in-column DNase treatment, according to the

manufacturer's instructions.

Bioanalyzer

Total RNA integrity was analyzed on the Bioanalyzer using either the RNA nano or RNA pico chips, depending on the available RNA amount. The area under the curve (AUC) was calculated in R with the "AUC" function from the DescTools package using the default settings (Signorell, 2021). To account for differences in the amount of RNA loaded, all samples were normalized to their total AUC. This was done by dividing the sample fluorescence by the sample AUC and multiplying by the mean AUC across all samples. In our hands, this method resulted in a more accurate normalization than using the sample concentration estimates supplied by the Bioanalyzer software.

5'-3' RT-qPCR

RNP transfection by iTOP was performed in 96-well plates, as described above. Besides LbuCas13a and either a targeting or a non-targeting guide RNA, an empty transfection control was always included. All ddCq values are relative to this empty transfection control. Total RNA was isolated and DNase-treated as described above. An anchored oligo(dT) primer was used for reverse transcription using SuperScript III reverse transcriptase (Invitrogen, 18080093), according to the manufacturer's protocol. qPCR was performed in triplicate using Bio-Rad iQ SYBR Green Supermix with 300 nM primers. The primer sequences used for reverse transcription and qPCR can be found in supplementary table 2.

Traditionally, the ratio between the Cq values of the 3' and 5' primer pair has been used to quantify transcript integrity (Nolan et al., 2006). However, this will results in different ratios for the same amount of degradation when expression levels are different ($\frac{20}{25} \neq \frac{25}{30}$). Therefore, to quantify the transcript integrity we used a modified version of the ddCq method where the 3' primer pair takes on the function of the housekeeping gene. First, to calculate the 5'-3' dCq values, the 3' Cq value was subtracted from the 5' Cq value for each sample. Next, to calculate the 5'-3' ddCq, the mean 5'-3' dCq value of the empty transfection controls was subtracted from the 5'-3' dCq values of the treatments. To avoid confusion with the method were the ratio between the 3' and 5' primer pair is used, we called the assay 5'-3' RT-qPCR here instead.

Western blot

First, the cells were put on ice and washed three times with ice cold PBS. Cells were collected in ice cold RIPA buffer and kept on ice for 20 minutes. The lysate was spun down at 16000g at 4°C for 20 minutes. The supernatant was transferred to a new tube and the protein concentration was measured using a BCA assay (Thermo Scientific, 23225). Before loading on gel, samples were mixed with Laemmli buffer (Bio-Rad, 1610747) and heat denatured at 95°C for 5 minutes. Twenty µg protein lysate was loaded per well, unless otherwise indicated. Samples were run on 4-20% Mini-PROTEAN precast gels (Bio-Rad, 4568094). The protein was then transferred to a PVDF membrane using a wet transfer at 300 mA for 90 minutes. After transfer, the membrane was blocked in blocking buffer (PBS with 5% w/v skim milk powder, 1% w/v BSA, and 0.1% Tween20) at room temperature for twenty minutes. Next, the membrane was stained overnight in blocking buffer with an anti-HA tag (ThermoFisher, 26183) and a GAPDH loading control antibody (ThermoFisher, MA5-15738), diluted at 10000x and 1000x respectively. Next, the membrane was washed three times for five minutes in PBST (PBS with 0.1% Tween20). The membrane was stained for two hours at room temperature with Goat Anti-Mouse IgG H&L (HRP) (Abcam, ab6789), diluted 1:1000 in blocking buffer. After staining the membrane was again washed three times for five minutes in PBST. Finally, ECL substrate (ThermoFisher, 32106) was applied to the membrane and the signal was detected on a Bio-Rad GelDoc Go System.

Nanopore cDNA sequencing

The experiment and subsequent library preparation consisted of seven steps: (1) RNP transfection, (2) RNA isolation, (3) rRNA depletion, (4) poly(A) tailing, (5) reverse transcription, (6) second strand synthesis and (7) barcode and sequencing adapter ligation. Manufacturer's instructions were followed for all kits unless otherwise noted.

To ensure enough RNA for sequencing, this experiment was performed in 6-well plates, plating 600 000 HAP1-dEGFP^{Y66S} cells per well the day before the transfections. One well was reserved as a no transfection control. iTOP was performed as described above, using an iTOP mixture volume of 1 mL per well. Each timepoint had three conditions: an empty transfection control, a transfection with LbuCas13a and a non-targeting guide RNA, and a transfection with LbuCas13a and a dEGFP targeting guide RNA (SP2). Cells were collected in TRIzol at 0, 50,

100, 200, 400 and 1440 minutes after iTOP. Total RNA was isolated and DNase-treated as

described above. RNA integrity was assessed on the Bioanalyzer (Supplementary figure 10A).

Next, 5 µg total RNA of each sample was used for rRNA depletion with the RiboMinus

Eukaryotic system v2 (Invitrogen, A15026). rRNA depletion was confirmed with the Bioanalyzer

(Supplementary figure 10B). In order to also detect cleaved RNA fragments without a poly-A tail,

we polyadenylated the rRNA depleted RNA with *E. coli* Poly(A) polymerase (NEB, M0276L) in

the presence of 1 mM ATP and 1 unit/uL RNasin® (Promega, N2515) for 30 minutes at 37^{o}C.

Afterwards, the RNA was cleaned with the Zymo RNA Clean & Concentrator -5 Kit (R1013).

Next, reverse transcription was performed using SuperScript IV RT (Invitrogen, 18090010) and

anchored oligo dT$_{20}$ primers (Supplementary table 2). The reverse transcription reaction was

incubated at 50^{o}C for 15 minutes followed by 80^{o}C for 10 minutes. To remove RNA before

second strand synthesis, 1 uL of RNase Cocktail (Thermo, AM2286) was added after reverse

transcription and incubated at 37^{o}C for 20 minutes. The NEBNext second strand synthesis kit

(NEB, E6111) with 100ng random hexamer primers (Promega, C1181) per sample was used for

second strand synthesis. Next, dsDNA was purified using AMPure XP beads (Beckman Coulter:

A63882) at a ratio of 2 beads to 1 sample. The NEBNext Ultra II End Repair/dA-tailing Module

(NEB, E7546S) was used before ligating Nanopore native barcodes (EXP-NBD104&114).

Finally, samples were pooled at equal mass and the sequencing adapter (SQK-LSK109) was

ligated. 1000 ng DNA was loaded on the Promethion flow cell (FLO-PRO002) and sequenced

for 3 days. Two independent replicates were performed.

Nanopore data analysis

During sequencing, the Nanopore MinKNOW software (version 21.11.7, Guppy version 5.1.13)

called bases in real-time by the using the 'Super-accurate' base calling setting, demultiplexed

samples and trimmed barcodes. For replicate 1, the 0 minute after transfection, dEGFP

targeting sample had a very low DNA yield after library preparation and consequently much

fewer reads than the other samples. Therefore, it was excluded from all analysis. Reads were

mapped to the transcriptome using minimap2 (Li, 2018). Transcriptome alignment was done

using the *-ax map-ont* arguments, against a merge of the Ensemble

Homo_sapiens.GRCh38.cdna.all (release 104), Ensemble Homo_sapiens.GRCh38.ncrna

(release 104), the entire pEF-1a_dEGFP[Y66S]-IRIS-PuroR-WPRE plasmid (Supplementary file 1)

and the 18S and 28S ribosomal RNA. Mapping rates were similar across samples, with dEGFP

targeting samples showing a small decrease at 500 and 1440 minutes after transfection

(Supplementary figure 15A). Only the primary alignment was used for analysis. To generate the

density plots in Figure 6A and Supplementary figure 11B, all histone transcripts and WDR74

were excluded. Histone transcripts and WDR74 were so abundant that they obscured all other

mRNAs in the density plot. The extremely high abundance of WDR74 was caused by a small

nuclear RNA mapping to WDR74-204. All analysis was performed in Rstudio, using data.table

(Dowle and Srinivasan, 2022) and ShortRead (Morgan et al., 2009) for data loading, Tidyverse

(Wickham et al., 2019) for data handling and ggplot2 (Wickham, 2010) for plotting. Raw

sequencing reads are available online at https://www.ncbi.nlm.nih.gov/sra/PRJNA912090. The

script used for calculating transcript coverages is available on

https://github.com/Geijsenlab/cas13.

RNA sequencing and analysis

For the RNA-seq experiment, HAP1-dEGFP^{Y66S} cells were transfected using iTOP, as described

above. The day before transfection, 600 000 cells per well had been plated in 6-well plates.

Treatments were: an empty transfection, a transfection with LbuCas13a but no guide RNA, a

transfection with LbuCas13a and a non-targeting guide RNA, and finally a transfection with

LbuCas13a and a dEGFP targeting guide RNA (SP2). All treatments were done in triplicates.

After 16 hours, cells were lysed in TRIzol reagent. Total RNA was isolated and DNase-treated

as described above. mRNA was then extracted using a NEBNext Poly(A) mRNA Magnetic

Isolation Module (NEB, E7490S) and RNA-seq libraries were prepared using a NEBNext Ultra

Directional RNA Library Prep Kit for Illumina (NEB, E7420L). RNA-seq libraries were sequenced

on an Illumina NextSeq500 Instrument with at least 10 million reads per library. Quality control

by FastQC (Andrews, 2010) and RSeQC (Wang et al., 2012) did not reveal any major issues.

An index was generated using the RefSeq GRCh37 assembly and reads were aligned and

quantified using STAR (Dobin et al., 2013) and featureCounts (Liao et al., 2014). Differentially

expressed genes were determined using DESeq2 (Love et al., 2014). Genes with an adjusted

P-value < 0.05 and log2(Fold Change) > |1| were regarded as differentially expressed genes,

and they were subjected to subsequent enrichment analysis. GO term and pathway enrichment

analysis was done using ShinyGO (Ge et al., 2020) V0.75, supplying all detected transcripts as

background. RNA-seq data is available online at the NCBI Gene expression omnibus

(GSE220759). The code of the differential gene expression analysis is available on

https://github.com/Geijsenlab/cas13.

Live cell imaging

The live cell imaging protocol used here was adapted from Wallberg *et al*. (Wallberg et al.,

2016). HAP1 and HAP1-dEGFP^{Y66S} cells were imaged in imaging medium (FluoroBrite™

DMEM, 10% FBS, 2.5 mM $CaCl_2$) supplemented with 2 µL CellTox™ Green (Promega, G8741)

and 5 µL Annexin V-AV647 (Invitrogen, A23204) per milliliter. Cells were seeded in 8-well µ-

Slides (ibidi, 80827) coated with a 1:100 dilution of Matrigel (Corning, 356231), plating 55 000

cells per well. The next day, iTOP was performed as described above, using an iTOP mixture

volume of 200 µL per well. After removing the iTOP buffer, wells were washed with 300 µL PBS

before adding 300 µL imaging medium. This extra PBS wash step prevents protein or salt

precipitation during imaging. After iTOP, cells were allowed to recover in the cell culture

incubator for up to four hours. Next, cells were moved to the Leica AF7000 and allowed to

equilibrate for one hour (37°C, 5% CO_2). During live imaging, pictures were taken every 15

minutes at four positions per well for 20 hours, using a 20x Dry N.A. 0.50 objective. Finally, the

amount of Annexin V and CellTox™ Green positive cells were counted with CellProfiler

(McQuin et al., 2018) using the IdentifyPrimaryObjects module.

Cell selection experiments

For the cell selection experiments, HAP1-EGFP cells were mixed with HAP1 cells at a ratio of

roughly 4:1. HAP1-EGFP were selected against by introducing LbuCas13a with an EGPF

targeting guide RNA (SP2) using iTOP. The percentage of EGFP positive cells was measured

by FACS on the BD FACSCanto II (BD Bioscience). To generate HAP1-EGFP lines with

different levels of EGFP expression, cells were sorted into fifteen different populations with

increasing EGFP intensities using the BD FACSJazz cell sorter (BD Bioscience). We sorted the

cells in complete cell culture medium. After sorting, the three populations with the lowest EGFP

intensity were cultured in medium without puromycin. The next three lowest populations were

cultured at 2 µg/mL puromycin, the following groups of three at 5 µg/mL, 10 µg/mL and finally 20

µg/mL. For the selection experiment, the different EGFP lines were mixed with wildtype HAP1

cells and subjected to selection as described above. Additionally, in parallel they were also

plated in a separate well. These parallel wells were harvested in TRIzol immediately after the

mixed population had been subjected to iTOP, and used to determine their EGFP expression

levels. Conventional RT-qPCR was performed as described above for 5'-3' RT-qPCR. qPCR

primers can be found in supplementary table 2.

To repair the dEGFP$^{\Delta fluor}$ mutation, cells were transfected with recombinant AsCas12a protein,

guide RNA (Supplementary table 3) and a LAHR repair template (Supplementary table 3) using

iTOP. For each well of a 96-well plate, 50 µL of iTOP mixture that contained 20 µL of

transfection supplement B, 10 µL of AsCas12a protein (150 µM), 10 µL of guide RNA (300 µM)

and 10 µL of 500 pmol repair template in nuclease-free water was used. Three days after iTOP

transfection, the cells that had not been edited were selected against by delivering LbuCas13a

and a dEGFP$^{\Delta fluor}$ targeting guide RNA (Supplementary table 1) using iTOP. Finally, the

percentage of EGFP positive cells was assessed on the BD FACSCanto II three days after

selection with LbuCas13a.

Data availability

The datasets and computer code produced in this study are available in the following

databases:

-    Nanopore sequencing data: NCBI Sequencing Read Archive PRJNA912090
(https://www.ncbi.nlm.nih.gov/sra/PRJNA912090)
-    RNA-seq data: Gene Expression Omnibus
(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE220759)
-    Scripts for nanopore sequencing coverage and RNA-seq analysis: GitHub
(https://github.com/Geijsenlab/cas13)

References

Abudayyeh OO, Gootenberg JS, Essletzbichler P, Han S, Joung J, Belanto JJ, Verdine V, Cox

DBT, Kellner MJ, Regev A, Lander ES, Voytas DF, Ting AY, Zhang F. 2017. RNA targeting

with CRISPR–Cas13. *Nature*. doi:10.1038/nature24049

Abudayyeh OO, Gootenberg JS, Konermann S, Joung J, Slaymaker IM, Cox DBT, Shmakov S,

Makarova KS, Semenova E, Minakhin L, Severinov K, Regev A, Lander ES, Koonin E V.,

Zhang F. 2016. C2c2 is a single-component programmable RNA-guided RNA-targeting

CRISPR effector. *Science (1979)* **353**. doi:10.1126/science.aaf5573

Ai Y, Liang D, Wilusz JE. 2022. CRISPR/Cas13 effectors have differing extents of off-target

effects that limit their utility in eukaryotic cells. *Nucleic Acids Res* **50**:1–16.

doi:10.1093/nar/gkac159

Aman R, Ali Z, Butt H, Mahas A, Aljedaani F, Khan MZ, Ding S, Mahfouz M. 2018. RNA virus

interference via CRISPR/Cas13a system in plants. *Genome Biol* **19**:1. doi:10.1186/s13059-

017-1381-1

Andrews S. 2010. FastQC: A Quality Control Tool for High Throughput Sequence Data [Online].

http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Arizti-sanz J, Freije CA, Stanton AC, Boehm CK, Petros BA, Siddiqui S, Shaw BM, Adams G,

Kosoko-Thoroddsen T-SF, Kemball ME, Gross R, Wronka L, Caviness K, Hensley LE,

Bergman NH, Macinnis BL, Lemieux JE, Sabeti PC, Myhrvold C. 2020. Integrated sample

inactivation, amplification, and Cas13-based detection of SARS-CoV-2 Jon. *bioRxiv*.

Auer H, Lyianarachchi S. 2003. Chipping away at the chip bias: RNA degradation in microarray

analysis. *Nat Genet* **35**:292–293.

Blanchard EL, Vanover D, Bawage SS, Tiwari PM, Rotolo L, Beyersdorf J, Peck HE, Bruno NC,

Hincapie R, Michel F, Murray J, Sadhwani H, Vanderheyden B, Finn MG, Brinton MA,

Lafontaine ER, Hogan RJ, Zurla C, Santangelo PJ. 2021. Treatment of influenza and

SARS-CoV-2 infections via mRNA-encoded Cas13a in rodents. *Nat Biotechnol* **39**:717–726. doi:10.1038/s41587-021-00822-w

Blighe K, Rana S, Lewis M. 2022. EnhancedVolcano: Publication-ready volcano plots with enhanced colouring and labeling.

Bot JF, van der Oost J, Geijsen N. 2022. The double life of CRISPR–Cas13. *Curr Opin Biotechnol* **78**:102789. doi:10.1016/j.copbio.2022.102789

Buchman AB, Brogan DJ, Sun R, Yang T, Hsu PD, Akbari OS. 2020. Programmable RNA Targeting Using CasRx in Flies. *CRISPR J* **3**:164–176. doi:10.1089/crispr.2020.0018

Consortium Rna. 2021. RNAcentral 2021: Secondary structure integration, improved sequence search and new member databases. *Nucleic Acids Res* **49**:D212–D220. doi:10.1093/nar/gkaa921

Cox DBT, Gootenberg JS, Abudayyeh OO, Franklin B, Kellner MJ, Joung J, Zhang F. 2017. RNA editing with CRISPR-Cas13. *Science (1979)* **358**:1019–1027. doi:10.1126/science.aaq0180

Crooks G, Hon G, Chandonia J, Brenner S. 2004. WebLogo: A Sequence Logo Generator. *Genome Res* **14**:1188–1190. doi:10.1101/gr.849004.1

D'Astolfo DS, Pagliero RJ, Pras A, Karthaus WR, Clevers H, Prasad V, Lebbink RJ, Rehmann H, Geijsen N. 2015. Efficient intracellular delivery of native proteins. *Cell* **161**:674–690. doi:10.1016/j.cell.2015.03.028

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras

TR. 2013. STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**:15–21.

doi:10.1093/bioinformatics/bts635

Dowle M, Srinivasan A. 2022. data.table: Extension of "data.frame."

East-Seletsky A, O'Connell MR, Burstein D, Knott GJ, Doudna JA, O'Connell MR, Burstein D,

Knott GJ, Doudna JA, O'Connell MR, Burstein D, Knott GJ, Doudna JA. 2017. RNA

Targeting by Functionally Orthogonal Type VI-A CRISPR-Cas Enzymes. *Mol Cell* **66**:373–

383. doi:10.1016/j.molcel.2017.04.008

East-Seletsky A, O'Connell MR, Knight SC, Burstein D, Cate JHD, Tjian R, Doudna JA. 2016.

Two distinct RNase activities of CRISPR-C2c2 enable guide-RNA processing and RNA

detection. *Nature* **538**:270–273. doi:10.1038/nature19802

Ewen-Campen B, Perrimon N. 2018. ovoD Co-selection: A method for enriching CRISPR/

Cas9-edited alleles in drosophila. *G3: Genes, Genomes, Genetics* **8**:2749–2756.

doi:10.1534/g3.118.200498

Fozouni P, Son S, Derby MD de L, Knott GJ, Gray CN, D'Ambrosio M V., Zhao C, Switz NA,

Kumar GR, Stephens SI, Boehm D, Tsoa C, Shu J, Bhuiya A, Armstrong M, Harris AR,

Chen P-Y, Osterloh JM, Meyer-Franke A, Joehnk B, Walcott K, Sil A, Langelier C, Pollard

KS, Crawford ED, Puschik AS, Phelps M, Kistler A, DeRisi JL, Doudna JA, Fletcher DA, Ott

M. 2021. Amplification-free detection of SARS-CoV-2 with CRISPR-Cas13a and mobile

phone microscopy. *Cell* **184**:323–333. doi:10.1016/j.cell.2020.12.001

Freije CA, Myhrvold C, Boehm CK, Lin AE, Welch NL, Carter A, Metsky HC, Luo CY,

Abudayyeh OO, Gootenberg JS, Yozwiak NL, Zhang F, Sabeti PC. 2019. Programmable

Inhibition and Detection of RNA Viruses Using Cas13. *Mol Cell* 1–12.

doi:10.1016/J.MOLCEL.2019.09.013

Ge SX, Jung D, Jung D, Yao R. 2020. ShinyGO: A graphical gene-set enrichment tool for

animals and plants. *Bioinformatics* **36**:2628–2629. doi:10.1093/bioinformatics/btz931

Gootenberg JS, Abudayyeh OO, Kellner MJ, Joung J, Collins JJ, Zhang F. 2018. Multiplexed

and portable nucleic acid detection platform with Cas13, Cas12a, and Csm6. *Science*

(1979) **444**. doi:10.1126/science.aaq0179

Gootenberg JS, Abudayyeh OO, Lee JW, Essletzbichler P, Dy AJ, Joung J, Verdine V, Donghia

N, Daringer NM, Freije CA, Myhrvold C, Bhattacharyya RP, Livny J, Regev A, Koonin E V.,

Hung DT, Sabeti PC, Collins JJ, Zhang F. 2017. Nucleic acid detection with CRISPR-

Cas13a/C2c2. *Science (1979)* **356**:438–442. doi:10.1126/science.aam9321

He B, Peng W, Huang J, Zhang H, Zhou Y, Yang X, Liu J, Li Z, Xu C, Xue M, Yang H, Huang P.

2020. Modulation of metabolic functions through Cas13d-mediated gene knockdown in

liver. *Protein Cell* **11**:518–524. doi:10.1007/s13238-020-00700-2

Huynh N, Depner N, Larson R, King-Jones K. 2020. A versatile toolkit for CRISPR-Cas13-based

RNA manipulation in Drosophila. *Genome Biol* **21**:1–29. doi:10.1186/s13059-020-02193-y

Jing X, Xie B, Chen L, Zhang N, Jiang Y, Qin H, Wang H, Hao P, Yang S, Li X. 2018.

Implementation of the CRISPR-Cas13a system in fission yeast and its repurposing for

precise RNA editing. *Nucleic Acids Res* **46**:e90. doi:10.1093/nar/gky433

Johnson TI, Minteer CJ, Kottmann D, Dunlop CR, Fernández SB de Q, Carnevalli LS, Wallez Y,

Lau A, Richards FM, Jodrell DI. 2021. Quantifying cell cycle-dependent drug sensitivities in

cancer using a high throughput synchronisation and screening approach. *EBioMedicine* **68**.

doi:10.1016/j.ebiom.2021.103396

Knott GJ, East-Seletsky A, Cofsky JC, Holton JM, Charles E, O'Connell MR, Doudna JA. 2017.

Guide-bound structures of an RNA-targeting A-cleaving CRISPR–Cas13a enzyme. *Nat*

Struct Mol Biol. doi:10.1038/nsmb.3466

Konermann S, Lotfy P, Brideau NJ, Oki J, Maxim N. 2018. Transcriptome engineering with

RNA-targeting Type VI-D CRISPR effectors. *Cell* 1–12. doi:10.1016/j.cell.2018.02.033

Koonin E V., Makarova KS, Zhang F. 2017. Diversity, classification and evolution of CRISPR-

Cas systems. *Curr Opin Microbiol* **37**:67–78.

Kushawah G, Hernandez-Huertas L, Abugattas-Nuñez del Prado J, Martinez-Morales JR,

DeVore ML, Hassan H, Moreno-Sanchez I, Tomas-Gallardo L, Diaz-Moscoso A, Monges

DE, Guelfo JR, Theune WC, Brannan EO, Wang W, Corbin TJ, Moran AM, Sánchez

Alvarado A, Málaga-Trillo E, Takacs CM, Bazzini AA, Moreno-Mateos MA. 2020. CRISPR-

Cas13d Induces Efficient mRNA Knockdown in Animal Embryos. *Dev Cell* **54**:805-817.e7.

doi:10.1016/j.devcel.2020.07.013

Li H. 2018. Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics* **34**:3094–

3100. doi:10.1093/bioinformatics/bty191

Li S, Li X, Xue W, Zhang L, Yang LZ, Cao SM, Lei YN, Liu CX, Guo SK, Shan L, Wu M, Tao X,

Zhang JL, Gao X, Zhang J, Wei J, Li J, Yang L, Chen LL. 2021. Screening for functional

circular RNAs using the CRISPR–Cas13 system. *Nat Methods* **18**:51–59.

doi:10.1038/s41592-020-01011-4

Li Y, Xu J, Guo X, Li Z, Cao L, Liu S, Guo Y, Wang G, Luo Y, Zhang Z, Wei X, Zhao Y, Liu T,

Wang X, Xia H, Kuang M, Guo Q, Li J, Chen L, Wang Y, Li Q, Wang F, Liu Q, You F. 2023.

The collateral activity of RfxCas13d can induce lethality in a RfxCas13d knock-in mouse

model. *Genome Biol* **24**:1–25. doi:10.1186/s13059-023-02860-w

Li Z, Li Zihan, Cheng X, Wang S, Wang X, Ma S, Lu Z, Zhang H, Zhao W, Chen Z, Yao Y,

Zhang C, Chao L, Li W, Fei T. 2023. Intrinsic targeting of host RNA by Cas13 constrains its

utility. *Nat Biomed Eng* **8**:177–192. doi:10.1038/s41551-023-01109-y

Liao Y, Smyth GK, Shi W. 2014. FeatureCounts: An efficient general purpose program for

assigning sequence reads to genomic features. *Bioinformatics* **30**:923–930.

doi:10.1093/bioinformatics/btt656

Liu G, Lin Q, Jin S, Gao C. 2022. The CRISPR-Cas toolbox and gene editing technologies. *Mol*

Cell **82**:333–347. doi:10.1016/j.molcel.2021.12.002

Liu L, Li X, Ma J, Li Z, You L, Wang J, Wang M, Zhang X, Wang Y. 2017a. The Molecular

Architecture for RNA-Guided RNA Cleavage by Cas13a. *Cell* **170**:714-726.e10.

doi:10.1016/j.cell.2017.06.050

Liu L, Li X, Wang J, Wang M, Chen P, Yin M, Li J, Sheng G, Wang Y. 2017b. Two Distant

Catalytic Sites Are Responsible for C2c2 RNase Activities. *Cell* **168**:121-134.e12.

doi:10.1016/j.cell.2016.12.031

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol* **15**. doi:10.1186/s13059-014-0550-8

Mahas A, Aman R, Mahfouz M. 2019. CRISPR-Cas13d mediates robust RNA virus interference

in plants. *Genome Biol* 1–16.

Makarova KS, Wolf YI, Alkhnbashi OS, Costa F, Shah SA, Saunders SJ, Barrangou R, Brouns

SJJ, Charpentier E, Haft DH, Horvath P, Moineau S, Mojica FJM, Terns RM, Terns MP,

White MF, Yakunin AF, Garrett RA, van der Oost J, Backofen R, Koonin E V. 2015. An

updated evolutionary classification of CRISPR–Cas systems. *Nat Rev Microbiol* **13**:722–

736. doi:10.1038/nrmicro3569

Makarova KS, Wolf YI, Iranzo J, Shmakov SA, Alkhnbashi OS, Brouns SJJ, Charpentier E,

Cheng D, Haft DH, Horvath P, Moineau S, Mojica FJM, Scott D, Shah SA, Siksnys V,

Terns MP, Venclovas Č, White MF, Yakunin AF, Yan W, Zhang F, Garrett RA, Backofen R,

van der Oost J, Barrangou R, Koonin E V. 2020. Evolutionary classification of CRISPR–

Cas systems: a burst of class 2 and derived variants. *Nat Rev Microbiol* **18**:67–83.

doi:10.1038/s41579-019-0299-x

Marraffini LA. 2015. CRISPR-Cas immunity in prokaryotes. *Nature* **526**.

doi:10.1038/nature15386

McQuin C, Goodman A, Chernyshev V, Kamentsky L, Cimini BA, Karhohs KW, Doan M, Ding L,

Rafelski SM, Thirstrup D, Wiegraebe W, Singh S, Becker T, Caicedo JC, Carpenter AE.

2018. CellProfiler 3.0: Next-generation image processing for biology. *PLoS Biol* **16**:1–17.

doi:10.1371/journal.pbio.2005970

Meeske AJ, Nakandakari-Higa S, Marraffini LA. 2019. Cas13-induced cellular dormancy

prevents the rise of CRISPR-resistant bacteriophage. *Nature*. doi:10.1038/s41586-019-

1257-5

Mohanraju P, Makarova KS, Zetsche B, Zhang F, Koonin E V., Oost J van der. 2016. Diverse

evolutionary roots and mechanistic variations of the CRISPR-Cas systems. *Science (1979)*

353. doi:10.1126/science.aad5147

Mohanraju P, Saha C, van Baarlen P, Louwen R, Staals RHJ, van der Oost J. 2022. Alternative

functions of CRISPR–Cas systems in the evolutionary arms race. *Nat Rev Microbiol*

0123456789. doi:10.1038/s41579-021-00663-z

Morgan M, Anders S, Lawrence M, Aboyoun P, Pagès H, Gentleman R. 2009. ShortRead: A

bioconductor package for input, quality assessment and exploration of high-throughput

sequence data. *Bioinformatics* **25**:2607–2608. doi:10.1093/bioinformatics/btp450

Myhrvold C, Freije CA, Gootenberg JS, Abudayyeh OO, Metsky HC, Kellner MJ, Barnes KG,

Chak B, Yozwiak NL, MacInnis BL, Zhang F, Sabeti PC, Durbin AF, Gehrke L, Bosch I,

Tan AL, Paul LM, Isern S, Michael SF, Parham LA, Garcia KF, Lorenzana I, Mondini A,

Nogueira ML. 2018. Field-deployable viral diagnostics using CRISPR-Cas13. *Science*

(1979) **448**:444–448. doi:10.1126/science.aas8836

Nolan T, Hands RE, Bustin SA. 2006. Quantification of mRNA using real-time RT-PCR. *Nat*

Protoc **1**:1559–1582. doi:10.1038/nprot.2006.236

Oost J Van Der, Westra ER, Jackson RN, Wiedenheft B. 2014. Unravelling the structural and

mechanistic basis of CRISPR – Cas systems. *Nat Rev Microbiol* **12**:479–492.

doi:10.1038/nrmicro3279

Özcan A, Krajeski R, Ioannidi E, Lee B, Gardner A, Makarova KS, Koonin E V, Abudayyeh OO,

Gootenberg JS. 2021. Programmable RNA targeting with the single-protein CRISPR

effector Cas7-11. *Nature* **597**. doi:10.1038/s41586-021-03886-5

Patchsung M, Jantarug K, Pattama A, Aphicho K, Suraritdechachai S, Meesawat P, Sappakhaw

K, Leelahakorn N, Ruenkam T, Wongsatit T, Athipanyasilp N, Eiamthong B, Lakkanasirorat

B, Phoodokmai T, Niljianskul N, Pakotiprapha D, Chanarat S, Homchan A, Tinikul R,

Kamutira P, Phiwkaow K, Soithongcharoen S, Kantiwiriyawanitch C, Pongsupasa V,

Trisrivirat D, Jaroensuk J, Wongnate T, Maenpuen S, Chaiyen P, Kamnerdnakta S,

Swangsri J, Chuthapisith S, Sirivatanauksorn Y, Chaimayo C, Sutthent R, Kantakamalakul

W, Joung J, Ladha A, Jin X, Gootenberg JS, Abudayyeh OO, Zhang F, Horthongkham N,

Uttamapinant C, Nucleic. 2020. Clinical validation of a Cas13-based assay for the detection

of SARS-CoV-2 RNA. *Nat Biomed Eng* **4**. doi:10.1038/s41551-020-00603-x

Powell JE, Lim CKW, Krishnan R, Mccallister TX, Saporito-magriña C, Zeballos MA, Mcpheron

GD, Gaj T. 2022. Targeted gene silencing in the nervous system with CRISPR-Cas13. *Sci*

Adv **8**.

Sharma G, Sharma AR, Bhattacharya M, Lee SS, Chakraborty C. 2021. CRISPR-Cas9: A

Preclinical and Clinical Perspective for the Treatment of Human Diseases. *Molecular*

Therapy **29**:571–586. doi:10.1016/j.ymthe.2020.09.028

Shi P, Murphy MR, Aparicio AO, Kesner JS, Fang Z, Chen Z, Trehan A, Guo Y, Wu X. 2023.

Collateral activity of the CRISPR/RfxCas13d system in human cells. *Commun Biol* **6**.

doi:10.1038/s42003-023-04708-2

Shi P, Murphy MR, Aparicio AO, Kesner JS, Fang Z, Chen Z, Trehan A, Wu X. 2021. RNA-

guided cell targeting with CRISPR/RfxCas13d collateral activity in human cells. *bioRxiv*.

Shy BR, Macdougall MS, Clarke R, Merrill BJ. 2016. Co-incident insertion enables high

efficiency genome engineering in mouse embryonic stem cells. *Nucleic Acids Res*

44:7997–8010. doi:10.1093/nar/gkw685

Signorell A et mult. al. 2021. DescTools: Tools for descriptive statistics.

Slaymaker IM, Mesa P, Kellner MJ, Kannan S, Brignole E, Koob J, Feliciano PR, Stella S,

Abudayyeh OO, Gootenberg JS, Strecker J, Montoya G, Zhang F. 2019. High-Resolution

Structure of Cas13b and Biochemical Characterization of RNA Targeting and Cleavage.

Cell Rep **26**:3741-3751.e5. doi:10.1016/j.celrep.2019.02.094

Smargon AA, Cox DBT, Pyzocha NK, Zheng K, Slaymaker IM, Gootenberg JS, Abudayyeh OA,

Essletzbichler P, Shmakov S, Makarova KS, Koonin E V., Zhang F. 2017. Cas13b Is a

Type VI-B CRISPR-Associated RNA-Guided RNase Differentially Regulated by Accessory

Proteins Csx27 and Csx28. *Mol Cell* **65**:618–630. doi:10.1016/j.molcel.2016.12.023

Swift GH, Peyton MJ, MacDonald RJ. 2000. Assessment of RNA quality by semi-quantitative

RT-PCR of multiple regions of a long ubiquitous mRNA. *Biotechniques* **28**:524–531.

doi:10.2144/00283rr01

Tieu V, Sotillo E, Bjelajac JR, Chen C, Malipatlolla M, Guerrero JA, Xu P, Quinn PJ, Fisher C,

Klysz D, Mackall CL, Qi L s. 2024. A versatile CRISPR-Cas13d platform for multiplexed

transcriptomic regulation and metabolic engineering in primary human T cells. *Cell*

187:1278–1295. doi:https://doi.org/10.1016/j.cell.2024.01.035

Tng PYL, Carabajal Paladino L, Verkuijl SAN, Purcell J, Merits A, Leftwich PT, Fragkoudis R,

Noad R, Alphey L. 2020. Cas13b-dependent and Cas13b-independent RNA knockdown of

viral sequences in mosquito cells following guide RNA expression. *Commun Biol* **3**.

doi:10.1038/s42003-020-01142-6

Tong H, Huang J, Xiao Q, He B, Dong X, Liu Y, Yang X, Han D, Wang Z, Wang X, Ying W,

Zhang R, Wei Y, Xu C, Zhou Y, Li Y, Cai M, Wang Q, Xue M, Li G, Fang K, Zhang H, Yang

H. 2022. High-fidelity Cas13 variants for targeted RNA degradation with minimal collateral

effects. *Nat Biotechnol*. doi:10.1038/s41587-022-01419-7

Tong H, Huang J, Xiao Q, He B, Dong X, Liu Y, Yang X, Wang Z, Yin W, Zhang R, Wei Y,

Wang X, Xu C, Zhou Y, Li Y, Cai M, Wang Q, Xue M, Li G, Fang K, Zhang H, Yang H.

2021. High-fidelity Cas13 variants for targeted RNA degradation with minimal collateral

effect. *bioRxiv* 2021.12.18.473271.

Wallberg F, Tenev T, Meier P. 2016. Time-lapse imaging of cell death. *Cold Spring Harb Protoc*

2016:251–258. doi:10.1101/pdb.prot087395

Wang L, Wang S, Li W. 2012. RSeQC: Quality control of RNA-seq experiments. *Bioinformatics*

28:2184–2185. doi:10.1093/bioinformatics/bts356

Wang Q, Liu X, Zhou J, Yang C, Wang G, Tan Y, Wu Y, Zhang S, Yi K, Kang C. 2019. The

CRISPR-Cas13a Gene-Editing System Induces Collateral Cleavage of RNA in Glioma

Cells. *Advanced Science* **1901299**. doi:10.1002/advs.201901299

Wang Q, Liu Y, Han C, Yang M, Huang F, Duan X, Wang S, Yu Y, Liu J, Yang H, Lu D, Zhao H,

Zhang Y, Qin Q. 2021. Efficient RNA Virus Targeting via CRISPR/CasRx in Fish. *J Virol*

95.

Wessels HH, Méndez-Mancilla A, Guo X, Legut M, Daniloski Z, Sanjana NE. 2020. Massively

parallel Cas13 screens reveal principles for guide RNA design. *Nat Biotechnol* **38**:722–727.

doi:10.1038/s41587-020-0456-9

Wessels HH, Stirn A, Méndez-Mancilla A, Kim EJ, Hart SK, Knowles DA, Sanjana NE. 2024.

Prediction of on-target and off-target activity of CRISPR–Cas13d guide RNAs using deep

learning. *Nat Biotechnol* **42**:628–637. doi:10.1038/s41587-023-01830-8

Wickham H. 2010. A Layered grammar of graphics. *Journal of Computational and Graphical*

Statistics **19**:3–28. doi:10.1198/jcgs.2009.07098

Wickham H. 2009. ggplot2 Elegant graphics for data analysis, Springer. doi:10.1007/978-0-387-

98141-3

Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G, Hayes A,

Henry L, Hester J, Kuhn M, Pedersen T, Miller E, Bache S, Müller K, Ooms J, Robinson D,

Seidel D, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H. 2019. Welcome to

the Tidyverse. *J Open Source Softw* **4**:1686. doi:10.21105/joss.01686

Xu C, Zhou Y, Xiao Q, He B, Geng G, Wang Z, Cao B, Dong X, Bai W, Wang Y, Wang X, Zhou

D, Yuan T, Huo X, Lai J, Yang H. 2021. Programmable RNA editing with compact

CRISPR–Cas13 systems from uncultivated microbes. *Nat Methods* **18**:499–506.

doi:10.1038/s41592-021-01124-4

Yan WX, Chong S, Zhang H, Makarova KS, Koonin E V, Cheng DR, Scott DA. 2018. Cas13d Is

a Compact RNA-Targeting Type VI CRISPR Effector Positively Modulated by a WYL-

Domain- Containing Accessory Protein. *Mol Cell* 1–13. doi:10.1016/j.molcel.2018.02.028

Zhang C, Konermann S, Brideau NJ, Griffin PR, Hsu PD, Lyumkis D, Zhang C, Konermann S,

Brideau NJ, Lotfy P, Wu X, Novick SJ. 2018. Structural Basis for the RNA-Guided

Ribonuclease Activity of CRISPR-Cas13d Article Structural Basis for the RNA-Guided

Ribonuclease Activity of CRISPR-Cas13d. *Cell* **175**:212-223.e17.

doi:10.1016/j.cell.2018.09.001

Zhang Y, Nguyen TM, Zhang XO, Wang L, Phan T, Clohessy JG, Pandolfi PP. 2021. Optimized

RNA-targeting CRISPR/Cas13d technology outperforms shRNA in identifying functional

circRNAs. *Genome Biol* **22**:1–22. doi:10.1186/s13059-021-02263-9

Zhang Z, Wang Q, Liu Q, Zheng Y, Zheng C, Yi K. 2019. Dual-Locking Nanoparticles Disrupt

the PD-1 / PD-L1 Pathway for Efficient Cancer Immunotherapy. *Advanced Materials*

1905751:1–10. doi:10.1002/adma.201905751

Zhao Z, Shang P, Sage F, Geijsen N. 2022. Ligation-assisted homologous recombination

enables precise genome editing by deploying both MMEJ and HDR. *Nucleic Acids Res*

50:e62–e62. doi:10.1093/nar/gkac118

Zhou H, Su J, Hu X, Zhou C, Li H, Chen Z, Xiao Q, Wang B, Wu W, Sun Y, Zhou Y, Tang C, Liu

F, Wang L, Feng C, Liu M, Li S, Zhang Y, Xu H, Yao H, Shi L, Yang H. 2020. Glia-to-

Neuron Conversion by CRISPR-CasRx Alleviates Symptoms of Neurological Disease in

Mice. *Cell* **181**:590–603. doi:10.1016/j.cell.2020.03.024

Zuker M. 2003. Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic*

Acids Res **31**:3406–3415. doi:10.1093/nar/gkg595

1013

1014

Acknowledgements

The authors thank Stefan van der Elst of the Hubrecht Institute FACS facility, the LUMC FACS
facility and the Utrecht sequencing facility (USEQ) for technical help. The novo Nordisk
Foundation Center for Stem Cell Medicine (reNEW) is supported by a Novo Nordisk Foundation
grant number NFF21CC0073729. This work was also supported by funding from NWO-TTW
and the Princes Beatrix Muscle Foundation. Z.Z. was supported, in part, by the China
Scholarship Council.

1022

Author contributions

1024 JFB experimental design, execution and help writing the manuscript, ZZ experimental design,
1025 execution and help writing the manuscript, ML technical assistance, DK technical assistance,
1026 PS experimental design and help writing the manuscript, NG conceptual and experimental
1027 design, project supervision and help writing the manuscript.

1028

Conflict of Interest

1030 NG is co-founder of NTrans Technologies and Divvly. No conflicting interest.

Figure 1. LbuCas13a exhibits strong collateral cleavage activity *in vitro* and in human cells. A Total RNA profiles at various timepoints after transfection of the LbuCas13a RNP (n=3). Both HAP1 and HAP1-dEGFP cells were transfected with LbuCas13a and a dEGFP targeting guide RNA. The bottom row is zoomed in to the region indicated by the dotted box in the 0 minute panel. Fluorescence values are corrected for differences in amount of loaded RNA. **B** Schematic depiction of the 5'-3' RT-qPCR assay. Green arrows indicate the 5' and 3' primer pairs used for qPCR. Red cross indicates no amplification is possible when there is a break between the poly-A tail and the 5' primer pair. RNA is depicted in black, cDNA in brown. **C** 5'-3' RT-qPCR assay (n=3) on samples from A. Line shows ggplot2's default LOESS regression (Wickham, 2009). Since wild type HAP1 control cells do not contain dEGFP, no 5'-3' RT-qPCR assay on dEGFP was performed for these cells.

J. F. Bot et al. Figure 2

Figure 2. LbuCas13a displays collateral cleavage in different cell lines. A Total RNA profiles after targeting the endogenous transcripts 18S rRNA, GAPDH and RPS19 respectively. Total RNA isolated 100 minutes after RNP transfection. Fluorescence values from the Bioanalyzer were normalized to the total area under the curve of each sample. n=3, except for the U2OS 18S rRNA, U2OS GAPDH and ARPE19 non-targeting samples, where n=2. **B** Quantification of total RNA cleavage in different cell lines, by taking the area under the curve of the region outlined by the grey boxes in A. P-values are calculated by unpaired Student's t-test. **C** 5'-3' RT-qPCR assays on samples from A (n=3). Dot and line show mean and standard error of the mean. P-values calculated by comparing GAPDH 5' with GAPDH 3' using an unpaired Student's t-test (ns = not significant, * = p-value < 0.05).

Figure 3. Cells enter apoptosis in response to collateral RNA cleavage. A Loss of cell viability when treated with LbuCas13a RNP targeting dEGFP RNA. Images taken 24 hours after transfection. Scale bar is 50 µm. **B** Target RNA specific induction of apoptosis in HAP1 cells by LbuCas13a. Total amount of Annexin V (early apoptosis marker) and CellTox Green (dead cell marker) positive cells during live imaging after transfection. CTRL is an empty transfection. SP1, SP2 and SP3 are transfections with LbuCas13a and three different guide RNAs all targeting dEGFP. Quantified using CellProfiler (McQuin et al., 2018).

J. F. Bot et al. Figure 4

Figure 4. LbuCas13a as a cell selection tool. A Schematic of the Cas13-based cell selection. **B** Depletion of EGFP positive HAP1 cells after targeting the EGFP transcript, as measured by FACS (n=3). Trendlines show a generalized additive model using formula $y \sim s(x, k = 6)$. **C** Multiple rounds of targeting improve depletion of EGFP positive HAP1 cells, as measured by FACS (n=3). Error bars show standard deviation. **D** Schematic showing creation of cell lines with different EGFP expression levels and subsequent selection experiments. **E** Target RNA expression level determines cell selection efficiency (n=3). Target RNA expression level was determined using RT-qPCR, and the percentage of GFP positive cells was determined using FACS. Trendlines created by ggplot2's default LOESS regression. **F** Enrichment of edited cells by targeting unedited cells with LbuCas13a. Error bars show standard deviation. P-values calculated by unpaired Welch's t-test (ns = not significant, **** = P-value < 0.0001).

Figure 5. Innate immune response genes are upregulated in response to collateral RNA cleavage.
A Many genes are upregulated in response to collateral RNA cleavage. RNA-Seq EnhancedVolcano (Blighe et al., 2022) plot of dEGFP targeting samples vs controls, n=3. Fold change and P-values were calculated using DESeq2, see methods. Dotted lines indicate adjusted P-value = 0.05 and |Log2 FC| = 1. **B** A ranked list of enriched GO biological processes using the significantly upregulated genes (adjusted P-value < 0.05 & Log2 FC > 1). Plot generated using ShinyGo (Ge et al., 2020).

Figure 6. Nanopore sequencing after collateral cleavage by LbuCas13a. A Density plot of read length versus reference transcript length of all reads mapped to protein coding transcripts, excluding histone genes and WDR74. Columns show minutes after transfection. Data from replicate 2, see Supplementary figure 5A for replicate 1. **B** Transcript coverage or the target transcript dEGFP. Reads were aligned to the entire plasmid sequence used to make this cell line (Supplementary file 1, pEF-1a_dEGFP-Y66S-IRIS-PuroR-WPRE). The start and stop codon are at position 3415 and 5316 respectively. The dotted lines indicate the protospacer position. Columns show minutes after transfection. The dEGFP targeting sample at 0 minutes after transfection of replicate 1 was excluded due to failed library preparation. **C** Transcript coverage of RPS19-202, a collateral RNA. RPS19-202 appears to have one major cleavage site.

Figure 7. LbuCas13a collateral RNA cleavage sites in human cells. A Transcript coverage of non-target RNAs, zoomed in at putative cleavage sites. **B** The predicted RNA structures at the putative cleavage sites from A. Red letters are the approximate location of the coverage dips, corresponding to the dashed lines in A. RNA structures for RPS19-202 and GAPDH-205 are the top predictions of Mfold (Zuker, 2003) using the default setting. 18S rRNA structure provided by RNAcentral.org (Consortium, 2021) (URS0000726FAB_9606). **C** Sequence logo around 100 manually annotated cleavage sites. The U at position 0 was enforced during potential cleavage site selection. Logo made using WebLogo (Crooks et al., 2004).

A

Figure 8. A timeline of events after RNA targeting with LbuCas13a in cells. A This plot provides an summary of the timing of events after RNA targeting with LbuCas13a. The target RNA cleavage timing takes into account both figures 1C and 6B. While figure 1C suggests target cleavage peaks at 0 minutes after delivery, figure 6B suggests it peaks at 50-200 minutes after delivery. Collateral RNA cleavage timing is based on figures 1C, 6A and 6C. For ribosomal RNA degradation see figure 1A. The early apoptosis timeline is based on figure 3B. The cell death timeline is based on figure 3B and 4B.