A systematic mapping of the genomic and proteomic variation associated with monogenic diabetes

Ksenia Kuznetsova[1,$], Jakub Vašíček[1], Dafni Skiadopoulou[1], Janne Molnes[1,2], Miriam Udler[3], Stefan Johansson[1,2], Pål Rasmus Njolstad[1,4], Alisa Manning[3,†], Marc Vaudel[1,5,6,†,$]

1. Mohn Center for Diabetes Precision Medicine, Department of Clinical Science, University of Bergen, Bergen, Norway
2. Department of Medical Genetics, Haukeland University Hospital, Bergen, Norway
3. Harvard Medical School, Mass General Hospital, Broad Institute, Cambridge and Boston, MA, USA
4. Children and Youth Clinic, Haukeland University Hospital, Bergen, Norway
5. Department of Genetics and Bioinformatics, Norwegian Institute of Public Health, Oslo, Norway
6. Computational Biology Unit (CBU), Department of Informatics, University of Bergen

[†] These authors jointly supervised the work
[$] To whom correspondence should be addressed

Summary

Monogenic diabetes is characterized as a group of diseases caused by rare variants in single genes. Multiple genes have been described to be responsible for monogenic diabetes, but the information on the variants is not unified among different resources. We have developed an automated pipeline for collecting and harmonizing data on genetic variants associated with monogenic diabetes. Furthermore, we have translated variant genetic sequences into protein sequences accounting for all protein isoforms and their variants. This allows researchers to consolidate information on variant genes and proteins associated with monogenic diabetes and facilitates their study using proteomics or structural biology. Our open and flexible implementation using Jupyter notebooks enables tailoring and modifying the pipeline and its application to other rare diseases.

Introduction

The most common forms of monogenic diabetes are maturity-onset diabetes of the young (MODY), neonatal diabetes (1), inherited lipodystrophies, mitochondrial diabetes, among others

(2). Today, international guidelines are available for the diagnostic and follow-up of patients with suspected MODY (3). These patients may now receive a molecular genetic diagnosis using diagnostic gene sequencing panels (panelapp.genomicsengland.co.uk/panels/472). This allows precise MODY subtyping and, depending on the diagnosis, the opportunity to avoid lifelong insulin medication and complications through lifestyle management or alternative treatment using oral antidiabetic drugs (4). Furthermore, because incorrect treatment due to misdiagnosis may change the future response to the correct medication, the timeline of diagnosis and care is thus crucial (4). It is estimated that around 77 % of monogenic diabetes cases remain undiagnosed (5). These patients and their relatives remain unaware of their familial condition and do not benefit from adapted care.

The first challenge in establishing a firm diagnosis in all cases of MODY is the mapping of all genes that can cause monogenic diabetes. To date, multiple genes have been discovered to have associations with familial forms of diabetes (2). Fourteen of them are notably often referred to as "the MODY genes" (3), although this list is subject to debate in the literature (6) and is being systematically assessed by international experts using established guidelines to determine gene-disease relationships (https://clinicalgenome.org/affiliation/40016/). A second challenge is the difficulty to evaluate the pathogenicity of genetic variants (7), for which there is also an on-going international effort to establish guidelines and provide expert variant curations in the ClinVar database (clinicalgenome.org/affiliation/50016) (8). Furthermore, the response to alternative treatment might differ between populations (9). One of the ways to address the challenge of precise diagnostics is to complement genetic screening with additional data, combining both molecular and clinical dimensions (10).

The recent advent of high-performance computational models for protein structures notably holds the promise to democratize the study of gene sequences (11). We aim to study the properties of the proteins encoded by genes carrying alleles suspected to cause monogenic diabetes. This can, for example, help us understand whether the structure and properties of the protein are affected by these alleles, hence shedding light on the pathogenicity of the variant investigated (12).

The adoption of these approaches is impaired by the difficulty of mapping variants associated with monogenic diabetes to the different forms of proteins that they encode. First, given that the variants associated with monogenic diabetes are rare, the coverage by genomic databases is low. Maintaining an updated list of variants requires constant monitoring of the literature by experts. Second, variants reported in the literature often lack standardization in their identifiers and coordinates, making it challenging to map them to a given genome build and requiring manual variant mapping. Third, inferring the consequences on protein sequences is still a daunting task for some variants (those alleles affecting splice sites and untranslated regions [UTRs], for example). Fourth, a given protein-coding sequence might encode different protein isoforms, which will produce different forms of proteins upon folding and post-translational modification (13), hence for a given variant multiple protein sequences need to be investigated. Mapping genetic variants associated with monogenic diabetes from genes to proteins is therefore not tractable and sustainable without automation using dedicated bioinformatic tools.

Here, we describe a new open-source modular pipeline based on Jupyter notebooks (eprints.soton.ac.uk/403913) that allows for the systematic collection of variants associated with monogenic diabetes and their mapping to Ensembl (14) and ClinVar (8). We demonstrate how the different genes associated with monogenic diabetes harbor variants of different levels of pathogenicity. Finally, we port the variant sequences to the protein level and provide the resulting sequences in a standard format that can readily be used for proteomic and structural proteomic analyses using mass spectrometry or protein structure modeling.

Methods

General architecture

The pipeline consists of seven independent modules written in Python using Jupyter notebooks (figure 1). The notebooks can be chained together as a pipeline, used as standalone applications, or integrated into other pipelines. First, the pipeline takes a list of genes and extracts exonic variants from Ensembl retaining only those variants that are predicted to affect protein sequences. Next, the program integrates the variants from ClinVar and maps them to

Ensembl. Similarly, variants are extracted from the literature, here using the literature mining by Rafique et al. (15), and mapped to Ensembl. Subsequently, the harmonized collection of variants is consolidated in a database stored in the form of a text file that can easily be parsed and reused. Finally, the table of variants is mapped to all the transcripts linked by Ensembl to the genes of interest to obtain protein sequences of all the possible isoforms encoded by these genes. In this last step, the DNA sequences are translated to amino acid sequences and stored as protein FASTA files. To visually inspect the results, a separate module allows overlaying all the variants that can possibly affect a given gene onto the corresponding amino acid sequences.

Figure 1. **General architecture of the pipeline.** *Module 1*: extract variants from Ensembl affecting genes associated with monogenic diabetes. *Module 2*: filter variant by consequence on the protein. *Module 3*: extract variants from ClinVar affecting genes associated with monogenic diabetes. *Modules 4* and *Module 5*: extract the variants from the literature using the mining by Rafique et al. *Module 6*: consolidate the variants in a single table. *Translation*: produce the possible variant protein sequences. MODY - maturity-onset diabetes of the young, MD - monogenic diabetes, ND - neonatal diabetes, API - application programming interface, PMID - identifiers of scientific publications from the PubMed database, "rs..." identifiers - identifiers of the genomic variants from the dbSNP database, that start from "rs".

Module 1 - Mining variants in Ensembl

In *Module 1*, a list of genes is mapped to Ensembl genes, transcripts, and exon identifiers using the Ensembl REST API (16). Subsequently, the Ensembl REST API is queried using the exon identifiers to return all variants in Ensembl overlapping with the corresponding regions along with their annotation (identifiers, coordinates, consequences, etc.). For multi-allelic variants, we treat every alternative allele as a different entry, and add it as a new line to the table produced.

Module 2 - Categorizing by consequence and pathogenicity

Module 2 takes as an input the table of exonic variants from Ensembl generated in *Module 1*. For each consequence type, the prevalence of variants with different levels of pathogenicity is computed and visualized as heat maps (figure 2A). The same is done for the data from ClinVar with all the variants regardless of the associated phenotype (figure 2B).

figure 2. Distribution of the consequence types within pathogenicity categories of the variants from Ensembl (A) and ClinVar (B). The heatmap shows the percentage of variants within pathogenicity categories for each consequence types. Arrows show the consequence types that were left in the Ensembl reference table created in *Module 2* after filtering. "Stop lost" is marked with a blue arrow as it has conflicting evidence in two sources (see detailed in the text).

By default, only variants whose consequence feature mostly *likely pathogenic* and *pathogenic* are retained for further analyses: 'missense_variant', 'protein_altering_variant', 'coding_sequence_variant', 'frameshift_variant', 'splice_donor_variant',

'splice_acceptor_variant', 'splice_donor_5th_base_variant', 'start_lost', 'stop_gained', 'stop_lost', 'inframe_deletion', 'inframe_insertion'.

Module 3. Mapping of the variants from ClinVar

Module 3 takes as an input three tables exported from ClinVar containing all the variants associated with three phenotypes: "MODY", "monogenic diabetes", and "neonatal diabetes". ClinVar annotates variants with different levels of pathogenicity: "pathogenic", "likely pathogenic", "uncertain significance", "likely benign" or "benign". The variants presenting dbSNP identifiers are passed to *Module 2*. The others are mapped to Ensembl using genomic coordinates and alleles, and predicted consequences are obtained with the Ensembl Variant Effect Predictor (VEP)(17) called on all variants using the Ensembl REST API. The variants with predicted consequences according to Ensembl are filtered in the same way as in *Module 2*. The variants where no consequence type was returned were all either short deletions, short insertions, or other short fragment replacements. This subset is directly passed to *Module 6*.

Module 4. Mapping variants from the literature (step 1 of 2)

Module 4 takes variants associated with monogenic diabetes according to the literature to cover the variants where the association with monogenic diabetes is not yet consolidated in ClinVar or Ensembl. As input we used the variants mined in the review by Rafique et al. (15) and provided as a supplementary table. The module uses the Vcfanno (18) library to annotate the variants with DNA coordinates and the dbSNP identifiers where possible. The Ensembl REST API is queried as in *Module 1*, and the results are passed to *Module 6*.

Module 5. Mapping variants from the literature (step 2 of 2)

For the variants that could not be mapped automatically in *Module 4*, the title of the publication as obtained from *Module 4* is queried against the Entrez Programming Utilities API (https://www.ncbi.nlm.nih.gov/books/NBK25500/) to return the PubMed (pubmed.ncbi.nlm.nih.gov) identifiers (PMIDs) of these articles. Note that some of the PMIDs were mapped and had to be added manually. Next, these PMIDs are used to query the same API and return the "rs" identifiers of the variants mentioned in these publications. Finally, the

variants are mapped to Ensembl using their identifiers and passed to *Module 6*. Note that not all these variants could be mapped automatically and those that did not map to Ensembl were formatted manually for input to *Module 6*.

Module 6 - Consolidation as table and VCF file

Module 6 combines all the tables produced by the previous modules and creates Venn diagrams showing the number of variants obtained from the different sources and their overlap. This table is then used to create a Variant Calling Format (VCF) file listing the site of all the variants.

We further categorized the variants in two levels of pathogenicity evidence. *Level 1* variants include the variants from ClinVar obtained in *Module 3* associated with three phenotypes: "MODY", "monogenic diabetes", or "neonatal diabetes" regardless of their reported pathogenicity, complemented with the variants extracted from the literature in *Module 4* and *Module 5*. *Level 2* variants undergo stricter filtration criteria. For the variants obtained from ClinVar in *Module 3*, all variants labeled as *benign*, *likely benign*, or of *unknown pathogenicity* were filtered out. For the variants obtained from the literature in *Module 4* and *Module 5*, variants in *BLK*, *KLF11*, and *PAX4* were removed as these genes were reported to lack association with MODY in more recent literature (6).

Translation of the variant sequences into protein sequences

The translation step was performed using the ProHap tool (https://github.com/ProGenNo/ProHap). Shortly, the variants from the table obtained in *Module 6* were mapped to canonical mRNA sequences from Ensembl to retrieve all the transcripts of the same gene with the annotation of start and stop codons. After mapping, all the sequences were translated to their amino acid sequences and written into a protein FASTA file.

Sequence overlay

All the variants from the resulting database were overlaid with the reference protein sequences obtained from Ensembl for all the transcripts of all genes. The variants are represented using two separate rows corresponding to the two levels of pathogenicity confidence (see example in figure 3). FASTA files are parsed using the Pyteomics library (19). The protein sequences and variants are plotted using the Matplotlib library (20).

figure 3. Examples of canonical isoforms of three proteins visualized with their variants positions: A. *HNF1A* gene product; B. *ABCC8* gene product; C. *KCNJ11* gene product. Each figure contains the visualization of the full sequence along with a zoomed-in fragment. The top row of blue dots represents the positions of the variants from the level 2 database, whilst the bottom, green, row represents the positions of the variants from the level 1 database. Being the most well-studied gene, *HNF1A* has an almost equal number of dots in both rows as most of the reported variants are revised and confirmed as pathogenic. *ABCC8* and *KCNJ11* have a lot of variants in the level 1 database that are not confirmed as pathogenic and, thus, are not in the level 2 database. The figure shows only one isoform of each protein as an example. Module 7 produces the visualization of all the protein isoforms. These figures can be seen in the supplementary materials on https://github.com/kuznetsovaks/MD_variants.git.

Results

Unlike more common forms of diabetes like type 2 diabetes (T2D), where large numbers of samples are available and federated initiatives consolidate information on genetic variants and their consequences in aggregated and harmonized forms (e.g. t2d.hugeamp.org), monogenic diabetes, as a rarer disease, relies on small cohorts and information on genetic variants is scattered in the literature and online databases. The aggregation and comparison of variants known to be associated with monogenic diabetes therefore currently relies on expert manual curation and annotation.

Mining variants in genes associated with monogenic diabetes

We mined variants based on a list of 101 genes associated with monogenic diabetes, aiming at being as comprehensive as possible. These 101 genes consist of: i. 14 MODY genes taken from OMIM (21) or other reviews on MODY such as (3); ii. 10 genes associated with neonatal diabetes, lipodystrophy, and insulin signaling taken from (2); iii. 77 genes having variants with any evidence of association with either MODY, neonatal diabetes, or just the condition referred to as "monogenic diabetes" by ClinVar.

Categorizing by consequence and pathogenicity

Since variants associated with monogenic diabetes are highly penetrant, we focused on exonic variants when mining Ensembl. Nevertheless, some variants reported in ClinVar and in the literature mapped to untranslated regions (UTR) and splice regions. In these cases we

decided whether to keep these variants or not based on the pathogenicity of variants in each consequence type category. To produce protein sequences, we focused on predicted consequences 'missense variant', 'protein altering variant', or 'coding sequence variant', and ignored variants on a transcript that were 'non-coding transcript variant' or 'synonymous variant'. For other types of consequences we focused on those presenting more than 80 % pathogenic and likely pathogenic variants: 'missense variant', 'protein altering variant', 'coding sequence variant', 'frameshift variant', 'splice donor variant', 'splice acceptor variant', 'stop retained', 'stop gained', 'inframe insertion', and 'inframe deletion' (figure 2). The 'stop lost' consequences yielded different prevalence of pathogenic or likely pathogenic variants when considering the consequences reported by Ensembl vs. ClinVar (figure 2), which can be explained by the differences between Ensembl and ClinVar. Here, the Ensembl dataset consisted of the exonic variants, but is not associated with any pathological phenotype. The ClinVar dataset, though, was not narrowed down to any particular regions in the genes of interest, but all the variants in it are associated with clinical conditions. Therefore, the Ensembl dataset is enriched with protein coding variants, whereas the ClinVar dataset is enriched with pathogenic variants. 'stop lost' consequences were included in our analysis. Altogether, the resulting table variants contained 69,256 unique variants located in 101 genes.

Variant association with monogenic diabetes

We could map 2,701 of these variants to variants associated with monogenic diabetes according to ClinVar or the literature, termed *level 1* variants thereafter: 2,220 (82 %) mapped uniquely to ClinVar, 136 (5 %) to publications reviewed by Rafique et al., and 345 (13 %) to both (figure 4). An effect on protein sequences was predicted for 2,624 (97 %) of them, resulting in 12,643 different protein sequences when accounting for all isoforms. The reason for some variants not reaching the final translated sequences is that some transcripts in Ensembl do not have translated products. These are included when selecting exons in *Module 1* but do not yield protein sequences. On the other hand, genetic variation can cause translation of UTRs and other normally untranslated regions (22), and genetic variation in the UTRs and splice regions might affect the translation of the proteins in an indirect way (23,24), but the effects of these

variants on amino acid sequences remain challenging to predict. We further filtered the variants to retain only the pathogenic and likely pathogenic variants, termed *level 2* variants thereafter, yielding 876 variants, of which 641 (73 %) mapped uniquely to ClinVar, 160 (18 %) to publications reviewed by Rafique et al., and 75 (9 %) to both (figure 4). An effect on protein sequences was predicted for 714 (82 %) of them, resulting in 3,776 distinct protein sequences.

In both cases, the overlap between variants from ClinVar and the literature is limited. This can be explained by the fact that the ClinVar dataset consisted of the variants associated with all kinds of monogenic diabetes (101 genes) and the Rafique et al. dataset consisted only of the MODY variants (14 genes). Thus, we re-ran the pipeline considering only MODY variants from ClinVar. The overlap of the ClinVar MODY dataset was then 18 % and 10 % for the level 1 and 2 variants respectively. While the overlap was improved, it is still limited. This illustrates the importance to consider different sources of information when studying monogenic diabetes, and rare diseases in general.

figure 4. Venn diagram representing the number of variants in two levels of the database and the number of variants taken from different sources. Level 1 database (at the left) consists of i. ClinVar variants associated with three phenotypes, i.e. "MODY", "monogenic diabetes", and "neonatal diabetes" mapped to Ensembl; ii. All variants from Rafique et al. mapped to Ensembl. Level 2 database (at the right) consists of i. ClinVar "pathogenic" + "likely pathogenic" variants; ii. Variants from Rafique et al. excluded *BLK*, *KLF11,* and *PAX4*.

Distribution of variants among the genes

We observed strong disparities in the number of variants associated with monogenic diabetes among the different genes. *HNF1A*, *GCK*, and *HNF1B* are the genes presenting most variants in both level 1 and level 2 variants, indicating that most of the monogenic diabetes-associated variants have been reported in these genes, as well as most of the ones confirmed as pathogenic or likely pathogenic (figure 3). Conversely, genes like *ABCC8* and *KCNJ11* feature many level 1 variants, only few are confirmed to be pathogenic or likely pathogenic (figure 3). In fact, of all the genes observed, just about a third bear more than 20 variants, and only around 5 % have more than 100 variants. Figure 5 represents the number of gene variants in both levels of the database, where the number of variants in the gene is more than 20. The table and figure "number of variants per gene" in the supplementary materials on GitHub represents the number of variants in each gene in both levels of the database. Besides the biological reasons that some genes have more association with monogenic diabetes than others, this effect can be due to study biases or heterogeneous levels of information on these variants. For example, the most common genes known to cause MODY (*HNF1A*, *GCK*, *HNF1B*, and *HNF4A*)(2) might have received more attention than others (e.g. *ABCC8* and *KCNJ11*), yielding a lower confirmation rate for the pathogenicity of the variants. At the same time, *ABCC8* is reported to be also associated with other types of diabetes (25) and definitely deserves attention from the perspective of MODY associations as this may bring the researchers closer to the overall understanding of diabetes causal mechanisms and more precise treatment outcome predictions. The variant distribution in the studied genes can be observed in the figures visualizing the variants in the protein sequences (figure 3 and others in the "figure" directory at https://github.com/kuznetsovaks/MD_variants.git).

figure 5. Number of variants in the genes with the largest number of variants. Selected are the genes in which the number of variants in the *level 1* database is more than 20. The full bars represent the number of variants in the *level 1* database, and the brown part of the bars represents the number of variants in the *level 2* database.

Discussion

In this work, we presented a computational pipeline that allows for systematic monogenic diabetes-associated variant collection and mapping. The sources of information on the genetic variation are not unified which makes mapping of the variants challenging. An automated and reproducible pipeline for variant mapping has been developed and is available for public use. A database of variant protein sequences was created for the gene products of variants associated with monogenic diabetes. All known variants reported to be associated with monogenic diabetes published by the beginning of 2023 have been included in the database. The database contains variants with two levels of clinical significance: variants ever reported as associated with monogenic diabetes and pathogenic variants. Here we were considering the variants pathogenic if they had *pathogenic* or *likely pathogenic* clinical significance according to ClinVar. All the monogenic diabetes-associated variants have been translated into protein products and can be compared to the canonical protein sequences. This will help predict the effect of genetic variation on the resulting protein structure and function.

The workflow is automated and aims to gather multiple variants from different sources and avoid their manual annotation. The implementation in Jupyter notebooks provides a good trade-off between automation and flexibility. For example, researchers can execute the entire pipeline as is, adapt it to specific use cases, execute only modules of interest, or completely change the set of genes to study another disease. The public availability, extensive documentation, and permissive license further enable the reuse of our work.

The collected genetic variants associated with monogenic diabetes have been translated to protein sequences and mapped to all known protein isoforms resulting in the collection of all predicted protein variant sequences. Our database is represented in the form of tables along with FASTA files and can be accessed both manually and automatically allowing implementation in various workflows. The localization of variants on proteins and protein domains can shed light on their possible consequences and pathogenicity. In turn, overlaying variant pathogenicity on protein sequences can help understanding protein function. The FASTA files can be

supplemented with other proteins and used for proteomic search of mass spectrometry data. The variant protein sequences can also be used in protein structure modeling using tools like Alphafold (11).

This work illustrates how the different genes associated with monogenic diabetes show very different levels of annotation. Besides well-investigated genes, featuring a high number of variants with unambiguous consequences, many understudied genes bear variants lacking evidence of pathogenicity.

Furthermore, some variants simply lack basic genomic annotation, and are reported as amino acid changes, e.g. "Gly292Argfs". An amino acid substitution cannot always be mapped to a single genetic variant. Furthermore, most of the genes encode several protein isoforms and knowing an amino acid change does not give information on which isoform it affects and how it maps to other isoforms. Thus, reporting single amino acid substitutions impairs their inclusion in genomic and bioinformatic studies. In this work, we manually curated these variants and were able to match some of them to genomic coordinates using other variants and aligning protein isoforms using the IsoAligner tool (26). For complex proteomic research in humans, it is important to account for common variation. In future work, we are going to combine our variant sequences with the database of human protein haplotypes (27).

Our work focused on single nucleotide variants (SNVs) or short indels and therefore does not cover larger mutations causing monogenic diabetes. Large genomic rearrangements, such as full deletion of the *HNF1B* gene (28) or full deletion of 17q12 locus, have been shown to cause HNF1B-MODY (MODY 5) and can be missed by conventional point mutation screening (28,29). These events cannot be directly translated to protein sequences and, as a result, are not reflected in our database.

The research of monogenic diabetes is a dynamically developing field, and new variants are being constantly reported from different cohorts. Now with the pipeline we have developed, we and others can easily update MODY variant collections as new variants are reported. Our pipeline can further be used altogether or in parts to study other diseases. This can enable researchers to collect automatically and reproducibly collect variants associated with

phenotypes of interest and consolidate them to a unified format. In research on rare diseases, the availability of flexible pipelines based on notebooks represents a good compromise between manual expert curation that lacks reproducibility and automated pipelines that cannot be tailored to the application.

Acknowledgments

This work was supported by the Research Council of Norway (project #301178 to MV), the University of Bergen, and the Novo Nordisk Foundation (project NNF20OC0063872 to SJ).

This research was funded, in whole or in part, by the Research Council of Norway 301178. A CC BY or equivalent license is applied to any Author Accepted Manuscript (AAM) version arising from this submission, in accordance with the grant's open access conditions.

Data Availability

The source code of all the modules is freely available as Jupyter notebooks at https://github.com/kuznetsovaks/MD_variants.git and the resulting tables are deposited along with all the figures and the results in the same repository. Large FASTA files are available at https://doi.org/10.6084/m9.figshare.21444963.v1 All our code and data are openly available under a permissive CC-BY license.

Competing interests

The authors declare no competing interests.

References

1. Beltrand J, Busiah K, Vaivre-Douret L, Fauret AL, Berdugo M, Cavé H, et al. Neonatal Diabetes Mellitus. Front Pediatr. 2020;8:540718.
2. Kavvoura FK, Owen KR. Monogenic diabetes. Medicine (Baltimore). 2019 Jan;47(1):16–21.
3. Aarthy R, Aston-Mourney K, Mikocka-Walus A, Radha V, Amutha A, Anjana RM, et al. Clinical features, complications and treatment of rarer forms of maturity-onset diabetes of the young (MODY) - A review. J Diabetes Complications. 2021 Jan 1;35(1):107640.
4. Shepherd MH, Shields BM, Hudson M, Pearson ER, Hyde C, Ellard S, et al. A UK nationwide

prospective study of treatment change in MODY: genetic subtype and clinical characteristics predict optimal glycaemic control after discontinuing insulin and metformin. Diabetologia. 2018 Dec 1;61(12):2520–7.

5. Sun BB, Kurki MI, Foley CN, Mechakra A, Chen CY, Marshall E, et al. Genetic associations of protein-coding variants in human disease. Nature. 2022 Mar;603(7899):95–102.

6. Laver TW, Wakeling MN, Knox O, Colclough K, Wright CF, Ellard S, et al. Evaluation of Evidence for Pathogenicity Demonstrates That BLK, KLF11, and PAX4 Should Not Be Included in Diagnostic Testing for MODY. Diabetes. 2022 May;71(5):1128–36.

7. Duzkale H, Shen J, McLaughlin H, Alfares A, Kelly MA, Pugh TJ, et al. A systematic approach to assessing the clinical significance of genetic variants. Clin Genet. 2013 Nov;84(5):453–63.

8. Landrum MJ, Lee JM, Benson M, Brown GR, Chao C, Chitipiralla S, et al. ClinVar: improving access to variant interpretations and supporting evidence. Nucleic Acids Res. 2018 Jan 4;46(D1):D1062–7.

9. Martagón AJ, Bello-Chavolla OY, Arellano-Campos O, Almeda-Valdés P, Walford GA, Cruz-Bautista I, et al. Mexican Carriers of the HNF1A p.E508K Variant Do Not Experience an Enhanced Response to Sulfonylureas. Diabetes Care. 2018 Aug 1;41(8):1726–31.

10. Tebani A, Afonso C, Marret S, Bekri S. Omics-Based Strategies in Precision Medicine: Toward a Paradigm Shift in Inborn Errors of Metabolism Investigations. Int J Mol Sci. 2016 Sep 14;17(9):1555.

11. Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models. Nucleic Acids Res. 2022 Jan 7;50(D1):D439–44.

12. Kind L, Raasakka A, Molnes J, Aukrust I, Bjørkhaug L, Njølstad PR, et al. Structural and biophysical characterization of transcription factor HNF-1A as a tool to study MODY3 diabetes variants. J Biol Chem. 2022 Apr;298(4):101803.

13. Aebersold R, Agar JN, Amster IJ, Baker MS, Bertozzi CR, Boja ES, et al. How many human proteoforms are there? Nat Chem Biol. 2018 Mar;14(3):206–14.

14. Cunningham F, Allen JE, Allen J, Alvarez-Jarreta J, Amode MR, Armean IM, et al. Ensembl 2022. Nucleic Acids Res. 2022 Jan 7;50(D1):D988–95.

15. Rafique I, Mir A, Saqib MAN, Naeem M, Marchand L, Polychronakos C. Causal variants in Maturity Onset Diabetes of the Young (MODY) – A systematic review. BMC Endocr Disord. 2021 Nov 11;21(1):223.

16. Yates A, Beal K, Keenan S, McLaren W, Pignatelli M, Ritchie GRS, et al. The Ensembl REST API: Ensembl Data for Any Language. Bioinforma Oxf Engl. 2015 Jan 1;31(1):143–5.

17. McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, et al. The Ensembl Variant Effect Predictor. Genome Biol. 2016 Jun 6;17(1):122.

18. Pedersen BS, Layer RM, Quinlan AR. Vcfanno: fast, flexible annotation of genetic variants. Genome Biol. 2016 Jun 1;17(1):118.

19. Levitsky LI, Klein JA, Ivanov MV, Gorshkov MV. Pyteomics 4.0: Five Years of Development of a Python Proteomics Framework. J Proteome Res. 2019;18(2):709–14.

20. Hunter JD. Matplotlib: A 2D Graphics Environment. Comput Sci Eng. 2007 May;9(3):90–5.

21. Hamosh A, Scott AF, Amberger JS, Bocchini CA, McKusick VA. Online Mendelian Inheritance in Man (OMIM), a knowledgebase of human genes and genetic disorders. Nucleic Acids Res. 2005 Jan 1;33(Database issue):D514-517.

22. Umer HM, Audain E, Zhu Y, Pfeuffer J, Sachsenberg T, Lehtiö J, et al. Generation of ENSEMBL-based proteogenomics databases boosts the identification of non-canonical peptides. Bioinforma Oxf Engl. 2021 Dec 14;btab838.

23. Hull J, Campino S, Rowlands K, Chan MS, Copley RR, Taylor MS, et al. Identification of

Common Genetic Variation That Modulates Alternative Splicing. PLOS Genet. 2007 Jun 15;3(6):e99.

24.   Steri M, Idda ML, Whalen MB, Orrù V. Genetic Variants in mRNA Untranslated Regions. Wiley Interdiscip Rev RNA. 2018 Jul;9(4):e1474.

25.   Haghverdizadeh P, Sadat Haerian M, Haghverdizadeh P, Sadat Haerian B. ABCC8 genetic variants and risk of diabetes mellitus. Gene. 2014 Jul 25;545(2):198–204.

26.   Hanimann J, Moch H, Zoche M, Kahraman A. IsoAligner: dynamic mapping of amino acid positions across protein isoforms [Internet]. F1000Research; 2022 [cited 2022 Nov 18]. Available from: https://f1000research.com/articles/11-382

27.   Vašíček J, Skiadopoulou D, Kuznetsova KG, Johansson S, Njølstad PR, Bruckner S, et al. Identifying Protein Haplotypes by Mass Spectrometry [Internet]. bioRxiv; 2022 [cited 2023 Jan 18]. p. 2022.11.21.517096. Available from: https://www.biorxiv.org/content/10.1101/2022.11.21.517096v1

28.   Bellanné-Chantelot C, Clauin S, Chauveau D, Collin P, Daumont M, Douillard C, et al. Large genomic rearrangements in the hepatocyte nuclear factor-1beta (TCF2) gene are the most frequent cause of maturity-onset diabetes of the young type 5. Diabetes. 2005 Nov;54(11):3126–32.

29.   Mefford HC, Clauin S, Sharp AJ, Moller RS, Ullmann R, Kapur R, et al. Recurrent Reciprocal Genomic Rearrangements of 17q12 Are Associated with Renal Disease, Diabetes, and Epilepsy. Am J Hum Genet. 2007 Nov;81(5):1057–69.