Rare disease research workflow using multilayer networks elucidates the molecular determinants of severity in Congenital Myasthenic Syndromes

Iker Núñez-Carpintero [1,*], Emily O'Connor [2,4,*], Maria Rigau [1,5,6], Mattia Bosio [1,7], Yoshiteru Azuma [8,9], Ana Topf [10,11], Rachel Thompson [2], Peter A.C. 't Hoen [12], Teodora Chamova [13], Ivailo Tournev [13,14], Velina Guergueltcheva [15], Steven Laurie [16], Sergi Beltran [16,17,18], Salvador Capella [1,7], Davide Cirillo [1,#], Hanns Lochmüller [2,3,4,16,19], Alfonso Valencia [1,20]

[1] *Barcelona Supercomputing Center (BSC), C/ Jordi Girona 29, 08034, Barcelona, Spain*

[2] *Children's Hospital of Eastern Ontario Research Institute; Ottawa, Canada*

[3] *Division of Neurology, Department of Medicine, The Ottawa Hospital; Ottawa, Canada*

[4] *Brain and Mind Research Institute, University of Ottawa, Ottawa, Canada*

[5] *MRC London Institute of Medical Sciences, Du Cane Road, London, W12 0NN, United Kingdom*

[6] *Institute of Clinical Sciences, Faculty of Medicine, Imperial College London, Hammersmith Hospital Campus, Du Cane Road, London, W12 0NN, United Kingdom*

[7] *Spanish National Bioinformatics Institute Unit, Structural Biology and BioComputing Programme, Spanish National Cancer Research Centre (CNIO), Madrid 28029, Spain*

[8] *Department of Human Genetics, Yokohama City University Graduate School of Medicine, Yokohama, Japan*

[9] *Department of Pediatrics, Aichi Medical University, Nagakute, Japan*

[10] *John Walton Muscular Dystrophy Research Centre, Translational and Clinical Research Institute, Newcastle University, Newcastle upon Tyne, United Kingdom*

[11] *Newcastle Hospitals NHS Foundation Trust, Newcastle upon Tyne, United Kingdom*

[12] *Center for Molecular and Biomolecular Informatics, Radboud Institute for Molecular Life Sciences, Radboud university medical center, Nijmegen, The Netherlands*

[13] *Department of Neurology, Expert Centre for Hereditary Neurologic and Metabolic Disorders, Alexandrovska University Hospital, Medical University-Sofia, Sofia, Bulgaria*

[14] *Department of Cognitive Science and Psychology, New Bulgarian University, Sofia 1618, Bulgaria*

[15] *Clinic of Neurology, University Hospital Sofiamed, Sofia University St. Kliment Ohridski, Sofia, Bulgaria.*

[16] *Centro Nacional de Análisis Genómico (CNAG-CRG), Center for Genomic Regulation, Barcelona Institute of Science and Technology (BIST), Barcelona, Catalonia, Spain*

[17] *Universitat Pompeu Fabra (UPF), Barcelona, Spain*

[18] *Departament de Genètica, Microbiologia i Estadística, Facultat de Biologia, Universitat de Barcelona (UB), Barcelona, Spain.*

[19] *Department of Neuropediatrics and Muscle Disorders, Medical Center – University of Freiburg, Faculty of Medicine, Freiburg, Germany*

[20] *ICREA, Pg. Lluís Companys 23, 08010, Barcelona, Spain*

* these authors contributed equally

corresponding author: davide.cirillo@bsc.es

Abstract

Exploring the molecular basis of disease severity in rare disease scenarios is a challenging
task provided the limitations on data availability. Causative genes have been described for
Congenital Myasthenic Syndromes (CMS), a group of diverse minority neuromuscular
junction (NMJ) disorders; yet a molecular explanation for the phenotypic severity differences
remains unclear. Here, we present a workflow to explore the functional relationships between
CMS causal genes and altered genes from each patient, based on multilayer network analysis
of protein-protein interactions, pathways and metabolomics.

Our results show that CMS severity can be ascribed to the personalized impairment of
extracellular matrix components and postsynaptic modulators of acetylcholine receptor
(AChR) clustering. Moreover, reducing expression of the zebrafish orthologue, we confirm
the effect on movement and NMJ morphology of a gene previously unknown to be a NMJ
interactor, USH2A.

This work showcase how coupling multilayer network analysis with personalized -omics
information provides molecular explanations to the varying severity of rare diseases; paving
the way for sorting out similar cases in other rare diseases.

Keywords: multi-omics data, network biology, multilayer networks, personalized medicine,
applied network science, network community analysis, rare diseases, congenital myasthenic
syndromes.

Introduction

Understanding phenotypic severity is crucial for prediction of disease outcomes, as well as for administration of personalized treatments. Different severity levels among patients presenting the same medical condition could be explained by characteristic relationships between diverse molecular entities (i.e. gene products, metabolites, etc) in each individual. In this setting, multi-omics data integration is becoming a promising tool for research, as it has the potential to gain complex insights of the molecular determinants underlying disease heterogeneity. However, even in a scenario where the level of biomedical detail available to study is growing in an exponential manner (Karczewski and Snyder, 2018), the analysis of the molecular determinants of disease severity is not typically adressed in rare disease research literature (Boycott et al. 2013), despite its obvious relevance at the medical and clinical level. Rare diseases represent a challenging setting for the application of precision medicine because, by definition, they affect a small number of patients, and therefore the data available for study is considerably limited in comparison to other conditions. Accordingly, leveraging the wealth of biomedical knowledge of diverse nature coming from publicly available databases have the potential to adress data limitations in rare diseases (Mitani and Haneuse, 2020) (Buphamalai et al, 2021). In this sense, multilayer networks can offer an holistic representation of biomedical data resources (Halu et al. 2017) (Gosak et al. 2018), which may allow to explore the biology related to a given disease independently of cohort sizes and their available omics data.

Here, in order to evaluate and demonstrate the potential of multilayer networks as means of assesing severity in rare disease scenarios, we provide an illustrative case where we develop a framework for analyzing a patient cohort affected by Congenital Myasthenic Syndromes (CMS), a group of inherited rare disorders of the neuromuscular junction (NMJ). Fatigable weakness is a common hallmark of these syndromes, that affects approximately 1 patient in 150,000 people worldwide. The inheritance of CMS is autosomal recessive in the majority of patients. CMS can be considered a relevant use case because, while patients share similar clinical and genetic features (Finsterer 2019), phenotypic severity of CMS varies greatly, with patients experiencing a range of muscle weakness and movement impairment. While over 30 genes are known to be monogenic causes of different forms of CMS (**Table 1**), these genes do not fully explain the ample range of observed severities, which has been suggested to be determined by additional factors involved in neuromuscular function (Thompson et al. 2019). Examples of CMS-related genes are AGRN, LRP4 and MUSK which code for proteins that mediate communication between the nerve ending and the muscle, which is crucial for formation and maintenance of the NMJ **(Figure 1)**.

In particular, the AGRN-LRP4 receptor complex activates MUSK by phosphorylation, inducing clustering of the acetylcholine receptor (AChR) in the postsynaptic membrane allowing the presynaptic release of acetylcholine (ACh) to trigger muscle contraction (Whicher, Philbin, and Aronson 2018). Additional evidence of CMS severity heterogeneity emerged within the NeurOmics and RD-Connect projects (Lochmüller et al. 2018) studying a small population (about 100 individuals) of gypsy ethnic origin from Bulgaria.

Location	Phenotype	Inheritance	Gene
2q31.1	CMS1A, slow-channel	AD	CHRNA1
2q31.1	CMS1B, fast-channel	AR, AD	
17p13.1	CMS2A, slow-channel	AD	CHRNB1
17p13.1	CMS2C, associated with acetylcholine receptor deficiency	AR	
2q37.1	CMS3 A, slow-channel	AD	CHRND
2q37.1	CMS3 B, fast-channel	AR	
2q37.1	CMS3 C, associated with acetylcholine receptor deficiency	AR	
17p13.2	CMS4 A, slow-channel	AR, AD	CHRNE
17p13.2	CMS4 B, fast-channel	AR	
17p13.2	CMS4 C, associated with acetylcholine receptor deficiency	AR	
3p25.1	CMS5	AR	COLQ
10q11.23	CMS6, presynaptic	AR	CHAT
1q32.1	CMS7, presynaptic	AD	SYT2
1p36.33	CMS8, with pre- and postsynaptic defects	AR	AGRN
9q31.3	CMS9, associated with acetylcholine receptor deficiency	AR	MUSK
4p16.3	CMS10	AR	DOK7
11p11.2	CMS11, associated with acetylcholine receptor deficiency	AR	RAPSN
2p13.3	CMS12, with tubular aggregates	AR	GFPT1
11q23.3	CMS13, with tubular aggregates	AR	DPAGT1
9q22.33	CMS14, with tubular aggregates	AR	ALG2
1p21.3	CMS15, without tubular aggregates	AR	ALG14
17q23.3	CMS16	AR	SCN4A
11p11.2	CMS17	AR	LRP4
20p12.2	CMS18	AD	SNAP25
10q22.1	CMS19	AR	COL13A1
2q12.3	CMS20, presynaptic	AR	SLC5A7
10q11.23	CMS21, presynaptic	AR	SLC18A3
2p21	CMS22	AR	PREPL
22q11.21	CMS23, presynaptic	AR	SLC25A1
15q23	CMS24, presynaptic	AR	MYO9A
12p13.31	CMS25, presynaptic	AR	VAMP1
3p21.31	CMS, related to GMPPB	AR	GMPBB
20q13.33	CMS, presynaptic	AR	LAMA5
3p21.31	CMS, with nephrotic syndrome	AR	LAMB2
8q24.3	CMS, with plectin defect	AR	PLEC
12q24.13	CMS, related to RPH3A	AR	RPH3A
9p13.3	CMS, presynaptic, related to MUNC13-1	AR	UNC13B
2q37.1	Escobar syndrome	AR	CHRNG

Table 1. Location, phenotype, inheritance and genes involved in CMS (adapted from https://omim.org/phenotypicSeries/PS601462 and http://www.musclegenetable.fr). AR: autosomal recessive; AD: autosomal dominant.

Figure 1. A schematic depiction of the main molecular activities of known CMS causal genes (Methods) taking place at the neuromuscular junction (NMJ) in the presynaptic terminal (in blue), synaptic cleft (in white), and skeletal muscle fiber (in yellow) (for a detailed description of this system see **Supplementary Information**).

All affected individuals shared the same causal homozygous mutation (a deletion within the AChR ε subunit, *CHRNE* c.1327delG (A. Abicht et al. 1999)), however, the severity of symptoms across this cohort varies considerably regardless of age, gender and initiated therapy, suggesting the existence of additional genetic causes for the diversity of disease phenotypes. By analyzing multi-omics data, we performed an in-depth characterization of 20 CMS patients, representing the two opposite ends of the spectrum observed in the wider cohort, aiming to investigate the molecular basis of the observed differences in the individual severity of the disease. Two CMS severity levels have been identified through extensive phenotyping, namely a severe disease phenotype (8 patients) and a not-severe disease phenotype (2 intermediate and 10 mild patients) (**Suppl. Table 1**). No demographic factor (age, sex) nor clinical tests (speech, mobility, respiratory dysfunctions, among others) show a significant association with the severity classes, with exception of Forced Vital Capacity (FVC) and shoulder lifting ability (two-tailed Fisher's exact test p-values of 0.0128 and 0.0418, respectively; **Suppl. Figure 1**). We sought to interrogate whether severity was

determined by additional genetic variations impacting neuromuscular activity, on top of the causative CHRNE mutation. We analyzed three main types of genetic variations: single nucleotide polymorphisms (SNPs), copy number variations (CNVs), and compound heterozygous variants (two recessive alleles located at different loci within the same gene in a given individual). The extensive analysis of the genomic information did not render any SNPs that could be considered a unique cause of disease severity by being common to all the cases. Nevertheless, a number of CNVs and compound heterozygous variants were found to appear exclusively in the different severity groups, in one or more patients. Moreover, the compound heterozygous variants of the severe group are enriched in pathways related to the extracellular matrix (ECM) receptors, which have been proposed as a target for CMS therapy (Ito and Ohno 2018). To investigate the functional relationship between these variants and CMS severity, **we designed an analytical workflow based on multilayer networks (Figure 2)**, allowing the integration of external biological knowledge to acquire deeper functional insights. A multilayer network consists of several layers of nodes and edges describing different aspects of a system (Kivelä et al. 2014). In biomedicine, this data representation has been used to study biomolecular interactions (Zitnik and Leskovec 2017) and diseases (Halu et al. 2017), facilitating integration and interpretation of heterogeneous sources of data.

Figure 2. Analytical workflow employed to address the severity of a cohort of patients affected by Congenital Myasthenic Syndromes (CMS). A multi-scale functional analysis approach, based on multilayer networks, was used to identify the functional relationships between genetic alterations obtained from omics data (Whole Genome Sequencing, WGS; RNA-sequencing, RNAseq) with known CMS causal genes. Modules of CMS linked genes are detected using graph community detection at a resolution range (γ) (Methods) where the most prominent changes in community structure occur. Modules that emerged from this analysis were characterized at single individual level.

Several established tools for network analysis have been recently adapted for multilayer
networks, such as random walk with restart (Valdeolivas et al. 2019), community detection
algorithms (Didier, Brun, and Baudot 2015) and node embeddings (Pio-Lopez et al. 2021).
By crossing patient genomic data with the information provided by a biomedical knowledge
multilayer network, we are able to describe the functional relationships of new genetic
modifiers responsible for the different phenotypic severity levels, showcasing the potential of
multilayer networks to provide support on the personalized analysis of rare disease patients.

Results

Variants do not segregate with patient severity
We first searched for variants able to segregate the disease phenotypes (severe and not-
severe) by analyzing a large panel of mutational events (mutations in isoforms, splicing sites,
small and long noncoding genes, promoters, TSS, predicted pathogenic mutations, loss of
function mutations, among others). We could not find one single mutation or combinations of
mutations that were able to completely segregate the two groups (**Supplementary**
Information) although partial segregation can be observed (**Suppl. Table 2**). As already
described for monogenic diseases (Kousi and Katsanis 2015) and cancer (Castro-Giner,
Ratcliffe, and Tomlinson 2015), we hypothesized that distinct weak disease-promoting
effects may represent patient-specific causes to CMS severity, which bring damage to sets of
genes that are functionally related. To find these causes, we sought to search for variants with
the potential to alter gene functions, such as CNVs and compound heterozygous variants,
which have been previously reported to be key to CMS (Angela Abicht, Müller, and
Lochmüller 1993; Richard et al. 2003; Bevilacqua et al. 2017; Yang et al. 2018).

Compound heterozygous variants are functionally related
In order to explore the hypothesis that disease severity in this cohort is due to variants in
patient-specific critical elements, we sought to identify potentially damaging compound
heterozygous variants and CNVs. We analyzed the gene lists associated with these mutations
to search for evidence of alterations in relevant pathways for the severe (n=8) and not-severe
cases (n=12). We first performed a functional enrichment analysis (Methods) of the genes
with CNVs found in the two groups. The set of affected genes in the severe group is
composed of 26 unique genes (10 private to the severe group), while the not-severe group
presented 86 unique genes (**Suppl. Table 3**). None of these gene sets showed any functional
enrichment. Moreover, none of these genes had been described as causal for CMS, and none
carried compound heterozygous variants. (**Suppl. Figure 2**).
As for compound heterozygous variants, the set of affected genes in the severe group is
composed of 112 unique genes (89 private to the severe group), while the not-severe group
resulted in 152 unique genes (**Suppl. Table 3**). We found that the severe group shows
significant enrichment in genes belonging to extracellular matrix (ECM) pathways, in
particular "ECM receptor interactions" (KEGG hsa04512, adjusted p-value 0.002337) and
"ECM proteoglycans" (Reactome R-HSA-30001787, adjusted p-value 0.001237), which are

the top-hit pathways when the 89 genes appearing only in the severe group are considered.
Both these pathways share common genes, namely *TNXB*, *LAMA2*, *TNC*, and *AGRN*. The
role of extracellular matrix proteins for the formation and maintenance of the NMJ has
recently drawn attention to the study of CMS (Beeson 2016; Rodríguez Cruz, Palace, and
Beeson 2018). In particular, within the genes linked with ECM pathways, *AGRN* and *LAMA2*
stand out for their implication in CMS and other rare neuromuscular diseases (Bertini et al.
2011; Nicole et al. 2014; Bönnemann et al. 2014). ECM-related pathways are not enriched in
the not-severe set of genes (KEGG hsa04512, adjusted p-value 0.6170). Moreover, top-hit
pathways of the not-severe set of genes are not explicitly related to ECM and not consistent
between Reactome and KEGG (Reactome "Susceptibility to colorectal cancer" R-HSA-
5083636, adjusted p-value 4.131e-7, genes MUC3A/5B/12/16/17/19; KEGG "Huntington's
disease" hsa05016, adjusted p-value 0.07103, genes REST, CREB3L4, CLTCL1,
DNAH2/8/10/11). These findings support our hypothesis that the severe patients might
present disruptions in NMJ functionally related genes that, combined with *CHRNE* causative
alteration, may be responsible for the worsening of symptoms.

CMS-specific monolayer and multilayer community detection

As disease-related genes tend to be interconnected (Menche et al. 2015), we sought to
analyze the relationships among the CMS linked genes (i.e. known CMS causal genes, and
severe and not-severe compound heterozygous variants and CNVs; Methods) using network
community clustering analysis. We employed the Louvain algorithm (Methods) to find
groups of interrelated genes in three monolayer networks that represents biological
knowledge contained in databases, separately: the Reactome database (Fabregat et al. 2018),
the Recon3D Virtual Metabolic Human database (Brunk et al. 2018) (both downloaded in
May 2018), and from the Integrated Interaction Database (IID) (Kotlyar et al. 2016)
(downloaded in October 2018) (**Suppl. Figure 3**). The last two networks, represent the
'metabolome' and the 'interactome' data, respectively. By measuring community similarity
(Methods), we observed that the same CMS linked genes did not form the same communities
across the different networks (**Suppl. Figure 4**). These results show that, although disease-
related genes are prone to form well-defined communities in distinct networks (Goh et al.
2007; Cantini et al. 2015), different facets of biological information (i.e. reactome,
metabolome, interactome) reflect diverse participation modalities of such genes into
communities. In order to deliver an integrated analysis of such heterogeneous information,
we further consider them as a multilayer network (Gosak et al. 2018).

Large-scale multilayer community detection of disease associated genes

We first sought to test the hypothesis that disease-related genes tend to be part of the same
communities also in a multilayer network. We used the curated gene-disease associations
database DisGeNET (Piñero et al. 2017), showing that disease-associated genes are
significantly found to be members of the same multilayer communities (Wilcoxon test p-
value < 0.001 in a range of resolution parameters described in the Methods). We pre-
processed DisGeNET database by filtering out diseases and disease groups with only one

associated gene (6,352 diseases), and those whose number of associated genes was more than
1.5 * interquartile range (IQR) of the gene associated per disease distribution (823 diseases
with more than 33 associated genes) (**Suppl. Figure 5A-B**). This procedure prevents a
possible analytical bias due to the higher amounts of genes annotated to specific disease
groups (e.g. entry C4020899, "Autosomal recessive predisposition", annotates 1445 genes).
We then retrieved the communities of each associated gene, excluding 428 genes not present
in our multilayer network and the diseases left with only one associated gene. The final
analysis comprised a total of 5,892 diseases with an average number of 7.38 genes per
disease. For each disease, we counted the number of times that the disease-associated genes
are found in the same multilayer communities, and compared the distribution of such
frequencies with that of balanced random associations (1000 randomizations). Results show
that disease-associated genes are significantly found in the same multilayer communities
across the resolution interval (**Suppl. Figure 5C**).

Modules within the CMS multilayer communities
We define a module as a group of CMS linked genes that are systematically found to be part
of the same multilayer community while increasing the multilayer network community
resolution parameter (Methods; Supplementary Information; **Figures 3-4**).

Figure 3. Identification of the largest module containing genes that are found in the same community
in the entire range of resolution parameters (Methods). In each module, genes are connected if they
are found in the same multilayer communities at *n* values of the resolution parameter γ within the
range under consideration ($\gamma \in (0,4]$). The arrows indicate the systematic increase of *n*. At *n* = 8, the
module contains genes that are always found in the same community in the entire range of resolution
(see Supplementary Information "Multilayer community detection analysis"). The largest modules
containing the CMS linked gene set (highlighted in pink), which includes known CMS causal genes,
severe-specific heterozygous compound variants and CNVs, are shown.

Within each of these communities, we identified smaller modules of CMS linked genes that
are specific to the severe and not-severe groups. We tested the significance of obtaining these
exact genes in the severe and not-severe largest modules upon severity class label shuffling
among all individuals (1000 randomizations). We found that 13 (p-value 0.034; on average 8
CMS linked genes and 5 causal genes) and 14 (p-value 0.026; on average 8 CMS linked
genes and 4 causal genes) are the minimum number of genes composing the modules that are
not expected to be found at random in the severe and not-severe largest components,
respectively (**Suppl. Figure 6)**. In the two groups, the significantly largest module that
contains known CMS causal genes is composed of 15 genes (**Figure 4**). 6 out of these 15 are
previously described CMS causal genes (Methods), namely the ECM heparan sulfate
proteoglycan agrin (*AGRN*); the cytoskeleton component plectin (*PLEC*), causative of
myasthenic disease (Forrest et al. 2010); the agrin receptor *LRP4*, key for AChR clustering at
NMJ (Barik et al. 2014) and causative of CMS by compound heterozygous variants
(Ohkawara et al. 2014); the ECM components *LAMA5* and *LAMB2* laminins, and *COL13A1*
collagen.

Figure 4. Largest module, containing known CMS causal genes, within the multilayer communities
of CMS linked genes that are specific to the not-severe (A) and severe (B) groups. In turquoise,
compound heterozygous variants; in yellow, CNVs; in pink, known CMS causal genes. Being a CMS
causal gene bearing compound heterozygous variants, AGRN is depicted using both turquoise and
pink.

All the other genes of the two modules are involved in a varied spectrum of muscular
dysfunctions, discussed in the following sections. As the location of the causal gene products
determine the most common classification of the disease (i.e. presynaptic, synaptic, and
postsynaptic CMS) (Rodríguez Cruz, Palace, and Beeson 2018), we determined class and
localization of the members of the found modules (**Table 2**). Laminins, well-known CMS
glycoproteins, are affected in both severe (*LAMA2*, *USH2A*) and not-severe (*LAMB4*) groups,
and are bound by specific receptors that are damaged in the not-severe group (*MCAM*)

(Dagur and McCoy 2015). Collagens, known CMS-related factors, are associated with the not-severe group (*COL6A5*), and bound by specific receptors that are damaged in the not-severe group (*MSR1*) (Gowen et al. 2000).

However, overall collagen biosynthesis is affected in both severe and not-severe groups. Indeed, metalloproteinases, damaged in the not-severe group, are responsible for the proteolytic processing of lysyl oxidases (*LOXL3*), which are implicated in collagen biosynthesis (Panchenko et al. 1996) and damaged in the severe group.

Alterations in proteoglycans (*AGRN, HSPG2, VCAN, COL15A1*) (Iozzo and Schaefer 2015), tenascins (*TNC, TNXB*) (Pedrosa-Domellöf, Virtanen, and Thornell 1995), and chromogranins (*CHGB*) (Andreose, Sala, and Fumagalli 1994) are specific of the severe group. We observed no genes associated with proteoglycan damage in the not-severe group, suggesting a direct involvement of ECM in CMS severity.

Personalized analysis of the severe cases

We sought to analyze the 15 genes of the largest module of the severe group in each one of the 8 patients, hereafter referred to using the WGS sample labels (**Suppl. Table 1)**. Overall, these genes have a varied range of expression levels in tissues of interest (**Suppl. Figure 7**), for instance in skeletal muscle *HSPG2, LAMA2, PLEC* and *LAMB2* show medium expression levels (9 to 107 TPM) while the others show low expression levels (0.6 to 9 TPM) (Methods). Patient 2, a 15 years old male, presents compound heterozygous variants in tenascin C (*TNC*), mediating acute ECM response in muscle damage (Sorensen et al. 2018), and CNVs (specifically, a partial heterozygous copy number loss) in usherin (*USH2A*), which has been associated with hearing and vision loss (Austin-Tse et al. 2018).

Patient 16, a 25 years old female, presents compound variants in tenascin XB (*TNXB*), which is mutated in Ehlers-Danlos syndrome, a disease that has already been reported to have phenotypic overlap with muscle weakness (Kirschner et al. 2005; N. C. Voermans and van Engelen 2008) and whose compound heterozygous variants have been reported for a primary myopathy case (Pénisson-Besnier et al. 2013; Nicol C. Voermans et al. 2014); and versican (*VCAN*), which has been suggested to modify tenascin C expression (Keller et al. 2012) and is upregulated in Duchenne muscular dystrophy mouse models (McRae et al. 2017).

Patient 13, a 26 years old male, presents compound mutations in laminin α2 chain (*LAMA2*), a previously reported gene related to various muscle disorders (Amin et al. 2019; Løkken et al. 2015, 2; Dimova and Kremensky 2018, 2) whose mutations cause reduction of neuromuscular junction folds (Rogers and Nishimune 2017), and collagen type XV α chain (*COL15A1*), which is involved in guiding motor axon development (Guillon, Bretaud, and Ruggiero 2016) and functionally linked to a skeletal muscle myopathy (Eklund et al. 2001; Muona et al. 2002).

Patient 12, a 49 years old female, presents compound mutations in chromogranin B4 (*CHGB*), potentially associated with amyotrophic lateral sclerosis early onset (Pampalakis et al. 2019). Patient 18, a 51 years old man, presents compound mutations in agrin (*AGRN*), a CMS causal gene that mediates AChR clustering in the skeletal fiber membrane (Huzé et al. 2009).

Activity localization	Class	CMS causal gene	Phenotype group		Function	Synaptic localization (Manual curation)	Localization (UniProt)
			Not-severe	Severe			
ECM (ECM)	Proteoglycans	AGRN	-	AGRN	Cell hydration and growth factor trapping	Pre- and postsynaptic (PMID: 29462312)	Synaptic basal lamina / ECM
		-	-	HSPG2		Basement membrane (PMID:30453502)	Basement membrane / ECM
		-	-	VCAN		ECM (PMID:29211034)	ECM
		-	-	COL15A1		Basement membrane (PMID:26937007)	ECM
	Collagens	COL13A1	-	-	Structural support	Basement membrane, post-synaptic (PMID: 30768864)	Post-synaptic cell membrane
		-	COL6A5	-		Basement membrane (PMID:23869615)	Extracellular matrix
	Laminins	LAMA5	-	-	Web-like structures	Pre-synaptic (PMID:28544784)	Basement membrane / ECM
		LAMB2	-	-		Basement membrane (PMID:27614294)	Basement membrane / ECM / Synaptic cleft
		-	LAMB4	-		Myenteric plexus basement membrane (PMID: 28595269)	Basement membrane / ECM
		-	-	LAMA2		Pre-synaptic (PMID:9396756)	Basement membrane / ECM
		-	-	USH2A		Neuronal projection of stereocilia (PMID:19023448)	Stereocilia membrane / Secreted (Extracellular region)
	Fibulins	-	HMCN1	-	Scaffolding	Glomerular Extracellular matrix (PMID: 29488390)	Basement membrane / ECM
	Tenascins	-	-	TNC	Anti-adhesion	Basement membrane (PMID: 29466693)	ECM / Perisynaptic ECM (Ensembl)
		-	-	TNXB		Basement membrane (PMID: 23768946)	ECM
	Enzymes	-	-	LOXL3	Collagen assembly	Basement membrane (PMID:26954549)	Secreted (extracellular region)

Localization	Category				Function	Synaptic localization	Localization
			ADAMTS9	-	Proteoglycan cleavage	Secreted to ECM (PMID:30626608)	ECM
			ADAM28			ECM (PMID:24613731)	Cell membrane / Secreted (extracellular region)
	Neuropeptides	-	-	CHGB	Regulatory peptides precursor	Pre- and postsynaptic (PMID:7526287)	Secreted (extracellular region)
	Others	-	ITIH5	-	Hyaluronic acid binding	ECM (PMID:27143355)	Secreted (extracellular region)
Cell surface	Receptors	-	MSR1	-	Proteoglycan and collagen binding	Macrophage surface Scavenger Receptor (PMID:12488451)	Plasma membrane
			MCAM			Plasma membrane (PMID:28923978)	Plasma membrane
		LRP4	-	-	Laminin binding	Post-synaptic (PMID:25319686)	Post-synaptic cell membrane
Cytoplasm	Cytoskeleton	PLEC	-	-	Structural support	Post-synaptic (PMID:20624679)	Post-synaptic cytoskeleton

Table 2. Localization and functions of proteins encoded by the genes found in the largest modules of the multilayer communities of severe and not-severe groups. In turquoise, compound heterozygous variants; in yellow, CNVs; in pink, known CMS causal genes. Synaptic localization was retrieved from manual curation and Uniprot database (Methods).

Patient 20, a 57 years old male, presents compound mutations in lysyl oxidase-like 3 (*LOXL3*), involved in myofiber extracellular matrix development by improving integrin signaling through fibronectin oxidation and interaction with laminins (Kraft-Sheleg et al. 2016), and perlecan (*HSPG2*), a gene whose deficiency leads to muscular hypertrophy (Xu et al. 2010), that is also mutated in Schwartz-Jampel syndrome (Stum et al. 2006), Dyssegmental dysplasia Silverman-Handmaker type (*DDSH*) (Arikawa-Hirasawa et al. 2002) and fibrosis (Lord et al. 2018), such as Patient 19, a 62 years old female. Furthermore, based on the estimated familial relatedness (Methods) and personal communication (February 2018, Teodora Chamova), patients 19 and 20 are siblings (**Suppl. Table 4**).

Functional consequences of variants in the severe-specific module
Studying the functional impact of the compound heterozygous variants in the severe-specific genes of the module, we observed that **in 6 of the 8 patients at least one of the variants is predicted to be deleterious by the Ensembl Variant Effect Predictor (VEP)** (McLaren et al. 2016) (Methods; **Suppl. Table 5**). For example, as for Patient 18, who presents 3 different variants in AGRN gene, only rs200607541 is predicted to be deleterious by VEP's Condel (score = 0.756), SIFT (score = 0.02), and PolyPhen (score = 0.925).

In particular, the variant (a C>T transition) presents an allele frequency (AF) of 4.56E-03 (gnomAD exomes) (Karczewski et al. 2020) and affects a region encoding a position related to a EGF-like domain (SMART:SM00181) and a Follistatin-N-terminal like domain (SMART:SM00274). Both of these domains are part of the Kazal domain superfamily which are specially found in the extracellular part of agrins (PFAM: CL0005) (Laskowski and Kato 1980). On the other hand, Patient 16 presents a total of 38 *TNXB* transcripts affected by three gene variants (rs201510617, rs144415985, rs367685759) that are all predicted to be deleterious by the three scoring systems, have allele frequencies of 3.17E-02, 4.83E-02 and 5.90E-03, respectively; and in overall, are affecting two conserved domains. The first consists of a fibrinogen related domain that is present in most types of tenascins (SMART:SM00186), while the second is a fibronectin type 3 domain (SMART:SM00060) that is found in various animal protein families such as muscle proteins and extracellular-matrix molecules (Bork and Doolittle 1992). Two of the severe patients (Patients 12 and 19) present severe-only specific compound heterozygous variants that are not predicted to be deleterious. However, one variant in the *CHGB* gene (rs742710, AF=1.07E-01), present in patient 12, has been previously reported to be potentially causative for amyotrophic lateral sclerosis early onset (Pampalakis et al. 2019). This gene has also been strongly suggested in literature as a possible marker for multiple sclerosis (Mo et al. 2013), and other related neural diseases like Parkinson's (Nilsson et al. 2009) and Alzheimer's disease (Y. Chen et al. 2019). As for patient 19, the variant rs146309392 (AF=8.40E-04) in the gene *HSPG2* has been previously referred to be causal of Dyssegmental dysplasia as a compound heterozygous mutation (Arikawa-Hirasawa et al. 2001). This variant, as pointed out before, is shared with sibling patient 20. One severe individual (Patient 3), a 37 years old female, does not carry compound heterozygous variants included in this module but others at a lower resolution parameter value (**Suppl. Figure 8; Suppl. Table 6**). Interestingly, most of the genes carrying severe-specific deleterious compound heterozygous variants in this patient (*CDH3*, *FAAP100*, *FCGBP*, *GFY*, *RPTN*) are not related to processes at the NMJ level (Hull et al. 2016; Ramanagoudr-Bhojappa et al. 2018; Johansson, Thomsson, and Hansson 2009; Swuec et al. 2017; Kaneko-Goto et al. 2013). Nevertheless, three of these variants occur in genes potentially involved in NMJ functionality. In particular, variants rs111709242 (AF=2.64E-03) and rs77975665 (AF=3.03E-02) affect gene *PPFIBP2*, which encodes a member of the liprin family (liprin-β) that has been described to control synapse formation and postsynaptic element development (Bernadzki et al. 2017; Astigarraga et al. 2010). Furthermore, the variant rs111709242 is predicted to be deleterious by the SIFT algorithm (see **Suppl. Table 6**). Moreover, variant rs151154986 (AF=2.18E-02) affects the acyl-CoA thioesterase *ACOT2*, which generate CoA and free fatty acids from acyl-CoA esters in peroxisomes (Grevengoed, Klett, and Coleman 2014). A role for CoA levels in skeletal muscle for this enzyme class has been previously described (Li et al. 2015). Interestingly, PPFIBP2 appears in modules at lower resolution parameter values associated with known CMS causal genes (e.g. *DOK7*, *RPSN*, *RPH3A*, *VAMP1*, *UNC13B*). Moreover, this patient presents high relatedness with three not-severe patients (Patients 8, 9, and 10) who in turn display a very high relatedness among them (**Suppl. Table 4**).

Potential pharmacological implications

Finding a genetic diagnosis might help select the appropriate medication for each patient. For instance, fluoxetine and quinine are used for treating the slow-channel syndrome, an autosomal dominant type of CMS caused by mutations affecting the ligand binding or pore domains of AChR, but this treatment should be avoided in patients with fast-channel CMS (Engel et al. 2015).

Within our cohort, 13 (7 mild, 2 moderate and 4 severe) out of 20 individuals from our CMS cohort are receiving a pharmacological treatment consisting of pyridostigmine, an acetylcholinesterase inhibitor used to treat muscle weakness in myasthenia gravis and CMS (Lee, Beeson, and Palace 2018). This treatment slows down acetylcholine hydrolysis, elevating acetylcholine levels at the NMJ, which eventually extends the synaptic process duration when the AChR are mutated. Although the severity could potentially be related to how well a patient responds to the standard treatment with the AchE inhibitors, we could not find a clear correlation between severity and pyridostigmine treatment (two-tailed Fisher's exact test p-value 0.356; **Suppl. Figure 1**).

In Addition to the causal mutation in *CHRNE*, our results indicate that severity is related to AChR clustering at the Agrin-Plectin-LRP4-Laminins axis level, suggesting the potential benefit of pharmaceutical intervention enhancing the downstream process of AChR clustering. For example, beta-2 adrenergic receptor agonists like ephedrine and salbutamol have been documented as capable of enhancing AChR clustering (Clausen, Cossins, and Beeson 2018) and proved to be successful in the treatment for severe AChR deficiency syndromes (Rodríguez Cruz et al. 2015) (Garg and Goyal, 2022). Furthermore, the addition of salbutamol in pyridostigmine treatments have been described as being able to ameliorate the possible secondary effects of pyridostigmine in the postsynaptic structure (Vanhaesebrouck et al. 2019).

Experimental validation of USH2A involvement at the NMJ

To determine the potential relevance of one of our identified potential modifiers with no previously published relationship to the NMJ, we analyzed its function using zebrafish. For this we chose *USH2A,* a gene associated with Usher syndrome and Retinitis pigmentosa in humans (OMIM ID 608400, https://omim.org/), which was identified as a copy number loss in patient 2. While we expect the phenotypic outcome (more severe disease) of this genetic difference to manifest when expressed in conjunction with the *CHRNE* mutation causing this patients' CMS, we hypothesized that knockdown of *USH2A* expression alone may cause detectable NMJ impairments. Therefore, we used a MO to knockdown the expression of the zebrafish orthologue; *ush2a,* and studied the effects on survival, development and NMJ function. Zebrafish *ush2a* is expressed from 1 to 5 dpf, as shown in **Suppl. Figure 9A**. Using a MO targeting the exon 3/intron 3 splice donor site we were able to decrease expression of *ush2a* with a 6 ng to 18 ng MO injection (**Suppl. Figure 9B**). Survival of control and *ush2a*-MO zebrafish was not significantly affected as compared to wildtype (WT) fish over 5 dpf (log-rank test, WT n = 574, control MO 4 ng n = 46, 6 ng n = 75, 18 ng n = 34, *ush2a*-MO 2 ng n = 72, 4 ng n = 68, 6 ng n = 360, 12 ng n = 288, 18 ng n = 139, **Suppl. Figure 9C**).

There were no obvious gross morphological differences between control MO and *ush2a*-MO
fish up to 5 dpf (representative images of 2 dpf fish shown in **Suppl. Figure 9D**). As length
is an indicator of developmental stage, we measured the length of 18 ng injected *ush2a*-MO
fish at 2 dpf and found a significant reduction in length as compared to controls ($p = 0.013$, t
= 2.59, df = 38, unpaired t-test, control MO n = 20, *ush2a*-MO n = 20, **Suppl. Figure 9E**).
Eye area can be reduced in zebrafish models of retinitis pigmentosa, the condition that
USH2A mutations are associated with in humans. We measured eye area in 2 dpf fish and
found it to be significantly reduced in 18 ng-injected *ush2a*-MO fish as compared to controls
($p = 0.0006$, t = 3.73 df = 38, unpaired t-test, control MO n = 20, *ush2a*-MO n = 20,
Supplementary Figure 9F).

Figure 5. Early movement behaviors in *ush2a*-MO zebrafish. (A) Chorion rotations per minute (burst
count), and (B) mean chorion rotation duration in seconds for control and *ush2a*-MO-injected
zebrafish at 1 days post fertilization (dpf). (C) Average velocity and (D) initial acceleration of control
and *ush2a*-MO zebrafish at 2 dpf in response to touch. Dashed line shows the median, dotted lines
show the quartiles, **p < 0.01, ****p < 0.0001, ns = not significant, Mann Whitney test (A and B),
unpaired t-test (C and D).

Eye area remains significantly different after normalizing for body length (data not shown).
CMS manifests as fatigable muscle weakness in patients and in developing zebrafish we can
study the ability of fish to perform repetitive, well-characterized movements during
development to determine whether impairments to the functioning of the neuromuscular
system may be present. We quantified the number and duration of chorion movements in 1
dpf fish following administration of a control or 18 ng *ush2a*-MO. This revealed a significant
decrease in the number of burst events performed per minute in knockdown fish as compared
to controls (p = 0.003, Mann Whitney test, control MO n = 84, *ush2a*-MO n = 74, **Figure**
5A). The average duration of each burst event was not significantly affected by loss of Ush2a
(p = 0.467, Mann Whitney test, control MO n = 72, *ush2a*-MO n = 49, **Figure 5B**). To
ascertain whether impairments to movement are present in the knockdown fish while
swimming free of the chorion, we also performed a touch response assay at 2 dpf. We
observed a significant decrease in average velocity of the fish injected with *ush2a*-MO as
compared to control MO in response to a touch stimulus (p < 0.0001, t = 4.42, df = 48,
unpaired t-test; n = 25, **Figure 5C**). There was no significant difference in acceleration of
ush2a-MO fish as compared to controls (p = 0.263, t = 1.13 df = 47, unpaired t-test; control
MO n = 24, *ush2a*-MO n = 25, **Figure 5D**).
To determine whether changes in movement are reflected at the level of gross NMJ structure,
analysis of NMJ morphology was performed on 2 dpf zebrafish (**Figure 6A**). A significant
decrease in the number of SV2-positive clusters per 100 µm^2 (representative of the pre-
synaptic motor neurons) was identified on the fast muscle fibers of *ush2a*-MO fish as
compared to controls (p = 0.0004, Mann Whitney test, control MO n = 11, *ush2a*-MO n = 15,
Figure 6B). SV2-positive clusters overlie postsynaptic AChRs to form NMJs and these
receptors can be detected with fluorophore-labelled α-bungarotoxin. Analysis of AChR
clusters revealed no significant differences in number per 100 µm^2 between the two
conditions (p = 0.217, Mann Whitney test, control MO n = 11, *ush2a*-MO n = 15, **Figure**
6C). Colocalization analysis revealed no significant differences in co-occurrence of SV2 and
AChR on fast muscle fibers (SV2 colocalization with AChRs: p = 0.371, t = 0.911, df = 24,
nested t-test, **Figure 6D** and AChR colocalization with SV2: p = 0.372, t = 0.909, df = 24,
control MO n = 11, *ush2a*-MO n = 15, nested t-test, **Figure 6E**). There was also no
significant difference in colocalization of SV2 with AChRs on slow muscle, however, a
significant reduction in co-occurrence of AChRs with SV2 is present on *ush2a*-MO slow
muscle (SV2 colocalization with AChRs: p = 0.516, t = 0.660, df = 24, nested t-test, **Figure**
6F and AChR colocalization with SV2: p = 0.002, t = 3.41, df = 24, control MO n = 11,
ush2a-MO n = 15, nested t-test, **Figure 6G**). Movement differences in zebrafish may also be
caused by changes in muscle growth and development. Therefore, we assessed 2 dpf fish for
gross phenotypic differences in muscle fiber orientation and structure using a phalloidin stain
to detect actin in muscles (**Suppl. Figure 10A)**. We identified no significant differences in
muscle fiber dispersion (organization) or myotome size between *ush2a*-MO and control-MO
zebrafish (p = 0.922, t = 0.099, df = 24 unpaired t-test and p = 985, t = 0.019, df = 24 nested
t-test, respectively. Control MO n = 11 and *ush2a*-MO n = 15, **Suppl. Figure 10B, C**).

Figure 6. Neuromuscular junction morphology in *ush2a*-MO zebrafish. (A) Representative images of neuromuscular junctions from control and *ush2a*-MO zebrafish at 2 days post fertilization (dpf). Acetylcholine receptors (AChRs) are stained with fluorophore bound α-bungarotoxin (aBt, cyan), and motor neurons detected with an antibody against synaptic vesicle protein 2 (SV2, magenta). Scale bar = 50 µm. (B) Number of SV2-positive clusters and (C) number of aBt-positive clusters per 100 μm^2. (D) Colocalization of SV2 with aBt and (E) colocalization of αBT with SV2 on fast muscle cells, using Mander's correlation coefficient (0 = no colocalization, 1 = full colocalization). (F) Colocalization of SV2 with aBt and (G) colocalization of aBt with SV2 on slow muscle cells at the myosepta, using Mander's correlation coefficient. Dashed line shows the median, dotted lines show the quartiles, **$p < 0.05$, ***$p < 0.001$, ns = not significant, nested t-test.

Discussion

In this work, we have developed a framework for the analysis of disease severity in scenarios heavily impacted by sample size. Presenting limited numbers of cases is one of the main obstacles for the application of precision medicine methods in rare disease research, as it critically affects the level of expected statistical power, a common hallmark in the analysis of minority conditions (Whicher, Philbin, and Aronson 2018). This fact difficults exploring the molecular relationships that define the inherently heterogeneous levels of disease severity observed in rare disease populations, making it an atypically addressed biomedical problem (Boycott et al. 2013). Our approach, based on the application of multilayer networks, enable the user to account for the many interdependencies that are not properly captured by a single source of information, effectively combining the available patient genomic information with general biomedical knowledge from relevant databases representing different aspects of molecular biology. The application to a relevant clinical case, where we tested the hypothesis that the severity of CMS is determined by patient-specific alterations that impact NMJ functionality, provided evidence on how the methodology is able to recover the molecular relationships between the candidate patient-specific genomic variants, the observed causal AChR mutation and previously described CMS causal genes **(Table 1).**

Our in-depth functional analysis focused on a cohort of 20 CMS patients, from a narrow, geographically isolated and ethnically homogenous population, who share the same causative mutation in the AChR ε subunit (CHRNE) but present with different levels of severity. The isolation and endogamy that characterize the population from which these patients come from might have favored the accumulation of damaging variants (Fareed and Afzal 2017; Petukhova et al. 2009), giving rise to the emergence of compound effects on relevant genes for CMS. This observation has previously been made in similar syndromes (Ohno 2003; Müller et al. 2004) and in a number of other neuromuscular diseases (Zhong et al. 2017; Wang et al. 2018). In CMS, compound heterozygous variants are known to be concentrated in CHRNE (Thompson et al. 2019). The initial analysis of compound heterozygous variants revealed a significant enrichment of functional categories that are specific to the severe cases, namely ECM functions. This suggests the existence of functional relationships between major actors of the NMJ that are affected by severity-associated damaging mutations. Such interactors include already known CMS causal genes (e.g. AGRN, LRP4, PLEC) as well as genes known to interact with them. While severity-specific compound heterozygous variants and CNVs are observed, demographic factors (e.g. sex, age), pharmacological treatment, and personalized omics data (e.g. variant calling, differential gene expression, allele specific expression, splicing isoforms) do not segregate with patient severity.

Therefore, this motivated the developing of our multilayer network community analysis to investigate the relationship between known CMS causal genes and severity-associated variants (compound heterozygous variants and CNVs), integrating pathways, metabolic reactions, and protein-protein interactions. Recently, we used a multilayer network as a means to perform dimensionality reduction tasks for patient stratification in medulloblastoma, a childhood brain tumor (Núñez-Carpintero et al. 2021). Here, we started by analyzing DisGeNET data in order to verify that disease-associated genes tend to belong

to the same multilayer communities. We then identified stable and significantly large gene modules within our CMS cohort's multilayer communities and mapped the corresponding damaging mutations back to the single patients, providing a personalized mechanistic explanation of severity differences. Given the difficulties of cohort recruitment for rare diseases, this approach could be used to investigate forms of CMS and other phenotypically variable rare diseases caused by a common mutation (Estephan et al. 2018).

Overall, the personalized analysis of these mutations suggests that CMS severity can be ascribed to the damage of specific molecular functions of the NMJ which, despite affecting individuals in a personalized manner, involve genes belonging to distinct classes and localizations, namely ECM components (proteoglycans, tenascins, chromogranins) and postsynaptic modulators of AChR clustering (LRP4, PLEC) (Table 2). Alterations of other genes related to the production of ECM components, such as laminins and collagen, are observed but are not specific to the severity levels.

Finding a personalized genetic diagnosis for phenotypic severity might help select the appropriate medication for each patient. For instance, fluoxetine and quinine are used for treating the slow-channel syndrome, an autosomal dominant type of CMS caused by mutations affecting the ligand binding or pore domains of AChR, but this treatment should be avoided in patients with fast-channel CMS (Engel et al. 2015). Within our cohort, 13 out of 20 individuals from our CMS cohort are receiving a pharmacological treatment consisting of pyridostigmine, an acetylcholinesterase inhibitor used to treat muscle weakness in myasthenia gravis and CMS (Lee, Beeson, and Palace 2018). Although the severity could potentially be related to how well a patient responds to the standard treatment with the AchE inhibitors, we could not find a clear correlation between severity and pyridostigmine treatment (two-tailed Fisher's exact test p-value 0.356; Suppl. Figure 1). Our results indicate that severity is related to AChR clustering at the Agrin-Plectin-LRP4-Laminins axis level, suggesting the potential benefit of pharmaceutical intervention enhancing the downstream process of AChR clustering. Strikingly, beta-2 adrenergic receptor agonists like ephedrine and salbutamol have been documented as capable of enhancing AChR clustering (Clausen, Cossins, and Beeson 2018) and proved to be successful in the treatment for severe AChR deficiency syndromes (Sadeh, Shen and Engel, 2011) (Rodríguez Cruz et al. 2015) (Vanhaesebrouck et al. 2019) (Garg and Goyal, 2022), but a strong molecular explanation for the observed favorable effects was still missing. This study reinforces explainability for the described successful usage of such treatments by relating CMS phenotypic severity with the normal development of AChR clusters at the motor neuron membrane.

Several of the genes identified in this analysis do not have previous associations with the NMJ, such as the Usher syndrome and Retinitis pigmentosa associated gene; USH2a, identified as a copy number loss in patient 2. To provide proof of principal for this gene acting as a potential modifier of CMS severity, we investigated whether knockdown of ush2a, the zebrafish orthologue, could result in NMJ defects. Both CRISPR and TALEN-mediated knockout of ush2a in zebrafish have previously revealed phenotypes consistent with Usher syndrome and Retinitis pigmentosa such as hearing loss and progressive visual impairments (Han et al. 2018).

However, neither study assessed impacts on muscle structure or movement of the fish. Zebrafish perform well-characterized movements throughout development, starting with spontaneous chorion rotations from approximately 17 hours post fertilization (hpf, the time at which primary motor axons start extending into the muscle) to 30 hp (Saint-Amant & Drapeau. 1998). We treated 1-cell-stage embryos with a high dose of MO to reduce expression of ush2a (or equivalent dose of a control MO) and found a decrease in the number of chorion rotations performed at 24 hpf. These movements are mediated at the level of the spinal cord and are independent of supraspinal inputs (Downes & Granato. 2006), thus implying an early defect in NMJ or muscle development, or in signal transduction in the spinal cord/peripheral nervous system. By 2 dpf zebrafish can respond to touch and do so by rapidly swimming at least 1 body-length away from the stimulus (Saint-Amant & Drapeau. 1998). In ush2a-MO fish the average swimming velocity was significantly slower than in controls, whereas the initial acceleration (proportional to the force of muscle contraction) was unaffected (Sztal et al 2016). This implies that the initial fast muscle response is not significantly affected at this time-point, but that loss of Ush2a at the NMJs of slow muscle may be impacting swimming. Defects in movement are reported in many other zebrafish models of CMS, such as those lacking Dok7 (Müller et al. 2010), Gfpt1 (Senderek et al. 2011) and Syt2 (Wen et al. 2010). Our motility findings are supported by the identification of a reduction in colocalization of AChRs with SV2-positive clusters on slow muscle fibers in 2 dpf fish, thus showing an increase in the number of AChRs that have not been contacted by a motor axon. We also identified an overall reduction in the number of SV2-positive clusters, which may be indicative of a defect or delay in development of the motor nervous system. Previous studies have commented on USH2A presence on the basement membranes of perineurium nerve fibers (Pearsall et al. 2002) (Schwaller et al. 2021), however, further functional studies will be required to determine the precise localization of the defect and whether loss of USH2a alone can impact NMJ signaling or whether co-occurrence with CHRNE CMS is required. Further functional work is also required to ascertain the importance of other potential modifiers identified in this study.

Our work represents a thorough study of a narrow population showing a differential accumulation of damaging mutations in patients with CMS who have varying phenotypic severities, building on the initial impact of CHRNE mutations on the NMJ. It is important to remark that CMS is of particular interest among rare diseases, since drugs that influence neuromuscular transmission can produce clear improvements in the affected patients (Engel 2007). In this sense, identifying meaningful molecular relationships between gene variants allow us to gain insight into the disease mechanisms through a multiplex biomedical framework, paving the way for a whole new set of computational approximations for rare disease research.

Acknowledgments

The authors acknowledge the donors and families, Daniel Rico (Newcastle University) for his contribution in early stages of the project, Anaïs Baudot (Aix Marseille Université and Barcelona Supercomputing Center) for her careful revision of the manuscript, Miguel Vázquez (Barcelona Supercomputing Center) for advising about Rbbt analysis, Jon Sánchez-Valle (Barcelona Supercomputing Center) and Núria Olvera (Barcelona Supercomputing Center and IDIBAPS) for the insightful discussions.

Funding

The NeurOmics and RD-Connect projects have been funded by the European Union's Seventh Framework Programme for research, technological development and demonstration under grant agreements no 2012-305121 and 2012-305444. I.N.C. was supported by a grant for pre-doctoral contracts for the training of doctors (Project ID: SEV-2015-0493-18-2) (Grant ID: PRE2018-083662) from the Spanish Ministry for Science, Innovation and Universities. E.O. was supported by an AFM-Téléthon postdoctoral fellowship for the duration of this work. H.L. receives support from the Canadian Institutes of Health Research (Foundation Grant FDN-167281), the Canadian Institutes of Health Research and Muscular Dystrophy Canada (Network Catalyst Grant for NMD4C), the Canada Foundation for Innovation (CFI-JELF 38412), and the Canada Research Chairs program (Canada Research Chair in Neuromuscular Genomics and Health, 950-232279). V.G. was a research fellow of the Alexander von Humboldt Foundation. D.C. was supported by the European Commission's Horizon 2020 Program, H2020-SC1-DTH-2018- 1, "iPC - individualizedPaediatricCure" (ref. 826121).

Author contributions

T.C., I.T. and V.G. collected and processed the biopsies; H.L. and R.T. coordinated data sharing; A.T., P.A.C.T., S.B. and S.C. coordinated and performed the omics data analysis with Y.A., S.L., M.R. and M.B.; E.O. performed the experiments in zebrafish; D.C. and A.V. coordinated the multilayer network analysis performed by I.N.C. All authors contributed to the writing and revising of the manuscript.

Ethics approval

This study was approved by the Ethics committee of Sofia Medical University (protocol 4/15-April-2013).

Competing Interest

None declared.

Methods

WGS and RNA-seq

Whole genome sequencing (WGS) data have been obtained from blood using the Illumina TruSeq PCR-free library preparation kit. Sample sequencing was performed with the HiSeqX sequencing platform (HiseqX v1 or v2 SBS kit, 2x150 cycles), with an average mean depth coverage $\geq$ 30X. Samples have been analyzed using the RD-Connect specific pipeline: BWA-mem for alignment; Picard for duplicate marking and GATK 3.6.0 for variant calling. RNA sequencing (RNA-seq) data have been obtained from fibroblasts, using Illumina TruSeq RNA Library Preparation Kit v2, sequencing with an average of 60M reads per sample (paired-end 2X125 cycles). Data has been processed with the following pipeline (Laurie et al. 2016): STAR 2.35a for alignment, RSEM 1.3.0 for quantification, and GATK 3.6.0 for variant calling. All analyses have been performed using the human genome GRCh37d5 as reference.

Copy number variants

Copy Number Variants (CNVs) have been extracted using ClinCNV (https://github.com/imgag/ClinCNV) by employing a set of Eastern European samples as a background control group. Out of the 569 autosomal CNVs we selected as potential candidates the CNVs of the following types that overlapped with protein-coding genes: 1) whole gene gains or losses, and 2) partial losses (deletions overlapping with exons but not with the whole gene). The list of potential candidates included 55 CNVs that created a total of 82 whole gene gains or losses and 28 partial losses.

Compound heterozygous variants

Compound heterozygous variants have been obtained by phasing the WGS variant calls with the RNA-seq aligned BAM files using phASER (Castel et al. 2016). At first, variants are imputed using Sanger Imputation Service with EAGLE2 pre-phasing step (Durbin 2014). PhASER is then applied to extend phased regions to gene-wide haplotypes. By accurately reflecting the muscle transcriptome, fibroblasts have been previously proved to be excellent and minimally invasive diagnostic tools for rare neuromuscular diseases (Gonorazky et al. 2019). We then annotated variants with eDiVA tool (www.ediva.crg.es) (Bosio et al. 2019), and removed all mutations with Genome Aggregation Database (gnomAD) (Lek et al. 2016) that show allele frequency > 3% globally, all variants outsde exonic and splicing regions using Ensembl annotation, all synonymous mutations, and all variants with read depth (coverage) smaller than 8. Afterwards we selected all genes with at least two hits on different alleles as genes affected by damaging compound heterozygous variants. Each sample has been processed individually throughout the whole process.

Monolayer community detection

We performed a network community detection analysis using the Louvain clustering algorithm (Blondel et al. 2008) implemented in R package igraph (https://igraph.org/) with default parameters. We carried out the analysis using three (monolayer) networks, obtained

from Reactome database (Fabregat et al. 2018), from the Recon3D Virtual Metabolic Human
database (Brunk et al. 2018) (both downloaded in May 2018), and from the Integrated
Interaction Database (IID) (Kotlyar et al. 2016) (downloaded in October 2018). The first
network consists of 10,618 nodes (genes) and 875,436 edges, representing shared pathways
between genes. The second network consists of 1,863 nodes (genes) and 902,188 edges,
representing shared reaction metabolites between genes. The third network consists of 18,018
nodes (genes) and 947,606 edges, representing aggregated protein-protein interactions from
all tissues. All gene identifiers of each network were converted to NCBI Entrez gene
identifiers using R packages AnnotationDbi v1.44.0 and org.Hs.eg.db v3.7.0
(http://bioconductor.org/). After detecting the community structure from each layer
independently, we retrieved the community membership of the genes of interest, henceforth
called "CMS linked genes", i.e. known CMS causal genes, and severe and not-severe
compound heterozygous variants and CNVs. We then defined a community similarity
measure as Jaccard Index, i.e. the number of shared genes of interest between the
communities divided by the sum of the total number of genes of each community.

Multilayer community detection
We constructed a multilayer gene network composed of the three monolayer networks
described in the previous section (Reactome, Virtual Metabolic Human and Integrated
Interaction Database). Each of these three networks represents one layer of the multilayer
network and, in general, three facets of fundamental molecular processes in the cell (**Suppl.**
Figure 11). The multilayer community detection analysis was performed by using MolTi
software (Didier, Brun, and Baudot 2015), which adapts the Louvain clustering algorithm
with modularity maximization to multilayer networks. The algorithm is parametrized by the
resolution (γ): the higher the value of γ, the smaller the size of the detected multilayer
communities. By varying the resolution parameter γ it is possible to uncover the modular
structure of network communities (Fortunato and Barthelemy 2007). By exploring a wide
range of resolution parameter values, we identified $\gamma=4$ (727 communities, each one
composed of 26.46 genes on average) as an extreme value before both size and number of the
detected multilayer communities stabilize (**Suppl. Figure 12**). The most dramatic changes in
number and composition of detected communities are observed in the resolution parameter
interval $\gamma \in (0,4]$. We, therefore, used this parameter interval to test the hypothesis that
disease-related genes consistently appear in the same multilayer communities, as well as to
identify modules containing CMS linked genes within them. In this analysis, we define a
module as a group of CMS linked genes that are systematically found to be part of the same
multilayer community while increasing the resolution parameter (see Supplementary
Information "Multilayer community detection analysis").

Additional analyses and code availability
We retrieved known CMS causal genes from the GeneTable of Neuromuscular Disorders
(http://www.musclegenetable.fr , version November 2018) (Bonne, Rivier, and Hamroun
2017). Segregation analysis of WGS data has been performed using Rbbt (Vázquez et al.

2010). DisGeNET database (Piñero et al. 2017) was downloaded in November 2018. The
association between CMS severity, demographic factors and clinical tests was assessed with a
two-tailed Fisher's test using R statistical environment (www.R-project.org). Networks were
rendered with Cytoscape (Saito et al. 2012). We used VCFtools (Danecek et al. 2011) to
compute familial relatedness Ω among patients, scaled to $-\log_2(2\Omega)$. We used Enrichr (E. Y.
Chen et al. 2013) for the functional enrichment analysis of the gene lists under study. We
used Ensembl Variant Effect Predictor (VEP) (McLaren et al. 2016) to assess the impact of
the compound heterozygous variants in the genes of the severe-specific largest module.
Expression levels in tissues of interest (GTEx and Illumina Body Map) were retrieved from
EBI Expression Atlas (www.ebi.ac.uk/) by filtering with the following keywords: 'nerve',
'muscle cell', 'fibroblast' and 'nervous system' (0.5 TPM default cutoff). We used
Expression Atlas expression level categories: low (0.5 to 10 TPM), medium (11 to 1000
TPM), and high (more than 1000 TPM) (Petryszak et al. 2016). Synaptic localization was
retrieved from the UniProt database (https://www.uniprot.org/). All code from the original
analysis is available for reproducibility purposes at: https://github.com/ikernunezca/CMS.
The analysis of multilayer community communties can also be performed using CmmD
(Nuñez-Carpintero et al., 2021) (https://github.com/ikernunezca/CmmD) with parameters:
resolution_start: 0, resolution_end: 4, interval: 0.5 and the CMS linked genes as nodelist.

Zebrafish morpholino injections
Zebrafish have one orthologue of human *USH2a: ush2a*, as identified using the UCSC
database (http://genome.ucsc.edu/, GRCz11/danRer11 assembly). We confirmed that *ush2a* is
expressed throughout the first 5 days post fertilization (dpf). Gene Tools LLC (USA) then
designed and synthesized an antisense morpholino oligonucleotide (MO) targeting the splice
donor site of exon 3/intron 3 of *ush2a* (5'-3' GAGAAATGCTGCTCACCTGTAGAGC,
ENSDART00000086201.5). We also obtained a control MO that targets a human beta-globin
mutation (5'-3' CCTCTTACCTCAGTTACAATTTATA). MOs were diluted to 2 ng/nl in
Danieau buffer (58 mM NaCl, 5 mM HEPES, 0.7 mM KCl, 0.6 mM $Ca(NO_3)_2$, 0.4 mM
$MgSO_4$; pH 7.6) and supplemented with 1% phenol red, before being injected into the yolk-
sac of 1-cell stage embryos. A range of doses between 6 and 18 ng per 1-cell stage embryo
were trialed for success in reducing *ush2a* expression and producing a measurable phenotypic
change. A dose of 18 ng per 1-cell stage embryo was selected for behavioral and
morphological analysis, as survival was not significantly affected for any dose tested.
Embryos were maintained at 28.5°C in blue water (system water with 0.1 µg/ml Methylene
Blue) for up to 5 dpf and survival recorded daily. At 2 dpf zebrafish were imaged using a
Leica EZ4 W stereomicroscope and eye size and length measured using Fiji (ImageJ).

Chorion movement analysis in zebrafish
At 1 dpf (24 hours post fertilization), zebrafish were recorded in their chorions for 1 minute
at 30 frames per second using a Leica EZ4 W stereomicroscope. Videos were analyzed using
DanioScope software (Noldus Information Technology Inc., Leesburg, VA) to automatically
assess duration of bursts and burst count/minute (bursts are full rotations performed by the

zebrafish within the chorion).

Touch response analysis

At 2 dpf, a touch response assay was performed as previously described (O'Connor et al.
2018). Only fish with a normal phenotype were used for movement analysis. Briefly, fish that
had not hatched from the chorion were enzymatically dechorionated with pronase (1 mg/ml,
Sigma) for 10 min in blue water, followed by 3x washes in blue water. An individual fish was
placed in a petri dish containing blue water and a Sony RX0 II (DSC-RX0M2) camera was
placed 20 cm above the petri dish. A ruler with 1 mm markings was used as a scale for
recordings. A gel loading pipette tip was used to touch the zebrafish on the back of the head
and the response recorded. Videos were imported into Fiji ImageJ (Schindelin et al. 2012) as
FFmpeg movies and movements analyzed using the Trackmate plugin (Tinevez et al. 2017).
Values for average speed were exported and used to derive initial acceleration.

RNA isolation, cDNA synthesis and RT-PCR in zebrafish

RNA was isolated from pools of around 20 2 dpf zebrafish (control MO and *ush2a* MO-
injected) following removal of chorions with pronase (Streptomyces griseus, Roche,1 mg/ml
in blue water). Zebrafish were washed 3 times with blue water, euthanized with a 1:1 ratio of
fresh system water:4 mg/ml tricaine methanesulfonate (Sigma). Fish were homogenized in
RLT buffer (RNeasy mini kit, Qiagen) using 5 mm stainless steel beads with a TissueLyser II
(Qiagen) at 25 Hz for 2 mins. RNA was then isolated following the RNeasy kit
manufacturer's instructions, including on-column DNase digestion. RNA was measured
using a Nanodrop ND-1000 and 1 µg used for cDNA synthesis according to manufacturer's
instructions (5X All-In-One RT MasterMix, abm). Reverse-transcriptase PCR (RT-PCR) was
performed to check for *ush2a* gene expression and knockdown success in MO-treated
embryos, using MyTaq™ DNA Polymerase (Meridian Bioscience) and primers as follows:
eef1a1l1 forward 5'-CTGGAGGCCAGCTCAAACATGG-3', reverse 5'-
CTTGCTGTCTCCAGCCACATTAC-3' and *ush2a* forward 5'-
CTGGGCACACTTGGCTCTAC -3', reverse 5'-TTCTTCAATCTCCCTGTTGGTT-3'.

Immunofluorescent staining, imaging and analysis of zebrafish neuromuscular junctions
and muscle fibers

Whole mount staining of 2 dpf zebrafish NMJs was performed as previously described
(O'Connor et al. 2019). Briefly, a mouse anti-synaptic vesicle protein 2 (SV2) antibody was
used to visualize motor neurons (1:200, AB2315387, Developmental Studies Hybridoma
Bank) and Alexa Fluor 488-α-bungarotoxin conjugate (1:1000, B13422, Invitrogen) was used
for visualizing acetylcholine receptors (AChRs). Phalloidin-iFluor 594 was used to visualize
filamentous actin within muscle fibers (1:1000, ab176757). Z-stack images encompassing the
depth of the midsection of the zebrafish tail were obtained using a 20× air objective on an
LSM800 confocal microscope. Analysis of NMJ structure was performed as previously
described (O'Connor et al. 2019), using Fiji (ImageJ, Madison, WI, USA). The number of
SV2-positive and α-bungarotoxin-positive clusters per 100 μm^2 were measured. Co-
localization analysis between SV2 and α-bungarotoxin was performed on maximum intensity

projections using the 'JACoP' Fiji plugin (Bolte & Cordelières, 2006). Briefly, each
fluorophore was subject to manual thresholding to remove background, and the Mander's
correlation coefficient calculated to give a value between 0 and 1, reflecting the degree of co-
occurrence of signals between both SV2 and α-bungarotoxin, and α-bungarotoxin with SV2.
For phalloidin-stained fish, average myotome size was measured, and degree of fiber
dispersion quantified using the directionality plugin. Data was collected from at least 4
myotomes per fish.

Statistics for zebrafish experiments

805
Statistical analysis was performed using GraphPad Prism software (v9.3.0). Outliers were
removed from data using the ROUT method (Q = 1 %). Cleaned data was tested for normal
distribution then depending on outcome either a nonparametric Mann-Whitney test or
parametric unpaired t-test were applied for behavioral studies and degree of dispersion. For
NMJ morphology experiments in which 4+ myotomes (technical replicates) per fish
(biological replicates) were analyzed, data was assessed for significance using a nested t-test
to avoid pseudoreplication. Statistical significance was taken as $p < 0.05$, degrees of freedom
(df) and t-value are given for all parametric tests, and n numbers listed in the results section.
Survival analysis was performed using the log-rank test comparing WT to each other
condition, and threshold for significance was corrected for multiple comparisons using the
Bonferroni method ($p < 0.006$). Zebrafish studies were blinded before image/video
acquisition and unblinded following analysis.

References

Abicht, A., R. Stucka, V. Karcagi, A. Herczegfalvi, R. Horváth, W. Mortier, U. Schara, et al.
1999. "A Common Mutation (Epsilon1267delG) in Congenital Myasthenic Patients of
Gypsy Ethnic Origin." *Neurology* 53 (7): 1564–69.
https://doi.org/10.1212/wnl.53.7.1564.
Abicht, Angela, Juliane Müller, and Hanns Lochmüller. 1993. "Congenital Myasthenic
Syndromes." In *GeneReviews®*, edited by Margaret P. Adam, Holly H. Ardinger,
Roberta A. Pagon, Stephanie E. Wallace, Lora JH Bean, Karen Stephens, and Anne
Amemiya. Seattle (WA): University of Washington, Seattle.
http://www.ncbi.nlm.nih.gov/books/NBK1168/.
Amin, Mutaz, Yousuf Bakhit, Mahmoud Koko, Mohamed Osama Mirgahni Ibrahim, M. A.
Salih, Muntaser Ibrahim, and Osheik A. Seidi. 2019. "Rare Variant in LAMA2 Gene
Causing Congenital Muscular Dystrophy in a Sudanese Family. A Case Report." *Acta*
Myologica: Myopathies and Cardiomyopathies: Official Journal of the
Mediterranean Society of Myology 38 (1): 21–24.
Andreose, J. S., C. Sala, and G. Fumagalli. 1994. "Immunolocalization of Chromogranin B,
Secretogranin II, Calcitonin Gene-Related Peptide and Substance P at Developing and
Adult Neuromuscular Synapses." *Neuroscience Letters* 174 (2): 177–80.
Arikawa-Hirasawa, Eri, William R. Wilcox, Alexander H. Le, Neil Silverman, Prasanthi
Govindraj, John R. Hassell, and Yoshihiko Yamada. "Dyssegmental Dysplasia,
Silverman-Handmaker Type, Is Caused by Functional Null Mutations of the Perlecan
Gene." *Nature Genetics* 27, no. 4 (April 2001): 431–34.
https://doi.org/10.1038/86941.
Arikawa-Hirasawa, Eri, Alexander H. Le, Ichizo Nishino, Ikuya Nonaka, Nicola C. Ho, Clair
A. Francomano, Prasanthi Govindraj, et al. 2002. "Structural and Functional
Mutations of the Perlecan Gene Cause Schwartz-Jampel Syndrome, with Myotonic
Myopathy and Chondrodysplasia." *American Journal of Human Genetics* 70 (5):
1368–75. https://doi.org/10.1086/340390.
Astigarraga, Sergio, Kerstin Hofmeyer, Reza Farajian, and Jessica E. Treisman. 2010. "Three
Drosophila Liprins Interact to Control Synapse Formation." *The Journal of*
Neuroscience: The Official Journal of the Society for Neuroscience 30 (46): 15358–
68. https://doi.org/10.1523/JNEUROSCI.1862-10.2010.
Austin-Tse, Christina A., Diana L. Mandelker, Andrea M. Oza, Heather Mason-Suares, Heidi
L. Rehm, and Sami S. Amr. 2018. "Analysis of Intragenic USH2A Copy Number
Variation Unveils Broad Spectrum of Unique and Recurrent Variants." *European*
Journal of Medical Genetics 61 (10): 621–26.
https://doi.org/10.1016/j.ejmg.2018.04.006.
Barik, Arnab, Yisheng Lu, Anupama Sathyamurthy, Andrew Bowman, Chengyong Shen, Lei
Li, Wen-cheng Xiong, and Lin Mei. 2014. "LRP4 Is Critical for Neuromuscular
Junction Maintenance." *The Journal of Neuroscience: The Official Journal of the*
Society for Neuroscience 34 (42): 13892–905.

https://doi.org/10.1523/JNEUROSCI.1733-14.2014.

Beeson, David. 2016. "Congenital Myasthenic Syndromes: Recent Advances." *Current Opinion in Neurology* 29 (5): 565–71. https://doi.org/10.1097/WCO.0000000000000370.

Bernadzki, Krzysztof M., Marta Gawor, Marcin Pęziński, Paula Mazurek, Paweł Niewiadomski, Maria J. Rędowicz, and Tomasz J. Prószyński. 2017. "Liprin-α-1 Is a Novel Component of the Murine Neuromuscular Junction and Is Involved in the Organization of the Postsynaptic Machinery." *Scientific Reports* 7 (1): 9116. https://doi.org/10.1038/s41598-017-09590-7.

Bertini, Enrico, Adele D'Amico, Francesca Gualandi, and Stefania Petrini. 2011. "Congenital Muscular Dystrophies: A Brief Review." *Seminars in Pediatric Neurology* 18 (4): 277–88. https://doi.org/10.1016/j.spen.2011.10.010.

Bevilacqua, Jorge A., Marian Lara, Jorge Díaz, Mario Campero, Jessica Vázquez, and Ricardo A. Maselli. 2017. "Congenital Myasthenic Syndrome Due to DOK7 Mutations in a Family from Chile." *European Journal of Translational Myology* 27 (3): 6832. https://doi.org/10.4081/ejtm.2017.6832.

Blondel, Vincent D., Jean-Loup Guillaume, Renaud Lambiotte, and Etienne Lefebvre. 2008. "Fast Unfolding of Communities in Large Networks." *Journal of Statistical Mechanics: Theory and Experiment* 2008 (10): P10008. https://doi.org/10.1088/17425468/2008/10/P10008.

Bolte, Sussane, and Fabrice P. Cordelières. "A Guided Tour into Subcellular Colocalization Analysis in Light Microscopy." *Journal of Microscopy* 224, no. 3 (2006): 213–32. https://doi.org/10.1111/j.1365-2818.2006.01706.x.

Bonne, Gisèle, François Rivier, and Dalil Hamroun. 2017. "The 2018 Version of the Gene Table of Monogenic Neuromuscular Disorders (Nuclear Genome)." *Neuromuscular Disorders: NMD* 27 (12): 1152–83. https://doi.org/10.1016/j.nmd.2017.10.005.

Bönnemann, Carsten G., Ching H. Wang, Susana Quijano-Roy, Nicolas Deconinck, Enrico Bertini, Ana Ferreiro, Francesco Muntoni, et al. 2014. "Diagnostic Approach to the Congenital Muscular Dystrophies." *Neuromuscular Disorders: NMD* 24 (4): 289–311. https://doi.org/10.1016/j.nmd.2013.12.011.

Bork, P., and R. F. Doolittle. 1992. "Proposed Acquisition of an Animal Protein Domain by Bacteria." *Proceedings of the National Academy of Sciences of the United States of America* 89 (19): 8990–94.

Bosio, Mattia, Oliver Drechsel, Rubayte Rahman, Francesc Muyas, Raquel Rabionet, Daniela Bezdan, Laura Domenech Salgado, et al. 2019. "EDiVA-Classification and Prioritization of Pathogenic Variants for Clinical Diagnostics." *Human Mutation*, April. https://doi.org/10.1002/humu.23772.

Boycott, Kym M., Megan R. Vanstone, Dennis E. Bulman, and Alex E. MacKenzie. 2013. "Rare-Disease Genetics in the Era of next-Generation Sequencing: Discovery to Translation." Nature Reviews Genetics 14 (10): 681–91. https://doi.org/10.1038/nrg3555.

Brunk, Elizabeth, Swagatika Sahoo, Daniel C. Zielinski, Ali Altunkaya, Andreas Dräger,

Nathan Mih, Francesco Gatto, et al. 2018. "Recon3D Enables a Three-Dimensional View of Gene Variation in Human Metabolism." *Nature Biotechnology* 36 (3): 272–81. https://doi.org/10.1038/nbt.4072.

Buphamalai, Pisanu, Tomislav Kokotovic, Vanja Nagy, and Jörg Menche. 2021. 'Network Analysis Reveals Rare Disease Signatures across Multiple Levels of Biological Organization'. Nature Communications 12 (1): 6306. https://doi.org/10.1038/s41467-021-26674-1.

Cantini, Laura, Enzo Medico, Santo Fortunato, and Michele Caselle. 2015. "Detection of Gene Communities in Multi-Networks Reveals Cancer Drivers." *Scientific Reports* 5 (December): 17386. https://doi.org/10.1038/srep17386.

Castel, Stephane E., Pejman Mohammadi, Wendy K. Chung, Yufeng Shen, and Tuuli Lappalainen. 2016. "Rare Variant Phasing and Haplotypic Expression from RNA Sequencing with PhASER." *Nature Communications* 7: 12817. https://doi.org/10.1038/ncomms12817.

Castro-Giner, Francesc, Peter Ratcliffe, and Ian Tomlinson. 2015. "The Mini-Driver Model of Polygenic Cancer Evolution." *Nature Reviews. Cancer* 15 (11): 680–85. https://doi.org/10.1038/nrc3999.

Chen, Edward Y., Christopher M. Tan, Yan Kou, Qiaonan Duan, Zichen Wang, Gabriela Vaz Meirelles, Neil R. Clark, and Avi Ma'ayan. 2013. "Enrichr: Interactive and Collaborative HTML5 Gene List Enrichment Analysis Tool." *BMC Bioinformatics* 14 (April): 128. https://doi.org/10.1186/1471-2105-14-128.

Chen, Yuewen, Jinying Xu, Xiaopu Zhou, Saijuan Liu, Yulin Zhang, Shuangshuang Ma, Amy K. Y. Fu, Nancy Y. Ip, and Yu Chen. 2019. "Changes of Protein Phosphorylation Are Associated with Synaptic Functions during the Early Stage of Alzheimer's Disease." *ACS Chemical Neuroscience* 10 (9): 3986–96. https://doi.org/10.1021/acschemneuro.9b00190.

Clausen, Lisa, Judith Cossins, and David Beeson. 2018. "Beta-2 Adrenergic Receptor Agonists Enhance AChR Clustering in C2C12 Myotubes: Implications for Therapy of Myasthenic Disorders." *Journal of Neuromuscular Diseases* 5 (2): 231–40. https://doi.org/10.3233/JND-170293.

Dagur, Pradeep K., and J. Philip McCoy. 2015. "Endothelial-Binding, Proinflammatory T Cells Identified by MCAM (CD146) Expression: Characterization and Role in Human Autoimmune Diseases." *Autoimmunity Reviews* 14 (5): 415–22. https://doi.org/10.1016/j.autrev.2015.01.003.

Danecek, P., A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E. Handsaker, et al. 2011. "The Variant Call Format and VCFtools." *Bioinformatics* 27 (15): 2156–58. https://doi.org/10.1093/bioinformatics/btr330.

Didier, Gilles, Christine Brun, and Anaïs Baudot. 2015. "Identifying Communities from Multiplex Biological Networks." *PeerJ* 3: e1525. https://doi.org/10.7717/peerj.1525.

Dimova, Ivanka, and Ivo Kremensky. 2018. "LAMA2 Congenital Muscle Dystrophy: A Novel Pathogenic Mutation in Bulgarian Patient." *Case Reports in Genetics* 2018: 3028145. https://doi.org/10.1155/2018/3028145.

Downes, Gerald B., and Michael Granato. 2006. "Supraspinal Input Is Dispensable to
Generate Glycine-Mediated Locomotive Behaviors in the Zebrafish Embryo."
Journal of Neurobiology 66, no. 5: 437–51. https://doi.org/10.1002/neu.20226.
Durbin, Richard. 2014. "Efficient Haplotype Matching and Storage Using the Positional
Burrows-Wheeler Transform (PBWT)." *Bioinformatics (Oxford, England)* 30 (9):
1266–72. https://doi.org/10.1093/bioinformatics/btu014.
Eklund, L., J. Piuhola, J. Komulainen, R. Sormunen, C. Ongvarrasopone, R. Fássler, A.
Muona, et al. 2001. "Lack of Type XV Collagen Causes a Skeletal Myopathy and
Cardiovascular Defects in Mice." *Proceedings of the National Academy of Sciences of*
the United States of America 98 (3): 1194–99.
https://doi.org/10.1073/pnas.031444798.
Engel, Andrew G. 2007. "The Therapy of Congenital Myasthenic Syndromes."
Neurotherapeutics: The Journal of the American Society for Experimental
NeuroTherapeutics 4 (2): 252–57. https://doi.org/10.1016/j.nurt.2007.01.001.
Engel, Andrew G., Xin-Ming Shen, Duygu Selcen, and Steven M. Sine. 2015. "Congenital
Myasthenic Syndromes: Pathogenesis, Diagnosis, and Treatment." *The Lancet.*
Neurology 14 (4): 420–34. https://doi.org/10.1016/S1474-4422(14)70201-7.
Estephan, Eduardo de Paula, Antonio Alberto Zambon, Paulo Eurípedes Marchiori, André
Macedo Serafim da Silva, Vitor Marques Caldas, Cristiane Araújo Martins Moreno,
Umbertina Conti Reed, et al. 2018. "Clinical Variability of Early-Onset Congenital
Myasthenic Syndrome Due to Biallelic RAPSN Mutations in Brazil." *Neuromuscular*
Disorders 28 (11): 961–64. https://doi.org/10.1016/j.nmd.2018.08.007.
Fabregat, Antonio, Steven Jupe, Lisa Matthews, Konstantinos Sidiropoulos, Marc Gillespie,
Phani Garapati, Robin Haw, et al. 2018. "The Reactome Pathway Knowledgebase."
Nucleic Acids Research 46 (D1): D649–55. https://doi.org/10.1093/nar/gkx1132.
Fareed, Mohd, and Mohammad Afzal. 2017. "Genetics of Consanguinity and Inbreeding in
Health and Disease." *Annals of Human Biology* 44 (2): 99–107.
https://doi.org/10.1080/03014460.2016.1265148.
Finsterer, Josef. 2019. "Congenital Myasthenic Syndromes." *Orphanet Journal of Rare*
Diseases 14 (1): 57. https://doi.org/10.1186/s13023-019-1025-5.
Forrest, Katharine, Jemima E. Mellerio, Stephanie Robb, Patricia J. C. Dopping-Hepenstal,
John A. McGrath, Lu Liu, Stefan J. A. Buk, Safa Al-Sarraj, Elizabeth Wraige, and
Heinz Jungbluth. 2010. "Congenital Muscular Dystrophy, Myasthenic Symptoms and
Epidermolysis Bullosa Simplex (EBS) Associated with Mutations in the PLEC1 Gene
Encoding Plectin." *Neuromuscular Disorders: NMD* 20 (11): 709–11.
https://doi.org/10.1016/j.nmd.2010.06.003.
Fortunato, S., and M. Barthelemy. 2007. "Resolution Limit in Community Detection."
Proceedings of the National Academy of Sciences 104 (1): 36–41.
https://doi.org/10.1073/pnas.0605965104.
Garg, Divyani, and Vinay Goyal. "Positive Response to Inhaled Salbutamol in Congenital
Myasthenic Syndrome Due to CHRNE Mutation." Muscle & Nerve n/a, no. n/a.
Accessed June 2, 2022. https://doi.org/10.1002/mus.27563.

Goh, K.-I., M. E. Cusick, D. Valle, B. Childs, M. Vidal, and A.-L. Barabasi. 2007. "The
Human Disease Network." *Proceedings of the National Academy of Sciences* 104
(21): 8685–90. https://doi.org/10.1073/pnas.0701361104.
Gonorazky, Hernan D., Sergey Naumenko, Arun K. Ramani, Viswateja Nelakuditi, Pouria
Mashouri, Peiqui Wang, Dennis Kao, et al. 2019. "Expanding the Boundaries of RNA
Sequencing as a Diagnostic Tool for Rare Mendelian Disease." *American Journal of*
Human Genetics 104 (3): 466–83. https://doi.org/10.1016/j.ajhg.2019.01.012.
Gosak, Marko, Rene Markovič, Jurij Dolenšek, Marjan Slak Rupnik, Marko Marhl, Andraž
Stožer, and Matjaž Perc. 2018. "Network Science of Biological Systems at Different
Scales: A Review." *Physics of Life Reviews* 24: 118–35.
https://doi.org/10.1016/j.plrev.2017.11.003.
Gowen, B. B., T. K. Borg, A. Ghaffar, and E. P. Mayer. 2000. "Selective Adhesion of
Macrophages to Denatured Forms of Type I Collagen Is Mediated by Scavenger
Receptors." *Matrix Biology: Journal of the International Society for Matrix Biology*
19 (1): 61–71.
Grevengoed, Trisha J., Eric L. Klett, and Rosalind A. Coleman. 2014. "Acyl-CoA
Metabolism and Partitioning." *Annual Review of Nutrition* 34: 1–30.
https://doi.org/10.1146/annurev-nutr-071813-105541.
Guillon, Emilie, Sandrine Bretaud, and Florence Ruggiero. 2016. "Slow Muscle Precursors
Lay Down a Collagen XV Matrix Fingerprint to Guide Motor Axon Navigation." *The*
Journal of Neuroscience: The Official Journal of the Society for Neuroscience 36 (9):
2663–76. https://doi.org/10.1523/JNEUROSCI.2847-15.2016.
Halu, Arda, Manlio De Domenico, Alex Arenas, and Amitabh Sharma. 2017. "The Multiplex
Network of Human Diseases." *BioRxiv*. https://doi.org/10.1101/100370.
Han, Shanshan, Xiliang Liu, Shanglun Xie, Meng Gao, Fei Liu, Shanshan Yu, Peng Sun, et
al. 2018. "Knockout of Ush2a Gene in Zebrafish Causes Hearing Impairment and
Late Onset Rod-Cone Dystrophy." *Human Genetics* 137, no. 10: 779–94.
https://doi.org/10.1007/s00439-018-1936-6.
Hull, Sarah, Gavin Arno, Anthony G. Robson, Suzanne Broadgate, Vincent Plagnol, Martin
McKibbin, Stephanie Halford, et al. 2016. "Characterization of CDH3-Related
Congenital Hypotrichosis With Juvenile Macular Dystrophy." *JAMA Ophthalmology*
134 (9): 992–1000. https://doi.org/10.1001/jamaophthalmol.2016.2089.
Huzé, Caroline, Stéphanie Bauché, Pascale Richard, Frédéric Chevessier, Evelyne Goillot,
Karen Gaudon, Asma Ben Ammar, et al. 2009. "Identification of an Agrin Mutation
That Causes Congenital Myasthenia and Affects Synapse Function." *American*
Journal of Human Genetics 85 (2): 155–67.
https://doi.org/10.1016/j.ajhg.2009.06.015.
Iozzo, Renato V., and Liliana Schaefer. 2015. "Proteoglycan Form and Function: A
Comprehensive Nomenclature of Proteoglycans." *Matrix Biology: Journal of the*
International Society for Matrix Biology 42 (March): 11–55.
https://doi.org/10.1016/j.matbio.2015.02.003.
Ito, Mikako, and Kinji Ohno. 2018. "Protein-Anchoring Therapy to Target Extracellular

Matrix Proteins to Their Physiological Destinations." *Matrix Biology: Journal of the*
International Society for Matrix Biology 68–69: 628–36.
https://doi.org/10.1016/j.matbio.2018.02.014.

Johansson, Malin E. V., Kristina A. Thomsson, and Gunnar C. Hansson. 2009. "Proteomic
Analyses of the Two Mucus Layers of the Colon Barrier Reveal That Their Main
Component, the Muc2 Mucin, Is Strongly Bound to the Fcgbp Protein." *Journal of*
Proteome Research 8 (7): 3549–57. https://doi.org/10.1021/pr9002504.

Kaneko-Goto, Tomomi, Yuki Sato, Sayako Katada, Emi Kinameri, Sei-ichi Yoshihara,
Atsushi Nishiyori, Mitsuhiro Kimura, et al. 2013. "Goofy Coordinates the Acuity of
Olfactory Signaling." *The Journal of Neuroscience: The Official Journal of the*
Society for Neuroscience 33 (32): 12987–96, 12996a.
https://doi.org/10.1523/JNEUROSCI.4948-12.2013.

Karczewski, Konrad J., and Michael P. Snyder. 2018. "Integrative Omics for Health and
Disease." Nature Reviews Genetics 19 (5): 299–310.
https://doi.org/10.1038/nrg.2018.4.

Karczewski, Konrad J., Laurent C. Francioli, Grace Tiao, Beryl B. Cummings, Jessica
Alföldi, Qingbo Wang, Ryan L. Collins, et al. "The Mutational Constraint Spectrum
Quantified from Variation in 141,456 Humans." *Nature* 581, no. 7809 (May 2020):
434–43. https://doi.org/10.1038/s41586-020-2308-7.

Keller, Kate E., Ying Ying Sun, Janice A. Vranka, Lauren Hayashi, and Ted S. Acott. 2012.
"Inhibition of Hyaluronan Synthesis Reduces Versican and Fibronectin Levels in
Trabecular Meshwork Cells." *PloS One* 7 (11): e48523.
https://doi.org/10.1371/journal.pone.0048523.

Kirschner, Janbernd, Ingrid Hausser, Yaqun Zou, Gudrun Schreiber, Hans-Jürgen Christen,
Susan C. Brown, Ingrun Anton-Lamprecht, Francesco Muntoni, Folker Hanefeld, and
Carsten G. Bönnemann. 2005. "Ullrich Congenital Muscular Dystrophy: Connective
Tissue Abnormalities in the Skin Support Overlap with Ehlers-Danlos Syndromes."
American Journal of Medical Genetics. Part A 132A (3): 296–301.
https://doi.org/10.1002/ajmg.a.30443.

Kivelä, Mikko, Alexandre Arenas, Marc Barthelemy, James P. Gleeson, Yamir Moreno, and
Mason A. Porter. 2014. "Multilayer Networks." *Journal of Complex Networks* 2 (3):
203–71. https://doi.org/10.1093/comnet/cnu016.

Kotlyar, Max, Chiara Pastrello, Nicholas Sheahan, and Igor Jurisica. 2016. "Integrated
Interactions Database: Tissue-Specific View of the Human and Model Organism
Interactomes." *Nucleic Acids Research* 44 (D1): D536-541.
https://doi.org/10.1093/nar/gkv1115.

Kousi, M., and N. Katsanis. 2015. "Genetic Modifiers and Oligogenic Inheritance." *Cold*
Spring Harbor Perspectives in Medicine 5 (6): a017145–a017145.
https://doi.org/10.1101/cshperspect.a017145.

Kraft-Sheleg, Ortal, Shelly Zaffryar-Eilot, Olga Genin, Wesal Yaseen, Sharon Soueid-
Baumgarten, Ofra Kessler, Tatyana Smolkin, et al. 2016. "Localized LoxL3-
Dependent Fibronectin Oxidation Regulates Myofiber Stretch and Integrin-Mediated

Adhesion.” *Developmental Cell* 36 (5): 550–61. https://doi.org/10.1016/j.devcel.2016.02.009.

Laskowski, M., and I. Kato. 1980. “Protein Inhibitors of Proteinases.” *Annual Review of Biochemistry* 49: 593–626. https://doi.org/10.1146/annurev.bi.49.070180.003113.

Laurie, Steve, Marcos Fernandez-Callejo, Santiago Marco-Sola, Jean-Remi Trotta, Jordi Camps, Alejandro Chacón, Antonio Espinosa, et al. 2016. “From Wet-Lab to Variations: Concordance and Speed of Bioinformatics Pipelines for Whole Genome and Whole Exome Sequencing.” *Human Mutation* 37, (12): 1263–71. https://doi.org/10.1002/humu.23114.

Lee, Manon, David Beeson, and Jacqueline Palace. 2018. “Therapeutic Strategies for Congenital Myasthenic Syndromes.” *Annals of the New York Academy of Sciences* 1412 (1): 129–36. https://doi.org/10.1111/nyas.13538.

Lek, Monkol, Konrad J. Karczewski, Eric V. Minikel, Kaitlin E. Samocha, Eric Banks, Timothy Fennell, Anne H. O’Donnell-Luria, et al. 2016. “Analysis of Protein-Coding Genetic Variation in 60,706 Humans.” *Nature* 536 (7616): 285–91. https://doi.org/10.1038/nature19057.

Li, Lei O., Trisha J. Grevengoed, David S. Paul, Olga Ilkayeva, Timothy R. Koves, Florencia Pascual, Christopher B. Newgard, Deborah M. Muoio, and Rosalind A. Coleman. 2015. “Compartmentalized Acyl-CoA Metabolism in Skeletal Muscle Regulates Systemic Glucose Homeostasis.” *Diabetes* 64 (1): 23–35. https://doi.org/10.2337/db13-1070.

Lochmüller, Hanns, Dorota M. Badowska, Rachel Thompson, Nine V. Knoers, Annemieke Aartsma-Rus, Ivo Gut, Libby Wood, et al. 2018. “RD-Connect, NeurOmics and EURenOmics: Collaborative European Initiative for Rare Diseases.” *European Journal of Human Genetics: EJHG* 26 (6): 778–85. https://doi.org/10.1038/s41431018-0115-5.

Løkken, Nicoline, Alfred Peter Born, Morten Duno, and John Vissing. 2015. “LAMA2-Related Myopathy: Frequency among Congenital and Limb-Girdle Muscular Dystrophies.” *Muscle & Nerve* 52 (4): 547–53. https://doi.org/10.1002/mus.24588.

Lord, Megan S., Fengying Tang, Jelena Rnjak-Kovacina, James G. W. Smith, James Melrose, and John M. Whitelock. 2018. “The Multifaceted Roles of Perlecan in Fibrosis.” *Matrix Biology: Journal of the International Society for Matrix Biology* 68–69: 150–66. https://doi.org/10.1016/j.matbio.2018.02.013.

McLaren, William, Laurent Gil, Sarah E. Hunt, Harpreet Singh Riat, Graham R. S. Ritchie, Anja Thormann, Paul Flicek, and Fiona Cunningham. 2016. “The Ensembl Variant Effect Predictor.” *Genome Biology* 17 (1): 122. https://doi.org/10.1186/s13059-016-0974-4.

McRae, Natasha, Leonard Forgan, Bryony McNeill, Alex Addinsall, Daniel McCulloch, Chris Van der Poel, and Nicole Stupka. 2017. “Glucocorticoids Improve Myogenic Differentiation In Vitro by Suppressing the Synthesis of Versican, a Transitional Matrix Protein Overexpressed in Dystrophic Skeletal Muscles.” *International Journal of Molecular Sciences* 18 (12). https://doi.org/10.3390/ijms18122629.

Menche, Jörg, Amitabh Sharma, Maksim Kitsak, Susan Dina Ghiassian, Marc Vidal, Joseph Loscalzo, and Albert-László Barabási. 2015. "Disease Networks. Uncovering Disease-Disease Relationships through the Incomplete Interactome." *Science (New York, N.Y.)* 347 (6224): 1257601. https://doi.org/10.1126/science.1257601.

Mitani, Aya A., and Sebastien Haneuse. 2020. "Small Data Challenges of Studying Rare Diseases." *JAMA Network Open* 3 (3): e201965. https://doi.org/10.1001/jamanetworkopen.2020.1965.

Mo, Michelle, Ha Thi Hoang, Stefan Schmidt, Robert B. Clark, and Barbara E. Ehrlich. 2013. "The Role of Chromogranin B in an Animal Model of Multiple Sclerosis." *Molecular and Cellular Neurosciences* 56 (September): 102–14. https://doi.org/10.1016/j.mcn.2013.04.003.

Müller, Juliane S., Angela Abicht, Hans-Jürgen Christen, Rolf Stucka, Ulrike Schara, Wilhelm Mortier, Angela Huebner, and Hanns Lochmüller. 2004. "A Newly Identified Chromosomal Microdeletion of the Rapsyn Gene Causes a Congenital Myasthenic Syndrome." *Neuromuscular Disorders: NMD* 14 (11): 744–49. https://doi.org/10.1016/j.nmd.2004.06.010.

Müller, Juliane S., Catherine D. Jepson, Steven H. Laval, Kate Bushby, Volker Straub, and Hanns Lochmüller. 2010. "Dok-7 Promotes Slow Muscle Integrity as Well as Neuromuscular Junction Formation in a Zebrafish Model of Congenital Myasthenic Syndromes." *Human Molecular Genetics* 19, no. 9: 1726–40. https://doi.org/10.1093/hmg/ddq049.

Muona, Anu, Lauri Eklund, Timo Väisänen, and Taina Pihlajaniemi. 2002. "Developmentally Regulated Expression of Type XV Collagen Correlates with Abnormalities in Col15a1(-/-) Mice." *Matrix Biology: Journal of the International Society for Matrix Biology* 21 (1): 89–102.

Nicole, Sophie, Amina Chaouch, Torberg Torbergsen, Stéphanie Bauché, Elodie de Bruyckere, Marie-Joséphine Fontenille, Morten A. Horn, et al. 2014. "Agrin Mutations Lead to a Congenital Myasthenic Syndrome with Distal Muscle Weakness and Atrophy." *Brain: A Journal of Neurology* 137 (Pt 9): 2429–43. https://doi.org/10.1093/brain/awu160.

Nilsson, Anna, Maria Fälth, Xiaoqun Zhang, Kim Kultima, Karl Sköld, Per Svenningsson, and Per E. Andrén. 2009. "Striatal Alterations of Secretogranin-1, Somatostatin, Prodynorphin, and Cholecystokinin Peptides in an Experimental Mouse Model of Parkinson Disease." *Molecular & Cellular Proteomics: MCP* 8 (5): 1094–1104. https://doi.org/10.1074/mcp.M800454-MCP200.

Núñez-Carpintero, Iker, Marianyela Petrizzelli, Andrei Zinovyev, Davide Cirillo and Alfonso Valencia. 2021. "The multilayer community structure of medulloblastoma." *iScience* 24. https://doi.org/10.1016/j.isci.2021.102365

O'Connor, Emily, Ana Töpf, René P Zahedi, Sally Spendiff, Daniel Cox, Andreas Roos and Hanns Lochmüller. 2018. "Clinical and research strategies for limb-girdle congenital myasthenic syndromes." *Annals of the New York Academy of Sciences* 1412, 102–112. https://doi.org/10.1111/nyas.13520

O'Connor, Emily, George Cairns, Sally Spendiff, David Burns, Stefan Hettwer, Armin Mäder, Juliane Müller, Rita Horvath, Clarke Slater, Andreas Roos, Hanns Lochmüller. 2019. "Modulation of Agrin and RhoA Pathways Ameliorates Movement Defects and Synapse Morphology in MYO9A-Depleted Zebrafish." *Cells* 8, no. 8, 848. https://doi.org/10.3390/cells8080848.

Ohkawara, Bisei, Macarena Cabrera-Serrano, Tomohiko Nakata, Margherita Milone, Nobuyuki Asai, Kenyu Ito, Mikako Ito, et al. 2014. "LRP4 Third β-Propeller Domain Mutations Cause Novel Congenital Myasthenia by Compromising Agrin-Mediated MuSK Signaling in a Position-Specific Manner." *Human Molecular Genetics* 23 (7): 1856–68. https://doi.org/10.1093/hmg/ddt578.

Ohno, K. 2003. "E-Box Mutations in the RAPSN Promoter Region in Eight Cases with Congenital Myasthenic Syndrome." *Human Molecular Genetics* 12 (7): 739–48. https://doi.org/10.1093/hmg/ddg089.

Pampalakis, Georgios, Konstantinos Mitropoulos, Georgia Xeromerisiou, Efthymios Dardiotis, Georgia Deretzi, Maria Anagnostouli, Theodora Katsila, Michail Rentzos, and George P. Patrinos. 2019. "New Molecular Diagnostic Trends and Biomarkers for Amyotrophic Lateral Sclerosis." *Human Mutation* 40 (4): 361–73. https://doi.org/10.1002/humu.23697.

Panchenko, M. V., W. G. Stetler-Stevenson, O. V. Trubetskoy, S. N. Gacheru, and H. M. Kagan. 1996. "Metalloproteinase Activity Secreted by Fibrogenic Cells in the Processing of Prolysyl Oxidase. Potential Role of Procollagen C-Proteinase." *The Journal of Biological Chemistry* 271 (12): 7113–19.

Pearsall, Nicole, Gautam Bhattacharya, Jim Wisecarver, Joe Adams, Dominic Cosgrove, and William Kimberling. 2002. "Usherin Expression Is Highly Conserved in Mouse and Human Tissues." Hearing Research 174, no. 1: 55–63. https://doi.org/10.1016/S0378-5955(02)00635-4.

Pedrosa-Domellöf, F., I. Virtanen, and L. E. Thornell. 1995. "Tenascin Is Present in Human Muscle Spindles and Neuromuscular Junctions." *Neuroscience Letters* 198 (3): 173–76.

Pénisson-Besnier, Isabelle, Valérie Allamand, Philippe Beurrier, Ludovic Martin, Joost Schalkwijk, Ivonne van Vlijmen-Willems, Corine Gartioux, et al. 2013. "Compound Heterozygous Mutations of the TNXB Gene Cause Primary Myopathy." *Neuromuscular Disorders: NMD* 23 (8): 664–69. https://doi.org/10.1016/j.nmd.2013.04.009.

Petryszak, Robert, Maria Keays, Y. Amy Tang, Nuno A. Fonseca, Elisabet Barrera, Tony Burdett, Anja Füllgrabe, et al. 2016. "Expression Atlas Update—an Integrated Database of Gene and Protein Expression in Humans, Animals and Plants." *Nucleic Acids Research* 44 (D1): D746–52. https://doi.org/10.1093/nar/gkv1045.

Petukhova, Lynn, Yutaka Shimomura, Muhammad Wajid, Prakash Gorroochurn, Susan E. Hodge, and Angela M. Christiano. 2009. "The Effect of Inbreeding on the Distribution of Compound Heterozygotes: A Lesson from Lipase H Mutations in Autosomal Recessive Woolly Hair/Hypotrichosis." *Human Heredity* 68 (2): 117–30.

https://doi.org/10.1159/000212504.

Piñero, Janet, Àlex Bravo, Núria Queralt-Rosinach, Alba Gutiérrez-Sacristán, Jordi Deu-Pons, Emilio Centeno, Javier García-García, Ferran Sanz, and Laura I. Furlong. 2017. "DisGeNET: A Comprehensive Platform Integrating Information on Human Disease-Associated Genes and Variants." *Nucleic Acids Research* 45 (D1): D833–39. https://doi.org/10.1093/nar/gkw943.

Pio-Lopez, Léo, Alberto Valdeolivas, Laurent Tichit, Élisabeth Remy, Anaïs Baudot. 2021. "MultiVERSE: a multiplex and multiplex-heterogeneous network embedding approach." *Scientific reports* 11(1): 8794. https://doi.org/10.1038/s41598-021-87987-1.

Ramanagoudr-Bhojappa, Ramanagouda, Blake Carrington, Mukundhan Ramaswami, Kevin Bishop, Gabrielle M. Robbins, MaryPat Jones, Ursula Harper, et al. 2018. "Multiplexed CRISPR/Cas9-Mediated Knockout of 19 Fanconi Anemia Pathway Genes in Zebrafish Revealed Their Roles in Growth, Sexual Development and Fertility." *PLoS Genetics* 14 (12): e1007821. https://doi.org/10.1371/journal.pgen.1007821.

Richard, P., K. Gaudon, F. Andreux, E. Yasaki, C. Prioleau, S. Bauché, A. Barois, et al. 2003. "Possible Founder Effect of Rapsyn N88K Mutation and Identification of Novel Rapsyn Mutations in Congenital Myasthenic Syndromes." *Journal of Medical Genetics* 40 (6): e81.

Rodríguez Cruz, Pedro M., Jacqueline Palace, Hayley Ramjattan, Sandeep Jayawant, Stephanie A. Robb, and David Beeson. 2015. "Salbutamol and Ephedrine in the Treatment of Severe AChR Deficiency Syndromes." *Neurology* 85 (12): 1043–47. https://doi.org/10.1212/WNL.0000000000001952.

Rodríguez Cruz, Pedro M., Jacqueline Palace, and David Beeson. 2018. "The Neuromuscular Junction and Wide Heterogeneity of Congenital Myasthenic Syndromes." *International Journal of Molecular Sciences* 19 (6). https://doi.org/10.3390/ijms19061677.

Rogers, Robert S., and Hiroshi Nishimune. 2017. "The Role of Laminins in the Organization and Function of Neuromuscular Junctions." *Matrix Biology: Journal of the International Society for Matrix Biology* 57–58: 86–105. https://doi.org/10.1016/j.matbio.2016.08.008.

Sadeh, Menachem, Xin-Ming Shen, and Andrew G. Engel. 2011. "Beneficial Effect of Albuterol in Congenital Myasthenic Syndrome with Epsilon-Subunit Mutations." *Muscle & Nerve* 44, no. 2: 289–91. https://doi.org/10.1002/mus.22153.

Saint-Amant, Louis and Pierre Drapeau. "Time Course of the Development of Motor Behaviors in the Zebrafish Embryo." 1998. *Journal of Neurobiology* 37, no. 4: 622–32. https://doi.org/10.1002/(sici)1097-4695(199812)37:4<622::aid-neu10>3.0.co;2-s.

Saito, Rintaro, Michael E Smoot, Keiichiro Ono, Johannes Ruscheinski, Peng-Liang Wang, Samad Lotia, Alexander R Pico, Gary D Bader, and Trey Ideker. 2012. "A Travel Guide to Cytoscape Plugins." *Nature Methods* 9 (11): 1069–76. https://doi.org/10.1038/nmeth.2212.

Schindelin, Johannes, Ignacio Arganda-Carreras, Erwin Frise, Verena Kaynig, Mark Longair, Tobias Pietzsch, Stephan Preibisch, Curtis Rueden, Stephan Saalfeld, Benjamin Schmid, Jean-Yves Tinevez, Daniel James White, Volker Hartenstein, Kevin Eliceiri, Pavel Tomancak and Albert Cardona. 2012. "Fiji - an Open Source platform for biological image analysis." *Nature Methods* 9, 10.1038/nmeth.2019. https://doi.org/10.1038/nmeth.2019

Schwaller, Fred, Valérie Bégay, Gema García-García, Francisco J. Taberner, Rabih Moshourab, Brennan McDonald, Trevor Docter, Johannes Kühnemund, Julia Ojeda-Alonso, Ricardo Paricio-Montesinos, Stefan G. Lechner, James F. A. Poulet, Jose M. Millan and Gary R. Lewin. 2021. "USH2A is a Meissner's Corpuscle Protein Necessary for Normal Vibration Sensing in Mice and Humans." *Nature Neuroscience* 24, no. 1: 74–81. https://doi.org/10.1038/s41593-020-00751-y.

Senderek, Jan, Juliane S Müller, Marina Dusl, Tim M. Strom, Velina Guergueltcheva, Irmgard Diepolder, Steven H. Laval, Susan Maxwell, Judy Cossins, Sabine Krause, Nuria Muelas, Juan J. Vilchez, Jaume Colomer, Cecilia Jimenez Mallebrera, Andres Nascimento, Shahriar Nafissi, Ariana Kariminejad, Yalda Nilipour, Bita Bozorgmehr, Hossein Najmabadi, Carmelo Rodolico, Jörn P Sieb, Ortrud K. Steinlein, Beate Schlotter, Benedikt Schoser, Janbernd Kirschner, Ralf Herrmann, Thomas Voit, Anders Oldfors, Christopher Lindbergh, Andoni Urtizberea, Maja von der Hagen, Angela Hübner, Jacqueline Palace, Kate Bushby, Volker Straub, David Beeson, Angela Abicht and Hanns Lochmüller 2011. "Hexosamine Biosynthetic Pathway Mutations Cause Neuromuscular Transmission Defect." *American Journal of Human Genetics* 88, no. 2: 162–72. https://doi.org/10.1016/j.ajhg.2011.01.008.

Sorensen, Jacob R., Caitlin Skousen, Alex Holland, Kyle Williams, and Robert D. Hyldahl. 2018. "Acute Extracellular Matrix, Inflammatory and MAPK Response to Lengthening Contractions in Elderly Human Skeletal Muscle." *Experimental Gerontology* 106: 28–38. https://doi.org/10.1016/j.exger.2018.02.013.

Stum, Morgane, Claire-Sophie Davoine, Savine Vicart, Léna Guillot-Noël, Haluk Topaloglu, Francisco Javier Carod-Artal, Hülya Kayserili, et al. 2006. "Spectrum of HSPG2 (Perlecan) Mutations in Patients with Schwartz-Jampel Syndrome." *Human Mutation* 27 (11): 1082–91. https://doi.org/10.1002/humu.20388.

Swuec, Paolo, Ludovic Renault, Aaron Borg, Fenil Shah, Vincent J. Murphy, Sylvie van Twest, Ambrosius P. Snijders, Andrew J. Deans, and Alessandro Costa. 2017. "The FA Core Complex Contains a Homo-Dimeric Catalytic Module for the Symmetric Mono-Ubiquitination of FANCI-FANCD2." *Cell Reports* 18 (3): 611–23. https://doi.org/10.1016/j.celrep.2016.11.013.

Sztal, Tamar E., Avnika A. Ruparelia, Caitlin Williams, and Robert J. Bryson-Richardson. 2016. "Using Touch-Evoked Response and Locomotion Assays to Assess Muscle Performance and Function in Zebrafish." *Journal of Visualized Experiments* : JoVE, no. 116: 54431. https://doi.org/10.3791/54431.

Thompson, Rachel, Anastasios Papakonstantinou Ntalis, Sergi Beltran, Ana Töpf, Eduardo de Paula Estephan, Kiran Polavarapu, Peter A. C. 't Hoen, Paolo Missier, and Hanns

Lochmüller. 2019. "Increasing Phenotypic Annotation Improves the Diagnostic Rate of Exome Sequencing in a Rare Neuromuscular Disorder" *Human Mutation*, June. https://doi.org/10.1002/humu.23792.

Tinevez, Jean-Yves, Nick Perry, Johannes Schindelin, Genevieve M. Hoopes, Gregory D. Reynolds, Emmanuel Laplantine, Sebastian Y. Bednarek, Spencer L. Shorte, and Kevin W. Eliceiri. "TrackMate: An Open and Extensible Platform for Single-Particle Tracking." *Methods (San Diego, Calif.)* 115 (February 15, 2017): 80–90. https://doi.org/10.1016/j.ymeth.2016.09.016.

Valdeolivas, Alberto, Laurent Tichit, Claire Navarro, Sophie Perrin, Gaëlle Odelin, Nicolas Levy, Pierre Cau, Elisabeth Remy, and Anaïs Baudot. 2019. "Random Walk with Restart on Multiplex and Heterogeneous Biological Networks." *Bioinformatics (Oxford, England)* 35 (3): 497–505. https://doi.org/10.1093/bioinformatics/bty637.

Vanhaesebrouck, An E, Richard Webster, Susan Maxwell, Pedro M Rodriguez Cruz, Judith Cossins, James Wickens, Wei-wei Liu, et al. 2019. "B2-Adrenergic Receptor Agonists Ameliorate the Adverse Effect of Long-Term Pyridostigmine on Neuromuscular Junction Structure." *Brain* 142, no. 12: 3713–27. https://doi.org/10.1093/brain/awz322.

Vázquez, Miguel, Rubén Nogales, Pedro Carmona, Alberto Pascual, and Juan Pavón. 2010. "Rbbt: A Framework for Fast Bioinformatics Development with Ruby." In *Advances in Bioinformatics*, edited by Miguel P. Rocha, Florentino Fernández Riverola, Hagit Shatkay, and Juan Manuel Corchado, 74:201–8. Berlin, Heidelberg: Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-13214-8_26.

Voermans, N. C., and B. G. van Engelen. 2008. "Differential Diagnosis of Muscular Hypotonia in Infants: The Kyphoscoliotic Type of Ehlers-Danlos Syndrome (EDS VI)." *Neuromuscular Disorders: NMD* 18 (11): 906; author reply 907. https://doi.org/10.1016/j.nmd.2008.05.016.

Voermans, Nicol C., Karin Gerrits, Baziel G. van Engelen, and Arnold de Haan. 2014. "Compound Heterozygous Mutations of the TNXB Gene Cause Primary Myopathy." *Neuromuscular Disorders: NMD* 24 (1): 88–89. https://doi.org/10.1016/j.nmd.2013.10.007.

Wang, Dan-Ni, Zhi-Qiang Wang, Yu-Qing Chen, Guo-Rong Xu, Min-Ting Lin, and Ning Wang. 2018. "Limb-Girdle Muscular Dystrophy Type 2I: Two Chinese Families and a Review in Asian Patients." *The International Journal of Neuroscience* 128 (3): 199–207. https://doi.org/10.1080/00207454.2017.1380640.

Wen, Hua, Michael W. Linhoff, Matthew J. McGinley, Geng-Lin Li, Glen M. Corson, Gail Mandel, and Paul Brehm. 2010. "Distinct Roles for Two Synaptotagmin Isoforms in Synchronous and Asynchronous Transmitter Release at Zebrafish Neuromuscular Junction." *Proceedings of the National Academy of Sciences of the United States of America* 107, no. 31: 13906–11. https://doi.org/10.1073/pnas.1008598107.

Whicher, Danielle, Sarah Philbin, and Naomi Aronson. 2018. "An Overview of the Impact of Rare Disease Characteristics on Research Methodology." *Orphanet Journal of Rare Diseases* 13 (1): 14. https://doi.org/10.1186/s13023-017-0755-5.

Xu, Zhuo, Naoki Ichikawa, Keisuke Kosaki, Yoshihiko Yamada, Takako Sasaki, Lynn Y.
Sakai, Hisashi Kurosawa, Nobutaka Hattori, and Eri Arikawa-Hirasawa. 2010.
"Perlecan Deficiency Causes Muscle Hypertrophy, a Decrease in Myostatin
Expression, and Changes in Muscle Fiber Composition." *Matrix Biology: Journal of*
the International Society for Matrix Biology 29 (6): 461–70.
https://doi.org/10.1016/j.matbio.2010.06.001.
Yang, Kunfang, Hongyi Cheng, Fang Yuan, Linyi Meng, Rongrong Yin, Yuanfeng Zhang,
Simei Wang, et al. 2018. "CHRNE Compound Heterozygous Mutations in Congenital
Myasthenic Syndrome: A Case Report." *Medicine* 97 (17): e0347.
https://doi.org/10.1097/MD.0000000000010347.
Zhong, Jingzi, Gang Chen, Yiwu Dang, Haixia Liao, Jiapeng Zhang, and Dan Lan. 2017.
"Novel Compound Heterozygous PLEC Mutations Lead to Early-onset Limb-girdle
Muscular Dystrophy 2Q." *Molecular Medicine Reports* 15 (5): 2760–64.
https://doi.org/10.3892/mmr.2017.6309.
Zitnik, Marinka, and Jure Leskovec. 2017. "Predicting Multicellular Function through Multi-
Layer Tissue Networks." *Bioinformatics (Oxford, England)* 33 (14): i190–98.
https://doi.org/10.1093/bioinformatics/btx252.