Multiomic interpretation of fungus-infected ant metabolomes during manipulated summit disease.

I. Will[1*], G. M. Attardo[2], C. de Bekker[1,3*]

1: University of Central Florida, Biology Department, USA

2: University of California – Davis, Entomology and Nematology Department, USA

3: Utrecht University, Biology Department, NL

* Corresponding authors: ian.will@knights.ucf.edu (IW), a.m.debekker@uu.nl (CdB)

ORCiD: 0000-0002-2396-6778 (IW), 0000-0001-6265-2969 (GMD), 0000-0002-7325-172X (CdB)

Abstract

Camponotus floridanus ants show altered behaviors followed by a fatal summiting phenotype

when infected with manipulating *Ophiocordyceps camponoti-floridani* fungi. Host summiting as a

strategy to increase transmission is also observed with parasite taxa beyond fungi, including aquatic and

terrestrial helminths and baculoviruses. The drastic phenotypic changes can sometimes reflect significant

physiological changes within host cells that span molecular levels from metabolites to nucleic acids.

Nevertheless, the underlying mechanisms still need to be fully characterized. To investigate the small

molecules producing summiting behavior, we infected *C. floridanus* ants with *O. camponoti-floridani* and

sampled their heads for LC-MS/MS when we observed the characteristic summiting phenotype. We link

this metabolomic data with our previous genomic and transcriptomic data to propose mechanisms that

underlie manipulated summiting behavior in "zombie ants." This "multiomic" evidence points toward the

dysregulation of neurotransmitter levels and neuronal signaling. We propose that these processes are

altered during infection and manipulation based on 1) differential expression of neurotransmitter synthesis

and receptor genes, 2) altered abundance of metabolites and neurotransmitters (or their precursors) with

known behavioral effects in ants and other insects, and 3) possible suppression of a connected immunity

pathway. We additionally report signals for metabolic activity during manipulation related to primary

metabolism, detoxification, and anti-stress protectants.

Introduction

Infections often lead to altered host phenotypes that can be adaptive for the host or parasite. Parasite-adaptive changes in animal host behavior are mostly referred to as manipulations. Apparent manipulations of host behavior and physiology occur in a broad range of host and parasite taxa. While descriptive examples are plenty, the underlying mechanisms are still in the process of being revealed [1–5].

The ant-manipulating *Ophiocordyceps* (zombie ant fungi) have been leveraged in laboratory studies to better understand the mechanistic basis of behavioral manipulation [5–7]. Infected ants display summit disease: a common parasitic manipulation observed in invertebrates where the manipulated individual occupies an elevated location and dies stereotypically [5,8]. These final, elevated, exposed summit positions can be adaptive for the parasite by either aiding trophic transmission or direct dispersal of infectious propagules. In the case of *Ophiocordyceps*, ants latch and bite onto their final summit perches until death, which is thought to promote development of the fungal fruiting body and dispersal of the spores therein [9–13]. Moreover, preceding this fatal change in behavior, infected ants also deviate from foraging trails and increase locomotor activity, reduce nestmate communication, and convulse [12–17].

These various changes may represent behaviors that can be easily coopted for manipulation by the fungal parasite [5]. As such, the parasite may be taking advantage of existing host processes and symptoms without relying on costly mechanisms to "rewire" their hosts [8,18]. A profound understanding of the fitness costs and benefits of these behaviors, for both the host and parasite, may reveal that some could also be adaptive for the host, at the individual or group level (perhaps especially so for eusocial animals, such as ants). Regardless of the ultimate "source" of the altered behaviors observed, physiological and molecular characterizations of hosts displaying modified behavior may offer a glimpse into how parasites (or hosts) modulate animal behavior, the coevolution of specialized host-parasite relationships, and undescribed properties of proteins and metabolites involved.

In the myrmecophilous *Ophiocordyceps*, investigations of phenotypes, genomes, transcriptomes, and metabolomes continue to bring various mechanistic hypotheses of behavioral manipulation into focus. The range of hypotheses comprise fungal neuromodulators and effector proteins, host circadian

rhythms and phototropism, insect hormones, fungal neuroprotectants, and ant muscular hyperactivity

[6,7,14,19–24]. Previous metabolomic studies, centered on the *Ophiocordyceps kimflemingiae* zombie ant

fungus and its host *Camponotus castaneus*, have identified various metabolites associated with

manipulation. These metabolomic signals also varied in a species-specific manner, changing when either

the parasite or host was exchanged for a different species [21,22,25]. However, these studies did not place the

results within the context of other types of "omics" data. Doing so would present the opportunity to

distinguish the most robust lines of evidence from the more stochastic signals that might differ across

experimental conditions and setups.

Here, we present a "multiomic" approach by conducting a metabolomic analysis of *Camponotus*

floridanus (Florida carpenter ant) manipulated by *Ophiocordyceps camponoti-floridani* (Florida zombie

ant fungus) interpreted in the context of our previous transcriptomic and genomic data [24]. To add to our

existing transcriptomic study, we collected metabolomic data using three liquid chromatography-tandem

mass spectrometry (LC-MS/MS) protocols. Our study compares whole head samples of healthy *C.*

floridanus ants and ants manipulated by *O. camponoti-floridani*, allowing direct comparison to the

transcriptomic ant-head data. Whole heads contain the chemical profiles of ant neural tissue, muscle, and

hemolymph in addition to fungal cells that all may contain metabolites of most interest [13,15,20–22,25,26].

We primarily focused our interpretation and discussion of the results on altered metabolic

pathways that are supported by a combination of metabolite and gene signals. This improves mechanistic

interrogation as the key processes of manipulation could involve specialized and general mechanisms,

host and parasite effectors, and changes from the genetic to chemical level. At the same time, combining

multiple omics approaches and datasets allowed us to identify consistent and robust signals for key

biological processes. To analyze our metabolomics data we also combined multiple statistical methods to

reduce the inherent bias that any given analysis introduces. Data that represent substantial biological

changes are more likely to be selected by multiple complementary tests. Our reporting, therefore,

highlights manipulation-associated metabolite features that were selected by multiple statistical methods.

With this, we propose multiple pathways and compounds that reflect, or drive, manipulated behavior in

Ophiocordyceps infected ants.

Results & discussion

82   **Infection mortality, observations of manipulation, and LC-MS/MS:** Similar to previous

laboratory infections with *O. camponoti-floridani* [24], we collected *C. floridanus* displaying manipulated

summiting between four hours before (zeitgeber time, ZT 20) to half an hour after dawn (ZT 0.5),

beginning three weeks after infection (Fig. 1). Sham-treated healthy ants showed no mortality until 29

days post injection (dpi), whereas the survival rate among infected ants gradually decreased from the

beginning of the infection experiment, with a steeper drop during the first half of the manipulation phase

starting in week three. As such, survival was markedly different between treatments (p = 4.1E-7, log-rank

test) (Fig. 1A). In total, three sham-treated ants died out of the 28 (11%) that survived the initial injection

procedure. Twenty-three infected ants were manipulated out of the 39 (59%) that survived the injection.

The remaining 16 infected ants died with fungal infection, indicated by fungal blastospores present upon

microscopic investigation, but without observed behavioral manipulation. From these experimental

infections, we sampled 20 manipulated ants and 20 healthy (sham) ants for LC-MS/MS. We also

produced three blank samples to contrast our treatment samples against. Using these samples, we

produced three LC-MS/MS datasets: "biogenic amines" (2,955 features), "polyphenols" (4,315 features),

and "lipids" (4,924 features).

Figure 1. Ant manipulations occur at stereotypical days and times.

a) *Ophiocordyceps camponoti-floridani* infection leads to significantly increased mortality compared to sham treatment (p = 4.1E-7, log-rank test). **b**) Manipulated ants most often appeared 21-27 dpi, but as late as 33 dpi. **c**) Summiting behavior always occurred around dawn, most often 2.5 h before lights turned on, at ZT 21.5.

Metabolomic feature selection and grouping for pathway enrichment analyses: We combined multiple analyses to stringently select "top features" and metabolic pathways for further biological interpretation. We identified metabolite features that distinguished healthy and manipulated ants by their *i*) being differentially abundant metabolites (DAMs), *ii*) loading values in principal component analyses (PCAs), or *iii*) importance for navigating decision trees (Boruta) [27] (Fig. 2A). Combining all three LC-MS/MS datasets, approximately 60% of features were DAMs, with most DAMs increasing in abundance in manipulated ants compared to healthy controls (Fig. 2B, Supplementary Table S1). Separation of treatment groups by PCA only clearly clustered along PC1 (ca. 60% to 80%), with PC2 (ca. 4% to 9%) and other PCs (not shown) describing little of the variation between treatments (Fig. 2C). Using Boruta random forests, we found ~40% of features to be "confirmed" as "important" for

distinguishing treatment types (Supplementary Table S1). We performed pathway analyses with top

features that satisfied at least two of the following criteria: the feature was *i)* found to be a DAM, *ii)*

among the top 50% of absolute PC1 loading values, or *iii)* confirmed by Boruta. Most top features that

satisfied at least two requirements (n = 6,468 features, 53% of a total 12,194 features), in fact satisfied all

three (n = 4,656, 72% of features passing two criteria) (Fig. 2D, Supplementary Table S1, Supplementary

Data S1). The feature set passing at least two filters contained 347 of the 760 (46%) features that were

functionally annotated. This many features passing our stringent selection approach is likely due to a

combination of small molecules related specifically to infection processes and general aspects of fungal

metabolism, as *O. camponoti-floridani* cells were only present in manipulated ants.

We analyzed the 6,468 top features with Mummichog and a gene set enrichment analysis

(GSEA)-based approach to detect enriched Kyoto Encyclopedia of Genes and Genomes (KEGG)

metabolic pathways (Fig. 2A) [28–30]. We found eight significantly enriched pathways that passed our p-

value, fold-enrichment, metabolite feature count, and enzyme differentially expressed gene (DEG) count

thresholds (Fig. 2A, Fig. 2D, Supplementary Data S2). These enzyme DEGs were identified by their

association with KEGG metabolic pathways and the previously published expression data from our

Ophiocordyceps-infection study in which the heads of manipulated ants were compared to those of

healthy ants using RNASeq [24]. Combining enriched pathways from these top features and the network

module analysis (see below) (Fig. 2A, Fig. 3, Table 1), most enzyme DEGs were downregulated both in

the ant (90%) and the fungus (62%). This degree of downregulation is higher than previously found for

transcriptome-wide DEGs during manipulation [24] (Supplementary Discussion S1). We infer a positive

correlation between DEG transcription and protein levels as the most parsimonious interpretation of the

data. Although, gene expression reflecting homeostatic responses to changes in functional protein

abundance or a lack of consistent correlation are alternative explanations, which future proteomic data

could help make clear.

The eight KEGG metabolic pathways enriched among top features included signals that suggest

altered primary metabolism related to amino sugars, galactose, valine and leucine, pyrimidines, and fatty

a **Feature sets selection:** analyses between healthy and manipulated ants

Boruta: random forest selection ("important")

DAMs: differential abundance (≥ |1.5|-fold change, p ≤ 0.05)

PCA: loading value (top 50%)

Top features passing ≥ 2 analyses (combined as 1 feature set)

WGCNA: module correlation (correlation ≥ |0.5|, p ≤ 0.05) → Manipulation-correlated modules (16 modules as 16 feature sets)

Pathway enrichment analyses: GSEA and/or Mummichog analyses on feature sets, based on *D. melanogaster* and *S. cerevisiae* KEGG databases

feature set

pathway enrichments

background features

Pathway enrichment selection: final metabolic pathway enrichments undergoing change during manipulation

- p ≤ 0.05
- ≥ 1.25-fold enrichment
- ≥ 3 metabolite features
- ≥ 3 differential expressed enzyme genes

b Biogenic amines   Polyphenols   Lipids

$-\log_{10}(p)$ $-\log_2$ (fold-change)   Decreased DAMs   Increased DAMs
(from healthy to manipulated)

c Biogenic amines   Polyphenols   Lipids

PC 2 (4.7%)   (9.1%)   (4.0%)

PC 1 (68.5%)   (58.6%)   (82.9%)

Manipulated Healthy

d DAM   PCA
950   **1229**   205
575   **4656**   **8**
83
Boruta

8 enriched KEGG pathways

Figure 2. Selection of top features based on three independent analyses.

a) We used three independent statistical tests to select a single top features for metabolic pathway enrichment

analyses. Additionally, we used a weighted co-abundance network approach (analogous to WGCNA used for gene

expression data) to form manipulation-correlated feature modules (resulting in 16 feature sets) for enrichment

analyses. Results from those enrichment analyses were subsequently subjected to multiple filters to arrive at a final

set of pathways and feature sets of interest. We also performed an enrichment analysis on a set of features only

present in manipulated samples, but we found no overrepresented pathways and this feature set is not depicted in the

figure. **b**) Volcano plots per dataset show DAMs that increased (red) or decreased (blue) at manipulation according

to a 1.5-fold change and $p \leq 0.05$ t-test threshold. **c**) PCAs for each dataset show distinct clustering of samples

between manipulated (orange) and healthy ants (gray) along PC1. d) Features that passed at least two selection

criteria (bold) were classified as top features and used for further pathway analyses. Top features were enriched for

eight KEGG metabolic pathways.

acids (Table 1). Changes in primary metabolism could reflect various processes, including diseased host

metabolism, parasite presence or acquisition of host resources, or host-parasite cross-talk [31–37].

Furthermore, the previous transcriptomics data prompted hypotheses connecting behavioral manipulation

to nutritional state and insulin-related signaling [24].

158   Other enriched pathways plausibly reflect manipulated behavior and infection in different aspects

of metabolism (Table 1). We found KEGG pathway "drug metabolism – other enzymes" enriched among

top features. Broadly suggesting antagonistic host-parasite chemical interactions, this pathway annotation

contains multiple sub-pathways corresponding to different toxic metabolites, which are difficult to resolve

as we lacked positive identification of these specific compounds in our data (Supplementary Discussion

S2). Additionally, as aspects of enriched "beta-alanine metabolism," we detected an apparent carnosine

feature and its constituent amino acids as DAMs (Supplementary Discussion S3). Although we lack

robust transcriptomic support for the synthesis of carnosine (Supplementary Discussion S3), the

combination of pathway enrichment and metabolite feature data indicated that carnosine is produced from

host amino acids during infection. This compound may act as a cytoprotectant during the stresses

behavioral manipulation [38–41]. We also further discuss enriched "glycerophospholipid metabolism" in

context with its multiomic signals and identified metabolites in detail below.

Figure 3. WGCNA network modules of metabolite features, their correlation to manipulation, and enriched pathways. In total we detected three, five, and eight network modules that were significantly correlated to manipulation (Student correlation ≥ |0.5|, p ≤ 0.05) in the biogenic amines, lipids, and polyphenol datasets, respectively. "Unsorted" modules contain all features that were not successfully correlated to other features in a network.

Table 1. Enriched KEGG pathways selected across analyzed feature sets.
For KEGG pathways that were found in Mummichog or GSEA analyses (p ≤ 0.05) to be of interest, we also required them to show a minimum 1.25-fold enrichment, have at least three features contributing to that enrichment, and the previous transcriptomics dataset [24] to contain at least three DEGs homologous to representative enzymes within that pathway. Pathways are listed alphabetically with all feature sets (network modules or the top features set) that were enriched for that pathway. Network modules names: biogenic amine (A1-A6), lipid (L1-L10), and polyphenol (P1-P19). Only 16 of these 35 modules were correlated to manipulation and appear in this table.

Enriched KEGG metabolic pathway	Feature sets
Alanine, aspartate and glutamate metabolism	P13

Pathway						
Amino sugar and nucleotide sugar metabolism	A1	P1	P2	P3	P18	top features
Aminoacyl-tRNA biosynthesis	L3	P2	P11			
Arginine and proline metabolism	A1	P2				
Arginine biosynthesis	A1					
Ascorbate and aldarate metabolism	A1	P13				
beta-Alanine metabolism	P2	top features				
Biosynthesis of unsaturated fatty acids	L2	L10	P3	top features		
Butanoate metabolism	P2					
Citrate cycle (TCA cycle)	P3	P13				
Cyanoamino acid metabolism	A1					
Cysteine and methionine metabolism	L1	P1	P18			
Drug metabolism - cytochrome P450	L3	P3				
Drug metabolism - other enzymes	A1	top features				
Fatty acid degradation	P3					
Folate biosynthesis	P2					
Fructose and mannose metabolism	P2	P18				
Galactose metabolism	A1	P18	top features			
Glutathione metabolism	A1					
Glycerolipid metabolism	P2					
Glycerophospholipid metabolism	P2	top features				
Glycine, serine and threonine metabolism	P2	P11				
Glycolysis or Gluconeogenesis	P3					
Glyoxylate and dicarboxylate metabolism	P13					
Histidine metabolism	A1					
Inositol phosphate metabolism	P2					
Insect hormone biosynthesis	A1	P3				
Lysine biosynthesis	A1					
Lysine degradation	P2					
Metabolism of xenobiotics by cytochrome P450	A1	P2				
Nicotinate and nicotinamide metabolism	P2					
Pantothenate and CoA biosynthesis	A1	P18				
Pentose and glucuronate interconversions	A1	P18				
Pentose phosphate pathway	A1	L1	P1	P2	P3	P18
Phenylalanine metabolism	A1	P2				
Purine metabolism	L1	P1	P18			
Pyrimidine metabolism	A1	P11	top features			
Pyruvate metabolism	P13					

Starch and sucrose metabolism	A1	P3	P18
Sulfur metabolism	P2		
Taurine and hypotaurine metabolism	L1	P1	
Tyrosine metabolism	P2	P3	
Ubiquinone and other terpenoid-quinone biosynthesis	P3		
Valine, leucine and isoleucine degradation	P18	top features	

Many metabolite features in our dataset were uniquely represented in manipulated samples: 454

features were never detected in healthy sham treatment ants and 1,781 additional features either were

detected in less than 50% of the healthy ant samples and/or their median peak value was less than two-

fold that of blank controls (2,235 features total). These features may represent uniquely produced fungal

molecules or ant molecules that are tightly upregulated with *Ophiocordyceps*-infection while being scarce

in healthy ants. Although we predicted that hallmarks of infection and manipulation processes would be

overrepresented among these unique features, we did not find any significant pathway enrichments. This

could indicate that these metabolites are not linked by common metabolic pathways and/or many of these

metabolites were not well annotated in the KEGG databases (e.g., species-specific metabolites).

Co-expression (or co-abundance) networks can suggest linked biological functions and activity in

the same metabolic processes [42–45]. To cluster metabolite features that similarly change abundance

between healthy and manipulated *C. floridanus* treatments, we used a weighted gene co-expression

network analysis (WGCNA) based approach (Fig. 2A, Fig. 3) [46]. This resulted in a total of six biogenic

amine (A1-A6), 10 lipid (L1-L10), and 19 polyphenol (P1-P19) modules. Of these network modules,

three biogenic amine, five lipid, and eight polyphenol modules were significantly correlated to

Ophiocordyceps manipulation (Fig. 3, Table 1). Due to our sparse annotation data obtained from mass

spectrometry database matching, we were not able to consistently compare hub features. Instead, we

considered all available annotations within a network module and Mummichog pathway enrichments

(Fig. 2A). Across all network modules significantly correlated to the transition from healthy to

manipulated ants, we found 44 unique enriched KEGG pathways that passed our selection criteria, which

included the eight found from the top features set (Table 1, Supplementary Data S2).

Biogenic monoamine neurotransmitters as putative proximate drivers of *Ophiocordyceps*

infection-related behaviors: A combination of pathway enrichments, annotated DAMs, and DEGs

suggest that shifts in metabolism associated with neurotransmitters relate to behavioral alterations in *C.*

floridanus infected and manipulated by *O. camponoti-floridani*. The monoamine neurotransmitters

dopamine, serotonin, and octopamine (analogous to vertebrate norepinephrine) have been implicated in

modulating locomotor, foraging, learning, social, reproductive, and aggressive behaviors in many insects,

and have been tested in social insects, including ants [47–51]. Serotonin, octopamine, and melanin derived

from L-3,4-dihydroxyphenylalanine (L-DOPA), or tyrosine and dopamine, have roles in insect immunity

as well [52,53]. Although we did not directly identify dopamine, serotonin, or octopamine in our

metabolomic data, we did measure features annotated as biosynthetically related compounds.

Furthermore, gene expression networks negatively correlated between the ant and fungus were enriched

for host neuronal function and parasite manipulation-associated secreted proteins. This correlation

suggested that parasite effectors promote, directly or indirectly, the downregulation of host gene modules

related to neuronal maintenance, circadian rhythm, olfaction, and memory [24].

Moreover, these neurotransmitters can affect behaviors relatable to *Ophiocordyceps* disease

phenotypes. In *Formica polyctena* wood ants and *Odontomachus kuroiwae* trap-jaw ants, administration

of dopamine and serotonin induced aggressive mandible-opening or biting behaviors and defensive

responses in the laboratory [54,55]. In the field, *Pogonomyrmex barbatus* harvester ant dopamine levels

positively correlated with foraging trips [56]. Similarly, pharmacological depletion of serotonin led to

reduced and impaired trail-following foraging behavior in *Pheidole dentata* ants [57]. In *Formica aquilonia*,

there is evidence for octopamine increasing aggressive behaviors [58]. However, increased aggression upon

octopamine treatment was not found in *F. polyctena* [54]. Octopamine additionally appears to modulate the

social behavior of trophallaxis in *Camponotus fellah*, while serotonin had no significant effect [59]. As such,

the exact effects of these neurotransmitters can differ by species and social or physiological context.

However, parallels exist with *Ophiocordyceps*-infected ants that exhibit changes in behavior that include

increased walking, deviation from foraging trails, reduced nestmate communication, and the final

manipulated bite [12–16].

Changes in dopamine metabolism possibly modifies ant behavior. A combination of metabolomic

and transcriptomic data indirectly suggest increased dopamine metabolism in manipulated ants. The

metabolomic signals consisted of changes in neurotransmitter precursor abundances and enrichment of

pathways in metabolite network modules correlated to manipulation. Phenylalanine and tyrosine are

precursors in synthesis pathways for dopamine and octopamine. KEGG pathway "phenylalanine

metabolism" was enriched in both network modules A1 and P2 ($p = 0.026$ and 0.028, fold-enrichment =

1.5 and 1.9, respectively), both of which were negatively correlated to manipulated ant samples.

Additionally, KEGG pathway "tyrosine metabolism" was enriched in modules P2 and P3 ($p = 0.030$ and

0.024, fold-enrichment = 1.4 and 1.6, respectively), both also negatively correlated to manipulation

(Figure 3, Table 1). Moreover, module P2 also contained the most informative compound annotations

regarding dopamine metabolism.

The immediate precursor to dopamine, L-DOPA, was 2.8-fold reduced in abundance in

manipulated ants ($p = 0.008$, top feature) (Fig. 4). L-tyrosine was also annotated in

Figure 4. Altered neurotransmitter precursor metabolism suggests suppressed ant immunity and increased
dopamine during infection and manipulation. Upregulation of enzyme genes converting tyrosine to L-DOPA to
dopamine coupled with reduced tyrosine and L-DOPA levels suggest increased concentrations of dopamine during
infection, which may, in turn, contribute to manipulated behavior. Additionally, cardol triene may be produced by
the fungus to suppress ant immunity by inhibiting melanin synthesis and thereby also increase availability of
tyrosine and L-DOPA for dopamine synthesis. Combined with these specific metabolites, phenylalanine and
tyrosine metabolism pathway enrichments further implicate these processes in manipulated behavior. Increases in
expression, abundance, or activity are indicated by solid lines and red color, decreases by dotted lines and blue
color. Metabolites are in rounded rectangles, genes are in rectangles. Annotated metabolites that were not DAMs
have gray color, while compounds with white color were not identified in the LC-MS/MS data. Hypothesized links
to ant (green) and fungal (orange) phenotypes are given in the shaded circles. Black arrows are shown when useful
for context to indicate that an enzymatic step had a corresponding enzyme gene in either the fungal or ant genome,
but it was not a DEG.

module P2 and is the precursor to L-DOPA. L-Tyrosine was a DAM with a 2.1-fold decrease in

abundance within the polyphenols dataset (p = 1E-4, top feature). An annotation for L-tyrosine was also

present in the network module A1, however, we did not identify this metabolite feature as a DAM in the

biogenic amines dataset (p = 0.154, 1.2-fold reduction). Outside of network modules A1, P2, and P3, we

also detected phenylalanine, the precursor to tyrosine. Phenylalanine showed a modest increase in

abundance during manipulation just below our 1.5-fold change threshold for classification as a DAM (p =

0.019, 1.4-fold increase) (Fig. 4). Decreases in L-tyrosine and L-DOPA are difficult to interpret, but

possibly indicate their conversion to produce high concentrations of dopamine. Alternatively, reduced

precursor abundances could reflect wide-spread reduced levels of metabolites, including dopamine.

However, our previous RNAseq data suggested an increased dopamine synthesis activity causing the

depletion of these dopamine precursors.

During manipulation, both ant hosts and fungal parasites upregulated an aromatic L-amino acid

decarboxylase (AADC) gene (Fig. 4). This enzyme catalyzes the final step of L-DOPA conversion to

dopamine [60]. The AADC enzyme requires vitamin B6 as a cofactor. We detected differential abundance

of an inactive vitamer form of B6 and a precursor for its *de novo* synthesis in module A5 (Supplementary

Discussion S4). However, the combination of our metabolomic and transcriptomic data do not provide for

a clear interpretation as to if active vitamin B6 is changing in abundance and in which organism(s).

In addition, a tyrosine 3-monooxygenase, which catalyzes the rate-limiting conversion of L-

tyrosine to L-DOPA, was upregulated by manipulated ants. Meanwhile, the ant homolog to Henna (i.e.,

DTPHu phenylalanine 4-monooxygenase), which converts phenylalanine to L-tyrosine was not

differentially expressed. This enzyme also converts peripheral tryptophan to serotonin's precursor in *D.*

melanogaster [61–63]. This consistent expression may help explain the only modest change of phenylalanine

during manipulation while its derivatives were found at lower concentrations.

S*hifts in dopamine and melanin metabolism could mediate host-pathogen immunological*

interactions during manipulation. Dopamine and its precursors can also be converted into melanin, a key

compound for insect wound response and immunity [52,64,65]. Additionally, fungi can synthesize melanin for

use in host-pathogen interactions and against competing microbes [66,67]. We found evidence that this

pathway for melanin production is suppressed in the ant during manipulation and, possibly, increased in

the fungus (Fig. 4).

We propose that the production of a fungal-derived compound may inhibit host tyrosinases

necessary to convert dopamine-pathway metabolites to melanin and hence simultaneously steer these

metabolites toward dopamine production. This metabolite feature was annotated as cardol triene (i.e., 5-

(8,11,14-pentadecatrienyl)resorcinol) and was a DAM only detected in manipulated ant samples (p =

0.0073, median peak value = 19,604, top feature) (Fig. 4). As cardol triene was never detected in healthy

ants, it plausibly is produced solely by the fungus, although we cannot discount the possibility that

production of this metabolite is produced by ants combatting infection and completely absent in healthy

individuals. We detected the putative cardol triene in six of 10 manipulation samples, which could

indicate that cardol triene levels are tightly regulated. Cardol triene has been shown to inhibit tyrosinase

oxidation of L-DOPA in vitro [68]. In insects, phenoloxidases are a type of tyrosinase that are key enzymes

that begin the conversion of dopamine and L-DOPA to pro-immune melanin production pathways [52,65,69].

As such, we hypothesize that fungal cardol triene chemically inhibits host phenoloxidase activity to

reduce host melanin metabolism.

Additionally, ants showed mixed transcriptional signals related to the melanin-immune response

by phenoloxidase and phenoloxidase activating factor genes (Fig. 4) [70]. Entomopathogen infections

inhibit phenoloxidase activity in mosquitoes and suppression of the activation of pro-phenoloxidase has

been hypothesized as a mechanism of fungal interference with insect host immunity [71,72]. In addition to

suppressing host melanin production, the fungus may be converting L-DOPA or related compounds into

fungal melanin precursors, as four upregulated and one downregulated tyrosinases were found during

manipulation (Fig. 4). Three of the upregulated fungal tyrosinases are putatively secreted and may

possibly interact with host tyrosine and its derivatives. Taken together, our metabolomics and previous

transcriptomics data not only suggest that altered dopamine metabolism could lead to increased dopamine

concentration that induces altered host behavior, but also result in weakened host immunity while

bolstering parasite pathogenicity (Fig. 4).

Serotonin and octopamine metabolism may also correlate to manipulated ant behavior.

Biosynthetically and phenotypically intertwined with dopamine, fluctuating serotonin and octopamine

concentrations may also contribute to manipulated ant behavior. Although the metabolomic signals for

these compounds were less robust than for dopamine, we found multiomic evidence that offers support

for hypotheses of increasing serotonin and decreasing octopamine (Supplementary Discussion S5). For

example, higher serotonin concentration is indicated by upregulation of the AADC enzyme shared

between dopamine and serotonin pathways [60] and at least one signal for the reduction of tryptophan,

which can be converted into the precursor of serotonin. Coupled with the increasing serotonin, we

additionally propose a hypothesis for increased kynurenic acid, which is also tied to tryptophan and may

act as a neuroprotectant for heavily diseased and manipulated ants (Supplementary Discussion S5).

Reduced octopamine is suggested by changes in tyramine metabolism evidenced by both reduced L-

tyrosine (a tyramine precursor) and a tyramine derivative (N-acetyltyramine) (Supplementary Discussion

S5).

Altered glycerophospholipid metabolism suggests increased fungal cell membrane activity

and ant accumulation of the excitatory neurotransmitter, acetylcholine: The KEGG pathway

"glycerophospholipid metabolism" was enriched in module P2 and among the top-features (p =0.024 and

0.030, fold-enrichment = 1.9 and 1.4, respectively). Of our annotated features, metabolic signals centered

on choline offered the most insight into altered glycerophospholipid metabolism. We found evidence for

two possible major effects, changes in cell membrane composition and the accumulation of acetylcholine.

Glycerophospholipids and choline-containing compounds have been associated with neurodegenerative

diseases, such as Alzheimer's or Parkinson's, as well as insect host-pathogen responses [73–75]. Choline is a

key component involved in the synthesis of phospholipids associated with forming cell membranes,

especially as incorporated through phosphatidylcholines [76].

Although, in fungi and insects, phosphatidylethanolamines are also a major cell membrane

component [77–79]. Using the lipids LC-MS/MS dataset we compared peak values of

phosphatidylethanolamines (n = 95 features) and phosphatidylcholines (n = 85 features) and found

phosphatidylethanolamines to indeed be abundant in our insect and fungal tissue samples. Within healthy

ants, phosphatidylethanolamines were only slightly more abundant (1.10-fold greater) than

phosphatidylcholines (t-test, p = 0.017, t = -2.657). Within fungus-manipulated ants, there was no strong

difference (t-test, p = 0.739, t = -0.339, phosphatidylethanolamines 1.05-fold higher).

Figure 5. Changing glycerophospholipid metabolism during manipulation primarily indicate altered

acetylcholine levels in the ant and cell membrane metabolism in the fungus.

Manipulated ant samples appeared to have two distinct groups, one with high acetylcholine and one with levels

similar to healthy ants. In manipulated ants, the putative acetylcholinesterase and two acetylcholine receptor genes

were downregulated, suggesting acetylcholine levels were accumulating. We hypothesize that increased

acetylcholine in manipulated ants is associated with the increased muscular activity during early manipulation (e.g.,

tremors, biting, and latching behavior). Once the ant is secured in a summited position and the mandible muscles

(and associated neurons) are further degraded, the fungus may be scavenging and converting the available

acetylcholine to choline, thereby rapidly reducing acetylcholine levels. The fungus may be converting choline to

phospholipids, such as phosphatidylcholines to remodel and build cell membranes. Although we detected multiple

phosphatidylcholine and phosphatidylethanolamine class features, we did not identify the singular metabolites 1,2-

diacyl-sn-glycero-3-phosphocholine and 1-acyl-2-acyl-sn-glycero-3-phosphoethanolamine, annotated as

"phosphatidylcholine" and "phosphatidylethanolamine" in the KEGG glycerophospholipid pathway, respectively.

The legend is the same as described in Figure 5, with the additional inclusion of genes that nearly qualified as DEGs

depicted in gray.

Choline was among the DAMs that increased in abundance during manipulation (p = 1.7E-11,

20.0-fold increase, top feature) (Fig. 5). We also found other DAMs related to choline within the

glycerophospholipid metabolism pathway. Glycerophosphocholine, a choline precursor, increased at

manipulation in both the amine and polyphenol datasets (p = 3.4E-11 and 1.2E-7, 5.9- and 7.7-fold

increase, respectively, top features). We also detected two DAMs interconnected between choline and

phosphatidylethanolamines: ethanolamine (p = 5.2E-5, 2.1-fold increase, top feature) and ethanolamine

phosphate (p =1.6E-9, 5.5-fold decrease, top feature) (Fig. 5).

Multiple phospholipase DEGS in both host and parasite indicate possible altered lipid metabolism

acting on molecules biosynthetically related to choline and ethanolamine compounds. Broadly,

phospholipases cleave phospholipids and have been implicated in cell signaling, fungal pathogenicity, and

cell membrane metabolism [80,81]. Although Menezes *et al.* 2023 note a preponderance phospholipase A2 in

the *Ophiocordyceps australis* genome, with possible functions in pathogenesis – our previous

transcriptomics data did not detect fungal type A2 DEGs during active manipulation. Rather, three

putative *O. camponoti-floridani* phospholipase C genes were DEGs, two upregulated and one

downregulated (Fig. 5). When cleaving either phosphatidylcholines or phosphatidylethanolamines they

can produce an intermediate 1,2-diacyl-sn-glycerol product. Additionally, the fungus expressed a putative

diacylglycerol cholinephosphotransferase gene 1.9-fold higher during manipulation, nearly qualifying as a

DEG. This enzyme combines 1,2-diacyl-sn-glycerol and CDP-choline to produce phosphatidylcholine.

Phosphatidylcholines could then be converted to glycerophosphocholine within two metabolic

steps, but this pathway appeared to be downregulated. Three genes putatively encoding the enzymes

driving this conversion were downregulated in the ant (phospholipases A1, A2, and B). In the fungus, a

phospholipase B-like gene was also downregulated (Fig. 5). The net effect appears to be fungal

composition and/or catabolism of phosphatidylethanolamines and synthesis of phosphatidylcholines (Fig.

5). Fungal phospholipase activity may also disrupt host responses and facilitate destruction of host tissues

[81]. Modification of the fungal cell membrane could also be necessary to accommodate transmembrane

proteins or secretory activity involving organelle membranes such as in the endoplasmic reticulum [83,84].

This altered membrane metabolism may be involved in pathogenesis and release of manipulation

effectors, but alternatively or additionally, relate to changing fungal growth [85,86]. Shortly after

manipulating its host, the fungus undergoes a switch from blastospore to hyphal growth and further

colonizes the host cadaver [12,15,20,26]. In line with this, we also hypothesize that signals related to gamma-

aminobutyric acid (GABA) and glutamate indicate altered fungal growth metabolism (see below,

Supplementary Discussion S6).

Choline can also be metabolized into acetylcholine, the most abundant excitatory

neurotransmitter in insects [87]. In bees, excessive activation of acetylcholine receptors can cause changes

in locomotor, navigational, foraging, and social behaviors [88]. Acetylcholine can be rapidly converted back

to choline by the activity of the acetylcholinesterase enzyme, which has been a target of enzyme-

inhibiting insecticides [89–92]. From this fast enzymatic activity, we intuit that acetylcholine levels could

reflect changes in physiology and behavior at a fine timescale since a subgroup of the manipulated ant

samples showed markedly higher acetylcholine levels than both healthy and other manipulated ants (Fig.

5, Supplementary Fig. 1). Speculatively, this could indicate that, for some fast-changing metabolites such

as acetylcholine, our sampling approach captured multiple subtly different physiological states during the

dynamic process of behavioral manipulation.

Acetylcholine showed a 2.7-fold mean increase at manipulation overall. However, it was not

identified as a DAM (p = 0.142) with the feature-peak value distribution showing two distinct groups.

Three manipulated ants had 5.5-fold higher acetylcholine peak values than the other seven that spanned a

similar range as control samples (Fig. 5). The previously collected transcriptomics data similarly showed

a broad distribution of acetylcholinesterase-2 expression in manipulated ants. Notably,

acetylcholinesterase-2 is the functionally dominant acetylcholinesterase enzyme in *Camponotus* and many

hymenopterans [93]. In the ant, an acetylcholinesterase-2-like gene was downregulated during active

manipulation overall, but two of the five samples had 2.5-fold lower acetylcholinesterase expression than

the other three (Supplementary Fig. S1) [24]. As the ant dies, acetylcholinesterase-2 expression begins to

rebound toward healthy ant levels (Supplementary Fig. S1). Additionally, the ant downregulated a

putative muscarinic acetylcholine receptor gar 2 that has been found in sensory and motor neurons of *C.*

elegans [94]. Nearly a DEG as well, a putative ant nicotinic acetylcholine receptor acr-16 subunit gene was

expressed 1.9-fold lower during manipulation. Possibly, receptor downregulation represents a

homeostatic response to the proposed elevated acetylcholine levels present during part of manipulated

summiting and biting.

421   Although acetylcholine has fast-acting regulatory effects on motor behavior, importantly,

422   glutamate is proposed as the direct primary excitatory neurotransmitter at insect neuromuscular junctions

423   [87,95,96]. Glutamate was nearly classified as a reduced DAM ($p = 0.007$, 1.4-fold decrease) while a related

424   compound, glutamine, did qualify ($p = 1.8E-6$ and $6.2E-11$, 3.3- and 3.8-fold decrease, in the amine and

425   polyphenol datasets, respectively), with transcriptomic evidence suggesting that the host and parasite may

426   be mounting contrasting regulatory responses [24] (Supplementary Discussion S6).

427   **Common metabolites between *Ophiocordyceps-Camponotus* manipulations across species-**

interactions: Previous metabolomic studies with *O. kimflemingiae* have identified many metabolites

associated with infection of *C. castaneus* and proposed links between these compounds and behavioral

manipulation [21,22,25]. Laboratory co-culture of *O. kimflemingiae* with brains from *C. castaneus* and other

ant species showed both shared metabolites and distinct compounds reflecting a combination of target

host-specific fungal secretions and ant species profile [25]. Other studies have compared the metabolomes of

O. kimflemingiae and generalist entomopathogenic *Cordyceps bassiana* infections, again finding both

overlap and differences in metabolomic profiles [21,22].

These studies detected ergothioneine, an amino acid produced by some fungi, but not animals, in

mixed ant-fungal tissue from manipulated ant brains and muscles [21,22]. Corroborating these findings, we

detected two ergothioneine signals (biogenic amine and polyphenol datasets) that were wholly absent or

with peak values near that of blank-collection samples in the healthy ants (p = 7.3E-15 and 4.6E-12, top

features). In addition to fungal metabolism, ergothioneine has been discussed as a possible

neuroprotectant to preserve the ant brain until manipulated summiting and biting are complete [21].

Supporting this line of reasoning, ergothioneine in the ant brain was not detected at elevated levels in

moribund ants infected by the generalist entomopathogen *C. bassiana* that consumes hosts more quickly,

without inducing summit disease [21]. Notably, *Ophiocordyceps* do not appear to typically reside directly

within the brain, suggesting ergothioneine may be secreted by the fungus [13,21,26,97 preprint].

Previous work with *O. kimflemingiae* and this work with *O. camponoti-floridani* also found

adenosine and adenosine-phosphate (AMP) to be differentially abundant between manipulation and

healthy and/or generalist-infected ants [21,22]. We found two features annotated as adenosine, one which was

not differentially abundant and one that was a reduced DAM in manipulated samples (p = 6.6E-4, 2.5-fold

decrease). In contrast, previous works found adenosine to increase in abundance in both manipulated ant

brains and mandible muscles [21,22]. Whether these contrasting results indicate species-specific patterns or

high variation is unclear. We also detected decreased AMP (p = 0.003, 2.1-decrease, top feature), as was

found in manipulated *C. castaneus* [21,22]. In *O. kimflemingiae* infection, a role for adenosine-mediated

neuromodulation was hypothesized, as was changing AMP levels possibly reflecting changes in

mitochondrial activity [21,22]. Cyclic AMP is involved in many pathways, including as a secondary

messenger in G-protein coupled receptor signaling that has been hypothesized to change during *O.*

camponoti-floridani behavioral manipulation with the previous transcriptomic study [24]. If changing

abundance of AMP reflects upon the cyclic form as well, this could relate to dysregulation of receptor

signaling. Metabolically related to adenosine, hypoxanthine and inosine were previously proposed to have

links to energetic and neurological processes in manipulated ants [21,22]. As in those studies, we also

detected increased levels in two of three hypoxanthine-annotated features at manipulation (p = 0.007 and

7.7E-6, 1.6-fold increase or absent in sham, the latter of which was a top feature). We also detected one

inosine and two inosine-monophosphate features increased in manipulated ants (p = 0.002, 3.8E-7, and

1.4E-5, 1.9-, 2.8-, 2.5-fold increase, top features). This could indicate the conversion of AMP and

adenosine to hypoxanthine primarily via inosine intermediates.

Conclusion

Infection and behavioral manipulation of hosts could involve many interacting molecular and

cellular processes. As these host-parasite interactions are better characterized, we can also gain insights

into specialist co-evolutions, the mechanistic basis of animal behavior, and the discovery of new bioactive

compounds. To unravel the mechanistic complexity of infection and manipulated host behavior, we

require functional tests built upon robust hypotheses. Towards this goal, we studied parasitic

manipulation of *C. floridanus* by *O. camponoti-floridani*, combining metabolomic, transcriptomic, and

genomic data.

We found evidence for altered biogenic monoamine neurotransmitter production during infection

and manipulation that may be proximate causes of manipulated behavior. Although, we did not identify

and measure these neurotransmitters directly. Going forward, now with evidence implicating these

neurotransmitters in behavioral manipulation, experiments focused on positively identifying them could

employ increased sample concentration (i.e, pooling multiple ant heads per sample) and recent

advancements in LC-MS/MS approaches to detect monoamines in insect brain tissue [98]. Additionally,

different metabolomic analyses may support discovery of specialized fungal effector metabolites as has

been observed in a fly-manipulating fungus and hypothesized in *Ophiocordyceps* [99,100 preprint].

However, by integrating transcriptomic data with the metabolite features we did identify, we

propose a scenario where the manipulated ant has elevated production of dopamine, with possible

increases of serotonin and decreases of octopamine. The fungus may also simultaneously suppress the

host's melanin and immunity pathways while metabolizing dopamine and its precursors for fungal

melanin production. Ants infected by *Ophiocordyceps* display enhanced locomotion/activity levels,

reduction of social behaviors, and a final summit latching behavior [12–16]. Elements of these behaviors have

already been associated with neurotransmitter signaling in ants, albeit phenotypic variation across species

and social or physiological context exists [54–59]. Intriguingly, aquatic gammarid crustaceans manipulated

by acanthocephalan worms display a phototactic summiting and clinging phenotype in the water column

associated with increased serotonin, decreased octopamine, and suppressed phenoloxidase-mediated

immunity [101–104]. In essence, our data indicate comparable biogenic amine level changes and immunity

suppression in manipulated *Camponotus,* which is in line with the premise that convergently evolved

manipulated summiting behaviors are established by overlapping mechanisms [5,105]. Such mechanisms are

likely specific to different behavioral manipulation phenotypes, but may include common players found

across phenotypes. For example, some parasitoid wasps induce a low-activity, defensive "bodyguard"

state in their insect hosts that has been hypothesized to be linked to increased, rather than decreased,

octopamine levels in the host [106]. Manipulators of vertebrate behavior are also thought to leverage

serotoninergic and dopaminergic processes to alter fish and mouse behaviors related to escape responses,

locomotion, and aggression [50,107–109]. How much of the alteration of neurotransmitter metabolism may be

a general response to fungal infection and how much results from specific manipulation by

Ophiocordyceps is difficult to pinpoint at this time. Possibly promoting both infection and manipulation,

cardol triene appears to be a fungal derived compound that could suppress ant immunity and increase

substrate availability for dopamine production. Also, *Ophiocordyceps* upregulated three putatively

secreted tyrosinases that possibly interact with host dopamine/melanin precursors. Although less typically

reported than their intracellular counterparts, extracellular fungal tyrosinases have also been found in

another fungus [110,111].

Changing glycerophospholipid metabolism during manipulation, largely centered on choline,

suggested fungal biosynthesis of cell membrane components and dysregulation of acetylcholine levels in

the ant. *Ophiocordyceps* may require increased choline-derived cell-membrane components used during

cell proliferation and membrane remodeling for structures such as lipid rafts or transmembrane proteins

important during infection, manipulation, and hyphal colonization of the host [83–86]. In some manipulated

ant samples, we detected elevated acetylcholine concentrations, which may correspond to reduced

acetylcholinesterase gene expression previously observed. Speculatively, transient high acetylcholine

levels could contribute to summiting and wandering behavior (i.e., increased locomotion), tremors, and

the hypercontracted biting that locks manipulated ants into place [12,15–17,20,25,87,88,112].

By inducing changes in neurotransmitters that regulate behavior in healthy ants, the fungus may

be impinging upon existing host systems to modify behavior rather than forcibly introducing novel

pathways [8,18]. Indeed, many of the hallmark phenotypes of infection by ant-manipulating *Ophiocordyceps*

are not so dissimilar from activities influenced by levels of dopamine, serotonin, octopamine, and

acetylcholine in eusocial insects: foraging/locomotor behavior (i.e., summiting and hyperactivity),

sociality (i.e., wandering and reduced communication), and the biting/aggression (the final "death grip"

bite) [47,48,59,49–51,54–58]. Changes in social behavior could not only be a symptom of disease, but contribute to

it as well. As socially isolated infected ants may less frequently exchange trophallactic fluid with

nestmates, these infected individuals would lose an important source of nutrition, communication, and

hormones [113] that may alter their metabolome. How exactly the fungus dysregulates physiological

pathways that influence behavior, such as insect hormone metabolism, and to what degree some level of

dysregulation is typical in many diseases is still unclear. These hypotheses can be further investigated by

functional tests (e.g., gene knockouts or dosing ants with small compounds) and LC-MS/MS validations

of key compounds using chemical standards and improved protocols.

Methods

Ant collection & husbandry: We used a single laboratory-housed wild-caught colony of *C.*

floridanus to produce all samples. This colony was collected in March 2021 in the University of Central

Florida arboretum and had several hundred individuals including minors, majors, and brood. Based on

methods in Will *et al.* 2020, we housed the colony in a 9.5 L plastic container (42 cm long × 29 cm wide)

lined with talcum powder (Fisher Scientific) and contained aluminum foil wrapped test-tubes (16 mm x

125 mm, Fisher Scientific) with moist cotton to serve as darkened, humid nest spaces. Ants fed ad libitum

on 15% sucrose solution, water, and sporadic frozen *Drosophila* feedings. Following collection, we

housed the ants indoors under 48 h constant light to prepare them for circadian entrainment to laboratory

conditions. Follow the "circadian reset," we transferred the colony to a climate-controlled incubator

(I36VL, Percival) cycling 12-12 h, light-dark, 28-20 C°, 65-80% RH daily cycles. Climate conditions

cycled in a ramping manner, with a 4 h ramp from lights-on to a 4 h hold at peak day conditions to a 4 h

ramp to lights-off. The colony housed under these conditions for approximately six weeks before

individuals were removed for the experiment, which was conducted in the same incubator.

Fungal strain collection and culturing: *O. camponoti-floridani* strain OJ2.1 was used to infect

all ants in this work. We isolated this fungal strain using mycelium extracted from the abdomen of a

recently manipulated and expired *C. floridanus* cadaver. This cadaver was found in the Ocala National

Forest, FL (29° 13' 41.7" N, 81° 41' 44.9" W), in March 2021 and permitted by letter of authorization

(United States Department of Agriculture, #2720-22). The isolation procedure, short ribosomal subunit

genetic validation, and culturing followed the protocols detailed in Will *et al.* 2020. In preparation for

experimental infections, we grew fungal cells for 10 days in T25 cell culture flasks with 20 mL Grace's

medium (Gibco) supplemented with 2.5% fetal bovine serum (Gibco) ("GF" medium), shaken at 50 rpm

in 28 C°. We then filtered the culture using two-layers of 12-ply gauze (Curity) to separate yeast-like

blastospores from hyphal cells. We adjusted the filtered blastospores to a working concentration of

approximately 3×10^6 cells/mL (optical density 660 nm = ca. 0.5) prior to injection.

Experimental ant infections: Prior to *Ophiocordyceps* infection and sham injection, we

separated ants from their colony into treatment groups. For the infection treatment, we gathered 51 ants,

of which we injected 41 individuals. For the control group, which received sham injections, we collected

38 ants, of which we sham-injected 30 individuals. The additional ants in each group did not receive

treatment other than a painted dot on their abdomens (Testors paint) to be able to recognize them. Our

previous experiments have shown that such a small group of untreated ants improves the recovery of the

treated group during the first few days of the experiment [14]. One day before infection treatments, we

removed the sugar and water from the ant box to reduce the amount of fluid in the ants and ease the

injection procedure. We injected each ant with 0.5 µL *Ophiocordyceps* blastospores (infection, n = 41) or

0.5 µL of GF (sham, n = 30) as detailed in Will *et al.* 2020. We placed the treated ants and their painted

"caretaker ants" in plaster-lined boxes with biting structures (*Tillandsia usneoides* tied to wooden sticks

with twine) to be able to observe *Ophiocordyceps*-induced summiting behavior.

Behavioral observations: After three dpi we removed ants that did not survive the treatment.

Starting day four, we screened for manipulated ants and deaths twice per day, at zeitgeber time (ZT) 0,

the beginning of subjective day (i.e., lights-on), and ZT 6, the mid-point of subjective day. At ten dpi, we

removed the painted caretaker ants as previous work has observed *C. floridanus* to aggressively interfere

with manipulated nestmates, presumably as a component of their social immunity and nest hygiene

[14,24,114]. Starting 15 dpi, we began more frequent opportunistic checks as behavioral manipulations of *C.*

floridanus that involve summiting had previously been observed starting 19 dpi in the laboratory [24]. After

discovering the first manipulation at 21 dpi, we checked multiple times, daily beginning at ZT 20 and

ending at ZT 14 (on the next subjective day). We selected this observation window to improve the odds of

collecting manipulations close to their onset since our previous laboratory infections with *O. camponoti-*

floridani showed that manipulated summiting synchronized to the ants' subjective pre-dawn hours [24].

Summiting behavior is characterized by infected individuals clinging and biting *Tillandsia* suspended in

the experiment box. The behavior has never been observed in sham-treated individuals. We screened non-

manipulated mortalities for fungal infection by crushing tissue from the dead ant in 50 µL of water,

briefly centrifuging, and examining 15 µL of the supernatant under a microscope. Notable levels of fungal

cells were always and only seen in the *Ophiocordyceps* treatment ants. We analyzed and plotted survival

data using the R packages survival and survminer in R (v 4.1) with R Studio (v. 2021.09.2) [115–118].

Sample collection for LC-MS/MS: We collected live manipulated ants displaying characteristic

summiting biting behavior for LC-MS/MS immediately upon discovery (n = 20). We subsequently

collected dpi-matched sham ants at ZT-21.5, as this was the most common time at which manipulated

ants were collected (n = 20) (see Results and Discussion). Manipulated ants found after death were

recorded for survival statistics but not used in the LC-MS/MS analyses (n = 3). For manipulated ant

collection, we held ants with forceps, clipped the biting substrate just above and below the manipulated

ant and submerged the ant in liquid nitrogen briefly until frozen. The sample was then placed in a pre-

chilled, sterile 2 mL microcentrifuge tube (USA Scientific) kept in liquid nitrogen until long-term storage

at -80 C°. We collected sham ant samples in a similar manner, lifting them from the experiment box with

forceps.

We also collected three blank samples at the beginning, middle, and end of manipulated ant

sampling, i.e., after the first, 10^{th}, and 20^{th} manipulated ants were found (n = 3). We made blanks by

briefly touching the forceps to the floor of the sham treatment housing boxes, submerging them in liquid

nitrogen, and swirling them in a microcentrifuge tube. We cleaned all tools and surfaces with 70% ethanol

and deionized water before sampling and between sample types.

We later prepared collected samples for LC-MS/MS by isolating ant heads. Each frozen ant was

quickly processed on a plastic petri dish (USA Scientific) on a bed of dry ice. Using forceps, we removed

biting substrate from the ant's mandibles if needed, snapped the head off, and placed it in a new

microcentrifuge tube. For blank samples, we dragged the forceps along the petri dish and swirled the

forceps inside the original collection tube. We cleaned the tools and petri dish with 70% ethanol and

deionized water between individuals and used a new petri dish between sample types. All samples and

tubes were kept constantly chilled on dry ice or in liquid nitrogen.

LC-MS/MS processing of ant head samples: All samples were sent on dry ice to the West

Coast Metabolomics Center (WCMC, University of California, Davis) for LC-MS/MS analysis. The

WCMC processed samples in three LC-MS/MS protocols broadly tailored toward different chemistries:

"biogenic amines," "polyphenols and flavonoids," and "complex lipids." From the 20 samples collected

per infection and sham treatments, the WCMC collected polyphenol data for ten individual ant heads.

They used the remaining ten heads per treatment for biphasic extractions, with each individual head used

for both biogenic amines and lipids. Blank-collection samples were shared across the three LC-MS/MS

chemistries. The WCMC produced quality-control (QC) pool samples consisting of all samples used per

protocol and additional blank-extraction controls (and therefore distinct from our experimental blanks

made during sample collection). These WCMC control samples were run at the beginning and every ten

samples thereafter per protocol. All LC-MS/MS protocols collected MS/MS data in a top-four data-

dependent-acquisition (DDA) mode in both electrospray ionization (ESI) positive and negative modes.

The WCMC processed samples for biogenic amines for use in hydrophilic interaction

chromatography (HILIC) with quadrapole time of flight (Q-TOF) mass spectrometry. Homogenized ant

head samples were extracted with a methanol and methyl-tert-butyl ether protocol [119]. The aqueous phase

was used for biogenic amines LC-MS/MS and the organic phase for lipids (see below). Extractions were

mixed with internal standards for normalization according to WCMC protocols to a final volume of 100

μL (Supplementary Data S3).

Samples for lipids were processed for use in charged surface hybrid (CSH) chromatography with

Q-TOF mass spectrometry. The organic phase per biphasic extraction (shared with biogenic amines, see

above) was taken for lipids LC-MS/MS and mixed with internal standards according to WCMC protocols

to a final resuspension volume of 110 μL (Supplementary Data S3).

Samples for polyphenols were processed for use in pentafluorophenyl (PFP) chromatography

with QExactive HF (QE) mass spectrometry. Homogenized ant head samples were extracted in a similar

fashion as described above and resuspended in a final 100 μL volume with internal standards

(Supplementary Data S3). Further details for all three LC-MS/MS protocols are provided in

Supplementary Methods.

Metabolomic data processing & feature filtering: The WCMC performed initial processing

steps on all samples to produce metabolomic feature lists consisting of mass/charge ratio (*m/z*), retention

time (rt), and value of quantification ion peaks. They aligned raw data with MS-DIAL (v 4.7.0) [120],

subtracted signals detected in the blank-extractions produced in the core facility, removed duplicate

signals and isotopes, normalized samples by internal standards, and assigned compound and adduct

annotations based on manual MS/MS matching in Massbank, Metlin, and NIST14 databases and, for

lipids, with LipidBlast (v 68) [121]. The resulting data consisted of several thousand features per dataset:

biogenic amines (5605 total with 141 annotated), polyphenols (6178 total with 158 annotated), and lipids

(7021 with 517 annotated) (Supplementary Data S4).

We applied additional feature quality control and data reduction methods to create a final list of

metabolite features to investigate. Features that had greater than 50% missing values within both sham

and infection treatments were removed. Similarly, we removed features with both median peak values

(sham and infection) less than double that of median blank sample peaks. Following removal based on

missingness and peak values, we removed features that we determined to have signals too variable for

robust analysis based on their coefficient of variation (CV%, or relative standard deviation). For the

biogenic amines and lipid datasets that had successful QC pool data returned, we applied a CV%

maximum threshold to remove peaks that varied greatly due to technical variation between QC pool

injections. We selected the CV% threshold as the lowest (i.e., strictest) value between 30% and a dataset

specific value determined from the distribution of CV% values. We calculated this data-specific value as

the 75^{th}-quartile + 1.5*interquartile-range (IQR), as is used to identify extreme values or outliers in data

distributions (e.g., boxplots). For the lipids, the 1.5*IQR method produced CV 25% as the cutoff,

removing 5.7% of features. For biogenic amines, the 1.5*IQR CV% was greater than 30%; therefore,

30% was used, removing 6.6% of features. The WCMC reported that all polyphenol QC pools failed on

acquisition. Therefore, we ranked each feature by its lowest CV% (i.e., from either CV%-sham or CV%-

manipulated) and removed the highest 6.1% of features. We selected this 6.1% value because it was the

mean of the CV% removal rates of the biogenic amine and lipid datasets. Although we cannot fully

discount biological variation in biogenic amine peak values within a treatment, by selecting the lowest

CV% per feature (whether from the sham or infection treatment), we sought to remove only features that

had high CV% regardless of treatment. After these filtering steps, we continued with 3268 biogenic amine

features (58% of initially collected), 4878 polyphenol features (79% of initial), and 6305 lipid features

(90% of initial).

We further applied a putative adduct calling step to consolidate unannotated features that may

represent different adducts of the same compound. Per LC-MS/MS dataset, we divided features by

treatment and ESI mode, and processed each of these subsets with Binner (v 1.0.0) [122]. Binner compares

metabolite features "binned" by rt and calls putative adduct relationships between these features based on

a table of possible adducts, feature *m/z*, and correlation of peak values across samples. Data were split by

treatment to avoid possibly conflating signals for different compounds found in only one of the treatments

as adducts of each other. Data were natural log transformed, no additional deisotoping step was

preformed, the mass tolerance was set to 0.005 Da (biogenic amines and lipids) or 0.003 Da

(polyphenols), and default settings were used elsewhere. For biogenic amines and polyphenols, we

searched for adducts used previously used by Binner [122]. For lipids, we supplemented this adduct table

with an additional adduct set publicly available online [123]. We then reunited data subsets to create feature

lists that included all peaks that were labeled as primary ions or remained unannotated in either treatment

group. In effect, this step only removed peaks that were determined to be putative adduct peaks in both

treatments, resulting in the final processed datasets for analysis of: 2955 biogenic amines (90% of QC

filtered peaks remained), 4315 polyphenols (88% of QC filtered peaks), and 4924 lipids (78% of QC

filtered peaks) (Supplementary Data S1).

682   **Metabolomic data analysis & feature selection:** To select a single set of top-features of interest

combined across all three datasets, we applied three statistical approaches (Fig. 2A). For features with

values equal to zero, we imputed zeros as 20% of the minimum value measured for that feature. Each

analysis we preformed was done independently per LC-MS/MS dataset (biogenic amines, lipids, and

polyphenols). We determined if a feature represented a differentially abundant metabolite (DAM) with a

t-test (corrected FDR, $p \le 0.05$) and minimum ±1.5-fold change in abundance between sham and infection

treatments. Principal component analyses always indicated that only PC1 separated samples by treatment

type (see Results). As such, we noted features ranked in the upper 50% of PC1 absolute loading values.

Given that our choice of threshold was defined as half of the data, we did not rely on this test solely as

merit for discussing a feature, but rather as a way to add confidence to results of other analyses. Both t-

tests and PCAs were performed through the MetaboAnalyst interface (v 5.0) [28]. Finally, we selected

features "confirmed" as "important" by Boruta (v 7.0.0, R package) random-forest based analyses with

900 maxRuns and 70,000 ntree settings [27]. Boruta selects "important" features by z-scores of mean

decrease accuracy in decision trees in comparison to randomized versions of the data [27]. This offers an

improvement over standard random-forest analyses for high dimensional data such as in metabolomics,

increasing the stability and robustness of the features selected [124].

Additionally, we created co-abundance network modules of metabolite features per LC-MS/MS

dataset using the WGCNA method [46] (Fig. 2A). Although WGCNA was developed primarily for gene

expression data, this technique has been used and evaluated for metabolomic data as well [42,125,126].

Heightened activity in a metabolic pathway could lead to many compounds increasing. But

simultaneously, their precursors may become depleted. Therefore, correlations within a network could be

both positive and negative. As such, we constructed unsigned WGCNA modules. All WGCNA modules

were constructed on feature data without imputing zeros, based on the biweight midcorrelation to

compare between treatment types (e.g., corFnc="bicor", robustY=F, maxPOutlier=0.05) (R package

WGCNA, v 1.67). Soft power thresholds were selected according to program recommendations as

follows: biogenic amines – 3, polyphenols – 10, and lipids – 11.

We analyzed different groups of selected features for metabolic pathway enrichments. We formed

feature sets that combined data across all three LC-MS/MS datasets based on individual top-feature

selection analyses (DAM t-tests, PCAs, and Boruta) or their presence only in manipulation samples.

Dataset-specific network modules were only analyzed with background features from the LC-MS/MS

dataset they belonged to (i.e., amines, polyphenols, or lipids) and if the module was correlated to

treatment (Student correlation $\geq |0.5|$, $p \leq 0.05$) (Fig. 2A).

We preformed Mummichog analyses on all the above feature sets [30]. This process matched 1,523

unique metabolite features to 6,725 KEGG annotated compounds. For enrichment analyses combining

features across datasets, we additionally used a technique based on GSEA ranking features by t-scores

returned by t-tests [127]. In network module pathway enrichment analyses, only Mummichog was used as it

sets a hard threshold to define a group of interest versus the background. Membership within or out of a

network module was simply defined and could be used for this division. All enrichment analyses were

conducted through the MetaboAnalyst interface with the following settings: mixed ion mode, 3 ppm mass

tolerance, retention time included, primary ions not enforced, all adducts tested, default currency

metabolites, and conducted once for both the *Drosophila melanogaster* and *Saccharomyces cerevisiae*

KEGG metabolic pathway and compound databases (Fig. 2A) [28]. As we sought to characterize the

composite metabolome of manipulated ant heads containing both insect and fungal tissue, we chose both

a well-studied insect and fungal model organism to improve our odds of matching annotations. We did

not use results from one database or the other to discriminate between a host (insect) or parasite (fungus)

signal. Neither of the databases would represent the full metabolic repertoire of their respective model

organisms and combining them could offer more complete data for comparison. Furthermore, we wished

to consider the possibility that due to the relationship of the host and parasite, either organism may be

producing metabolites canonically associated with the other as part of its strategy in this antagonism.

We selected significantly enriched pathways based on one of the hypergeometric enrichment

results reported by MetaboAnalyst. When using Mummichog only (WGCNA module feature sets), we

used the gamma-null adjusted p-value (≤ 0.05). When including a GSEA as a joint analysis (top-features

or features present in manipulation only), we used the combined p-value (≤ 0.05), calculated with

MetaboAnalyst using the Fisher method. Of the overrepresented pathways selected by the enrichment

analyses ($p \leq 0.05$), we further imposed thresholds for: (*i*) a minimum of three selected features in the

pathway, (*ii*) 1.25-fold enrichment, and (*iii*) three differentially expressed genes (DEGs) (see below)

encoding associated enzymes to discard weak signals that would likely be difficult to interpret

biologically in a multiomic pathway-level context (Fig. 2A). For cases when a pathway was selected

using both *D. melanogaster* and *S. cerevisiae* KEGG databases, a mean p-value and fold-enrichment is

reported.

Multiomics integration of genomic and transcriptomic data: We interpreted our LC-MS/MS

data in light of previously collected transcriptomic data when possible [24]. This RNAseq data was collected

with a similar experimental framework, including comparisons of gene expression of both *C. floridanus*

and *O. camponoti-floridani* between a control (uninfected ant or fungal culture) and manipulated ant

samples. Identification of DEGs (Cuffdiff analysis corrected FDR $p \leq 0.05$, minimum two-fold change,

minimum 4 RPKM) and relevant statistics are reported in Will *et al.* 2020 [128]. For each enriched

metabolic pathway, we retrieved gene sequence data for enzymes associated with that pathway in KEGG

(by enzyme nomenclature EC number) using KEGGREST (v 1.30.1) [129]. For each enzyme EC, we

extracted the first representative gene listed for *D. melanogaster* and *S. cerevisiae*. We searched for

reciprocal best matches to those genes in both the *C. floridanus* and *O. camponoti-floridani* genomes

using Proteinortho (v 5) with default settings [130]. While this non-exhaustive approach would not identify

every possible *C. floridanus* or *O. camponoti-floridani* gene with homology to known enzymes of

interest, it did assist with initially selecting pathway enrichments for deeper analysis. Per pathway, we

required that at least three DEGs combined from both transcriptomes of the ant and fungus had homologs

to the enzymes associated with that pathway. This selection criterion followed the pathway-level thinking

to require at least three contributing metabolite features matched by Mummichog to consider an

enrichment result for discussion. Metabolic pathway relationships were based on KEGG data and

supplemented by other sources where cited.

Data Availability

Genomic and transcriptomic data referenced here are available in Will *et al.* 2020 and the

GenBank accessions reported therein. Processed metabolomic data are given with this publication as

supplementary information. Raw LC-MS/MS peak data have been uploaded to MassIVE (UPLOAD URL

PENDING).

References

1.   Libersat, F., Kaiser, M. & Emanuel, S. Mind Control: How Parasites Manipulate Cognitive

Functions in Their Insect Hosts. *Front. Psychol.* **9**, 572 (2018).

2.   Heil, M. Host Manipulation by Parasites: Cases, Patterns, and Remaining Doubts. *Front. Ecol.*

Evol. **4**, 80 (2016).

3.   Thomas, F., Poulin, R. & Brodeur, J. Host manipulation by parasites: A multidimensional

phenomenon. *Oikos* **119**, 1217–1223 (2010).

4.   Moore, J. An overview of parasite-induced behavioral alterations - and some lessons from bats. *J.*

Exp. Biol. **216**, 11–7 (2013).

5.   de Bekker, C., Beckerson, W. C. & Elya, C. Mechanisms behind the Madness: How Do Zombie-

Making Fungal Entomopathogens Affect Host Behavior To Increase Transmission? *MBio* (2021)

doi:10.1128/mBio.01872-21.

6.   de Bekker, C. *Ophiocordyceps*–ant interactions as an integrative model to understand the

molecular basis of parasitic behavioral manipulation. *Curr. Opin. Insect Sci.* **33**, 19–24 (2019).

7.   de Bekker, C., Merrow, M. & Hughes, D. P. From behavior to mechanisms: An integrative

approach to the manipulation by a parasitic fungus (*Ophiocordyceps unilateralis s.l.*) of its host

ants (*Camponotus spp.*). *Integr. Comp. Biol.* **52**, 166–176 (2014).

8.   Lovett, B., St. Leger, R. J. & de Fine Licht, H. H. Going gentle into that pathogen-induced

goodnight. *J. Invertebr. Pathol.* **174**, 107398 (2020).

9.   Loreto, R. G. *et al.* Evidence for convergent evolution of host parasitic manipulation in response to

environmental conditions. *Evolution (N. Y).* 211144 (2018) doi:10.1101/211144.

10.   Will, I., Linehan, S., Jenkins, D. G. & de Bekker, C. Natural history and ecological effects on the

establishment and fate of Florida carpenter ant cadavers infected by the parasitic manipulator

Ophiocordyceps camponoti-floridani. Funct. Ecol. **00**, 1–14 (2022).

11.   Andriolli, F. S. *et al.* Do zombie ant fungi turn their hosts into light seekers? *Behav. Ecol.* **30**, 609–

616 (2019).

12.   Andersen, S. B. *et al.* The Life of a Dead Ant: The Expression of an Adaptive Extended

Phenotype. *Am. Nat.* **174**, 424–433 (2009).

13.   Hughes, D. P. *et al.* Behavioral mechanisms and morphological symptoms of zombie ants dying

from fungal infection. *BMC Ecol.* **11**, 13 (2011).

14.   Trinh, T., Ouellette, R. & de Bekker, C. Getting lost: the fungal hijacking of ant foraging

behaviour in space and time. *Anim. Behav.* **181**, 165–184 (2021).

15.   de Bekker, C. *et al.* Gene expression during zombie ant biting behavior reflects the complexity

800   underlying fungal parasitic behavioral manipulation. *BMC Genomics* **16**, 620 (2015).

801   16.   Pontoppidan, M. B., Himaman, W., Hywel-Jones, N. L., Boomsma, J. J. & Hughes, D. P.

802   Graveyards on the move: The spatio-temporal distribution of dead *Ophiocordyceps*-infected ants.

803   *PLoS One* **4**, e4835 (2009).

804   17.   Sakolrak, B., Blatrix, R., Sangwanit, U. & Kobmoo, N. Experimental infection of the ant

805   *Polyrhachis furcata* with *Ophiocordyceps* reveals specificity of behavioural manipulation. *Fungal*

806   *Ecol.* **33**, 122–124 (2018).

807   18.   Adamo, S. A. Turning your victim into a collaborator: exploitation of insect behavioral control

808   systems by parasitic manipulators. *Curr. Opin. Insect Sci.* **33**, 25–29 (2019).

809   19.   Kobmoo, N. *et al.* A genome scan of diversifying selection in *Ophiocordyceps* zombie-ant fungi

810   suggests a role for enterotoxins in co-evolution and host specificity. *Mol. Ecol.* **27**, 3582–3598

811   (2018).

812   20.   Mangold, C. A., Ishler, M. J., Loreto, R. G., Hazen, M. L. & Hughes, D. P. Zombie ant death grip

813   due to hypercontracted mandibular muscles. *J. Exp. Biol.* **222**, jeb200683 (2019).

814   21.   Loreto, R. G. & Hughes, D. P. The metabolic alteration and apparent preservation of the zombie

815   ant brain. *J. Insect Physiol.* **118**, 103918 (2019).

816   22.   Zheng, S. *et al.* Specialist and Generalist Fungal Parasites Induce Distinct Biochemical Changes in

817   the Mandible Muscles of Their Host. *Int. J. Mol. Sci.* **20**, (2019).

818   23.   de Bekker, C. & Das, B. Hijacking time: How *Ophiocordyceps* fungi could be using ant host

819   clocks to manipulate behavior. *Parasite Immunol.* **44**, e12909 (2022).

820   24.   Will, I. *et al.* Genetic Underpinnings of Host Manipulation by *Ophiocordyceps* as Revealed by

821   Comparative Transcriptomics. *G3 (Bethesda).* **10**, 2275–2296 (2020).

822   25.   de Bekker, C. *et al.* Species-specific Ant Brain Manipulation by a Specialized Fungal Parasite.

823   *BMC Evol. Biol.* **14**, 166 (2014).

26.   Fredericksen, M. A. *et al.* Three-dimensional visualization and a deep-learning model reveal

complex fungal parasite networks in behaviorally manipulated ants. *Proc. Natl. Acad. Sci.* **114**,

201711673 (2017).

27. Kursa, M. B. & Rudnicki, W. R. Feature Selection with the Boruta Package. *J. Stat. Softw.* **36**, 1–

13 (2010).

28. Pang, Z. *et al.* MetaboAnalyst 5.0: narrowing the gap between raw spectra and functional insights.

Nucleic Acids Res. **49**, W388–W396 (2021).

29. Kanehisa, M. & Goto, S. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res.*

28, 27 (2000).

30. Li, S. *et al.* Predicting Network Activity from High Throughput Metabolomics. *PLOS Comput.*

Biol. **9**, e1003123 (2013).

31. Prior, K. F. *et al.* Synchrony between daily rhythms of malaria parasites and hosts is driven by an

essential amino acid. *Wellcome Open Res.* **6**, (2021).

32. Pellon, A. *et al.* Role of Cellular Metabolism during Candida-Host Interactions. *Pathog. 2022,*

Vol. 11, Page 184 **11**, 184 (2022).

33. Ene, I. V., Brunke, S., Brown, A. J. P. & Hube, B. Metabolism in Fungal Pathogenesis. *Cold*

Spring Harb. Perspect. Med. **4**, (2014).

34. Lee, D.-K. *et al.* Comparison of primary and secondary metabolites for suitability to discriminate

the origins of Schisandra chinensis by GC/MS and LC/MS. *Food Chem.* **141**, 3931–3937 (2013).

35. Bing, X. *et al.* Unravelling the relationship between the tsetse fly and its obligate symbiont

Wigglesworthia□: transcriptomic and metabolomic landscapes reveal highly integrated

physiological networks. *Proc. R. Soc. B Biol. Sci.* **284**, 20170360 (2017).

36. Wangi, Y. *et al.* Symbiotic bracovirus of a parasite manipulates host lipid metabolism via

tachykinin signaling. *PLOS Pathog.* **17**, e1009365 (2021).

37. Kitowski, A. & Bernardes, G. J. L. A Sweet Galactose Transfer: Metabolic Oligosaccharide

Engineering as a Tool To Study Glycans in Plasmodium Infection. *ChemBioChem* **21**, 2696–2700

(2020).

38.   Wang-Eckhardt, L., Bastian, A., Bruegmann, T., Sasse, P. & Eckhardt, M. Carnosine synthase

deficiency is compatible with normal skeletal muscle and olfactory function but causes reduced

olfactory sensitivity in aging mice. *J. Biol. Chem.* **295**, 17100–17113 (2020).

39.   Jukić, I. *et al.* Carnosine, Small but Mighty—Prospect of Use as Functional Ingredient for

Functional Food Formulation. *Antioxidants 2021, Vol. 10, Page 1037* **10**, 1037 (2021).

40.   Dolan, E. *et al.* Comparative physiology investigations support a role for histidine-containing

dipeptides in intracellular acid–base regulation of skeletal muscle. *Comp. Biochem. Physiol. Part*

A Mol. Integr. Physiol. **234**, 77–86 (2019).

41.   Bellia, F., Vecchio, G., Cuzzocrea, S., Calabrese, V. & Rizzarelli, E. Neuroprotective features of

carnosine in oxidative driven diseases. *Mol. Aspects Med.* **32**, 258–266 (2011).

42.   Bartzis, G. *et al.* Estimation of metabolite networks with regard to a specific covariable:

applications to plant and human data. *Metabolomics* **13**, 129 (2017).

43.   Carlson, M. R. J. *et al.* Gene connectivity, function, and sequence conservation: Predictions from

modular yeast co-expression networks. *BMC Genomics* **7**, 1–15 (2006).

44.   Stuart, J. M., Segal, E., Koller, D. & Kim, S. K. A gene-coexpression network for global discovery

of conserved genetic modules. *Science (80-.).* **302**, 249–255 (2003).

45.   Gillis, J. & Pavlidis, P. "Guilt by Association" Is the Exception Rather Than the Rule in Gene

Networks. *PLOS Comput. Biol.* **8**, e1002444 (2012).

46.   Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network analysis.

BMC Bioinformatics **9**, 559 (2008).

47.   Kamhi, J. F., Arganda, S., Moreau, C. S. & Traniello, J. F. A. Origins of aminergic regulation of

behavior in complex insect social systems. *Front. Syst. Neurosci.* **11**, 74 (2017).

48.   Verlinden, H. Dopamine signalling in locusts and other insects. *Insect Biochem. Mol. Biol.* **97**, 40–

52 (2018).

49.   Aonuma, H. & Watanabe, T. Changes in the content of brain biogenic amine associated with early

colony establishment in the Queen of the ant, *Formica japonica*. *PLoS One* **7**, e43377–e43377

(2012).

50.   Adamo, S. A., Linn, C. E. & Beckage, N. E. Parasites: evolution's neurobiologists. *J. Exp. Biol.*
216, 3–10 (2013).

51.   Adamo, S. A. Norepinephrine and octopamine: linking stress and immune function across phyla.
ISJ **5**, 12–19 (2008).

52.   González-Santoyo, I. & Córdoba-Aguilar, A. Phenoloxidase: a key component of the insect
immune system. *Entomol. Exp. Appl.* **142**, 1–16 (2012).

53.   Geun, S. K., Nalini, M., Kim, Y. & Lee, D. W. Octopamine and 5-hydroxytryptamine mediate
hemocytic phagocytosis and nodule formation via eicosanoids in the beet armyworm, *Spodoptera*
exigua. *Arch. Insect Biochem. Physiol.* **70**, 162–176 (2009).

54.   Szczuka, A. *et al.* The effects of serotonin, dopamine, octopamine and tyramine on behavior of
workers of the ant *Formica polyctena* during dyadic aggression tests. *Acta Neurobiol. Exp. (Wars).*
73, 495–520 (2013).

55.   Aonuma, H. Serotonergic control in initiating defensive responses to unexpected tactile stimuli in
the trap-jaw ant *Odontomachus kuroiwae*. *J. Exp. Biol.* **223**, (2020).

56.   Friedman, D. A. *et al.* The Role of Dopamine in the Collective Regulation of Foraging in
Harvester Ants. *iScience* **8**, 283 (2018).

57.   Muscedere, M. L., Johnson, N., Gillis, B. C., Kamhi, J. F. & Traniello, J. F. A. Serotonin
modulates worker responsiveness to trail pheromone in the ant *Pheidole dentata*. *J. Comp.*
Physiol. A Neuroethol. Sensory, Neural, Behav. Physiol. **198**, 219–227 (2012).

58.   Yakovlev, I. K. Effects of Octopamine on Aggressive Behavior in Red Wood Ants. *Neurosci.*
Behav. Physiol. 2018 483 **48**, 279–288 (2018).

59.   Boulay, R., Soroker, V., Godzinska, E. J., Hefetz, A. & Lenoir, A. Octopamine reverses the
isolation-induced increase introphallaxis in the carpenter ant *Camponotus fellah*. *Journgal Exp.*
Biol. **203**, 513–520 (2000).

60.   Dakshinamurti, K. *et al.* Vitamin B6: Effects of Deficiency, and Metabolic and Therapeutic

Functions. *Handb. Famine, Starvation, Nutr. Deprivation* 1–23 (2017) doi:10.1007/978-3-319-

40007-5_81-1.

61.   Neckameyer, W. S. & White, K. A single locus encodes both phenylalanine hydroxylase and

tryptophan hydroxylase activities in Drosophila. *J. Biol. Chem.* **267**, 4199–4206 (1992).

62.   Coleman, C. M. & Neckameyer, W. S. Serotonin synthesis by two distinct enzymes in *Drosophila*

melanogaster. *Arch. Insect Biochem. Physiol.* **59**, 12–31 (2005).

63.   Neckameyer, W. S., Coleman, C. M., Eadie, S. & Goodwin, S. F. Compartmentalization of

neuronal and peripheral serotonin synthesis in Drosophila melanogaster. *Genes, Brain Behav.* **6**,

756–769 (2007).

64.   Sugumaran, M. & Barek, H. Critical Analysis of the Melanogenic Pathway in Insects and Higher

Animals. *Int. J. Mol. Sci.* **17**, (2016).

65.   Wang, Y. *et al.* Activation of *Aedes aegypti* prophenoloxidase-3 and its role in the immune

response against entomopathogenic fungus. *Insect Mol. Biol.* **26**, 552 (2017).

66.   Nosanchuk, J. D. & Casadevall, A. Impact of melanin on microbial virulence and clinical

resistance to antimicrobial compounds. *Antimicrob. Agents Chemother.* **50**, 3519–3528 (2006).

67.   Nosanchuk, J. D., Stark, R. E. & Casadevall, A. Fungal melanin: What do we know about

structure? *Front. Microbiol.* **6**, 1463 (2015).

68.   Zhuang, J. X. *et al.* Irreversible Competitive Inhibitory Kinetics of Cardol Triene on Mushroom

Tyrosinase. *J. Agric. Food Chem.* **58**, 12993–12998 (2010).

69.   Gorman, M. J. & Arakane, Y. Tyrosine hydroxylase is required for cuticle sclerotization and

pigmentation in *Tribolium castaneum*. *Insect Biochem. Mol. Biol.* **40**, 267–273 (2010).

70.   Piao, S. *et al.* Crystal structure of a clip-domain serine protease and functional roles of the clip

domains. *EMBO J.* **24**, 4404 (2005).

71.   Ramirez, J. L., Muturi, E. J., Dunlap, C. & Rooney, A. P. Strain-specific pathogenicity and

subversion of phenoloxidase activity in the mosquito *Aedes aegypti* by members of the fungal

entomopathogenic genus *Isaria*. *Sci. Reports 2018 81* **8**, 1–12 (2018).

72.   Feng, P., Shang, Y., Cen, K. & Wang, C. Fungal biosynthesis of the bibenzoquinone oosporein to

930   evade insect immunity. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 11365–11370 (2015).

931   73.   Walter, A. *et al.* Glycerophosphocholine is elevated in cerebrospinal fluid of Alzheimer patients.

932   *Neurobiol. Aging* **25**, 1299–1303 (2004).

933   74.   Hoxmeier, J. C. *et al.* Analysis of the metabolome of *Anopheles gambiae* mosquito after exposure

934   to *Mycobacterium ulcerans. Sci. Reports 2015 51* **5**, 1–8 (2015).

935   75.   Alecu, I. & Bennett, S. A. L. Dysregulated lipid metabolism and its role in α-synucleinopathy in

936   Parkinson's disease. *Front. Neurosci.* **13**, 328 (2019).

937   76.   Li, Z. & Vance, D. E. Thematic Review Series: Glycerolipids. Phosphatidylcholine and choline

938   homeostasis. *J. Lipid Res.* **49**, 1187–1194 (2008).

939   77.   Dawaliby, R. *et al.* Phosphatidylethanolamine Is a Key Regulator of Membrane Fluidity in

940   Eukaryotic Cells. *J. Biol. Chem.* **291**, 3658 (2016).

941   78.   Cassilly, C. D. & Reynolds, T. B. PS, It's Complicated: The Roles of Phosphatidylserine and

942   Phosphatidylethanolamine in the Pathogenesis of *Candida albicans* and Other Microbial

943   Pathogens. *J. Fungi* **4**, (2018).

944   79.   Batrakov, S. G., Konova, I. V., Sheichenko, V. I., Esipov, S. E. & Galanina, L. A. Two unusual

945   glycerophospholipids from a filamentous fungus, *Absidia corymbifera. Biochim. Biophys. Acta -*

946   *Mol. Cell Biol. Lipids* **1531**, 169–177 (2001).

947   80.   Pokotylo, I. *et al.* The plant non-specific phospholipase C gene family. Novel competitors in lipid

948   signalling. *Prog. Lipid Res.* **52**, 62–79 (2013).

949   81.   Keyhani, N. O. Lipid biology in fungal stress and virulence: Entomopathogenic fungi. *Fungal*

950   *Biol.* **122**, 420–429 (2018).

951   82.   Menezes, T. A. de *et al.* Unraveling the Secrets of a Double-Life Fungus by Genomics:

952   *Ophiocordyceps australis* CCMB661 Displays Molecular Machinery for Both Parasitic and

953   Endophytic Lifestyles. *J. Fungi 2023, Vol. 9, Page 110* **9**, 110 (2023).

83.   Shyu, P. *et al.* Membrane phospholipid alteration causes chronic ER stress through early

degradation of homeostatic ER-resident proteins. *Sci. Reports 2019 91* **9**, 1–15 (2019).

84.   van der Veen, J. N. *et al.* The critical role of phosphatidylcholine and phosphatidylethanolamine

metabolism in health and disease. *Biochim. Biophys. Acta - Biomembr.* **1859**, 1558–1572 (2017).

85.   Markham, P., Robson, G. D., Bainbridge, B. W. & Trinci, A. P. J. Choline: Its role in the growth

of filamentous fungi and the regulation of mycelial morphology. *FEMS Microbiol. Rev.* **10**, 287–

300 (1993).

86.   Wang, J. *et al.* Phospholipid homeostasis plays an important role in fungal development, fungicide

resistance and virulence in *Fusarium graminearum. Phytopathol. Res.* **1**, 1–12 (2019).

87.   Gauthier, M. *Insect Nicotinic Acetylcholine Receptors: State of the Art on Insect Nicotinic*

Acetylcholine Receptor Function in Learning and Memory. (2010).

88.   Grünewald, B. & Siefert, P. Acetylcholine and Its Receptors in Honeybees: Involvement in

Development and Impairments by Neonicotinoids. *Insects* **10**, (2019).

89.   Colovic, M. B., Krstic, D. Z., Lazarevic-Pasti, T. D., Bondzic, A. M. & Vasic, V. M.

Acetylcholinesterase Inhibitors: Pharmacology and Toxicology. *Curr. Neuropharmacol.* **11**, 315

(2013).

90.   Fournier, D., Bride, J.-M., Hoffmann, F. & Karch, F. Acetylcholinesterase two types of

modifications confer resistance to insecticide. *J. Biol. Chem.* **267**, 14270–14274 (1992).

91.   Singh, K. D. *et al.* Biochemical efficacy, molecular docking and inhibitory effect of 2, 3-

dimethylmaleic anhydride on insect acetylcholinesterase. *Sci. Reports 2017 71* **7**, 1–11 (2017).

92.   Quinn, D. M. Acetylcholinesterase: Enzyme Structure, Reaction Dynamics, and Virtual Transition

States. *Chem. Reiviews* **87**, 955–979 (1987).

93.   Kim, Y. H. & Lee, S. H. Which acetylcholinesterase functions as the main catalytic enzyme in the

Class Insecta? *Insect Biochem. Mol. Biol.* **43**, 47–53 (2013).

94.   Lee, Y. S. *et al.* Characterization of GAR-2, a Novel G Protein-Linked Acetylcholine Receptor

from *Caenorhabditis elegans. J. Neurochem.* **75**, 1800–1809 (2000).

95.   Pflüger, H. J. & Duch, C. Dynamic neural control of insect muscle metabolism related to motor

behavior. *Physiology* **26**, 293–303 (2011).

96.   Johansen, J., Halpern, M. E., Johansen, K. M. & Keshishian, H. Stereotypic morphology of

glutamatergic synapses on identified muscle cells of *Drosophila larvae*. *J. Neurosci.* **9**, 710

(1989).

97.   Das, B., Brachmann, A. & Bekker, C. de. Both behavior-manipulating and non-manipulating

entomopathogenic fungi affect rhythmic gene expression in carpenter ant foragers upon infection.

bioRxiv 2023.01.19.524837 (2023) doi:10.1101/2023.01.19.524837.

98.   Davla, S. *et al.* An LC-MS/MS method for simultaneous analysis of up to six monoamines from

brain tissues. *J. Chromatogr. B* **1216**, 123604 (2023).

99.   Naundrup, A. *et al.* Pathogenic fungus uses volatiles to entice male flies into fatal matings with

infected female cadavers. *ISME J. 2022 1610* **16**, 2388–2397 (2022).

100.   Beckerson, W. C., Krider, C., Mohammad, U. A. & Bekker, C. de. 28 Minutes Later: Investigating

the role of aflatrem-like compounds in *Ophiocordyceps* parasite manipulation of zombie ants.

bioRxiv 2022.09.08.507134 (2023) doi:10.1101/2022.09.08.507134.

101.   Herbison, R. E. H. Lessons in Mind Control: Trends in Research on the Molecular Mechanisms

behind Parasite-Host Behavioral Manipulation. *Front. Ecol. Evol.* **5**, 102 (2017).

102.   Lefèvre, T. *et al.* Invasion of the body snatchers: the diversity and evolution of manipulative

strategies in host-parasite interactions. *Adv. Parasitol.* **68**, 45–83 (2009).

103.   Helluy, S. Parasite-induced alterations of sensorimotor pathways in gammarids: collateral damage

of neuroinflammation? *J. Exp. Biol.* **216**, 67–77 (2013).

104.   Helluy, S. & Holmes, J. C. Serotonin, octopamine, and the clinging behavior induced by the

parasite *Polymorphus paradoxus* (Acanthocephala) in *Gammarus lacustris* (Crustacea). *Can. J.*

Zool. **68**, 1214–1220 (1990).

105.   Herbison, R. *et al.* A molecular war: convergent and ontogenetic evidence for adaptive host

manipulation in related parasites infecting divergent hosts. *Proc. R. Soc. B Biol. Sci.* **286**,

20191827 (2019).

106.   Mohan, P. & Sinu, P. A. Is direct bodyguard manipulation a parasitoid-induced stress sleep? A

1008   new perspective. *Biol. Lett.* **18**, (2022).

1009   107.   Lafferty, K. D. & Shaw, J. C. Comparing mechanisms of host manipulation across host and

1010   parasite taxa. *J. Exp. Biol.* **216**, 56–66 (2013).

1011   108.   Shaw, J. C. *et al.* Parasite manipulation of brain monoamines in California killifish (*Fundulus*

1012   *parvipinnis*) by the trematode *Euhaplorchis californiensis*. *Proc. R. Soc. B Biol. Sci.* **276**, 1137–

1013   1146 (2009).

1014   109.   Prandovszky, E. *et al.* The Neurotropic Parasite *Toxoplasma gondii* Increases Dopamine

1015   Metabolism. *PLoS One* **6**, e23866 (2011).

1016   110.   Gasparetti, C., Nordlund, E., Jänis, J., Buchert, J. & Kruus, K. Extracellular tyrosinase from the

1017   fungus *Trichoderma reesei* shows product inhibition and different inhibition mechanism from the

1018   intracellular tyrosinase from *Agaricus bisporus*. *Biochim. Biophys. Acta - Proteins Proteomics*

1019   **1824**, 598–607 (2012).

1020   111.   Selinheimo, E. *et al.* Production and characterization of a secreted, C-terminally processed

1021   tyrosinase from the filamentous fungus *Trichoderma reesei*. *FEBS J.* **273**, 4322–4335 (2006).

1022   112.   Evans, H. C., Elliot, S. L. & Hughes, D. P. *Ophiocordyceps unilateralis:* A keystone species for

1023   unraveling ecosystem functioning and biodiversity of fungi in tropical forests? *Commun. Integr.*

1024   *Biol.* **4**, 598–602 (2011).

1025   113.   Leboeuf, A. C. *et al.* Oral transfer of chemical cues, growth proteins and hormones in social

1026   insects. *Elife* **5**, (2016).

1027   114.   Diez, L., Lejeune, P. & Detrain, C. Keep the nest clean: survival advantages of corpse removal in

1028   ants. *Biol. Lett.* **10**, (2014).

1029   115.   R Core Team. R: A language and environment for statistical computing. at https://www.r-

1030   project.org/ (2021).

1031   116.   RStudio Team. RStudio: Integrated Development for R. at http://www.rstudio.com/ (2015).

1032   117.   Therneau, T. A Package for Survival Analysis in S. at (2015).

1033   118.   Kassambara, A., Kosinski, M., Biecek, P. & Fabian, S. Drawing Survival Curves using 'ggplot2'

1034   [R package survminer version 0.4.4]. at https://cloud.r-

1035   project.org/web/packages/survminer/index.html (2019).

1036   119.   Matyash, V., Liebisch, G., Kurzchalia, T. V., Shevchenko, A. & Schwudke, D. Lipid extraction by

1037   methyl-tert-butyl ether for high-throughput lipidomics. *J. Lipid Res.* **49**, 1137–1146 (2008).

1038   120.   Tsugawa, H. *et al.* MS-DIAL: data-independent MS/MS deconvolution for comprehensive

1039   metabolome analysis. *Nat. Methods 2015 126* **12**, 523–526 (2015).

1040   121.   Kind, T. *et al.* LipidBlast in silico tandem mass spectrometry database for lipid identification. *Nat.*

1041   *Methods 2013 108* **10**, 755–758 (2013).

1042   122.   Kachman, M. *et al.* Deep annotation of untargeted LC-MS metabolomics data with Binner.

1043   *Bioinformatics* **36**, 1801–1806 (2020).

1044   123.   Fiehn Lab. ESI-MS-adducts-2020. https://fiehnlab.ucdavis.edu/images/files/software/ESI-MS-

1045   adducts-2020.xls (2020).

1046   124.   Acharjee, A., Larkman, J., Xu, Y., Cardoso, V. R. & Gkoutos, G. V. A random forest based

1047   biomarker discovery and power analysis framework for diagnostics research. *BMC Med.*

1048   *Genomics* **13**, 1–14 (2020).

1049   125.   DiLeo, M. V., Strahan, G. D., den Bakker, M. & Hoekenga, O. A. Weighted Correlation Network

1050   Analysis (WGCNA) Applied to the Tomato Fruit Metabolome. *PLoS One* **6**, e26683 (2011).

1051   126.   Zhang, G. *et al.* Integration of Metabolomics and Transcriptomics Revealed a Fatty Acid Network

1052   Exerting Growth Inhibitory Effects in Human Pancreatic Cancer. *Clin. Cancer Res.* **19**, 4983

1053   (2013).

1054   127.   Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for

1055   interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 15545–15550

1056   (2005).

1057   128.   Trapnell, C. *et al.* Differential gene and transcript expression analysis of RNA-seq experiments

1058   with TopHat and Cufflinks. *Nat. Protoc.* **7**, 562–578 (2012).

1059   129.   Tenebaum, D. & Maintainer, B. KEGGREST: Client-side REST access to the Kyoto Encyclopedia

1060   of Genes and Genomes (KEGG). at (2021).

1061   130.   Lechner, M. *et al.* Proteinortho: Detection of (Co-)orthologs in large-scale analysis. *BMC*

1062   *Bioinformatics* **12**, 1–9 (2011).

1063

1064

Acknowledgments

We would like to thank Sophia Vermeulen for assisting with testing fungal strain OJ2.1 and

Biplabendu Das for their help with ant infections and observations. We also thank Devin Burris for help

with ant monitoring. Jordan Dowell provided insightful discussions on LC-MS/MS methods and

interpretation for which we are grateful. The Genetics Society of America provided travel support for

dissemination of early results of this work. Research funding for this project came from NSF (NSF-

CAREER IOS-1941546 awarded to CdB).

1072

1073   **Author Contributions**

1074   CdB and IW conceived the study with methodological guidance from GMA. IW performed the

1075   experiment and analyses. All authors contributed to interpretation, writing, and revision and have

1076   approved this manuscript for submission.

1077

1078   **Additional Information**

1079   **Competing Interests**

1080   The authors declare no competing interests.

Main text figure and table legends

Figure 2. Ant manipulations occur at stereotypical days and times.

A) *Ophiocordyceps camponoti-floridani* infection leads to significantly increased mortality compared to sham treatment (p = 4.1E-7, log-rank test). B) Manipulated ants most often appeared 21-27 dpi, but as late as 33 dpi. C) Summiting behavior always occurred around dawn, most often 2.5 h before lights turned on, at ZT 21.5.

Figure 2. Selection of top features based on three independent analyses.

A) We used three independent statistical tests to select a single top features for metabolic pathway enrichment analyses. Additionally, we used a weighted co-abundance network approach (similar to WGCNA used for gene expression data) to form manipulation-correlated feature modules (resulting in 16 feature sets) for enrichment analyses. Results from those enrichment analyses were subsequently subjected to multiple filters to arrive at a final set of pathways and feature sets of interest. We also performed an enrichment analysis on a set of features only present in manipulated samples, but we found no overrepresented pathways and this feature set is not depicted in the figure. B) Volcano plots per dataset show DAMs that increased (red) or decreased (blue) at manipulation according to a 1.5-fold change and p ≤ 0.05 t-test threshold. C) PCAs for each dataset show distinct clustering of samples between manipulated (orange) and healthy ants (gray) along PC1. D) Features that passed at least two selection criteria (bold) were classified as top features and used for further pathway analyses. Top features were enriched for eight KEGG metabolic pathways.

Figure 3. WGCNA network modules of metabolite features, their correlation to manipulation, and enriched pathways. In total we detected three, five, and eight network modules that were significantly correlated to manipulation (Student correlation ≥ |0.5|, p ≤ 0.05) in the biogenic amines, lipids, and polyphenol datasets, respectively. "Unsorted" modules contain all features that were not successfully correlated to other features in a network.

Figure 4. Altered neurotransmitter precursor metabolism suggests suppressed ant immunity and increased dopamine during infection and manipulation. Upregulation of enzyme genes converting tyrosine to L-DOPA to dopamine coupled with reduced tyrosine and L-DOPA levels suggest increased concentrations of dopamine during infection, which may, in turn, contribute to manipulated behavior. Additionally, cardol triene may be produced by the fungus to suppress ant immunity by inhibiting melanin synthesis and thereby also increase availability of tyrosine and L-DOPA for dopamine synthesis. Combined with these specific metabolites, phenylalanine and tyrosine metabolism pathway enrichments further implicate these processes in manipulated behavior. Increases in expression, abundance, or activity are indicated by solid lines and red color, decreases by dotted lines and blue color. Metabolites are in rounded rectangles, genes are in rectangles. Annotated metabolites that were not DAMs have gray color, while compounds with white color were not identified in the LC-MS/MS data. Hypothesized links to ant (green) and fungal (orange) phenotypes are given in the shaded circles. Black arrows are shown when useful for context to indicate that an enzymatic step had a corresponding enzyme gene in either the fungal or ant genome,

1118   but it was not a DEG.

1119

Figure 5. Changing glycerophospholipid metabolism during manipulation primarily indicate altered

acetylcholine levels in the ant and cell membrane metabolism in the fungus.

Manipulated ant samples appeared to have two distinct groups, one with high acetylcholine and one with levels

similar to healthy ants. In manipulated ants, the putative acetylcholinesterase and two acetylcholine receptor genes

were downregulated, suggesting acetylcholine levels were accumulating. We hypothesize that increased

acetylcholine in manipulated ants is associated with the increased muscular activity during early manipulation (e.g.,

tremors, biting, and latching behavior). Once the ant is secured in a summited position and the mandible muscles

(and associated neurons) are further degraded, the fungus may be scavenging and converting the available

acetylcholine to choline, thereby rapidly reducing acetylcholine levels. The fungus may be converting choline to

phospholipids, such as phosphatidylcholines to remodel and build cell membranes. Although we detected multiple

phosphatidylcholine and phosphatidylethanolamine class features, we did not identify the singular metabolites 1,2-

diacyl-sn-glycero-3-phosphocholine and 1-acyl-2-acyl-sn-glycero-3-phosphoethanolamine, annotated as

"phosphatidylcholine" and "phosphatidylethanolamine" in the KEGG glycerophospholipid pathway, respectively.

The legend is the same as described in Figure 5, with the additional inclusion of genes that nearly qualified as DEGs

depicted in gray.

1135

Table 2. Enriched KEGG pathways selected across analyzed feature sets.

For KEGG pathways that were found in Mummichog or GSEA analyses ($p \leq 0.05$) to be of interest, we also

required them to show a minimum 1.25-fold enrichment, have at least three features contributing to that enrichment,

and the previous transcriptomics dataset [24] to contain at least three DEGs homologous to representative enzymes

within that pathway. Pathways are listed alphabetically with all feature sets (network modules or the top features set)

that were enriched for that pathway. Network modules names: biogenic amine (A1-A6), lipid (L1-L10), and

polyphenol (P1-P19). Only 16 of these 35 modules were correlated to manipulation and appear in this table.

1143

Supplementary Data 1. LC-MS/MS feature data used for analysis.

Supplementary Data 2. KEGG pathway enrichments and annotated features per feature group.

Supplementary Data 3. Internal standards used by the WCMC.

Supplementary Data 4. Original WCMC data files.

Supplementary Figure 1. Acetylcholine and acetylcholinesterase data.

Supplementary Table 1. Number of significant features per statistical analysis.

Supplementary Discussion 1. Metabolic pathway DEGs.

Supplementary Discussion 2. Drug metabolism pathway enrichments.

Supplementary Discussion 3. Beta-alanine pathway enrichment.

Supplementary Discussion 4. Vitamin B6 metabolism.

Supplementary Discussion 5. Serotonin and octopamine metabolism.

Supplementary Discussion 6. Glutamine and GABA metabolism.

Supplementary Methods. Expanded LC-MS/MS methods.