The catalytic activity of microRNA Argonautes plays a modest role in microRNA star strand cleavage in *C. elegans*

Kasuen Kotagama[1,3], Acadia L. Grimme[1,2,3], Bing Yang[1], Rima M. Sakhawala[1,2], Guoyun Yu[1], Katherine McJunkin[1,4]

[1]Laboratory of Cellular and Developmental Biology, NIDDK Intramural Research Program, 50 South Drive, Bethesda, MD 20892, USA

[2]Johns Hopkins University Department of Biology, 3400 N. Charles Street, Baltimore, MD 21218, USA.

[3]Authors contributed equally.

[4]Correspondence: mcjunkin@nih.gov

Abstract

Many Argonaute proteins can cleave RNA ("slicing") as part of the microRNA-induced silencing complex (miRISC), even though miRNA-mediated target repression is generally independent of target cleavage. Here we use *C. elegans* to examine the role of miRNA-guided slicing in organismal development. We find that unwinding and decay of the miRNA star strand is weakly defective in the absence of slicing, with the largest effect observed in embryos. Argonaute-Like Gene 2 (ALG-2) is more dependent on slicing for unwinding than ALG-1. The miRNAs that displayed the greatest (albeit very minor) dependence on slicing for unwinding tend to form stable duplexes with their star strand, and in some cases, lowering duplex stability alleviates dependence on slicing. While a few miRNA guide strands are reduced in slicing mutants, the basis of this is unclear since changes were not dependent on EBAX-1, a factor in the Target-Directed miRNA Degradation (TDMD) pathway. Gene expression and gross phenotype were wild type in slicing mutants, indicating that 1) slicing is not required for target regulation and 2) the weak unwinding defect does not significantly reduce the pool of functional miRISC. Thus, in contrast to previous work, miRISC slicing in *C. elegans* is dispensable in standard laboratory conditions.

Introduction

Multiple classes of small RNAs bind to Argonaute proteins to form an RNA-induced silencing complex (RISC) that represses expression of complementary RNAs. The PIWI domain of Argonaute resembles RNase H and in many Argonautes retains the ability to cleave RNA ("slicing") (1–5). While the role for Argonaute-mediated slicing is well-defined in some small RNA pathways, the role of slicing in the animal microRNA (miRNA) pathway is less clear.

Catalytic activity requires a tetrad of conserved residues (DEDH) (2, 3, 6, 7), with other parts of the protein also contributing supporting roles to catalysis (8–11). The catalytic or "slicing" activity

of Argonaute proteins can play a role at three phases of the RISC life cycle: small RNA biogenesis, the unwinding step of RISC assembly, and target RNA cleavage by mature RISC.

The role for slicing in small RNA biogenesis is most prominent in the PIWI-interacting RNA (piRNA) pathway. Slicing by Aubergine (Aub) and Argonaute-3 (Ago3) is required for piRNA biogenesis in flies (12, 13), and slicing activity of Mili is required for piRNA biogenesis in mice (14, 15). Slicing is dispensable for biogenesis of many other classes of small RNAs, including nearly all miRNAs (see below).

In RISC assembly, a small RNA duplex is first loaded into Argonaute. This is followed by unwinding, in which the "passenger" or "star" strand of the duplex dissociates from the complex, leaving the single-stranded "guide" bound to Argonaute as mature RISC. Slicing of the star strand can promote unwinding (16–19). The extent to which slicing of the star strand occurs and contributes to unwinding depends on the extent of base-pairing of the small RNA duplex (20–22). Thus, star strands of fully complementary siRNA duplexes are efficiently cleaved (16–19, 23). In contrast, star strands of bulged miRNA duplexes are generally inefficiently cleaved, and their dissociation is usually slicing-independent (24).

In mature RISC, complementary targets occupy the same position as the small RNA star strand and are likewise cleaved by Argonaute according to similar parameters. RISCs containing the Piwi proteins cleave complementary transposon-derived and genic transcripts (12, 25–29). In *C. elegans*, target cleavage by the Argonaute CSR-1 is crucial for regulation of germline and maternal mRNA targets (30–32). High complementarity generally promotes target cleavage by slicing-competent Argonautes, but the degree of complementarity required and the efficiency of slicing is Argonaute-dependent and incompletely understood (33–35).

In contrast to the clear role for Argonaute-mediated slicing in small RNA biogenesis, RISC assembly, and target cleavage in some animal small RNA pathways, the necessity for slicing in the miRNA pathway is less clear. The catalytic tetrad and slicing activity are conserved in many Argonautes in the miRNA pathway, despite the fact that biogenesis, unwinding, and target repression are slicing-independent for the vast majority of miRNAs. What biological roles does slicing play in the miRNA pathway that maintain selective pressure on the critical catalytic residues?

Biogenesis of animal miRNAs is generally carried out by sequential Microprocessor- and Dicer-mediated cleavage steps, and is independent of Argonaute-mediated slicing, with only one known exception. miR-451 is generated in a Dicer-independent manner that requires cleavage of its unusual precursor hairpin by Argonaute (36, 37).

Unwinding of animal miRNAs is also largely slicing-independent and is aided by a wedging function of Argonaute's N domain (24). One exception to this is the miR-486 guide:star duplex which is completely complementary (like an siRNA) and requires Argonaute to cleave the star strand for unwinding and activation of miR-486-RISC (38). Interestingly both slicing-dependent miRNAs are required for erythropoiesis, and rescuing expression of both miR-451 and miR-486 bypasses the requirement for Ago2-mediated slicing in erythropoiesis (38). Nonetheless, miR-451

and miR-486 are not conserved outside of vertebrates (39), so these described roles for slicing do not explain why miRNA Argonautes would retain their catalytic activity in other animals.

What about target cleavage? In animals, miRNAs bind their targets with incomplete complementarity, which is sufficient for translational repression and mRNA decay, and the vast majority of endogenous miRNA targets are repressed in a GW182-dependent manner that does not require target cleavage, with a few known exceptions (40–46). Because miRISC can efficiently promote target repression in the absence of slicing, slicing may be completely dispensable for target repression, even though it occurs in a limited number of miRNA:target interactions (42, 43, 45). One complication to answering this question in mammals is the multiple small RNA populations bound by the slicing-competent miRNA Argonaute Ago2. Outside of the vertebrate hematopoietic system, catalytic activity of Ago2 is also required for trophectoderm development, a function that is presumed to be due to Ago2's role in binding endogenous siRNAs (47). In other animals – such as in *C. elegans* and *Drosophila melanogaster* – endo-siRNAs and miRNAs are bound by distinct Argonaute proteins, so conservation of siRNA function would not exert selective pressure on the miRNA Argonautes. The dedicated roles of Argonaute proteins in these systems also makes these models excellent contexts for assessing the roles of slicing in the miRNA pathway.

A study by Bouasker and Simard previously examined the role of slicing in the miRNA pathway in *C. elegans* (48). By rescuing null mutations and RNAi of the miRNA Argonautes with overexpression of catalytically-inactivated Argonautes, Bouasker and Simard observed strong phenotypic effects, suggesting that catalytic activity in the miRNA pathway is important to animal physiology (48). Although small RNAs and mRNAs were not profiled globally, northern blots of select miRNAs showed incomplete precursor cleavage of some miRNAs in the context of overexpression of catalytic-dead Argonaute, suggesting a role for slicing in miRNA biogenesis (48).

We set out to revisit the role of the catalytic tetrad in *C. elegans* miRNA Argonautes using CRISPR/Cas9-mediated genome editing and deep sequencing of small RNAs and mRNAs. Here we show that slicing plays a role in star strand ejection and decay, and that this is more prominent in the Argonaute ALG-2 than ALG-1. However, the observed effects are only modest when using genome editing to inactivate slicing. No impact on target regulation was observed by RNA-seq. Consistent with this, animal physiology was like wild type. Thus, revisiting this question with state-of-the-art techniques shows that the dramatic effects observed in the previous study may have been due to overexpression (as necessitated by transgenic technologies at that time). This work demonstrates a modest role for miRNA-guided RNA cleavage in an invertebrate organism. Selective pressure to retain catalytic activity of miRNA Argonautes may be subtle or only apparent in harsher non-laboratory "real world" conditions.

Methods

C. elegans culture
Nematode culture was performed on NGM plates seeded with OP50 (49). Worms were maintained at 20°C except where otherwise noted. Embryo samples were harvested by hypochlorite treatment. For L4 and young adult samples, L1s were hatched overnight in M9 supplemented with cholesterol, then grown on NGM with OP50 for 48h at 20°C (for L4s) or for 44h at 25°C (for young adults). Brood size assays were conducted at 25°C after maintaining the strains at 25°C for one generation before the assay.

Genome editing
Guide RNAs were designed as close to the site of the edit as possible, while also favoring high GC content and favorable CRISPRscan scores (50, 51). Alt-R crRNAs (IDT) were pre-annealed with tracrRNA at 10μM by heating to 95°C for 5 minutes in IDT duplex buffer, then cooling to room temperature. Annealed guide RNAs were then assembled with Cas9 (IDT) at 2 μM Cas9 and 4 μM of total guide RNAs. Injection mixes also included donor molecules with 35nt homology arms for homology-directed repair (52); donor molecules were 100-150ng/μl (since all donors were relatively short). As a marker of CRISPR efficiency, a guide RNA and donor for *dpy-10* was included in all injection mixes (53). For *mir-63* and *mir-72* mutations, two rounds of CRISPR were performed. First, the *mir-63* and *mir-72* precursors were deleted and replaced by a cleavage site for an efficient crRNA (54) either in the wild type or the *alg-1(AEDH); alg-2(AEDH)* background (strains MCJ488 and MCJ511). In these backgrounds, CRISPR was performed again to reintroduce the mutant versions of the *mir-63* and *mir-72* precursors. Strains, alleles and oligos used in this study are listed in Supplemental Tables S1 and S2.

Quantitative miRNA-Taqman
For each sample, 9.96ng of total RNA was used to perform 5μL reverse transcription reactions using the TaqMan MicroRNA Reverse Transcription kit (ThermoFisher). After RT, 1.33μL of 1:4-diluted RT reaction was used in 5μL qPCR reactions with Taqman miRNA probes and Taqman Universal Mastermix II with UNG (ThermoFisher). Technical triplicates were run for each reaction on the Applied Biosystems QuantStudio Pro 6. Standard curves were generated by similarly performing miRNA-Taqman on a dilution series of an RNA oligo of the sequence of interest (Horizon Discoveries).

Deep sequencing library construction and data analysis
Samples were snap frozen, then resuspended in Trizol and vortexed for 15 minutes at room temperature. Thereafter, total RNA was purified according to the Trizol manufacturer's instructions (ThermoFisher).

For preparing small RNA libraries, RNA oligos listed in Table S2 were spiked into total RNA at a concentration of 1pg/μl in 100ng/μl total RNA. Small RNA libraries were prepared using the NEBNext Small RNA Library Prep kit (NEB), with some modifications. Size selection was only performed after reverse transcription, using 8% urea gels to purify 65-75nt products. RT products were treated with RNAse H for 30 min at 37°C prior to gel purification. After excision, products were eluted overnight in 700μl 0.3M NaCl in TE. Products were then purified by isopropanol

precipitation, washed in 80% ethanol, and resuspended in 40μl 50mM Tris-HCl 75mM KCl 3mM MgCl2 pH 8.3 before proceeding with the NEBNext protocol. Samples were sequenced on a HiSeq 3000 or NextSeq 550 (Illumina). Raw data can be accessed under SRA Bioproject accession number PRJNA922944.

Small RNA libraries were trimmed using Cutadapt 3.4 (55), then mapped using bowtie2 2.4.4 with settings "--no-unal --end-to-end –sensitive" (56). BAM files were sorted and indexed using samtools 1.13, then assigned to miRNAs using htseq 0.13.5 (57, 58). For genome mapping and miRNA assignment, custom reference files were used which included the spiked in RNA oligos. Statistical analysis was performed using DEseq2 and GraphPad Prism (59). For spike-in normalization (Figure S1), size factors for DEseq2 analysis were calculated by dividing spike-in reads in each sample by the median number of spike-in reads per sample. For all other analyses, default DEseq2 normalization was performed using all endogenous miRNA-mapping reads (spike-in reads were excluded from analysis).

For RNAseq, libraries were prepared by Psomagen using the Truseq Stranded Total RNA Library Prep Kit and sequenced on a NovaSeq6000 S4 (Illumina). Low quality bases and adaptors were trimmed from sequence reads using Cutadapt v2.7 with parameters "–q 20 -minimum-length 25" (55). Resulting reads were mapped to WS277 using HISAT2 v2.1.0 with default parameters (60). Aligned reads were counted using subread featureCounts v1.6.4 with default parameters, except that "-s" option was used to specify appropriate strands (61). Differential expression analysis was done with DESeq2 v1.22.1 (ref4) in v3.5.1 R version (59).

To calculate predicted miRNA:star duplex structures and calculate minimum free energy (MFE), RNAcofold was used with default parameters, as part of the ViennaRNA package (version 2.4.18).

Reagents
The following reagents were used: Alt-R crRNAs (IDT), Cas9 (IDT), Trizol (ThermoFisher), TaqMan MicroRNA Reverse Transcription kit (ThermoFisher), Taqman miRNA probes (ThermoFisher), Taqman Universal Mastermix II with UNG (ThermoFisher), NEBNext Small RNA Library Prep kit (NEB), RNA oligonucleotides (Horizon Discoveries). The following specialized instruments were used: QuantStudio Pro 6 (Applied Biosystems), HiSeq 3000 (Illumina), NextSeq 550 (Illumina), NovaSeq6000 S4 (Illumina).

Biological Resources
All strains used in the study are listed in Table S1.

Statistical Analyses
Statistical analysis for deep sequencing datasets was performed using DEseq2 using default settings except when spike-ins were used as noted (59). For analysis of standard data types, the Analysis function in GraphPad Prism was used to select and perform the appropriate statistical test. At least three (in most cases four) replicates of each genotype were used for all experiments.

Data Availability/Sequence Data Resources:
All deep sequencing raw data can be accessed under SRA Bioproject PRJNA922944.

Data Availability/Novel Programs, Software, Algorithms
Not applicable.

Web Sites/Data Base Referencing:
MirGeneDB 2.1 (https://mirgenedb.org/), miRBase (https://www.mirbase.org/), WormBase (https://wormbase.org/)

Results

Mutation of catalytic tetrad in both miRNA Argonautes does not impact physiology.

To revisit the role of miRISC-mediated slicing in invertebrates, we used CRISPR/Cas9-mediated genome editing to mutate the DEDH catalytic tetrad to AEDH in the major *C. elegans* miRNA Argonautes, ALG-1 and ALG-2 (Figure 1A). This mutation has previously been shown to completely inactivate catalytic activity in orthologous Argonautes *in vitro* and *in vivo* (3, 36). (The germline-specific miRNA Argonaute ALG-5 lacks conservation in any of the positions corresponding to DEDH and therefore likely lacks catalytic activity (Figure 1A).)

Mutant strains were superficially wild type. Negligible lethality was observed. Brood size assays showed that single *alg-1(AEDH)* or *alg-2(AEDH)* or double *alg-1(AEDH); alg-2(AEDH)* mutants all had wild-type like fecundity (Figure 1B). In comparison, *alg-1(null)* animals had compromised fecundity, as previously observed (62, 63). Therefore, *alg-1(AEDH)* and *alg-2(AEDH)* largely retain wild type function and do not mimic loss-of-function alleles.

These results are in contrast to a previous study in which catalytic dead versions of Argonaute were over-expressed in an Argonaute loss-of-function background (48). In that case, catalytic-dead versions of Argonaute only partially rescued viability and provided no rescue of fertility (all animals were dead or sterile). Taken together, this suggests that the previous study was impacted by expression level of the catalytic dead Argonaute and that, in fact, slicing is not required for grossly normal physiology and fitness.

Some miRNA star strands are slightly elevated in the absence of slicing.

Previous work in other models showed that the catalytic activity of Argonaute in the miRNA pathway is crucial for the biogenesis of a Dicer-independent miRNA and the proper loading of an siRNA-like miRNA (36–38). To determine the impact of catalytically inactive Argonautes on *C. elegans* miRNAs, we profiled miRNA star and guide strands from embryos, L4 stage larvae and young adults.

Exogenous RNA oligos were spiked in prior to library prep to control for potential large-scale changes in miRNA populations. Spike-in normalization (Figure S1) and normalization to all endogenous miRNA reads (Figure 2) yielded similar results, demonstrating that wholesale changes in small RNA population did not occur. Normalization to all endogenous miRNAs resulted in slightly lower variance across replicates and thus yielded slightly more significantly altered small

RNAs (Figure S1 and Figure 2); thus, we proceeded with this type of normalization for all other analyses.

Embryo samples showed the greatest number of significantly altered miRNAs. Fifteen miRNA strands were upregulated, all of which are miRNA star strands (Figure 2A, Tables S3-S5). Subtle star strand upregulation was also widespread such that star strands were upregulated overall compared to miRNA guide strands in embryos (Figure 2B-C). Four and two star strands were also upregulated in L4 larvae and adults, respectively (Figure 2A, Tables S6-S7). The upregulation of star strands suggests that slicing promotes unwinding, as previously observed for siRNAs (16, 18, 23, 64). The amplitude of these changes is modest compared to the guide:star strand ratio, which is ≥10-fold for most miRNAs (65). Therefore, the putative slicing-dependent unwinding of these star strands occurs alongside a more dominant slicing-independent pathway. This low amplitude of star strand elevation suggests that only a very small proportion of miRNA guide strand function would be impaired and may not result in observable guide strand loss-of-function.

Very few miRNAs were downregulated (Figure 2, Table S5-S7). These were mostly miRNA guide strands, including *mir-243-3p, mir-1821-3p,* and *mir-1829a-5p*, which were each downregulated in more than one developmental stage. Again, the amplitude of changes is modest, suggesting that slicing is not strictly required for the biogenesis of these miRNAs (Figure 2). The mechanism of the downregulation of these guide strands is not understood, though one possible model is explored below.

Slicing plays a greater role in ALG-2 unwinding than that of ALG-1.

We sought to understand why embryo samples displayed more widespread elevation of star strands upon inactivation of slicing than other stages. We hypothesized that the larger effect in embryos might be due to a difference in relative expression of the Argonaute paralogs ALG-1 and ALG-2. ALG-2 is expressed earlier in embryogenesis than ALG-1, and its expression decreases after embryogenesis (63, 66). Furthermore, the most abundant embryonic miRNA family, *mir-35-42*, largely depends on ALG-2 expression for its stabilization (63, 67, 68). Together these data suggest that the ratio of ALG-2:ALG-1 is higher in embryos than later stages. Therefore, the larger effect of slicing deficiency in embryos may be due to a greater dependence of ALG-2 unwinding on slicing than that of ALG-1.

To test this hypothesis, we profiled small RNAs from embryos that were single mutants, with slicing inactivated in either ALG-1 or ALG-2. Single *alg-1(AEDH)* mutant embryos displayed very few significant changes in miRNA abundance (Figure 3A, Table S8). In contrast, *alg-2(AEDH)* single mutants showed nearly as many significant changes as *alg-1(AEDH); alg-2(AEDH)* double mutant embryos (Figure 3A, Table S9). Below the threshold of significance, star strands were globally slightly elevated in the *alg-2(AEDH)* single mutants, again mimicking the *alg-1(AEDH); alg-2(AEDH)* double mutant (Figure 3B). Therefore, the elevation of star strands in *alg-1(AEDH); alg-2(AEDH)* is largely due to the sensitivity of ALG-2 to catalytic inactivation. Samples harvested from later stages may show subtler changes in miRNA abundance due to a

higher ratio of ALG-1:ALG-2 expression coupled with a lower sensitivity of ALG-1 to inactivation of slicing.

High duplex stability favors a greater role for slicing in unwinding.

While many miRNA star strands were elevated in embryos, very few were elevated in adults and L4, and these were largely consistent across both stages. We further examined the properties of these guide:star duplexes to determine if any distinctive feature might predict the extent to which a miRNA duplex depends on slicing for unwinding.

The star strands *mir-63-5p* and *mir-72-3p* were upregulated across all three stages examined. We observed that *mir-63* and *mir-72* both had relatively low minimum free energy (MFE) of the guide:star duplex compared to other miRNAs (they are both in the lowest 15% of MFEs across all miRNA duplexes) (Figure 4A-B). We also observed that the guide strands of these duplexes displayed a high rate of 3' end trimming (within the top 12% of all guide strands) (69) (Figure 4A).

One possible model is that the high stability of the guide:star duplex slows slicing-independent unwinding (22, 70, 71). The presence of the star strand could favor a structural conformation in which the 3' end of the guide strand is not bound by the PAZ domain, leaving it vulnerable to exonucleolytic trimming (72, 73). Thus, a longer dwell time of the star strand could lead to higher prevalence of trimmed guide strand molecules. The slow rate of slicing-independent unwinding and thereby long dwell time of the star strand could allow the inefficient slicing-dependent star strand ejection mechanism to play a relatively larger (though still minor) role in miRISC maturation.

To test this model, we performed CRISPR to make mutations that are predicted to preserve overall base pairing structure while decreasing duplex stability (i.e. increasing MFE) (Figure 4B). These mutations consisted of one to three mutations on each strand of the miRNA duplex, replacing G:C base pairs with A:U base pairs (Figure 4C). Mutations were introduced near the middle of the duplex (avoiding the terminal four base pairs) in an attempt to preserve thermodynamic asymmetry and strand selection (74–76).

We then examined the mutant duplex miRNAs in an otherwise wild type or the *alg-1(AEDH); alg-2(AEDH)* background to determine whether decreasing duplex stability also reduces the dependence of star strand unwinding on slicing. To measure both star strand stabilization and strand selection, we performed qPCR in L4 samples calibrated with standard curves of synthetic RNA oligos to enable absolute quantification (Figure S2). For both wild type *mir-72* and *mir-63*, qPCR confirmed upregulation of the star strand in the *alg-1(AEDH); alg-2(AEDH)* mutant background with no concomitant effect on the guide strand (Figure 4E-F). For *mir-72*, either one or two mutations that reduced thermodynamic stability of the guide:star duplex also abrogated the effect of *alg-1(AEDH); alg-2(AEDH)* on the star strand abundance (Figure 4E). This suggests that the dependence of the wild type duplex on slicing for complete unwinding is due to high thermodynamic stability of the miRNA:star duplex. For both *mir-72* mutant duplexes, the guide

strand selection remained high, arguing against an alternative model in which star strand effects are altered by increased star strand loading into Argonaute (Figure 4G).

For *mir-63*, two mutations in the duplex (*mir-63_mut2*) did not abolish the increased star strand abundance observed in the *alg-1(AEDH); alg-2(AEDH)* background (Figure 4F). A third mutation in the duplex of *mir-63* (*mir-63_mut3*) caused a switch in strand selection that favored loading of *mir-63-5p* over the erstwhile guide strand *mir-63-3p*, making it difficult to interpret the effects of this mutation in the *alg-1(AEDH); alg-2(AEDH)* background (Figure 4H).

Overall, the *mir-72* experiments show that duplex thermodynamic stability is one factor that influences dependence on slicing for unwinding (Figure 4E). At the same time, the weak correlation between thermodynamic stability and dependence on slicing (color and MFE - Figure 4A) and the lack of effect of two mutations in *mir-63* (Figure 4F) together show that other unknown factors also influence dependence on slicing for unwinding.

Minor changes in star strand abundance do not result in changes in target repression.

As discussed above, physiology of the *alg-1(AEDH); alg-2(AEDH)* double mutant animals is grossly normal (Figure 1). Next, we performed RNA-seq to determine whether changes in miRNA abundance (Figure 2) or absence of target cleavage led to a more subtle molecular phenotype. Negligible changes in gene expression were observed. No genes were significantly altered in expression in L4 stage *alg-1(AEDH); alg-2(AEDH)* animals compared to wild type, whereas only one or two were altered in embryo and adult, respectively (Figure 5, Tables S10-S12). These changes were both down-regulation of protein-coding genes (not an up-regulation that might be expected from de-repression of miRNA target genes).

We further examined the gene *gpap-1*, which was down-regulated in both embryo and adult *alg-1(AEDH); alg-2(AEDH)* animals (Figure 5). By close examination of the RNA-seq data, we observed a single-nucleotide polymorphism in the *gpap-1* 5' UTR in *alg-1(AEDH); alg-2(AEDH)* samples (Figure S3). Thus, this polymorphism is very likely the basis of these gene expression changes, rather than the mutations in *alg-1* and *alg-2*. No homology was observed between the guide RNAs used to generate the *alg-1* and *alg-2* alleles and the region containing the SNP in *gpap-1*. Thus, we posit that this SNP is due to genetic drift and fixation that occurred during the passaging and homozygosing of the *alg-1(AEDH); alg-2(AEDH)* strain. Therefore, nearly no gene expression changes arise as a result of *alg-1(AEDH); alg-2(AEDH)*. Overall, this suggests that target mRNA cleavage plays little or no role in regulation of gene expression by miRNAs in *C. elegans*. Furthermore, the role that slicing plays in miRISC maturation is so minor that it does not drive changes in gene expression.

Down-regulation of select guide strands in the absence of slicing is *ebax-1*-independent.

In addition to elevation of star strands in the *alg-1(AEDH); alg-2(AEDH)* mutants, decreased levels of a few miRNA guide strands were also observed. The three guide strands that were

downregulated in two or more developmental stages were *mir-243-3p*, *mir-1829a-5p*, and *mir-1821-3p*. We sought to formulate a model for why these guides might be downregulated in *alg-1(AEDH); alg-2(AEDH)* based on the features of these miRNAs. *mir-243-3p* was previously shown to be largely loaded into RDE-1 and to act as an siRNA to target a fully complementary site in the 3' UTR of Y47H10A.5 (68). Highly complementary targets can drive target-directed miRNA decay (TDMD); in this process, target cleavage usually does not occur due to a region of central mismatches in the miRNA:target duplex (72, 77–89). A minor portion of *mir-243* is bound by ALG-1, and we wondered whether the mild decrease in *mir-243* (as well as *mir-1829a-5p* and *mir-1821-3p*) in *alg-1(AEDH); alg-2(AEDH)* could be due to enhanced TDMD in the absence of target cleavage. In this model, TDMD would be less active when ALG-1 is wild type and can cleave the target; when cleavage is inactivated in the *alg-1(AEDH); alg-2(AEDH)* context, TDMD would be more potent, leading to decreased *mir-243* (and other miRNA guide strands) in this mutant.

We tested this model by inactivating this putative TDMD in two ways: 1) by introducing a null mutation in the TDMD effector EBAX-1 (known as ZSWIM8 in mammals) and 2) by deleting the binding site for *mir-243* in the Y47H10A.5 3' UTR (68, 90, 91). If the model is correct, then either of these contexts should abrogate the effect of the *alg-1(AEDH); alg-2(AEDH)* mutations on *mir-243* abundance. Comparing *ebax-1(null)* young adults to wild type, we corroborate the stabilization of *mir-788, mir-247, mir-42, mir-230, mir-797, mir-43 and mir-253* in this mutant as previously observed (Figure S4, Table S13) (91). (Note that *mir-35-41* are not highly expressed in these samples, which do not contain embryos, in contrast to the previous studies of *ebax-1(null)* which examined gravid adults or embryos (91, 92).)

Next, *ebax-1(null)* was introduced to the *alg-1(AEDH); alg-2(AEDH)* background, and miRNA abundances were compared to *ebax-1(null)* single mutants. Very similar amplitude in changes were observed when comparing *ebax-1(null); alg-1(AEDH); alg-2(AEDH)* to *ebax-1(null)* (Figure 6A, Table S14) as when comparing *alg-1(AEDH); alg-2(AEDH)* to wild type (Figure 6B). The correlation of log$_2$(fold changes) across all abundant miRNAs in these two comparisons was fairly strong (Figure 6C, Pearson's r = 0.6376) since it was similar to the correlation between two independent experiments of the same genotypes (each experiment including four biological replicates, Figure S4B, Table S15, Pearson's r = 0.7173). This demonstrates that overall the observed effects of *alg-1(AEDH); alg-2(AEDH)* are not EBAX-1-dependent.

When examining the guide strands that were decreased in *alg-1(AEDH); alg-2(AEDH)*, similar amplitude of downregulation was observed for *mir-243-3p*, *mir-1829a-5p*, and *mir-1821-3p* in the *ebax-1(null)* context (Figure 6A,D). However, this change for *mir-243-3p* and *mir-1821-3p* did not reach statistical significance in the second set of experiments in the wild type background (Figure 6B), and *mir-243-3p* likewise did not reach significance in the *ebax-1(null)* background (Figure 6A). This largely reflects the moderate amplitude of the downregulation (as well as noise and stringency introduced by correction due to multiple testing). Nonetheless, despite the failure to reach arbitrary statistical cutoffs, the amplitude of downregulation is similar in the *ebax-1(null)* context (Figure 6A,D), and our interpretation is that this downregulation is EBAX-1-independent.

Thus, the model that this change is due to enhanced TDMD in the catalytic dead mutant is incorrect.

The *mir-243* binding site mutant (hereafter *Y47(mir-243bs_del)*) further refuted the model. The decrease of *mir-243* was still observed when comparing *Y47(mir-243bs_del); alg-1(AEDH); alg-2(AEDH)* to *Y47(mir-243bs_del)* single mutants (Figure 6D-F, Table S16). This is further evidence negating a role for TDMD in the decrease of select miRNA guide strands in the slicing mutant (also see Discussion).

Discussion

Here, we demonstrate an unexpectedly modest role for slicing in the miRNA pathway in *C. elegans*. Unlike in vertebrates, no miRNA species are strictly dependent upon slicing for their biogenesis or unwinding. Furthermore, target gene expression is unchanged as measured by RNAseq. This results in normal development and reproduction in standard laboratory conditions. This work thus overturns the current literature with regard to the importance of miRNA-guided slicing in *C. elegans* (48). *C. elegans* is the first animal to our knowledge in which slicing by miRNA Argonautes is conserved but dispensable on an organismal level.

Concurrently to this work, the Simard lab also generated and performed extensive phenotypic analysis of ALG-1 and ALG-2 slicing mutants (Pal, et al. *Co-submitted*). By examining compound mutants and heat stress, Pal et al. observe fitness phenotypes associated with catalytic inactivation of ALG-1 and/or ALG-2. Thus, although phenotypes of slicing mutants are much milder than previously thought, they are still apparent in sensitized genetic and physical conditions. These stress-associated phenotypes may provide an experimental framework for understanding the selection pressure for maintaining the catalytic activity of ALG-1 and ALG-2 in the long-term.

Molecularly, the basis of these phenotypes may be related to impaired miRNA:star duplex unwinding. We observed a measurable effect on duplex unwinding, represented by a modest overall stabilization of star strands in embryos (Figure 7). Why is this more prominent in *C. elegans* than other previously studied organisms? By examining single mutant strains, catalytic-dead ALG-2 appears to have a greater unwinding defect than catalytic-dead ALG-1. ALG-2 may be more dependent upon slicing for unwinding than previously-examined miRNA Argonautes from other organisms. This may be due to either lower efficiency of slicing-independent unwinding or to higher efficiency of slicing of bulged miRNA:star duplexes. Differences in the N terminal domain or non-catalytic residues in the PIWI domain could contribute to either reduced slicing-independent unwinding or enhanced slicing, whereas differences in the PAZ domain could also impair slicing-independent unwinding (8–11, 24, 93–95). This opens up new areas of investigation into the biochemical properties of ALG-2 compared to ALG-1 or Argonautes from other species.

Some miRNA star strands were elevated in catalytic-dead mutants in all examined contexts, not only in embryos. The duplexes from which these star strands are derived (*mir-63* and *mir-72*) have fairly high thermodynamic stability. For *mir-72*, we find that this duplex stability is indeed a determinant of its dependence on slicing for unwinding. For *mir-63*, lowering duplex stability did

not rescue unwinding in the absence of slicing. Thus, duplex stability is *one* of the factors that determines the extent of reliance on slicing for unwinding, as previously observed in other settings (20, 22, 71). However, as illustrated by the case of *mir-63* and the efficient slicing-independent unwinding of even more stable duplexes (Figure 4A), other uncharacterized factors also control the extent of slicing-dependent unwinding. Still, even for *mir-63* and *mir-72,* slicing-dependent unwinding is the minor pathway; only modest elevation in star strands is observed, and the vast majority of miRISC matures to the guide-only bound state and properly represses all targets (Figure 7).

A more mysterious observation in this study was the decreased levels of a few guide strands. While we explored a model of enhanced TDMD in the absence of slicing, this was not borne out. However, conventional TDMD has yet to be demonstrated in *C. elegans* (69, 92, 96), so the pathways of slicing and TDMD may not be competing in this context as might be expected in other systems (72, 77–91). Because of the modest reduction of these few guide strands, we posit that slicing is not central to their biogenesis. However, slicing could represent one of multiple biogenesis pathways for these guides or somehow regulate their canonical biogenesis. Interestingly, at least two of the three downregulated guides are primarily loaded into an alternative Argonaute: *mir-243-3p* into RDE-1 and *mir-1829a-5p* into ALG-5 (63, 68). Could ALG-1/2 cleavage products load into other Argonautes? Precedents for this type of interaction include the loading of Aub and Ago3 cleavage products into each other in the piRNA ping-pong pathway (97). The secondary structures of *mir-243* and *mir-1829a* do not provide a clear parallel, but future experiments may nonetheless reveal some similarity.

Overall, this study of miRISC-mediated slicing in a model organism outside the vertebrate lineage showed a surprisingly mild effect on physiology. Although laboratory conditions facilitate the reproducibility to observe clear molecular signatures, they do not recapitulate challenges encountered in the wild. Harsher growth conditions used in a parallel study revealed modest fitness phenotypes associated with loss of miRISC slicing (Pal, et al. *Co-submitted*). Robustness of miRISC maturation at a broad range of temperatures may contribute to selection pressure on catalytic residues in the wild. Moreover, our study examined samples derived from whole animals. In the long term, tissue-specific studies may reveal more salient roles for miRISC-mediated slicing not apparent in our study.

Data Availability All deep sequencing raw data can be accessed under SRA Bioproject PRJNA922944. Reviewer link:
https://dataview.ncbi.nlm.nih.gov/object/PRJNA922944?reviewer=u39pd248iat8khairclnt9ge21

Funding
This work was supported by the NIDDK Intramural Research Program (ZIA DK075147).

Acknowledgments
We thank WormBase, the NIDDK Genomics Core, the NCI Genomics Core, and NIH High Performance Computing. Strains are regularly received from and deposited to the CGC, which is

funded by NIH Office of Research Infrastructure Programs (P40 OD010440). Thank you to Anna Zinovyeva, Martin Simard, and members of the McJunkin lab for helpful discussions.

References

1. Liu,J., Carmell,M.A., Rivas,F. V., Marsden,C.G., Thomson,J.M., Song,J.J., Hammond,S.M., Joshua-Tor,L. and Hannon,G.J. (2004) Argonaute2 is the catalytic engine of mammalian RNAi. *Science (80-.).*, **305**, 1437–1441.

2. Song,J.J., Smith,S.K., Hannon,G.J. and Joshua-Tor,L. (2004) Crystal structure of argonaute and its implications for RISC slicer activity. *Science (80-.).*, **305**, 1434–1437.

3. Rivas,F. V., Tolia,N.H., Song,J.J., Aragon,J.P., Liu,J., Hannon,G.J. and Joshua-Tor,L. (2005) Purified Argonaute2 and an siRNA form recombinant human RISC. *Nat. Struct. Mol. Biol.*, **12**, 340–349.

4. Rand,T.A., Ginalski,K., Grishin,N. V. and Wang,X. (2004) Biochemical identification of Argonaute 2 as the sole protein required for RNA-induced silencing complex activity. *Proc. Natl. Acad. Sci. U. S. A.*, **101**, 14385–14389.

5. Meister,G., Landthaler,M., Patkaniowska,A., Dorsett,Y., Teng,G. and Tuschl,T. (2004) Human Argonaute2 mediates RNA cleavage targeted by miRNAs and siRNAs. *Mol. Cell*, **15**, 185–197.

6. Nakanishi,K., Weinberg,D.E., Bartel,D.P. and Patel,D.J. (2012) Structure of yeast Argonaute with guide RNA. *Nat. 2012 4867403*, **486**, 368–374.

7. Yuan,Y.R., Pei,Y., Ma,J.B., Kuryavyi,V., Zhadina,M., Meister,G., Chen,H.Y., Dauter,Z., Tuschl,T. and Patel,D.J. (2005) Crystal structure of A. aeolicus argonaute, a site-specific DNA-guided endoribonuclease, provides insights into RISC-mediated mRNA cleavage. *Mol. Cell*, **19**, 405–419.

8. Faehnle,C.R., Elkayam,E., Haase,A.D., Hannon,G.J. and Joshua-Tor,L. (2013) The Making of a Slicer: Activation of Human Argonaute-1. *Cell Rep.*, **3**, 1901–1909.

9. Hauptmann,J., Dueck,A., Harlander,S., Pfaff,J., Merkl,R. and Meister,G. (2013) Turning catalytically inactive human Argonaute proteins into active slicer enzymes. *Nat. Struct. Mol. Biol.*, **20**, 814–817.

10. Hauptmann,J., Kater,L., Löffler,P., Merkl,R. and Meister,G. (2014) Generation of catalytic human Ago4 identifies structural elements important for RNA cleavage. *RNA*, **20**, 1532–1538.

11. Schürmann,N., Trabuco,L.G., Bender,C., Russell,R.B. and Grimm,D. (2013) Molecular dissection of human Argonaute proteins by DNA shuffling. *Nat. Struct. Mol. Biol.*, **20**, 818–826.

12. Brennecke,J., Aravin,A.A., Stark,A., Dus,M., Kellis,M., Sachidanandam,R. and Hannon,G.J.

(2007) Discrete small RNA-generating loci as master regulators of transposon activity in Drosophila. *Cell*, **128**, 1089–1103.

13. Gunawardane,L.S., Saito,K., Nishida,K.M., Miyoshi,K., Kawamura,Y., Nagami,T., Siomi,H. and Siomi,M.C. (2007) A slicer-mediated mechanism for repeat-associated siRNA 5' end formation in Drosophila. *Science*, **315**, 1587–1590.

14. Aravin,A.A., Sachidanandam,R., Girard,A., Fejes-Toth,K. and Hannon,G.J. (2007) Developmentally regulated piRNA clusters implicate MILI in transposon control. *Science*, **316**, 744–747.

15. De Fazio,S., Bartonicek,N., Di Giacomo,M., Abreu-Goodger,C., Sankar,A., Funaya,C., Antony,C., Moreira,P.N., Enright,A.J. and O'carroll,D. (2011) The endonuclease activity of Mili fuels piRNA amplification that silences LINE1 elements. *Nature*, **480**, 259–263.

16. Matranga,C., Tomari,Y., Shin,C., Bartel,D.P. and Zamore,P.D. (2005) Passenger-strand cleavage facilitates assembly of siRNA into Ago2-containing RNAi enzyme complexes. *Cell*, **123**, 607–620.

17. Miyoshi,K., Tsukumo,H., Nagami,T., Siomi,H. and Siomi,M.C. (2005) Slicer function of Drosophila Argonautes and its involvement in RISC formation. *Genes Dev.*, **19**, 2837–2848.

18. Leuschner,P.J.F., Ameres,S.L., Kueng,S. and Martinez,J. (2006) Cleavage of the siRNA passenger strand during RISC assembly in human cells. *EMBO Rep.*, **7**, 314–320.

19. Rand,T.A., Petersen,S., Du,F. and Wang,X. (2005) Argonaute2 cleaves the anti-guide strand of siRNA during RISC activation. *Cell*, **123**, 621–629.

20. Kawamata,T., Seitz,H. and Tomari,Y. (2009) Structural determinants of miRNAs for RISC loading and slicer-independent unwinding. *Nat. Struct. Mol. Biol.*, **16**, 953–960.

21. Shin,C. (2008) Cleavage of the Star Strand Facilitates Assembly of Some MicroRNAs into Ago2-containing Silencing Complexes in Mammals. *Mol. Cells*, **26**, 308–313.

22. Park,J.H. and Shin,C. (2015) Slicer-independent mechanism drives small-RNA strand separation during human RISC assembly. *Nucleic Acids Res.*, **43**, 9418–9433.

23. Steiner,F.A., Okihara,K.L., Hoogstrate,S.W., Sijen,T. and Ketting,R.F. (2009) RDE-1 slicer activity is required only for passenger-strand cleavage during RNAi in Caenorhabditis elegans. *Nat. Struct. Mol. Biol.*, **16**, 207–211.

24. Kwak,P.B. and Tomari,Y. (2012) The N domain of Argonaute drives duplex unwinding during RISC assembly. *Nat. Struct. Mol. Biol.*, **19**, 145–151.

25. Wu,P.H., Fu,Y., Cecchini,K., Özata,D.M., Arif,A., Yu,T., Colpan,C., Gainetdinov,I., Weng,Z. and Zamore,P.D. (2020) The evolutionarily conserved piRNA-producing locus pi6 is required for male mouse fertility. *Nat. Genet.*, **52**, 728–739.

26. Goh,W.S.S., Falciatori,I., Tam,O.H., Burgess,R., Meikar,O., Kotaja,N., Hammell,M. and Hannon,G.J. (2015) piRNA-directed cleavage of meiotic transcripts regulates spermatogenesis. *Genes Dev*, **29**, 1032–1044.

27. Vourekas,A., Zheng,Q., Alexiou,P., Maragkakis,M., Kirino,Y., Gregory,B.D. and

Mourelatos,Z. (2012) Mili and Miwi target RNA repertoire reveals piRNA biogenesis and function of Miwi in spermiogenesis. *Nat. Struct. Mol. Biol.*, **19**, 773–781.

28. Zhang,P., Kang,J.Y., Gou,L.T., Wang,J., Xue,Y., Skogerboe,G., Dai,P., Huang,D.W., Chen,R., Fu,X.D., *et al.* (2015) MIWI and piRNA-mediated cleavage of messenger RNAs in mouse testes. *Cell Res.*, **25**, 193–207.

29. Reuter,M., Berninger,P., Chuma,S., Shah,H., Hosokawa,M., Funaya,C., Antony,C., Sachidanandam,R. and Pillai,R.S. (2011) Miwi catalysis is required for piRNA amplification-independent LINE1 transposon silencing. *Nature*, **480**, 264–267.

30. Aoki,K., Moriguchi,H., Yoshioka,T., Okawa,K. and Tabara,H. (2007) In vitro analyses of the production and activity of secondary small interfering RNAs in C. elegans. *EMBO J.*, **26**, 5007–5019.

31. Gerson-Gurwitz,A., Wang,S., Sathe,S., Green,R., Yeo,G.W., Oegema,K. and Desai,A. (2016) A Small RNA-Catalytic Argonaute Pathway Tunes Germline Transcript Levels to Ensure Embryonic Divisions. *Cell*, **165**, 396–409.

32. Quarato,P., Singh,M., Cornes,E., Li,B., Bourdon,L., Mueller,F., Didier,C. and Cecere,G. (2021) Germline inherited small RNAs facilitate the clearance of untranslated maternal mRNAs in C. elegans embryos. *Nat. Commun.*, **12**.

33. Park,M.S., Sim,G.Y., Kehling,A.C. and Nakanishi,K. (2020) Human Argonaute2 and Argonaute3 are catalytically activated by different lengths of guide RNA. *Proc. Natl. Acad. Sci. U. S. A.*, **117**, 28576–28578.

34. Nakanishi,K. (2022) Anatomy of four human Argonaute proteins. *Nucleic Acids Res.*, 10.1093/nar/gkac519.

35. Arif,A., Bailey,S., Izumi,N., Anzelon,T.A., Ozata,D.M., Andersson,C., Gainetdinov,I., MacRae,I.J., Tomari,Y. and Zamore,P.D. (2022) GTSF1 accelerates target RNA cleavage by PIWI-clade Argonaute proteins. *Nature*, **608**, 618–625.

36. Cheloufi,S., dos Santos,C.O., Chong,M.M. and Hannon,G.J. (2010) A dicer-independent miRNA biogenesis pathway that requires Ago catalysis. *Nature*, **465**, 584–589.

37. Cifuentes,D., Xue,H., Taylor,D.W., Patnode,H., Mishima,Y., Cheloufi,S., Ma,E., Mane,S., Hannon,G.J., Lawson,N.D., *et al.* (2010) A novel miRNA processing pathway independent of Dicer requires Argonaute2 catalytic activity. *Science*, **328**, 1694–1698.

38. Jee,D., Yang,J.S., Park,S.M., Farmer,D.T., Wen,J., Chou,T., Chow,A., McManus,M.T., Kharas,M.G. and Lai,E.C. (2018) Dual Strategies for Argonaute2-Mediated Biogenesis of Erythroid miRNAs Underlie Conserved Requirements for Slicing in Mammals. *Mol. Cell*, **69**, 265-278.e6.

39. Fromm,B., Billipp,T., Peck,L.E., Johansen,M., Tarver,J.E., King,B.L., Newcomb,J.M., Sempere,L.F., Flatmark,K., Hovig,E., *et al.* (2015) A Uniform System for the Annotation of Vertebrate microRNA Genes and the Evolution of the Human microRNAome. *http://dx.doi.org/10.1146/annurev-genet-120213-092023*, **49**, 213–242.

40. Yekta,S., Shih,I.-H. and Bartel,D.P. (2004) MicroRNA-directed cleavage of HOXB8 mRNA.

Science, **304**, 594–6.

41. Davis,E., Caiment,F., Tordoir,X., Cavaillé,J., Ferguson-Smith,A., Cockett,N., Georges,M. and Charlier,C. (2005) RNAi-mediated allelic trans-interaction at the imprinted Rtl1/Peg11 locus. *Curr. Biol.*, **15**, 743–749.

42. Karginov,F. V., Cheloufi,S., Chong,M.M.W., Stark,A., Smith,A.D. and Hannon,G.J. (2010) Diverse endonucleolytic cleavage sites in the mammalian transcriptome depend upon microRNAs, Drosha, and additional nucleases. *Mol. Cell*, **38**, 781–788.

43. Shin,C., Nam,J.W., Farh,K.K.H., Chiang,H.R., Shkumatava,A. and Bartel,D.P. (2010) Expanding the microRNA targeting code: functional sites with centered pairing. *Mol. Cell*, **38**, 789–802.

44. Dallaire,A., Frédérick,P.-M. and Simard,M.J. (2018) Somatic and Germline MicroRNAs Form Distinct Silencing Complexes to Regulate Their Target mRNAs Differently. *Dev. Cell*, **47**, 239-247.e4.

45. Park,J.H., Ahn,S., Kim,S., Lee,J., Nam,J.W. and Shin,C. (2013) Degradome sequencing reveals an endogenous microRNA target in C. elegans. *FEBS Lett.*, **587**, 964–969.

46. Moran,Y., Fredman,D., Praher,D., Li,X.Z., Wee,L.M., Rentzsch,F., Zamore,P.D., Technau,U. and Seitz,H. (2014) Cnidarian microRNAs frequently regulate targets by cleavage. *Genome Res.*, **24**, 651.

47. Stein,P., Rozhkov,N. V., Li,F., Cárdenas,F.L., Davydenk,O., Vandivier,L.E., Gregory,B.D., Hannon,G.J. and Schultz,R.M. (2015) Essential Role for Endogenous siRNAs during Meiosis in Mouse Oocytes. *PLoS Genet.*, **11**, 1–19.

48. Bouasker,S. and Simard,M.J. (2012) The slicing activity of miRNA-specific Argonautes is essential for the miRNA pathway in C. elegans. *Nucleic Acids Res.*, **40**, 10452–10462.

49. Stiernagle,T. (2006) Maintenance of C. elegans. *WormBook*, 10.1895/wormbook.1.101.1.

50. Yang,B., Schwartz,M. and McJunkin,K. (2020) In vivo CRISPR screening for phenotypic targets of the mir-35-42 family in C. elegans. *Genes Dev.*, **34**, 1227–1238.

51. Moreno-Mateos,M.A., Vejnar,C.E., Beaudoin,J.D., Fernandez,J.P., Mis,E.K., Khokha,M.K. and Giraldez,A.J. (2015) CRISPRscan: designing highly efficient sgRNAs for CRISPR-Cas9 targeting in vivo. *Nat. Methods*, **12**, 982–988.

52. Paix,A., Wang,Y., Smith,H.E., Lee,C.Y.S., Calidas,D., Lu,T., Smith,J., Schmidt,H., Krause,M.W. and Seydoux,G. (2014) Scalable and versatile genome editing using linear DNAs with microhomology to Cas9 sites in Caenorhabditis elegans. *Genetics*, **198**, 1347–1356.

53. Arribere,J.A., Bell,R.T., Fu,B.X.H., Artiles,K.L., Hartman,P.S. and Fire,A.Z. (2014) Efficient marker-free recovery of custom genetic modifications with CRISPR/Cas9 in Caenorhabditis elegans. *Genetics*, **198**, 837–46.

54. Duan,Y., Choi,S., Nelson,C. and Ambros,V. (2020) Engineering essential genes with a 'jump board' strategy using CRISPR/Cas9. *microPublication Biol.*, **2020**.

55. Martin,M. (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal*, **17**, 10–12.

56. Langmead,B., Trapnell,C., Pop,M. and Salzberg,S.L. (2009) Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.*, **10**, R25.

57. Danecek,P., Bonfield,J.K., Liddle,J., Marshall,J., Ohan,V., Pollard,M.O., Whitwham,A., Keane,T., McCarthy,S.A., Davies,R.M., *et al.* (2021) Twelve years of SAMtools and BCFtools. *Gigascience*, **10**.

58. Anders,S., Pyl,P.T. and Huber,W. (2015) HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics*, **31**, 166–169.

59. Love,M.I., Huber,W. and Anders,S. (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.*, **15**, 1–21.

60. Kim,D., Paggi,J.M., Park,C., Bennett,C. and Salzberg,S.L. (2019) Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.*, **37**, 907–915.

61. Liao,Y., Smyth,G.K. and Shi,W. (2014) featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics*, **30**, 923–930.

62. Grishok,A., Pasquinelli,A.E., Conte,D., Li,N., Parrish,S., Ha,I., Baillie,D.L., Fire,A., Ruvkun,G. and Mello,C.C. (2001) Genes and mechanisms related to RNA interference regulate expression of the small temporal RNAs that control C. elegans developmental timing. *Cell*, **106**, 23–34.

63. Brown,K.C., Svendsen,J.M., Tucci,R.M., Montgomery,B.E. and Montgomery,T.A. (2017) ALG-5 is a miRNA-associated Argonaute required for proper developmental timing in the Caenorhabditis elegans germline. *Nucleic Acids Res.*, **45**, 9093–9107.

64. Miyoshi,K., Tsukumo,H., Nagami,T., Siomi,H. and Siomi,M.C. (2005) Slicer function of Drosophila Argonautes and its involvement in RISC formation. *Genes Dev.*, **19**, 2837–2848.

65. Kim,H., Kim,J., Kim,K., Chang,H., You,K. V. and Kim,N. (2019) Bias-minimized quantification of microRNA reveals widespread alternative processing and 3' end modification. *Nucleic Acids Res.*, **47**, 2630–2640.

66. Vasquez-Rifo,A., Jannot,G., Armisen,J., Labouesse,M., Bukhari,S.I., Rondeau,E.L., Miska,E.A. and Simard,M.J. (2012) Developmental characterization of the microRNA-specific C. elegans Argonautes alg-1 and alg-2. *PLoS One*, **7**, e33750.

67. Alvarez-Saavedra,E. and Horvitz,H.R. (2010) Many families of C. elegans microRNAs are not essential for development or viability. *Curr. Biol.*, **20**, 367–373.

68. Corrêa,R.L., Steiner,F.A., Berezikov,E. and Ketting,R.F. (2010) MicroRNA-directed siRNA biogenesis in Caenorhabditis elegans. *PLoS Genet.*, **6**, e1000903.

69. Vieux,K.-F., Prothro,K.P., Kelley,L.H., Palmer,C., Maine,E.M., Veksler-Lublinsky,I. and McJunkin,K. (2021) Screening by deep sequencing reveals mediators of microRNA tailing in C. elegans. *Nucleic Acids Res.*, **49**, 11167–11180.

70. Kawamata,T., Seitz,H. and Tomari,Y. (2009) Structural determinants of miRNAs for RISC loading and slicer-independent unwinding. *Nat. Struct. Mol. Biol.*, **16**, 953–960.

71. Gu,S., Jin,L., Zhang,F., Huang,Y., Grimm,D., Rossi,J.J. and Kay,M.A. (2011) Thermodynamic stability of small hairpin RNAs highly influences the loading process of different mammalian Argonautes. *Proc. Natl. Acad. Sci. U. S. A.*, **108**, 9208–9213.

72. Sheu-Gruttadauria,J., Pawlica,P., Klum,S.M., Wang,S., Yario,T.A., Schirle Oakdale,N.T., Steitz,J.A. and MacRae,I.J. (2019) Structural Basis for Target-Directed MicroRNA Degradation. *Mol. Cell*, **75**, 1243-1255.e7.

73. Yang,A., Shao,T.-J., Bofill-De Ros,X., Lian,C., Villanueva,P., Dai,L. and Gu,S. (2020) AGO-bound mature miRNAs are oligouridylated by TUTs and subsequently degraded by DIS3L2. *Nat. Commun.*, **11**, 2765.

74. Khvorova,A., Reynolds,A. and Jayasena,S.D. (2003) Functional siRNAs and miRNAs exhibit strand bias. *Cell*, **115**, 209–216.

75. Schwarz,D.S., Hutvágner,G., Du,T., Xu,Z., Aronin,N. and Zamore,P.D. (2003) Asymmetry in the assembly of the RNAi enzyme complex. *Cell*, **115**, 199–208.

76. Medley,J.C., Panzade,G. and Zinovyeva,A.Y. (2021) microRNA strand selection: Unwinding the rules. *Wiley Interdiscip. Rev. RNA*, **12**.

77. Cazalla,D., Yario,T., Steitz,J.A. and Steitz,J. (2010) Down-regulation of a host microRNA by a Herpesvirus saimiri noncoding RNA. *Science*, **328**, 1563–6.

78. de la Mata,M., Gaidatzis,D., Vitanescu,M., Stadler,M.B., Wentzel,C., Scheiffele,P., Filipowicz,W. and Grosshans,H. (2015) Potent degradation of neuronal miRNAs induced by highly complementary targets. *EMBO Rep.*, **16**, 500–511.

79. Pawlica,P., Moss,W.N. and Steitz,J.A. (2016) Host miRNA degradation by Herpesvirus saimiri small nuclear RNA requires an unstructured interacting region. *RNA*, **22**, 1181–9.

80. Kingston,E.R. and Bartel,D.P. (2021) Ago2 protects Drosophila siRNAs and microRNAs rom target-directed degradation, even in the absence of 2-O-methylation. *RNA*, **27**, 710–724.

81. Kingston,E.R., Blodgett,L.W. and Bartel,D.P. (2022) Endogenous transcripts direct microRNA degradation in Drosophila, and this targeted degradation is required for proper embryonic development. *Mol. Cell*, **82**, 3872-3884.e9.

82. Ameres,S.L., Horwich,M.D., Hung,J.-H., Xu,J., Ghildiyal,M., Weng,Z. and Zamore,P.D. (2010) Target RNA-directed trimming and tailing of small silencing RNAs. *Science*, **328**, 1534–1539.

83. Libri,V., Helwak,A., Miesen,P., Santhakumar,D., Borger,J.G., Kudla,G., Grey,F., Tollervey,D. and Buck,A.H. (2012) Murine cytomegalovirus encodes a miR-27 inhibitor disguised as a target. *Proc. Natl. Acad. Sci. U. S. A.*, **109**, 279–284.

84. Marcinowski,L., Tanguy,M., Krmpotic,A., Rädle,B., Lisnić,V.J., Tuddenham,L., Chane-Woon-Ming,B., Ruzsics,Z., Erhard,F., Benkartek,C., *et al.* (2012) Degradation of cellular

mir-27 by a novel, highly abundant viral transcript is important for efficient virus replication in vivo. *PLoS Pathog.*, **8**, e1002510.

85. Lee,S., Song,J., Kim,S., Kim,J., Hong,Y., Kim,Y., Kim,D., Baek,D. and Ahn,K. (2013) Selective degradation of host MicroRNAs by an intergenic HCMV noncoding RNA accelerates virus production. *Cell Host Microbe*, **13**, 678–690.

86. Ghini,F., Rubolino,C., Climent,M., Simeone,I., Marzi,M.J. and Nicassio,F. (2018) Endogenous transcripts control miRNA levels and activity in mammalian cells by target-directed miRNA degradation. *Nat. Commun.*, **9**, 3119.

87. Bitetti,A., Mallory,A.C., Golini,E., Carrieri,C., Carreño Gutiérrez,H., Perlas,E., Pérez-Rico,Y.A., Tocchini-Valentini,G.P., Enright,A.J., Norton,W.H.J., *et al.* (2018) MicroRNA degradation by a conserved target RNA regulates animal behavior. *Nat. Struct. Mol. Biol.*, **25**, 244–251.

88. Kleaveland,B., Shi,C.Y., Stefano,J. and Bartel,D.P. (2018) A Network of Noncoding Regulatory RNAs Acts in the Mammalian Brain. *Cell*, **174**, 350-362.e17.

89. Li,L., Sheng,P., Li,T., Fields,C.J., Hiers,N.M., Wang,Y., Li,J., Guardia,C.M., Licht,J.D. and Xie,M. (2021) Widespread microRNA degradation elements in target mRNAs can assist the encoded proteins. *Genes Dev.*, **35**, 1595–1609.

90. Han,J., LaVigne,C.A., Jones,B.T., Zhang,H., Gillett,F. and Mendell,J.T. (2020) A ubiquitin ligase mediates target-directed microRNA decay independently of tailing and trimming. *Science*, **370**, eabc9546.

91. Shi,C.Y., Kingston,E.R., Kleaveland,B., Lin,D.H., Stubna,M.W. and Bartel,D.P. (2020) The ZSWIM8 ubiquitin ligase mediates target-directed microRNA degradation. *Science*, **370**, eabc9359.

92. Donnelly,B.F., Yang,B., Grimme,A.L., Vieux,K.F., Liu,C.Y., Zhou,L. and McJunkin,K. (2022) The developmentally timed decay of an essential microRNA family is seed-sequence dependent. *Cell Rep.*, **40**.

93. Gu,S., Jin,L., Huang,Y., Zhang,F. and Kay,M.A. (2012) Slicing-independent RISC activation requires the argonaute PAZ domain. *Curr. Biol.*, **22**, 1536–1542.

94. Zinovyeva,A.Y., Bouasker,S., Simard,M.J., Hammell,C.M. and Ambros,V. (2014) Mutations in Conserved Residues of the C. elegans microRNA Argonaute ALG-1 Identify Separable Functions in ALG-1 miRISC Loading and Target Repression. *PLoS Genet.*, **10**.

95. Zinovyeva,A.Y., Veksler-Lublinsky,I., Vashisht,A.A., Wohlschlegel,J.A., Ambros,V.R. and Horvitz,H.R. (2015) Caenorhabditis elegans ALG-1 antimorphic mutations uncover functions for Argonaute in microRNA guide strand selection and passenger strand disposal. *Proc. Natl. Acad. Sci. U. S. A.*, **112**, E5271–E5280.

96. Chatterjee,S., Fasler,M., Büssing,I. and Grosshans,H. (2011) Target-mediated protection of endogenous microRNAs in C. elegans. *Dev. Cell*, **20**, 388–96.

97. Czech,B. and Hannon,G.J. (2016) One Loop to Rule Them All: The Ping-Pong Cycle and piRNA-Guided Silencing. *Trends Biochem. Sci.*, **41**, 324–337.

Figure 1. Mutation of catalytic triad in both miRNA Argonautes does not impact physiology.
A) Schematic of miRNA Argonaute genes in *C. elegans* with positions encoding residues of catalytic tetrad in ALG-1 and ALG-2 highlighted. ALG-5 residues in equivalent positions are not conserved. *Yellow position highlights aspartate that is mutated to alanine in "AEDH" mutants. B) Fecundity assay. Each dot represents one brood; four replicates conducted shown side by side for each genotype. (Only three replicates were done for *alg-1(null)*.) All mutants were compared to wild type be one-way ANOVA with Dunnett's test. Only *alg-1(null)* shows significant difference from wild type. ****p<0.0001. Assay was performed at 25̊C.

Figure 2. Some miRNA star strands are slightly elevated in the absence of slicing. A) MA plot showing average abundance on X-axis and $\log_2$(fold change) on Y-axis. miRNAs showing significant changes in *alg-1(AEDH); alg-2(AEDH)* versus wild type are shown in purple (miRNA guide strands) or green (star strands) (DEseq2 p_{adj}<0.05). Three (L4) or four (embryo and young adult) biological replicates of each genotype were analyzed. (B) MA plot as in (A), with all guide strands shown in black and all star strands in light blue. (C) Summary of $\log_2$(fold change) values for all guides or star strands in each stage in *alg-1(AEDH); alg-2(AEDH)* versus wild type. $\log_2$(fold change) of star strands were compared to those of guide strands in the same sample by one-way ANOVA followed by Sidak's multiple comparisons test. ****p<0.0001

Figure 3. Slicing plays a greater role in star strand ejection from ALG-2 than that of ALG-1. A) MA plot showing average abundance on X-axis and $\log_2$(fold change) on Y-axis. miRNAs showing significant changes in *alg-1(AEDH)* or *alg-2(AEDH)* versus wild type embryos are shown in purple (miRNA guide strands) or green (star strands) (DEseq2 $p_{adj} < 0.05$). Four biological replicates of each genotype were analyzed. (B) MA plot as in (A), with all guide strands shown in black and all star strands in light blue.

Figure 4. High duplex stability favors a greater role for slicing in unwinding. A) Percent trimming of guide strand and duplex stability (minimum free energy) of each miRNA duplex. Frequency of star strand misregulation in *alg-1(AEDH); alg-2(AEDH)* among three stages (embryo, L4, adult) is color-coded. B) Histogram of duplex stabilities with bins containing *mir-63* and *mir-72* and their mutant variants indicated. A-B) Only miRNAs with sufficient abundance for trimming analysis in (69) are shown. C-D) Predicted structure of *mir-63* and *mir-72* and engineered mutations in each duplex. E-F) Absolute quantification by qPCR of *mir-63* and *mir-72* guide and star strands in L4 samples (three biological replicates for each genotype). G-H) Ratio of guide to star strand calculated from absolute quantification. E-H) Unpaired t-tests, **p-value < 0.01, *p-value < 0.05.

Figure 5. Minor changes in star strand abundance do not result in changes in target repression. MA plots showing average abundance (of all genes > 10 reads per million) on X-axis and log2(fold change) on Y-axis in *alg-1(AEDH); alg-2(AEDH)* versus wild type. Only 1, 0, and 2 genes are significantly altered in expression in embryo, L4 and adult, respectively (DEseq2 p_{adj}<0.05). Three (L4) or four (embryo and young adult) biological replicates of each genotype were analyzed.

Figure 6. Down-regulation of select guide strands in the absence of slicing is EBAX-1-independent. A-B, E) MA plots showing average abundance on X-axis and $\log_2$(fold change) on Y-axis. miRNAs showing significant changes are shown in light blue (DEseq2 $p_{adj}<0.05$). Genotypes being compared are indicated in graph title. C) Correlation of changes induced by *alg-1(AEDH); alg-2(AEDH)* in wild type (X-axis) or *ebax-1(null)* background (Y-axis). D) Level of *mir-243* in each sample after default DEseq2 normalization in pairwise comparisons shown. One-way ANOVA was performed, and p-values are from post hoc Sidak's multiple comparisons test. C,F) Only miRNAs with baseMean > 50 in wild type experiment 1 are included. (A-F) All samples are young adults. Four biological replicates of each genotype were analyzed.

Figure 7. Model of observed effects on miRNA star strands in slicing mutants. A small portion of miRISC is matured by the slicing-dependent unwinding pathway. In *alg-1(AEDH);alg-2(AEDH)* mutant backgrounds, this small portion of miRISC retains the star strand (leading to elevated star strand levels). The major portion of miRISC is matured in a slicing-independent manner, thus leaving overall miRISC function largely intact in the mutant background. Slicing-dependent unwinding is most apparent in embryos in which ALG-2 is highly expressed and for certain thermodynamically-stable miRNA:star duplexes like *mir-63* and *mir-72*. For most miRNAs in L4 and adult samples, slicing-dependent unwinding is negligible.

List of Supplemental Materials

Figure S1. Spike in normalization supports minor changes in miRNA abundances
Figure S2. Standard curves for absolute miRNA quantification by qPCR
Figure S3. Polymorphism in *gpap-1* 5' UTR likely contributes to decreased expression
Figure S4. Reproducibility of miRNA changes in *ebax-1(null)* and across two *alg-1(AEDH); alg-2(AEDH)* experiments

Table S1. Strains used in this study
Table S2. Alleles generated and oligonucleotides used in this study
Table S3. Samples used for deep sequencing
Table S4. Raw miRNA and spike-in reads for all samples
Table S5. Results of small RNA sequencing analysis in embryo samples*: alg-1(AEDH); alg-2(AEDH)* versus wild type
Table S6. Results of small RNA sequencing analysis in L4 samples: *alg-1(AEDH); alg-2(AEDH)* versus wild type
Table S7. Results of small RNA sequencing analysis in adult samples: *alg-1(AEDH); alg-2(AEDH)* versus wild type (experiment 1 samples)
Table S8. Results of small RNA sequencing analysis in embryo samples: *alg-1(AEDH)* versus wild type
Table S9. Results of small RNA sequencing analysis in embryo samples: *alg-2(AEDH)* versus wild type
Table S10. Results of RNA-seq analysis in embryo samples: *alg-1(AEDH); alg-2(AEDH)* versus wild type
Table S11. Results of RNA-seq analysis in L4 samples: *alg-1(AEDH); alg-2(AEDH)* versus wild type
Table S12. Results of RNA-seq analysis in adult samples: *alg-1(AEDH); alg-2(AEDH)* versus wild type
Table S13. Results of small RNA sequencing analysis in adult samples: *ebax-1(null)* versus wild type
Table S14. Results of small RNA sequencing analysis in adult samples: *ebax-1(null); alg-1(AEDH); alg-2(AEDH) versus ebax-1(null)*
Table S15. Results of small RNA sequencing analysis in adult samples: *alg-1(AEDH); alg-2(AEDH)* versus wild type (experiment 2 samples)
Table S16. Results of small RNA sequencing analysis in adult samples: *Y47(mir-243bs_del); alg-1(AEDH); alg-2(AEDH)* versus *Y47(mir-243bs_del)*

Supplemental Figures

Figure S1. Spike in normalization supports minor changes in miRNA abundances. A) MA plot showing average abundance on X-axis and log$_2$(fold change) on Y-axis. miRNAs showing significant changes in *alg-1(AEDH); alg-2(AEDH)* versus wild type are shown in purple (miRNA guide strands) or green (star strands) (DEseq2 p$_{adj}$<0.05). Reads were normalized to exogenous spike-ins (See methods and Table S2). Three (L4) or four (embryo and young adult) biological replicates of each genotype were analyzed. (B) MA plot as in (A), with all guide strands shown in black and all star strands in light blue. (C) Summary of log$_2$(fold change) values for all guides or star strands in each stage in *alg-1(AEDH); alg-2(AEDH)* versus wild type. Log$_2$(fold change) of star strands were compared to those of guide strands in the same sample by one-way ANOVA followed by Sidak's multiple comparisons test. ***p<0.0001

mir-63-3p	y = -1.0665x + 6.4959	R² = 0.9907
mir-63-5p	y = -1.0633x + 6.0536	R² = 0.9816
mir-63-3p_mut2	y = -1.0993x + 7.2333	R² = 0.9988
mir-63-5p_mut2	y = -1.0436x + 6.4644	R² = 0.9944
mir-63-3p_mut3	y = -1.074x + 5.6509	R² = 0.9990
mir-63-5p_mut3	y = -1.26x + 8.1393	R² = 0.9660

mir-72-3p	y = -1.1342x + 8.7734	R² = 0.9998
mir-72-5p	y = -1.1278x + 9.12	R² = 0.9933
mir-72-3p_mut1	y = -1.0592x + 7.7451	R² = 0.9996
mir-72-5p_mut1	y = -1.0799x + 5.5879	R² = 0.9992
mir-72-3p_mut2	y = -1.0636x + 7.2479	R² = 0.9998
mir-72-5p_mut2	y = -1.0377x + 9.2635	R² = 0.9995

Figure S2. Standard curves for absolute miRNA quantification by qPCR. Synthetic RNA oligonucleotides were diluted to indicated concentrations, and 1.66µl was used in 5µl RT reactions and downstream miRNA-Taqman as indicated in methods. Average of three technical replicates is shown. Highest and lowest concentrations were excluded from the curve if they were outside the linear range. All experimental sample concentrations fell within the linear range of the respective assay.

Figure S3. Polymorphism in *gpap-1* 5' UTR likely contributes to decreased expression. Only one gene was identified as mis-regulated across more than one developmental stage in the *alg-1(AEDH);alg-2(AEDH)* mutant strain. Close examination of RNA-seq data revealed that this gene, *gpap-1*, contains a polymorphism in its 5' UTR in the mutant strain.

Figure S4. Reproducibility of miRNA changes in *ebax-1(null)* and across two *alg-1(AEDH); alg-2(AEDH)* experiments. A) MA plots showing average abundance on X-axis and $\log_2$(fold change) on Y-axis. miRNAs showing significant changes are highlighted in light blue or purple (DEseq2 $p_{adj}<0.05$); upregulated miRNAs that were previously noted in Shi, et al are in purple. Four biological replicates of each genotype were analyzed. B) Correlation of changes induced by *alg-1(AEDH); alg-2(AEDH)* in wild type experiment 1 (X-axis) or experiment 2 (Y-axis). Each experiment consists of four biological replicates of each genotype. Only miRNAs with baseMean > 50 in wild type experiment 1 are included. (A-B) All samples are from young adults.