Title: Plasticity in the functional properties of NMDA receptors improves network

stability during severe energy stress

Abbreviated Title: State-dependent NMDAR plasticity improves hypoxia tolerance

Authors: Nikolaus Bueschke[1†], Lara Amaral-Silva[1†], Min Hu[2], Alvaro Alvarez[2], Joseph M. Santin[1]*

Author Affiliations: [1]University of Missouri, Columbia, MO 65201; [2]University of North Carolina-Greensboro, Greensboro, NC 27402

***Email for correspondence**: santinj@missouri.edu
[†] Indicates co-first authorship

Number of figures: 7

Number of Words in Abstract: 219

Number of Words in Introduction: 634
Number of Words in Discussion: 1,499

Conflict of Interests: The authors declare no conflicts of interest.

Acknowledgment: We would like to thank the National Institutes of Health (R15NS112920-01A1, R01NS114514) the U.S. Department of Defense (W911NF2010275) for funding to JMS.

Abstract

Brain energy stress leads to neuronal hyperexcitability followed by a rapid loss of

function and cell death. In contrast, the frog brainstem switches into a state of extreme

metabolic resilience that allows them to maintain motor function during hypoxia as they

emerge from hibernation. NMDA receptors (NMDARs) are Ca^{2+}-permeable glutamate

receptors that contribute to the loss of homeostasis during hypoxia. Therefore, we

hypothesized that hibernation leads to plasticity that reduces the role of NMDARs within

neural networks to improve function during energy stress. To test this, we assessed a

circuit with a large involvement of NMDAR synapses, the brainstem respiratory network

of female bullfrogs, *Lithobates catesbeianus*. Contrary to our expectations, hibernation

did not alter the role of NMDARs in generating network output, nor did it affect the

amplitude, kinetics, and hypoxia sensitivity of NMDAR currents. Instead, hibernation

strongly reduced NMDAR Ca^{2+} permeability and enhanced desensitization during

repetitive stimulation. Under severe hypoxia, the normal NMDAR profile caused network

hyperexcitability within minutes, which was mitigated by blocking NMDARs. After

hibernation, the modified complement of NMDARs protected against hyperexcitability,

as disordered output did not occur for at least one hour in hypoxia. These findings

uncover state-dependence in the plasticity of NMDARs, whereby multiple changes to

receptor function improve neural performance during energy stress without interfering

with its normal role during healthy activity.

Significance Statement:

Neural circuits lose homeostasis during severe energy stress, and NMDA-glutamate

receptors play a major role in this response. In contrast, frogs have the remarkable

capacity to use plasticity that improves circuit function from minutes to hours during

hypoxia, likely as an adaptation to survive emergence from hibernation. We found this

occurs, in part, through modification of NMDA receptors that renders them less

permeable to Ca^{2+} and more likely to desensitize during high activity states. These

NMDA receptor modifications do not influence normal network function but protect

against hyperexcitability caused by hypoxia. This work points to endogenous plasticity

mechanisms that improve network function during energy stress without altering circuit

function when the brain is well-oxygenated.

Introduction:

Brain function requires high rates of ATP synthesis (Bordone et al., 2019). The

sudden loss of oxygen triggers hyperexcitability that leads to ion dysregulation and cell

death, termed "excitotoxicity" (Buck & Pamenter, 2018). A key contributor to

excitotoxicity involves the activation of NMDA-glutamate receptors (NMDARs), causing

pathological Ca^{2+} influx and saturation of intracellular Ca^{2+} buffering systems

(Szydlowska & Tymianski, 2010; White & Reynolds, 1995). As rates of ATP synthesis

wane, active transport of Ca^{2+} out of the cell becomes increasingly difficult, which

culminates in cell death through reactive oxygen species, pro-death signaling pathways,

and mitochondrial damage (Wu & Tymianski, 2018). Therefore, hyperexcitability caused

by overstimulation of NMDARs and subsequent Ca^{2+} influx represents a critical step in

the loss of homeostasis that drives neurological issues during hypoxia.

Unlike most mammals, some species experience variable oxygen tensions in

their environments and have evolved strategies to survive brain hypoxia (Larson et al.,

2014). Survival strategies often involve entry into a hypometabolic state associated with

arrested synaptic transmission to conserve energy (Buck & Pamenter, 2018). However,

some animals must remain active during hypoxia (Czech-Damal et al., 2014; Larson et

al., 2014; Larson & Park, 2009). This presents the challenge of maintaining circuit

function while avoiding hyperexcitability and excitotoxicity. An extreme example of this

problem is embodied by the respiratory network of American bullfrogs. This network

generates rhythmic output through mechanisms that involve AMPA and NMDA-

glutamate receptors (Kottick et al., 2013) and, therefore, requires ongoing aerobic

metabolism (Adams et al., 2021). However, for several months each year, frogs

hibernate in ice-covered ponds without breathing air, using only skin gas exchange. As

a consequence, blood oxygen falls dramatically, which may be as low as 1-3 mmHg

(Tattersall & Ultsch, 2008). Low oxygen in this environment does not pose an immediate

threat due to reduced metabolic rates in the cold (Tattersall & Boutilier, 1997). However,

during emergence at warm temperatures, this life sustaining network, must restart

activity on the background of severe hypoxia. If this was not already difficult enough, an

additional problem lies in the fact that frogs are not generally considered to be strongly

"hypoxia-tolerant," and low O_2 levels present during emergence cannot power brainstem

circuits (Adams et al., 2021). To overcome these challenges, hibernation induces

metabolic plasticity at synapses that improve network function during severe hypoxia

from a few minutes to several hours (Bueschke et al., 2021, Amaral-Silva and Santin,

2023; Hu and Santin, 2022). Therefore, hibernation in frogs provides insight into

plasticity that shifts a typically "hypoxia-intolerant" circuit into a state that functions

remarkably well during severe energy stress.

As in mammals, hypoxia in the frog brain induces network hyperexcitability that

disrupts patterned output, followed by a swift loss of function (Adams et al., 2021;

Bueschke et al., 2021a). Thus, this network must engage mechanisms that constrain

excitability to maintain activity with a limited energy supply upon emergence from

hibernation. Many hypoxia-tolerant vertebrates have low levels of NMDARs or suppress

NMDARs in hypoxia to conserve energy and avoid excitotoxicity (Bickler et al., 2000;

Bickler & Buck, 2007; Wilkie et al., 2008). Therefore, we hypothesized that shifting from

"hypoxia intolerance" to "functional hypoxia tolerance" involves reduced NMDAR

function within the network. To test this hypothesis, we assessed the NMDAR tone of

the respiratory network, NMDAR currents using whole-cell voltage-clamp, and NMDAR

subunit composition using single-cell quantitative PCR. In contrast to our hypothesis,

hibernation did not influence the role of NMDARs in the network, current amplitude,

kinetics, and hypoxia sensitivity. We instead found that hibernation decreased the Ca^{2+}-

permeability of NMDARs and enhanced desensitization, serving to reduce Ca^{2+} influx

and lower depolarizing drive during high activity states. These modifications oppose the

loss of homeostasis driven by the normal profile of NMDARs. Overall, we identified

NMDAR plasticity that improves network activity during energy stress without

influencing their contribution in well-oxygenated conditions.

Materials and Methods:

Animal husbandry and ethical approval:

The use of animals was approved by the Institutional Animal Care and Use Committee (IACUC) at the University of North Carolina at Greensboro (Protocol #19-006). Female American bullfrogs (*Lithobates catesbeianus*) were ordered from Phrog Farm (Twin Falls, ID, USA). Frogs were randomly assigned to one of two groups: control or hibernation and were placed in designated plastic tubs containing dechlorinated water bubbled with room air. Control frogs were kept at ambient room temperatures (20˚C) with access to wet and dry areas and fed pellets provided by Phrog Pharm once a week. Plastic tubs containing frogs assigned for aquatic overwintering were placed into low-temperature incubators (Thermo Fisher Scientific, Waltham, MA, USA) that were gradually reduced from 20˚ to 4˚ C over the course of 7 days. Once at overwintering temperatures, screens were placed below water level to ensure frogs' cutaneous respiration as the primary oxygen exchange, and they were kept at 4˚ C for 30 days before use. Frogs exhibit metabolic suppression at these temperatures, consistent with hibernation (Tattersall & Ultsch, 2008). Therefore, we refer to this group as "hibernation." All frogs were kept under a 12:12 hour light/dark cycle.

Dissection of Brainstem:

Frogs were deeply anesthetized with isoflurane (1 mL) in a sealed 1 L container until loss of toe-pinch response. They were then rapidly decapitated, and heads were submerged for dissection in chilled artificial cerebrospinal fluid (aCSF; 104 mM NaCl, 4 mM KCl, 1.4 mM $MgCl_2$, 7.5 mM D-glucose, 1 mM NaH_2PO_4, 40 mM $NaHCO_3$, and 2.5

mM $CaCl_2$; (Bueschke et al., 2021), which was bubbled with 98.5% O_2 and 1.5% CO_2

for oxygenation, resulting in aCSF with a pH of 7.9 ± 0.1. These CO_2 and pH values are

normal for frogs at room temperature (Howell et al., 1970). The brainstem-spinal cord

was rapidly exposed, and the forebrain was crushed. Brainstem-spinal cord nerve roots

were carefully trimmed, allowing its' excision from the cranium, which was followed by

the dura membrane removal. Following all dissections, the preparations were held at

room temperature ($22\pm1^{\circ}C$) and all experiments were performed at this temperature.

Extracellular nerve root recordings:

119   For experiments that assessed motor output from the intact network, we

assessed extracellular motor output from the vagus nerve rootlets, which innervates the

glottal dilator muscle to control airflow in and out of the lung. Dissected preparations

were pinned ventral side up in 6 mL Petri dishes coated with Sylgard 184 (Dow Inc.

Midland, MI, USA). Brainstem-spinal cords were continuously superfused with

oxygenated aCSF using a peristaltic pump (Watson Marlow, Falmouth, CNL, UK).

Population activity from the vagal nerve root (CNX) was recorded using a suction

electrode attached to a fire-polished borosilicate glass pulled from a horizontal pipette

puller (Sutter Instrument, Novato, CA, USA). Extracellular signals were amplified

(x1000) and filtered (low-pass, 1000 Hz; high pass, 100 Hz) using an AM-Systems 1700

amplifier (Sequim, WA, USA), then digitized using Powerlab 8/35 (ADInstruments,

Dunedin, New Zealand). The raw signal was integrated and rectified (100 ms τ) using

LabChart data acquisition system (ADInsruments). After 4 hours of decapitation, a

stable signal was recorded for ~30 minutes, and the preparation was challenged with

hypoxic aCSF (98.5% N_2 balanced with CO_2) or NMDAR antagonist, D-AP5 (50 µM). All

preparations included in this study produced rhythmic motor output associated with lung

breathing.

Motoneuron labeling and slice preparation:

For experiments that assessed NMDA receptor currents in motoneurons from

brain slices, we isolated the 4th root of the glossopharyngeal-vagus complex and

attached a fire-polished borosilicate glass pipette to it. Tetramethylrhodamine dextran

dye was then added to the tip of the pipette (Invitrogen, Waltham, MA, USA) in contact

to the nerve root for at least 2 hours to diffuse to the soma. We then sliced the

brainstem-spinal cord preparations at 300 μM using a vibratome (Technical Products

International series 1000, St Louis, MO, USA). During dye loading and slicing the tissue

was maintained in regular aCSF (described above).

NMDAR Currents:

Slices were superfused with an extracellular solution (76.5 mM NaCl, 2.2 mM

KCl, 7.5 mM D-Glucose, 10 mM HEPES, 300 μM $CdCl_2$, 20 mM TEA-Cl, 250 nM TTX,

10 μM DNQX) containing two different Ca^{2+} concentrations. High Ca^{2+} had 10 mM of

$CaCl_2$, and low Ca^{2+} 1 mM of $CaCl_2$. Sucrose was added to maintain consistent

osmolarity (330 mOsm). Thus, 60mM was added to the high Ca^{2+} solution, and 89mM

was added to the low Ca^{2+} solution. $MgCl_2$ was excluded to prevent Mg^{2+} block of

NMDARs. Extracellular solutions were oxygenated by bubbling 98.5% O_2 and 1.5%

CO_2.

Labeled vagal neurons were approached by glass pipettes (2–4 MΩ resistance)

using a micromanipulator (MP-285/ MPC-200, Sutter Instruments, Novato, CA, USA)

attached to a head stage (CV203BU, Molecular Devices, San Jose, CA, USA). Positive

pressure was applied to the pipette while approaching the cell and quickly removed,

gentle negative pressure was used to form a >1GΩ, and the whole-cell access was

obtained by breaking the seal with rapid negative pressure. We did not observe any

obvious differences in cell viability and ability to obtain patch clamp recordings in high

and low Ca^{2+} solutions. The solution filling the patch pipette (76.5 mM K-gluconate, 10

mM D-glucose, 10 mM HEPES, 1 mM Na_2-ATP, 0.1 mM Na_2-GTP, 2 mM $MgCl_2$, 30 mM

TEA-Cl) was designed with Na^+ and K^+ concentrations equal and opposite to the

extracellular solution to generate a reversal (E_{rev}) potential of ~0 mV without Ca^{2+}

(assuming no difference in ion selectivity between Na^+ and K^+) as seen previously

(Jatzke et al., 2002).

To evoke NMDAR currents, we focally applied NMDA and glycine to the cell

body. For this, a borosilicate pipette pulled with a tip diameter of ~5 μm was filled with

NMDA (1 mM) and glycine (10 μM) (both from Hellobio, Princeton, NJ, USA) dissolved

in the extracellular solution. The pipette wasdriven by a Picospritzer II, and the focal

solution was applied during 10-20 ms onto the soma of labeled vagal motoneurons

(General Valve Corporation, Fairfield, NJ, USA) to activate NMDARs. Glycine is a co-

agonist of the NMDAR but did not elicit a response when applied alone in control or

hibernation neurons.

NMDAR E_{rev} was recorded by a voltage clamp step protocol with a Δ+5 mV step

between -20 mV and 40 mV. In 3 recordings, E_{rev} was more hyperpolarized, and those

cells were stepped from -40 mV to 20 mV in 5 mV increments (control, n=16;

hibernation, n=16). NMDAR desensitization was assessed by puffing at a rate of 0.5 Hz

for a total of ten pulses at -20 mV (control, n=12; hibernation, n=14). Hypoxia was

applied to brain slices by bubbling extracellular solution with 98.5% of N_2 and 1.5% of

CO_2 for ~5 minutes before recording. All desensitization protocols and single activation

amplitude and kinetics experiments were performed on neurons held at -20 mV. All data

were acquired in pClamp 11 software using Axopatch 200B amplifier and Axon Digidata

1550B digitizer (Molecular Devices, San Jose, CA, USA).

NMDAR subunits electrophysiology:

Vagus motoneurons were recorded using patch-clamp as described above. For

this protocol, the pipette solution contained 110 mM K-gluconate, 2 mM $MgCl_2$, 10 mM

HEPES, 1 mM Na_2-ATP, 0.1 mM Na_2-GTP, and 2.5 mM EGTA. Once gaining whole cell

access the solution bathing the slice was changed from regular aCSF to 104 mM NaCl,

4 mM KCl, 7.5 mM D-glucose, 1 mM NaH_2PO_4, 40 mM $NaHCO_3$, 2.5 mM $CaCl_2$, 250 nM

TTX, 10 µM DNQX, 3 µM Glycine, and 2 µM Strychnine. NMDA currents were elicited

once every minute by focal application of NMDA (1 mM) and glycine (3 µM) diluted in

the aCSF in 500 ms pulses. The cell was continuously monitored in voltage clamp, and

after observing a stable current in 3 consecutive NMDA puffs (~ 10-15 min of recording),

we applied a specific NMDA subunit inhibitor in aCSF.

The subunit GluN2A was inhibited using 1 µM TCN 201 (Tocris Bioscience,

Bristol, UK) (Bettini et al., 2010; Edman et al., 2012); GluN2B was inhibited by 2 µM RO

25-6981 (Tocris Bioscience, Bristol, UK) (Abrahamsson et al., 2017; France et al.,

2017); GluN2C/GluN2D was inhibited using 10 µM QZN 46 (Tocris Bioscience, Bristol,

UK) (Hansen and Traynelis, 2011); and GluN3 was blocked by 30 µM TK30 (4-(2,4-

dichlorobenzoyl)-1H-pyrrole-2-carboxylic acid; Santa Cruz Biotechnology, Dallas, TX,

USA) (Kvist et al., 2013; Christian et al., 2021). A time control experiment was

performed recording NMDAR currents while maintaining the slice in the aCSF described

above with no inhibitor added.

Electrophysiology Data analysis:

<u>Extracellular motor output</u>: Burst amplitude and frequency were determined by

averaging amplitude of fictive breaths selected in a 5-minute window prior to D-AP5

application. Amplitude and frequency in the presence of NMDAR antagonist were

analyzed for 5 minutes after burst amplitude stabilized. Peak amplitude was determined

using the Peak Analysis extension in LabChart (ADInstruments, Dunedin, New

Zealand). To characterize chaotic bursting behavior during hypoxia, we analyzed the

peak of the largest "non-respiratory" motor burst that clearly disrupted patterned

network output associated with respiratory activity and normalized it to the background

nerve signal value at the baseline to allow comparisons across groups.

<u>NMDAR Ca^{2+} permeability</u>: Relative Ca^{2+} permeability was determined by the

degree of shift in the NMDAR reversal potential (E_{rev}) based on the mean data of the

population in different Ca^{2+} concentrations in extracellular recording solution. The

underlying premise is that receptors that are permeable to Ca^{2+} have a depolarizing shift

in E_{rev}, while Ca^{2+}-impermeable channels do not (Jatzke et al., 2002). We evoked

NMDAR currents using low (1 mM) and high (10 mM) Ca^{2+} and measured the peak

current evoked by NMDA-glycine at each voltage step. We then plotted the current-

voltage relationship and interpolated E_{rev} through the x-intercept of the line of best fit

from 3-5 plot points near the intercept where I=0 pA. The currents elicited by NMDA-

glycine were small near E_{rev} and were unlikely to be affected by voltage errors due to

the series resistance (R_s). However, the holding current at depolarized voltages was

often substantial, even in the presence of TEA to block outward currents. Therefore, we

corrected the holding voltage for R_s errors. For this, we measured R_s at each voltage

step based on the peak of the transient current and corrected the holding voltage by the

voltage error caused by R_s. Currents were measured using the Peak Analysis extension

in LabChart. Averaging/decimation at 0.05-0.1 ms was applied to the traces prior to

peak analysis to filter out high-frequency spontaneous synaptic activity. This level of

filtering was chosen as it did not alter the amplitude of the NMDAR current.

<u>Single NMDAR currents and desensitization</u>: Single NMDAR currents were

measured using similar methods as the NMDAR Ca^{2+} permeability, where peak currents

induced by NMDA and glycine were assessed at -20 mV. Decay time constant was

determined between 90% and 10% of the peak height and calculated using the Peak

Analysis tool in LabChart. Desensitization was assessed by measuring the peak current

evoked at each step. The baseline for each current in the series was taken as the

recovered current after proceeding puff.

NMDAR subunit inhibition: The current amplitude was analyzed using peak

analysis in LabChart. The average of the last 3 currents before inhibitor application was

compared to currents recorded after 10 minutes of exposure to the drug. A set of time

control experiments compared the 3 first stable currents (parameter used to decide to

apply the inhibitor) to 10 min after that.

Single-cell Real-time quantitative PCR:

We used single-cell quantitative PCR (qPCR) to determine mRNA expression for

NMDARs subunits using the same cell harvesting, RNA extraction, cDNA synthesis,

and preamplification procedures detailed in Pellizzari et al., 2023.

PCR primers for open reading frames of the genes that code for NMDA

glutamate receptor subunits, *Grin1, Grin2a, Grin2b, Grin2c, Grin2d, Grin3a, and Grin3b,*

were designed from sequences found in the coding DNA sequence for *Lithobates*

catesbeianus (Table 1 and Table 2). For this, we used annotated amino acid sequences

for GluN subunits from *Rana temporaria* as a query in the *Lithobates catesbeianus*

amino acid database. This search revealed peptide sequences with high amino acid

sequence conservation. We then performed a reciprocal BLAST against the entire

nonredundant protein database using hits from *Lithobates catesbeianus* to verify the

identity of the target. Accession numbers were then used to identify the open reading

frame in the CDS to design PCR primers. Only *Grin1, Grin2a, and Grin3a* were

identified in the bullfrog CDS, likely due to low coverage and/or poor assembly of the *L.*

catesbeiansus genome (Hammond et al., 2017). For *Grin2b, Grin2c, Grin2d, and*

Grin3b, we found the regions of the coding sequence in *Rana temporaria* (a species

closely related to *L. catesbeianus*) and *N. parkeri* (more distantly related frog species)

with high similarity to design PCR primers. Our rationale was that close sequence

identity at the nucleotide level between these two species (~98%) would allow us to

design primers for use in *Lithobates catesbeianus*.

Primer sets were validated with a series of four four-fold dilutions of brainstem

cDNA. All primer sets used here produced efficiencies greater than 80% (Table 1 and

Table 2) and a single peak in the melt curve in a SYBR Green assay, suggesting the

amplification of a single PCR product. As we observed in Pellizzari et al., 2023 some

primer sets that produced one peak in the melt curve using bulk brain cDNA as the input

material showed multiple peaks in the melt curve after single-cell preamplification.

When this occurred, these primer sets were redesigned.

All neurons in this experiment were assessed for the expression of each of the 7

Grin subtypes. *Grin2b, Grin2c, Grin2d, and Grin3b* were run using SYBR Green assays,

and *Grin1, Grin2a, and* Grin3a were run in one multiplex assay. For SYBR Green

assays, quantitative PCR was run in 10 µL reaction volumes containing 2.5 µM forward

and reverse primers and followed the instructions of the 2X SYBR Green Mastermix

(Applied Biosystems, ThermoFisher Scientific, Waltham, MA). Assays were run on 96-

well plates on an Applied Biosystems QuantStudio 3 (Applied Biosystems,

ThermoFisher Scientific, Waltham, MA) using the following cycling conditions according

to the SYBR Green instructions: 50°C-2m, 95 °C-10 m, 95 °C-15 s, 60 °C-1 m. Following

40 cycles of PCR (95°C-15 s, 60°C-1 m), melt curves for all PCR products were

acquired by increasing the temperature in increments of 0.3°C for 5 seconds from 60°C

to 95 °C. For multiplex assays, we ran triplexed probed-based assays. For this, we used

the same primer concentration as described for SYBR Green assays, 312.5 nM reporter

probes, and followed the instructions of the 5X PerfeCTa qPCR Toughmix mastermix

(Quanta Bio). 18s ribosomal RNA was run to ensure the quality of the sample and for

normalization of copy number to account for the possibility of different amounts on

cDNA input and efficiency in the cDNA synthesis reaction.

Absolute quantitation of transcript abundance was estimated through copy

number standard curves as previous described (Santin & Schulz, 2019). We normalized

absolute copy number by a normalization factor using 18s Cq values to account for the

possibility of different amounts of harvested cytoplasm and efficiencies of the cDNA

synthesis reaction across samples (Garcia et al., 2018). We ran qPCR for all *Grin*

subunits on 19 control neurons and 20 hibernation neurons. In some neurons, *Grin*

subunits appeared to be absent, or to be very lowly expressed. However, it is also

possible a lack of detection represents a false negative due to stochasticity in the cDNA

synthesis reaction due low RNA input quantities associated with the single cell. Thus,

we included *Grin* genes in analysis that had C_q values <29.5 after preamplification. The

only exception to this was for *Grin2D.* The abundance was consistently low for most

samples; therefore, we included all data points in the analysis for *Grin2D*.

Statistical analysis:

Statistical analysis was performed using GraphPad Prism 9 (GraphPad Software,

San Diego, CA, USA). Comparisons between independent 2 groups were carried out

with a two-tailed unpaired t-test. With 3 or more groups, we used a one-way ANOVA. In

experiments with a "before-after" design, we used a two-tailed paired t-test. In

experiments with two main effects, we used a two-way ANOVA. One-way and two-way

ANOVA were followed up with Holm-Sidak multiple comparison test when appropriate.

Individual data points were presented in addition to mean±SD or with box-and-whisker

plots, on the latter, boxes represent the interquartile range, and whiskers indicate

maximum and minimum values in the data set. Significant was accepted when p<0.05.

Results:

Results:

Function of the brainstem respiratory network in severe hypoxia increases by

hours after animals emerge from hibernation (Bueschke et al., 2021; Amaral-Silva &

Santin, 2023). To determine if reductions in NMDAR function play a role in this

response, we assessed the NMDAR tone in *in vitro* brainstem-spinal cord preparations.

This preparation produces rhythmic output associated with breathing that can be

recorded from the cranial nerve X rootlet (Fig. 1A). NMDARs are involved in transmitting

synaptic input from the respiratory rhythm generator to the motor pools (Kottick et al.,

2013). If the contribution of NMDAR was reduced following hibernation, we expected to

observe a lower sensitivity to the block of NMDARs using D-AP5 relative to controls. D-

APV reduced the amplitude of the motor output by ~50%, corroborating previous results

demonstrating that NMDAR transmission plays a large role in recruiting motoneurons to

the population burst (Kottick et al., 2013). However, sensitivity of the motor amplitude to

D-APV did not change after hibernation (Fig. 1B-C, $t_{(13)}$ = 0.9325, p = 0.37, unpaired t-

test). In addition, sensitivity of respiratory burst frequency to D-APV, which provides an

assessment of rhythm generator function, did not change after hibernation (Fig. 1D, $t_{(13)}$

= 1.699, p = 0.11, unpaired t-test). These results show that the overall contribution of

NMDARs to network function is unchanged following hibernation.

Although the contribution of NMDAR to network function did not change, we

hypothesized that functional properties of the NMDAR may play a role in improving

robustness in hypoxia. For example, plasticity in current amplitude or kinetics (*i.e.*,

smaller currents with fast deactivation times) may lower ionic fluxes through NMDARs,

and, therefore, play a role in energy conservation. Moreover, some hypoxia-tolerant

species contain NMDARs that are inhibited by hypoxia, offsetting excitotoxicity during

energetic stress (Bickler et al., 2000). Therefore, we measured NMDARs currents in

identified vagal motoneurons from brain slices using whole-cell voltage clamp in

response to focal application of the NMDAR co-agonists, 1 mM NMDA and 10 µM

glycine (Fig. 2A). Current amplitude and deactivation time constants at -20 mV did not

change in response to hibernation (Fig 2B-D, amplitude: $t_{(24)} = 1.502$, $p = 0.1461$,

unpaired t-test; deactivation time constant: $t_{(24)} = 0.4516$, $p = 0.6556$, unpaired t-test). In

addition, superfusion of aCSF bubbled with a hypoxic gas mixture (0% O_2) for 5 minutes

did not consistently alter the amplitude or deactivation time in controls (Fig 2E,

amplitude: $t_{(4)} = 1.455$, $p = 0.2193$, paired t-test; deactivation time constant: $t(4) = 1.980$,

$p = 0.1189$, paired t-test) or following hibernation (Fig 2F, amplitude: $t_{(3)} = 0.2362$,

$p=0.8285$, paired t-test; deactivation time constant: $t_{(3)} = 0.1515$, $p=0.8892$, paired t-

test). Therefore, hibernation does not alter whole-cell NMDAR currents, as well as their

deactivation kinetics and sensitivity to acute hypoxia.

Ca^{2+} influx through NMDARs plays a role in network dysfunction during hypoxia

(Szydlowska and Tymianski, 2010). Thus, we assessed the Ca^{2+} permeability of

NMDARs before and after hibernation, as Ca^{2+} selectivity may increase or decrease

without obvious changes in the whole-cell current amplitude and kinetics (Murphy et al.,

2014; Skeberdis et al., 2006). For this, we measured the reversal potential of the

NMDAR current (E_{rev}) in low Ca^{2+} (1 mM) and high Ca^{2+} (10 mM). Changes in E_{rev} during

exposure to high Ca^{2+} concentrations provide a way to assess the relative Ca^{2+}

permeability of ligand-gated ion channels (Jatzke et al., 2002). Consistent with the

canonical role of NMDAR as a Ca^{2+} permeable channel, raising extracellular Ca^{2+}

depolarized E_{rev} of in control neurons (Fig. 3A top). Divalent cations can adhere to the

outer edge of cell membranes and alter the surface charge, making the apparent

voltage at the membrane more depolarized with increasing concentrations of divalent

cations (Hille, 2001). This is important to consider, given that we found a depolarizing

shift in E_{rev} in response to high Ca^{2+}. However, charge screening is unlikely to affect our

interpretation since E_{rev} shows a <2 mV difference between 1 mM and 10 mM Ca^{2+} on

Ca^{2+}-impermeable kainite receptors (Jatzke et al., 2002). Thus, most of the ~13 mV

depolarization we observe between low and high Ca^{2+} in control neurons likely reflects

its true Ca^{2+} permeability and not changes in the surface charge. After hibernation, the

change in E_{rev} during high Ca^{2+} appeared to be far smaller (Fig. 3A bottom). Indeed,

there was a significant interaction between Ca^{2+} concentration and group on E_{rev} (Fig.

3C, $F_{(1,60)}$ = 7.735, p = 0.0072, two-way ANOVA), with a strong increase in E_{rev} by high

Ca^{2+} in the control group and no significant change in E_{rev} in high Ca^{2+} after hibernation.

Therefore, hibernation leads to a reduction in the Ca^{2+} permeability of the NMDAR, with

minimal influence on the whole-cell current amplitude and deactivation kinetics.

We next investigated the dynamic properties of NMDAR currents. Desensitization

describes the degree to which a receptor loses responsiveness to the agonist in

response to continued exposure, which may reflect a mechanism to dynamically reduce

NMDAR currents when the network is in a high activity state. For this, we simulated

NMDAR receptor activation that likely occurs during large phasic motor activation that

disrupts normal rhythmic output in severe hypoxia (Fig. 4A) (Adams et al., 2021). Thus,

we applied NMDAR agonists every 2 seconds for a total of 10 pulses per neuron to

assess desensitization in a physiologically-relevant way (Fig 4B). We observed a

moderate degree of desensitization in controls, whereby the 10th pulse in the series

produced a current that was ~60% of the baseline amplitude. NMDARs from hibernators

also desensitized, but the final puff elicited a current that was significantly smaller than

controls, at ~20% of the initial value (Fig. 4C, $t_{(24)}$ = 2.946, p=0.0071, unpaired t-test). In

addition, a two-way ANOVA for relative current amplitude across all puffs shows a main

effect of group, indicating that currents from hibernators were more desensitized across

the entire experimental protocol (Fig. 4D, $F_{(1, 24)}$ = 7.084, p=0.0137, two-way ANOVA).

Therefore, NMDAR currents not only become less permeable to Ca^{2+} after hibernation,

but they also pass less current during repetitive activation that otherwise causes

network output to lose homeostasis.

NMDARs are heterotetramers composed of an obligatory GluN1 subunit and a

combination of GluN2A-D and/or GluN3A-B protein subunits (Paoletti et al., 2013). To

gain insight into the NMDAR subunits that correspond these physiological changes, we

used inhibitors for the subunits GluN2A, GluN2B, GluN2C/GluN2D, and GluN3 to

uncover the physiological participation of NMDA subunits in neurons from controls and

hibernators. In controls, we observed a significant inhibition of the total NMDAR current

when the GluN2B and GluN2C/GluN2D subunits were inhibited compared to time

control, shown by a current decrease of 44% for GluN2B (t_{17}=4.915, p=0.0001 t-test)

and 29% for GluN2C/D (t_{15}=2.916, p=0.0106 t-test). It is important to appreciate that

time controls underwent a slight increase in current amplitude during the experimental

time course (Fig 5A); thus, the decreases induced by each drug represent an

underestimation of their true contribution to the total current. After hibernation GluN2B

maintained its participation in the NMDA current, showing a decrease relative to time

control when inhibited (t_{22}=3.103, p=0.0052 t-test), which did not differ from control

neurons (t_{20}=1.427, p=0.1689 t-test; Fig. 5C). However, GluN2C/GluN2D no longer

contributed to the NMDA current after hibernation (t_{19}=0.2833, p=0.7800 t-test

compared to time control), showing a difference from the control group (t_{15}=2.612,

p=0.0196 t-test, Fig. 5D). The subunits GluN2A (Fig. 5B) and GluN3 (Fig. 5E) did not

seem to have a consistent functional contribution to the NMDA current in control or

overwintered frogs. Therefore, the GluN2B subunit appears to be the main GluN2

subunit participating in the NMDA current, while GluN2C/D plays a smaller role and is

lost after hibernation.

To understand the potential for transcriptional control of NMDAR subunits, we

followed up physiology experiments with single-cell quantitative PCR (qPCR) to

measure the mRNA expression of all 7 NMDAR subunits that encode the NMDAR (the

Grin gene family). These data are summarized in Fig. 6. Consistent with the dominant

contribution of GluN2B to the total NMDAR current, *Grin2B* appeared to be the most

abundant *Grin* transcript for both controls and hibernators, along with *Grin1* that codes

for the obligatory GluN1 subunit. Pharmacologically, QZN 46 is selective for both

GluN2C and GluN2D. However, most neurons lacked mRNA expression of *Grin2D* (Fig

6), indicating the GluN2C is the main functional subunit blocked by QZN 46 in control

neurons. Consistent with our functional results, after hibernation we observed a

significant decrease in the mean abundance *Grin2C* (p=0.0003, Mann-Whitney U test).

Surprisingly, although we did not observe a contribution from the GluN3 subunit to the

NMDAR in either group, we observed mRNA expression of both *Grin3A* and *Grin3B*,

and each of these subunits decreased after hibernation (*Grin3A*: t_{34}=2.920, p=0.0062,

unpaired t test; *Grin3B*: $t_{21.62}$=2.597, p=0.0166, Welch's t-test). These data suggest that

a transcriptional program influences the NMDAR subunit composition in concert with

modification of whole-cell receptor current properties.

What are the functional impacts of NMDAR modifications on the ability of the

network to function during hypoxia? In control brainstems, the first several minutes of

hypoxia leads to large, uncoordinated output that interferes with the normal patterned

activity of the respiratory network (Fig. 7A$_1$). This type of chaotic activity is thought to

arise, at least in part, due to motor hyperexcitability induced by energy stress, as it does

not occur when the brainstem is well-oxygenated (Adams et al., 2021; Bueschke et al.,

2021). To determine the extent to which NMDARs play a role in this response, we

exposed a group of brainstem preparations to D-AP5 and then measured the degree of

large amplitude output that disrupted normal activity shortly after the onset of hypoxia.

In the presence of D-AP5 (Fig. 7A$_2$), the mean amplitude of this chaotic "non-

respiratory" activity was strongly blunted relative to controls. In addition, the small

degree of disruption was often not sufficient to disrupt patterned output during hypoxia

as shown in Fig. 7A$_2$. These results show that the normal profile of NMDARs (with a

permeability to Ca^{2+} and relatively less desensitization) contribute to chaotic output that

disrupts normal function of the network during hypoxia.

As disruptive motor activity during hypoxia arose largely due to NMDARs, this

type of activity provided a sensitive indicator of how the modified NMDAR profile

influenced network excitability in hypoxia. Strikingly, the chaotic, large amplitude, non-

respiratory bursting in hypoxia that was caused largely by NMDARs did not occur in any

hibernation preparation (Fig. 7B$_1$). We also applied D-AP5 to a group of hibernation

preparations. The addition of D-AP5 did not influence network function during hypoxia,

as there was no disruptive activity to suppress (Fig. 7B$_2$). These results are summarized

in Fig. 7C by a one-way ANOVA for control, control+D-AP5, hibernation, and

hibernation+D-AP5 ($F_{(3,31)}$=15.25, p<0.0001, one-way ANOVA). Post hoc tests reveal

that blocking NMDARs in control preparations suppresses the large non-respiratory

bursting in hypoxia (p=0.0002, Holm-Sidak multiple comparisons test). This is not

different than the degree of large amplitude bursting during hypoxia from hibernation

preparations without (p=0.6824, Holm-Sidak multiple comparisons test) or with D-AP5

(p=0.6824, Holm-Sidak multiple comparisons test). These results show that control

preparations without functional NMDARs (because they have been blocked by D-AP5)

behave similar to circuits with intact NMDARs after they have become less permeable

to Ca^{2+} and desensitize more strongly. Thus, modifications in NMDAR function act in a

way to prevent hyperexcitable network states during energy stress while maintaining

their role in network output under well-oxygenated conditions.

Discussion:

During energy stress, NMDAR activation disrupts circuit output in a wide range of species, including amphibians (Fig 7). To emerge from hibernation in an ice-covered pond, frogs must restart critical behaviors, including breathing, on the background of extraordinarily low oxygen levels (Tattersall & Ultsch, 2008; Ultsch et al., 2004). The challenge lies in the fact that most frogs are not generally considered to be a strongly "hypoxia-tolerant" species. Metabolically, a suite of plasticity mechanisms arise during hibernation and improve the fuel supply needed to operate brainstem synapses under anaerobic conditions (Bueschke et al., 2021a; Hu & Santin, 2022, Amaral-Silva & Santin, 2023). Here, we tested the hypothesis that these animals also use physiological mechanisms to avoid hyperexcitable network states induced by energy stress. We identified two key modifications to NMDARs that offset hyperexcitable output in hypoxia: reduced Ca^{2+} permeability (Fig. 3) and enhanced desensitization (Fig. 4).

These NMDAR modifications are consistent with plasticity that acts to maintain network excitability and reduce energy consumption in hypoxia. First, enhancing desensitization reduces depolarizing drive during repetitive stimulation that normally occurs during hypoxia (Fig. 4). As NMDARs contribute to chaotic bursting that disrupts patterned network output (Fig. 4A, Fig. 7A), stronger desensitization during high activity states likely acts as a potent brake on excitability. Second, switching to NMDARs with lower Ca^{2+} permeability likely incurs a cost savings. Here, roughly 50% of the population motor output that drives breathing is generated through NMDARs (Fig 1), which means for every breath the animal takes, increases in intracellular Ca^{2+} from NMDARs must be cleared to maintain ion homeostasis. Although the regulation of intracellular Ca^{2+} is

multifaceted, the plasma membrane Ca^{2+} ATPase plays a large role in maintaining Ca^{2+}

homeostasis in active neurons (Malci et al., 2022; Schmidt et al., 2017). This pump

consumes more than 3 times the ATP as the Na^+/K^+ ATPase, as it extrudes 1 Ca^{2+} ion

per ATP hydrolyzed and transports Ca^{2+} against a larger electrochemical gradient

compared to Na^+ and K^+. Thus, in addition to minimizing well-established pathways for

Ca^{2+}-induced excitotoxicity (Szydlowska and Tymianski, 2010), lowering the Ca^{2+}

permeability of NMDAR may also reduce the metabolic burden of Ca^{2+} regulation in

hypoxia, allowing neurons to allocate energy to other processes required to maintain

homeostasis.

Interestingly, these modifications do not alter the normal role of the NMDARs in

generating motor output under well-oxygenated conditions (Fig. 1). One reason for this

may be that NMDAR currents in this study were measured at the cell body. Thus, the

plasticity in NMDAR function we observe may be localized to extra-synaptic regions,

which are well known to elicit the pathological actions of NMDARs during hypoxia and

ischemia (Tu et al., 2010; Zhou et al., 2013). It is not yet known if hibernation influences

NMDARs at respiratory-related synapses. However, if hibernation alters the properties

of synaptic NMDARs, reduced Ca^{2+} permeability and enhanced desensitization likely

have little influence on network output since these modifications do not alter the whole-

cell current amplitude and kinetics over the timescale of individual networks population

bursts (<1 s burst every ~10 s) (Fig. 1A). Altogether, NMDAR plasticity appears to

contribute to a state of resilience during metabolic stress without influencing normal

functioning of the network under healthy conditions.

513   What cellular mechanisms influence the functional changes we observe?

514 NMDAR receptors are heterotetrameric complexes composed of an obligatory GluN1

515 subunit and GluN2A-D and GluN3A-B subunits, which determine receptor properties

516 (Paoletti et al., 2013). Heterologous expression of GluN3 with GluN1 renders the

517 receptor impermeable to Ca^{2+} and highly responsive to glycine (Chatterton et al., 2002).

518 Although our single-cell qPCR data show altered mRNA expression of genes that code

519 for the GluN3 subunit, our functional studies show that it does not contribute to the

520 NMDAR current, indicating that changes in GluN3 are not responsible for decreases in

521 Ca^{2+} permeability after hibernation. Interestingly, in NMDAR complexes containing

522 GluN2, Ca^{2+} permeability is not an intrinsic property of the protein, but rather, is caused

523 by phosphorylation of the GluN2B subunit (Skeberdis et al., 2006; Murphy et al., 2014).

524 As GluN2B had the greatest functional role and mRNA expression (Fig. 5 and Fig. 6),

525 we speculate that alterations in the phosphorylation state of GluN2B may regulate Ca^{2+}

526 permeability. For desensitization, many factors influence this property, including subunit

527 composition (Vicini et al., 1998), protein binding partners (Sornarajah et al., 2008), and

528 intracellular signaling pathways (Alagarsamy et al., 1999). Most relevant to our results,

529 GluN1/GluN2C heterodimers undergo relatively weak desensitization (Dravid et al.,

530 2008; Alsaloum et al., 2016). We observed a decrease in *Grin2C* mRNA expression and

531 a reduced contribution of GluN2C to the whole-cell current after hibernation. These

532 results suggest that transcriptional control of GluN2C expression may shift neurons to

533 favor subunit combinations with greater degrees of desensitization, such as

534 GluN1/GluN2B (Vicini et al., 1998). As a caveat, we acknowledge that attributes of

535 NMDAR currents may represent the sum of many potential combinations of GluN

subunits; although we did not find significant functions of GluN3 and GluN2A, some

individual cells within the population did appear to respond to their inhibitors, suggesting

that variable combinations of NMDAR subunits compose the total whole cell current. In

addition, receptor expression may vary depending on the localization within the cell

(e.g., synaptic vs. extrasynaptic). This complicates the connection between single cell

mRNA abundances and NMDAR function measured at the cell body. Nevertheless,

combining functional and molecular approaches point to mechanisms that alter the

functional properties of the NMDARs to improve function during energy stress.

The frog brainstem has a remarkable ability to improve its function during

hypoxia and ischemia after hibernation (Bueschke et al., 2021). This response involves

enhancing anaerobic glucose metabolism at active synapses (Amaral-Silva & Santin,

2023; Bueschke et al., 2021a). Dominant hypotheses for neural circuit evolution posit

that natural selection acts on the efficiency of synapses, optimizing the ratio of

information transfer to ATP consumption (Harris et al., 2015; Quintela-López et al.,

2022). However, synapses here can switch into a state that supports network activity for

hours with ~1/15th the amount of ATP through adjustments in synaptic metabolism that

maximize glycolytic ATP production (Amaral-Silva & Santin, 2023). Our results suggest

that physiological modifications help to lower the cost of network activity to match the

dramatically reduced rate of energy production while maintaining seemingly normal

output. Given that this circuit makes physiological and energetic modifications that allow

it to run on very little energy, these results demonstrate some neural circuits may reside

far from their "optimal" efficiency. This begs the question of why animals would

suppress such a state until it is critical for survival (*e.g.,* emerging from an ice-covered

pond after months of hibernation).

Although our results show strong support for NMDAR plasticity as a potential

energy-saving mechanism, they also hint at a potential cost of maintaining an "ultra-

high" efficiency network state. NMDARs play a key role in plasticity, whereby Ca^{2+} influx

through the receptor potentiates synaptic strength in a wide range of systems, including

the respiratory motoneurons in this species (Bueschke et al., 2021b). NMDAR

modifications we observe likely incur an energy savings but have features that appear

to be incompatible with NMDAR-dependent plasticity: strongly reduced Ca^{2+}

permeability and greatly enhanced desensitization which quickly reduces channel

activity upon repetitive stimulation. Indeed, Ca^{2+} impermeable NMDARs cause memory

impairments and reduce long-term potentiation in mice (Conde-Dusman et al., 2021;

Hurley et al., 2022). Therefore, we suggest a trade-off exists between metabolic

robustness and the capacity for plasticity within circuits. It seems sensible for an animal

to favor an extreme degree of energetic resilience to restart critical behaviors in

severely hypoxic conditions. However, when animals are not faced with energy stress,

as is the case for most vertebrates most of the time, it is clearly beneficial for the brain

to adapt to changes in the environment through plasticity. In support of the relationship

between plasticity and metabolic resilience, the rodent hippocampus shows a dorsal-

ventral gradient in damage by ischemia (Ashton et al., 1989), which mirrors the capacity

for synaptic plasticity owing to the degree of Ca^{2+}-permeability of the NMDARs (Hurley

et al., 2022). Therefore, hypotheses for selective pressures shaping the evolution of

neural circuits may need to incorporate how synapses have balanced the need for

plasticity and robust function during energy stress.

In conclusion, a failure to balance energy supply and physiological demands of

neurons leads to disordered circuit activity. We identified a circuit that has the capacity

to modify NMDARs to reduce Ca^{2+} influx and constrain excitability in hypoxia without

altering their normal contribution to network function. These findings represent a state-

dependent form of NMDAR plasticity that likely plays an adaptive role by promoting

coordinated network activity in a circuit that cannot typically maintain homeostasis in low

oxygen. These results provide insight into natural mechanisms that reduce neural circuit

reliance on high rates of energy production. Yet, they also point to a potential trade-off

between energetic robustness and plasticity that requires Ca^{2+} influx through NMDARs.

Uncovering mechanisms that shift the balance between these two states may inform

novel neuroprotection strategies with high clinical relevance.

Author Contributions: Conception and Design; JMS. Performed Experiments; NB,

MH, LAS, AA. Analyzed Data; NB, JMS. Wrote Manuscript; JMS, NB. Edited

manuscript; JS, NB, MH, LAS, AA. Approved Final version of the manuscript; JS, NB,

MH, LAS, AA.

References

Abrahamsson T, Chou CYC, Li SY, Mancino A, Costa RP, Brock JA, Nuro E, Buchanan KA, Elgar D, Blackman AV (2017). Differential regulation of evoked and spontaneous release by presynaptic NMDA receptors. *Neuron* 96:839-855. e835. https://doi.org/10.1016/j.neuron.2017.09.030

Adams, S., Zubov, T., Bueschke, N., & Santin, J. M. (2021). Neuromodulation or energy failure? Metabolic limitations silence network output in the hypoxic amphibian brainstem. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, *320*(2), R105–R116. https://doi.org/10.1152/ajpregu.00209.2020

Alagarsamy, S., Marino, M. J., Rouse, S. T., Gereau, R. W., Heinemann, S. F., & Conn, P. J. (1999). Activation of NMDA receptors reverses desensitization of mGluR5 in native and recombinant systems. *Nature Neuroscience*, *2*(3), 234–240. https://doi.org/10.1038/6338

Alsaloum M, Kazi R, Gan Q, Amin J, Wollmuth LP (2016). A molecular determinant of subtype-specific desensitization in ionotropic glutamate receptors. *Journal of Neuroscience* 36, 2617-2622. https://doi.org/10.1523/JNEUROSCI.2667-15.2016

Amaral-Silva, L., Santin, J.M., (2023). Synaptic modifications transform neural networks to function without oxygen. *BMC Biology,* 21(54).

Ashton, D., Van Reempts, J., Haseldonckx, M., & Willems, R. (1989). Dorsal-ventral gradient in vulnerability of CA1 hippocampus to ischemia: A combined histological and electrophysiological study. *Brain Research*, *487*(2), 368–372. https://doi.org/10.1016/0006-8993(89)90842-1

Beesley, S., Sullenberger, T., & Kumar, S. S. (2020). The GluN3 subunit regulates ion selectivity within native N-methyl-d-aspartate receptors. *IBRO Reports*, *9*, 147–156. https://doi.org/10.1016/j.ibror.2020.07.009

Bettini E, Sava A, Griffante C, Carignani C, Buson A, Capelli AM, Negri M, Andreetta F, Senar-Sancho SA, Guiral L (2010). Identification and characterization of novel NMDA receptor antagonists selective for NR2A-over NR2B-containing receptors. *Journal of Pharmacology and Experimental Therapeutics* 335, 636-644. https://doi.org/10.1124/jpet.110.172544

Bickler, P. E., & Buck, L. T. (2007). Hypoxia Tolerance in Reptiles, Amphibians, and Fishes: Life with Variable Oxygen Availability. *Annual Review of Physiology*, *69*(1), 145–170. https://doi.org/10.1146/annurev.physiol.69.031905.162529

Bickler, P. E., Donohoe, P. H., & Buck, L. T. (2000). Hypoxia-Induced Silencing of
NMDA Receptors in Turtle Neurons. *The Journal of Neuroscience*, *20*(10), 3522–
3528. https://doi.org/10.1523/JNEUROSCI.20-10-03522.2000
Bordone, M. P., Salman, M. M., Titus, H. E., Amini, E., Andersen, J. V., Chakraborti, B.,
Diuba, A. V., Dubouskaya, T. G., Ehrke, E., Espindola de Freitas, A., Braga de
Freitas, G., Gonçalves, R. A., Gupta, D., Gupta, R., Ha, S. R., Hemming, I. A.,
Jaggar, M., Jakobsen, E., Kumari, P., … Seidenbecher, C. I. (2019). The
energetic brain – A review from students to students. *Journal of Neurochemistry*,
151(2), 139–165. https://doi.org/10.1111/jnc.14829
Buck, L. T., & Pamenter, M. E. (2018). The hypoxia-tolerant vertebrate brain: Arresting
synaptic activity. *Comparative Biochemistry and Physiology Part B: Biochemistry
and Molecular Biology*, *224*, 61–70. https://doi.org/10.1016/j.cbpb.2017.11.015
Bueschke, N., Amaral-Silva, L., Adams, S., & Santin, J. M. (2021a). Transforming a
neural circuit to function without oxygen and glucose delivery. *Current Biology*,
31(24), R1564–R1565. https://doi.org/10.1016/j.cub.2021.11.003
Bueschke, N., Amaral-Silva, L., Hu, M., & Santin, J. M. (2021b). Lactate ions induce
synaptic plasticity to enhance output from the central respiratory network. *The
Journal of Physiology*, *599*(24), 5485–5504. https://doi.org/10.1113/JP282062
Chatterton, J. E., Awobuluyi, M., Premkumar, L. S., Takahashi, H., Talantova, M., Shin,
Y., Cui, J., Tu, S., Sevarino, K. A., Nakanishi, N., Tong, G., Lipton, S. A., &
Zhang, D. (2002). Excitatory glycine receptors containing the NR3 family of
NMDA receptor subunits. *Nature*, *415*(6873), 793–798.
https://doi.org/10.1038/nature715
660

661 Christian DT, Stefanik MT, Bean LA, Loweth JA, Wunsch AM, Funke JR, Briggs CA,
662   Lyons J, Neal D, Milovanovic M (2021). GluN3-containing NMDA receptors in the
663   rat nucleus accumbens core contribute to incubation of cocaine craving. Journal
664   of Neuroscience 41, 8262-8277. https://doi.org/10.1523/JNEUROSCI.0406-
665   21.2021
666

Conde-Dusman, M. J., Dey, P. N., Elía-Zudaire, Ó., Rabaneda, L. G., García-Lira, C.,
Grand, T., Briz, V., Velasco, E. R., Andero, R., Niñerola, S., Barco, A., Paoletti,
P., Wesseling, J. F., Gardoni, F., Tavalin, S. J., & Perez-Otaño, I. (2021). Control
of protein synthesis and memory by GluN3A-NMDA receptors through inhibition
of GIT1/mTORC1 assembly. *ELife*, *10*, e71575.
https://doi.org/10.7554/eLife.71575
Czech-Damal, N. U., Geiseler, S. J., Hoff, M. L. M., Schliep, R., Ramirez, J.-M., Folkow,
L. P., & Burmester, T. (2014). The role of glycogen, glucose and lactate in
neuronal activity during hypoxia in the hooded seal (Cystophora cristata) brain.
Neuroscience, *275*, 374–383. https://doi.org/10.1016/j.neuroscience.2014.06.024
Dravid, S. M., Prakash, A., & Traynelis, S. F. (2008). Activation of recombinant
NR1/NR2C NMDA receptors: NR1/NR2C receptor activation. *The Journal of
Physiology*, *586*(18), 4425–4439. https://doi.org/10.1113/jphysiol.2008.158634
680

Edman S, McKay S, Macdonald L, Samadi M, Livesey M, Hardingham G, Wyllie D (2012). TCN 201 selectively blocks GluN2A-containing NMDARs in a GluN1 co-agonist dependent but non-competitive manner. *Neuropharmacology* 63, 441-449. https://doi.org/10.1016/j.neuropharm.2012.04.027

France G, Fernández-Fernández D, Burnell ES, Irvine MW, Monaghan DT, Jane DE, Bortolotto ZA, Collingridge GL, Volianskis A (2017). Multiple roles of GluN2B-containing NMDA receptors in synaptic plasticity in juvenile hippocampus. *Neuropharmacology* 1127 6-83. https://doi.org/10.1016/j.neuropharm.2016.08.010

Garcia, V. B., Abbinanti, M. D., Harris-Warrick, R. M., & Schulz, D. J. (2018). Effects of Chronic Spinal Cord Injury on Relationships among Ion Channel and Receptor mRNAs in Mouse Lumbar Spinal Cord. *Neuroscience, 393*, 42–60. https://doi.org/10.1016/j.neuroscience.2018.09.034

Hammond, S. A., Warren, R. L., Vandervalk, B. P., Kucuk, E., Khan, H., Gibb, E. A., Pandoh, P., Kirk, H., Zhao, Y., Jones, M., Mungall, A. J., Coope, R., Pleasance, S., Moore, R. A., Holt, R. A., Round, J. M., Ohora, S., Walle, B. V., Veldhoen, N., … Birol, I. (2017). The North American bullfrog draft genome provides insight into hormonal regulation of long noncoding RNA. *Nature Communications, 8*(1), 1433. https://doi.org/10.1038/s41467-017-01316-7

Hansen KB, Traynelis SF (2011). Structural and mechanistic determinants of a novel site for noncompetitive inhibition of GluN2D-containing NMDA receptors. *Journal of Neuroscience* 31, 3650-3661. https://doi.org/10.1523/JNEUROSCI.5565-10.2011

Harris, J. J., Jolivet, R., Engl, E., & Attwell, D. (2015). Energy-Efficient Information Transfer by Visual Pathway Synapses. *Current Biology, 25*(24), 3151–3160. https://doi.org/10.1016/j.cub.2015.10.063

Hille, B. (2001). *Ion channels of excitable membranes* (3rd ed). Sinauer.

Howell, B., Baumgardner, F., Bondi, K., & Rahn, H. (1970). Acid-base balance in cold-blooded vertebrates as a function of body temperature. *American Journal of Physiology-Legacy Content, 218*(2), 600–606. https://doi.org/10.1152/ajplegacy.1970.218.2.600

Hu, M., & Santin, J. M. (2022). Transformation to ischaemia tolerance of frog brain function corresponds to dynamic changes in mRNA co-expression across metabolic pathways. *Proceedings of the Royal Society B: Biological Sciences, 289*(1979), 20221131. https://doi.org/10.1098/rspb.2022.1131

Hurley, E. P., Mukherjee, B., Fang, L., Barnes, J. R., Nafar, F., Hirasawa, M., & Parsons, M. P. (2022). *Atypical NMDA Receptors Limit Synaptic Plasticity in the Adult Ventral Hippocampus* [Preprint]. Neuroscience. https://doi.org/10.1101/2022.10.05.510966

Jatzke, C., Junryo Watanabe, & Wollmuth, L. P. (2002). Voltage and concentration dependence of Ca^{2+} permeability in recombinant glutamate receptor subtypes.

The Journal of Physiology, 538(1), 25–39.
https://doi.org/10.1113/jphysiol.2001.012897

Kottick, A., Baghdadwala, M. I., Ferguson, E. V., & Wilson, R. J. A. (2013). Transmission of the respiratory rhythm to trigeminal and hypoglossal motor neurons in the American Bullfrog (Lithobates catesbeiana). *Respiratory Physiology & Neurobiology, 188*(2), 180–191.
https://doi.org/10.1016/j.resp.2013.06.008

Kvist T, Greenwood JR, Hansen KB, Traynelis SF, Bräuner-Osborne H (2013). Structure-based discovery of antagonists for GluN3-containing N-methyl-D-aspartate receptors. *Neuropharmacology* 75:324-336.
https://doi.org/10.1016/j.neuropharm.2013.08.003

Larson, J., Drew, K. L., Folkow, L. P., Milton, S. L., & Park, T. J. (2014). No oxygen? No problem! Intrinsic brain tolerance to hypoxia in vertebrates. *Journal of Experimental Biology, 217*(7), 1024–1039. https://doi.org/10.1242/jeb.085381

Larson, J., & Park, T. J. (2009). Extreme hypoxia tolerance of naked mole-rat brain. *NeuroReport, 20*(18), 1634–1637.
https://doi.org/10.1097/WNR.0b013e32833370cf

Malci, A., Lin, X., Sandoval, R., Gundelfinger, E. D., Naumann, M., Seidenbecher, C. I., & Herrera-Molina, R. (2022). Ca2+ signaling in postsynaptic neurons: Neuroplastin-65 regulates the interplay between plasma membrane Ca2+ ATPases and ionotropic glutamate receptors. *Cell Calcium, 106*, 102623.
https://doi.org/10.1016/j.ceca.2022.102623

Murphy, J. A., Stein, I. S., Lau, C. G., Peixoto, R. T., Aman, T. K., Kaneko, N., Aromolaran, K., Saulnier, J. L., Popescu, G. K., Sabatini, B. L., Hell, J. W., & Zukin, R. S. (2014). Phosphorylation of Ser1166 on GluN2B by PKA Is Critical to Synaptic NMDA Receptor Function and Ca $^{2+}$ Signaling in Spines. *The Journal of Neuroscience, 34*(3), 869–879. https://doi.org/10.1523/JNEUROSCI.4538-13.2014

Paoletti, P., Bellone, C., & Zhou, Q. (2013). NMDA receptor subunit diversity: Impact on receptor properties, synaptic plasticity and disease. *Nature Reviews Neuroscience, 14*(6), 383–400. https://doi.org/10.1038/nrn3504

Pellizzari, S., Hu, M., Amaral-Silva, L., Saunders, S., Santin, J.M. (2023). Neuron populations use variable combinations of short-term feedback mechanisms to stabilize firing rate. *PLOS Biology*, accepted for publication.

Quintela-López, T., Shiina, H., & Attwell, D. (2022). Neuronal energy use and brain evolution. *Current Biology, 32*(12), R650–R655.
https://doi.org/10.1016/j.cub.2022.02.005

Santin, J. M., & Schulz, D. J. (2019). Membrane Voltage Is a Direct Feedback Signal That Influences Correlated Ion Channel Expression in Neurons. *Current Biology, 29*(10), 1683-1688.e2. https://doi.org/10.1016/j.cub.2019.04.008

Schmidt, N., Kollewe, A., Constantin, C. E., Henrich, S., Ritzau-Jost, A., Bildl, W., Saalbach, A., Hallermann, S., Kulik, A., Fakler, B., & Schulte, U. (2017).

Neuroplastin and Basigin Are Essential Auxiliary Subunits of Plasma Membrane Ca2+-ATPases and Key Regulators of Ca2+ Clearance. *Neuron*, *96*(4), 827–838.e9. https://doi.org/10.1016/j.neuron.2017.09.038

Skeberdis, V. A., Chevaleyre, V., Lau, C. G., Goldberg, J. H., Pettit, D. L., Suadicani, S. O., Lin, Y., Bennett, M. V. L., Yuste, R., Castillo, P. E., & Zukin, R. S. (2006). Protein kinase A regulates calcium permeability of NMDA receptors. *Nature Neuroscience*, *9*(4), 501–510. https://doi.org/10.1038/nn1664

Sornarajah, L., Vasuta, O. C., Zhang, L., Sutton, C., Li, B., El-Husseini, A., & Raymond, L. A. (2008). NMDA Receptor Desensitization Regulated by Direct Binding to PDZ1-2 Domains of PSD-95. *Journal of Neurophysiology*, *99*(6), 3052–3062. https://doi.org/10.1152/jn.90301.2008

Szydlowska, K., & Tymianski, M. (2010). Calcium, ischemia and excitotoxicity. *Cell Calcium*, *47*(2), 122–129. https://doi.org/10.1016/j.ceca.2010.01.003

Tattersall, G. J., & Boutilier, R. G. (1997). Balancing hypoxia and hypothermia in cold-submerged frogs. *The Journal of Experimental Biology*, *200*(Pt 6), 1031–1038. https://doi.org/10.1242/jeb.200.6.1031

Tattersall, G. J., & Ultsch, G. R. (2008). Physiological Ecology of Aquatic Overwintering in Ranid Frogs. *Biological Reviews*, *83*(2), 119–140. https://doi.org/10.1111/j.1469-185X.2008.00035.x

Tu, W., Xu, X., Peng, L., Zhong, X., Zhang, W., Soundarapandian, M. M., Belal, C., Wang, M., Jia, N., Zhang, W., Lew, F., Chan, S. L., Chen, Y., & Lu, Y. (2010). DAPK1 Interaction with NMDA Receptor NR2B Subunits Mediates Brain Damage in Stroke. *Cell*, *140*(2), 222–234. https://doi.org/10.1016/j.cell.2009.12.055

Ultsch, G. R., Reese, S. A., & Stewart, E. R. (2004). Physiology of hibernation inRana pipiens: Metabolic rate, critical oxygen tension, and the effects of hypoxia on several plasma variables. *Journal of Experimental Zoology*, *301A*(2), 169–176. https://doi.org/10.1002/jez.a.20014

Vicini, S., Wang, J. F., Li, J. H., Zhu, W. J., Wang, Y. H., Luo, J. H., Wolfe, B. B., & Grayson, D. R. (1998). Functional and Pharmacological Differences Between Recombinant *N* -Methyl- D -Aspartate Receptors. *Journal of Neurophysiology*, *79*(2), 555–566. https://doi.org/10.1152/jn.1998.79.2.555

White, R. J., & Reynolds, I. J. (1995). Mitochondria and Na+/Ca2+ exchange buffer glutamate-induced calcium loads in cultured cortical neurons. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *15*(2), 1318–1328.

Wilkie, M. P., Pamenter, M. E., Alkabie, S., Carapic, D., Shin, D. S. H., & Buck, L. T. (2008). Evidence of anoxia-induced channel arrest in the brain of the goldfish (Carassius auratus). *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, *148*(4), 355–362. https://doi.org/10.1016/j.cbpc.2008.06.004

Wu, Q. J., & Tymianski, M. (2018). Targeting NMDA receptors in stroke: New hope in neuroprotection. *Molecular Brain*, *11*(1), 15. https://doi.org/10.1186/s13041-018-0357-8

Zhou, X., Hollern, D., Liao, J., Andrechek, E., & Wang, H. (2013). NMDA receptor-mediated excitotoxicity depends on the coactivation of synaptic and extrasynaptic

815   receptors. *Cell Death & Disease, 4*(3), e560–e560.
816   https://doi.org/10.1038/cddis.2013.82

818 **Tables**

Table 1. Primer Sequences for SYBR Green qPCR Assays			
Target	Forward	Reverse	Efficiency
Grin2b	GAGTGGGAGGACAGAACTGC	TGGCAGACTTCACAGCGGAT	81%
Grin2c	GGTGCTGCAAGTGGAAACTG	GGATGTAGCGGGCACTCTTT	100%
Grin2d	TGTTCGCATGGGAGCATCTT	GCCTTTGGATCCTCAGCACT	95%
Grin3b	CCAGCGATCTTCCTTGGTGT	TGTCCAACCAGCCAGTGAAA	84%
18S rRNA	CAGGCCGGTCGCCTGAATAC	GGCCCCAGTTCCGAAAACCA	101%

Table 2. Primer Sequences for Probe-based qPCR Assays			
Target	**Forward Primer**	**Reverse Primer**	**Efficiency**
Grin1	TGGTGGCGGTAATGTTGTATC	TCAGTGCATCCTCCTCTTCTTC	91%
Grin2a	TGACCCATTGCCGAAGTTG	GCATTCGGTCTTCATCAATGTCA	91%
Grin3a	GGCCAGACCAAGCACAAAG	CACAAAGGGTGGCTCAATCAAG	96%
Prober Sequences		**Probe-Quencher**	
Grin1	CAGGTTCAGTCCTTTTGGCCGGTT	FAM-BHQ1	
Grin2a	ACCCAGTTATACTGGTCACTACTCAACCA	ROX-BHQ2	
Grin3a	CAGAAGTCATTTCCAACAACCACCTCGA	VIC-BHQ1	

     Figures and Legends

Figure 1. Hibernation does not alter the NMDAR tone of respiratory motor output. (A)
Schematic of the in vitro brainstem preparation. Raw motor output associated with breathing
can be recorded from the cut cranial nerve X rootlet (CNX). Raw signals are then rectified and
integrated for analysis. (B) Representative respiratory motor bursts in baseline conditions and
after block of NMDA-glutamate receptors with 50 μM D-AP5. Control animals are shown on the
top and hibernation is shown on the bottom. (C) Box and whisker plot comparing burst
amplitude in D-AP5 relative to control between controls (left, green, n = 8) and hibernation
(right, blue, n = 7) brainstems, showing no significant change in D-AP5 sensitivity (unpaired t-
test). (D) Box and whisker plots comparing burst frequency changes in response to D-AP5.
There was no significant difference between the change in burst frequency induced by D-AP5
across groups. Dots represent each data point of individual experiments.

Figure 2. Hibernation does not alter the whole-cell NMDAR current amplitude, decay kinetics, and hypoxia sensitivity in identified respiratory motoneurons. (A) Schematic illustrating the backfill labeling process to identify motoneurons and the setup to focally apply NMDAR agonists (1 mM NMDA and 10 µM glycine) to the cell body. (B) Traces showing typical responses to focal application of NMDA and glycine. (C-D) NMDAR current amplitude and decay time constant is unaltered between control (n = 11 neurons from N = 5 animals) and hibernation (n = 13 neurons from N = 4 animals) neurons following unpaired t tests. Boxes show interquartile range and whiskers represent min and max values with dots showing individual data points. (E-F) Individual NMDAR amplitude and decay values before and after ~5 minutes of hypoxia (0% O$_2$). Paired t test results suggest hypoxia does not change either NMDAR variable in control (n = 5 neurons from N = 4 animals) or hibernation (n=4 neurons from N = 4 animals) neurons. Line represents the "before" and "after" response for individual neurons in response to hypoxia.

Figure 3. Hibernation decreases Ca²⁺ permeability of NMDARs. (A-B) Raw whole-cell
voltage clamp traces held at a range of voltages (-20 to +40 mV, Δ5 mV) responding to NMDAR
activation *via* focal application of NMDAR agonists. Control (A) is shown on the top in green,
and hibernation (B) is shown in blue. Reversal potentials (E_{rev}) in these example traces are
highlighted to visualize the typical reversal potentials in 1 mM and 10 mM Ca²⁺ from control and
hibernation motoneurons. (C) Results of NMDAR E_{rev} (n = 16 neurons, all groups) show
significance in 2-way ANOVA interaction (p = 0.0072), indicating that hibernation affects the
response to Ca²⁺. Pairwise comparisons reveal a significant depolarization of E_{rev} by 10 mM
Ca²⁺ in control neurons but not hibernation neurons. We used 4 animals for hibernation in low
Ca²⁺ and 5 animals for all other groups. Dots represent each data point of individual cells in
each group, with boxes representing interquartile range and whiskers displaying min and max
E_{rev} values. E_{rev} values shown in this plot are corrected for calculated series resistance errors
based on the holding current. ****signifies p<0.0001

Figure 4. Hibernation enhances NMDAR desensitization in response to repetitive activation. (A) Example recording of population motor output from a control brainstem preparation ~5 minutes into hypoxia. This illustrates the transition from normal patterned output to chaotic, higher frequency output with large amplitude "non-respiratory" bursting activity that occurs during hypoxia. (B) Representative traces of vagal motoneuron response to pulses of NMDAR/glycine onto the soma at 0.5 Hz. Controls under a moderate degree of desensitization (green, top), while hibernation neurons (bottom, blue) tended to have smaller current amplitudes by the end of the experiment. (C) Final (10th) pulse current amplitude relative to initial pulse. Hibernation motoneurons (n = 14 neuron from N = 4 animals) had a significantly lower final amplitude than controls (n = 12 neurons, N = 5 animals), suggesting that NMDARs after hibernation are more susceptible to desensitization. (D) Plot showing all data from the experimental series. Two-way ANOVA reveals a main effect of group and puff number. A significant group effect indicates that NMDAR currents throughout the experimental series were overall smaller after hibernation, consistent with enhanced desensitization. Dots represent each data point from individual cells, boxes represent the interquartile range, and lines display mean with min and max values.

Figure 5. Physiological contributions from NMDAR subunits and changes after hibernation. Change in NMDAR current amplitude after 10 minutes of recording (A, Time control), inhibition of GluN2A (B, 1 µM TCN 201), GluN2B (C, 2 µM RO 25-6981), GluN2C/GluN2D (D, 10 µM QZN 46), and GluN3 (E, 30 µM TK30). Left - change in current amplitude (expressed as % of baseline) in control neurons compared to after hibernation. Right- Example traces of a baseline current overlapped with a current after 10 min of recording (A) or after 10 min of the inhibitor exposure (B-E) in neurons from frogs in control and hibernation conditions. The "after drug" or time controls traces are slightly offset from their respective baseline recordings to enhance visibility. Number of cells (n) and frogs (N) used in these experiments: Time control in control n= 10, N= 4, in hibernation n=11, N=7; GluN2A inhibition in control n= 9, N= 3, in hibernation n=9, N=6; GluN2B inhibition in control n= 9, N= 3, in hibernation n=13, N=6; GluN2C/GluN2D inhibition in control n= 7, N= 3, in hibernation n=10, N=6; GluN3 inhibition in control n= 7, N= 2, in hibernation n=10, N=9.

Figure 6. Single-cell qPCR shows that hibernation changes in the NMDAR subunit profile in motoneurons. Single-cell mRNA abundances for all 7 *Grin* family members. The dominant subunits were *Grin1*, which encodes the obligatory NMDAR subunit, GluN1, as well *Grin2B* and *Grin3B.* Hibernation led to decreases in the abundance of *Grin2C*, *Grin3A*, and *Grin3B.* Sample sizes for each gene are as follows for control and hibernation. *Grin1,* n=19 control, n=20 hibernation; *Grin2A*, n=16 control, n=16 hibernation; *Grin2B*, n=18 control, n=20 hibernation; *Grin2C,* n=19 control, n=20 hibernation; *Grin2D,* n=19 control, n=20 hibernation; *Grin3A,* n=18 control, n=18 hibernation; *Grin3B*; n=19 control, n=20 hibernation. Control cells came from N=6 animals and Hibernation cells came from N=5 animals. * p<0.05; **p<0.01; ****; p<0.001.

Figure 7. Modified NMDAR profile after hibernation protects against network hyperexcitability during hypoxia. Extracellular recordings of motor nerve output from the brainstem-spinal cord preparation. (A_1) Shows the typical response to hypoxia in controls, whereby large amplitude bursting disrupts patterned output from the respiratory network. (A_2) shows motor output during hypoxia in a preparation that was pretreated with D-AP5 to block NMDARs. In the presence of D-AP5, large amplitude bursting is less severe and often does not fully disrupt the respiratory rhythm. This indicates the normal profile of NMDARs contributes to disruptive motor activity in hypoxia. (B_1) Shows patterned motor output from a hibernation preparation over the same timescale as controls. Despite intact NMDARs, there is no observable disruptive motor output. (B_2) A hibernation preparation pretreated with D-AP5, also showing disrupted motor output. (C) Summary data of the degree of large amplitude non-respiratory bursting from n=11 controls, n=9 control+D-AP5, n=8 hibernation, and n=7 hibernation+D-AP5 analyzed by a one-way ANOVA. Non-respiratory bursting was smaller on average after the application of D-AP5, and this was not statistically different from hibernation preparations with or without D-AP5.