1

3   A lizard is never late: squamate genomics as a recent catalyst for understanding sex

4   chromosome and microchromosome evolution

Brendan J. Pinto[1,2,3,*], Tony Gamble[3,4,5], Chase H. Smith[6], Melissa A. Wilson[1,2,7]

1. School of Life Sciences, Arizona State University, Tempe, AZ USA

2. Center for Evolution and Medicine, Arizona State University, Tempe, AZ USA

3. Department of Zoology, Milwaukee Public Museum, Milwaukee, WI USA

4. Department of Biological Sciences, Marquette University, Milwaukee WI USA

5. Bell Museum of Natural History, University of Minnesota, St Paul, MN USA

6. Department of Integrative Biology, University of Texas at Austin, Austin, TX, USA

7. Center for Mechanisms of Evolution, Biodesign Institute, Tempe, AZ USA

Corresponding author email: brendanjohnpinto@gmail.com

Abstract

In 2011, the first high-quality genome assembly of a squamate reptile (lizard or

snake) was published for the green anole. Dozens of genome assemblies were

subsequently published over the next decade, yet these assemblies were largely

inadequate for answering fundamental questions regarding genome evolution in

squamates due to their lack of contiguity or annotation. As the "genomics age" was

beginning to hit its stride in many organismal study systems, progress in squamates

was largely stagnant following the publication of the green anole genome. In fact, *zero*

high-quality (chromosome-level) squamate genomes were published between the years

2012–2017. However, since 2018, an exponential increase in high-quality genome

assemblies has materialized with 24 additional high-quality genomes published for

species across the squamate tree of life. As the field of squamate genomics is rapidly

evolving, we provide a systematic review from an evolutionary genomics perspective.

We collated a near-complete list of publicly available squamate genome assemblies

from more than half-a-dozen international and third-party repositories and systematically

evaluated them with regard to their overall quality, phylogenetic breadth, and usefulness

for continuing to provide accurate and efficient insights into genome evolution across

squamate reptiles. This review both highlights and catalogs the currently available

genomic resources in squamates and their ability to address broader questions in

vertebrates, specifically sex chromosome and microchromosome evolution, while

addressing why squamates may have received less historical focus and has caused

their progress in genomics to lag behind peer taxa.

History and Background

Genome sequencing has revolutionized biology in every group of organisms;

however, some organismal groups have better representation, genomically, than others.

In the intervening years between the first lizard karyotype (Tellyesniczky, 1897) and first

published lizard genome (Alföldi et al. 2011), many questions have been raised where

squamate reptiles stand to provide unique insight into the patterns and processes of

genome evolution including those character states shared with other organismal groups

(e.g. Perry et al. 2021; Pinto et al. 2019a) and those unique to squamates (e.g. Gamble

2019). Namely, squamates provide an invaluable model system for two areas of active

research: (1) the evolution of sex chromosomes (Gamble et al. 2015a) and (2) the

evolution and function of microchromosomes (Perry et al. 2020). We start by briefly

reviewing the development of the history of squamate genomics since its inception.

The argument for why sequencing lizard genomes is necessary, as a departure

from human- and laboratory model-centric taxa, was first made in 2005 (Losos et al.

2005). Five years later, the green anole (*Anolis carolinensis*) genome appeared on

NCBI and the paper published the following year (Alföldi et al, 2011). However,

genomics in squamate reptiles (lizards and snakes) has lagged behind most other

vertebrate groups and *all* other amniote lineages (Hotaling et al. 2021). Another seven

years passed until the second high-quality squamate genome was made available

through the intervention of the DNAZoo sequencing initiative, with the re-scaffolding of

the Burmese python (*Python bivittatus*) genome into a chromosome-level assembly

(Figure 1; Castoe et al. 2013; Dudchenko et al. 2017 & 2018). Herein, we roughly define

"high-quality" genomes as those scaffolded into representative chromosomal linkage

groups (scaffolds) but acknowledge that this ignores the contiguity of the primary

assembly (contigs), which is possibly more important for assembly accuracy and

suggest readers incorporate this metric when both assembling/publishing new

assemblies or choosing an available assembly for use. As of July 12th, 2022, we had

identified 73 'publicly available' genome assemblies across squamate reptiles, 81% of

which were published in the last 5 years (2018–present). Further, it's been as many

years since the last review of squamate genomics (Deakin and Ezaz, 2019). Due to this

lag behind other vertebrate groups, such as birds—who recently surpassed 500

genome assemblies (Bravo et al. 2021) for the approximately 11,162 available bird

species, squamates have largely been overlooked as a key evolutionary group for

genomics studies, with ~11,300 species until just recently represented by the lone

Anolis carolinensis genome (Hotaling et al. 2021; Rhie et al. 2020; Uetz et al. 2022).

Thus, to help refresh this mindset, we provide an up-to-date review to acclimate

scientists, from taxonomically-focused biologists to computational biologists, on the

state of genomics within squamate reptiles—a key, yet understudied, model group to

address important biological questions in an evolutionary context.

Squamate Genomics Today

In Appendix I, we aggregated a near-complete list of squamate genome

assemblies and assembly information to (1a) characterize why squamate genome

assemblies have lagged behind other groups and (1b) identify specific taxonomic

groups within the field that are lacking, (2) interrogate various assembly metrics across

taxa to identify potential trends in data generation and assembly, and (3) discuss how

currently available squamate genomes, although lacking in phylogenetic density

(number of taxa), still possess the phylogenetic breadth to revise how we think about

vertebrate genomics, specifically (3a) sex chromosome evolution and (3b)

microchromosome evolution. As of mid-2022 (the data collection cutoff date for this

manuscript), among all available squamate genome assemblies, snakes outnumbered

all others combined (37 snake vs. 34 lizard assemblies). However, when accounting for

only high-quality assemblies the numbers reverse (9 snake vs. 16 lizard assemblies).

Importantly, all but one of these assemblies was published in the last five years (Figure

1).

One important factor in the historical lag in squamate genomics behind other

amniotic groups is likely, at least in part, due to faith placed in large-scale sequencing

initiatives that have then prioritized other groups. In short, the future of high-quality

squamate genome generation is in the hands of those with a keen interest in reptiles.

Large-scale sequencing initiatives with large resource pools, such as the Vertebrate

Genome Project (VGP) consortium, have largely neglected this speciose group of

amniotes (Genome 10K Community of Scientists, 2009). For scale, according to the

IUCN Red List (i.e. Uetz et al. 2022), there are more non-avian reptiles (11,690) than

avian (birds; 11,162)—even approaching twice as many species of squamates (11,300)

than mammals (6,578)—however, as of mid-2022 of the 129 amniote genomes

available through the VGP 33% (43/129) were birds and 21% (27/129) were mammals,

with a staggering 1.5% (2/129) and 3% (4/129) for squamates and non-avian reptiles,

respectively (https://hgdownload.soe.ucsc.edu/hubs/VGP). Without changes to these

trends, there appears to be little hope for squamate genomes to be generated *en masse*

through these types of initiatives. Funding agencies appear to be responding to this

need and funding genome projects by smaller research groups who are excited about,

and committed to, assembling reptile genomes (authors pers. obs.).

One issue that continues to inhibit accurate characterization and analysis of

squamate genomes broadly, is the lack of centralization, or even a semi-centralization,

of the available genomic resources (Appendix I). While most genomes have made their

way to NCBI's GenBank or other international government-sponsored analogs (e.g.,

ENI, CNCB), many remain scattered throughout unincorporated repositories that remain

difficult to track down *a priori* (e.g., Figshare, GigaDB, DNAZoo, etc.). However, we

believe this issue is larger than researchers simply not wanting to centralize these data

for broader ease of access. From a researcher perspective, submitting a genome to

GenBank (or similar repository) is a non-trivial task and becomes extremely

cumbersome when attempting to accompany the genome assembly with annotation

information generated "in-house". Indeed, while it is a trivial task to upload a gzipped

FASTA and GFF file to a third-party data repository (e.g. Figshare), or even simply a

FASTA genome file to Genbank, uploading the GenBank-specific formatted

assembly/annotation has multiple challenges. For example, most annotation programs

don't generate the required files for downstream use, and it then falls on researchers to

then generate these files post hoc, opt for a third-party data repository to save

significant time and effort, or some hybrid between the two—with the assembly

cataloged on a government server and the annotation housed in a third-party repository.

Although there are a few available programs that attempt to bridge this gap by piping

necessary annotation software together, they are not without their difficulties (Banerjee

et al. 2021; Cantarel et al. 2008; Hoff et al. 2019; Palmer, 2018). There remains a need

to centralize genome assemblies with consistent, high-quality annotation information. At

present, the ideal situation appears to be submitting a genome to NCBI and inquiring

with RefSeq about providing annotation, which will generally provide high-quality

genome annotations assuming sufficient RNAseq data is available. However, this

avenue can only progress once the genome has been publicly released and can take

many months due to an ever-growing queue. Accompanying high-quality genome

assemblies with complimentary genome annotation is essential for drawing significant

biological insights from new high-quality, reference genome assemblies. Thus, we must

forewarn that although the increased quality of DNA sequencing technologies and

genome assembly tools have caused a 'boom' in genome assembly generation across

the tree of life, the subfield of squamate genomics may 'bust' under its own weight if

steps are not taken soon to address the laborious nature of genome annotation and

data dissemination. We see potential avenues for cloud computing to lessen this burden

for individual research groups as databases, such as NCBI's Sequence Read Archive

(SRA), move to becoming available on the cloud (https://anvilproject.org/ncpi). It is

widely known that NCBI's GenBank, for example, provides extensive curation services

and continues to expand its functions and utility, such as recently adding the NCBI

Datasets (https://www.ncbi.nlm.nih.gov/datasets/) for querying data across studies and

the Comparative Genome Viewer (https://ncbi.nlm.nih.gov/genome/cgv) for

understanding synteny across reference assemblies. We hypothesize that these

functions will only increase in utility if the activation energy for data uploading were to be

reduced in some way.

(1a) Why have squamate genomics lagged behind other groups?

Two major factors appear to have synchronously contributed to the lag in

squamate genome sequencing relative to other vertebrate groups: genome size and

funding. For most vertebrate groups, genomic investigations have benefited from either

small genome sizes (i.e. could accomplish more with less) and/or substantial funding

models (i.e. could accomplish more with more). For example, birds (637 assemblies

representing 11,162 species; Bravo et al. 2021) and fishes (594 assemblies

representing 32,000 species; Randhawa & Pawar, 2021) each possess some of the

smallest vertebrate genomes described—most within the ~0.4-1.4Gb range. While far

larger genome sizes occur in mammals (~2.5-3.5Gb), applied funding from health and

agricultural sources (far exceeding that allocated to other vertebrate groups, such as

squamates) have offset similar phenomena in the field of mammal genome sequencing

(Supplemental Table 1). In the most extreme case, amphibian genomes are even larger

and suffer more greatly than squamates due to this form of genome size bias, however,

further extrapolation here is beyond the scope of the current article. At a glance,

squamates have an average genome size of 1.73Gb (N=71), ranging from 1.1Gb in

Crotalus pyrrhus assembly (Gilbert et al. 2014) to 2.86Gb in *Sceloporus occidentalis*.

However, this estimate is fraught with bias due to an overabundance of low-quality

short-read assemblies that likely skew the genome size estimates lower than reality

(Supplemental Figures 2 and 3). We can roughly account for this by discarding all

genome size estimates from primary assemblies derived from short-read technologies,

assuming long-read primary assemblies are better representations of the repeat content

within a genome (Rhoads and Au, 2015). This provides a revised estimate of the

approximate average genome size in squamates of 1.86Gb (N=17), ranging from

1.39Gb in *Lacerta agilis* to 2.86Gb in *S. occidentalis*. Thus, the larger genome sizes in

squamates, albeit on average still ~0.8-1Gb smaller than mammals, combined with less

overall funding than mammalian taxa, has likely led to a stagnation in high-quality

genome assemblies in squamates—that is until the cost of sequencing decreased

exponentially over the past five years (Wetterstrand, 2021). Thus, as sequencing costs

have declined exponentially, requiring less funding to accomplish more sequencing, the

subfield of squamate genomics has finally erupted and is beginning to flourish (Figure

1).

(1b) What taxonomic groups remain unsampled?

An overarching theme of the current state of squamate genomics is that, while

few groups are adequately represented in terms of genomic resources (such as elapid

snakes – 15 genomes from 395 species), most squamate groups are in dire need of

additional high-quality genomic resources (such as geckos, from our very biased point-

of-view, which include 2,186 species, but only six genomes). However, there are many

extremely diverse and evolutionarily important groups that are completely absent, such

as chameleons (222 species), amphisbaenids (182 species), and scincomorphs (1,886

species) (Figure 2). In fact, approximately five years ago, a high-quality multi-tissue

transcriptome was published for the veiled chameleon (*Chamaeleo calyptratus*) with an

accompanying call for additional genomic resources to be generated for this extremely

interesting clade (Pinto et al. 2019b). However, to date, there has yet to be a single

genome assembly of any quality, made publicly available for a chameleon—with a

similar situation at play in scincomorphs and many other squamate families (Figure 2).

Indeed, of the 46 squamate families that appear in Figure 2, 31 families including all

chameleons and scincomorphs—occurring globally, except Antarctica—have no publicly

available reference genomes. Future directions in squamate genomics should focus on

including these missing taxa as important players in the investigations in the genomics

of vertebrates.

(2) Trends in data generation and assembly

Regardless of sequencing methodology implemented, most empiricists have

become aware that the quality of sample collection and preparation can "make or break"

a genome assembly experiment. This includes every stage of sample preparation up to

its conversion from bases to bytes, including, but not limited to: tissue selection,

dissection, storage, extraction, library prep, sequencing, and assembly (Dahn et al.

2022; Pinto et al. 2022 & 2023). Many squamate species are rare/hard to collect, have a

limited distribution (Meiri et al. 2018), and lack material in museum collections adequate

for long-read sequencing or chromatic-contact sequencing (HiC). Most species will need

new specimens to be collected specifically for a genome sequencing project.

Fortunately, relative to some other animal groups (for example, freshwater bivalves

(Smith, 2021) and *Xiphophorus* fishes (author's pers. obs.)), squamate DNA appears to

remain remarkably stable throughout this process, which provides some relief for field

collection and sub-optimal tissue conditions—preferring blood or liver tissue when

available (Dahn et al. 2022; Pinto et al. 2022 & 2023). These factors prime squamates

to benefit from recent advances in sequencing technology – like long, accurate

sequencing reads – that have opened new doors in genome assembly. Just as the

publication of the first human reference genome at the turn of the century signaled the

beginnings of the "genomics age", the recent publication of the complete human

telomere-to-telomere (T2T) genome assembly has signaled a 'rebirth' of the genomics

age, where now all model systems can be subject to high-quality reference genomes for

relatively low cost, including squamates (Nurk et al. 2022; Pinto et al. 2022, 2023; Rhie

et al. 2021; Sun et al. 2021). Recent advances in increasing contiguity of primary

genome assemblies has been driven by third generation sequencing technologies,

including Pacific Biosciences (PacBio) and Oxford Nanopore platforms (e.g. Nurk et al.

2022; Peona et al. 2020). For the past two years, PacBio High Fidelity (HiFi) reads have

shown that high-accuracy reads (~20 kb; phred quality scores ~20+) can outperform

longer reads with lower accuracy (~40 kb+; phred quality scores ~10) in many cases—

certainly at the cost-per-base (Lang et al. 2020; Peona et al. 2020; Vollger et al. 2020).

However, recent data also confirmed that some genomic regions require ultra-long read

lengths to overcome extremely long stretches of repetitive DNA, some lengths of which

may still be unachievable, but certainly enforces a hard ceiling for the 'coverage-to-

contiguity' ratio at around 30X when using HiFi data alone (Pinto et al. 2023; Sun et al

2021). That said, Oxford Nanopore's forthcoming Q20 chemistry (Kit 14 with the

R10.4.1 flowcell) may provide the missing link in completing Telomere-to-Telomere

(T2T) genome assemblies that makes them more approachable to squamate

researchers on a tight budget.

One way of accelerating genome assembly generation across squamates would

be to decentralize sequencing and assembly. This is currently how squamate genomics

has advanced and assisting the field in this endeavor is one goal of this manuscript.

However, it is far from as decentralized as one might imagine. Indeed, when delving into

where lepidosaur (squamates and the tuatara) genome assemblies are derived from,

the research group doing the sequencing (inferred via first and last authorships), and

the vast majority of assemblies come from research groups in the 'global north' (75%;

55/73) and China (21%; 15/73); totaling 96% of all assemblies (70/73). This leaves only

three assemblies having been generated by the rest of the global community.

Importantly, these numbers *do not* account for the middle-author contributions to

projects made by members of the global south, which are no doubt significant. In

response to these types of devastating numbers, organizations, such as GetGenome

(getgenome.net), may be helpful in reducing this disparity between the global north and

south. Importantly, organizations like this that formally empower groups to conduct this

work in-house, instead of outsourcing to a consortium, will likely produce greater

innovation using these data in the long run (e.g. Hofstra et al. 2020) and could help

more broadly mitigate the current state of scientific exclusion of the global south within

the subfield. As such, it is important to note that since the subfield of squamate

genomics is relatively young we are in an optimal position to lead an equitable

globalization effort moving forward with regard to data generation and usage—an

important steppingstone for the herpetological field more broadly.

Practical considerations regarding sex chromosomes in squamate genomics

More generally, as genome sequencing technologies are capable of producing

both long and accurate sequence reads, an important step to genome assembly is

producing fully phased, or haplotype-resolved, genome assemblies in place of

traditional chimeric assembly where alleles are assembled together (Cheng et al. 2021

& 2022). This may allow for the resolution of divergent genomic regions of biological

importance, such as polyploid genomes, heterozygous inversions, alternative splice

variants, and sex chromosomes (XY or ZW). Indeed, once haplotype-resolved genomes

become common within squamates, sex chromosomes within the assemblies will be

phased—as they are in the human genome and some others (Nurk et al. 2022; Webster

et al. 2019).

Once both sex chromosomes (X and Y or Z and W) are present in the reference

assembly, researchers will need to specifically assess and account for the sex

chromosome complement when conducting bioinformatic experiments, such as read

mapping and variant calling (Carey et al. 2022; Olney et al. 2020; Pinto et al. 2023b;

Webster et al. 2019). To effectively account for sex chromosome complement in an

assembly, haplotypes of the sex chromosomes must be resolved. In tandem with this

paper, the first attempt at generating a haplotype-resolved genome of a squamate, a

temperature-dependent sex determining gecko—the leopard gecko, *Eublepharis*

macularius, was published (Pinto et al. 2023). With this along with other recent

assemblies, the advent of reference quality, phased genomes for squamate taxa has

become achievable for the average research group and bodes well for the future study

of sex chromosomes across squamates.

(3a) Genomics and sex chromosomes in squamates

Squamates are an invaluable model system for studying sex chromosome

evolution. Within their ranks all three major modes of vertebrate sex determination

occur: environmentally determined sex (temperature dependence) and genetic sex

determination (both male heterogamety [XX/XY] and female heterogamety [ZZ/ZW]

systems), with multiple independent transitions among the three mechanisms (Gamble

et al. 2015a; Stöck et al. 2021). Studying squamates provides a powerful system to

better understand the gaps in our knowledge of sex chromosome evolution broadly;

specifically questions such as, (I) are some linkage groups more likely to be recruited as

a sex determining role than others?, (II) are ancient sex chromosome systems an

evolutionary trap that species cannot escape?, and (III) how do mechanisms of dosage

balance and compensation between the sexes evolve? (e.g. Gamble et al. 2015a;

Kratochvíl et al. 2021; Nielsen et al. 2019; Rupp et al. 2017). We explore these topics

framed by how modern squamate genomics stand to help answer these questions.

(I) Are some linkage groups more likely to be recruited as a sex determining role

than others?

The identification and characterization of sex chromosome systems are perhaps

the most well-reviewed aspect of squamate genomics—whose study has also been

intimately associated with the advent of genomics in squamates—with progress

increasing exponentially in recent years (Gamble, 2010; Gamble et al. 2015a, 2017,

2018; Kratochvíl et al. 2021; Pinto et al. 2022; Stöck et al. 2021). Four species-rich

clades, with known, conserved sex chromosome systems – pleurodonts (iguanas, spiny

lizards, and anoles, excluding corytophanids); caenophidian snakes; skinks; and

lacertids – make up approximately 60% of squamate species (Rovatsos et al. 2014,

2015, 2019a; Nielsen et al. 2019; Kostmann et al. 2021). The remaining 40% of

squamate species are in clades with varying levels of sex chromosome conservation,

although transitions are likely common in many of these groups (Gamble et al. 2015a;

Gamble et al. 2017; Nielsen et al. 2018; Keating et al. 2022; Pinto et al. 2022). Given

the available data it has been suggested that linkage group recruitment as sex

chromosomes is nonrandom, i.e. some linkage groups are more likely to be recruited as

a sex chromosome than others (Kratochvíl et al. 2021). However, the pattern was weak

and the discovery of additional linkage groups acting as sex chromosomes in geckos

and dibamids requires a re-evaluation (Pensabene et al. 2023; Pinto et al. 2022;

Rovatsos et al. 2022). Additionally, inferences that all taxa within a clade share an

ancestral sex chromosome, i.e. knowing 5% of taxa sex chromosome systems and

inferring that we know ~60%, is drawn from Occam's razor using sparse sampling

(Kostmann et al. 2021), but in squamate sex chromosome evolution, where sex

chromosome turnovers are commonplace, this kind of assumption has been shown to

be untrue (e.g. Gamble et al. 2017). Thus, it stands to reason that this 60% figure may

be an overestimate. Fortunately, recent advances in DNA sequencing technologies

have allowed us to sample more broadly and ask finer-scale questions about how sex

chromosomes originate, degenerate, and turnover (e.g. Acosta et al. 2019; Gamble et

al. 2015a; 2017; 2018; Keating et al. 2020; Kostmann et al. 2021; Nielsen et al. 2018;

2019; 2020; Pinto et al. 2022; Rovatsos et al. 2019b; 2022); so the intertwined nature of

developing squamate genomics and sex chromosome evolution presents great promise

for future work in identifying and characterizing sex chromosome linkage groups across

squamates.

The most conclusive evidence of shared ancestry of a sex chromosome system

is the identification of a conserved primary sex determiner (or primary sex determining

gene; PSD) among focal taxa (such as *Sry* in therian mammals; Graves, 2008).

However, no prior publication has yet assembled the X/Z chromosome and then
identified a putative PSD in a squamate, until now. Indeed, high-quality genome
assemblies and annotations are only recently allowing us to confidently implicate
putative PSDs in squamates. The first example to our knowledge being the Puerto
Rican leaf-litter gecko, *Sphaerodactylus townsendi* (Box 1). It's worth noting that
previous implications of PSDs in squamates (*Pogona vitticeps* (*Sr1*) and anguimorphs
(*Amh*): *Varanus komodoensis* and *Heloderma suspectum*) were based on incomplete
catalogs of Z-linked genes (Deakin et al, 2016; Rovatsos et al. 2019; Webster et al.
2023). In an ideal world, assembling both the complete X/Z and Y/W would yield the
best possible candidate PSD. Beyond implicating a candidate PSD, one downstream
issue that is in the process of being overcome is that even upon the
identification/confirmation of a putative PSD, we have limited capability to perform
functional tests to confirm a putative sex-determining gene. Although there is significant
progress happening on this front with the first successful gene editing in an *Anolis* lizard
and a gecko (Rasys et al. 2019; Abe et al. 2023). High-quality genome assemblies and
annotations are crucial to expanding the utility of functional genomic tools in squamates.
Thus, although high-quality genomes are now allowing us to better characterize putative
PSDs in squamates, we're still a few years away from using gene editing to confirm
these putative PSDs in different squamate species.

Box 1: *Sex chromosomes and sex determination in squamates, a case for high-quality genome annotations*

In 2021, the chromosome-level genome of *Sphaerodactylus townsendi* helped elucidate the dynamic evolution of sex chromosomes within this genus of geckos (Pinto et al. 2022). However, when examining the annotated gene content within the identified sex determining region (SDR) in the initial annotation [MPM_Stown_v2.2], we found no sign of a putative sex determining gene (gene known to have a consequential role in the vertebrate sex determining pathway). Through collaboration with NCBI RefSeq, this genome was re-annotated using only existing RNAseq data (i.e. no new transcriptomic data was generated between annotations) using the NCBI Eukaryotic Genome Annotation Pipeline. NCBI Annotation Release 100 of MPM_Stown_v2.3 provided significant improvements to the annotation quality (BUSCO completeness of annotated peptides from 61.5% to 92.5% using BUSCO [v5.1.2]). When re-examining the SDR of *S. townsendi* using this new annotation, a candidate primary sex determining gene (PSD) became clear, anti-Müllerian hormone receptor 2 (*AMHR2*). Indeed, *AMHR2* has been identified as the independently-evolved primary sex determining gene in at least two groups of fish, fugu and ayu (Kamiya et al. 2012; Nakamoto et al. 2021) and its inactivation causes male-to-female sex reversal in the Northern Pike (Pan et al. 2022). This example supports that—even without generation of additional data—high-quality annotations can be generated for divergent species with minimal available transcriptomic data. However, it is also likely that this high-level of quality for genome annotation is beyond the reach of many (if not most) biology-focused research groups, as it was for us (Pinto et al. 2022). We suggest that this may serve as motivation for the generation of the development of additional genome annotation pipelines or the adaptation of existing pipelines to be more approachable 'lay-empiricists' interested in answering these fundamental types of questions in newer model systems.

(II) Are ancient sex chromosome systems an evolutionary trap that species
cannot escape?

Early sex chromosome work highlighting mammalian and avian taxa suggested
that perhaps ancient sex chromosome systems may become so entangled in the
biology of the organisms' development that it served as an "evolutionary trap", to which
there was little chance of escape (Bull, 1983; Bull and Charnov, 1985; Gamble et al.
2015a; Nielsen et al. 2019; Pokorná and Kratochvíl, 2009). Indeed, many taxa that
possess an ancient, ancestral sex chromosome system appear to remain evolutionarily
ensnared within it, including mammals (XY), birds (ZW), *Drosophila* (XY), lepidopterans
(ZW), and "advanced" snakes (ZW) (Bachtrog et al. 2014; Rovatsos et al. 2015; Gamble
et al. 2017; Graves, 2008; Ohno, 1967; Webster et al. 2023). To our knowledge, there
are few empirical examples of taxa escaping old, degenerated sex chromosome
systems (Rovatsos et al. 2019c; Terao et al. 2022). However, one possible example
within squamates are the basilisk and casque-headed lizards (Corytophanidae) that
possess a different sex chromosome system than all other pleurodonts. Phylogenetic
uncertainty plagues this claim as a conclusive case of escaping the trap and more work
is needed (Acosta et al. 2019; Nielsen et al. 2019). However, as more and more
transitions among sex-determining systems have been identified it is unclear whether all
sex chromosomes are destined to become traps. Because they have a variety of sex-
determining systems with numerous transitions among them (Gamble et al. 2015a;
Gamble et al. 2017; Pokorná and Kratochvíl 2009; Ezaz et al. 2009) squamates are an
excellent model to investigate this question.

(III) How do mechanisms of dosage balance and compensation between the
sexes evolve?
Perhaps the scarcest data available regarding sex chromosomes in squamates
lies in how these animals deal with gene dosage changes that evolve in response to the
degeneration of the sex-limited sex chromosome. In many well-characterized animal
model systems, such as the XY systems of mammals and fruit flies or the ZW systems
in birds and moths, differences in gene copy number between the sex chromosomes
can result in myriad disparate outcomes (Bachtrog et al. 2014; Gu and Walters, 2017;

Vicoso and Bachtrog, 2009). Sex chromosome dosage work contains two interrelated

questions specific to genes within the non-recombining region of the sex chromosomes,

(1) what is the gene dosage between the sexes, relative to each other, known as

dosage balance, and (2) what is the gene dosage of the sex chromosomes in each sex

relative to the ancestral (autosomal) condition, known as dosage compensation (Gu and

Walters, 2017). For instance, in mammals and moths there are mechanism(s) to

globally silence one of the two X/Z chromosomes in homogametic individuals to balance

the dosage between the sexes; however, although global expression between the sexes

is equal, expression in both sexes is lower than the ancestral condition. In other words,

mammals and moths possess dosage balance mechanisms, but not those for dosage

compensation. Meanwhile, sex chromosomes in fruit flies are both balanced and

compensated for, and birds are neither balanced or compensated (Gu and Walters,

2017). Because the outcomes of changes in gene dosage are disparate across taxa,

more naturally occurring 'evolutionary experiments' are desperately needed to better

understand the underpinnings of these phenomena.

Due to the lability of sex chromosomes across squamates, they may again play a

pivotal role in deciphering the broader mechanistic underpinnings of sex chromosome

gene dosage. Indeed, a unique characteristic of squamates relative to most other

amniotes is that, due to the high rates of sex chromosome turnover, one can more

easily infer the ancestral, autosomal gene expression level of multiple sex chromosome

systems using closely related species (e.g. Keating, 2022) instead of using distant

proxies, which may introduce additional uncertainty (e.g. Webster et al. 2023). This

concept has been used across taxonomic groups to elucidate the evolutionary history of

a variety of traits (e.g. Blount et al. 2018; Sackton and Clark 2019; Smith et al. 2020;

Neemuchwala et al. 2023). In squamates specifically, this concept has been used to

study many other independently-evolved traits such as adhesive digits (Gamble et al.

2012) and photic activity patterns (Gamble et al. 2015b; Pinto et al. 2019c), among

many others. Thus, to more effectively study how dosage compensation mechanisms

evolve in amniotes, squamates are an important model system to utilize. However, to-

date the lack of genomic resources have especially hindered these investigations. This

is a direct result of the lack of high-quality genomic resources available for squamates

prior to 2018—because knowing how genes are linked together is necessary

information to investigate dosage (Keating et al. 2022; Vicoso et al. 2013; Webster et al.

2023).

As stated in the introduction, the only high-quality genome available prior to 2018

was the *Anolis carolinensis* genome (Alföldi et al. 2011), as such, we know that *Anolis*

carolinensis possesses both dosage balance and compensation (Marin et al. 2017;

Rupp et al. 2017). Clever application of the *Anolis* genome to similar analyses in snakes

also identified relatively early on that caenophidian, so-called "advanced", snakes, like

birds, lack both dosage balance and compensation (Vicoso et al. 2013), which was later

confirmed using additional high-quality snake resources (Schield et al. 2019). More

recently, conceptually similar approaches to those used by Vicoso et al. (2013) have led

to an increase in transcriptomic data mapped to a distant relative genome to elucidate

presence/absence of dosage balance in corytophanid (Pleurodonta), pygopodid

(Gekkota), and anguimorph lizards (Nielsen et al. 2019; Rovatsos et al. 2019b; 2021).

Additional work, including additional genomic data, have led to additional findings that

both anguimorphs and diplodactylids (Gekkota), similarly to birds and snakes, appear to

lack both dosage balance and compensation (Keating, 2022; Webster et al. 2023).

Indeed, given the sheer diversity within Squamata our knowledge of how these animals

handle dosage differences between the sexes is exceptionally sparse.

Trending with previous sections regarding the necessity of high-quality

annotations to accompany high-quality genome assemblies (e.g. Box 1) also apply *ad*

infinitum to studying sex chromosome dosage. When examining dosage there are

essentially two scales one can use to examine differences between the sexes (a) global

and (b) positional scales. (a) Global can only be used to study dosage balance at a

broad scale, where comparing gene expression differences between males and females

on different linkage groups (e.g. Nielsen et al. 2019; Rovatsos et al. 2019b; 2021).

However, as the name might imply, this scale provides little insight into the fine-scale

processes of sex chromosome evolution. Indeed, when a high-quality reference is

available for a given species (or close relative) one can conduct (b) finer-scale,

positional examinations of gene expression across the pseudo-autosomal (PAR)

boundary and decrease noise from 'misplaced' genes that are no longer linked in the

focal taxon, even if they are in a distant relative such as chicken or *Anolis* (Schield et al.

2019; Webster et al. 2023). Further, when examining expression on smaller

chromosomes with relatively few genes, missing genes due to poor annotation quality

can decrease statistical power to detect changes in dosage significantly (e.g. Keating,

2022; Webster et al. 2023). Thus in addition to addressing broader questions, such as

those discussed above, high-quality annotations are necessary to accompany new

reference genomes being generated to better understand how sex chromosome dosage

evolves, identify putative sex determining genes (Box 1), and more generally to better

characterize the "sexomes" of squamate reptiles (Stöck et al. 2021).

(3b) Microchromosome evolution

In chicken, early 20[th] century cytologists identified 12 easily-distinguishable large

chromosomes and an additional 18+ smaller, dot-like chromosomes; Dr. Nettie Stevens

notably prefaced this finding in her laboratory notebook with, "impossible to tell how

many small ones" (Boring, 1923; Hance, 1924). Later work coined the term

"microchromosomes" to describe these 'innumerable' small chromosomes and their

larger counterparts as "macrochromosomes" (Newcomer, 1957; Ohno, 1961;

Yamashina, 1944). However, no universally agreed upon definition of a

microchromosome has yet to be established in the literature, certainly not since the

advent of high-quality genome assemblies in reptiles (Boring, 1923; Fillon, 1998;

Newcomer, 1957; Ohno, 1961). Indeed, at the advent of genome sequencing in birds,

chicken chromosomes were arbitrarily grouped as macrochromosomes (1-5),

intermediate chromosomes (5-10), and microchromosomes (11+) (Hillier et al. 2004).

Subsequent studies have either used these criteria, grouping macrochromosomes and

intermediate chromosomes as macrochromosomes, ranging in size from ~23Mb to

~200Mb (O'Connor et al. 2018), or established their own criteria for an arbitrary cutoff,

such as 10Mb, 30Mb, or 50Mb (Karawita et al. 2022; Perry et al. 2021; Srikulnath et al.

2021; Waters et al. 2021). However, these arbitrary categorizations—enforced across
vertebrates—make direct comparisons between taxa difficult and may encourage
spurious correlations from these artifacts. These factors, among others, warrant a re-
analysis of "what is a microchromosome?" and "why are they important?" and we
demonstrate how squamate genomics provides vital insight into these questions.

Microchromosomes, no matter how they are defined, are present in most
vertebrate groups (Srikulnath et al. 2021). However, their evolution remains murky –
they have either been inherited from a common ancestor and lost independently
multiple times or gained and lost independently multiple times. Since
microchromosomes have historically been inhibitively difficult to assemble prior to long-
read sequencing technologies, studies detailing finer-scale analyses have been lacking.
Importantly, studies have lacked proper controls in an evolutionary context. No analysis
to-date of microchromosomes using genomic sequence data has included, and
specifically accounted for, the two squamate lineages that are known to not possess
microchromosomes, i.e. geckos and lacertids (Deakin & Ezaz, 2019; Olmo, 1986; Olmo
et al. 1990; Pinto et al. 2022; Srikulnath 2013; Tellyesniczky, 1897). Indeed, past
studies excluding these groups have shown that microchromosomes have a set of
distinct properties relative to macrochromosomes, including higher GC content, higher
gene density, and a distinct nuclear architecture (Perry et al. 2021; Srikulnath et al.
2021). Here, we take a fresh look across vertebrates (mostly reptiles) as a primer to
better understand the biology of microchromosomes and their evolution.

Are microchromosomes conserved across reptiles?

Microchromosomes were likely present in the ancestor of all reptiles, including
birds (Waters et al. 2021). However, within squamates, the hypothesis that the MRCA
possessed microchromosomes has never been explicitly examined with synteny
analyses including both geckos and lacertids. Support for an ancestral lack of
microchromosomes in squamates would appear as strong conservation of linkage
groups between geckos and lacertids regarding microchromosome fusions, which we
do not see (Figure 3). Instead, we observe lineage-specific fusions of

microchromosomes to different macrochromosomes in geckos and lacertids.

Furthermore, there is a near 1:1 relationship of microchromosomal synteny across

snakes, teiids, and anguimorphs—spanning the phylogenetic breadth of non-gekkotan

squamates and extending to birds (Figure 3). Thus, geckos and lacertids have most

likely lost microchromosomes twice independently. Additionally, when losing

microchromosomes in both taxa it is apparent that, although their absolute size tends to

fluctuate between taxa, their relative sizes tend to stay the same (i.e. small

chromosomes tend to stay small)—unless they become fused to other chromosomes,

which contrasts the patterns seen in some birds, such as chicken—which has gained

multiple microchromosomes relative to the inferred ancestral karyotype (O'Connor et al.

2018). Given the currently available data, these additions to the microchromosome

evolution discourse provide some insight into the evolutionary processes involved in the

gains/losses of microchromosomes in certain vertebrate lineages.

What is a microchromosome?

A null prediction of genomic composition of a chromosome might suggest, since

the majority of an animal's DNA is noncoding—all else being equal—that smaller

chromosomes should have higher gene density. Similarly, GC-biased gene conversion

may also lead to overall higher GC content on smaller chromosomes (Fullerton et al.

2001)—since smaller chromosomes also have less space to recombine this GC bias

should, in-turn, scale with chromosome size. Therefore, to truly deviate from this null

expectation, a "microchromosome" should deviate from what's observed from closely

related species that don't possess microchromosomes. These expectations are

supported by a strong linear relationship between chromosome size and gene

content/GC content in species without microchromosomes, which is exactly what we

see in the gecko (Pearson's r: gene content = 0.865***/GC content = -0.781***), lacertid

(gene = 0.699**/GC = -0.727**), alligator (gene = 0.936***/ GC = -0.720*), and even

human (gene = 0.857***/GC = -0.598*) (Supplemental Figure 1, panels A–D).

With a null expectation between chromosome size, gene and GC content

established, we examine deeper when/if deviations occur in taxa that possess

microchromosomes. We find that non-avian reptiles do not deviate from the expectation
of GC content based solely on chromosome size. Evidence for this observation is two-
fold, (1) the overall range of GC content remains constant in non-avian reptiles, from
about 42-52% genome-wide, and (2) the correlation between GC content chromosome
size remains constant (Pearson's r >-0.6) and significant in all taxa except the snake
(Supplemental Figure 1, panels E–H)., which has distinct distribution of data—
compared to all other taxa—with an apparent break between chromosomal GC content
between 40-42% (Supplemental Figure 1, panel G). Since the extremely high GC
content and presence of immensely small microchromosomes (<10Mb) in birds are both
independently derived since their divergence with their closest extant relatives
(crocodilians and testudines, respectively), it's difficult to draw broader conclusions from
analyzing bird genomes alone. In this context, birds also do not appear to deviate from
the expectation set by other vertebrates, however, the shear diminutiveness of their
microchromosomes appears to have caused them to increase GC content much higher
than non-avian vertebrates have attained, ranging from 40-63% (Supplemental Figure
1, panels I–L). For birds, this excessive GC content in microchromosomes may be
related to the presence of both endothermy and microchromosomes, as higher GC
content is associated with thermostability of DNA molecules (Bernardi and Bernardi,
1986).

Since the advent of HiC in squamates (2018–present, e.g. Pinto et al. 2022;
Shield et al. 2019) new understandings of microchromosome evolution have begun to
emerge, yet are still being explored at a fundamental level. As the lines between macro-
and micro-chromosomes somewhat blur in the age of chromosome-level genome
assemblies, recent work has begun exploring the nuclear organization of
microchromosomes in reptiles (Perry et al. 2021). Specifically, HiC data implicates a
distinct intra-cellular compartmentalization of microchromosomes in the nucleus (Perry
et al. 2021; Waters et al. 2021). Importantly, previous investigations were either missing
data from geckos and lacertids or used arbitrary cutoffs to infer the presence of
microchromosomes when they weren't present (Figure 4).

It is difficult to generalize across study systems when using arbitrary numerical

cutoffs for what makes a microchromosome in different taxa. We briefly explored this

concept in available bird data (chicken, zebra finch, and black swan; Supplemental

Figure 1, panels I–K). Specifically, we used two arbitrary cutoffs to group macro/micro

chromosomes (1) microchromosomes <30Mb and (2) microchromosomes <10Mb. We

can see that chromosomes <10Mb possess far more extreme values of gene and GC

content than those >10Mb more-or-less meeting the *a priori* expectations of

microchromosomal composition. However, using a <30Mb cutoff is more representative

of the original karyotypic 'definition' of a microchromosome (Boring, 1923). Importantly,

when investigating the correlation between chromosome size, GC content, and

chromosomal interaction within a single species, the black swan showed a

disassociation between chromosome size and (1) higher GC content and (2) chromatin

conformation that are both generally associated with microchromosomes (Figure 4a;

Supplemental Figure 1, panel K). We see that although a <30Mb cutoff is representative

of the karyotypic definition of microchromosome, only chromosomes at a <15Mb cutoff

appear to be enriched for the predicted microchromosomal interaction that 'true'

microchromosomes are expected to possess (Figure 4a; Perry et al. 2021). Thus, it is

unclear how to best navigate categorizing chromosomes as macro/micro and the

downstream implications on studying the innate properties of these entities.

These inconsistencies bring up a logical conflict as to the nomenclature of

microchromosomes. At this point, there are two equally valid ways to 'define' a

microchromosome, (a) the historical definition of small dot-like chromosomes that are

difficult to pair cytogenetically (e.g. Boring, 1923; Hance, 1924) or (b) a grouping of

relatively small chromosomes within a genome that possess a distinct nuclear

organization (Figure 4; Perry et al. 2021; Waters et al. 2021). It's important to note that,

like in the tegu (Figure 4b), these definitions do not necessarily conflict, however, like in

the swan (Figure 4a), they may. By either definition, it is clear that some taxa possess

microchromosomes and others do not (Figure 4; Olmo et al. 1990; Perry et al. 2021;

Pinto et al. 2022; Srikulnath et al. 2021). Thus, it is important to resolve these conflicts

by using specific language that conveys these intricacies. We suggest that rather than

attempt to redefine what a microchromosome is *a posteriori*, we qualify the evidence

weighted to how we describe microchromosomes. Specifically, at least until we better

understand the nuclear function of the observed nuclear organizations of

microchromosomes, we can retain the historical definition (1) of microchromosomes and

specifically preface those microchromosomes that are isolated in the nucleus as

"organized microchromosomes". For an example under this framework, the black swan

(Figure 4a), all chromosomes <30Mb (10-28) are microchromosomes, but only

chromosomes <15Mb would likely be considered "organized microchromosomes";

however, in the tegu (Figure 4b) all microchromosomes would be considered organized

microchromosomes. This type of classification may help clarify communication

regarding microchromosomes and any potential functional role these sequestered

microchromosomal foci may have across taxa. Further investigations into the evolution

of microchromosomes are necessary and likely ongoing, however, to fully understand

how microchromosomes evolve the field will need access to additional genomic data

from across squamates.

611   Conclusion

In conclusion, as prices in genome sequencing continue to fall, squamate

genomics will exponentially increase (see also Card et al. 2023). However, keeping up

with this progress will not be a trivial task. We show here that the currently available

reference genomes, however sparse, are phylogenetically broad enough to make

significant contributions to our understanding of genome evolution in vertebrates and

additional data will only serve to deepen this understanding. We caution that high-

quality genomes without high-quality annotations are limited in their utility to the broader

field, but this is an area that needs additional attention from both funding sources,

program developers, and empiricists; we see potential for cloud computing as a

resource for this work. Current work in sex chromosome and microchromosome

evolution (among others) stand to make great strides in coming years as high-quality

genomic data become more prevalent in additional taxa. Thus, squamate genomics as

a field has blossomed in recent years and this presents a bright outlook for the future of

genomics of these often overlooked, yet speciose and charismatic animals.

Methods

We compiled a near-complete list of all available lepidosaur genome assemblies from GenBank (NCBI), Ensembl (EVI), DNA Zoo (Dudchenko et al. 2017), National Genomics Data Center (CNCB), and individual paper data repositories (e.g. Figshare and GigaDB). We noted the disclosed technologies used to acquire the assembly (from either the database, when available, or the primary article) whether each assembly had an accompanying annotation file available from the download source. We then downloaded each assembly to confirm its existence/availability and calculated basic statistics on each using assembly-stats [v1.0.1] (https://github.com/sanger-pathogens/assembly-stats). We then conducted a literature search to identify the sex determining system of each species (if known), the linkage group (in *Gallus gallus*), the sex-determining region location (if known), and putative sex determining genes. We used this information to assess whether each assembly was considered to be "chromosome-level" or not (in squamates generally, if the scaffold L50 <8 but varies by species) and analyzed this subset using BUSCO [v5.1.2] (Simão et al. 2015) on the gVolante web server [v2.0.0] (Nishimura et al. 2017). Further, for the four genome assemblies of species that were not annotated, which also had an outdated assembly that was annotated, we used Liftoff [v1.6.3] (Shumate and Salzberg, 2021) and uploaded them to an archived repository for public availability and reuse (https://doi.org/10.6084/m9.figshare.20201099). All counts of number of species per clade were collected from Reptile Database (Uetz et al. 2022)

To better understand the genomic composition of reptiles, we used the aforementioned information to best inform which taxa would be the most informative to three downstream analyses. (1) We compiled summary information for representative genomes across amniotes with per chromosome information for number of genes and GC content from NCBI (a lizard, *Podarcis muralis*; two birds, *Gallus gallus* and *Taeniopygia guttata*; a turtle, *Mauremys mutica*; and human, *Homo sapiens*), with a few exceptions that were not directly available through NCBI including gene numbers for other representative squamates: *Shinisaurus crocodilurus* (Liftoff), *Sphaerodactylus townsendi*, and *Naja naja*. In addition, we calculated gene number and GC content for *Alligator sinensis* (Liftoff) and GC content only for *Cygnus atratus*. For each species, we

conducted Bayesian and frequentist correlation analyses using JASP (JASP Team,

2022) between each of three variables: GC content, gene number, and chromosome

size (the latter two normalized by dividing each by the mean value for each). (2) From a

representative number of taxa with chromosome-level reference genomes, we

generated a synteny map across squamates, rooted with chicken. We generated

corresponding peptide files from each genome using gffread (Pertea and Pertea, 2020)

and calculated the synteny map using Genespace (Lovell et al. 2022). (3) We then

collated chromatin-contact information from the DNAZoo for a bird (*Cygnus atratus*) and

squamate (*Salvator merianae*) and used the recently published contact map from the

leopard gecko, *Eublepharis macularius* (Pinto et al. 2023). For *Podarcis muralis*, we had

to generate a contact map (that had not been previously published) from the published

genome data (PRJNA515813; Andrade et al. 2019). We used Juicer and 3D-DNA to

generate the contact map (Dudchenko et al. 2017) and generated images (Figure 4) at

a standardized resolution using Juicebox Assembly Tools [v1.11.08] (Durand et al.

2016).

Data Availability

As indicated above, our cutoff for finding and including additional genomes to the dataset for this paper was July 12th, 2022 and the information used for this study is summarized in Appendix I. However, we have continued aggregating genomes to the summary table beyond this date and have appended them to a live document available here (https://drpintothe2nd.weebly.com/squamates.html). We will continue updating this spreadsheet for the foreseeable future, likely either until genomes become too numerous to keep up with or a better resource is made available. Please feel free to reach out to B.J.P. via email to incorporate additional resources or make corrections to the list.

Acknowledgements

The authors would like to dedicate this manuscript to Kari Pinto (1960-2022), an avid naturalist, who passed away suddenly during the preparation of this manuscript. This publication was supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number R35GM124827 to MAW and NSF IOS 2151318 to TG.

References

Abe, T., Kaneko, M., & Kiyonari, H. (2023). A reverse genetic approach in geckos with the CRISPR/Cas9 system by oocyte microinjection. *Developmental Biology*. 497:26-32.

Alföldi, J., Di Palma, F., Grabherr, M., Williams, C., Kong, L., Mauceli, E., ... Lindblad-Toh, K. (2011). The genome of the green anole lizard and a comparative analysis with birds and mammals. *Nature*. 477(7366):587-591.

Andrade, P., Pinho, C., Pérez i de Lanuza, G., Afonso, S., Brejcha, J., Rubin, C.-J., Wallerman, O., Pereira, P., Sabatino, S. J., Bellati, A., Pellitteri-Rosa, D., Bosakova, Z., Bunikis, I., Carretero, M. A., Feiner, N., Marsik, P., Paupério, F., Salvi, D., Soler, L., … Carneiro, M. (2019). Regulatory changes in pterin and carotenoid genes underlie balanced color polymorphisms in the wall lizard. *PNAS*. 116(12):5633-5642.

Bachtrog D, Mank JE, Peichel CL, Kirkpatrick M, Otto SP, Ashman TL, Hahn MW, Kitano J, Mayrose I, Ming R, Perrin N, Ross L, Valenzuela N, Vamosi JC. 2014. Sex determination: Why so many ways of doing it? *PLoS Biol*. 12: e1001899.

Blount, Z. D., Lenski, R. E., & Losos, J. B. (2018). Contingency and determinism in evolution: Replaying life's tape. *Science*. 362(6415).

Bravo, G. A., Schmitt, C. J., Edwards, S. V. (2021). What have we learned from the first 500 avian genomes? *Annu Rev Ecol Evol Syst*. 52(1):611-639.

Bernardi, G., Bernardi, G. (1986). Compositional constraints and genome evolution. *Journal of Molecular Evolution*. 24(1–2):1-11.

Boring, A. M. (1923). Notes by N. M. Stevens on chromosomes of the domestic chicken. Science. 58:73-74.

Bull JJ. 1983 Evolution of sex determining mechanisms. Menlo Park, CA: Benjamin Cummings Publishing Company, Inc

Bull J, Charnov E. 1985. On irreversible evolution. Evolution 39, 1149–1155.

Carey, S. B., Lovell, J. T., Jenkins, J., Leebens-Mack, J., Schmutz, J., Wilson, M. A., & Harkess, A. (2022). Representing sex chromosomes in genome assemblies. *Cell Genomics*, 100132.

Cantarel, B. L., Korf, I., Robb, S. M., Parra, G., Ross, E., Moore, B., ... Yandell, M. (2008). MAKER: an easy-to-use annotation pipeline designed for emerging model organism genomes. *Genome Research*. 18(1):188-196.

Card, D. C., Jennings, W. B., & Edwards, S. V. (2023). Genome Evolution and the Future of Phylogenomics of Non-Avian Reptiles. *Animals*. 13(3):471.

Carey, S. B., Lovell, J. T., Jenkins, J., Leebens-Mack, J., Schmutz, J., Wilson, M. A., & Harkess, A. (2022). Representing sex chromosomes in genome assemblies. *Cell Genomics*. 2(5).

Cheng, H., Concepcion, G. T., Feng, X., Zhang, H., Li, H. (2021). Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. *Nature Methods*. 18(2):170–175.

Cheng, H., Jarvis, E. D., Fedrigo, O., Koepfli, K. P., Urban, L., Gemmell, N. J., Li, H. (2022). Haplotype-resolved assembly of diploid genomes without parental data. *Nature Biotechnology*. 1-4.

Dahn, H. A., Mountcastle, J., Balacco, J., Winkler, S., Bista, I., Schmitt, A. D., Pettersson, O. V., Formenti, G., Oliver, K., Smith, M., Tan, W., Kraus, A., Mac, S., Komoroske, L. M., Lama, T., Crawford, A. J., Murphy, R. W., Brown, S., Scott, A. F., …

Fedrigo, O. (2022). Benchmarking ultra-high molecular weight DNA preservation methods for long-read and long-range sequencing. *GigaScience*. 11.

Deakin, J. E., Edwards, M. J., Patel, H., O'Meally, D., Lian, J., Stenhouse, R., Ryan, S., Livernois, A.M., Azad, B., Hollelely, C.E., et al. (2016). Anchoring genome sequence to chromosomes of the central bearded dragon (Pogona vitticeps) enables reconstruction of ancestral squamate macrochromosomes and identifies sequence content of the Z chromosome. *BMC Genomics*. 17, 1-15.

Deakin, J. E., Ezaz, T. (2019). Understanding the Evolution of Reptile Chromosomes through Applications of Combined Cytogenetics and Genomics Approaches. Cytogenetic and Genome Research, 157(1–2), 7–20.

Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC, Shamim MS, et al. (2017). *De Novo* assembly of the *Aedes aegypti* genome using Hi-C yields chromosome-length scaffolds. *Science*. 356:92-95.

Durand, N.C., Shamim, M.S., Machol, I., Rao, S.S., Huntley, M.H., Lander, E.S., Aiden, E.L. (2016). Juicer provides a one-click system for analyzing loop-resolution Hi-C experiments. *Cell Syst*. 3:95-98.

Ezaz, T., Sarre, S.D., O'Meally, D., Graves, J.A.M., and Georges, A. (2009). Sex chromosome evolution in lizards: Independent origins and rapid transitions. *Cytogenet Genome Res*. 127:249–260.

Gamble, T. (2010). A review of sex determining mechanisms in geckos (Gekkota: Squamata). *Sexual Development*. 4(1–2):88-103.

Gamble, T., Greenbaum, E., Jackman, T. R., Russell, A. P., & Bauer, A. M. (2012). Repeated origin and loss of adhesive toepads in geckos. *PloS One*. 7(6): e39429.

Gamble, T., Coryell, J., Ezaz, T., Lynch, J., Scantlebury, D. P., Zarkower, D. (2015a). Restriction site-associated DNA sequencing (RAD-seq) reveals an extraordinary number of transitions among gecko sex-determining systems. *Molecular Biology and Evolution*. 32(5):1296-1309.

Gamble, T., Greenbaum, E., Jackman, T. R., & Bauer, A. M. (2015). Into the light: diurnality has evolved multiple times in geckos. *Biological Journal of the Linnean Society*. 115(4):896-910.

Gamble, T., Castoe, T. A., Nielsen, S. V., Banks, J. L., Card, D. C., Schield, D. R., Schuett, G. W., Booth, W. (2017). The discovery of XY sex chromosomes in a *Boa* and *Python*. *Current Biology*. 27(14):2148-2153.

Gamble, T., McKenna, E., Meyer, W., Nielsen, S. V., Pinto, B. J., Scantlebury, D. P., Higham, T. E. (2018). XX/XY sex chromosomes in the South American dwarf gecko (*Gonatodes humeralis*). *Journal of Heredity*. 109(4):462-468.

Gamble, T. 2019. Duplications in corneous beta protein genes and the evolution of gecko adhesion. *Integrative and Comparative Biology*. 59:193–202.

Genome 10K Community of Scientists. (2009). Genome 10K: a proposal to obtain whole-genome sequence for 10 000 vertebrate species. *Journal of Heredity*, *100*(6), 659-674.

Gilbert, C., Meik, J. M., Dashevsky, D., Card, D. C., Castoe, T. A., & Schaack, S. (2014). Endogenous hepadnaviruses, bornaviruses and circoviruses in snakes. *Proceedings of the Royal Society B: Biological Sciences*. 281(1791):20141122.

Gu, L., & Walters, J. R. (2017). Evolution of Sex Chromosome Dosage Compensation in Animals: A Beautiful Theory, Undermined by Facts and Bedeviled by Details. Genome Biology and Evolution. 9(9):2461–2476.

Fillon, V. (1998). The chicken as a model to study microchromosomes in birds: a review. *Genet Sel Evol*. 30(3):209-219.

Fullerton, S.M., Bernardo Carvalho, A., Clark, A.G. (2001). Local rates of recombination are positively correlated with GC content in the human genome. *Mol Biol Evol*. 18:1139-1142.

Hance, R. T. (1924). The somatic chromosomes of the chick and their possible sex relations. *Science*. 59:424-425.

Hoff, K.J., Lomsadze, A., Borodovsky, M. and Stanke, M. (2019). Whole-Genome Annotation with BRAKER. *Methods Mol Biol*. 1962:65-95.

Hofstra, B., Kulkarni, V. V., Munoz-Najar Galvez, S., He, B., Jurafsky, D., McFarland, D. A. (2020). The diversity–innovation paradox in science. *PNAS*. 117(17):9284-9291.

Hotaling, S., Kelley, J. L., & Frandsen, P. B. (2021). Toward a genome sequence for every animal: Where are we now?. *PNAS*. 118(52):e2109019118.

JASP Team. (2022). JASP (Version 0.16.2) https://jasp-stats.org/

Kamiya, T., Kai, W., Tasumi, S., Oka, A., Matsunaga, T., Mizuno, N., Fujita, M., Suetake, H., Suzuki, S., Hosoya, S., Tohari, S., Brenner, S., Miyadai, T., Venkatesh, B., Suzuki, Y., Kikuchi, K. (2012). A Trans-Species Missense SNP in Amhr2 Is Associated with Sex Determination in the Tiger Pufferfish, *Takifugu rubripes* (Fugu). *PLoS Genetics*. 8(7):e1002798.

Karawita, A. C., Cheng, Y., Chew, K. Y., Challgula, A., Kraus, R., Mueller, R. C., Tong, M. Z. W., Hulme, K. D., Beielefeldt-Ohmann, H., Steele, L. E., Wu, M., Sng, J., Noye, E., Bruxner, T. J., Au, G. G., Lowther, S., Blommaert, J., Suh, A., McCauley, A. J., … Short, K. R. (2022). The swan genome and transcriptome: it's not all black and white. *BioRxiv*.

Keating, S. E., Griffing, A. H., Nielsen, S. V., Scantlebury, D. P., & Gamble, T. (2020). Conserved ZZ/ZW sex chromosomes in Caribbean croaking geckos (*Aristelliger*: Sphaerodactylidae). Journal of Evolutionary Biology, 33(9), 1316–1326.

Keating SE. (2022). Evolution of sex chromosomes in geckos (Reptilia: Gekkota). Unpublished dissertation. Marquette University.

Keating, S. E., Greenbaum, E., Johnson, J. D., & Gamble, T. (2022). Identification of a cis-sex chromosome transition in banded geckos (Coleonyx, Eublepharidae, Gekkota). Journal of Evolutionary Biology, 35(12), 1675-1682.

Kostmann A, Kratochvíl L, Rovatsos M. (2021). Poorly differentiated XX/XY sex chromosomes are widely shared across skink radiation. *Proc Roy Soc B*. 288:20202139.

Kumar S, Stecher G, Suleski M, Hedges SB. 2017. TimeTree: a resource for timelines, timetrees, and divergence times. *Mol Biol Evol*. 34:1812-1819.

Lang, D., Zhang, S., Ren, P., Liang, F., Sun, Z., Meng, G., ... & Liu, S. (2020). Comparison of the two up-to-date sequencing technologies for genome assembly: HiFi reads of Pacific Biosciences Sequel II system and ultralong reads of Oxford Nanopore. *Gigascience*. 9(12):giaa123.

Li, R., Li, Y., Zheng, H., Luo, R., Zhu, H., Li, Q., Qian, W., Ren, Y., Tian, G., Li,
J., Zhou, G., Zhu, X., Wu, H., Qin, J., Jin, X., Li, D., Cao, H., Hu, X., Blanche, H., …
Wang, J. (2010). Building the sequence map of the human pan-genome. *Nature*
Biotechnology. 28(1):57-63.
Losos J, Braun E, Brown D, Clifton S, Edwards S, Gibson-Brown J, ... Warren W.
2005. Proposal to sequence the first reptilian genome: the green anole lizard, *Anolis*
carolinensis. *NHGRI White Paper*.
Lovell, J. T., Sreedasyam, A., Schranz, M. E., Wilson, M., Carlson, J. W.,
Harkess, A., Emms, D., Goodstein, D. M., Schmutz, J. (2022). GENESPACE tracks
regions of interest and gene copy number variation across multiple genomes. *ELife*. 11.
Marin, R., Cortez, D., Lamanna, F., Pradeepa, M. M., Leushkin, E., Julien, P.,
Liechti, A., Halbert, J., Brüning, T., Mössinger, K., Trefzer, T., Conrad, C., Kerver, H. N.,
Wade, J., Tschopp, P., & Kaessmann, H. (2017). Convergent origination of a
Drosophila-like dosage compensation mechanism in a reptile lineage. Genome
Research, 27(12), 1974–1987.
Meiri, S., Bauer, A. M., Allison, A., Castro-Herrera, F., Chirio, L., Colli, G., ... &
Roll, U. (2018). Extinct, obscure or imaginary: the lizard species with the smallest
ranges. *Diversity and Distributions*, 24(2), 262-273.
Nakamoto, M., Uchino, T., Koshimizu, E., Kuchiishi, Y., Sekiguchi, R., Wang, L.,
Sudo, R., Endo, M., Guiguen, Y., Schartl, M., Postlethwait, J. H., Sakamoto, T. (2021).
A Y-linked anti-Müllerian hormone type-II receptor is the sex-determining gene in ayu,
Plecoglossus altivelis. *PLoS Genetics*. 17(8):e1009705.
Neemuchwala, S., Johnson, N.A., Pfeiffer, J.M., Lopes-Lima, M., Gomes-dos-
Santos, A., Froufe, E., Hillis, D.M., Smith, C.H. 2023. Coevolution with host fishes
shapes parasitic life histories in a group of freshwater mussels (Unionidae: Quadrulini).
Bulletin of the Society of Systematic Biologists. *In press*.
Newcomer, E. H. (1957). The mitotic chromosomes of the domestic fowl. *Journal*
of Heredity. 48(5):227-234.
Nielsen, S. V., Banks, J. L., Diaz, R. E., Trainor, P. A., & Gamble, T. (2018).
Dynamic sex chromosomes in Old World chameleons (Squamata: Chamaeleonidae).
Journal of Evolutionary Biology, 31(4), 484–490.
Nielsen, S. V., Guzmán-Méndez, I. A., Gamble, T., Blumer, M., Pinto, B. J.,
Kratochvíl, L., & Rovatsos, M. (2019). Escaping the evolutionary trap? Sex chromosome
turnover in basilisks and related lizards (Corytophanidae: Squamata). *Biology letters*,
15(10), 20190498.
Nielsen, S. V., Pinto, B. J., Guzmán-Méndez, I. A., & Gamble, T. (2020). First
Report of Sex Chromosomes in Night Lizards (Scincoidea: Xantusiidae). Journal of
Heredity, 111(3), 307–313.
Nishimura O, Hara Y, Kuraku S. (2017). gVolante for standardizing completeness
assessment of genome and transcriptome assemblies. *Bioinformatics*. 33:3635-3637.
Nurk, S., Koren, S., Rhie, A., Rautiainen, M., Bzikadze, A. V., Mikheenko, A.,
Vollger, M. R., Altemose, N., Uralsky, L., Gershman, A., Aganezov, S., Hoyt, S. J.,
Diekhans, M., Logsdon, G. A., Alonge, M., Antonarakis, S. E., Borchers, M., Bouffard,
G. G., Brooks, S. Y., … Phillippy, A. M. (2022). The complete sequence of a human
genome. *Science*. 376(6588):44–53.

O'Connor RE, Kiazim L, Skinner B, Fonseka G, Joseph S, Jennings R, Larkin
DM, Griffin DK. (2019). Patterns of microchromosome organization remain highly
conserved throughout avian evolution. *Chromosoma*. 128:21–29.
Ohno, S. (1961). Sex chromosome and microchromosomes of *Gallus*
domesticus. *Chromosoma*. 11:484-498.
Ohno S. 1967. Sex chromosomes and sex-linked genes. Berlin, Germany:
Springer Verlag.
Olmo E. (1986). A. Reptilia. In: John B (ed) Animal cytogenetics 4 Chordata 3.
Gebriider Borntraeger, Berlin-Stuttgart.
Olmo E, Odierna G, Capriglione T, Cardone A. (1990). DNA and chromosome
evolution in lacertid lizards. In: Olmo E, editor Cytogenetics of Amphibians and Reptiles.
Berlin, Germany: Birkhauser Verlag. p. 181–204.
Olney, K. C., Brotman, S. M., Andrews, J. P., Valverde-Vesling, V. A., & Wilson,
M. A. (2020). Reference genome and transcriptome informed by the sex chromosome
complement of the sample increase ability to detect sex differences in gene expression
from RNA-Seq data. *Biology of Sex Differences*. 11(1).
Palmer B. (2020). Funannotate v1.8.1: Eukaryotic genome annotation pipeline.
https://doi.org/10.5281/zenodo.4054262
Pan, Q., Herpin, A., & Guiguen, Y. (2022). Inactivation of the Anti-Müllerian
Hormone Receptor Type 2 (*amhrII*) Gene in Northern Pike (*Esox lucius*) Results in
Male-To-Female Sex Reversal. *Sexual Development*. 1-6.
Pensabene, E., Yurchenko, A., Kratochvíl, L., & Rovatsos, M. (2023).
Madagascar Leaf-Tail Geckos (Uroplatus spp.) Share Independently Evolved
Differentiated ZZ/ZW Sex Chromosomes. *Cells*, 12(2), 260.
Peona, V., Blom, M. P. K., Xu, L., Burri, R., Sullivan, S., Bunikis, I., Liachko, I.,
Haryoko, T., Jønsson, K. A., Zhou, Q., Irestedt, M., & Suh, A. (2020). Identifying the
causes and consequences of assembly gaps using a multiplatform genome assembly of
a bird-of-paradise. *Molecular Ecology Resources*. 21(1):263-286.
Perry, B. W., Schield, D. R., Adams, R. H., Castoe, T. A. (2021).
Microchromosomes exhibit distinct features of vertebrate chromosome structure and
function with underappreciated ramifications for genome evolution. *Molecular Biology*
and Evolution. 38(3):904-910.
Pertea, G., Pertea, M. (2020). GFF Utilities: GffRead and GffCompare.
F1000Research. 9:304.
Pinto, B. J., Nielsen, S. V., & Gamble, T. (2019a). Transcriptomic data support a
nocturnal bottleneck in the ancestor of gecko lizards. *Molecular Phylogenetics and*
Evolution. 141:106639.
Pinto, B. J., Card, D. C., Castoe, T. A., Diaz, R. E., Nielsen, S. V., Trainor, P. A.,
Gamble, T. (2019b). The transcriptome of the veiled chameleon (*Chamaeleo*
calyptratus): A resource for studying the evolution and development of vertebrates.
Developmental Dynamics. 248(8):702–708.
Pinto, B. J., Nielsen, S. V., & Gamble, T. (2019c). Transcriptomic data support a
nocturnal bottleneck in the ancestor of gecko lizards. *Molecular Phylogenetics and*
Evolution. 141:106639.
Pinto, B. J., Keating, S. E., Nielsen, S. V., Scantlebury, D. P., Daza, J. D.,
Gamble, T. (2022). Chromosome-level genome assembly reveals dynamic sex

chromosomes in neotropical leaf-litter geckos (Sphaerodactylidae: *Sphaerodactylus*). *Journal of Heredity*. 113(3):272–287.

Pinto BJ, Gamble T, Smith CH, Keating SE, Havird J, Chiari Y. (2023a). The revised leopard gecko (*Eublepharis macularius*) reference genome provides insight into the considerations of genome phasing and assembly. *Journal of Heredity. In press.*

Pinto BJ, O'Connor B, Schatz MC, Zarate S, Wilson MA. (2023b). Concerning the eXclusion in human genomics: The choice of sex chromosome representation in the human genome drastically affects number of identified variants. *BioRxiv*.

Pokorná, M., & Kratochvíl, L. (2009). Phylogeny of sex-determining mechanisms in squamate reptiles: are sex chromosomes an evolutionary trap? *Zoological Journal of the Linnean Society*, 156(1), 168–183.

Randhawa, S. S., & Pawar, R. (2021). Fish genomes: Sequencing trends, taxonomy and influence of taxonomy on genome attributes. *Journal of Applied Ichthyology*. 37(4):553-562.

Rasys, A. M., Park, S., Ball, R. E., Alcala, A. J., Lauderdale, J. D., Menke, D. B. (2019). CRISPR-Cas9 gene editing in lizards through microinjection of unfertilized oocytes. *Cell Reports*. 28(9):2288-2292.

Revell LJ, 2012. Phytools: an R package for phylogenetic comparative biology (and other things). *Methods Ecol Evol*. **3:**217–223.

Rhoads A, Au KF. (2015). PacBio Sequencing and Its Applications. *Genom Proteom Bioinform*. 13(5):278–289.

Rovatsos, M., Pokorná, M., Altmanová, M., & Kratochvíl, L. (2014). Cretaceous park of sex determination: sex chromosomes are conserved across iguanas. Biology Letters, 10(3), 20131093.

Rovatsos, M., Vukić, J., Lymberakis, P., & Kratochvíl, L. (2015). Evolutionary stability of sex chromosomes in snakes. Proceedings of the Royal Society B: Biological Sciences, 282(1821), 20151992.

Rovatsos, M., Vukić, J., Mrugała, A., Suwala, G., Lymberakis, P., & Kratochvíl, L. (2019a). Little evidence for switches to environmental sex determination and turnover of sex chromosomes in lacertid lizards. Scientific Reports, 9(1).

Rovatsos M, Rehák I, Velenský P, Kratochvíl L. (2019b). Shared Ancient Sex Chromosomes in Varanids, Beaded Lizards, and Alligator Lizards. *Mol Biol Evol*. **36:** 1113-20.

Rovatsos, M., Farkačová, K., Altmanová, M., Johnson Pokorná, M., & Kratochvíl, L. (2019c). The rise and fall of differentiated sex chromosomes in geckos. Molecular Ecology, 28(12), 3042–3052.

Rovatsos, M., Gamble, T., Nielsen, S. V., Georges, A., Ezaz, T., & Kratochvíl, L. (2021). Do male and female heterogamety really differ in expression regulation? Lack of global dosage balance in pygopodid geckos. Philosophical Transactions of the Royal Society B: Biological Sciences, 376(1833), 20200102.

Rovatsos, M., Galoyan, E., Spangenberg, V., Vassilieva, A., & Kratochvíl, L. (2022). XX/XY sex chromosomes in a blind lizard (Dibamidae): Towards understanding the evolution of sex determination in squamates. *Journal of Evolutionary Biology*, 35(12), 1791–1796.

Rupp, S. M., Webster, T. H., Olney, K. C., Hutchins, E. D., Kusumi, K., & Wilson Sayres, M. A. (2017). Evolution of dosage compensation in *Anolis carolinensis*, a reptile

with XX/XY chromosomal sex determination. *Genome Biology and Evolution*. 9(1):231-240.

Sackton, T. B., & Clark, N. (2019). Convergent evolution in the genomics era: new insights and directions. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 374(1777):20190102.

Schield, D. R., Card, D. C., Hales, N. R., Perry, B. W., Pasquesi, G. M., Blackmon, H., Adams, R. H., Corbin, A. B., Smith, C. F., Ramesh, B., Demuth, J. P., Betrán, E., Tollis, M., Meik, J. M., Mackessy, S. P., & Castoe, T. A. (2019). The origins and evolution of chromosomes, dosage compensation, and mechanisms underlying venom regulation in snakes. Genome Research, 29(4), 590–601.

Shumate A, Salzberg SL. 2021. Liftoff: Accurate mapping of gene annotations. *Bioinformatics*. **12:**1639-1643.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*. **31:**3210-3212.

Smith, C. H. (2021). A High-Quality Reference Genome for a Parasitic Bivalve with Doubly Uniparental Inheritance (Bivalvia: Unionida). *Genome Biology and Evolution*. 13(3).

Smith, C. H., Pfeiffer, J. M., Johnson, N. A. (2020). Comparative phylogenomics reveal complex evolution of life history strategies in a clade of bivalves with parasitic larvae (Bivalvia: Unionoida: Ambleminae). *Cladistics* 36(5):505-520.

Srikulnath K. 2013. The dynamics of chromosome evolution in reptiles. *Thai J Genet*. 6:77–79.

Srikulnath, K., Ahmad, S. F., Singchat, W., Panthum, T. (2021). Why do some vertebrates have microchromosomes?. *Cells*. 10(9):2182.

Sun, Y., Shang, L., Zhu, Q. H., Fan, L., Guo, L. (2021). Twenty years of plant genome sequencing: Achievements and challenges. Trends in Plant Science.

Terao, M., Ogawa, Y., Takada, S., Kajitani, R., Okuno, M., Mochimaru, Y., Matsuoka, K., Itoh, T., Toyoda, A., Kono, T., Jogahara, T., Mizushima, S., & Kuroiwa, A. (2022). Turnover of mammal sex chromosomes in the *Sry*-deficient Amami spiny rat is due to male-specific upregulation of *Sox9*. Proceedings of the National Academy of Sciences, 119(49).

Uetz P, Freed P, Aguilar R, Hošek J. (eds.) 2022. The Reptile Database, http://www.reptile-database.org, accessed August 26, 2022.

Vicoso, B., Bachtrog, D. (2009). Progress and prospects toward our understanding of the evolution of dosage compensation. *Chromosome Research*. 17(5).

Vicoso, B., Emerson, J. J., Zektser, Y., Mahajan, S., & Bachtrog, D. (2013). Comparative Sex Chromosome Genomics in Snakes: Differentiation, Evolutionary Strata, and Lack of Global Dosage Compensation. PLoS Biology, 11(8), e1001643.

Vollger, M. R., Logsdon, G. A., Audano, P. A., Sulovari, A., Porubsky, D., Peluso, P., ... & Eichler, E. E. (2020). Improved assembly and variant detection of a haploid human genome using single-molecule, high-fidelity long reads. *Annals of Human Genetics*. 84(2):125-140.

Webster, T. H., Couse, M., Grande, B. M., Karlins, E., Phung, T. N., Richmond, P. A., ... Wilson, M. A. (2019). Identifying, understanding, and correcting technical

1008 artifacts on the sex chromosomes in next-generation sequencing data. *Gigascience*.
1009 8(7):giz074.
1010   Webster, T.H., Vannan, A., Pinto, B.J., Denbrock, G., Morales, M., Dolby, G.A.,
1011 Fiddes, I.T., DeNardo, D.F., Wilson, M.A. (2023). Incomplete dosage compensation and
1012 lack of dosage balance in the ZZ/ZW Gila monster (*Heloderma suspectum*) revealed by
1013 de novo genome assembly. *In review*.
1014   Wetterstrand, K.A. (2021) DNA Sequencing Costs: Data from the NHGRI
1015 Genome Sequencing Program (GSP). Available at:
1016 www.genome.gov/sequencingcostsdata. Accessed 10-10-2022.
1017   Yamashina, M. Y. (1944). Karyotype Studies in Birds I. Comparative morphology
1018 of chromosomes in seventeen races of Domestic fowl. *Cytologia*. 13(3-4):270-296.

Figure 1: Chronological breakdown of genome assemblies published per-year and proportion of the assemblies that are chromosome-level (top panel) or annotated (bottom panel). Importantly, not all chromosome-level genomes are annotated and most chromosome-level assemblies that improve a previously annotated assembly do not publish updated annotations.

Figure 2: Breakdown of total number of published genome assemblies (bar graph) per phylogenetic group (family or superfamily) co-plotted with number of species in each clade (branch colors and parenthetical numbers). Phylogeny from TimeTree using a representative species from each clade (Kumar et al. 2017), species counts from the Reptile Database (Uetz et al. 2022), and plotted using phytools (Revell, 2012).

Species per clade

1   3280

Sphenodon (1)
Bipedidae (3)
Amphisbaenidae (183)
Rhineuridae (1)
Lacertidae (362)
Gymnophthalmidae (275)
Teiidae (172)
Agamidae (556)
Chamaeleonidae (222)
Iguanidae (44)
Phrynosomatidae (170)
Crotaphytidae (12)
Dactyloidae (437)
Corytophanidae (11)
Leiocephalidae (29)
Tropiduridae (142)
Liolaemidae (338)
Opluridae (8)
Polychrotidae (8)
Xenosauridae (13)
Helodermatidae (5)
Anguidae (86)
Shinisauridae (1)
Lanthanotidae (1)
Varanidae (83)
Anomalepididae (21)
Aniliidae (1)
Viperidae (374)
Homalopsidae (56)
Colubroidea (3,279)
Elapidae (389)
Uropeltoidea (70)
Pythonidae (38)
Loxocemidae (1)
Boidae (67)
Leptotyphlopidae (141)
Typhlopidae (275)
Xantusiidae (37)
Gerrhosauridae (38)
Cordylidae (68)
Scincidae (1,744)
Eublepharidae (44)
Pygopodidae (46)
Carphodactylidae (32)
Diplodactylidae (161)
Sphaerodactylidae (229)
Gekkonidae (1,481)
Phyllodactylidae (158)
Dibamidae (25)

Laterata

Iguania

Anguimorpha

Snakes

Scincomorpha

Geckos

0   2   4   6   8   10   12   14

Genome Assemblies

Figure 3: Time-calibrated phylogeny of squamate reptiles pruned to include only species with high-quality genome assemblies (rooted with chicken, *Gallus gallus*). Branches leading to major phylogenetic groups labeled, those with multiple taxa are highlighted. Phylogeny obtained from TimeTree (Kumar et al. 2017) and plotted using GeneSpace (Lovell et al. 2022) and FigTree [v1.4.4]. It's apparent that microchromosomes are homologous in squamates that possess them (*Salvator*, *Naja*, and *Shinisaurus*), while different linkage group fusions have led to their loss in taxa that lack them (*Sphaerodactylus* and *Podarcis*).

Gallus gallus
Paroedura picta
Sphaerodactylus townsendi
Eublepharis macularius
Podarcis muralis
Lacerta agilis
Zootoca vivipara
Salvator merianae
Python bivittatus
Crotalus viridis
Hydrophis curtus
Hydrophis cyanocinctus
Thamnophis elegans
Pantherophis guttatus
Pituophis catenifer
Arizona elegans
Naja naja
Anolis sagrei
Anolis carolinensis
Sceloporus tristichus
Sceloporus undulatus
Phrynosoma platyrhinos
Pogona vitticeps
Shinisaurus crocodilurus

Geckos

Laterata

Snakes

Iguania

Anguimorpha

300 250 200 150 100 50 0

Chromosomes scaled by gene rank order

5k genes

Figure 4: HiC contact maps for representative reptile taxa demonstrating the presence of microchromosomes in (A) birds and (B) teiids, or absence of microchromosomes in (C) geckos and (D) lacertids. Microchromosomes denoted with black bars to the bottom and right of the respective contact map. Annotation of microchromosomal organization denoted via top-right bracket in (A) and (B).

Supplemental Materials:

Supplemental Figure 1: Bayesian analyses of chromosomal make-up across vertebrate taxa. (A) Gecko: *Sphaerodactylus townsendi*, (B) Wall lizard: *Podarcis muralis*, (C) Chinese alligator: *Alligator sinensis*, (D) Human: *Homo sapiens*, (E) Crocodile lizard: *Shinisaurus crocodilurus*, (F) Argentine black and white tegu: *Salvator merianae*, (G) Indian cobra: *Naja naja*, (H) Yellowpond turtle: *Mauremys mutica*, (I) Chicken: *Gallus gallus*, (J) Zebra finch: *Taeniopygia guttata*, (K) Black swan: *Cygnus atratus*, (L) Southern platyfish: *Xiphophorus maculatus*.

Results

Bayesian Correlation

Bayesian Pearson Correlations

			n	Pearson's r	BF₁₀
norm_size	-	norm_gene	17	0.865***	2879.882
norm_size	-	gene_density	17	-0.477	1.702
norm_size	-	GC content (%)	17	-0.781***	161.651
norm_gene	-	gene_density	17	-0.024	0.301
norm_gene	-	GC content (%)	17	-0.476	1.688
gene_density	-	GC content (%)	17	0.721**	40.141

* BF₁₀ > 10, ** BF₁₀ > 30, *** BF₁₀ > 100

Bayesian Correlation Matrix Plot

B Results

Bayesian Correlation

Bayesian Pearson Correlations

			n	Pearson's r	BF$_{10}$
norm_size	-	norm_gene	19	0.699**	48.582
norm_size	-	gene_density	19	-0.438	1.464
norm_size	-	GC content (%)	19	-0.727**	90.447
norm_gene	-	gene_density	19	0.277	0.525
norm_gene	-	GC content (%)	19	-0.243	0.454
gene_density	-	GC content (%)	19	0.734***	105.664

* BF$_{10}$ > 10, ** BF$_{10}$ > 30, *** BF$_{10}$ > 100

Bayesian Correlation Matrix Plot

Bayesian Correlation

Bayesian Pearson Correlations

			Pearson's r	BF$_{10}$
norm_size	-	norm_gene	0.936***	111257.209
norm_size	-	gene_density	-0.538	2.591
norm_size	-	GC content (%)	-0.720*	28.773
norm_gene	-	gene_density	-0.291	0.536
norm_gene	-	GC content (%)	-0.600	4.958
gene_density	-	GC content (%)	0.850***	847.880

* BF$_{10}$ > 10, ** BF$_{10}$ > 30, *** BF$_{10}$ > 100

Bayesian Correlation Matrix Plot

D

Results

Bayesian Correlation

Bayesian Pearson Correlations

			n	Pearson's r	BF$_{10}$
norm_size	-	norm_gene	23	0.857***	86563.421
norm_size	-	gene_density	23	-0.443	2.134
norm_size	-	GC content (%)	23	-0.598*	18.642
norm_gene	-	gene_density	23	0.031	0.261
norm_gene	-	GC content (%)	23	-0.183	0.359
gene_density	-	GC content (%)	23	0.879***	402621.999

* BF$_{10}$ > 10, ** BF$_{10}$ > 30, *** BF$_{10}$ > 100

Bayesian Correlation Matrix Plot

Results

Bayesian Independent Samples T-Test

Bayesian Mann-Whitney U Test

	BF$_{-0}$	W	Rhat
gene_density	0.508	17.000	1.006
GC content (%)	0.703	15.000	1.000

Note. For all tests, the alternative hypothesis specifies that the location of group *Macro* is smaller than the location of group *Micro*.
Note. Result based on data augmentation algorithm with 5 chains of 1000 iterations.

Descriptives

Raincloud Plots

Bayesian Correlation

Bayesian Pearson Correlations

			n	Pearson's r	BF$_{10}$
norm_size	-	norm_gene	16	0.950***	501947.382
norm_size	-	gene_density	16	-0.294	0.543
norm_size	-	GC content (%)	16	-0.687*	16.134
norm_gene	-	gene_density	16	-0.058	0.315
norm_gene	-	GC content (%)	16	-0.572	3.635
gene_density	-	GC content (%)	16	0.764**	69.420

* BF$_{10}$ > 10, ** BF$_{10}$ > 30, *** BF$_{10}$ > 100

Bayesian Correlation Matrix Plot

Bayesian Independent Samples T-Test

	BF−0	W	Rhat
gene_density	5.712	9.000	1.004
GC content (%)	15.370	0.000	1.001

Note. For all tests, the alternative hypothesis specifies that the location of group *Macro* is smaller than the location of group *Micro*.
Note. Result based on data augmentation algorithm with 5 chains of 1000 iterations.

Descriptives

Raincloud Plots

Bayesian Correlation

Bayesian Pearson Correlations

			n	Pearson's r	BF10
norm_size	-	norm_gene	19	0.962***	1.089e+8
norm_size	-	gene_density	19	-0.662*	23.654
norm_size	-	GC content (%)	19	-0.780***	369.298
norm_gene	-	gene_density	19	-0.582	6.845
norm_gene	-	GC content (%)	19	-0.745***	140.687
gene_density	-	GC content (%)	19	0.908***	162973.383

* BF10 > 10, ** BF10 > 30, *** BF10 > 100

Bayesian Correlation Matrix Plot

Bayesian Independent Samples T-Test

Bayesian Mann-Whitney U Test

	BF$_{-0}$	W	Rhat
gene_density	21.345	1.000	1.005
GC content (%)	24.951	1.000	1.000

Note. For all tests, the alternative hypothesis specifies that the location of group *Macro* is smaller than the location of group *Micro* .
Note. Result based on data augmentation algorithm with 5 chains of 1000 iterations.

Descriptives

Raincloud Plots

Bayesian Correlation

Bayesian Pearson Correlations

			n	Pearson's r	BF$_{10}$
norm_size	-	norm_gene	19	0.992***	7.758e+12
norm_size	-	gene_density	19	-0.551	4.591
norm_size	-	GC content (%)	19	-0.609	9.944
norm_gene	-	gene_density	19	-0.474	2.005
norm_gene	-	GC content (%)	19	-0.545	4.249
gene_density	-	GC content (%)	19	0.952***	1.741e+7

* BF$_{10}$ > 10, ** BF$_{10}$ > 30, *** BF$_{10}$ > 100

Bayesian Correlation Matrix Plot

Bayesian Independent Samples T-Test

Bayesian Mann-Whitney U Test

	BF$_{-0}$	W	Rhat
gene_density	7.625	25.000	1.007
GC content (%)	17.355	15.000	1.003

Note. For all tests, the alternative hypothesis specifies that the location of group *Macro* is smaller than the location of group *Micro*.
Note. Result based on data augmentation algorithm with 5 chains of 1000 iterations.

Descriptives

Raincloud Plots

Bayesian Correlation

Bayesian Pearson Correlations

			n	Pearson's r	BF$_{10}$
norm_size	-	norm_gene	26	0.925***	5.066e+8
norm_size	-	gene_density	26	-0.595**	31.421
norm_size	-	GC content (%)	26	-0.633**	73.342
norm_gene	-	gene_density	26	-0.324	0.836
norm_gene	-	GC content (%)	26	-0.490	5.190
gene_density	-	GC content (%)	26	0.817***	50912.759

* BF$_{10}$ > 10, ** BF$_{10}$ > 30, *** BF$_{10}$ > 100

Bayesian Correlation Matrix Plot

Results

Bayesian Independent Samples T-Test

Bayesian Mann-Whitney U Test

Microchromosomes <30Mb

	BF-₀	W	Rhat
gene_density	20.169	0.000	1.007
GC content (%)	36.521	0.500	1.041

Note. For all tests, the alternative hypothesis specifies that the location of group *Macro* is smaller than the location of group *Micro* .
Note. Result based on data augmentation algorithm with 5 chains of 1000 iterations.

Descriptives

Raincloud Plots

Bayesian Independent Samples T-Test

Bayesian Mann-Whitney U Test

Microchromosomes <10Mb

	BF-₀	W	Rhat
gene_density	1508.004	6.000	1.003
GC content (%)	378.898	2.500	1.006

Note. For all tests, the alternative hypothesis specifies that the location of group *Macro* is smaller than the location of group *Micro* .
Note. Result based on data augmentation algorithm with 5 chains of 1000 iterations.

Descriptives

Raincloud Plots

Bayesian Correlation

Bayesian Pearson Correlations

			n	Pearson's r	BF₁₀
norm_size	-	norm_gene	40	0.986***	2.908e+27
norm_size	-	gene_density	40	-0.266	0.744
norm_size	-	GC content (%)	40	-0.642***	2921.784
norm_gene	-	gene_density	40	-0.302	1.114
norm_gene	-	GC content (%)	40	-0.674***	11004.147
gene_density	-	GC content (%)	40	0.448*	11.276

* BF₁₀ > 10, ** BF₁₀ > 30, *** BF₁₀ > 100

Bayesian Correlation Matrix Plot

Bayesian Independent Samples T-Test

Bayesian Mann-Whitney U Test

	BF₋₀	W	Rhat
gene_density	50.297	1.000	1.062
GC content (%)	217.699	0.000	1.020

Note. For all tests, the alternative hypothesis specifies that the location of group *Macro* is smaller than the location of group *Micro* .
Note. Result based on data augmentation algorithm with 5 chains of 1000 iterations.

Descriptives

Raincloud Plots

Bayesian Independent Samples T-Test

Bayesian Mann-Whitney U Test

	BF₋₀	W	Rhat
gene_density	574.148	3.000	1.005
GC content (%)	631.566	9.500	1.000

Note. For all tests, the alternative hypothesis specifies that the location of group *Macro* is smaller than the location of group *Micro* .
Note. Result based on data augmentation algorithm with 5 chains of 1000 iterations.

Descriptives

Raincloud Plots

Bayesian Correlation

Bayesian Pearson Correlations

			n	Pearson's r	BF₁₀
norm_size	-	norm_gene	40	0.983***	5.665e+25
norm_size	-	gene_density	40	-0.606***	779.978
norm_size	-	GC content (%)	40	-0.728***	162609.166
norm_gene	-	gene_density	40	-0.626***	1566.868
norm_gene	-	GC content (%)	40	-0.749***	584207.309
gene_density	-	GC content (%)	40	0.932***	1.594e+15

* BF₁₀ > 10, ** BF₁₀ > 30, *** BF₁₀ > 100

Bayesian Correlation Matrix Plot

Results

Bayesian Independent Samples T-Test

Bayesian Mann-Whitney U Test

Microchromosomes <30Mb

	BF$_{-0}$	W	Rhat
GC content (%)	20.148	1.000	1.005

Note. For all tests, the alternative hypothesis specifies that the location of group *Macro* is smaller than the location of group *Micro* .
Note. Result based on data augmentation algorithm with 5 chains of 1000 iterations.

Descriptives

Raincloud Plots

GC content (%)

Bayesian Independent Samples T-Test

Bayesian Mann-Whitney U Test

Microchromosomes <10Mb

	BF$_{-0}$	W	Rhat
GC content (%)	56.836	0.000	1.010

Note. For all tests, the alternative hypothesis specifies that the location of group *Macro* is smaller than the location of group *Micro* .
Note. Result based on data augmentation algorithm with 5 chains of 1000 iterations.

Descriptives

Raincloud Plots

GC content (%)

Bayesian Correlation

Bayesian Pearson Correlations

			Pearson's r	BF$_{10}$
norm_size	-	GC content (%)	-0.660***	241.027

* BF$_{10}$ > 10, ** BF$_{10}$ > 30, *** BF$_{10}$ > 100

Bayesian Correlation Matrix Plot

Results

Bayesian Correlation

Bayesian Pearson Correlations

			n	Pearson's r	BF$_{10}$
norm_size	-	norm_gene	24	0.685***	156.865
norm_size	-	gene_density	24	-0.245	0.475
norm_size	-	GC content (%)	24	-0.574*	14.579
norm_gene	-	gene_density	24	0.528	7.026
norm_gene	-	GC content (%)	24	-0.369	1.121
gene_density	-	GC content (%)	24	0.226	0.431

* BF$_{10}$ > 10, ** BF$_{10}$ > 30, *** BF$_{10}$ > 100

Bayesian Correlation Matrix Plot

Supplemental Figure 2: Beeswarm plots splitting genome assemblies by phylogenetic group (Infraorder). (A) Genome size per group for all assemblies (left panel) and long-read only assemblies (right panel). (B) Scaffold N50 and (C) scaffold L50 for all assemblies in each group. Scaffold N50/L50 statistics are capped by physical chromosome sizes within the species in high-quality assemblies, i.e. taxa with macro-/microchromosomes have larger potential N50's and lower potential L50's.

Supplemental Figure 3: Beeswarm plots splitting genome sizes by technology used to generate the primary assembly (contigs). According to total assembly data (B), snakes appear to have significantly smaller genomes than other squamates. However, (C) when accounting for the extreme bias of short-read assemblies in snakes, this difference disappears. *Long = PacBio and/or ONT, Short = Illumina, Sanger, 454, etc.*

Genome Size by Sequencing Technology

Genome Size: Snakes vs. Non-snakes

Supplemental Table 1: Squamates vs mammal funding source comparison. (A) Most high-quality squamate genomes are generated using soft money from start-up/personal funds or (generally smaller) basic science foundation grants, compared to mammals that receive most of their funding from health and agricultural funding agencies rather than basic/personal funding sources. (B) Breakdown of genomes and references used to generate the funding information shown in A.

A. Total Numbers			
Taxon	**Basic Science***	**Applied Science****	**Private/Personal*****
Squamate	10	4	6
Mammal	2	18	0

B. Numbers Breakdown		
Squamates	**Funding**	**Citation**
Sphaerodactylus townsendi	Personal***	Pinto et al. 2022
Eublepharis macularius	Personal***	Pinto et al. 2022
Paroedura picta	Personal***	Hara et al. 2018
Pogona vitticeps	Personal***	Georges et al. 2015
Phrynosoma platyrhinos	Personal***	Koochekian et al. 2022
Pantherophis guttatus	Personal***	Ullate-Agote and Tzika, 2020
Anolis carolinensis	Applied**	Alföldi et al. 2011
Lacerta agilis	Applied**	VGP, *unpublished*
Python bivittatus	Applied**	Castoe et al. 2013
Thamnophis elegans	Applied**	VGP, *unpublished*
Shinisaurus crocodilurus	Basic*	Xie et al. 2022
Sceloporus tristichus (SNOW)	Basic*	Bedoya & Leache, 2021
Sceloporus tristichus (HOL)	Basic*	Bedoya & Leache, 2021
Sceloporus undulatus	Basic*	Westfall et al. 2021
Anolis sagrei	Basic*	Geneva et al. 2021
Podarcis muralis	Basic*	Andrade et al. 2019
Zootoca vivipara	Basic*	Yurchenko et al. 2020
Crotalus viridis	Basic*	Schield et al. 2019
Hydrophis cyanocinctus	Basic*	Li et al. 2021
Hydrophis curtus	Basic*	Li et al. 2021
Mammals	**Funding**	**Citation**
Ochotona princeps	Basic*	Sjodin et al. 2021
Ursus arctos	Basic*	Armstrong et al. 2022
Odocoileus virginianus	Applied**	London et al. 2022
Bos taurus	Applied**	Zimin et al. 2009
Mus musculus	Applied**	European Bioinformatics Institute, 2002
Homo sapiens	Applied**	Nurk et al. 2022
Rattus norvegicus	Applied**	Weiss et al. 2004
Oryctolagus cuniculus	Applied**	Carneiro et al. 2014
Equus caballus	Applied**	Wade et al. 2009
Canis lupus	Applied**	Lindblad-Toh et al. 2005
Felis catus	Applied**	Pontius et al. 2007
Pan troglodytes	Applied**	Chimpanzee Sequencing and Analysis Consortium, 2005
Macaca mulatta	Applied**	Gibbs et al. 2007
Ovis aries	Applied**	Jiang et al. 2014

Sus scrofa	Applied**	Groenen et al. 2012
Monodelphis domestica	Applied**	Mikkelsen et al. 2007
Peromyscus maniculatus	Applied**	Harringmeyer and Hoekstra, 2022
Capra hircus	Applied**	Bickhart et al. 2017
Pongo abelii	Applied**	Kronenberg et al. 2018
Microcebus murinus	Applied**	Larsen et al. 2017

Key:
* National Science Foundation, USA (NSF) or equivalent.
** National Institutes of Health (NIH)/United States department of Agriculture (USDA) or equivalent.
*** Laboratory startup funds, private granting agencies, or equivalent.

Table References:

Alföldi, J., Di Palma, F., Grabherr, M., Williams, C., Kong, L., Mauceli, E., ... & Lindblad-Toh, K. (2011). The genome of the green anole lizard and a comparative analysis with birds and mammals. Nature, 477(7366), 587-591.

Andrade, P., Pinho, C., Pérez i de Lanuza, G., Afonso, S., Brejcha, J., Rubin, C.-J., Wallerman, O., Pereira, P., Sabatino, S. J., Bellati, A., Pellitteri-Rosa, D., Bosakova, Z., Bunikis, I., Carretero, M. A., Feiner, N., Marsik, P., Paupério, F., Salvi, D., Soler, L., … Carneiro, M. (2019). Regulatory changes in pterin and carotenoid genes underlie balanced color polymorphisms in the wall lizard. Proceedings of the National Academy of Sciences, 116(12), 5633–5642. https://doi.org/10.1073/pnas.1820320116

Armstrong, E. E., Perry, B. W., Huang, Y., Garimella, K. V., Jansen, H. T., Robbins, C. T., ... & Kelley, J. L. (2022). A beary good genome: Haplotype-resolved, chromosome-level assembly of the brown bear (Ursus arctos). Genome biology and evolution, 14(9), evac125.

Bedoya, A. M., & Leaché, A. D. (2021). Characterization of a pericentric inversion in plateau fence lizards (Sceloporus tristichus): evidence from chromosome-scale genomes. G3, 11(2), jkab036.

Bedoya, A. M., & Leaché, A. D. (2021). Characterization of a pericentric inversion in plateau fence lizards (Sceloporus tristichus): evidence from chromosome-scale genomes. G3, 11(2), jkab036.

Bickhart, D. M., Rosen, B. D., Koren, S., Sayre, B. L., Hastie, A. R., Chan, S., ... & Smith, T. P. (2017). Single-molecule sequencing and chromatin conformation capture enable de novo reference assembly of the domestic goat genome. Nature genetics, 49(4), 643-650.

Carneiro, M., Rubin, C. J., Di Palma, F., Albert, F. W., Alföldi, J., Barrio, A. M., ... & Andersson, L. (2014). Rabbit genome analysis reveals a polygenic basis for phenotypic change during domestication. Science, 345(6200), 1074-1079.

Castoe, T. A., De Koning, A. J., Hall, K. T., Card, D. C., Schield, D. R., Fujita, M. K., ... & Pollock, D. D. (2013). The Burmese python genome reveals the molecular basis for extreme adaptation in snakes. Proceedings of the National Academy of Sciences, 110(51), 20645-20650.

Chimpanzee Sequencing and Analysis Consortium. (2005). Initial sequence of the chimpanzee genome and comparison with the human genome. Nature, 437(7055), 69-87.

European Bioinformatics Institute. (2002). Initial sequencing and comparative analysis of the mouse genome. Nature, 420(6915), 520-562.

Geneva, A. J., Park, S., Bock, D., de Mello, P., Sarigol, F., Tollis, M., ... & Losos, J. B. (2021). Chromosome-scale genome assembly of the brown anole (Anolis sagrei), a model species for evolution and ecology. bioRxiv.

Georges, A., Li, Q., Lian, J., O'Meally, D., Deakin, J., Wang, Z., ... & Zhang, G. (2015). High-coverage sequencing and annotated assembly of the genome of the Australian dragon lizard Pogona vitticeps. Gigascience, 4(1), s13742-015.

Gibbs, R. A., Rogers, J., Katze, M. G., Bumgarner, R., Weinstock, G. M., Mardis, E. R., ... & Smith, D. G. (2007). Evolutionary and biomedical insights from the rhesus macaque genome. science, 316(5822), 222-234.

Groenen, M. A., Archibald, A. L., Uenishi, H., Tuggle, C. K., Takeuchi, Y., Rothschild, M. F., ... & Schook, L. B. (2012). Analyses of pig genomes provide insight into porcine demography and evolution. Nature, 491(7424), 393-398.

Hara Y, Takeuchi M, Kageyama Y, Tatsumi K, Hibi M, Kiyonari H, Kuraku S. 2018. Madagascar ground gecko genome analysis characterizes asymmetric fates of duplicated genes. BMC Biology. 16: 40.

Harringmeyer, O. S., & Hoekstra, H. E. (2022). Chromosomal inversion polymorphisms shape the genomic landscape of deer mice. Nature ecology & evolution, 1-15.

Jiang, Y., Xie, M., Chen, W., Talbot, R., Maddox, J. F., Faraut, T., ... & Dalrymple, B. P. (2014). The sheep genome illuminates biology of the rumen and lipid metabolism. Science, 344(6188), 1168-1173.

Koochekian, N., Ascanio, A., Farleigh, K., Card, D. C., Schield, D. R., Castoe, T. A., & Jezkova, T. (2022). A chromosome-level genome assembly and annotation of the desert horned lizard, Phrynosoma platyrhinos, provides insight into chromosomal rearrangements among reptiles. GigaScience, 11.

Kronenberg, Z. N., Fiddes, I. T., Gordon, D., Murali, S., Cantsilieris, S., Meyerson, O. S., ... & Eichler, E. E. (2018). High-resolution comparative analysis of great ape genomes. Science, 360(6393), eaar6343.

Larsen, P. A., Harris, R. A., Liu, Y., Murali, S. C., Campbell, C. R., Brown, A. D., ... & Worley, K. C. (2017). Hybrid de novo genome assembly and centromere characterization of the gray mouse lemur (Microcebus murinus). BMC biology, 15(1), 1-17.

Li, A., Wang, J., Sun, K., Wang, S., Zhao, X., Wang, T., Xiong, L., Xu, W., Qiu, L., Shang, Y., Liu, R., Wang, S., & Lu, Y. (2021). Two Reference-Quality Sea Snake Genomes Reveal Their Divergent Evolution of Adaptive Traits and Venom Systems. Molecular Biology and Evolution, 38(11), 4867–4883. https://doi.org/10.1093/molbev/msab212

Li, A., Wang, J., Sun, K., Wang, S., Zhao, X., Wang, T., Xiong, L., Xu, W., Qiu, L., Shang, Y., Liu, R., Wang, S., & Lu, Y. (2021). Two Reference-Quality Sea Snake Genomes Reveal Their Divergent Evolution of Adaptive Traits and Venom Systems. Molecular Biology and Evolution, 38(11), 4867–4883. https://doi.org/10.1093/molbev/msab212

Lindblad-Toh, K., Wade, C. M., Mikkelsen, T. S., Karlsson, E. K., Jaffe, D. B., Kamal, M., ... & Lander, E. S. (2005). Genome sequence, comparative analysis and haplotype structure of the domestic dog. Nature, 438(7069), 803-819.

London, E. W., Roca, A. L., Novakofski, J. E., & Mateus-Pinilla, N. E. (2022). A De Novo Chromosome-level Genome Assembly of the White-tailed Deer, Odocoileus virginianus. Journal of Heredity.

Mikkelsen, T. S., Wakefield, M. J., Aken, B., Amemiya, C. T., Chang, J. L., Duke, S., ... & Lindblad-Toh, K. (2007). Genome of the marsupial Monodelphis domestica reveals innovation in non-coding sequences. Nature, 447(7141), 167-177.

Nurk, S., Koren, S., Rhie, A., Rautiainen, M., Bzikadze, A. V., Mikheenko, A., ... & Phillippy, A. M. (2022). The complete sequence of a human genome. Science, 376(6588), 44-53.

Pinto, B. J., Gamble, T., Smith, C.H., Keating, S.E., Havird, J., Chiari, Y. (2022). The revised leopard gecko (Eublepharis macularius) reference genome 'raises the bar' for squamate genomics. In prep.

Pinto, B. J., Keating, S. E., Nielsen, S. V., Scantlebury, D. P., Daza, J. D., & Gamble, T. (2022). Chromosome-level genome assembly reveals dynamic sex chromosomes in Neotropical leaf-litter geckos (Sphaerodactylidae: Sphaerodactylus). Journal of Heredity.

Pontius, J. U., Mullikin, J. C., Smith, D. R., Team, A. S., Lindblad-Toh, K., Gnerre, S., ... & NISC Comparative Sequencing Program. (2007). Initial sequence and comparative analysis of the cat genome. Genome research, 17(11), 1675-1689.

Rhie, A., McCarthy, S. A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva, M., Chow, W., Fungtammasan, A., Kim, J., Lee, C., Ko, B. J., Chaisson, M., Gedman, G. L., Cantin, L. J., Thibaud-Nissen, F., Haggerty, L., Bista, I., Smith, M., … Jarvis, E. D. (2021). Towards complete and error-free genome assemblies of all vertebrate species. Nature, 592(7856), 737–746. https://doi.org/10.1038/s41586-021-03451-0

Rhie, A., McCarthy, S. A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva, M., Chow, W., Fungtammasan, A., Kim, J., Lee, C., Ko, B. J., Chaisson, M., Gedman, G. L., Cantin, L. J., Thibaud-Nissen, F., Haggerty, L., Bista, I., Smith, M., … Jarvis, E. D. (2021). Towards complete and error-free genome assemblies of all vertebrate species. Nature, 592(7856), 737–746. https://doi.org/10.1038/s41586-021-03451-0

Schield, D. R., Card, D. C., Hales, N. R., Perry, B. W., Pasquesi, G. M., Blackmon, H., Adams, R. H., Corbin, A. B., Smith, C. F., Ramesh, B., Demuth, J. P., Betrán, E., Tollis, M., Meik, J. M., Mackessy, S. P., & Castoe, T. A. (2019). The origins and evolution of chromosomes, dosage compensation, and mechanisms underlying venom regulation in snakes. Genome Research, 29(4), 590–601. https://doi.org/10.1101/gr.240952.118

Sjodin, B. M., Galbreath, K. E., Lanier, H. C., & Russello, M. A. (2021). Chromosome-level reference genome assembly for the American pika (Ochotona princeps). Journal of Heredity, 112(6), 549-557.

Ullate-Agote, A., & Tzika, A. C. (2021). Characterization of the Leucistic Texas Rat Snake Pantherophis obsoletus. Frontiers in Ecology and Evolution, 9. https://doi.org/10.3389/fevo.2021.583136

Wade, C. M., Giulotto, E., Sigurdsson, S., Zoli, M., Gnerre, S., Imsland, F., ... & Lindblad-Toh, K. (2009). Genome sequence, comparative analysis, and population genetics of the domestic horse. Science, 326(5954), 865-867.

Weiss Robert B., Dunn Diane M., Green Eric D., Blakesley Robert W., Bouffard Gerard G., de Jong, P. J., Osoegawa, K., Zhu, B., Marra, M., ... & Collins, F. (2004). Genome sequence of the Brown Norway rat yields insights into mammalian evolution. Nature, 428(6982), 493-521.

Westfall, A. K., Telemeco, R. S., Grizante, M. B., Waits, D. S., Clark, A. D., Simpson, D. Y., ... & Schwartz, T. S. (2021). A chromosome-level genome assembly for the Eastern Fence Lizard (Sceloporus undulatus), a reptile model for physiological and evolutionary ecology. GigaScience, 10(10), giab066.

Xie HX, Liang XX, Chen ZQ, Li WM, Mi CR, Li M, ... & Du WG. 2022. Ancient demographics determine the effectiveness of genetic purging in endangered lizards. Mol Biol Evol. 39: msab359.

Yurchenko, A. A., Recknagel, H., & Elmer, K. R. (2020). Chromosome-Level Assembly of the Common Lizard (Zootoca vivipara) Genome. Genome Biology and Evolution, 12(11), 1953–1960. https://doi.org/10.1093/gbe/evaa161

Zimin, A. V., Delcher, A. L., Florea, L., Kelley, D. R., Schatz, M. C., Puiu, D., ... & Salzberg, S. L. (2009). A whole-genome assembly of the domestic cow, Bos taurus. Genome biology, 10(4), 1-10.