Evidence against a cortical module for processing communicative gaze

Marius Görner[1,2,3,*,+], Peter W Dicke[1], and Peter Thier[1,4,*]

[1] Hertie Institute for Clinical Brain Research, Thier Lab, Tübingen, Germany

[2] GTC of Neuroscience, Tübingen, Germany

[3] IMPRS for Cognitive and Systems Neuroscience, Tübingen, Germany

[4] Werner Reichardt Centre for Integrative Neuroscience, Tu□bingen Germany

[+]current address: Max Planck Institute of Psychiatry, Department Emotion Research, Munich, Germany

[*] Corresponding authors: Peter Thier (thier@uni-tuebingen.de), Marius Görner (mariusgoerner@gmail.com)

Author contribution: MG designed the study, collected the data, analyzed the data, wrote the manuscript; PD collected the data; PT designed the study, wrote the manuscript

Classification: Biological Sciences/Neuroscience

Keywords: Social neuroscience, Theory of Mind, Joint attention, Gaze following, fMRI, domain specificity, domain generality

Competing Interest Statement: The authors have no conflicts of interest.

This PDF file includes:

Main Text

Figures 1 to 7, S1-4

Tables 1 to 3, S1-2

Abstract. Where we look lets others know what we are interested in and allows them to join our focus of attention. In several studies our group investigated the neuronal basis of gaze following behavior in humans and macaques and described a *gaze following patch* (GFP) in the posterior temporal cortex as being of central importance for this function. To our knowledge, this makes the GFP the most promising neurobiological correlate of Simon Baron-Cohen's *eye direction detector* (EDD), an integral part of his *mindreading system*. With the latter, Baron-Cohen proposed a set of domain-specific neurocognitive modules implementing a functional framework he suggested to be necessary to establish a Theory of Mind - the attribution of mental states to others. The tenet of domain-specificity requires that the EDD processes exclusively eye-like stimuli. In this preregistered fMRI study, we aimed to critically test if the GFP fulfills this criterion. Contrary to previous studies, the present results clearly demonstrate that GFP activity is not exclusively correlated with the perception of other people's gaze direction and thus does not fulfill the criterion of domain-specificity. Furthermore, we show that it cannot be functionally dissociated from the MT+ complex, with which it overlaps anatomically. In the context of our previous studies which were specifically designed to capture the full behavioral relevance of active gaze following and to reveal the EDD, the results of the present study provide evidence that such a domain specific module does not exist. Therefore, the plausibility of Baron-Cohen's *mindreading model* as a set of domain-specific modules is put into question.

Significance statement. The presented results demonstrate that a brain area previously thought to be a domain-specific cortical module specialized in processing other people's focus of visual attention is, in fact, domain-general. More specifically, we show that it is indistinguishable from cortical areas involved in the processing of visual motion. These findings add to a growing body of evidence arguing against domain-specific accounts in social neuroscience which posit that the functional building blocks of social behaviors are facilitated by dedicated neurocognitive modules.

Introduction

The direction of our gaze plays a fundamental role when we interact with a fellow human. Just by observing the other person's gaze averting her eyes from one towards a new location, we can tell that her attention has been grabbed by something else, inevitably also attracting our own attention. Thereby we establish *joint attention* - the congruence of attentional foci. Further, by mapping our own beliefs, desires and intentions associated with the object of the other´s attention enables us to develop a *theory of (the other´s) mind* (ToM) which allows us to predict the other´s behavior. With his *mindreading system*, Simon Baron-Cohen suggested a functional framework which is implemented by four domain-specific neuro-cognitive modules underpinning the route towards a full-fledged TOM. In this, the *eye direction detector* (EDD) is the node that contributes the information on gaze direction which is integrated with information from an *intentionality detector* (ID) and relayed to the *shared attention mechanism* (SAM), a term often used as an alternative to *joint attention* in order to emphasize the agent´s awareness of the congruence of their attention foci. Finally, the latter passes its information to the *theory of mind mechanism* (ToMM)[1–3]. He based this idea on Fodor's concept of modularity, which implies that the modular components have distinct and specific neural architectures. These architectures would consist of populations of neurons that exclusively process the respective information; in the case of the EDD this means eye-like stimuli with their typical shape and motion properties[1,4]. Apart from Baron-Cohen's concept, the idea of modularity is widespread in the field of social cognitive neuroscience[5–8] but has also been criticized from early on[9–11].

Given the ecological importance of judging gaze directions – whether for predator avoidance, prey detection or other forms of social communication[12] – it seems indeed conceivable that a specialized neural mechanism evolved to optimize processing in order to facilitate rapid behavioral responses to the other´s gaze (see[13] for a recent review on the evolutionary perspective on gaze-following). How fast you can detect a staring predator may decide upon life and death. Support for this idea is provided by behavioral studies showing that responses to gaze stimuli in both humans and macaques appear to be reflex-like, i.e. they are fast and cannot be suppressed[14–16,6,17–19]. Perrett and colleagues were the first to describe neurons in the superior temporal sulcus (STS) that are tuned to varying gaze, head and body orientations[20], yet without testing these neurons for behavioral relevance. Other studies have reported related results on the representation of gaze stimuli in the temporal cortex of monkeys and humans, in the latter emphasizing the right temporal cortex[21–23], which in general dominates the processing of information provided by faces[24] (see[25–27] for reviews).

In a series of behavioral, fMRI, and electrophysiological studies our group investigated the neural basis of gaze-following behavior in macaques and humans. In this work we were able to identify a circumscribed region in the posterior temporal cortex of humans and macaques, which we baptized the *gaze following patch* (*GFP*). This region is comparatively more active during gaze-following than

in control conditions, characterized by very similar visual and motor demands, yet lacking the need to

follow gaze[19,28–34,8]. This characteristic makes the GFP an obvious neural correlate of Baron-Cohen's

EDD. The distinctive feature of our experimental design was that it required *active* gaze-following,

i.e. participants did not only passively observe a face gazing in a certain direction as in the previous

work mentioned but had to utilize it to identify a target among distractors and to subsequently shift

their own visual attention to it. In the control condition that we used the target had to be identified

based on the iris-color[31] or the identity[8] of the stimulus face. We found that the GFP — revealed by

contrasting gaze-following and iris-color matching — does not overlap with any of the neighboring

elements of the face patch system[5,31,35], albeit positioned close to the middle face patch. Moreover, it

is located close to the middle temporal area (MT), well-known to be instrumental in the processing of

visual motion. In fact the human GFP is overlapping with parts of the MT+ complex delineated by the

neurosynth.org database[36] as well as by our own pilot experiments involving low-level motion stimuli

in humans (see Figure 2 and S1). Visual motion may indeed have been a critical stimulus component

in our work as the gaze stimuli used were based on a portrait in which the eye axes switched from

straight ahead to averted, thereby giving the impression of a gaze saccade – i.e., high speed motion of

the eyes. Although the control condition involved the same motion component, it was behaviorally

irrelevant as it was the iris-color that participants had to exploit to identify the target with the

matching color. Thus, it may well be that the critical aspect of the stimulus that triggered the GFP

activity was not the actual retrieval and use of gaze direction but the need to attend to the associated

movement of the eyes, activating generic motion specific cortex.

Hence, in the present, pre-registered study[37], we tested whether the GFP needs to be fused with

the visual motion processing system, thereby depriving the GFP of its property of domain-specificity

assumed by the earlier work[8,28,29,31,38]. This seems to be the case: our results clearly show that not only

the observation of an eye saccade is accompanied by activity in the GFP. Rather, a stimulus that bears

no visual resemblance to eyes, but has a movement pattern comparable to a saccade, also elicits

activity. Furthermore, we show that the activity in the GFP cannot be distinguished from activity in a

ROI delineated by low-level visual motion stimuli, as already suggested by the anatomical

congruence (see Figure 2). Given that all of our experiments have been designed to reveal an EDD as

an example of a cortical module underpinning domain specificity, we propose that our results have

strong implications beyond the ontological status of the GFP. Considering that the GFP —the best

candidate of a cortical realization of the EDD so far—failed a critical test of domain specificity

suggests that gaze-following may not have a domain-specific neural basis at all, but instead be

grounded in domain general cortical processing serving a much wider range of perceptual functions.

Figure 1 Illustration of experiment 1 and 2. In the example of experiment 1 the cubes indicate the middle target and the
gaze as well as the iris-color the left target. Depending on the instruction, the participant will have to choose the respective
target. In experiment 2 the two stimulus components gaze-shift and cube-rotation are presented separately. In the experiment
the schematic face shown here is replaced by a photo of a person.

Figure 2 Anatomical locations of motion related areas and the GFP. Shown are transversal cuts from   –   to
in steps of   . Locations given by the neurosynth[36] database when searching for "visual motion" are outlined in
white. The result of the MT localizer of Pilot 1 is outlined in cyan and the GFP from Marquardt et al [31] is outlined in green.

Results

Confirmatory

Experiment 1

In each trial of the first experiment the participants got to see a gaze shift as well as a change of the iris color of the demonstrator face, and a rotation of cubes that were positioned on the face. Auditory instructions indicated which of the cues should be used to identify the target (see Figure 1). For each participant we used the BOLD contrast between trials in which they had to use the gaze shift and trials in which they had to use the change of the iris color to identify the GFP-ROI. For all participants except sub-06 we could identify individual GFP-ROIs, for sub-06 we used a spherical ROI that was centered on the local maximum closest to the reference coordinates of the GFP (Figure 3/ S1 for a depiction of the ROIs). Using these as masks, we extracted the beta values of the *gaze-shift* and *cubes-rotation* conditions for each run of each participant (Figure 3).

Figure 3 For each participant beta values extracted from the GFP-ROIs are shown grouped by hemisphere and condition. Individual ROIs are shown on top of each boxplot For all but participant 6, the GFP could be identified individually. In both conditions the GFP is consistently activated with beta values larger than 0. Beta values related to the *cube-rotation* are consistently larger than those related to the *gaze-shift* condition.

Statistical analysis supports a difference in the activity associated with the two conditions, but with larger activity in the *cubes-rotation* condition than in the *gaze-shift* condition (see Table 1, mixed effects model with *run* as group-level factor, *BFs* are   transformed).

Table 1 Statistical results regarding differences between conditions (Table S1, Q1).

LH	-0.65	$[-0.87, -0.4]$	-11.02
RH	-1.15	$[-1.48, -0.82]$	-88.93

The non-logarithmically transformed Bayes Factor for the right hemisphere exceeds numerical
precision, fluctuating between positive and negative values, meaning that the posterior probability of
$\Delta\beta_{GFP-right}^{Gaze-Cubes} = 0$, i.e., no difference between the BOLD responses in the conditions.

Testing for differences between the left and the right GFP within each condition the data support
the existence of a difference for the *cube-rotation* with *moderate* evidence but equal activation related
to *gaze-shift* with *strong* evidence (see Table 2, mixed effects model with *run* as group-level factor,
BFs are log transformed).

Table 2 Statistical results regarding differences between hemispheres (Table S1, Q2).

	$\Delta\beta^{RH-LH}$	$CI^{95\%}$	BF_{01}^{Q2}
Gaze	0.01	$[-0.28, 0.31]$	1.41
Cubes	0.54	$[0.21, 0.88]$	-0.65

Experiment 2

Unlike in the first experiment, in this second experiment participants saw only either the cubes or
the demonstrator face shifting eye-gaze in a given trial. Using the individual GFP-ROIs from
experiment 1 and the individual MT+ ROIs from the motion localizer tasks (Figure S1 shows all
individual ROIs) we estimated hemodynamic response functions (*HRF*) for each participant
individually as well as the group level *HRFs* grouped by ROI, hemisphere, condition and target (left,
middle, right) (see Figure 7).

There were clear responses for all conditions in all ROIs and hemispheres with slightly larger
amplitudes in the *gaze-only* related responses. The temporal patterns are indistinguishable when
comparing within each ROI across conditions, as well as when comparing within conditions across
ROIs. Especially when comparing the responses of one condition across ROIs of the same
hemisphere, across columns in Figure 4, there is virtually no difference. One characteristic of the
responses that stands out is that in all plots (except for the right GFP) there is one curve that is
different from the other two. The difference is in the amplitude as well as the timing of the peak and
trough. For the left ROIs it is the response related to the right target and for the right ROIs the one

related to the left target. This pattern corresponds to the prediction based on the assumption that the
responses in the ROIs are lateralized, which is a property of area MT (Table S1, Q3.1 & 4.1).

To quantify the similarity between the responses to the *gaze-only* and *cubes-only* conditions we
computed the correlation between pairs of responses (Table S1, Q3&4). The correlations were
computed for each participant individually. The relevant pairs for which we computed the correlations
are shown in Table 3. Correlations between *HRF*s were only computed within each ROI (GFP or
MT+), however, across hemispheres and conditions. To directly compare the GFP with MT+ the
obtained correlation coefficients were then analyzed for their correlation, i.e., asking the question if
the correlation coefficients for the pairs of *HRF*s have similar values in the GFP and MT+ (Table S1,
Q5). Figure 5 shows the results of the direct comparisons between the two ROIs (Table S1, Q3-5).
Data of the participant for whom we could not identify an individual GFP-ROI are shown in the
scatter plot of Figure 5 with reduced opacity. We did not include this participant's data in the
statistical analysis presented here because the fact the ROI could not be inferred directly from the
gaze-shift minus *iris-color* contrast image casts doubt on whether there was a task-related signal at all.
However, including this subject changed the results only quantitatively. Our central finding is that
both ROIs showed the same correlation pattern. We found *strong* evidence that both, the GFP and
MT+, exhibited differences between the groups reflecting conditions (black and red bar plots in

Figure 5; Table S1, Q3.2 & 4.2). Questions 3.1 & 4.1, asking whether the ROIs show lateralized

responses independent of the experimental condition, cannot be answered with sufficient evidence

(see *Methods - Transparent Changes*); the GFP data, however, suggests with *moderate* evidence that

there is no difference between the groups reflecting lateralized responses (; Figure 5, top

left, blue and yellow bars). The similarity of the correlation patterns for the two ROIs is also

expressed in their direct comparison. The correlation between the correlation patterns for the GFP and

MT+ is 0.75, meaning that if a given pair of HRFs is (not) correlated in the GFP it is also (not)

correlated in MT+ (Figure 5, scatter plot, Table S1, Q5).

Figure 5 Results of the correlation analysis. The boxplots illustrate the correlation coefficients grouped according to the

two predictions (blue-yellow: Q3.1/ 4.1; red-black: Q3.2/ 4.2) for the two ROIs (top-left: GFP; bottom-right: MT+). The

inlets show the $\log_{10}$ transformed Bayes Factors. The scatter plot shows the correlation coefficients based on the two ROIs

independent of the groupings. It can be seen that if a pair of HRFs is correlated in one ROI, it tends to be correlated in the

other as well. The inlet shows the correlation coefficient between the two ROIs as well as the respective Bayes Factor

indicating *extreme evidence* in favor of a correlation different from 0. The data points plotted with a lower opacity come

from the subject for whom it was not possible to identify an individual GFP-ROI and whose data was not included in the

quantitative analysis.

Exploratory

To explore brain wide qualitative differences between the activity related to the *gaze-shift* and the

cube-rotation conditions we looked for brain areas where one of the experimental conditions elicited

positive β-values and the respective other condition non-significant (, *uncorrected*) or

negative *β*-values. Further, we applied a cluster size threshold with a minimum of 15 adjacent voxels

to remove noise and focus on broader activity patterns. Areas shaded in green in Figure 6 are those

that carried positive *β*-values during *gaze-shift* following and non-significant/ negative *β*-values

during *cube-rotation* following. Areas shaded in purple have the opposite pattern. Finally, the

uncolored areas are those in which the *β*-values were not significant or both conditions evoked a

response of the same sign.

In the context of our experiment, areas exclusively associated with gaze shifts tended to be

located in more medial parts of the occipital cortex and in the temporal cortex, whereas areas

exclusively associated with cube rotations tended to be located in lateral parts of the occipital cortex

and extend into parietal areas. More specifically, hotspots related to gaze shifts were located in the

medial parts of early visual areas BA 17, 18 and 19. Furthermore, there were hotspots in BA 37 (more

medial) in both hemispheres, in BA 23 and in the temporal cortices with hotspots in BA 41, 22 and

21. Areas exclusively activated in the *cube-rotation* condition were located in early visual areas BA

18 and 19, as well as in BA 37 (more lateral). In the parietal cortex, right BA 39 (extending into BA

22) and bihemispheric BA 40 and BA 7 exhibited hotspots.

Figure 6 Exploratory whole brain analysis of group-level activity patterns exclusively associated with *gaze-shift* and

cube-rotation following. First, the *β*-maps were thresholded with p=0.05 (uncorrected) and a minimum cluster-size of 15

voxels. Then all voxels that were positively correlated with one condition were removed from the *β*-map of the other

condition. I.e. all areas shaded in green represent activations that were positively correlated only with the *gaze-shift*

condition, and all areas shaded in purple were positively correlated only with the *cube-rotation* condition. The area outlined

in bright green is the GFP-ROI based on data from Marquardt et al. [31]. The areas outlined in white the MT+-ROI based on

the neurosynth database.

Discussion

It has been debated for many years whether there are neurocognitive modules that underlie complex (social) behaviors[39,9–11,40]. The discussion was sparked by Jeremy Fodor's influential thoughts on the modularity of the mind. But in fact, Fodor distinguished between *input systems* and *central systems* and proposed that only the former were modular in the strict sense[4]. Ignoring this distinction, social cognitive neuroscience has proposed central neurocognitive modules processing different aspects of social interactions. Simon Baron-Cohen's *mindreading system* is a particularly influential case in point.

In this study, we set out to critically reexamine the assumption of a modular architecture of mind-reading by questioning the existence of an EDD, a putative cortical module underpinning the perception of gaze direction. Such a module has been suggested to be accommodated by a distinct region in the occipitotemporal cortex (the *gaze following patch*, GFP)[1,2,28,31,34,38]. In order to challenge the assumed specificity of the GFP for the other´s gaze, we tested its response to a stimulus that carried directional information as well as visual motion while not resembling eyes. As a first step, we reproduced the GFP in each experimental subject by computing the functional contrast *gaze-shift following* minus *iris-color* that had been used in previous studies to delineate it. We then compared the *β*-values associated with the control condition lacking eye features, *cube-rotation following* with those related to *gaze-shift following* for the GFP-ROIs. This comparison clearly showed that the GFP is active not only when observing a gaze shift of a demonstrator, but also when observing a non-biological stimulus that has similarity to eyes, while moving in a comparable way and having a comparable behavioral relevance in the context of the experimental task. In fact, the *β*-values associated with the *cube-rotation* condition were consistently larger than those related to the *gaze-shift* condition for all subjects. This result speaks against the hypothesis that the GFP is a cortical module that processes others' gaze direction in a domain-specific manner. We can think of two reasons why the activity associated with the *cube-rotation* condition is even greater: either the detection of the cube rotation was more difficult – potentially due to interference with the face/ gaze stimulus in the background –, or the motion energy of the cube rotation was larger. The latter could be due to the fact that the cube rotation was animated by several frames, whereas the gaze shift was animated by only two frames. Whichever option may hold, both would indicate even more strongly that the GFP is not a dedicated EDD, since its activity scales either with the difficulty of detecting visual motion or with the energy of the motion stimulus. The fact that the activations associated with cube rotations were consistently greater than those associated with gaze shifts for all participants also rules out a potential alternative explanation for the activation of the GFP during cube-rotation trials: face pareidolia. This is a perceptual effect where one sees a face in a pattern made from arbitrary elements that resembles the structural properties of a face[41,42]. In this sense participants may have started to perceive eyes in the cubes after being exposed to the experiment for a while. If face

pareidolia was the cause, however, the activation associated with cube rotations would be expected to
be smaller and more variable across participants[43]. In a recent fMRI study[44] we had already
demonstrated that a gaze cue which does not involve any apparent visual motion is not sufficient to
reproduce the GFP. The present study provides an explanation for this finding.

Having learned that gaze shifts are not the only stimuli, GFP activity is positively correlated with,
we investigated whether it can be functionally dissociated from brain areas specialized in the
processing of visual motion, delineated by previous work resorting to a large spectrum of visual
motion stimuli and usually summarized under terms like MT+ or the "visual motion complex"[45–48]
(see Figure 2). To this end, in Experiment 2 we first determined an MT+ ROI for each participant
using a standard MT+ localization paradigm involving passive viewing of moving and static dots. We
compared the responses to *cube-rotation* and *gaze-shift following* within and between the GFP and the
MT+ ROIs. Importantly, in Experiment 2 we had modified the paradigm so that participants only saw
either the gaze shift or the cube rotation in each trial, whereas in Experiment 1 they were presented
together (see Figure 1). We introduced this modification to be able to explain a potential outcome of
Experiment 1, namely that the GFP would be active during *cube-rotation* trials but with smaller *β*-
values than during *gaze-shift* trials. This hypothetical result could have been explained by the
observation that behavioral responses to observed gaze shifts are reflexive and occur even if the task
requires to ignore them[14,49,17,19]. By separating the stimuli, we therefore ensured that any BOLD
response we would observe is a genuine response to the stimulus in question. However, because our
final results of Experiment 1 yielded larger *β*-values related to *cube-rotation*, falsifying the hypothesis
that the GFP is a cortical module dedicated to the processing of others' gaze-direction, we did not
continue with data collection until all quantitative analyses of experiment 2 reached the set threshold
(see *Transparent Changes*). To analyze Experiment 2, we estimated the hemodynamic response
functions (HRF) for the different trial types (grouped by stimulus and target ID) and ROIs and
computed the correlation between them (see Table 3 for a list of all pairings and Figure 7 for a
conceptual illustration). This method protected the analysis from effects that might arise from
differences in the visual complexity of the stimuli, as it relies purely on temporal dynamics. A
property of the experimental design led us to predict a specific response pattern that would occur if
the GFP corresponded to area MT: As illustrated in Figure 7 during each trial of our experiments two
motion events occurred, first the gaze shift/ cube rotation towards the target and second the reset to
the neutral orientations at the end of each trial. Importantly, the second motion event happened during
the period in which the participants looked at the respective target object, meaning that it occurred in
their visual periphery. As is well-known, the areas of the motion complex show lateralized responses
to peripheral input. Therefore, we predicted that if ROIs consist of voxels which are part of MT+, they
should exhibit sustained responses in the contralateral as compared to the ipsilateral hemisphere
relative to the second motion event (see Table S1 and[37]). This would give rise to a specific correlation

pattern among the pairs (yellow and blue groups in Figure 5 and Table 3). However, our data do not provide sufficient evidence to confirm or refute this prediction as we terminated data collection before having reached the set threshold, as explained above. Instead, we found *strong evidence* in favor of the alternative prediction that the correlations depend on the condition. That is, correlations between BOLD responses during trials of the same condition were slightly stronger than between those of different conditions (black and red groups in Figure 5 and Table 3). However, it is essential that this was the case for both ROIs and not just for the GFP, as the property of domain-specificity would require. The fact that both ROIs showed this feature suggests, firstly, that GFP and MT+ cannot be distinguished on the basis of their response properties in our experiment, and secondly, that the two stimuli induce similar – the cross-condition correlations are well above 0.5 on average (Figure 5, red bars) – yet, distinct HRFs within a brain area known to process also low-level visual motion.

Why did our data from the MT+ ROIs not match the pattern that we predicted based on properties of the motion complex regarding lateralization? We think the reason is that areas MST and FST and others that join area MT in the motion complex may comprise neurons with large receptive fields with no or only weak lateralization[47]. This is why they can be expected to exhibit activity evoked by both the central and the peripheral motion stimuli without much difference between hemispheres. Hence their contribution to the BOLD signals in the two ROIs may most probably have covered the much more specific contribution of area MT. Yet, irrespective of our inability to emphasize area MT, the similarity of the response patterns for both ROIs supports the localization of the GFP within MT+. This is also indicated by the strong correlation between the two ROIs, i.e., testing whether an HRF pair correlated (or not) in one ROI was also correlated (or not) in the other ROI (see Figure 5, scatter plot; Table S1, Q5).

Interestingly, the exploratory group level fits of the HRFs shown in Figure 4 reveal the pattern that was predicted based on lateralized responses independent of the experimental condition. Especially in the left hemisphere, BOLD responses during trials in which the target was on the right side were significantly reduced as compared to responses during trials in which the target was in the middle or on the left side (non-overlapping $CI^{95\%}$). Importantly, this pattern was present in both ROIs (top and bottom row in Figure 4) and for both experimental conditions (green and purple curves in Figure 4). In the right hemisphere the differences were less pronounced ($CI^{95\%}$ do overlap), but met the prediction in that the opposing trend to that observed in the left hemisphere was present. Furthermore, when comparing the group level HRF estimates across ROIs and experimental conditions qualitatively, the similarity between all HRFs is striking.

Particularly interesting are the present results also in the light of studies which have suggested that biological motion is represented as a special category in parts of the temporal cortex[50–53]. While others have already suggested that these findings may merely represent quantitative rather than

qualitative effects[54] our results provide further evidence that it is rather task difficulty and context that

determines the extent of activation of visual motion processing areas and not categorical membership.

We base this argument on the fact that the GFP ROI that was defined through biologically meaningful

motion, i.e. eye-movements, nevertheless exhibited larger β-values in relation to cube-rotations in

experiment 1, but not in experiment 2 in which the stimuli were presented separately (see smaller

cube-rotation related amplitudes in Figure 4). We think that two aspects are capable of explaining this

seemingly contradictory finding, namely visual complexity and the ecological relevance of faces.

Both may create a situation in experiment 1 in which the cube rotation is harder to detect (visual

complexity of the face in the background) or utilize (attentional disengagement from the ecologically

more relevant face), thus requiring a boost of the perceptual signal related to the rotating cubes

relative to the gaze-cue. In contrast, Experiment 2 did not necessitate such a boost, as there was no

competition between the stimuli.

If it is not the posterior temporal cortex that is tuned to socially guided spatial attention, are there

other areas of the brain that are? In an exploratory analysis of experiment 1 we addressed this question

by analyzing differences in the global brain activity associated with the *gaze-shift* and the *cube-*

rotation conditions. To this end, we screened the β-maps for voxels exclusively associated with gaze-

shifts and cube-rotations, respectively. Exclusively means that all voxels displayed in green in Figure

6 were positively correlated with the *gaze-shift* condition on the one hand, and negatively or not

significantly correlated with the *cube-rotation* condition on the other (vice versa for the purple shaded

voxels). In this comparison activity associated with the gaze-shifts was predominantly located in early

visual areas, likely due to the higher visual complexity of the behaviorally relevant stimulus. Some

smaller hotspots were in the temporal cortex (BA 21, 22 and 41) which are functionally associated

with auditory processing and speech perception (based on neurosynth[36] associations), possibly

reflecting the need to process the verbal instructions. But why should there be larger activity evoked

by the instruction to exploit gaze? The reason might be that we as social beings are well trained to use

other peoples' gaze. Hence, the activations associated with the instructions may not be suppressed by

the less demanding gaze cues, yet by the cube cues whose geometrical interpretation may be more

challenging. Figure S2 and S3 provide evidence supporting this explanation, as it shows that in most

participants the *cubes-rotation* related β-values are negative for the two example ROIs considered.

The most remarkable distinction of the activity exclusively related to the *cube-rotation* condition is

the prominent activation of parietal areas (BA 39, 40, 7). This is surprising insofar as the hallmark of

gaze-following is the redirection of visuospatial attention to specific spatial locations, a behavior

known to involve the dorsal attention network (DAN)[55–59]. However, here, our results revealed that

the IPS, a central node of the DAN, is not activated in the *gaze-following* condition, but in the *cube-*

rotation condition, which is identical in behavioral and geometrical terms, the very area is active (see

also Figure S4). Thus, it seems that gaze-following behavior as required in this experiment does not

depend on the dorsal attention network which is at odds with previous findings[60,25,21]. One potential

explanation for the absence of parietal activity might be that while extracting directional information

from the cube-rotation requires some form of mental rotation, gaze-shift following does not. Mental

rotation and viewpoint-dependent object recognition has been reported to involve parietal activity[61,62].

However, this does not explain why parietal activity has been observed in relation to gaze-following

in previous studies[28,63]. A difference between the current paradigm and previous versions may offer an

insight. Namely, while in previous studies participants had to delay their response to the cues until a

dedicated go-signal appeared after a fixed time, in the present version participants were asked to

respond to the cue immediately. Thus, previous designs[28,31] offered a constant period between the cue

and the go-signal in which the planning of the response saccade unfolded which might have triggered

a noticeable modulation of the BOLD signal in the posterior parietal cortex[64,65]. Thus, previously

observed gaze-following related parietal activity may have been related to intentional planning, an

aspect not present in the present experimental design.

In conclusion the findings of the present study force us to abandon previous hypotheses about the

modular processing and the categorical specificity of the gaze of others as a spatial reference[1,3,8].

Rather, we suggest that on the perceptual level core pathways, for example those which process visual

motion, are exploited. Importantly, these conclusions have to be seen in the context of previous

studies that discovered and investigated the GFP[8,28,29, 30,31,32,33,34,38]. Because of the small sample size,

the exploratory analysis of the present study is limited and only serves to illustrate global activity

patterns. However, any other specific brain area that might be an EDD, aside from the GFP, would

have been discovered in any of the previous studies which were designed to capture the full

behavioral relevance of active gaze following. Since the GFP was the only direct candidate neural

substrate for the EDD, and we show here that its response is not specific to gaze stimuli, the broader

conclusion that a domain-specific EDD does not exist is justified. This result is in line with recent

findings which suggest that other perceptual categories, like faces[66,67] or actions [68] have no modular or

domain-specific representation, as well. At other levels of processing, there again appear to be

differences between the processing of others' gaze and geometric objects in terms of spatial cues, as

indicated by the lack of parietal activity in the case of the former. Whether the reason for this is a

more fundamental difference between these two types of cues (see the 'association hypothesis'

suggested by Vecera and Rizzo[69]) or merely a consequence of overtraining to exploit the gaze of

others cannot be answered here.

Materials and Methods

Transparent changes

Because experiment 1 already provided clear evidence regarding the main question of this study (Q6a, see Table S1) we stopped data collection before reaching the evidence threshold set in the registered research plan[37] for Q3.1 & 4.1.

We added group-level fits of the HRFs for experiment 2 not described in the research plan[37] (Figure 4).

The data of participant 6 was not included in the analysis regarding questions 3, 4 and 5 (see Table S1) because for this participant no individual GFP ROI could be delineated. Including this participant's data, however, only had a quantitative effect on the results.

In the research plan[37] we stated that results regarding Q3.1&2 that deviate from the predictions of lateralized responses would invalidate the analysis of experiment 2. This statement was based on the assumption that the motion localizer paradigm would delineate only those areas of the MT+ complex that are strictly lateralized. However, this assumption was not justified because we did not limit the low-level motion stimulus to contralateral stimulation but stimulated the whole visual field. Thus, the delineated ROI arguably comprises neighboring motion sensitive areas like MST and FST with no or the opposite lateralization. In the present manuscript we deleted this restriction, changed Table S1 accordingly and provide an interpretation of the results.

Ethical Approval

The research presented here was approved by the Ethics Board of the Medical Faculty, University of Tübingen, Germany and was conducted in accordance with the principles of human research ethics of the Declaration of Helsinki. Written consent was obtained from all participants, and they were compensated with a lump sum of 40€.

Design

Sampling plan

We employed Bayesian methods for analysis. In line with suggested guidelines for registered reports (e.g.[70]) we included further participants the relevant Bayes Factors reached the threshold for *strong evidence* (≥ 10) in favor of the null- or the alternative hypothesis. Note that throughout this manuscript Bayes Factors are reported as log_{10} transforms, although this is not specified everywhere to improve readability. Furthermore, Bayes Factors are always reported from the perspective of the null hypothesis (BF_{01}), this implies that values above 1 correspond to *strong evidence* in favor of the null hypothesis and values below -1 in favor of the alternative hypothesis. The exclusion of data was

based solely on the observed behavior of the subject during the experiment. We observed the
participants performing the tasks with a camera and recorded their answers, i.e., the direction of their
response saccades, in writing for each trial. This was necessary since eye-tracking quality within the
fMRI scanner is often not stable. We sampled data from 7 right-handed participants, 4 females and 3
males with a mean age of 29.6 $\pm$ 11.2 years. In accordance with the registered sampling plan[37], we
excluded the data of two participants before conducting an analysis because they had incorrect
responses of more than 10% in at least two runs. We excluded the data from one run of another
participant because more than 10 % of the answers were incorrect.

Setup and general procedure

The study consisted of two experimental and one localizer tasks distributed over two separate
fMRI sessions on different days. We acquired MR images using a 3T scanner (Siemens Magnetom
Prisma) with a 20-channel phased array head coil. The head of the subjects was fixed inside the head
coil by using plastic foam cushions to avoid head movements. An AutoAlign sequence was used to
standardize the alignment of images across sessions and subjects. A high-resolution T1-weighted
anatomic scan (MP-RAGE, 176 × 256 × 256 voxel, voxel size 1 × 1×1 mm) and local field
maps were acquired. Functional scans were conducted using a T2*-weighted echo-planar multibanded
2D sequence (multiband factor = 2, TE = 35 ms, TR = 1500 ms, flip angle = 70°) which
covered the whole brain (44 × 64 × 64 voxel, voxel size 3 × 3 × 3 mm, interleaved slice
acquisition, no gap).

Participants were positioned in the scanner and viewed the stimulus monitor that was placed at its
head-end via a mirror system. The distance between the participant and the monitor was ~190 cm and
it covered ~20° of the field of view in the horizontal and ~12° in the vertical. In the session with
experiment 1 participants wore air pressure headphones. The stimuli and data collection was
controlled via a custom software package[71].

Experiment 1

In each trial of Experiment 1 (see Figure 1) participants saw a photograph of the face of a female
with superimposed cubes and a red fixation dot positioned on the nasion. The size of the cubes
matched the size of the irises, and they were arranged so that they are equidistant from the nasion in
the vertical direction as the eyes are in the horizontal direction. Each trial began with a verbal
instruction delivered over headphones during which the face looked towards the participant, her iris
color was grayish, and the cubes pointed straight ahead with one of their vertices. The verbal
instruction was either *Gaze*, *Cubes,* or *Iris* indicating which of the upcoming cues had to be used to
identify the target rectangle among three differently colored rectangles at the bottom of the stimulus
image. After a variable ITI between 2 and 5 s, during which participants had to fixate the central

fixation dot, the face made a gaze shift towards one of the rectangles, her iris color changed to match
one of the target colors and the cubes rotated so that the frontal vertex pointed towards one of the
targets. The participants had to respond to these cues as quickly as possible with a saccade to the
target and maintain the fixation until the start of the next trial. The start of the next trial was indicated
by resetting the cues to their initial state and by the next instruction. Each trial lasted between 5.5 and
8.5 s and participants completed 7 runs of 81 trials each. During each run, targets and instructions
were randomized. Participants were allowed to rest between trials.

Experiment 2

486 In Experiment 2, the face and the cubes were presented separately (*gaze-only*, *cubes-only*). Here,
487 the *Iris-color* condition was omitted (see Figure 1). Therefore, the instruction at the beginning of each
488 trial was not necessary and was omitted. In each run, only one type of cue (*Gaze* or *Cubes*) was
489 presented, but the targets were randomized, and participants completed 4 runs of 81 trials each.
490 Otherwise, the task and the temporal sequence were the same as in Experiment 1.

Motion localizer

To localize the complex of motion processing areas (MT+) participants performed a standard
motion localizer task in which they alternately viewed dot motion and a static dot pattern while
maintaining fixation on a central fixation dot. Both the moving and the static dots were presented 15
times per run for 15 seconds each. In total, participants performed 6 runs. The dot field had a size of 7
deg and consisted of 200 dots with a dot size of 0.1 deg and a lifetime of 500 ms. The lifetimes of
individual dots were asynchronous. During the motion period, individual dots moved at a speed of 1.5
deg/s. In 3 out of the 6 runs the movement direction of individual dots was randomized (noise level of
100%) and in the other 3 runs, the dots moved coherently (noise level of 0%) in a direction randomly
drawn without replacement from a vector specifying 45 possible directions, uniformly sampling 360
deg.

Registered Analysis

503 Table S1 provides an overview of all preregistered questions and related hypotheses, while a
504 detailed description can be found in reference[37]. Some changes were necessary and are reported in the
505 *Transparent Changes* section.

Preprocessing

We used the software package *fMRIPrep* (ver. 20.2.1)[72,73] (RRID:SCR_016216) which is based
on *Nipype* 1.5.1[74] (RRID:SCR_002502) to preprocess the fMRI data. The following sections

describing the preprocessing are boilerplates automatically generated by *fMRIPrep* and inserted
unchanged.

Anatomical data preprocessing

A total of 1 T1-weighted (T1w) images were found within the input BIDS dataset. The T1w
image was corrected for intensity non-uniformity (INU) with `N4BiasFieldCorrection`[75], distributed with
ANTs 2.3.3[76] (RRID:SCR_004757), and used as T1w-reference throughout the workflow. The T1w-
reference was then skull-stripped with a *Nipype* implementation of the `antsBrainExtraction.sh` workflow
(from ANTs), using OASIS30ANTs as target template. Brain tissue segmentation of cerebrospinal
fluid (CSF), white-matter (WM) and gray-matter (GM) was performed on the brain-extracted T1w
using `fast`[77] (FSL 5.0.9, RRID:SCR_002823). Brain surfaces were reconstructed using `recon-all`[78]
(FreeSurfer 6.0.1, RRID:SCR_001847), and the brain mask estimated previously was refined with a
custom variation of the method to reconcile ANTs-derived and FreeSurfer-derived segmentations of
the cortical gray-matter of Mindboggle[79] (RRID:SCR_002438). Volume-based spatial normalization
to one standard space (MNI152NLin2009cAsym) was performed through nonlinear registration with
`antsRegistration` (ANTs 2.3.3), using brain-extracted versions of both T1w reference and the T1w
template. The following template was selected for spatial normalization: *ICBM 152 Nonlinear*
Asymmetrical   template   version   2009c[80]   (RRID:SCR_008796;   TemplateFlow   ID:
MNI152NLin2009cAsym).

Functional data preprocessing

For each of the runs found per subject (across all tasks and sessions), the following preprocessing
will be performed. First, a reference volume and its skull-stripped version were generated by aligning
and averaging 1 single-band references (SBRefs). A B0-nonuniformity map (or *fieldmap*) was
estimated based on a phase-difference map calculated with a dual-echo GRE (gradient-recall echo)
sequence, processed with a custom workflow of *SDCFlows* inspired by the `epidewarp.fsl` script[81] and
further improvements in HCP Pipelines[82]. The *fieldmap* was then co-registered to the target EPI
(echo-planar imaging) reference run and converted to a displacements field map (amenable to
registration tools such as ANTs) with FSL's `fugue` and other *SDCflows* tools. Based on the estimated
susceptibility distortion, a corrected EPI (echo-planar imaging) reference was calculated for a more
accurate co-registration with the anatomical reference. The BOLD reference was then co-registered to
the T1w reference using `bbregister` (FreeSurfer) which implements boundary-based registration[83]. Co-
registration was configured with six degrees of freedom. Head-motion parameters with respect to the
BOLD reference (transformation matrices, and six corresponding rotation and translation parameters)
are estimated before any spatiotemporal filtering using `mcflirt`[84] (FSL 5.0.9). BOLD runs were slice-
time corrected using `3dTshift` from AFNI 20160207[85] (RRID:SCR_005927). First, a reference volume
and its skull-stripped version were generated using a custom methodology of *fMRIPrep*. The BOLD

time-series (including slice-timing correction when applied) were resampled onto their original, native

space by applying a single, composite transform to correct for head-motion and susceptibility

distortions. These resampled BOLD time-series will be referred to as *preprocessed BOLD in original*

space, or just *preprocessed BOLD*. The BOLD time-series were resampled into standard space,

generating a *preprocessed BOLD run in MNI152NLin2009cAsym space*. Several confounding time-

series were calculated based on the *preprocessed BOLD*: framewise displacement (FD), DVARS and

three region-wise global signals. FD was computed using two formulations following Power (absolute

sum of relative motions[86]) and Jenkinson (relative root mean square displacement between affines[84]).

FD and DVARS are calculated for each functional run, both using their implementations in *Nipype*

(following the definitions by Power[86]). The three global signals are extracted within the CSF, the

WM, and the whole-brain masks. Additionally, a set of physiological regressors were extracted to

allow for component-based noise correction (*CompCor*[87]). Principal components are estimated after

high-pass filtering the *preprocessed BOLD* time-series (using a discrete cosine filter with 128s cut-

off) for the two *CompCor* variants: temporal (tCompCor) and anatomical (aCompCor). tCompCor

components are then calculated from the top 2% variable voxels within the brain mask. For

aCompCor, three probabilistic masks (CSF, WM and combined CSF+WM) are generated in

anatomical space. The implementation differs from that of Behzadi et al.[87] in that instead of eroding

the masks by 2 pixels on BOLD space, the aCompCor masks are subtracted a mask of pixels that

likely contain a volume fraction of GM. This mask is obtained by dilating a GM mask extracted from

the FreeSurfer's *aseg* segmentation, and it ensures components are not extracted from voxels

containing a minimal fraction of GM. Finally, these masks are resampled into BOLD space and

binarized by thresholding at 0.99 (as in the original implementation). Components are also calculated

separately within the WM and CSF masks. For each CompCor decomposition, the *k* components with

the largest singular values are retained, such that the retained components' time series are sufficient to

explain 50 percent of variance across the nuisance mask (CSF, WM, combined, or temporal). The

remaining components are dropped from consideration. The head-motion estimates calculated in the

correction step were also placed within the corresponding confounds file. The confound time series

derived from head motion estimates and global signals were expanded with the inclusion of temporal

derivatives and quadratic terms for each[88]. Frames that exceeded a threshold of 0.5 mm FD or 1.5

standardised DVARS were annotated as motion outliers. All resamplings can be performed with *a*

single interpolation step by composing all the pertinent transformations (i.e., head-motion transform

matrices, susceptibility distortion correction when available, and co-registrations to anatomical and

output spaces). Gridded (volumetric) resamplings were performed using `antsApplyTransforms` (ANTs),

configured with Lanczos interpolation to minimize the smoothing effects of other kernels[89]. Non-

gridded (surface) resamplings were performed using `mri_vol2surf` (FreeSurfer).

Copyright Waiver

The above boilerplate text was automatically generated by *fMRIPrep* with the express intention
that users should copy and paste this text into their manuscripts *unchanged*. It is released under the
CC0 license.

Run- and first-level GLMs, ROI definition

For each participant we computed GLMs (general linear models) for each run individually (run-
level) as well as a GLM for the combination of all runs (first-level) to obtain the respective β-images
for each condition using the Python package *nilearn*[90]. To mitigate the effects of motion artifacts and
other noise sources, the nuisance regressors *global_signal*, *csf*, white_matter, *trans_x*, *trans_y*,
trans_z, *rot_x*, *rot_y*, *rot_z* and their respective first derivatives estimated by *fMRIPrep* were included
in the design matrices. As a model for the *HRF* we used the *glover + derivative + dispersion* model
provided by *nilearn* and included a polynomial drift model of order 3 to remove slow drifts in the
data. Further, we masked the data with each run's mask image (or the average for the first-level GLM)
provided by *fMRIPrep* and applied smoothing with *fwhm* = 5 mm.

To define the GFP and the MT+ ROI the resulting first-level β-images were used to compute the
contrasts *gaze-shift* minus *iris-color* (experiment 1) and *motion* minus *static* for the localizer task,
respectively. For each participant these ROIs were determined individually by applying the most
liberal statistical threshold that allowed to differentiate a separable activity patch at the location of the
reference coordinates. These are for the GFP $(x = -55, \ y = -68, \ z = 1)$ and $(x = 50, \ y = -64,$
$z = -3)$ [31] and for MT+ $(x = -45, \ y = 70, \ z = 3)$ and $(x = 46, \ y = -67, \ z = 4)$. The latter
coordinates are the center of the output of a neurosynth.org database [36] search for *visual motion*. All
coordinates reported in this study refer to the MNI space. The GFP-ROI was not identifiable for one
participant in the way described above. For this one we constructed a spherical ROI with a radius of 8
mm located at the individual local maximum closest to the reference coordinates.

Question 1/ 2: Comparison of gaze-shift and cube-rotation related activity in the left and right GFP

We extracted each participant's run-level *gaze-shift* and *cube-rotation* related β-values from the
GFP and averaged across voxels. Thus, we obtained 6 values per hemisphere and participant for each
condition. These were fed into a Bayesian hierarchical regression model with *run* as group-level
intercepts and slopes, and condition (*Question 1*) or hemisphere (*Question 2*) as the population-level
effect. To avoid model fitting problems, *participant* was not added as a group-level effect becausethe
number of participants was 5 as it is recommended by different sources (e.g. Bolker et al.[91]).
Modeling was carried out in R[92] using the *brms*[93] package. Priors for the generative model were tuned
to yield reasonable prior and posterior predictions for the pilot data (reported in reference[37]). As
marginal prior for the effect of condition/ hemisphere on the β-values we used normal distribution

with $\mu = 0$, $\sigma = 1$. The *R* code of the model and the priors are reported in the Supplementary

Materials section. The model provided an effect estimate, denoted as $\Delta\beta_{GFP-left}^{Gaze-Cubes}$. A positive value

indicates that the gaze-shift related activity is larger than the cube-rotation related activity, while a

negative value indicates the opposite. Bayes Factors were computed via the Savage-Dickey density

ratio evaluating the ration between the posterior and the prior at the parameter value of zero[94] and

log_{10} transformed. The terminology used in this study to describe the amount of evidence is adopted

from Wetzels et al[95].

Question 3-5: HRF estimation and correlation analysis

For each participant and ROI, we estimated the *HRF* for each condition and target of experiment

2. This yielded 24 HRFs estimates in total for each participant (2 ROIs × 2 hemispheres × 2

conditions × 3 targets). To do so, the mean BOLD signals from the ROIs was extracted and linear

deconvolution was applied to disentangle the responses to the different conditions/ targets. Prior to

signal extractions, the BOLD images were denoised using the same confounds as for the GLMs (see

above), smoothed with *fwhm* = 5 mm, and standardized. For the deconvolution, we resorted to the

nideconv package[96] and used 9 *fourier* regressors as the basis set which allows a larger flexibility than

the canonical HRF while being less complex than a FIR model.

For each ROI, we computed linear correlation coefficients (employing the function

correlationBF[97]) between all relevant pairs (see Table 3) of the responses to the left, middle and right

target within and across the conditions of experiment 2. In total, we obtained the correlation

coefficients of 34 HRF-pairs for each participant and ROI. These were grouped by the expected

correlation if the responses to the second motion cue in the end of trials were lateralized (color code

of the circles in Table 3) and by within- and cross-condition pairings (color code of the dots in Table

3). Figure 7 shows an example. These groups were analyzed for differences using Bayesian

hierarchical models. The correlation coefficients were modeled as gaussian distributions within the

limit of possible values of correlation coefficients – between -1 and 1. Priors for the generative model

were tuned to yield reasonable prior and posterior predictions for the pilot data (see reference[37]). As

marginal priors for the effect of the respective groupings we used a normal distribution with $\mu = 0$

and $\sigma = 2$. The first model provided the effect estimate as $\Delta\rho_{GFP}^{low-high}$ which is negative if the

prediction based on lateralization is met. The second model will provide the effect estimate as

$\Delta\rho_{GFP}^{within-cross}$ which is positive if the prediction based on condition is met. Bayes Factors were

computed via the Savage-Dickey method and log_{10} transformed.

Finally, if the correlation of a pair of HRFs is high in one ROI, we tested whether it is then also

high in the other ROI (Q5). We did so by computing the correlation between the ROIs and the Bayes

Factor testing against no correlation using the *correlationBF* function.

Figure 7 Illustration for Q3 and Q4. Shown are 4 example HRFs from one pilot subject to illustrate the analysis and predictions of experiment 2. In the illustrated gaze-only trial the participant looked at the right target such that the second motion event occurred in her left visual hemifield triggering a larger and longer lasting response in the right hemisphere compared to the left hemisphere (green curves). In the illustrated cube-only trial the participant fixated the left target leading to the opposite pattern (purple curves). Computing the correlations between the responses yields results that are in line with the blue and yellow predictions, indicating that the low-level motion components in the stimuli trigger lateralized responses. The shaded areas represent the 95% CI of the HRF estimates. See Table 3 for a full list of the predictions for all relevant response pairs.

Table 3 Predictions of magnitudes of correlations. GO = gaze-only, CO = Cubes-only; l – left target, m – middle target and r – right target. Correlations coefficients will be computed for all marked pairs. For pairs marked by blue circles correlation coefficients are expected to be higher than for those marked by yellow circles if responses are lateralized and triggered by low-level motion components (Q3.1/ 4.1). If, however, responses reflect the conditions (Q3.2/ 4.2), then correlation coefficients for pairs marked by dark gray dots will be greater than those marked by red dots. Target-2 trials (middle) are only paired with itself to consistently maintain lateralization with respect to the left and right targets. The analysis will be carried out on data of the GFP and MT+ (ROI = [GFP, MT]). Left and right refers to the hemisphere.

Exploratory Analysis

To explore brain wide differences between *gaze-shift* and *cubes-rotation* following related activity we computed second level □-maps for the two conditions based on the non-smoothed first level □-maps from experiment 1. The resulting second level map was smoothed using a 8 mm kernel. To assess qualitative as well as quantitative differences between activity related to the conditions in a first step using the thresholded □-maps (, uncorrected, minimum cluster size of 15 voxels) we created masks that excluded all significantly positively correlated voxels. In the second step we applied to the thresholded □-maps of respective other condition. This means we removed all voxels that are positively correlated with the *cube-rotation* condition from the □-map of the *gaze-shift* condition and vice versa. We applied a cluster threshold of 15 voxels to the results once more to partially account for the very liberal alpha threshold used here. This allowed us to create □-maps that contain only genuinely activated areas for the two conditions (Figure 6).

Acknowledgements

We thank Dr. Friedemann Bunjes for his invaluable technical assistance.

We acknowledge support from the German Research Foundation (DFG) project TH425/17-1 and from the Open Access Publication Fund of the University of Tübingen.

Data Availability Plan

Unprocessed data organized in accordance with the BIDS format is publicly available at doi:10.18112/openneuro.ds005903.v1.0.1.

Any code used for analysis is publicly available at https://github.com/maalaria/fMRIus/tree/gaze-motion.

References

1. 689 Baron-Cohen, S. How to build a baby that can read minds: Cognitive mechanisms in mind

690 reading. *Curr. Psychol. Cogn.* **13**, 513–552 (1994).

2. 691 Perrett, D. I. & Emery, N. J. Understanding the intentions of others from visual signals:

692 neuropsychological evidence. *Cah. Psychol. Cogn.* **13**, 683–694 (1994).

3. 693 Baron-Cohen, S. The Empathizing System: a revision of the 1994 model of the Mindreading

694 System. (2005).

4. 695 Fodor, J. A. *The Modularity of Mind: An Essay on Faculty Psychology*. (MIT Press, 1983).

5. 696 Kanwisher, N., McDermott, J. & Chun, M. M. The fusiform face area: a module in human

697 extrastriate cortex specialized for face perception. *J. Neurosci.* **17**, 4302–11 (1997).

6. 698 Batki, A., Baron-Cohen, S., Wheelwright, S., Connellan, J. & Ahluwalia, J. Is there an innate

699 gaze module? Evidence from human neonates. *Infant Behavior and Development* **23**, 223–229

700 (2000).

7. 701 Deen, B., Koldewyn, K., Kanwisher, N. & Saxe, R. Functional Organization of Social Perception

702 and Cognition in the Superior Temporal Sulcus. *Cereb. Cortex* **25**, 4596–4609 (2015).

8. 703 Ramezanpour, H. & Thier, P. Decoding of the other's focus of attention by a temporal cortex

704 module. *PNAS* **117**, 2663–2670 (2020).

9. 705 Panksepp, J. & Panksepp, J. The seven sins of evolutionary psychology. *Evolution and Cognition*

706 **6**, (2000).

10. 707 De Waal, F. B. M. Evolutionary Psychology: The Wheat and the Chaff. *Curr Dir Psychol Sci* **11**,

708 187–191 (2002).

11. 709 Westlin, C. *et al.* Improving the study of brain-behavior relationships by revisiting basic

710 assumptions. *Trends in Cognitive Sciences* **27**, 246–257 (2023).

12. 711 Emery, N. J. The eyes have it: The neuroethology, function and evolution of social gaze.

712 *Neuroscience & Biobehavioral Reviews* **24**, 581–604 (2000).

13. 713 Zeiträg, C., Jensen, T. R. & Osvath, M. Gaze following: A socio-cognitive skill rooted in deep

time. *Front Psychol* **13**, 950935 (2022).

14. Driver, J. *et al.* Gaze perception triggers reflexive visuospatial orienting. *Vis. cogn.* **6**, 509–540

(1999).

15. Langton, S. R. H. & Bruce, V. Reflexive visual orienting in response to the social attention of

others. *Visual Cognition* **6**, 541–567 (1999).

16. Langton, S. R. H. & Bruce, V. You must see the point: Automatic processing of cues to the

direction of social attention. *J. Exp. Psychol. Hum. Percept. Perform.* **26**, 747–757 (2000).

17. Deaner, R. O. & Platt, M. L. Reflexive Social Attention in Monkeys and Humans. *Current*

Biology **13**, 1609–1613 (2003).

18. Friesen, C. K., Ristic, J. & Kingstone, A. Attentional Effects of Counterpredictive Gaze and

Arrow Cues. **30**, 319–329 (2004).

19. Marciniak, K., Dicke, P. W. & Thier, P. Monkeys head-gaze following is fast, precise and not

fully suppressible. *J. Cogn. Neurosci.* **20**, 108–119 (2008).

20. Perrett, D. I. *et al.* Visual Cells in the Temporal Cortex Sensitive to Face View and Gaze

Direction. *Proc. R. Soc. B Biol. Sci.* **223**, 293–317 (1985).

21. Hoffman, E. A. & Haxby, J. V. Distinct representations of eye gaze and identity in the distributed

human neural system for face perception. *Nat. Neurosci.* **3**, 80–84 (2000).

22. Heywood, C. A. & Cowey, A. The role of the 'face-cell' area in the discrimination and

recognition of faces by monkeys. *Philos Trans R Soc Lond B Biol Sci* **335**, 31–37; discussion 37-

38 (1992).

23. Akiyama, T. *et al.* Gaze but not arrows: A dissociative impairment after right superior temporal

gyrus damage. *Neuropsychologia* **44**, 1804–1810 (2006).

24. Kingstone, A., Friesen, C. K. & Gazzaniga, M. S. Reflexive joint attention depends on lateralized

cortical connections. *Psychol Sci* **11**, 159–166 (2000).

25. Nummenmaa, L. & Calder, A. J. Neural mechanisms of social attention. *Trends Cogn. Sci.* **13**,

135–143 (2009).

26. Itier, R. J. & Batty, M. Neural bases of eye and gaze processing: The core of social cognition.

Neuroscience & Biobehavioral Reviews **33**, 843–863 (2009).

27. Hadders-Algra, M. Human face and gaze perception is highly context specific and involves

bottom-up and top-down neural processing. *Neuroscience & Biobehavioral Reviews* **132**, 304–

323 (2022).

28. Materna, S., Dicke, P. W. & Thier, P. Dissociable Roles of the Superior Temporal Sulcus and the

Intraparietal Sulcus in Joint Attention: A Functional Magnetic Resonance Imaging Study. *J.*

Cogn. Neurosci. **20**, 108–119 (2008).

29. Materna, S., Dicke, P. W. & Thier, P. The posterior superior temporal sulcus is involved in social

communication not specific for the eyes. *Neuropsychologia* **46**, 2759–2765 (2008).

30. Laube, I., Kamphuis, S., Dicke, P. W. & Thier, P. Cortical processing of head- and eye-gaze cues

guiding joint social attention. *Neuroimage* **54**, 1643–1653 (2011).

31. Marquardt, K., Ramezanpour, H., Dicke, P. W. & Thier, P. Following Eye Gaze Activates a Patch

in the Posterior Temporal Cortex That Is Not Part of the Human 'Face Patch' System. *eNeuro* **4**,

1–10 (2017).

32. Kraemer, P. M., Görner, M., Ramezanpour, H., Dicke, P. W. & Thier, P. Frontal, Parietal, and

Temporal Brain Areas Are Differentially Activated When Disambiguating Potential Objects of

Joint Attention. *eNeuro* **7**, (2020).

33. Kamphuis, S., Dicke, P. W. & Thier, P. Neuronal substrates of gaze following in monkeys. *Eur.*

J. Neurosci. **29**, 1732–1738 (2009).

34. Marciniak, K., Atabaki, A., Dicke, P. W. & Thier, P. Disparate substrates for head gaze following

and face perception in the monkey superior temporal sulcus. *Elife* **3**, 1–18 (2014).

35. Tsao, D. Y., Moeller, S. & Freiwald, W. A. Comparing face patch systems in macaques and

humans. *Proc. Natl. Acad. Sci.* **105**, 19514–19519 (2008).

36. Neurosynth. https://neurosynth.org/.

37. Görner, M., Dicke, P. W. & Thier, P. Is there a brain area dedicated to socially guided spatial

attention? *OSF* **registration (https://archive.org/details/osf-registrations-7kjyg-v1)**, (2023).

38. Chong, I., Ramezanpour, H. & Thier, P. Causal manipulation of gaze-following in the macaque

temporal cortex. *Progress in Neurobiology* **226**, 102466 (2023).

39. Cosmides, L. & Tooby, J. Origins of domain specificity: The evolution of functional

organization. in *Mapping the Mind: Domain Specificity in Cognition and Culture* (eds.

Hirschfeld, L. A. & Gelman, S. A.) 85–116 (Cambridge University Press, Cambridge, 1994).

doi:10.1017/CBO9780511752902.005.

40. Saxe, R. & Baron-Cohen, S. Editorial: The neuroscience of theory of mind. *Social Neuroscience*

1, 1–9 (2006).

41. Hadjikhani, N., Kveraga, K., Naik, P. & Ahlfors, S. P. Early (M170) activation of face-specific

cortex by face-like objects. *Neuroreport* **20**, 403–407 (2009).

42. Takahashi, K. & Watanabe, K. Gaze cueing by pareidolia faces. *Iperception* **4**, 490–492 (2013).

43. Zhou, L.-F. & Meng, M. Do you see the "face"? Individual differences in face pareidolia. *Journal*

of Pacific Rim Psychology **14**, e2 (2020).

44. Görner, M., Ramezanpour, H., Dicke, P. & Thier, P. Gaze and Arrows: does the Gaze Following

Patch in the posterior temporal cortex differentiate social and symbolic spatial cues? *eNeuro*

(2024) doi:10.1523/ENEURO.0065-24.2024.

45. Watson, J. D. G. *et al.* Area V5 of the Human Brain: Evidence from a Combined Study Using

Positron Emission Tomography and Magnetic Resonance Imaging. *Cereb Cortex* **3**, 79–94

(1993).

46. Tootell, R. *et al.* Functional analysis of human MT and related visual cortical areas using

magnetic resonance imaging. *J. Neurosci.* **15**, 3215–3230 (1995).

47. Huk, A. C., Dougherty, R. F. & Heeger, D. J. Retinotopy and Functional Subdivision of Human

Areas MT and MST. *J. Neurosci.* **22**, 7195–7205 (2002).

48. Born, R. T. & Bradley, D. C. STRUCTURE AND FUNCTION OF VISUAL AREA MT. *Annu.*

Rev. Neurosci. **28**, 157–189 (2005).

49. Bulletin, P., Friesen, C. K. & Kingstone, A. The eyes have it! Reflexive orienting is triggered by

nonpredictive gaze. *Psychon. Bull. Rev.* **5**, 490–495 (1998).

50. Oram, M. W. & Perrett, D. I. Responses of Anterior Superior Temporal Polysensory (STPa)

Neurons to 'Biological Motion' Stimuli. *J Cogn Neurosci* **6**, 99–116 (1994).

51. Grossman, E. *et al.* Brain areas involved in perception of biological motion. *J Cogn Neurosci* **12**,

711–720 (2000).

52. Servos, P. *et al.* The neural substrates of biological motion perception: an fMRI study. *Cereb.*

cortex **12**, 772–782 (2002).

53. Puce, A. & Perrett, D. Electrophysiology and brain imaging of biological motion. *Philos Trans R*

Soc Lond B Biol Sci **358**, 435–445 (2003).

54. Grossman, E. D. & Blake, R. Brain activity evoked by inverted and imagined biological motion.

Vision Res. **41**, 1475–1482 (2001).

55. Butterworth, G. & Jarrett, N. What minds have in common is space: Spatial mechanisms serving

joint visual attention in infancy. *British journal of developmental psychology* **9**, 55–72 (1991).

56. Moll, H. & Tomasello, M. 12- and 18-month-old infants follow gaze to spaces behind barriers.

Developmental Science **7**, F1–F9 (2004).

57. Todorović, D. Geometrical basis of perception of gaze direction. *Vision Res.* **46**, 3549–3562

(2006).

58. Corbetta, M. & Shulman, G. L. Control of goal-directed and stimulus-driven attention in the

brain. *Nature Reviews Neuroscience* **3**, 201–215 (2002).

59. Corbetta, M., Kincade, J. M., Ollinger, J. M., McAvoy, M. P. & Shulman, G. L. Voluntary

orienting is dissociated from target detection in human posterior parietal cortex. *Nat Neurosci* **3**,

292–297 (2000).

60. Calder, A. J. *et al.* Separate Coding of Different Gaze Directions in the Superior Temporal Sulcus

and Inferior Parietal Lobule. *Current Biology* **17**, 20–25 (2007).

61. Alivisatos, B. & Petrides, M. Functional activation of the human brain during mental rotation.

Neuropsychologia **35**, 111–118 (1997).

62. Gauthier, I. *et al.* BOLD Activity during Mental Rotation and Viewpoint-Dependent Object

Recognition. *Neuron* **34**, 161–171 (2002).

63. Shepherd, S. V., Klein, J. T., Deaner, R. O., Platt, M. L. & Shepard, R. N. Mirroring of attention

by neurons in macaque parietal cortex. *Proc. Natl. Acad. Sci.* **106**, 9489–9494 (2009).

64. Stetson, C. & Andersen, R. A. Early planning activity in frontal and parietal cortex in a simplified

task. *J Neurophysiol* **113**, 3915–3922 (2015).

65. Andersen, R. A., Aflalo, T. & Kellis, S. From thought to action: The brain-machine interface in

posterior parietal cortex. *Proc Natl Acad Sci U S A* **116**, 26274–26279 (2019).

66. Vinken, K., Prince, J. S., Konkle, T. & Livingstone, M. S. The neural code for "face cells" is not

face-specific. *Science Advances* **9**, (2023).

67. Shi, Y. *et al.* Rapid, concerted switching of the neural code in inferotemporal cortex. *bioRxiv*

2023.12.06.570341 (2023) doi:10.1101/2023.12.06.570341.

68. Karakose-Akbiyik, S., Caramazza, A. & Wurm, M. F. A shared neural code for the physics of

actions and object events. *Nat Commun* **14**, 3316 (2023).

69. Vecera, S. P. & Rizzo, M. Eye gaze does not produce reflexive shifts of attention: Evidence from

frontal-lobe damage. *Neuropsychologia* **44**, 150–159 (2006).

70. Registered Reports | Nature. https://www.nature.com/nature/for-authors/registered-reports.

71. Bunjes, F. & Gukelberger, J. open source measurement system nrec. (2022).

72. Esteban, O. *et al.* fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nat Methods* **16**,

111–116 (2019).

73. Esteban, O. *et al.* fMRIPrep: a robust preprocessing pipeline for functional MRI. Zenodo

https://doi.org/10.5281/zenodo.7430291 (2022).

74. Gorgolewski, K. *et al.* Nipype: A Flexible, Lightweight and Extensible Neuroimaging Data

Processing Framework in Python. *Frontiers in Neuroinformatics* **5**, (2011).

75. Tustison, N. J. *et al.* N4ITK: Improved N3 Bias Correction. *IEEE Transactions on Medical*

Imaging **29**, 1310–1320 (2010).

76. Avants, B. B., Epstein, C. L., Grossman, M. & Gee, J. C. Symmetric diffeomorphic image

registration with cross-correlation: Evaluating automated labeling of elderly and

neurodegenerative brain. *Medical Image Analysis* **12**, 26–41 (2008).

77. Zhang, Y., Brady, M. & Smith, S. Segmentation of brain MR images through a hidden Markov

random field model and the expectation-maximization algorithm. *IEEE Transactions on Medical*

Imaging **20**, 45–57 (2001).

78. Dale, A. M., Fischl, B. & Sereno, M. I. Cortical Surface-Based Analysis: I. Segmentation and

Surface Reconstruction. *NeuroImage* **9**, 179–194 (1999).

79. Klein, A. *et al.* Mindboggling morphometry of human brains. *PLOS Computational Biology* **13**,

e1005350 (2017).

80. Fonov, V., Evans, A., McKinstry, R., Almli, C. & Collins, D. Unbiased nonlinear average age-

appropriate brain templates from birth to adulthood. *NeuroImage* **47**, S102 (2009).

81. Epidewarp.

82. Glasser, M. F. *et al.* The minimal preprocessing pipelines for the Human Connectome Project.

NeuroImage **80**, 105–124 (2013).

83. Greve, D. N. & Fischl, B. Accurate and robust brain image alignment using boundary-based

registration. *NeuroImage* **48**, 63–72 (2009).

84. Jenkinson, M., Bannister, P., Brady, M. & Smith, S. Improved Optimization for the Robust and

Accurate Linear Registration and Motion Correction of Brain Images. *NeuroImage* **17**, 825–841

(2002).

85. Cox, R. W. & Hyde, J. S. Software tools for analysis and visualization of fMRI data. *NMR in*

Biomedicine **10**, 171–178 (1997).

86. Power, J. D. *et al.* Methods to detect, characterize, and remove motion artifact in resting state

fMRI. *NeuroImage* **84**, 320–341 (2014).

87. Behzadi, Y., Restom, K., Liau, J. & Liu, T. T. A component based noise correction method

(CompCor) for BOLD and perfusion based fMRI. *NeuroImage* **37**, 90–101 (2007).

88. Satterthwaite, T. D. *et al.* An improved framework for confound regression and filtering for

control of motion artifact in the preprocessing of resting-state functional connectivity data.

NeuroImage **64**, 240–256 (2013).

89. Lanczos, C. Evaluation of Noisy Data. *Journal of the Society for Industrial and Applied*

Mathematics Series B Numerical Analysis **1**, 76–85 (1964).

90. Nilearn, contributors. Nilearn: Statistics for NeuroImaging in Python.

91. Bolker, B. *GLMM FAQ*. http://bbolker.github.io/mixedmodels-misc/glmmFAQ.html.

92. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for

Statistical Computing (2021).

93. Bürkner, P.-C. brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal of*

Statistical Software **80**, 1–28 (2017).

94. Faulkenberry, T. J. Savage-Dickey Density Ratio for computing Bayes Factors. *The Book of*

Statistical Proofs https://statproofbook.github.io/P/bf-sddr.html (2020).

95. Wetzels, R., van Ravenzwaaij, D. & Wagenmakersq, E.-J. Bayesian Analysis. in *The*

Encyclopedia of Clinical Psychology 1–11 (John Wiley & Sons, Ltd, 2015).

96. Hollander, G. de & Knapen, T. nideconv. (2017).

97. Morey, R. D. *et al.* BayesFactor. (2022).

Supplementary material

 Table S1

Question	Null Hypothesis	Outcome Measures	Sampling plan	Analysis Plan	Interpretation of outcomes
Q1 Is the GFP-ROI differentially activated during *Gaze* and *Cubes* in experiment 1?	$H_0^{Q1.1}: \Delta\beta_{GFP-left}^{Gaze-Cubes} = 0$ $H_0^{Q1.2}: \Delta\beta_{GFP-right}^{Gaze-Cubes} = 0$	Beta estimates of BOLD signal	$BF_{01/10}^{Q1} \geq 10$	Bayesian hierarchical model	**1.** $\Delta\beta_{GFP}^{Gaze-Cubes} \leq 0$: no difference between *Gaze* and *Cubes* related activity or less *Gaze* related activity in the GFP. Support against the hypothesis that the GFP is a specialized module for gaze-following. **2** $\Delta\beta_{GFP}^{Gaze-Cubes} > 0$: The GFP is stronger activated during *Gaze* trials than during *Cubes* trials. Either the GFP is a specialized gaze-following module or it is a subpart of MT+ reflecting retinotopic locations. These possibilities are disentangled in Q6.
Q2 Is there a hemispheric dominance for *Gaze* or *Cubes*?	$H_0^{Q2.1}: \Delta\beta_{Gaze}^{RH-LH} = 0$ $H_0^{Q2.2}: \Delta\beta_{Cubes}^{RH-LH} = 0$	Beta estimates of BOLD signal	$BF_{01/10}^{Q2} \geq 10$	Bayesian hierarchical model	**1.** $\Delta\beta_{Gaze/Cubes}^{RH-LH} = 0$: no difference between the left and right hemisphere. **2.** $\Delta\beta_{Gaze/Cubes}^{RH-LH} > 0$: The right hemisphere is stronger activated than the left hemisphere. Complies with the right hemispheric dominance of visual face processing. **3.** $\Delta\beta_{Gaze/Cubes}^{RH-LH} < 0$: The left hemisphere is stronger activated than the right hemisphere.
Q3.1 Does target ID in conjunction with hemisphere affect the level of correlations in MT+.	$H_0^{Q3.1}: \Delta\rho_{MT+}^{low-high} = 0$	Correlation coefficients of HRF estimates of BOLD signals	$BF_{01/10}^{Q3.1} \geq 10$	Bayesian hierarchical model	**1.** $\Delta\rho_{MT+}^{low-high} = 0$: no lateralized response in MT+. ROI does not only contain araes with contralateral receptive fields or the peripheral motion cue is not of relevance. **2.** $\Delta\rho_{MT+}^{low-high} < 0$: the result matches the prediction based on lateralization. **3.** $\Delta\rho_{MT+}^{low-high} > 0$: reverse effect as predicted (implausible). Invalidates analysis of experiment 2.

Q3.2 Does condition affect the level of correlation in MT+. Q3.1 and 3.2 are positive controls for the second part of the analysis.	$H_0^{Q3.2}: \Delta\rho_{MT+}^{within-across} = 0$	Correlation coefficients between HRF estimates of BOLD signals	$BF_{01/10}^{Q3.2} \geq 10$	Bayesian hierarchical model	**1.** $\Delta\rho_{MT+}^{within-across} = 0$: correlations within conditions and across conditions are equally strong. Matches the prediction for a visual motion area. **2.** $\Delta\rho_{MT+}^{within-across} < 0$: correlations within conditions smaller than across conditions (implausible). **3.** $\Delta\rho_{MT+}^{within-across} > 0$: correlations within conditions larger than across conditions. Indicates a differential representation of the conditions within a domain-general brain area.
Q4.1 Does target ID in conjunction with hemisphere affect the level of correlations in the GFP	$H_0^{Q4.1}: \Delta\rho_{GFP}^{low-high} = 0$	Correlation coefficients between HRF estimates of BOLD signals	$BF_{01/10}^{Q4.1} \geq 10$	Bayesian hierarchical model	**1.** $\Delta\rho_{GFP}^{low-high} = 0$: no lateralized response in GFP. Given that $\Delta\rho_{MT+}^{low-high} < 0$, we will conclude that the GFP is functionally different from MT+. See Q6 for full interpretation. **2.** $\Delta\rho_{GFP}^{low-high} < 0$: the result matches the prediction based on lateralization. Given that $\Delta\rho_{MT+}^{low} < 0$, we will conclude that the GFP and MT+ are functionally homologous. See Q6 for full interpretation. **3.** $\Delta\rho_{GFP}^{low-high} > 0$: reverse effect as prediction (implausible). See Q6 for full interpretation.
Q4.2 Does condition affect the level of correlation in the GFP	$H_0^{Q4.2}: \Delta\rho_{GFP}^{within-across} = 0$	Correlation coefficients of HRF estimates of BOLD signals	$BF_{01/10}^{Q4.2} \geq 10$	Bayesian hierarchical model	**1.** $\Delta\rho_{GFP}^{within-across} = 0$: both conditions elicit the same responses in GFP. Indicating that it is not a specialized module for the processing of gaze. See Q6 for full interpretation. **2.** $\Delta\rho_{GFP}^{within-across} < 0$: lower correlation within- than across-conditions (implausible). **3.** $\Delta\rho_{GFP}^{within-across} > 0$: across-conditions pairs are less correlated than within-condition pairs. Supporting that the GFP is a specialized module if $\Delta\rho_{MT+}^{within-across} \leq 0$. See

					Q6 for full interpretation.
Q5 are the correlation coefficients from the GFP correlated with those from MT+	$H_0^{Q4}: \rho_{MT+/GFP} = 0$	Correlation coefficients	$BF_{01/10}^{Q5} \geq 10$	Bayesian correlation test	**1.** $\rho_{MT+/GFP} = 0$: no correlation between MT+ and GFP. GFP is functionally segregated from MT+. **2.** $\rho_{MT+/GFP} > 0$: GFP is (part of) MT+ **3.** $\rho_{MT+/GFP} < 0$: GFP and MT+ are anticorrelated (implausible).
Q6 Is the GFP (a) a specialized module for the processing of gaze and (b) functionally segregated from MT+?	-	-	-	Combination of Q1,3,4,5	In order to conclude that the GFP is a specialized module the following results must occur: (a)  Q1: $\Delta\beta_{GFP}^{Gaze-Cubes} > 0$ (b)  Q3.1: $\Delta\rho_{MT+}^{low-high} < 0$  Q3.2: $\Delta\rho_{MT+}^{within-across} = 0$  Q4.1: $\Delta\rho_{GFP}^{low-high} = 0$  Q4.2: $\Delta\rho_{GFP}^{within-across} > 0$  Q5: $\rho_{MT+/GFP} = 0$
Q7 Is the GFP identical with or a part of MT+?	-	-	-	Combination of Q1,3,4,5,6	In order to conclude that the relevant stimulus component triggering activity in the GFP is motion and that it cannot be distinguished from MT+ the following results must occur:  Q1: $\Delta\beta_{GFP}^{Gaze-Cubes} \leq 0$  Q3.1/ 4.1:

					$\Delta\rho_{MT+}^{low-high} < 0 > \Delta\rho_{GFP}^{low-high}$
					Q3.2/ 4.2:
					$\Delta\rho_{MT+}^{within-across} = 0 = \Delta\rho_{GFP}^{within-across}$
					Q5: $\rho_{MT+/GFP} > 0$

891 *Model specification*

892 **Table S2.** *R* code in *brms* syntax for the models and the priors.

	Formula	Priors
Q1/2	$\beta \sim x + (x \mid \text{run})$, family=gaussian(link="identity"), x = {condition, hemisphere}	normal(0, 5), class='Intercept' normal(0, 1), class='b', coef={'conditionGaze', hemisphereRH} normal(.5, .5), class='sd', coef='Intercept', group='run' normal(.3, .5), class='sd', coef='conditionGaze', group='run' lognormal(.5, .5), class='sigma'
Q3/4	$r \mid \text{trunc(lb=-1, ub=1)} \sim x$, family='gaussian', x = {lateralization, condition}	normal(0, 1), class='Intercept' normal(0, 2), class='b' lognormal(0, .5), class='sigma'

895 **Figure S1** Outlines of the GFP (green) and MT+ (blue) ROIs for each participant.

Language ROIs

Figure S2 Broca area *β*-values extracted from a ROI representing the Broca area. For all participants the values associated with the cube-rotation condition are smaller, and negative, than those associated with the gaze-shift condition. I.e. the activation of language areas does not reflect an activation of language areas as a simple BOLD contrast between the *gaze-shift* and *cube-rotation* conditions would suggest.

Figure S3 Middle temporal gyrus Here the picture is less clear than for the Broca area. Subject 03 shows positive *β*-values associated with the *gaze-shift* condition while subject 05 shows negative *β*-values for the *cube-rotation* condition. The reason for this discrepancy might be the chosen ROIs are not actually representing comparable brain areas. For example it might be that the ROI used for subject 03 coincides with face-responsive areas in the temporal cortex while the ROI for subject 05 does not.

910   **Figure S4 Intraparietal sulcus** The ROIs were defined based on the contrast *gaze-shift following* minus *cube-rotation*

911   *following*. The *ß*-values related to the *gaze-shifts* (and *iris-color mapping*) are on average not different from 0.