Task-specific patterns of odorant receptor expression in worker antennae indicates a sensory filter regulating division of labor in ants

Marcel A. Caminer[1]*, Romain Libbrecht[1,2], Megha Majoe[1], David V. Ho[3], Peter Baumann[3,4], Susanne Foitzik[1]

[1]Institute of Organismic and Molecular Evolution, Johannes Gutenberg University Mainz, Germany

[2]Institut de Recherche sur la Biologie de l'Insecte, UMR 7261, CNRS, University of Tours, France

[3]Institute of Developmental and Neurobiology, Johannes Gutenberg University Mainz, Germany

[4]Institute of Molecular Biology, Mainz, Germany

*Corresponding author

E-mail: marcelcaminer@gmail.com (MAC)

Abstract

Division of labor (DOL) is a characteristic trait of insect societies, where tasks are generally performed by groups of specialized individuals. In social insects, young workers perform duties within the safety of the nest (e.g., brood care), while older ones undertake riskier tasks (e.g., foraging for food). This DOL remains dynamic, and workers may switch back and forth when colony needs require. Theoretical models propose that workers differ in their thresholds to take on certain tasks when confronted to task-related stimuli, resulting in variation in their response to such stimuli, task-specialization, and thus DOL. Such models assume that workers differ in how they respond to task-related information rather than in how they perceive such information. Here, we test the hypothesis that DOL rather stems from workers differing in their efficiency to detect task-related stimuli. We used transcriptomics to compare gene expression in the antennae and in the brain between nurses and foragers in the ant *Temnothorax longispinosus*. We found that seven times as many genes were differentially expressed between the behavioral phenotypes in the antennae compared to the brain. Moreover, nearly half of all odorant receptors genes were differentially expressed, with an overrepresentation of the 9-exon gene subfamily upregulated in the antennae of nurses. These findings suggest that nurses and foragers differ in how they perceive their olfactory environment, and task-related signals. The results of this study support the hypothesis that a sensory filter in the antennae

42 predisposes workers to specialize in specific tasks, and may improve our understanding

43 of DOL in insect societies.

45 **Keywords:** social insects | transcriptomics | behavioral specialization | odorant

46 perception

48 Introduction

Division of labor (DOL) is an important organizing principle of complex biological systems

that arose independently during three of the major evolutionary transitions [1]. DOL was

originally formulated in the context of the production process in human societies [2], but

specialization to specific tasks is also found within cells and across many organisms [1].

Examples range from bacteria, where clonal populations are divided into subpopulations

focusing on different activities [3–6] to multicellular organisms with differentiation of cells

into different tissues and organs, and individuals performing specific roles in animal

societies [7,8]. To understand the evolution of complex life it is therefore essential to

investigate the mechanisms that underlie DOL.

DOL in insect societies results from individuals specializing in the performance of

specific tasks. In addition to the reproductive DOL between fertile queens and functionally

sterile workers, there is a behavioral DOL among workers that specialize in tasks such as

brood care, foraging, nest building, and defense [9–11]. Several factors can affect task

specialization, including age [12,13], nutrition [14,15], morphology [16], genotype [17–19],

experience [20], and colony size [21–23]. In most social insect species, workers display

an age polyethism by which younger individuals tend to perform intranidal tasks, while

older individuals are more likely to undertake activities such as nest-defense and foraging

outside the nest [24,25]. Yet, such specialization among workers remains flexible, as

foragers can revert to perform brood care when needed [26,27].

Several molecular mechanisms have been implicated in the regulation of DOL.

Task specialization is associated with transcriptional changes in the worker brain [28–33].

Molecular pathways such as the insulin/insulin-like signaling (IIS), vitellogenin (Vg) and

juvenile hormone (JH) pathways are involved in the regulation of worker behavior [34–

39]. Functional manipulations have confirmed that the expression of key genes in the

worker brain controls task specialization [40–42]. Behavioral variation among workers is

also associated with signaling of biogenic amines (e.g., dopamine, octopamine, tyramine,

serotonin), which act as neurotransmitters or neuromodulators involved in the modulation

of the responsiveness to task-associated stimuli [43–46].

The self-organization and collective behavior in insect societies are maintained via

the exchange of chemical information [47]. Social insects communicate primarily through

glandular pheromones and complex mixtures of long-chain hydrocarbons on their cuticle.

These cuticular hydrocarbons (CHC) facilitate recognition of nestmates, developmental

stages, castes, sexes, and species [48,49]. Social insects perceive chemical information

via different types of sensilla on their antennae [50–52]. Decoding the identities of

chemical compounds relies on odorant receptors (OR) located within each sensillum

[53,54]. OR are transmembrane proteins expressed in the dendrites of olfactory receptor

neurons (ORN). The largely conserved OR coreceptor (Orco) of insects [55] is required

for odorant recognition in the dendritic membrane: it forms an ion channel with specific

OR, which determines the sensitivity and specificity of the ORN [56]. Odorant molecules

penetrate through the antenna cuticular pores and are transported by odorant-binding

proteins to the ORN membrane, where they interact with receptors, leading to the

generation of action potentials [57,58]. ORN axons relay signals from the sensilla to the

glomeruli of the antennal lobes in the insect brain, which are the first processing unit for

olfactory information. Then, ORN make synaptic contact with the projection neurons and

local neurons, which transfer information to the central brain [59–61].

Several lines of evidence indicate that OR genes play central roles in the regulation

of social life of insects. First, social insect species typically harbor large numbers of OR

genes [62–65]. Second, some OR gene families have specifically expanded during social

evolution, such as the 9-exon subfamily in ants, which appears to serve an important

function in the perception of CHC [64,66–70]. Third, species that evolved socially parasitic

strategies resulting in reduced behavioral repertoires show a strong and convergent

reduction in the number of OR genes [71]. Finally, experimentally produced mutant ants

that lack the *Orco* gene (coding for the co-receptor necessary for OR to properly function)

show impaired social behavior [72,73].

Current DOL models bring together the chemical nature of social insect

communication and variation among workers in their response to chemical cues. They

posit that flexible response thresholds to task-related chemical stimuli serve as regulators

of worker specialization [74]. Workers take on a particular task when the stimulus intensity

exceeds their individual threshold for this task. Therefore, individuals with lower

thresholds for a given task are more likely to perform it than those with higher thresholds

[75]. Individual response decisions and task performance are modulated via numerous

parameters on different time scales [7,76–78], and despite extensive research on DOL

and task allocation, these mechanisms are not fully understood.

As indicated by the name, "response threshold models" typically assume that

individuals differ in their response to specific signals, not necessarily in their ability or

efficiency to detect such signals. This implies that such thresholds and the associated

responses are set in the central nervous system, possibly via molecular pathways in the

brain that correlate with behavioral variation [74,79,80]. We propose that DOL models

would benefit from considering odor sensitivity as a potential upstream sensory filter that

may affect task specialization. Along these lines, we hypothesize that behavioral variation

among workers may also stem from their ability and/or efficiency in detecting different

signals. For example, we propose that individuals that specialize in brood care do so

because they are better at detecting brood cues, rather than (or in addition to) being more

likely to respond to similar levels of brood cues. To test this hypothesis that a sensory

filter regulates inter-individual behavioral variation, and thus the DOL in social insects, we

investigated transcriptional differences in the brain and antennae of workers that

specialize in either brood care or foraging behavior in the ant *Temnothorax longispinosus*.

We found that i) behavioral variation was associated with important transcriptomic

changes in the antennae, ii) these changes included a large proportion of the OR gene

repertoire, and iii) individuals specializing in brood care overexpressed an OR gene family

putatively involved in detecting social cues. These findings support our hypothesis that

the peripheral nervous system, acting as a sensory filter, plays an important role in

regulating behavioral differences between workers and thus in the DOL of social insects.

Results

Identification of *Temnothorax longispinosus* behavioral phenotypes

Task specialization in the ant *T. longispinosus* is neither genetically fixed nor rigid, but changes with age and in response to colony needs [38]. We conducted behavioral observations of seven *T. longispinosus* laboratory colonies to identify individuals that specialize in brood care behavior (hereafter referred to as nurses), and others that specialize in foraging (hereafter referred to as foragers) (Fig 1*A*; see Methods section for more details). Individuals identified as nurses interacted with the brood in 54% ± 22% (mean ± sd) of the observations, and were recorded outside the nest in 1% ± 3% of the observations. On the contrary, foragers were found outside the nest in 42% ± 26% of the observations, but interacted with the brood in only 2% ± 4% of the observations.

Larger task-associated transcriptomic changes in the antennae than in the brain

To investigate transcriptomic variation between nurses and foragers, we used RNA-seq to generate seven nurse and seven forager brain and antenna samples, each consisting of pooled tissue from seven workers of a single colony. Of the 14,837 protein-coding genes annotated in the *T. longispinosus* genome, we found that 91% (13,494) and 92% (13,683) were expressed (FPKM > 0) in the brain and the antennae, respectively. Using an adjusted p-value of 0.05 as significance threshold, we detected 333 differentially

expressed genes (DEG) in the brain (217 upregulated in nurses, 116 in foragers), and 2,254 in the antennae (1,234 upregulated in nurses, 1,020 in foragers; Dataset S1; Fig 1*B*). We found an overlap of 162 DEG between the two tissues, including 143 DEG that showed differences in the same direction across tissues. Principal component analysis (PCA) plots indicate that both brain and antenna samples cluster by behavioral phenotype rather than the colony of origin (Fig 1 *C-D*).

OR gene expression differs between antennae of nurses and foragers

To investigate whether odorant perception differs between nurses and foragers, we focused our attention on the expression of OR genes in the antennae. We found that all 419 OR genes in the genome of *T. longispinosus* [71] were expressed in the antennae, and that 48% (202/419) of them were differentially expressed between nurses and foragers. Specifically, 65 OR genes were upregulated in nurses (15% of all OR genes), and 137 in foragers (32% of all ORs) (Fig 2). Then, we studied which OR subfamilies were preferentially expressed in nurses and foragers. The 65 OR genes overexpressed in nurses belonged to three OR subfamilies, while the 137 OR genes upregulated in foragers were distributed among 19 OR subfamilies. Foragers overexpressed 26, 8, and 8 OR genes from the L, P, and H subfamilies, respectively, while no OR genes from these subfamilies were upregulated in nurses (Dataset S2). We found that 61% (26/43) of the genes from the L subfamily were overexpressed in foragers, which represents a significant overrepresentation (Fisher's test, odds ratio = 3.63, p-value < 0.001). For the P and H subfamilies we did not find such an overrepresentation, likely due to lower gene

numbers (P subfamily: Fisher's test, odds ratio = 3.62, p-value = 0.06; H subfamily: Fisher's test, odds ratio = 1.87, p-value = 0.19). On the other hand, we found that 82% (53/65) of the OR genes overexpressed in nurses belong to the 9-exon subfamily. This results in an overrepresentation of the 9-exon subfamily in genes that were overexpressed in nurses (Fisher's test, odds ratio = 20.98, p-value < 0.001), with 46% (53/114) of this subfamily being overexpressed in nurses. In contrast, only 6% (8/137) of the 9-exon subfamily was overexpressed in foragers, which is less than expected by chance (Fisher's test, odds ratio = 0.10, p-value < 0.001). We also found that nurses overexpressed *Orco* compared to foragers (FDR p-value = 0.001).

Genes in biogenic amine pathways vary in expression between nurses and foragers

Because biogenic amines are implicated in the regulation of behavior, and they may affect sensory perception [45,81–83], we screened our lists of DEG in the brain and antennae for genes in biogenic amine pathways. We found five genes associated with biogenic amine signaling in the antennae, and none in the brain. Genes in the serotonin (*5-hydroxytryptamine*, DBV15_11483), tyramine (*tyramine beta-hydroxylase*, DBV15_00422) and octopamine (*octopamine receptor* LOC112465659) pathways were upregulated in foragers, while nurses showed a higher expression of genes in the dopamine (*dopamine 1-like receptor 2*, DBV15_07611) and octopamine (*octopamine beta2 receptor*, DBV15_10418) pathways (Fig 3 *A-E*).

Association between behavioral variation and molecular pathways in brain and antennae

Our analysis revealed expression differences between the brains of nurses and foragers for genes involved in the regulation of task specialization in social insects. Previous studies have found that *Vg* genes and the associated JH and IIS/TOR pathways are important endocrine networks that play central roles in the regulation of lifespan, fertility and behavior in bees and ants [35,84–89]. Therefore, we searched our RNA-seq data for task-associated expression of genes involved in the metabolism, biosynthesis, and regulation of these pathways in the brain and antennae. We found evidence that multiple genes from these pathways are differentially expressed between nurses and foragers (Table S1). Among the most interesting genes upregulated in the brain of nurses were *conventional Vg* (LOC112466671), *Vg-like A* (named/classified as per [42]; DBV15_03138), *venom carboxylesterase-6-like* (DBV15_11528), and *protein takeout* (DBV15_08771), while *allatostatin A-like* (LOC112454443) and *insulin-like growth factor I* (*IGF1*) (LOC112454447) were overexpressed in the brain of foragers. Remarkably, *venom carboxylesterase-6-like* and *IGF1* were overexpressed in the antennae of nurses and foragers respectively. Additionally, high expression levels of the zinc-finger transcription factor *Krüppel homolog-1* (*Kr-h1*) were detected in the antennae of foragers (DBV15_06330) (Fig 3*F*). The expression of this gene has been correlated with caste and behavioral differences in the brain of social insects [86,90–93], but little is known about its function and gene targets in the antennae.

We performed a GO enrichment analysis to gain a deeper understanding of the biological processes represented in the lists of DEG. This analysis detected many

enrichments based on single one or few genes, and we mention below a few interesting

processes with the number of genes driving the enrichment in parentheses (see Table S2

for complete list). Genes that were upregulated in the brain of nurses were enriched for

biological processes such as *translation* (19 genes), *cellular iron ion homeostasis* (2

genes), and *catabolic process* (6 genes); while *translation* (22 genes), *endocytosis* (6

genes), *glucose metabolic process* (4 genes), and *regulation of cell cycle* (3 genes) were

enriched in the antennae. In the list of genes that were overexpressed in the brain of

foragers, we found that enriched processes included *peptide metabolic process* (4

genes), *methionyl-tRNA aminoacylation* (1 gene), *positive regulation of type I interferon*

production (1 gene), and *cell redox homeostasis* (1 gene); while *phosphatidylinositol*

phosphorylation (4 genes), *inositol phosphate dephosphorylation* (3 genes), *carbohydrate*

metabolic process (17 genes), *innate immune response* (2 genes), and *nucleotide*

catabolic process (3 genes) were enriched in the antennae.

Discussion

In this study, we tested the hypothesis that variation among social insect workers in their ability to detect different chemical signals regulates task specialization, and thus DOL in insect societies. To do so, we compared the brain and antenna transcriptomes of nurses and foragers in the ant *T. longispinosus.* We report several lines of evidence that support our hypothesis. First, we found almost seven times as many genes to be differentially expressed between nurses and foragers in the antennae as in the brain, indicating that peripheral sensory organs may be more important than the central nervous system in the task specialization process of social insect workers. Second, we found half the OR genes in the *T. longispinosus* genomes are differentially expressed between nurse and forager antennae, suggesting that behavioral specialization is associated with different sensory filters that result in specific perceptions of the chemical environment. Third, our analyses revealed that nurses and foragers upregulated distinct families of OR genes, indicating that their sensory filters may target different types of chemical cues, possibly adjusted to the tasks they perform. Finally, we detected several genes in multiple biogenic pathways to be differentially expressed between the brains of nurses and foragers, potentially involved in fine-tuning the sensitivity of the odorant filters.

Many organisms have evolved sensory filters to focus on only a subset of all environmental cues. In insects, the peripheral olfactory filtering system enables individuals to detect and discriminate odors that convey ecologically relevant information used to mediate important behaviors such as courtship, locomotion and navigation to avoid predators and locate food or nesting sites [94]. For example, the mosquito *Anopheles gambiae* strongly responds to odorant components of its vertebrate hosts that

provide meals [95]. Sensory filters have been selected for because they limit the amount

of information perceived by the organism, and thus the energy and time required by the

brain to process it. Within insect colonies, individuals typically exhibit many morphological

and physiological traits associated with increased efficiency in task specialization. They

can be understood as adaptations that allow individuals to become more proficient at their

task due to learning, training, or the perceiving valuable task-related information, allowing

the colony to avoid the cost of task switching [96]. In this context, it is interesting to find

that *T. longispinosus* workers may have different sensory filters that would serve as a

basis for their task specialization, as workers would mostly perceive chemical cues that

pertain to their tasks. Our findings indicate that the sensory filter of ant workers is

dynamic, and its changes may underlie their behavioral maturation.

Age, genetic background, social environment, and individual experience produce

behavioral variation among workers that results in division of labor [13,20,23,97]. In our

study, age in particular likely differed between nurses and foragers. However, previous

experiments designed to disentangle gene expression associated with behavioral

specialization, age and fertility showed that behavioral specialization is more strongly

associated with gene expression than age and fertility in *T. longispinosus* [38]. We

propose that the factors known to regulate task specialization could drive different OR

expression patterns, which in turn would produce behavioral variation via contrasting

abilities to detect different sets of odors. This hypothesis is supported by several studies

(reviewed in [98]) showing that the physiological condition of an animal can influence the

level of receptor expression, including mating status, oviposition, feeding, circadian

rhythm, experience, and aging. Alternatively, we cannot exclude that the exposure to

different odors that is associated with performing different tasks may at least in part have

affected OR gene expression in the antennae. However, such an effect is unlikely to

explain the large-scale variation in gene expression, as there is very limited evidence that

the mere exposure to odors influences the expression of the gene coding for the OR that

binds this odor [99,100].

According to our sensory filter hypothesis, ant workers would be expected to

primarily detect chemical cues that correspond to their tasks. We found that the 9-exon

subfamily of OR was overrepresented in genes that were upregulated in the antennae of

nurses. Interestingly, recent studies have shown a rapid expansion of the 9-exon

subfamily in ants, and several lines of evidence indicate that OR genes from this subfamily

mediate complex social interactions in ant colonies [65–67,69]. First, comparisons of

antennal transcriptomes revealed that 9-exon OR genes are expressed more frequently

in workers than in males, suggesting a role in social communication among workers

[68,69]. Second, representative OR from the 9-exon subfamily can detect CHC extracts

from several castes [70]. According to McKenzie et al. [69], 9-exon OR genes were first

expressed in solitary ancestors of aculeate wasps and facilitated CHC discrimination,

likely for prey or mate recognition, with a lineage giving rise to the ancestors of ants.

Moreover, OR genes of the 9-exon subfamily were convergently lost in socially parasitic

ant species that lost the ability to perform brood care or foraging [71], suggesting that they

are essential for the performance of these worker tasks. Our finding of an overexpression

of 9-exon OR genes in the antennae of nurses is in line with these previous studies, and

suggests that many of these receptors have important functions within the nest, such as

sensing chemical cues from the larvae, queen, or other workers, and/or that they are less

important for sensing task-related stimuli or other signals outside the nest. Among the OR

genes overexpressed in nurses was also the co-receptor *Orco*, which is widely expressed

in olfactory sensory neurons and nearly unchanged in sequence in distant insect taxa

[101–103]. The overexpression of *Orco* may indicate a generally more active odorant

sensory system in the antennae of nurses compared to foragers.

The behavioral transition from nursing to foraging may be triggered by a lower

efficiency in detecting brood cues via the downregulation of specific OR genes (e.g., from

the 9-exon subfamily). This would result in ants moving farther away from the brood, and

this change in spatial location may trigger the behavioral transition to outside tasks

[104,105]. In addition to being less efficient at detecting brood cues, the sensory filter of

foragers may also become fine-tuned to detect a more diverse set of odors. Foragers

overexpressed a greater number of OR gene subfamilies compared to nurses (19 and 3

for foragers and nurses, respectively), which may indicate that the olfactory system of

foragers could be adjusted to the more diverse chemical environment outside the nest.

Similar to the 9-exon subfamily, the L subfamily has also been expanded in social insects

[62,64,65], and along with the P and H subfamilies, it has been lost in socially parasitic

ants [71]. Interestingly, the OR genes from the L, P, and H subfamilies have been

upregulated in foragers, and thus may have a task-specific function, such as recognition

of chemical cues related to environmental perception or recruitment cues outside the

nest. To date, however, there is no information on which odors are recognized by these

OR gene subfamilies.

Finding task-specific variation in OR gene expression raises the question as to

which molecular mechanisms regulate those changes. Previous studies in insects

revealed that olfaction-guided behavior is mediated by biogenic amine receptors in the

antenna, and their expression is involved in fine-tuning the sensitivity of the olfactory

system [82,98]. For example, modified concentration cAMP and intracellular Ca^{2+} levels

due to octopamine-induced signal transduction in the moth *Manduca sexta* [106] activate

Orco, leading to changes in ORN sensitivity [107,108]. Our results suggest that the

biogenic amine signaling pathway may modulate the sensory filtering function of insect

antennae and alter sensitivity to various signals. We found that genes encoding tyramine

and its precursor, octopamine, are upregulated in forager antennae, similar to genes

involved in serotonin signaling. Tyramine and dopamine (which was upregulated in

nurses) have been implicated in modulating taste and olfactory receptor neurons, while

serotonin may serve as a neurotransmitter and neurohormone in antennal vessels and

mechanosensory organs [82]. Serotonin influences foraging activity [81] and regulates

food intake in many animals [109–111]. Dopamine signaling also plays an important role

in controlling the insect circadian clock and mediating clock-controlled behavioral

phenotypes such as locomotion [112,113]. In our focal species *T. longispinosus*, inside

workers were found to exhibit a stronger circadian rhythmicity than foragers, which may

be regulated via differences in the acetylation of histone proteins [114].

Changes in behavior and olfactory sensitivity in insects could be related to the

expression of genes involved in IIS, target of rapamycin (TOR), JH and Vg pathways,

according to age, circadian rhythm, mating and feeding status [98]. For example, appetite

state in *D. melanogaster* is signaled by insulin, which upregulates a peptide receptor on

the olfactory receptor cells that innervate the DM1 glomerulus. Activation of the DM1

glomerulus is enough to drive the fly to reach for food [115]. Recent studies have shown

that pheromone release and odor sensitivity appear to be under JH control in

Schistocerca gregaria and *Locusta migratoria*, which could lead to behavioral changes

[116–120]. Finally, an experimental downregulation of *Vg-like A* in *T. longispinosus*

workers resulted in decreased brood care behavior and a lower sensitivity to brood-

related chemical cues, suggesting changes in odor perception and olfactory-driven

decision making [42]. Our results reveal that genes associated with all these pathways

were differentially expressed in the brain and antennae between nurses and foragers,

predicting a link between their role in task-associated behavioral changes and the

regulation of odor perception. Given the central role of IIS, Vg, JH, and TOR pathways in

regulating division of labor in social insects, and our finding of task-associated patterns

of the antennal expression of genes from multiple biogenic amines, we hypothesize that

these modulators and hormones could be involved in the regulation of the olfactory filter.

How the detailed molecular mechanisms of these pathways in the brain are causally

linked to the complex changes in olfactory perception in the antennae should therefore

be investigated next.

Conclusion

Our transcriptomic analyses of the brain and antennae of *T. longispinosus* nurses and

foragers provide support to our hypothesis that behavioral variation and task

specialization in ant workers are regulated via differences in olfactory perception. We

predict that antennal physiology acts as sensory filters that limit the type and amount of

chemical information passed to the brain. This would allow workers to target relevant

chemical information from the environment and discriminate signal from noise without

using energetically costly processing by the central nervous system. We argue that this

sensory filter is flexible and regulated through changes in physiological conditions such

as age, nutrition, and hormones. Variation among workers in their efficiency to detect

specific chemical cues would result in task specialization and division of labor. Our study

opens novel avenues of research to better understand the role of sensory filters in

controlling DOL in insect societies.

Materials and Methods

Sample collection and behavioral determination

A total of seven colonies of the ant *T. longispinosus* were selected with an average colony size of 110 ± 31.5 workers (mean ± SD, Dataset S3). The ants were collected in the forests of the Edmund Niles Huyck Preserve, Rensselllearville, NY, USA (42°31′41.0″N 74°09′38.8″W), in June of 2018 with permission. Upon collection, we housed each colony in a plaster-floored nesting box (43 cm × 28 cm × 10 cm) divided into three chambers containing a single slide nest, in which the colony relocated. A slide nest is an artificial nesting site comprised of a small Plexiglas cavity sandwiched between two glass microscope slides. Colonies were established at the Johannes Gutenberg University in Mainz, Germany, under a 14 h:10 h light:dark photoperiod at 18°C to a 22°C temperature. We provided honey and water ad libitum and fed crickets to the colony twice a week. To allow for visible behavioral division of labor between workers of the two behavioral phenotypes, we marked, observed and recaptured ants from inside and outside the nest. We defined foragers as workers that perform outside-nest tasks, including foraging for food, while nurses remained inside the dark nest and cared for the ant brood. A total of 69 workers inside (from the brood pile) and 76 workers outside the nest were marked with fine colored metal wires (0.02 mm Elektrisola, Eckenhagen, Germany) between the petiole and post-petiole. We performed behavioral observations every two hours, four times a day for five days (total = 20 scans), in which we noted down how many times an individual spent performing brood care and foraging behavior and the position in the nest (Table S3). Based on these behavioral observations, the marked individuals found outside the nest, exploring the surroundings for food or water, were identified as foragers.

These workers usually do not care for the brood and do not frequently reside on brood

piles, as ant colonies organize themselves spatially in a way that reduces contact

between foragers and brood [121]. We identified nurses as workers that remain inside the

nest in direct contact with brood and were unlikely to leave the nest. Foragers were found

in 42% ± SD 26% of the observations outside of the nest, whereas nurses spend only 1%

± SD 3% outside. In contrast, nurses were interacting with the brood in 54% ± SD 22% of

the observations, whereas we found that foragers only did this only in 2% ± SD 4% of the

observations. We scanned the behavior of workers over 20 observations, albeit earlier

studies have shown that a single observation allows to group *T. longispinosus* and other

ants reliably into nurses and foragers that differ in behavior [121,122], gene expression

[38] and CHC composition [42]. Furthermore, spatial location can alone can predict

behavior in *Temnothorax* workers [123]. We focused in this study on individuals highly

specialized on either foraging or brood care. Workers that performed both tasks regularly

were not included in this study. After all observations were completed, the marked nurses

and foragers were collected, directly frozen in liquid nitrogen and stored at -80 °C until

further processing for dissection and pooling according to behavioral state and colony.

RNA extraction and sequencing

For RNA extraction, we removed both antennae and stored them in a 1.5 ml Eppendorf

tube containing 50 µl TRIzol (Invitrogen), cut the head off and fixed it on a slide with

melted dental wax. We then made an incision around the head with a surgical scalpel and

removed the head capsule with forceps to expose the intact brain. Finally, we carefully

pulled the brain out of the head capsule and removed the remains of other tissues that

were connected to it. The dissected brain was transferred to a 1.5 ml Eppendorf tube

containing 20µl PBS. Each dissection was completed in less than 5 minutes to prevent

RNA degradation. We dissected brain and antennae tissues from 48 nurses and 49

foragers. We pooled the brains and antennae from seven workers from each behavioral

state and colony. The only exception was the "GO" colony (NY18 E110), for which we

pooled only six brains and twelve antennae from six nurses (Dataset S3) due to the loss

of one sample during the dissection process. Immediately after dissection of each brain

and antennae, the Eppendorf tubes were kept on dry ice while we dissected the remaining

individuals. Brain and antennae tissues were homogenized with a pestle. Sample brains

were transferred separately to a 1.5 ml Eppendorf tube containing 50µl of TRIzol. We

added 50µl chloroform to each brain and antenna samples, cautiously inverted for 30 s

and then centrifuged samples at 12,000 g for 15 min at 4°C. We collected the resulting

supernatant and precipitated RNA with 25µl 70% ethanol. We conducted the subsequent

RNA extraction with the RNeasy Mini Kit (Qiagen), following the manufacturer's

instruction. The resulting 28 samples (14 brains and 14 antennae) were stored at -80°C

until library preparation.

RNA-seq libraries were prepared by Novogene Company Limited, Cambridge, UK,

using the NEBNext Ultra RNA Library Prep Kit for Illumina according to the manufacturer's

protocol. After amplification and purification, 28 libraries were sequenced on an Illumina

NovaSeq 6000 S4 flow cell platform using a paired-end 150 bp. Approximately 43 million

raw reads were generated from each library (Dataset S3).

Gene expression analyses

Raw data obtained from Novogene were checked using FastQC v.0.11.9 [124], and

Illumina adapters were removed using Trimmomatic v.0.36 [125]. The protein-coding

genes of *T. longispinosus* (GCA_004794745.1; [126] and the congener *T. curvispinosus*

(GCA_003070985.1) were retrieved from the NCBI database and transferred to the

recently published *T. longispinosus* genome [71] using the Liftoff v.1.6.1 tool [127]. In total,

10,029 of 13,061 (~77%) annotated protein-coding genes were transferred from the

original *T. longispinosus* assembly (genes identified as "DBV15") and 4,808 were

transferred from *T. curvispinosus* (genes identified as "LOC"), for a total of 14,837. For

gene expression analysis, reads were mapped to our *T. longispinosus* genome assembly,

and the read counts table was generated using STAR 2.7.0 [128] with default settings.

Detailed mapping statistics for each sample is available in Dataset S3. We used the

deseq2 v1.16.1 package to identify differentially expressed genes [129]. To avoid biased

results due to low read counts, we removed from the counting matrix those genes for

which less than 10 of the reads mapped to at least 6 of our 14 samples (n - 1 of the

smallest sample size). Then, we conducted a differential gene expression analysis with

DESeq2 [130]. We began with comparisons between nurses and foragers using the

~Colony+Task model, followed by a likelihood ratio test (LRT) approach, with colony ID

as a fixed factor. Genes were considered differentially expressed if the false discovery

rate (FDR), using Benjamini-Hochberg procedure, had an adjusted p-value of ≤ 0.05. The

resulting lists of DEG refer to genes that are overexpressed and underexpressed in

foragers compared to nurses. We used the online tool Venny v.2.1

(https://bioinfogp.cnb.csic.es/tools/venny) to generate a Venn diagram containing the

DEG associated with task and tissues. Separation of differentially expressed genes by

task was visualized by performing principal component analysis (PCA) on transformed

reads of filtered transcriptomes from all contigs using the plotPCA function provided by

DESeq2. Finally, ORs genes that were upregulated in each behavioral phenotype was

visualized in a volcano plot using the EnhancedVolcano package in R.

Identification of behavior candidate genes and odorant receptors

We used gene annotations based on a BlastX search of the *T. longispinosus*

transcriptome compared to a list of different invertebrate proteomes (i.e., *Acromyrmex*

echinatior, Apis mellifera, Camponotus floridanus, Drosophila melanogaster,

Harpegnathos saltator, Odontomachus brunneus, Temnothorax curvispinosus)

downloaded from the NCBI database with an E-value of 1e-5 and below. Clusters

containing more than one sequence match per species were reduced to a single

specimen based on the highest blast score. We constructed orthogroups across all of the

above species using OrthoFinder [131], including amino acid sequences from the *T.*

longispinosus proteome [126], and retained orthogroups containing caste DEG (Dataset

S4) to again compare potential behavioral candidate genes previously identified as

involved in regulating the division of labor in social insects [38,41,87,93,120,132–135].

GO enrichment analysis was performed with TopGO v.2.44.0 using a Fisher's exact test

for the different gene sets compared to the whole genome with the weight01 algorithm

[136]. Only annotated GO terms with an adjusted p-value of ≤ 0.05 were considered

significantly enriched. All enrichment analyses were performed in RStudio v.1.4.1106

[137].

Odorant receptor (OR) protein sets were clustered across multiple ant species

using OrthoFinder to derive orthologous groups and identify subfamilies for each OR in

T. longispinosus. To associate orthogroups with previously identified OR subfamilies, we

used OR annotation in *Atta cephalotes, Acromyrmex echinatior* from Engsontia et al. [65],

and *Camponotus floridanus*, *Harpegnathos saltator*, and *Solenopsis invicta* from Zhou et

al. [64,68]. Missing subfamily information was labeled as "unassigned" (Dataset S2).

Acknowledgments

We thank Diego Páez-Moscoso for many helpful discussions and comments throughout

the study; Marah Stoldt and Carlotta Martelli for providing helpful comments on the

manuscript; and Marion Kever and Jenny Fuchs for technical assistance. This project was

funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)

– GRK2526/1 – Projectnr. 407023052.

Data. RNAseq data that support the findings of this study have been deposited in

GenBank with the BioProject accession codes PRJNA926589

(http://www.ncbi.nlm.nih.gov/bioproject/926589).

References

1. Maynard Smith J, Szathmáry E. The major evolutionary transitions in evolution. Oxford University Press, New York; 1995.
2. Smith A. An Inquiry into the Nature and Causes of the Wealth of Nations. Univ Chicago Press. 1776;1:1092.
3. Veening JW, Igoshin OA, Eijlander RT, Nijland R, Hamoen LW, Kuipers OP. Transient heterogeneity in extracellular protease production by *Bacillus subtilis*. Mol Syst Biol. 2008;4(184):1–15.
4. Marlow VL, Cianfanelli FR, Porter M, Cairns LS, Kim Dale J, Stanley-Wall NR. The prevalence and origin of exoproteaseproducing cells in the *Bacillus subtilis* biofilm. Microbiol (United Kingdom). 2014;160:56–66.
5. van Gestel J, Vlamakis H, Kolter R. From Cell Differentiation to Cell Collectives: *Bacillus subtilis* Uses Division of Labor to Migrate. PLoS Biol. 2015;13(4):1–29.
6. West SA, Cooper GA. Division of labour in microorganisms: An evolutionary perspective. Nat Rev Microbiol. 2016;14(11):716–23.
7. Oster GF, Wilson EO. Caste and ecology in the social insects. Acta Biotheor. 1979;28(3):234–5. Available from: https://doi.org/10.1007/BF00046355
8. Wilson EO. The ergonomics of caste in the social insects. Am Nat. 1968;102(923):41–66.
9. Wilson EO. Behavioral discretization and the number of castes in an ant species. Behav Ecol Sociobiol. 1976;1(2):141–54.
10. Herbers JM. Social Social Organization in *Leptothorax* ants: within-and between-species patterns. Psyche J. Entomol. 1983;90:52489. Available from: https://doi.org/10.1155/1983/52489
11. Pinter-Wollman N, Hubler J, Holley JA, Franks NR, Dornhaus A. How is activity distributed among and within tasks in *Temnothorax ants*? Behav Ecol Sociobiol. 2012;66(10):1407–20.
12. Seeley TD. Adaptive significance of the age polyethism schedule in honeybee colonies. Behav Ecol Sociobiol. 1982;11(4):287–93.
13. Tripet F, Nonacs P. Foraging for work and age-based polyethism: the roles of age and previous experience on task choice in ants. Ethology. 2004;110(11):863–77.
14. Blanchard GB, Orledge GM, Reynolds SE, Franks NR. Division of labour and seasonality in the ant *Leptothorax albipennis*: worker corpulence and its influence on behaviour. Anim Behav. 2000;59(4):723–38.
15. Toth AL, Robinson GE. Worker nutrition and division of labour in honeybees. Anim Behav. 2005;69(2):427–35.
16. Wetterer JK. The ecology and evolution of worker size-distribution in leaf-cutting ants (Hymenoptera: Formicidae). Sociobiology. 1999;34(1):119–44.
17. Fewell JH, Page RE. Genotypic variation in foraging responses to environmental stimuli by honey bees, *Apis mellifera*. Experientia. 1993;49(12):1106–12.
18. Graham S, Myerscough MR, Jones JC, Oldroyd BP. Modelling the role of intracolonial genetic diversity on regulation of brood temperature in honey bee (*Apis mellifera* L.) colonies. Insectes Soc. 2006;53(2):226–32.
19. Gove R, Hayworth M, Chhetri M, Rueppell O. Division of labour and social insect

colony performance in relation to task and mating number under two alternative
response threshold models. Insectes Soc. 2009;56(3):319–31.

20. Ravary F, Lecoutey E, Kaminski G, Châline N, Jaisson P. Individual experience
alone can generate lasting division of labor in ants. Curr Biol. 2007;17(15):1308–
12.

21. Gautrais J, Theraulaz G, Deneubourg JL, Anderson C. Emergent polyethism as a
consequence of increased colony size in insect societies. J Theor Biol.
2002;215(3):363–73.

22. Merkle D, Middendorf M. Dynamic polyethism and competition for tasks in
threshold reinforcement models of social insects. Adapt Behav. 2004;12(3–
4):251–62.

23. Jeanson R, Fewell JH, Gorelick R, Bertram SM. Emergence of increased division
of labor as a function of group size. Behav Ecol Sociobiol. 2007;62(2):289–98.

24. Wakano JY, Nakata K, Yamamura N. Dynamic model of optimal age polyethism
in social insects under stable and fluctuating environments. J Theor Biol.
1998;193(1):153–65.

25. Iwasa Y, Yamaguchi S. Task allocation in a cooperative society: specialized
castes or age-dependent switching among ant workers. Sci Rep. 2020;10(1):1–9.

26. Robinson EJH, Feinerman O, Franks NR. Flexible task allocation and the
organization of work in ants. Proc R Soc B Biol Sci. 2009;276(1677):4373–80.

27. Shimoji H, Kasutani N, Ogawa S, Hojo MK. Worker propensity affects flexible task
reversion in an ant. Behav Ecol Sociobiol. 2020;74(7).

28. Ben-Shahar Y, Robichon A, Sokolowski MB, Robinson GE. Influence of gene
action across different time scales on behavior. Science. 2002;296(5568):741–4.

29. Ben-Shahar Y, Leung HT, Pak WL, Sokolowski MB, Robinson GE. cGMP-
dependent changes in phototaxis: a possible role for the foraging gene in honey
bee division of labor. J Exp Biol. 2003;206(14):2507–15.

30. Whitfield CW, Cziko AM, Robinson GE. Gene expression profiles in the brain
predict behavior in individual honey bees. Science (80-). 2003;302(5643):296–9.

31. Ingram KK, Oefner P, Gordon DM. Task-specific expression of the foraging gene
in harvester ants. Mol Ecol. 2005;14(3):813–8.

32. Lucas C, Sokolowski MB. Molecular basis for changes in behavioral state in ant
social behaviors. Proc Natl Acad Sci U S A. 2009;106(15):6351–6.

33. Zayed A, Robinson GE. Understanding the relationship between brain gene
expression and social behavior: lessons from the honey bee. Annu Rev Genet.
2012;46:591–615.

34. Sullivan JP, Jassim O, Fahrbach SE, Robinson GE. Juvenile hormone paces
behavioral development in the adult worker honey bee. Horm Behav.
2000;37(1):1–14.

35. Ament SA, Corona M, Pollock HS, Robinson GE. Insulin signaling is involved in
the regulation of worker division of labor in honey bee colonies. Proc Natl Acad
Sci U S A. 2008;105(11):4226–31.

36. Dolezal AG, Brent CS, Hölldobler B, Amdam G V. Worker division of labor and
endocrine physiology are associated in the harvester ant, *Pogonomyrmex*
californicus. J Exp Biol. 2012;215(3):454–60.

37. Corona M, Libbrecht R, Wurm Y, Riba-Grognuz O, Studer RA, Keller L.

Vitellogenin underwent subfunctionalization to acquire caste and behavioral specific expression in the harvester ant *Pogonomyrmex barbatu*. PLoS Genet. 2013;9(8).

38. Kohlmeier P, Alleman AR, Libbrecht R, Foitzik S, Feldmeyer B. Gene expression is more strongly associated with behavioural specialization than with age or fertility in ant workers. Mol Ecol. 2019;28(3):658–670.

39. Libbrecht R, Corona M, Wende F, Azevedo DO, Serrão JE, Keller L. Interplay between insulin signaling, juvenile hormone, and vitellogenin regulates maternal effects on polyphenism in ants. Proc Natl Acad Sci U S A. 2013;110(27):11050–5.

40. Marco Antonio DS, Guidugli-Lazzarini KR, Do Nascimento AM, Simões ZLP, Hartfelder K. RNAi-mediated silencing of vitellogenin gene function turns honeybee (*Apis mellifera*) workers into extremely precocious foragers. Naturwissenschaften. 2008;95(10):953–61.

41. Gospocic J, Shields EJ, Glastad KM, Lin Y, Penick CA, Yan H, et al. The neuropeptide corazonin controls social behavior and caste Identity in ants. Cell. 2017;170(4):748-759.e12.

42. Kohlmeier P, Feldmeyer B, Foitzik S. *Vitellogenin-like* A–associated shifts in social cue responsiveness regulate behavioral task specialization in an ant. PLoS Biol. 2018;16(6):1–26.

43. Kamhi JF, Traniello JFA. Biogenic amines and collective organization in a superorganism: Neuromodulation of social behavior in ants. Brain Behav Evol. 2013;82(4):220–36.

44. Schulz D, Robinson G. Biogenic amines and division of labor in honey bee colonies: behaviorally related changes in the antennal lobes and age-related changes in the mushroom bodies. 1999;481–8.

45. Mercer AR, Menzel R. The effects of biogenic amines on conditioned and unconditioned responses to olfactory stimuli in the honeybee *Apis mellifera*. J Comp Physiol. 1982;145(3):363–8.

46. Scheiner R, Entler B V., Barron AB, Scholl C, Thamm M. The effects of fat body tyramine level on gustatory responsiveness of honeybees (*Apis mellifera*) differ between behavioral castes. Front Syst Neurosci. 2017;11(August):1–8.

47. Ferguson ST, Bakis I, Zwiebel LJ. Advances in the study of olfaction in eusocial ants. Insects. 2021;12(3)252.

48. Martin S, Drijfhout F. A review of ant cuticular hydrocarbons. J Chem Ecol. 2009;35(10):1151–61.

49. van Wilgenburg E, Symonds MRE, Elgar MA. Evolution of cuticular hydrocarbon diversity in ants. J Evol Biol. 2011;24(6):1188–98.

50. Slifer EH. The Structure of Arthropod Chemoreceptors. Annu Rev Entomol. 1970;15(1):121–42.

51. Altner H, Prillinger L. Ultrastructure of invertebrate chemo-, thermo-, and hygroreceptors and its functional significance. Int Rev Cytol. 1980;67:69–139.

52. Hallberg E, Hansson BS. Arthropod sensilla: morphology and phylogenetic considerations. Microsc Res Tech. 1999;47(6):428–39.

53. Wicher D, Miazzi F. Functional properties of insect olfactory receptors: ionotropic receptors and odorant receptors. Cell Tissue Res. 2021;383(1):7–19.

54. Leal WS. Odorant reception in insects: Roles of receptors, binding proteins, and

degrading enzymes. Annu Rev Entomol. 2013;58:373–91.

55.   Sánchez-Gracia A, Vieira FG, Almeida FC, Rozas J. Comparative genomics of
the major chemosensory gene families in arthropods. eLS. 2011.

56.   Breer H, Fleischer J, Pregitzer P, Krieger J. Molecular mechanism of insect
olfaction: olfactory receptors In: Picimbon JF, editors. Olfactory concepts of insect
control-alternative to insecticides. 2019;15:93–114.

57.   Jacquin-Joly E, Lucas P. Pheromone reception and transduction: mammals and
insects illustrate converging mechanisms across phyla. Curr Top Neurochem.
2005;4:75–105.

58.   Wilson RK, Mainen ZF. Early events in olfactory processing. Annu Rev Neurosci.
2006;29:163–201.

59.   Anton S, Homberg U. Antennal lobe structure. Insect Olfaction. 1999;97–124.

60.   Hansson BS, Anton S. Function and morphology of the antennal lobe: new
developments. Annu Rev Entomol. 2000;45:203–31.

61.   de Belle JS, Kanzaki R. Protocerebral Olfactory Processing. Insect Olfaction.
1999;243–81.

62.   Saad R, Cohanim AB, Kosloff M, Privman E. Neofunctionalization in ligand
binding sites of ant olfactory receptors. Genome Biol Evol. 2018;10(9):2490–500.

63.   Oxley PR, Ji L, Fetter-Pruneda I, McKenzie SK, Li C, Hu H, et al. The genome of
the clonal raider ant *Cerapachys Biroi*. Curr Biol. 2014;24(4):451–8.

64.   Zhou X, Rokas A, Berger SL, Liebig J, Ray A, Zwiebel LJ. Chemoreceptor
evolution in Hymenoptera and Its Implications for the evolution of eusociality.
Genome Biol Evol. 2015;7(8):2407–16.

65.   Engsontia P, Sangket U, Robertson HM, Satasook C. Diversification of the ant
odorant receptor gene family and positive selection on candidate cuticular
hydrocarbon receptors. BMC Res Notes. 2015;8(1):1–13.

66.   Smith CD, Zimin A, Holt C, Abouheif E, Benton R, Cash E, et al. Draft genome of
the globally widespread and invasive Argentine ant (*Linepithema humile*). Proc
Natl Acad Sci. 2011;108(14):5673–8.

67.   Smith CR, Smith CD, Robertson HM, Helmkampf M, Zimin A, Yandell M, et al.
Draft genome of the red harvester ant *Pogonomyrmex barbatus*. Proc Natl Acad
Sci U S A. 2011;108(14):5667–72.

68.   Zhou X, Slone JD, Rokas A, Berger SL, Liebig J, Ray A, et al. Phylogenetic and
transcriptomic analysis of chemosensory receptors in a pair of divergent ant
species reveals sex-specific signatures of odor coding. PLoS Genet. 2012;8(8).

69.   McKenzie SK, Fetter-Pruneda I, Ruta V, Kronauer DJC. Transcriptomics and
neuroanatomy of the clonal raider ant implicate an expanded clade of odorant
receptors in chemical communication. Proc Natl Acad Sci U S A.
2016;113(49):14091–6.

70.   Pask GM, Slone JD, Millar JG, Das P, Moreira JA, Zhou X, et al. Specialized
odorant receptors in social insects that detect cuticular hydrocarbon cues and
candidate pheromones. Nat Commun. 2017;8(1):1–10.

71.   Jongepier E, Séguret A, Labutin A, Feldmeyer B, Gstöttl C, Foitzik S, et al.
Convergent loss of chemoreceptors across independent origins of slave-making
in ants. Mol Biol Evol. 2022;39(1):1–14.

72.   Trible W, Olivos-Cisneros L, McKenzie SK, Saragosti J, Chang N-C, Matthews

BJ, et al. *orco* Mutagenesis causes loss of antennal lobe glomeruli and impaired
social behavior in ants. Cell. 2017;170(4):727–735.e10.

73. Yan H, Opachaloemphan C, Mancini G, Yang H, Gallitto M, Mlejnek J, et al. An
engineered *orco* mutation produces aberrant social behavior and defective neural
development in ants. Cell. 2017;170(4):736–747.e9.

74. Beshers S, Fewell JH. Models of division of labor in social insects. Annu. Rev.
Entomol. 2001;46, 413–40.

75. Lichocki P, Tarapore D, Keller L, Floreano D. Neural networks as mechanisms to
regulate division of labor. Am Nat. 2012;179(3):391–400.

76. Naug D. From division of labor to collective behavior: behavioral analyses at
different levels. Behav Ecol Sociobiol. 2016;70(7):1113–5.

77. Jeanson R, Weidenmüller A. Interindividual variability in social insects - proximate
causes and ultimate consequences. Biol Rev. 2014;89(3):671–87.

78. Ulrich Y, Kawakatsu M, Tokita CK, Saragosti J, Chandra V, Tarnita CE, et al.
Response thresholds alone cannot explain empirical patterns of division of labor
in social insects. PLoS Biol. 2021;19(6):1–21.

79. Bonabeau E, Sobkowski A, Theraulaz G, Deneubourg JL. Adaptive task allocation
inspired by a model of division of labor in social insects. Biocomput Emergent
Comput. 1997;36–45.

80. Bonabeau E, Theraulaz G, Deneubourg JL. Fixed response thresholds and the
regulation of division of labor in insect societies. Bull Math Biol. 1998;60(4):753–
807.

81. Schulz DJ, Elekonich MM, Robinson GE. Biogenic amines in the antennal lobes
and the initiation and maintenance of foraging behavior in honey bees. J
Neurobiol. 2003;54(2):406–16.

82. Zhukovskaya MI, Polyanovsky AD. Biogenic amines in insect antennae. Front
Syst Neurosci. 2017;11:1–9.

83. Cook CN, Mosqueiro T, Brent CS, Ozturk C, Gadau J, Pinter-Wollman N, et al.
Individual differences in learning and biogenic amine levels influence the
behavioural division between foraging honeybee scouts and recruits. J Anim Ecol.
2019;88(2):236–46.

84. Azevedo DO, De Paula SO, Zanuncio JC, Martinez LC, Serrão JE. Juvenile
hormone downregulates vitellogenin production in *Ectatomma tuberculatum*
(Hymenoptera: Formicidae) sterile workers. J Exp Biol. 2016;219(1):103–8.

85. Rodrigues MA, Flatt T. Endocrine uncoupling of the trade-off between
reproduction and somatic maintenance in eusocial insects. Curr Opin insect Sci.
2016;16:1–8.

86. Jedlicka P, Ernst UR, Votavová A, Hanus R, Valterová I. Gene expression
dynamics in major endocrine regulatory pathways along the transition from
solitary to social life in a bumblebee, *Bombus terrestris*. Front Physiol. 2016;7:1–
20.

87. Opachaloemphan C, Mancini G, Konstantinides N, Parikh A, Mlejnek J, Yan H, et
al. Early behavioral and molecular events leading to caste switching in the ant
Harpegnathos. Genes Dev. 2021;35(5–6):410–24.

88. Norman VC, Hughes WOH. Behavioural effects of juvenile hormone and their
influence on division of labour in leaf-cutting ant societies. J Exp Biol.

2016;219(1):8–11.

89.   Robinson GE. Regulation of honey bee age polyethism by juvenile hormone.
Behav Ecol Sociobiol. 1987;20(5):329–38.

90.   Fussnecker B, Grozinger C. Dissecting the role of Kr-h1 brain gene expression in
foraging behavior in honey bees (*Apis mellifera*). Insect Mol Biol. 2008;17(5):515–
22.

91.   Grozinger CM, Robinson GE. Endocrine modulation of a pheromone-responsive
gene in the honey bee brain. J Comp Physiol A Neuroethol Sensory, Neural,
Behav Physiol. 2007;193(4):461–70.

92.   Shpigler H, Patch HM, Cohen M, Fan Y, Grozinger CM, Bloch G. The
transcription factor Krüppel homolog 1 is linked to hormone mediated social
organization in bees. BMC Evol Biol. 2010;10(1):120.

93.   Gospocic J, Glastad KM, Sheng L, Shields EJ, Berger SL, Bonasio R. Kr-h1
maintains distinct caste-specific neurotranscriptomes in response to socially
regulated hormones. Cell. 2021;184(23):5807-5823.e14.

94.   von der Emde, G; Warrant E. The ecology of animal senses matched filters for
cconomical Sensing. 1st ed. Springer International Publishing; 2016.

95.   Carey AF, Wang G, Su CY, Zwiebel LJ, Carlson JR. Odorant reception in the
malaria mosquito *Anopheles gambiae*. Nature. 2010;464(7285):66–71.

96.   Dornhaus A. Specialization does not predict individual efficiency in an ant. PLoS
Biol. 2008;6(11):2368–75.

97.   Libbrecht R, Keller L. Genetic compatibility affects division of labor in the
argentine ant *Linepithema humile*. Evolution. 2013;67(2):517–24.

98.   Gadenne C, Barrozo RB, Anton S. Plasticity in insect olfaction: to smell or not to
smell? Annu Rev Entomol. 2016;61:317–33.

99.   Zhou S, Stone EA, Mackay TFC, Anholt RRH. Plasticity of the chemoreceptor
repertoire in *Drosophila melanogaster*. PLoS Genet. 2009;5(10).

100. Von Der Weid B, Rossier D, Lindup M, Tuberosa J, Widmer A, Col JD, et al.
Large-scale transcriptional profiling of chemosensory neurons identifies receptor-
ligand pairs in vivo. Nat Neurosci. 2015;18(10):1455–63.

101. Benton R, Sachse S, Michnick SW, Vosshall LB. Atypical membrane topology and
heteromeric function of *Drosophila* odorant receptors in vivo. PLoS Biol.
2006;4(2):240–57.

102. Krieger J, Klink O, Mohl C, Raming K, Breer H. A candidate olfactory receptor
subtype highly conserved across different insect orders. J Comp Physiol A
Neuroethol Sensory, Neural, Behav Physiol. 2003;189(7):519–26.

103. Larsson MC, Domingos AI, Jones WD, Chiappe ME, Amrein H, Vosshall LB.
Or83b encodes a broadly expressed odorant receptor essential for *Drosophila*
olfaction. Neuron. 2004;43(5):703–14.

104. Mersch DP, Crespi A, Keller L. Tracking individuals shows spatial fidelity is a key
regulator of ant social organization. Science. 2013;340(6136):1090–3.

105. Richardson TO, Kay T, Braunschweig R, Journeau OA, Rüegg M, McGregor S, et
al. Ant behavioral maturation is mediated by a stochastic transition between two
fundamental states. Curr Biol. 2021;31(10):2253-2260.e3.

106. Dacks AM, Dacks JB, Christensen TA, Nighorn AJ. The cloning of one putative
octopamine receptor and two putative serotonin receptors from the tobacco

hawkmoth, *Manduca sexta*. Insect Biochem Mol Biol. 2006;36(9):741–7.

107. Getahun MN, Olsson SB, Lavista-Llanos S, Hansson BS, Wicher D. Insect
odorant response sensitivity is tuned by metabotropically autoregulated olfactory
receptors. PLoS One. 2013;8(3):1–9.

108. Stengl M, Funk NW. The role of the coreceptor Orco in insect olfactory
transduction. J Comp Physiol A Neuroethol Sensory, Neural, Behav Physiol.
2013;199(11):897–909.

109. Blundell JE, Halford JCG. Serotonin and appetite regulation: implications for the
pharmacological treatment of obesity. CNS Drugs. 1998;9(6):473–95.

110. French AS, Simcock KL, Rolke D, Gartside SE, Blenau W, Wright GA. The role of
serotonin in feeding and gut contractions in the honeybee. J Insect Physiol.
2014;61(1):8–15.

111. Voigt JP, Fink H. Serotonin controlling feeding and satiety. Behav Brain Res.
2015;277:14–31.

112. Beninger RJ. The role of dopamine in locomotor activity and learning. Brain Res
Rev. 1983;6(2):173–96.

113. Liang X, Ho MCW, Zhang Y, Li Y, Wu MN, Holy TE, et al. Morning and evening
circadian pacemakers independently drive premotor centers via a specific
dopamine relay. Neuron. 2019;102(4):843-857.e4.

114. Libbrecht R, Nadrau D, Foitzik S. A role of histone acetylation in the regulation of
circadian rhythm in ants. iScience. 2020;23(2):100846.

115. Root CM, Ko KI, Jafari A, Wang JW. Presynaptic facilitation by neuropeptide
signaling mediates odor-driven food search. Cell. 2011;145(1):133–44.

116. Anton S, Dufour MC, Gadenne C. P Plasticity of olfactory-guided behaviour and
its neurobiological basis: lessons from moths and locusts. Entomol. Exp. Appl.
2007:1–11.

117. Tawfik AI, Osir EO, Hassanali A, Ismail SH. Effects of juvenile hormone treatment
on phase changes and pheromone production the desert locust, *Schistocerca*
gregaria (Forskal) (Orthoptera: Acrididae). J Insect Physiol. 1997;43(12):1177–82.

118. Tawfik AI, Treiblmayr K, Hassanali A, Osir EO. Time-course haemolymph juvenile
hormone titres in solitarious and gregarious adults of *Schistocerca gregaria*, and
their relation to pheromone emission, CA volumetric changes and oocyte growth.
J Insect Physiol. 2000;46(7):1143–50.

119. Wiesel G, Tappermann S, Dorn A. Effects of juvenile hormone and juvenile
hormone analogues on the phase behaviour of *Schistocerca gregaria* and
Locusta migratoria. J Insect Physiol. 1996;42(4):385–95.

120. Guo W, Song J, Yang P, Chen X, Chen D, Ren D, et al. Juvenile hormone
suppresses aggregation behavior through influencing antennal gene expression in
locusts. PLoS Genet. 2020;16(4):1–18.

121. Pamminger T, Foitzik S, Kaufmann KC, Schützler N, Menzel F. Worker
personality and its association with spatially structured division of labor. PLoS
One. 2014;9(1):1–8.

122. Kohlmeier P, Negroni MA, Kever M, Emmling S, Stypa H, Feldmeyer B, et al.
Intrinsic worker mortality depends on behavioral caste and the queens' presence
in a social insect. Sci Nat. 2017;104(3–4).

123. Sendova-Franks A, Franks NR. Task allocation in ant colonies within variable

environments (a study of temporal polyethism: Experimental). Bull. Math.
Biol.1993;55(1):75–96.

124. Andrews S. A quality control tool for high throughput sequence data. 2016.
Available from: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

125. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics. 2014;30(15):2114–20.

126. Kaur R, Stoldt M, Jongepier E, Feldmeyer B, Menzel F, Bornberg-Bauer E, et al.
Ant behaviour and brain gene expression of defending hosts depend on the
ecological success of the intruding social parasite. Philos Trans R Soc B Biol Sci.
2019;374:1769

127. Shumate A, Salzberg SL. Liftoff: accurate mapping of gene annotations.
Bioinformatics. 2021;37(12):1639–43.

128. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:
ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15–21.

129. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.

130. Love MI, Anders S, Huber W. Differential analysis of count data - the DESeq2
package. Genome Biology. 2014;15:550.

131. Emms DM, Kelly S. OrthoFinder: solving fundamental biases in whole genome
comparisons dramatically improves orthogroup inference accuracy. Genome Biol.
2015;16(1):157.

132. Das B, de Bekker C. Time-course RNASeq of *Camponotus floridanus* forager and
nurse ant brains indicate links between plasticity in the biological clock and
behavioral division of labor. BMC Genomics. 2022;23(1):1–23.

133. Alleman A, Stoldt M, Feldmeyer B, Foitzik S. Tandem-running and scouting
behaviour are characterized by up-regulation of learning and memory formation
genes within the ant brain. Mol Ecol. 2019;28:2342–2359.

134. Korb J, Meusemann K, Aumer D, Bernadou A, Elsner D, Feldmeyer B, et al.
Comparative transcriptomic analysis of the mechanisms underpinning ageing and
fecundity in social insects. Philos Trans R Soc B Biol Sci. 2021;376(1823).

135. Qiu B, Dai X, Li P, Larsen RS, Li R, Price AL, et al. Canalized gene expression
during development mediates caste differentiation in ants. Nat Ecol Evol.
2022;6(11):1753–65.

136. Alexa A, Rahnenfuhrer J. topGO: Enrichment Analysis for Gene Ontology. R
package version 2.50.0. 2018. Available from:
https://bioconductor.org/packages/release/bioc/html/topGO.html

137. RStudio Team. RStudio: Integrated Development for R. RStudio, PBC, Boston,
MA. 2020. Available from: http://www.rstudio.com/

Figures and Tables

Fig 1. Variation in behavior, brain and antenna gene expression between nurses and foragers. (*A*) Boxplot showing behavioral differences between ants selected for transcriptomic analysis. Each black dots represent individual ants. For differential expression analysis, each sample contains the pooled RNA of seven ants of the respective behavioral phenotype. (*B*) Venn diagrams showing the number of DEG that were overexpressed in nurses (red) and foragers (turquoise) for both antenna and brain tissues. Principal component analysis (PCA) plots based on all expressed genes for (*C*) antenna and (*D*) brain samples. The color of each sample represents task (red = nurse, turquoise = forager), and the shape the colony of origin. The ellipses were defined by a

881 95% confidence interval and show how similar/different the samples are within task

882 groups.

Fig 2. OR expression in the antennae differs between nurses and foragers. (*A*)

Among the DEG overexpressed in nurses and foragers, the most represented OR

subfamilies are indicated by different colors. The red dotted line represents the

significance threshold of our differential expression analysis. (*B,C*). Boxplots representing

the expression of (*B*) *Orco* and *OR257*, the gene with the highest expression difference

in terms of log$_2$ FoldChange in the 9-exon subfamily, and (*C*) *OR041*, *OR045* and *OR318*,

the genes with the highest expression difference in terms of log$_2$ FoldChange in the H, L

and P subfamilies, respectively.

Figure 3. Gene expression differences between nurse and forager antennae.

Boxplots representing the expression of (*A-E*) selected genes in biogenic amines

pathways, as well as (*F*) one behavioral candidate gene (Table S1). Gene identity is

shown in parentheses.

Supporting Information

Table S1. List of candidate genes related to behavior phenotypes.

Table S2. List of enriched GO biological process terms of the DEG in brain and antenna.

Table S3. Behaviors and positions annotations during nest scans. Nursing was defined as the number of observations a worker antennating, grooming, feeding or carrying a brood item, and its position inside the nest (on the brood or near to the brood pile). Foraging was defined as the number of observations an individual was found outside the nest, collecting food or water.

Dataset S1. Gene list with caste-differential expression, NCBI BlastX results, and gene read count of 28 RNA-seq samples.

Dataset S2. OR subfamilies and orthogroups from the DEG lists between nurses and foragers in the antenna.

Dataset S3. Summary of ant brain and antenna transcriptome data sets and mapping results.

Dataset S4. List of orthogroups from the DEG list between nurses and foragers in _Temnothorax longispinosus_.