Task-specific patterns of odorant receptor expression in worker antennae

# indicates a sensory filter regulating division of labor in ants

Marcel A. Caminer[1]*, Romain Libbrecht[1,2], Megha Majoe[1], David Vuong Ho[1,3], Susanne Foitzik[1]

[1]Institute of Organismic and Molecular Evolution, Johannes Gutenberg University Mainz, Germany
[2]current address: Institut de Recherche sur la Biologie de l'Insecte, UMR 7261, CNRS, University of
Tours, France
[3]Institute of Developmental and Neurobiology, Johannes Gutenberg University Mainz, Germany &
Institute of Molecular Biology, IMB
11

*correspondence: marcelcaminer@gmail.com

Author Contributions: M.A.C., R.L. and S.F. designed research; M.A.C. performed research; M.A.C.,

M.M. and D.H. contributed analytic tools; M.A.C analyzed data; and M.A.C., R.L., M.M, and S.F. wrote the

paper.

Competing Interest Statement: The authors declare no competing interest.

Classification: Biological Sciences. Evolution.

Keywords: social insects | transcriptomics | behavioral specialization | odorant perception

This PDF file includes:

Main Text

Figures 1 to 3

Tables S1 to S3

SI References

Legends for Datasets S1 to S4

Abstract

Division of labor (DOL) is a characteristic trait of insect societies, where tasks are generally performed by groups of specialized individuals. In social insects, young workers perform duties within the safety of the nest (e.g., brood care), while older ones undertake riskier tasks (e.g., foraging for food). This DOL remains dynamic, and workers may switch back and forth when colony needs require. Theoretical models propose that workers differ in their thresholds to take on certain tasks when confronted to task-related stimuli, resulting in variation in their response to such stimuli, task-specialization, and thus DOL. Such models assume that workers differ in how they respond to task-related information rather than in how they perceive such information. Here, we test the hypothesis that DOL rather stems from workers differing in their efficiency to detect task-related stimuli. We used transcriptomics to compare gene expression in the antennae and in the brain between nurses and foragers in the ant *Temnothorax longispinosus*. We found that seven times as many genes were differentially expressed between the behavioral phenotypes in the antennae compared to the brain. Moreover, nearly half of all odorant receptors genes were differentially expressed, with an overrepresentation of the 9-exon gene subfamily upregulated in the antennae of nurses. These findings indicate that nurses and foragers differ in how they perceive their olfactory environment, and task-related signals. The results of this study support the hypothesis that a sensory filter in the antennae predisposes workers to specialize in specific tasks, and may improve our understanding of DOL in insect societies.

Significance Statement

We tested the hypothesis that behavioral variation and task specialization in insect societies are regulated by differences in olfactory perception. We show that physiological differences between the most distinct behavioral phenotypes of nurses and foragers are less pronounced in the central nervous system than in the peripheral sensory organs. Nurses mostly activate odorant receptors from the 9-exon subfamily in the antenna that are predicted to be responsible for perception of chemical signals from nestmates, while foragers overexpress odorant receptors from diverse subfamilies. Our results demonstrate that workers performing different tasks also live in different sensory environments, suggesting that a sensory filter may play an important role in regulating division of labor in insect societies.

Main Text

Introduction

Division of labor (DOL) is an important organizing principle of complex biological systems that arose independently during three of the major evolutionary transitions [1]. DOL was originally formulated in the context of the production process in human societies [2], but specialization to specific tasks is also found within cells and across many organisms [1]. Examples range from bacteria, where clonal populations are divided into subpopulations focusing on different activities [3–6] to multicellular organisms with differentiation of cells into different tissues and organs, and individuals performing specific roles in animal societies [7, 8]. To understand the evolution of complex life it is therefore essential to investigate the mechanisms that underlie DOL.

DOL in insect societies results from individuals specializing in the performance of specific tasks. In addition to the reproductive DOL between fertile queens and functionally sterile workers, there is a behavioral DOL among workers that specialize in tasks such as brood care, foraging, nest building, and defense [9–11]. Several factors can affect task specialization, including age [12, 13], nutrition [14, 15], morphology [16], genotype [17–19], experience [20], and colony size [21–23]. In most social insect species, workers display an age polyethism by which younger individuals tend to perform intranidal tasks, while older individuals are more likely to undertake activities such as nest-defense and foraging outside the nest [24, 25]. Yet, such specialization among workers remains flexible, as foragers can revert to perform brood care when needed [26, 27].

Several molecular mechanisms have been implicated in the regulation of DOL. Task specialization is associated with transcriptional changes in the worker brain [28–33]. Molecular pathways such as the insulin/insulin-like signaling (IIS), vitellogenin (Vg) and juvenile hormone (JH) pathways are involved in the regulation of worker behavior [34–39]. Functional manipulations have confirmed that the expression of key genes in the worker brain controls task specialization [40–42]. Behavioral variation among workers is also associated with signaling of biogenic amines (e.g., dopamine, octopamine, tyramine, serotonin), which act as neurotransmitters or neuromodulators involved in the modulation of the responsiveness to task-associated stimuli [43–46].

The self-organization and collective behavior in insect societies are maintained via the exchange

of chemical information (47). Social insects communicate primarily through glandular pheromones and

complex mixtures of long-chain hydrocarbons on their cuticle. These cuticular hydrocarbons (CHC) facilitate

recognition of nestmates, developmental stages, castes, sexes, and species (48, 49). Social insects

perceive chemical information via different types of sensilla on their antennae (50–52). Decoding the

identities of chemical compounds relies on odorant receptors (OR) located within each sensillum (53, 54).

OR are transmembrane proteins expressed in the dendrites of olfactory receptor neurons (ORN). The

largely conserved OR coreceptor (Orco) of insects (55) is required for odorant recognition in the dendritic

membrane: it forms an ion channel with specific OR, which determines the sensitivity and specificity of the

ORN (56). Odorant molecules penetrate through the antenna cuticular pores and are transported by

odorant-binding proteins to the ORN membrane, where they interact with receptors, leading to the

generation of action potentials (57, 58). ORN axons relay signals from the sensilla to the glomeruli of the

antennal lobes in the insect brain, which are the first processing unit for olfactory information. Then, ORN

make synaptic contact with the projection neurons and local neurons, which transfer information to the

central brain (59–61).

Several lines of evidence indicate that OR genes play central roles in the regulation of social life of

insects. First, social insect species typically harbor large numbers of OR genes (62–65). Second, some OR

gene families have specifically expanded during social evolution, such as the 9-exon subfamily in ants,

which appears to serve an important function in the perception of CHC (64, 66–70). Third, species that

evolved socially parasitic strategies resulting in reduced behavioral repertoires show a strong and

convergent reduction in the number of OR genes (71). Finally, experimentally produced mutant ants that

lack the *Orco* gene (coding for the co-receptor necessary for OR to properly function) show impaired social

behavior (72, 73).

Current DOL models bring together the chemical nature of social insect communication and

variation among workers in their response to chemical cues. They posit that flexible response thresholds

to task-related chemical stimuli serve as regulators of worker specialization (74). Workers take on a

particular task when the stimulus intensity exceeds their individual threshold for this task. Therefore,

individuals with lower thresholds for a given task are more likely to perform it than those with higher

thresholds (75). Individual response decisions and task performance are modulated via numerous

parameters on different time scales (7, 76–78), and despite extensive research on DOL and task allocation,

these mechanisms are not fully understood.

As indicated by the name, "response threshold models" typically assume that individuals differ in

their response to specific signals, not necessarily in their ability or efficiency to detect such signals. This

implies that such thresholds and the associated responses are set in the central nervous system, possibly

via molecular pathways in the brain that correlate with behavioral variation (74, 79, 80). We propose that

DOL models would benefit from considering odor sensitivity as a potential upstream sensory filter that may

affect task specialization. Along these lines, we hypothesize that behavioral variation among workers may

also stem from their ability and/or efficiency in detecting different signals. For example, we propose that

individuals that specialize in brood care do so because they are better at detecting brood cues, rather than

(or in addition to) being more likely to respond to similar levels of brood cues. To test this hypothesis that a

sensory filter regulates inter-individual behavioral variation, and thus the DOL in social insects, we

investigated transcriptional differences in the brain and antennae of workers that specialize in either brood

care or foraging behavior in the ant *Temnothorax longispinosus*. We found that i) behavioral variation was

associated with important transcriptomic changes in the antennae, ii) these changes included a large

proportion of the OR gene repertoire, and iii) individuals specializing in brood care overexpressed an OR

gene family putatively involved in detecting social cues. These findings support our hypothesis that the

peripheral nervous system, acting as a sensory filter, plays an important role in regulating behavioral

differences between workers and thus in the DOL of social insects.

Results

Identification of **Temnothorax longispinosus** *behavioral phenotypes*

Task specialization in the ant *T. longispinosus* is neither genetically fixed nor rigid, but changes with age

and in response to colony needs (38). We conducted behavioral observations of seven *T. longispinosus*

laboratory colonies to identify individuals that specialize in brood care behavior (hereafter referred to as

nurses), and others that specialize in foraging (hereafter referred to as foragers) (Fig. 1*A*; see Methods

section for more details). Individuals identified as nurses interacted with the brood in 54% ± 22% (mean ±

sd) of the observations, and were recorded outside the nest in 1% ± 3% of the observations. On the

contrary, foragers were found outside the nest in 42% ± 26% of the observations, but interacted with the

brood in only 2% ± 4% of the observations.

Larger task-associated transcriptomic changes in the antennae than in the brain

To investigate transcriptomic variation between nurses and foragers, we used RNA-seq to generate seven

nurse and seven forager brain and antenna samples, each consisting of pooled tissue from seven workers

of a single colony. Of the 14,837 protein-coding genes annotated in the *T. longispinosus* genome, we found

that 91% (13,494) and 92% (13,683) were expressed (FPKM > 0) in the brain and the antennae,

respectively. Using an adjusted p-value of 0.05 as significance threshold, we detected 333 differentially

expressed genes (DEG) in the brain (217 upregulated in nurses, 116 in foragers), and 2,254 in the antennae

(1,234 upregulated in nurses, 1,020 in foragers; Dataset S1; Fig. 1*B*). We found an overlap of 162 DEG

between the two tissues, including 143 DEG that showed differences in the same direction across tissues.

Principal component analysis (PCA) plots indicate that both brain and antenna samples cluster by

behavioral phenotype rather than the colony of origin (Fig.1 *C-D*).

OR gene expression differs between antennae of nurses and foragers

To investigate whether odorant perception differs between nurses and foragers, we focused our attention

on the expression of OR genes in the antennae. We found that all 419 OR genes in the genome of *T.*

longispinosus (71) were expressed in the antennae, and that 48% (202/419) of them were differentially

expressed between nurses and foragers. Specifically, 65 OR genes were upregulated in nurses (15% of all

OR genes), and 137 in foragers (32% of all ORs) (Fig. 2). Then, we studied which OR subfamilies were

preferentially expressed in nurses and foragers. The 65 OR genes overexpressed in nurses belonged to

three OR subfamilies, while the 137 OR genes upregulated in foragers were distributed among 19 OR

subfamilies. Foragers overexpressed 26, 8, and 8 OR genes from the L, P, and H subfamilies, respectively,

while no OR genes from these subfamilies were upregulated in nurses (Dataset S2). We found that 61%

(26/43) of the genes from the L subfamily were overexpressed in foragers, which represents a significant

overrepresentation (Fisher's test, odds ratio = 3.63, p-value < 0.001). For the P and H subfamilies we did

not find such an overrepresentation, likely due to lower gene numbers (P subfamily: Fisher's test, odds ratio

= 3.62, p-value = 0.06; H subfamily: Fisher's test, odds ratio = 1.87, p-value = 0.19). On the other hand, we

found that 82% (53/65) of the OR genes overexpressed in nurses belong to the 9-exon subfamily. This

results in an overrepresentation of the 9-exon subfamily in genes that were overexpressed in nurses

(Fisher's test, odds ratio = 20.98, p-value < 0.001), with 46% (53/114) of this subfamily being overexpressed

in nurses. In contrast, only 6% (8/137) of the 9-exon subfamily was overexpressed in foragers, which is

less than expected by chance (Fisher's test, odds ratio = 0.10, p-value < 0.001). We also found that nurses

overexpressed *Orco* compared to foragers (FDR p-value = 0.001).

Genes in biogenic amine pathways vary in expression between nurses and foragers

Because biogenic amines are implicated in the regulation of behavior, and they may affect sensory

perception (45, 81–83), we screened our lists of DEG in the brain and antennae for genes in biogenic amine

pathways. We found five genes associated with biogenic amine signaling in the antennae, and none in the

brain. Genes in the serotonin (*5-hydroxytryptamine*, DBV15_11483), tyramine (*tyramine beta-hydroxylase*,

DBV15_00422) and octopamine (*octopamine receptor* LOC112465659) pathways were upregulated in

foragers, while nurses showed a higher expression of genes in the dopamine (*dopamine 1-like receptor 2*,

DBV15_07611) and octopamine (*octopamine beta2 receptor*, DBV15_10418) pathways (Fig. 3 *A-E*).

Association between behavioral variation and molecular pathways in brain and antennae

Our analysis revealed expression differences between the brains of nurses and foragers for genes involved

in the regulation of task specialization in social insects. Previous studies have found that *Vg* genes and the

associated JH and IIS/TOR pathways are important endocrine networks that play central roles in the

regulation of lifespan, fertility and behavior in bees and ants (35, 84–89). Therefore, we searched our RNA-

seq data for task-associated expression of genes involved in the metabolism, biosynthesis, and regulation

of these pathways in the brain and antennae. We found evidence that multiple genes from these pathways

are differentially expressed between nurses and foragers (Table S1). Among the most interesting genes

upregulated in the brain of nurses were *conventional Vg* (LOC112466671), *Vg-like A* (named/classified as

per 27; DBV15_03138), *venom carboxylesterase-6-like* (DBV15_11528), and *protein takeout*

(DBV15_08771), while *allatostatin A-like* (LOC112454443) and *insulin-like growth factor I* (*IGF1*)

(LOC112454447) were overexpressed in the brain of foragers. Remarkably, *venom carboxylesterase-6-*

like and *IGF1* were overexpressed in the antennae of nurses and foragers respectively. Additionally, high

expression levels of the zinc-finger transcription factor *Krüppel homolog-1* (*Kr-h1*) were detected in the

antennae of foragers (DBV15_06330) (Fig. 3*F*). The expression of this gene has been correlated with caste

and behavioral differences in the brain of social insects (86, 90–93), but little is known about its function

and gene targets in the antennae.

We performed a GO enrichment analysis to gain a deeper understanding of the biological

processes represented in the lists of DEG. This analysis detected many enrichments based on single one

or few genes, and we mention below a few interesting processes with the number of genes driving the

enrichment in parentheses (see Table S2 for complete list). Genes that were upregulated in the brain of

nurses were enriched for biological processes such as *translation* (19 genes), *cellular iron ion homeostasis*

(2 genes), and *catabolic process* (6 genes); while *translation* (22 genes), *endocytosis* (6 genes), *glucose*

metabolic process (4 genes), and *regulation of cell cycle* (3 genes) were enriched in the antennae. In the

list of genes that were overexpressed in the brain of foragers, we found that enriched processes included

peptide metabolic process (4 genes), *methionyl-tRNA aminoacylation* (1 gene), *positive regulation of type*

I interferon production (1 gene), and *cell redox homeostasis* (1 gene); while *phosphatidylinositol*

phosphorylation (4 genes), *inositol phosphate dephosphorylation* (3 genes), *carbohydrate metabolic*

process (17 genes), *innate immune response* (2 genes), and *nucleotide catabolic process* (3 genes) were

enriched in the antennae.

Discussion

216
In this study, we tested the hypothesis that variation among social insect workers in their ability to detect

different chemical signals regulates task specialization, and thus DOL in insect societies. To do so, we

compared the brain and antenna transcriptomes of nurses and foragers in the ant *T. longispinosus.* We

report several lines of evidence that support our hypothesis. First, we found almost seven times as many

genes to be differentially expressed between nurses and foragers in the antennae as in the brain, indicating

that peripheral sensory organs may be more important than the central nervous system in the task

specialization process of social insect workers. Second, we found half the OR genes in the *T. longispinosus*

genomes are differentially expressed between nurse and forager antennae, suggesting that behavioral

specialization is associated with different sensory filters that result in specific perceptions of the chemical

environment. Third, our analyses revealed that nurses and foragers upregulated distinct families of OR

genes, indicating that their sensory filters may target different types of chemical cues, possibly adjusted to

the tasks they perform. Finally, we detected several genes in multiple biogenic pathways to be differentially

expressed between the brains of nurses and foragers, potentially involved in fine-tuning the sensitivity of

the odorant filters.

Many organisms have evolved sensory filters to focus on only a subset of all environmental cues.

In insects, the peripheral olfactory filtering system enables individuals to detect and discriminate odors that

convey ecologically relevant information used to mediate important behaviors such as courtship, locomotion

and navigation to avoid predators and locate food or nesting sites (94). For example, the mosquito

Anopheles gambiae strongly responds to odorant components of its vertebrate hosts that provide meals

(95). Sensory filters have been selected for because they limit the amount of information perceived by the

organism, and thus the energy and time required by the brain to process it. Within insect colonies,

individuals typically exhibit many morphological and physiological traits associated with increased efficiency

in task specialization. They can be understood as adaptations that allow individuals to become more

proficient at their task due to learning, training, or the perceiving valuable task-related information, allowing

the colony to avoid the cost of task switching (96). In this context, it is interesting to find that *T. longispinosus*

workers may have different sensory filters that would serve as a basis for their task specialization, as

workers would mostly perceive chemical cues that pertain to their tasks. Our findings indicate that the

sensory filter of ant workers is dynamic, and its changes may underlie their behavioral maturation.

Age, genetic background, social environment, and individual experience produce behavioral

variation among workers that results in division of labor (13, 20, 23, 97). In our study, age in particular likely

differed between nurses and foragers. We propose that the factors known to regulate task specialization

actually drive different OR expression patterns, which in turn produce behavioral variation via contrasting

abilities to detect different sets of odors. This hypothesis is supported by several studies (reviewed in 97)

showing that the physiological condition of an animal can influence the level of receptor expression,

including mating status, oviposition, feeding, circadian rhythm, experience, and aging. Alternatively, we

cannot exclude that the exposure to different odors that is associated with performing different tasks may

at least in part have affected OR gene expression in the antennae. However, such an effect is unlikely to

explain the large-scale variation in gene expression, as there is very limited evidence that the mere

exposure to odors influences the expression of the gene coding for the OR that binds this odor (99, 100).

According to our sensory filter hypothesis, ant workers would be expected to primarily detect

chemical cues that correspond to their tasks. We found that the 9-exon subfamily of OR was

overrepresented in genes that were upregulated in the antennae of nurses. Interestingly, recent studies

have shown a rapid expansion of the 9-exon subfamily in ants, and several lines of evidence indicate that

OR genes from this subfamily mediate complex social interactions in ant colonies (65–67, 69). First,

comparisons of antennal transcriptomes revealed that 9-exon OR genes are expressed more frequently in

workers than in males, suggesting a role in social communication among workers (68, 69). Second,

representative OR from the 9-exon subfamily can detect CHC extracts from several castes (70). According

to McKenzie et al. (69), 9-exon OR genes were first expressed in solitary ancestors of aculeate wasps and

facilitated CHC discrimination, likely for prey or mate recognition, with a lineage giving rise to the ancestors

of ants. Moreover, OR genes of the 9-exon subfamily were convergently lost in socially parasitic ant species

that lost the ability to perform brood care or foraging (71), suggesting that they are essential for the

performance of these worker tasks. Our finding of an overexpression of 9-exon OR genes in the antennae

of nurses is in line with these previous studies, and suggests that many of these receptors have important

functions within the nest, such as sensing chemical cues from the larvae, queen, or other workers, and/or

that they are less important for sensing task-related stimuli or other signals outside the nest. Among the OR

genes overexpressed in nurses was also the co-receptor *Orco*, which is widely expressed in olfactory

sensory neurons and nearly unchanged in sequence in distant insect taxa (101–103). The overexpression

of *Orco* may indicate a generally more active odorant sensory system in the antennae of nurses compared

to foragers.

The behavioral transition from nursing to foraging may be triggered by a lower efficiency in detecting

brood cues via the downregulation of specific OR genes (e.g., from the 9-exon subfamily). This would result

in ants moving farther away from the brood, and this change in spatial location may trigger the behavioral

transition to outside tasks (104, 105). In addition to being less efficient at detecting brood cues, the sensory

filter of foragers may also become fine-tuned to detect a more diverse set of odors. Foragers overexpressed

a greater number of OR gene subfamilies compared to nurses (19 and 3 for foragers and nurses,

respectively), which may indicate that the olfactory system of foragers could be adjusted to the more diverse

chemical environment outside the nest. Similar to the 9-exon subfamily, the L subfamily has also been

expanded in social insects (62, 64, 65), and along with the P and H subfamilies, it has been lost in socially

parasitic ants (71). Interestingly, the OR genes from the L, P, and H subfamilies have been upregulated in

foragers, and thus may have a task-specific function, such as recognition of chemical cues related to

environmental perception or recruitment cues outside the nest. To date, however, there is no information

on which odors are recognized by these OR gene subfamilies.

Finding task-specific variation in OR gene expression raises the question as to which molecular

mechanisms regulate those changes. Previous studies in insects revealed that olfaction-guided behavior is

mediated by biogenic amine receptors in the antenna, and their expression is involved in fine-tuning the

sensitivity of the olfactory system (82, 98). For example, modified concentration cAMP and intracellular

Ca^{2+} levels due to octopamine-induced signal transduction in the moth *Manduca sexta* (106) activate Orco,

leading to changes in ORN sensitivity (107, 108). Our results suggest that the biogenic amine signaling

pathway may modulate the sensory filtering function of insect antennae and alter sensitivity to various

signals. We found that genes encoding tyramine and its precursor, octopamine, are upregulated in forager

antennae, similar to genes involved in serotonin signaling. Tyramine and dopamine (which was upregulated

in nurses) have been implicated in modulating taste and olfactory receptor neurons, while serotonin may

serve as a neurotransmitter and neurohormone in antennal vessels and mechanosensory organs (82).

Serotonin influences foraging activity (81) and regulates food intake in many animals (109–111). Dopamine

signaling also plays an important role in controlling the insect circadian clock and mediating clock-controlled

behavioral phenotypes such as locomotion (112, 113). In our focal species *T. longispinosus*, inside workers

were found to exhibit a stronger circadian rhythmicity than foragers, which may be regulated via differences

in the acetylation of histone proteins (114).

Changes in behavior and olfactory sensitivity in insects could be related to the expression of genes

involved in IIS, target of rapamycin (TOR), JH and Vg pathways, according to age, circadian rhythm, mating

and feeding status (98). For example, appetite state in *D. melanogaster* is signaled by insulin, which

upregulates a peptide receptor on the olfactory receptor cells that innervate the DM1 glomerulus. Activation

of the DM1 glomerulus is enough to drive the fly to reach for food (115). Recent studies have shown that

pheromone release and odor sensitivity appear to be under JH control in *Schistocerca gregaria* and *Locusta*

migratoria, which could lead to behavioral changes (116–120). Finally, an experimental downregulation of

Vg-like A in *T. longispinosus* workers resulted in decreased brood care behavior and a lower sensitivity to

brood-related chemical cues, suggesting changes in odor perception and olfactory-driven decision making

(42). Our results reveal that genes associated with all these pathways were differentially expressed in the

brain and antennae between nurses and foragers ,which confirms their role in task-associated behavioral

changes and regulation of odor perception. Given the central role of IIS, Vg, JH, and TOR pathways in

regulating division of labor in social insects, and our finding of task-associated patterns of the antennal

expression of genes from multiple biogenic amines, we hypothesize that these modulators and hormones

are involved in the regulation of the olfactory filter. How the detailed molecular mechanisms of these

pathways in the brain are causally linked to the complex changes in olfactory perception in the antennae

should therefore be investigated next.

Conclusion

Our transcriptomic analyses of the brain and antennae of *T. longispinosus* nurses and foragers provide

support to our hypothesis that behavioral variation and task specialization in ant workers are regulated via

differences in olfactory perception. We predict that antennal physiology acts as sensory filters that limit the

type and amount of chemical information passed to the brain. This would allow workers to target relevant

chemical information from the environment and discriminate signal from noise without using energetically

costly processing by the central nervous system. We argue that this sensory filter is flexible and regulated

through changes in physiological conditions such as age, nutrition, and hormones. Variation among workers

in their efficiency to detect specific chemical cues would result in task specialization and division of labor.

Our study opens novel avenues of research to better understand the role of sensory filters in controlling

DOL in insect societies.

Materials and Methods

Sample collection and behavioral determination

A total of seven colonies of the ant *T. longispinosus* were selected with an average colony size of 110 ± 31.5 workers (mean ± SD, Dataset S3). The ants were collected in the forests of the Edmund Niles Huyck Preserve, Rensselearville, NY, USA (42°31′41.0″N 74°09′38.8″W), in June of 2018 with permission. Upon collection, we housed each colony in a plaster-floored nesting box (43 cm × 28 cm × 10 cm) divided into three chambers containing a single slide nest, in which the colony relocated. A slide nest is an artificial nesting site comprised of a small Plexiglas cavity sandwiched between two glass microscope slides. Colonies were established at the Johannes Gutenberg University in Mainz, Germany, under a 14 h:10 h light:dark photoperiod at 18°C to a 22°C temperature. We provided honey and water ad libitum and fed crickets to the colony twice a week. To allow for visible behavioral division of labor between workers of the two behavioral phenotypes, we marked, observed and recaptured ants from inside and outside the nest. We defined foragers as workers that perform outside-nest tasks, including foraging for food, while nurses remained inside the dark nest and cared for the ant brood. A total of 69 workers inside (from the brood pile) and 76 workers outside the nest were marked with fine colored metal wires (0.02 mm Elektrisola, Eckenhagen, Germany) between the petiole and post-petiole. We performed behavioral observations every two hours, four times a day for five days (total = 20 scans), in which we noted down how many times an individual spent performing brood care and foraging behavior and the position in the nest (Table S3). Based on these behavioral observations, the marked individuals found outside the nest, exploring the surroundings for food or water, were identified as foragers. These workers usually do not care for the brood and do not frequently reside on brood piles, as ant colonies organize themselves spatially in a way that reduces contact between foragers and brood (121). We identified nurses as workers that remain inside the nest in direct contact with brood and were unlikely to leave the nest. Foragers were found in 42% ± SD 26% of the observations outside of the nest, whereas nurses spend only 1% ± SD 3% outside. In contrast, nurses were interacting with the brood in 54% ± SD 22% of the observations, whereas we found that foragers only did this only in 2% ± SD 4% of the observations. We focused in this study on individuals highly specialized on either foraging or brood care. Workers that performed both tasks regularly were not included in this study.

After all observations were completed, the marked nurses and foragers were collected, directly frozen in

liquid nitrogen and stored at -80 °C until further processing for dissection and pooling according to

behavioral state and colony.

RNA extraction and sequencing

For RNA extraction, we removed both antennae and stored them in a 1.5 ml Eppendorf tube containing 50

µl TRIzol (Invitrogen), cut the head off and fixed it on a slide with melted dental wax. We then made an

incision around the head with a surgical scalpel and removed the head capsule with forceps to expose the

intact brain. Finally, we carefully pulled the brain out of the head capsule and removed the remains of other

tissues that were connected to it. The dissected brain was transferred to a 1.5 ml Eppendorf tube containing

20µl PBS. Each dissection was completed in less than 5 minutes to prevent RNA degradation. We dissected

brain and antennae tissues from 48 nurses and 49 foragers. We pooled the brains and antennae from

seven workers from each behavioral state and colony. The only exception was the "GO" colony (NY18

E110), for which we pooled only six brains and twelve antennae from six nurses (Dataset S3) due to the

loss of one sample during the dissection process. Immediately after dissection of each brain and antennae,

the Eppendorf tubes were kept on dry ice while we dissected the remaining individuals. Brain and antennae

tissues were homogenized with a pestle. Sample brains were transferred separately to a 1.5 ml Eppendorf

tube containing 50µl of TRIzol. We added 50µl chloroform to each brain and antenna samples, cautiously

inverted for 30 s and then centrifuged samples at 12,000 g for 15 min at 4°C. We collected the resulting

supernatant and precipitated RNA with 25µl 70% ethanol. We conducted the subsequent RNA extraction

with the RNeasy Mini Kit (Qiagen), following the manufacturer's instruction. The resulting 28 samples (14

brains and 14 antennae) were stored at -80°C until library preparation.

RNA-seq libraries were prepared by Novogene Company Limited, Cambridge, UK, using the

NEBNext Ultra RNA Library Prep Kit for Illumina according to the manufacturer's protocol. After amplification

and purification, 28 libraries were sequenced on an Illumina NovaSeq 6000 S4 flow cell platform using a

paired-end 150 bp. Approximately 43 million raw reads were generated from each library (Dataset S3).

Gene expression analyses

Raw data obtained from Novogene were checked using FastQC v.0.11.9 (122), and Illumina adapters were

removed using Trimmomatic v.0.36 (123). The protein-coding genes of *T. longispinosus*

(GCA_004794745.1; (124) and the congener *T. curvispinosus* (GCA_003070985.1) were retrieved from the

NCBI database and transferred to the recently published *T. longispinosus* genome (71) using the Liftoff

v.1.6.1 tool (125). In total, 10,029 of 13,061 (~77%) annotated protein-coding genes were transferred from

the original *T. longispinosus* assembly (genes identified as "DBV15") and 4,808 were transferred from *T.*

curvispinosus (genes identified as "LOC"), for a total of 14,837. For gene expression analysis, reads were

mapped to our *T. longispinosus* genome assembly, and the read counts table was generated using STAR

2.7.0 (126) with default settings. Detailed mapping statistics for each sample is available in Dataset S3. We

used the deseq2 v1.16.1 package to identify differentially expressed genes (127). To avoid biased results

due to low read counts, we removed from the counting matrix those genes for which less than 10 of the

reads mapped to at least 6 of our 14 samples (n - 1 of the smallest sample size). Then, we conducted a

differential gene expression analysis with DESeq2 (128). We began with comparisons between nurses and

foragers using the ~Colony+Task model, followed by a likelihood ratio test (LRT) approach, with colony ID

as a fixed factor. Genes were considered differentially expressed if the false discovery rate (FDR), using

Benjamini-Hochberg procedure, had an adjusted p-value of ≤ 0.05. The resulting lists of DEG refer to genes

that are overexpressed and underexpressed in foragers compared to nurses. We used the online tool Venny

v.2.1 (https://bioinfogp.cnb.csic.es/tools/venny) to generate a Venn diagram containing the DEG associated

with task and tissues. Separation of differentially expressed genes by task was visualized by performing

principal component analysis (PCA) on transformed reads of filtered transcriptomes from all contigs using

the plotPCA function provided by DESeq2. Finally, ORs genes that were upregulated in each behavioral

phenotype was visualized in a volcano plot using the EnhancedVolcano package in R.

Identification of behavior candidate genes and odorant receptors

We used gene annotations based on a BlastX search of the *T. longispinosus* transcriptome compared to a

list of different invertebrate proteomes (i.e., *Acromyrmex echinatior, Apis mellifera, Camponotus floridanus,*

Drosophila melanogaster, Harpegnathos saltator, Odontomachus brunneus, Temnothorax curvispinosus)

downloaded from the NCBI database with an E-value of 1e-5 and below. Clusters containing more than one

sequence match per species were reduced to a single specimen based on the highest blast score. We

constructed orthogroups across all of the above species using OrthoFinder (129), including amino acid

sequences from the *T. longispinosus* proteome (124), and retained orthogroups containing caste DEG

(Dataset S4) to again compare potential behavioral candidate genes previously identified as involved in

regulating the division of labor in social insects (38, 41, 87, 93, 120, 130–133). GO enrichment analysis

was performed with TopGO v.2.44.0 using a Fisher's exact test for the different gene sets compared to the

whole genome with the weight01 algorithm (134). Only annotated GO terms with an adjusted p-value of ≤

0.05 were considered significantly enriched. All enrichment analyses were performed in RStudio v.1.4.1106

(135).

Odorant receptor (OR) protein sets were clustered across multiple ant species using OrthoFinder

to derive orthologous groups and identify subfamilies for each OR in *T. longispinosus*. To associate

orthogroups with previously identified OR subfamilies, we used OR annotation in *Atta cephalotes,*

Acromyrmex echinatior from Engsontia et al. (65), and *Camponotus floridanus*, *Harpegnathos saltator*, and

Solenopsis invicta from Zhou et al. (64, 68). Missing subfamily information was labeled as "unassigned"

(Dataset S2).

Acknowledgments

We thank Diego Páez-Moscoso for many helpful discussions and comments throughout the study; Marah

Stoldt and Carlotta Martelli for providing helpful comments on the manuscript; and Marion Kever and Jenny

Fuchs for technical assistance. This project was funded by the Deutsche Forschungsgemeinschaft (DFG,

German Research Foundation) – GRK2526/1 – Projectnr. 407023052.

References

1.   J. Maynard Smith, E. Szathmáry, The major evolutionary transitions in evolution. (Oxford
University Press, New York, US, 1995).
2.   A. Smith, An Inquiry into the Nature and Causes of the Wealth of Nations. (Chicago University
Press, 1776).
3.   J. W. Veening, *et al.*, Transient heterogeneity in extracellular protease production by *Bacillus*
subtilis. *Mol. Syst. Biol.* **4**, 1–15 (2008).
4.   V. L. Marlow, *et al.*, The prevalence and origin of exoproteaseproducing cells in the *Bacillus*
subtilis biofilm. *Microbiol.* **160**, 56–66 (2014).
5.   J. van Gestel, H. Vlamakis, R. Kolter, From cell differentiation to cell collectives: *Bacillus subtilis*
uses division of labor to migrate. *PLoS Biol.* **13**, 1–29 (2015).
6.   S. A. West, G. A. Cooper, Division of labour in microorganisms: an evolutionary perspective. *Nat.*
Rev. Microbiol. **14**, 716–723 (2016).
7.   G. F. Oster, E. O. Wilson, Caste and ecology in the social insects. *Acta Biotheor.* **28**, 234–235
(1979).
8.   E. O. Wilson, The ergonomics of caste in the social insects. *Am. Nat.* **102**, 41–66 (1968).
9.   E. O. Wilson, Behavioral discretization and the number of castes in an ant species. *Behav. Ecol.*
Sociobiol. **1**, 141–54 (1976).
10. J. M. Herbers, Social Organization in *Leptothorax* ants: within-and between-species patterns.
Psyche J. Entomol. **90**, 52489 (1983).
11. N. Pinter-Wollman, J. Hubler, J. A. Holley, N. R. Franks, A. Dornhaus, How is activity distributed
among and within tasks in *Temnothorax* ants? *Behav. Ecol. Sociobiol.* **66**, 1407–1420 (2012).
12. T. D. Seeley, Adaptive significance of the age polyethism schedule in honeybee colonies. *Behav.*
Ecol. Sociobiol. **11**, 287–293 (1982).
13. F. Tripet, P. Nonacs, Foraging for work and age-based polyethism: the roles of age and previous
experience on task choice in ants. *Ethology* **110**, 863–877 (2004).
14. G. B. Blanchard, G. M. Orledge, S. E. Reynolds, N. R. Franks, Division of labour and seasonality
in the ant *Leptothorax albipennis*: worker corpulence and its influence on behaviour. *Anim. Behav.*
59, 723–738 (2000).
15. A. L. Toth, G. E. Robinson, Worker nutrition and division of labour in honeybees. *Anim. Behav.* **69**,
427–435 (2005).
16. J. K. Wetterer, The ecology and evolution of worker size-distribution in leaf-cutting ants
(Hymenoptera: Formicidae). *Sociobiology* **34**, 119–144 (1999).
17. J. H. Fewell, R. E. Page, Genotypic variation in foraging responses to environmental stimuli by
honey bees, *Apis mellifera*. *Experientia* **49**, 1106–1112 (1993).
18. S. Graham, M. R. Myerscough, J. C. Jones, B. P. Oldroyd, Modelling the role of intracolonial
genetic diversity on regulation of brood temperature in honey bee (*Apis mellifera* L.) colonies.
Insectes Soc. **53**, 226–232 (2006).
19. R. Gove, M. Hayworth, M. Chhetri, O. Rueppell, Division of labour and social insect colony
performance in relation to task and mating number under two alternative response threshold
models. *Insectes Soc.* **56**, 319–331 (2009).
20. F. Ravary, E. Lecoutey, G. Kaminski, N. Châline, P. Jaisson, Individual Experience Alone Can
Generate Lasting Division of Labor in Ants. *Curr. Biol.* **17**, 1308–1312 (2007).
21. J. Gautrais, G. Theraulaz, J. L. Deneubourg, C. Anderson, Emergent polyethism as a
consequence of increased colony size in insect societies. *J. Theor. Biol.* **215**, 363–373 (2002).
22. D. Merkle, M. Middendorf, Dynamic polyethism and competition for tasks in threshold
reinforcement models of social insects. *Adapt. Behav.* **12**, 251–262 (2004).
23. R. Jeanson, J. H. Fewell, R. Gorelick, S. M. Bertram, Emergence of increased division of labor as
a function of group size. *Behav. Ecol. Sociobiol.* **62**, 289–298 (2007).
24. J. Y. Wakano, K. Nakata, N. Yamamura, Dynamic model of optimal age polyethism in social
insects under stable and fluctuating environments. *J. Theor. Biol.* **193**, 153–165 (1998).
25. Y. Iwasa, S. Yamaguchi, Task allocation in a cooperative society: specialized castes or age-
dependent switching among ant workers. *Sci. Rep.* **10**, 1–9 (2020).
26. E. J. H. Robinson, O. Feinerman, N. R. Franks, Flexible task allocation and the organization of
work in ants. *Proc. R. Soc. B Biol. Sci.* **276**, 4373–4380 (2009).

27.   H. Shimoji, N. Kasutani, S. Ogawa, M. K. Hojo, Worker propensity affects flexible task reversion in
495   an ant. *Behav. Ecol. Sociobiol.* **74** (2020).
496   28.   Y. Ben-Shahar, A. Robichon, M. B. Sokolowski, G. E. Robinson, Influence of gene action across
497   different time scales on behavior. *Science.* **296**, 741–744 (2002).
498   29.   Y. Ben-Shahar, H. T. Leung, W. L. Pak, M. B. Sokolowski, G. E. Robinson, cGMP-dependent
499   changes in phototaxis: A possible role for the foraging gene in honey bee division of labor. *J. Exp.*
500   *Biol.* **206**, 2507–2515 (2003).
501   30.   C. W. Whitfield, A. M. Cziko, G. E. Robinson, Gene expression profiles in the brain predict
502   behavior in individual honey bees. *Science.* **302**, 296–299 (2003).
503   31.   K. K. Ingram, P. Oefner, D. M. Gordon, Task-specific expression of the foraging gene in harvester
504   ants. *Mol. Ecol.* **14**, 813–818 (2005).
505   32.   C. Lucas, M. B. Sokolowski, Molecular basis for changes in behavioral state in ant social
506   behaviors. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 6351–6356 (2009).
507   33.   A. Zayed, G. E. Robinson, Understanding the relationship between brain gene expression and
508   social behavior: lessons from the honey bee. *Annu. Rev. Genet.* **46**, 591–615 (2012).
509   34.   J. P. Sullivan, O. Jassim, S. E. Fahrbach, G. E. Robinson, Juvenile hormone paces behavioral
510   development in the adult worker honey bee. *Horm. Behav.* **37**, 1–14 (2000).
511   35.   S. A. Ament, M. Corona, H. S. Pollock, G. E. Robinson, Insulin signaling is involved in the
512   regulation of worker division of labor in honey bee colonies. *Proc. Natl. Acad. Sci. U. S. A.* **105**,
513   4226–4231 (2008).
514   36.   A. G. Dolezal, C. S. Brent, B. Hölldobler, G. V. Amdam, Worker division of labor and endocrine
515   physiology are associated in the harvester ant, *Pogonomyrmex californicus*. *J. Exp. Biol.* **215**,
516   454–460 (2012).
517   37.   M. Corona, *et al.*, Vitellogenin underwent subfunctionalization to acquire caste and behavioral
518   specific expression in the harvester ant *Pogonomyrmex barbatus*. *PLoS Genet.* **9** (2013).
519   38.   P. Kohlmeier, A. R. Alleman, R. Libbrecht, S. Foitzik, B. Feldmeyer, Gene expression is more
520   strongly associated with behavioural specialization than with age or fertility in ant workers. *Mol.*
521   *Ecol.* (2019) https:/doi.org/10.1111/mec.14971.
522   39.   R. Libbrecht, *et al.*, Interplay between insulin signaling, juvenile hormone, and vitellogenin
523   regulates maternal effects on polyphenism in ants. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 11050–
524   11055 (2013).
525   40.   P. Kohlmeier, B. Feldmeyer, S. Foitzik, *Vitellogenin-like A*–associated shifts in social cue
526   responsiveness regulate behavioral task specialization in an ant. *PLoS Biol.* **16**, 1–26 (2018).
527   41.   D. S. Marco Antonio, K. R. Guidugli-Lazzarini, A. M. Do Nascimento, Z. L. P. Simões, K.
528   Hartfelder, RNAi-mediated silencing of vitellogenin gene function turns honeybee (*Apis mellifera*)
529   workers into extremely precocious foragers. *Naturwissenschaften* **95**, 953–961 (2008).
530   42.   J. Gospocic, *et al.*, The neuropeptide corazonin controls social behavior and caste Identity in ants.
531   *Cell* **170**, 748-759.e12 (2017).
532   43.   J. F. Kamhi, J. F. A. Traniello, Biogenic amines and collective organization in a superorganism:
533   neuromodulation of social behavior in ants. *Brain. Behav. Evol.* **82**, 220–236 (2013).
534   44.   D. Schulz, G. Robinson, Biogenic amines and division of labor in honey bee colonies: behaviorally
535   related changes in the antennal lobes and age-related changes in the mushroom bodies. *J. Comp.*
536   *Physiol. A Sens.* **5**, 481–488 (1999).
537   45.   A. R. Mercer, R. Menzel, The effects of biogenic amines on conditioned and unconditioned
538   responses to olfactory stimuli in the honeybee *Apis mellifera*. *J. Comp. Physiol.* **145**, 363–368
539   (1982).
540   46.   R. Scheiner, B. V. Entler, A. B. Barron, C. Scholl, M. Thamm, The effects of fat body tyramine
541   level on gustatory responsiveness of honeybees (*Apis mellifera*) differ between behavioral castes.
542   *Front. Syst. Neurosci.* **11**, 1–8 (2017).
543   47.   S. T. Ferguson, I. Bakis, L. J. Zwiebel, Advances in the study of olfaction in eusocial ants. *Insects*
544   **12** (2021).
545   48.   S. Martin, F. Drijfhout, A review of ant cuticular hydrocarbons. *J. Chem. Ecol.* **35**, 1151–1161
546   (2009).
547   49.   E. van Wilgenburg, M. R. E. Symonds, M. A. Elgar, Evolution of cuticular hydrocarbon diversity in
548   ants. *J. Evol. Biol.* **24**, 1188–1198 (2011).
549   50.   E. H. Slifer, The Structure of Arthropod Chemoreceptors. *Annu. Rev. Entomol.* **15**, 121–142

550   (1970).

51. H. Altner, L. Prillinger, Ultrastructure of invertebrate chemo-, thermo-, and hygroreceptors and its functional significance. *Int. Rev. Cytol.* **67**, 69–139 (1980).

52. E. Hallberg, B. S. Hansson, Arthropod sensilla: morphology and phylogenetic considerations. *Microsc. Res. Tech.* **47**, 428–439 (1999).

53. D. Wicher, F. Miazzi, Functional properties of insect olfactory receptors: ionotropic receptors and odorant receptors. *Cell Tissue Res.* **383**, 7–19 (2021).

54. W. S. Leal, Odorant reception in insects: roles of receptors, binding proteins, and degrading enzymes. *Annu. Rev. Entomol.* **58**, 373–391 (2013).

55. A. Sánchez-Gracia, F. G. Vieira, F. C. Almeida, J. Rozas, Comparative genomics of the major chemosensory gene families in arthropods. *eLS* (2011) https:/doi.org/10.1002/9780470015902.a0022848.

56. H. Breer, J. Fleischer, P. Pregitzer, J. Krieger, "Molecular mechanism of insect olfaction: olfactory receptors" In: Picimbon JF, editors. *Olfactory concepts of insect control-alternative to insecticides*. 93–114 (2019).

57. E. Jacquin-Joly, P. Lucas, Pheromone reception and transduction: mammals and insects illustrate converging mechanisms across phyla. *Curr Top Neurochem* **4**, 75–105 (2005).

58. R. K. Wilson, Z. F. Mainen, Early events in olfactory processing. *Annu. Rev. Neurosci.* **29**, 163–201 (2006).

59. S. Anton, U. Homberg, Antennal lobe structure. *Insect Olfaction*, 97–124 (1999).

60. B. S. Hansson, S. Anton, Function and morphology of the antennal lobe: new developments. *Annu. Rev. Entomol* **45**, 203–231 (2000).

61. J. S. de Belle, R. Kanzaki, Protocerebral Olfactory Processing. *Insect Olfaction*, 243–281 (1999).

62. R. Saad, A. B. Cohanim, M. Kosloff, E. Privman, Neofunctionalization in ligand binding sites of ant olfactory receptors. *Genome Biol. Evol.* **10**, 2490–2500 (2018).

63. P. R. Oxley, *et al.*, The genome of the clonal raider ant *Cerapachys Biroi*. *Curr. Biol.* **24**, 451–458 (2014).

64. X. Zhou, *et al.*, Chemoreceptor Evolution in Hymenoptera and its implications for the evolution of eusociality. *Genome Biol. Evol.* **7**, 2407–2416 (2015).

65. P. Engsontia, U. Sangket, H. M. Robertson, C. Satasook, Diversification of the ant odorant receptor gene family and positive selection on candidate cuticular hydrocarbon receptors. *BMC Res. Notes* **8**, 1–13 (2015).

66. C. D. Smith, *et al.*, Draft genome of the globally widespread and invasive Argentine ant (*Linepithema humile*). *Proc. Natl. Acad. Sci.* **108**, 5673–5678 (2011).

67. C. R. Smith, *et al.*, Draft genome of the red harvester ant *Pogonomyrmex barbatus*. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 5667–5672 (2011).

68. X. Zhou, *et al.*, Phylogenetic and transcriptomic analysis of chemosensory receptors in a pair of divergent ant species reveals sex-specific signatures of odor coding. *PLoS Genet.* **8** (2012).

69. S. K. McKenzie, I. Fetter-Pruneda, V. Ruta, D. J. C. Kronauer, Transcriptomics and neuroanatomy of the clonal raider ant implicate an expanded clade of odorant receptors in chemical communication. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 14091–14096 (2016).

70. G. M. Pask, *et al.*, Specialized odorant receptors in social insects that detect cuticular hydrocarbon cues and candidate pheromones. *Nat. Commun.* **8**, 1–10 (2017).

71. E. Jongepier, *et al.*, Convergent loss of chemoreceptors across independent origins of slave-making in ants. *Mol. Biol. Evol.* **39**, 1–14 (2022).

72. W. Trible, *et al.*, *orco* Mutagenesis causes loss of antennal lobe glomeruli and impaired social behavior in ants. *Cell* **170**, 727–735.e10 (2017).

73. H. Yan, *et al.*, An engineered *orco* mutation produces aberrant social behavior and defective neural development in ants. *Cell* **170**, 736–747.e9 (2017).

74. S. Beshers, J. H. Fewell, Models of Division of Labor in Social Insects. *Annu. Rev. Entomol.* **46**, 413–40 (2001).

75. P. Lichocki, D. Tarapore, L. Keller, D. Floreano, Neural networks as mechanisms to regulate division of labor. *Am. Nat.* **179**, 391–400 (2012).

76. D. Naug, From division of labor to collective behavior: behavioral analyses at different levels. *Behav. Ecol. Sociobiol.* **70**, 1113–1115 (2016).

77. R. Jeanson, A. Weidenmüller, Interindividual variability in social insects - proximate causes and

ultimate consequences. *Biol. Rev.* **89**, 671–687 (2014).

78. Y. Ulrich, *et al.*, Response thresholds alone cannot explain empirical patterns of division of labor in
social insects. *PLoS Biol.* **19**, 1–21 (2021).

79. E. Bonabeau, A. Sobkowski, G. Theraulaz, J. L. Deneubourg, Adaptive task allocation inspired by
a model of division of labor in social insects. *Biocomput. Emergent Comput.* **97**, 36–45 (1997).

80. E. Bonabeau, G. Theraulaz, J. L. Deneubourg, Fixed response thresholds and the regulation of
division of labor in insect societies. *Bull. Math. Biol.* **60**, 753–807 (1998).

81. D. J. Schulz, M. M. Elekonich, G. E. Robinson, Biogenic amines in the antennal lobes and the
initiation and maintenance of foraging behavior in honey bees. *J. Neurobiol.* **54**, 406–416 (2003).

82. M. I. Zhukovskaya, A. D. Polyanovsky, Biogenic amines in insect antennae. *Front. Syst. Neurosci.*
11, 1–9 (2017).

83. C. N. Cook, *et al.*, Individual differences in learning and biogenic amine levels influence the
behavioural division between foraging honeybee scouts and recruits. *J. Anim. Ecol.* **88**, 236–246
(2019).

84. D. O. Azevedo, S. O. De Paula, J. C. Zanuncio, L. C. Martinez, J. E. Serrão, Juvenile hormone
downregulates vitellogenin production in *Ectatomma tuberculatum* (Hymenoptera: Formicidae)
sterile workers. *J. Exp. Biol.* **219**, 103–108 (2016).

85. M. A. Rodrigues, T. Flatt, Endocrine uncoupling of the trade-off between reproduction and somatic
maintenance in eusocial insects. *Curr. Opin. insect Sci.* **16**, 1–8 (2016).

86. P. Jedlicka, U. R. Ernst, A. Votavová, R. Hanus, I. Valterová, Gene expression dynamics in major
endocrine regulatory pathways along the transition from solitary to social life in a bumblebee,
Bombus terrestris. Front. Physiol. **7**, 1–20 (2016).

87. C. Opachaloemphan, *et al.*, Early behavioral and molecular events leading to caste switching in
the ant *Harpegnathos. Genes Dev.* **35**, 410–424 (2021).

88. V. C. Norman, W. O. H. Hughes, Behavioural effects of juvenile hormone and their influence on
division of labour in leaf-cutting ant societies. *J. Exp. Biol.* **219**, 8–11 (2016).

89. G. E. Robinson, Regulation of honey bee age polyethism by juvenile hormone. *Behav. Ecol.*
Sociobiol. **20**, 329–338 (1987).

90. B. Fussnecker, C. Grozinger, Dissecting the role of Kr-h1 brain gene expression in foraging
behavior in honey bees (*Apis mellifera*). *Insect Mol. Biol.* **17**, 515–522 (2008).

91. C. M. Grozinger, G. E. Robinson, Endocrine modulation of a pheromone-responsive gene in the
honey bee brain. *J. Comp. Physiol. A Neuroethol. Sensory, Neural, Behav. Physiol.* **193**, 461–470
(2007).

92. H. Shpigler, *et al.*, The transcription factor Krüppel homolog 1 is linked to hormone mediated social
organization in bees. *BMC Evol. Biol.* **10** (2010).

93. J. Gospocic, *et al.*, Kr-h1 maintains distinct caste-specific neurotranscriptomes in response to
socially regulated hormones. *Cell* **184**, 5807-5823.e14 (2021).

94. E. von der Emde, G; Warrant, *The Ecology of Animal Senses Matched Filters for Economical*
Sensing, 1st ed (Springer International Publishing, 2016).

95. A. F. Carey, G. Wang, C. Y. Su, L. J. Zwiebel, J. R. Carlson, Odorant reception in the malaria
mosquito *Anopheles gambiae. Nature* **464**, 66–71 (2010).

96. A. Dornhaus, Specialization does not predict individual efficiency in an ant. *PLoS Biol.* **6**, 2368–
2375 (2008).

97. R. Libbrecht, L. Keller, Genetic compatibility affects division of labor in the argentine ant
Linepithema humile. Evolution **67**, 517–524 (2013).

98. C. Gadenne, R. B. Barrozo, S. Anton, Plasticity in insect olfaction: to smell or not to smell? *Annu.*
Rev. Entomol. **61**, 317–333 (2016).

99. S. Zhou, E. A. Stone, T. F. C. Mackay, R. R. H. Anholt, Plasticity of the chemoreceptor repertoire
in *Drosophila melanogaster. PLoS Genet.* **5** (2009).

100. B. Von Der Weid, *et al.*, Large-scale transcriptional profiling of chemosensory neurons identifies
receptor-ligand pairs in vivo. *Nat. Neurosci.* **18**, 1455–1463 (2015).

101. R. Benton, S. Sachse, S. W. Michnick, L. B. Vosshall, Atypical membrane topology and
heteromeric function of *Drosophila* odorant receptors in vivo. *PLoS Biol.* **4**, 240–257 (2006).

102. J. Krieger, O. Klink, C. Mohl, K. Raming, H. Breer, A candidate olfactory receptor subtype highly
conserved across different insect orders. *J. Comp. Physiol. A Neuroethol. Sensory, Neural, Behav.*
Physiol. **189**, 519–526 (2003).

103. M. C. Larsson, *et al.*, Or83b encodes a broadly expressed odorant receptor essential for
Drosophila olfaction. *Neuron* **43**, 703–714 (2004).
104. D. P. Mersch, A. Crespi, L. Keller, Tracking individuals shows spatial fidelity is a key regulator of
ant social organization. *Science.* **340**, 1090–1093 (2013).
105. T. O. Richardson, *et al.*, Ant behavioral maturation is mediated by a stochastic transition between
two fundamental states. *Curr. Biol.* **31**, 2253-2260.e3 (2021).
106. A. M. Dacks, J. B. Dacks, T. A. Christensen, A. J. Nighorn, The cloning of one putative
octopamine receptor and two putative serotonin receptors from the tobacco hawkmoth, *Manduca*
sexta. *Insect Biochem. Mol. Biol.* **36**, 741–747 (2006).
107. M. N. Getahun, S. B. Olsson, S. Lavista-Llanos, B. S. Hansson, D. Wicher, Insect odorant
response sensitivity is tuned by metabotropically autoregulated olfactory receptors. *PLoS One* **8**,
1–9 (2013).
108. M. Stengl, N. W. Funk, The role of the coreceptor Orco in insect olfactory transduction. *J. Comp.*
Physiol. A Neuroethol. Sensory, Neural, Behav. Physiol. **199**, 897–909 (2013).
109. J. E. Blundell, J. C. G. Halford, Serotonin and appetite regulation: Implications for the
pharmacological treatment of obesity. *CNS Drugs* **9**, 473–495 (1998).
110. A. S. French, *et al.*, The role of serotonin in feeding and gut contractions in the honeybee. *J. Insect*
Physiol. **61**, 8–15 (2014).
111. J. P. Voigt, H. Fink, Serotonin controlling feeding and satiety. *Behav. Brain Res.* **277**, 14–31
(2015).
112. R. J. Beninger, The role of dopamine in locomotor activity and learning. *Brain Res. Rev.* **6**, 173–
196 (1983).
113. X. Liang, *et al.*, Morning and evening circadian pacemakers independently drive premotor centers
via a specific dopamine relay. *Neuron* **102**, 843-857.e4 (2019).
114. R. Libbrecht, D. Nadrau, S. Foitzik, A role of histone acetylation in the regulation of circadian
rhythm in ants. *iScience* **23**, 100846 (2020).
115. C. M. Root, K. I. Ko, A. Jafari, J. W. Wang, Presynaptic facilitation by neuropeptide signaling
mediates odor-driven food search. *Cell* **145**, 133–144 (2011).
116. S. Anton, M. C. Dufour, C. Gadenne, Plasticity of olfactory-guided behaviour and its
neurobiological basis: lessons from moths and locusts. *Entomol. Exp. Appl.* **123**, 1–11 (2007).
117. A. I. Tawfik, E. O. Osir, A. Hassanali, S. H. Ismail, Effects of juvenile hormone treatment on phase
changes and pheromone production the desert locust, *Schistocerca gregaria* (Forskal)
(Orthoptera: Acrididae). *J. Insect Physiol.* **43**, 1177–1182 (1997).
118. A. I. Tawfik, K. Treiblmayr, A. Hassanali, E. O. Osir, Time-course haemolymph juvenile hormone
titres in solitarious and gregarious adults of *Schistocerca gregaria*, and their relation to pheromone
emission, CA volumetric changes and oocyte growth. *J. Insect Physiol.* **46**, 1143–1150 (2000).
119. G. Wiesel, S. Tappermann, A. Dorn, Effects of juvenile hormone and juvenile hormone analogues
on the phase behaviour of *Schistocerca gregaria* and *Locusta migratoria*. *J. Insect Physiol.* **42**,
385–395 (1996).
120. W. Guo, *et al.*, Juvenile hormone suppresses aggregation behavior through influencing antennal
gene expression in locusts. *PLoS Genet.* **16**, 1–18 (2020).
121. T. Pamminger, S. Foitzik, K. C. Kaufmann, N. Schützler, F. Menzel, Worker personality and its
association with spatially structured division of labor. *PLoS One* **9**, 1–8 (2014).
122. S. Andrews, A quality control tool for high throughput sequence data (2016).
123. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics **30**, 2114–2120 (2014).
124. R. Kaur, *et al.*, Ant behaviour and brain gene expression of defending hosts depend on the
ecological success of the intruding social parasite. *Philos. Trans. R. Soc. B Biol. Sci.* (2019)
https:/doi.org/10.1098/rstb.2018.0192.
125. A. Shumate, S. L. Salzberg, Liftoff: accurate mapping of gene annotations. *Bioinformatics* **37**,
1639–1643 (2021).
126. A. Dobin, *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).
127. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome Biol.* **15**, 550 (2014).
128. M. I. Love, S. Anders, W. Huber, Differential analysis of count data - the DESeq2 package.
Genome Biology **15**: 550 (2014).

718
129. D. M. Emms, S. Kelly, OrthoFinder: solving fundamental biases in whole genome comparisons
dramatically improves orthogroup inference accuracy. *Genome Biol.* **16**, 157 (2015).
130. B. Das, C. de Bekker, Time-course RNASeq of *Camponotus floridanus* forager and nurse ant
brains indicate links between plasticity in the biological clock and behavioral division of labor. *BMC*
Genomics **23**, 1–23 (2022).
131. A. Alleman, M. Stoldt, B. Feldmeyer, S. Foitzik, Tandem-running and scouting behaviour are
characterized by up-regulation of learning and memory formation genes within the ant brain. *Mol.*
Ecol. **28**, 2342–2359 (2019).
132. J. Korb, *et al.*, Comparative transcriptomic analysis of the mechanisms underpinning ageing and
fecundity in social insects. *Philos. Trans. R. Soc. B Biol. Sci.* **376** (2021).
133. B. Qiu, *et al.*, Canalized gene expression during development mediates caste differentiation in
ants. *Nat. Ecol. Evol.* **6**, 1753–1765 (2022).
134. A. Alexa, J. Rahnenführer, Gene set enrichment analysis with topGO. R package version 2.28.0
(2016).
135. RStudio Team, RStudio: Integrated Development for R. *RStudio, PBC, Boston, MA* (2020).
734

Figures and Tables

Figure 1. Variation in behavior, brain and antenna gene expression between nurses and foragers.

(*A*) Boxplot showing behavioral differences between ants selected for transcriptomic analysis. Each black

dots represent individual ants. For differential expression analysis, each sample contains the pooled RNA

of seven ants of the respective behavioral phenotype. (*B*) Venn diagrams showing the number of DEG that

were overexpressed in nurses (red) and foragers (turquoise) for both antenna and brain tissues. Principal

component analysis (PCA) plots based on all expressed genes for (*C*) antenna and (*D*) brain samples. The

color of each sample represents task (red = nurse, turquoise = forager), and the shape the colony of origin.

The ellipses were defined by a 95% confidence interval and show how similar/different the samples are

within task groups.

Figure 2. OR expression in the antennae differs between nurses and foragers. (*A*) Among the DEG overexpressed in nurses and foragers, the most represented OR subfamilies are indicated by different colors. The red dotted line represents the significance threshold of our differential expression analysis. (*B*,*C*). Boxplots representing the expression of (*B*) *Orco* and *OR257*, the gene with the highest expression difference in terms of log$_2$FC in the 9-exon subfamily, and (*C*) *OR041*, *OR045* and *OR318*, the genes with the highest expression difference in terms of log$_2$FC in the H, L and P subfamilies, respectively.

754 **Figure 3. Gene expression differences between nurse and forager antennae.** Boxplots representing

755 the expression of (*A-E*) selected genes in biogenic amines pathways, as well as (*F*) one behavioral

756 candidate gene (Supplementary Table S1). Gene identity is shown in parentheses.

Supporting Information

Table S1. List of candidate genes related to behavior phenotypes.

Gene Annotation	Locus (*T. longispinosus*)	Organism reference	Locus reference	Orthogroup	Task	Tissue	Pathways and genes	References
protein takeout isoform	DBV15_08771	*Camponotus floridanus*	LOC105257386	OG0007297	Nurse	Brain	JH: signalling	(1)
vitellogenin-1 (Vg conventional)	LOC112466671	*Harpegnathos saltator*	LOC105181726	OG0003549	Nurse	Brain	Vitellogenin	(1–4)
vitellogenin-2-like (Vg like-A)	DBV15_03138	*H. saltator*	LOC105191299	OG0009030	Nurse	Brain	Vitellogenin	(2, 3)
farnesyl pyrophosphate synthase	LOC112466955	*Drosophila melanogaster*	Fpps	OG0000401	Nurse	Brain	JH: biosynthesis	(5)
venom carboxylesterase-6 (Jhe1)	DBV15_11528	*H. saltator*	LOC112590013	OG0002262	Nurse	Brain/Antenna	JH: regulation of JH biosynthesis	(1, 3)
ribosomal protein S6	DBV15_00490	*D. melanogaster*	RpS6	OG0007546	Nurse	Brain/Antenna	IIS / TOR	(5)
dopamine 1-like receptor 2	DBV15_07611	*D. melanogaster*	Dop1R2	OG0001377	Nurse	Antenna	bioagenic amines	(6)
octopamine beta2 receptor	DBV15_10418	*D. melanogaster*	Octbeta2R	OG0000466	Nurse	Antenna	bioagenic amines	(6)

aldehyde dehydrogenase	DBV15_01763	*D. melanogaster*	Aldh-III	OG0000175	Nurse	Antenna	JH: biosynthesis	(5)
cyclin G	DBV15_03888	*D. melanogaster*	CycG	OG0003395	Nurse	Antenna	IIS / TOR	(5)
happyhour	DBV15_11653	*D. melanogaster*	hppy	OG0000352	Nurse	Antenna	IIS / TOR	(5)
insulin receptor isoform (InRL)	DBV15_10224	*H. saltator*	LOC105183944	OG0001487	Nurse	Antenna	IIS	(3)
Pi3K21B	DBV15_03081	*D. melanogaster*	Pi3K21B	OG0002238	Nurse	Antenna	IIS	(5)
Ras oncogene at 64B	DBV15_02802	*D. melanogaster*	Ras64B	OG0007075	Nurse	Antenna	IIS / TOR	(5)
sarcoplasmic calcium-binding protein 2	DBV15_05072	*D. melanogaster*	Scp2	OG0005773	Nurse	Antenna	JH: metabolism	(5)
shaggy	DBV15_09601	*D. melanogaster*	sgg	OG0000404	Nurse	Antenna	IIS / TOR	(3, 5)
allatostatin A	LOC112454443	*D. melanogaster*	AstA	OG0007104	Forager	Brain	JH: regulation of JH biosynthesis	(5)
insulin-like growth factor I (IGF1)	LOC112454447	*H. saltator*	LOC105186969	OG0004678	Forager	Brain/Antenna	IIS	(3, 4)
5-hydroxytryptamine (serotonin) receptor 2A	DBV15_11483	*D. melanogaster*	5-HT2A	OG0000533	Forager	Antenna	bioagenic amines	(6)
octopamine receptor in mushroom bodies	LOC112465659	*D. melanogaster*	Oamb	OG0001501	Forager	Antenna	bioagenic amines	(6)

tyramine beta hydroxylase	DBV15_00422	*D. melanogaster*	Tbh	OG0006646	Forager	Antenna	bioagenic amines	(6)
density regulated protein	DBV15_10283	*D. melanogaster*	DENR	OG0004331	Forager	Antenna	IIS / TOR	(5)
dreadlocks	DBV15_03530	*D. melanogaster*	dock	OG0001746	Forager	Antenna	IIS / TOR	(5)
juvenile hormone-inducible protein 26	DBV15_02220	*D. melanogaster*	JhI-26	OG0000832	Forager	Antenna	JH: regulation of JH biosynthesis	(5)
Krueppel homolog 1-like	DBV15_06330	*H. saltator*	LOC112589664	OG0008125	Forager	Antenna	JH: signalling	(3, 4)
rapamycin-insensitive companion of Tor	DBV15_07973	*D. melanogaster*	rictor	OG0003860	Forager	Antenna	IIS / TOR	(5)
SAPK-interacting protein 1	DBV15_07343	*D. melanogaster*	Sin1	OG0005622	Forager	Antenna	IIS / TOR	(5)

Table S2. List of enriched GO biological process terms of the DEG in the brain and the antenna.

Gene Set (DEG)	Tissue	GO ID	Term	Annotated	Significant	Expected	Fisher (adjusted *p*-value)
Nurse	Brain	GO:0006412	translation	73	19	1.79	0.000
Nurse	Brain	GO:0006879	cellular iron ion homeostasis	5	2	0.12	0.006
Nurse	Brain	GO:0009056	catabolic process	155	6	3.79	0.024
Nurse	Brain	GO:0030155	regulation of cell adhesion	1	1	0.02	0.025
Nurse	Brain	GO:0010040	response to iron (II) ion	1	1	0.02	0.025
Nurse	Brain	GO:0030245	cellulose catabolic process	1	1	0.02	0.025
Nurse	Brain	GO:0009204	deoxyribonucleoside triphosphate catabolic process	1	1	0.02	0.025
Nurse	Brain	GO:0006559	L-phenylalanine catabolic process	2	1	0.05	0.048
Nurse	Brain	GO:0006572	tyrosine catabolic process	2	1	0.05	0.048
Nurse	Brain	GO:0007160	cell-matrix adhesion	2	1	0.05	0.048
Nurse	Brain	GO:0006546	glycine catabolic process	2	1	0.05	0.048
Nurse	Brain	GO:0019236	response to pheromone	2	1	0.05	0.048
Nurse	Brain	GO:0030334	regulation of cell migration	2	1	0.05	0.048
Forager	Brain	GO:0006518	peptide metabolic process	89	4	0.96	0.001

Forager	Brain	GO:0006431	methionyl-tRNA aminoacylation	1	1	0.01	0.011
Forager	Brain	GO:0032481	positive regulation of type I interferon production	1	1	0.01	0.011
Forager	Brain	GO:0045454	cell redox homeostasis	2	1	0.02	0.022
Forager	Brain	GO:0046855	inositol phosphate dephosphorylation	4	1	0.04	0.043
Nurse	Antenna	GO:0006412	translation	73	22	8.84	0.000
Nurse	Antenna	GO:0006897	endocytosis	15	6	1.82	0.006
Nurse	Antenna	GO:0006006	glucose metabolic process	10	4	1.21	0.006
Nurse	Antenna	GO:0051726	regulation of cell cycle	9	3	1.09	0.014
Nurse	Antenna	GO:0015012	heparan sulfate proteoglycan biosynthetic process	3	2	0.36	0.040
Nurse	Antenna	GO:0006886	intracellular protein transport	63	13	7.63	0.047
Forager	Antenna	GO:0046854	phosphatidylinositol phosphorylation	6	4	0.57	0.001
Forager	Antenna	GO:0046855	inositol phosphate dephosphorylation	4	3	0.38	0.003
Forager	Antenna	GO:0005975	carbohydrate metabolic process	81	17	7.74	0.004
Forager	Antenna	GO:0045087	innate immune response	2	2	0.19	0.009
Forager	Antenna	GO:0009166	nucleotide catabolic process	4	3	0.38	0.025
Forager	Antenna	GO:0007229	integrin-mediated signaling pathway	9	3	0.86	0.045

| Forager | Antenna | GO:0006166 | purine ribonucleoside salvage | 4 | 2 | 0.38 | 0.047 |

Table S3. Behaviors and positions annotations during nest scans. Nursing was defined as the number of observations a worker antennating, grooming, feeding or carrying a brood item, and its position inside the nest (on the brood or near to the brood pile). Foraging was defined as the number of observations an individual was found outside the nest, collecting food or water.

Position	
Inside the nest	On the brood pile
	Near the brood pile
	Surroundings of the nest
	Near the entrance
Outside the nest	Chamber 1
	Chamber 2
	Chamber 3
Behavior	
Brood care	Antennation, grooming, feeding or carrying the brood
Nestmate care	Antennation, grooming, feeding or carrying a nestmate
Forager	Collecting food/drinking water outside the nest
Others	Resting
	Walking
	Being groomed or fed
	Grooming itself

Dataset S1 (separate file). Gene list with caste-differential expression, NCBI BlastX results, and gene read count of 28 RNA-seq samples.

Dataset S2 (separate file). OR subfamilies and orthogroups from the DEG lists between nurses and foragers in the antenna.

Dataset S3 (separate file). Summary of ant brain and antenna transcriptome data sets and mapping results.

Dataset S4 (separate file). List of orthogroups from the DEG list between nurses and foragers in *Temnothorax longispinosus*.

SI References

1.   B. Das, C. de Bekker, Time-course RNASeq of *Camponotus floridanus* forager and nurse ant brains indicate links between plasticity in the biological clock and behavioral division of labor. *BMC Genomics* **23**, 1–23 (2022).
2.   P. Kohlmeier, B. Feldmeyer, S. Foitzik, *Vitellogenin-like A*–associated shifts in social cue responsiveness regulate behavioral task specialization in an ant. *PLoS Biol.* **16**, 1–26 (2018).
3.   J. Gospocic, *et al.*, Kr-h1 maintains distinct caste-specific neurotranscriptomes in response to socially regulated hormones. *Cell* **184**, 5807-5823.e14 (2021).
4.   C. Opachaloemphan, *et al.*, Early behavioral and molecular events leading to caste switching in the ant *Harpegnathos*. *Genes Dev.* **35**, 410–424 (2021).
5.   J. Korb, *et al.*, Comparative transcriptomic analysis of the mechanisms underpinning ageing and fecundity in social insects. *Philos. Trans. R. Soc. B Biol. Sci.* **376** (2021).
6.   L. Gramates, *et al.*, FlyBase: a guided tour of highlighted features. *Genetics* **220**, iyac035 (2022).