1 **A highly discriminatory RNA strand-specific assay to facilitate analysis of the role of *cis-***

2 **acting elements in foot-and-mouth disease virus replication**

3 Samuel J. Dobson, Joseph C. Ward, Morgan R. Herod, David J. Rowlands, Nicola J. Stonehouse

4 School of Molecular and Cellular Biology and Astbury Centre for Structural Molecular Biology, Faculty

5 of Biological Sciences, United Kingdom

Abstract

Foot-and-mouth-disease virus (FMDV), the etiological agent responsible for foot-and-mouth disease

(FMD), is a member of the genus *Aphthovirus* within the *Picornavirus* family. In common with all

picornaviruses, replication of the single-stranded positive-sense RNA genome involves synthesis of a

negative-sense complementary strand that serves as a template for the synthesis of multiple positive-

sense progeny strands. We have previously employed FMDV replicons to examine viral RNA and

protein elements essential to replication, however, the factors affecting differential strand production

remain unknown. Replicon-based systems require transfection of high levels of RNA, which can

overload sensitive techniques such as qPCR preventing discrimination of specific strands. Here, we

describe a method in which replicating RNA is labelled *in vivo* with 5-ethynyl uridine. The modified

base is then linked to a biotin tag using click chemistry, facilitating purification of newly synthesised

viral genomes or anti-genomes from input RNA. This selected RNA can then be amplified by strand-

specific qPCR, thus enabling investigation of the consequences of defined mutations on the relative

synthesis of negative-sense intermediate and positive-strand progeny RNAs. We apply this new

approach to investigate the consequence of mutation of viral *cis*-acting replication elements and

provide direct evidence for their roles in negative-strand synthesis.

Keywords

23 Foot-and-mouth disease virus; FMDV; replication; strand-specific assay; negative sense; cre

Introduction

The *Aphthovirus* foot-and-mouth disease virus (FMDV) is a member of the *Picornaviridae* family of

single-stranded positive sense RNA viruses. FMDV is the etiological agent of foot-and-mouth disease

(FMD) and comprises six serotypes; A, O, Asia 1, Southern African Territories (SAT) 1, SAT 2 and SAT 3,

which are together responsible for endemic infection in large parts of Africa, Asia and the Middle East

(1). Whilst FMD is rarely fatal, its consequences of reduced animal productivity, restriction of trade

and slaughter of infected and at-risk animals result in severe economic losses (2, 3). The threat to

maintenance of virus-free status in non-endemic regions is exacerbated by the high transmissibility of

FMDV. Control measures following introduction of FMD into non-endemic regions, including strict

movement restriction and mass culling, can result in costs of £billions (3-5). The antigenic diversity of

FMDV adds to the challenges of disease control, with little cross-protection provided by strain-specific

vaccines and no effective therapeutics currently available (2, 6-10). There is therefore a need to better

understand the viral lifecycle in order to develop novel methods of control.

The genome of FMDV is approximately 8.4 kb and encodes a single polyprotein that is processed by

viral proteases to generate the proteins required for genome replication and encapsidation. Primary

cleavage of the polyprotein occurs at three positions to generate four products, the leader protease

(L^{pro}), the viral structural protein precursor P1-2A and the non-structural protein precursors P2 (2BC)

and P3 ($3AB_{1,2,3}CD$). Cleavage of L^{pro} occurs auto-catalytically, while release of the P1-2A precursor

from the polyprotein occurs through a co-translational 2A-dependent ribosome skipping mechanism.

The P1-2A region contains the viral structural proteins VP1, VP3 and VP0 (which is cleaved into VP2

and VP4 during virion maturation). Subsequent processing of the P2 and P3 precursors generates the

viral non-structural proteins 2B, 2C and 3A, $3B_{1,2+3}$ (VPg), 3C protease ($3C^{pro}$) and 3D polymerase ($3D^{pol}$)

(11-21). This processing is thought to be mediated by $3C^{pro}$ through alternative *cis-* and *trans-*

pathways that appear to co-ordinate replication (17, 22, 23).

The single viral open reading frame (ORF) is flanked by untranslated regions (UTRs) at the 5' and 3'

ends of the genome. The FMDV 5' UTR is longest amongst known picornaviruses and contains several

highly structured domains that are essential to the virus lifecycle (24-30). The S fragment is a ~360

nucleotide hairpin-loop that occupies the same location (at the extreme 5' end of the 5' UTR) as the

cloverleaf found in enteroviruses (31-34). It appears to be involved in immune modulation and viral

replication via interactions with host and viral proteins (27, 35, 36). Adjacent to the S-fragment is a

poly-C sequence of unknown function, followed by two to four RNA pseudoknots, which appear to be

involved in assembly of infectious virions in addition to genome replication (24, 37-39). Adjacent to

the pseudoknots is a small hairpin-loop known as the *cis-*acting replication element (*cre*) that in other

picornaviruses is located within the ORF (25, 40). Whilst the location of the *cre* appears not important,

it has an indispensable role in replication (25), acting as a template for the uridylylation of VPg by $3D^{pol}$

(41-43). VPg$_p$U$_p$U primes genome replication by $3D^{pol}$ (22, 44, 45). Finally, the internal ribosome entry

site (IRES) is located immediately upstream of the ORF and is essential for initiation of translation (26,
29, 30, 46).

Picornavirus genome replication is initiated by the synthesis of a negative-sense copy of the infecting
genome followed by assembly of a 'replicative form' (47-49). This complex comprises the negative
strand template RNA and several nascent progeny positive strand RNAs together with $3D^{pol}$ and other
viral and cellular proteins (50, 51). There is evidence that in the native state of the replicative form the
nascent daughter strands are not collapsed onto the negative template strand but are only transiently
associated at the site of transcription (50). The synthesis of both negative and positive RNA strands
appears to be primed with VPg_pU_pU (41).

While it is well documented that the *cre* element is essential for viral replication, it is not firmly
established whether this is required to produce VPgpUpU for the priming of both negative- and
positive-sense RNA molecules during intra-cellular virus replication. Indeed, it has been reported that
priming of negative-strand RNA synthesis can occur in a *cre*-independent manner during cell-free
replication of poliovirus, possibly via the poly(A) tail (52-54). As mutation of the *cre* is lethal to progeny
virus production, methods that facilitate the initiation of replicative events but are not reliant on
infection (such as transfection of *in vitro* transcribed viral RNA) are useful for study of the initial steps
in replication. Whilst assays to distinguish positive- and negative-strand RNAs have been developed
for several viruses including FMDV, o'nyong-nyong virus, dengue virus, murine norovirus and
chikungunya virus, their applications have so far been limited to studies involving infectious virus
production (55-59).

Replicons are mini-genomes in which the structural genes are replaced with a reporter gene to
facilitate the study of RNA replication independent of other aspects of the virus lifecycle (60). In
addition to their application in the dissection of the molecular details of replication, replicons permit
the study of viruses that require high-containment facilities, such as FMDV, at lower laboratory
containment. However, because the delivery of RNA by transfection is an inefficient process, large
quantities of *in vitro* transcribed RNA are used in order to ensure that sufficient cells are transfected.
This can overload subsequent strand-specific assays. Here, we describe a strand-specific qPCR assay
using FMDV replicons to determine the effects of mutations on the synthesis of both negative and
positive strands. We have applied this method to determine the role of the *cre* element in the initiation
of synthesis of both positive- and negative-strand RNAs.

Materials and methods

Cell lines and maintenance

Baby hamster kidney (BHK-21) cells were purchased from ATCC (LGC Standard) and were maintained in Dulbecco's modified Eagle's medium (DMEM) with glutamine, supplemented with 10% (v/v) foetal bovine serum (FBS), 50 U/mL penicillin and 50 µg/mL streptomycin. Growth medium and supplements were purchased from Sigma-Aldrich, Merck.

In vitro transcription

The construction, linearisation and purification of FMDV replicon plasmids has been described previously (24, 60). Linear plasmids were transcribed *in vitro* using the T7 RiboMAX Express Large Scale RNA Production System (Promega) using 'half sized' reactions whereby 250 ng linear plasmid was added to a reaction mixture using half the volume suggested for all reagents in the manufacturer's protocol. Reactions were incubated at 37°C for 1.5 hours prior to addition of 1 µL DNase I and further incubation at 37°C for 20 minutes. Resulting RNA was purified using the RNA Clean & Concentrator-25 kit (Zymo Research) prior to quantification by NanoDrop (Thermo Fisher) and confirmation of integrity by denaturing MOPS-formaldehyde gel electrophoresis.

Replication assays

BHK-21 cells were seeded into 6-well plates at a density of 1×10^6 cells/well and incubated for 16 hours. After incubation, cells were transfected with *in vitro* transcribed replicon RNA using Lipofectamine 2000 (Thermo Fisher). RNA (0.5 µg/cm^2) and Lipofectamine 2000 (1 µg RNA:3 µL reagent) were added separately to 50 µL opti-MEM (Thermo Fisher) and incubated for 10 minutes at room temperature. RNA and reagent were mixed and incubated for 20 minutes at room temperature before diluting with phenol red free DMEM (Thermo Fisher) supplemented with 10% (v/v) FBS and GlutaMAX (Thermo Fisher) diluted to 1x. Cells were washed 1x using PBS, before addition of transfection mixture. Cells were imaged over time using the 10X objective of an Incucyte S3 live cell imaging system to detect phase and green fluorescence. Analysis was performed with the inbuilt Incucyte 2021A analysis suite using surface fit segmentation, threshold minimum (green calibrated units, GCU) 8.0, edge split sensitivity -30, minimum area filter 50 µm^2 to determine green fluorescent cells. Following incubation for 6 hours, cells were washed 1x with PBS and extracted by addition of 1 mL TRI reagent (Merck) directly to the monolayer. Harvested cell extracts were stored at -20°C until further processing.

RNA extraction and cDNA synthesis

Total RNA was TRI reagent extracted from BHK-21 cells using Phasemaker tubes (Invitrogen, Thermo Fisher) and associated manufacturer protocol. An additional 75% EtOH RNA pellet wash step was performed to ensure removal of trace contaminants prior to RNA solubilisation using nuclease-free H_2O. Contaminating DNA was removed using the TURBO DNA-*free* kit (Thermo Fisher) following manufacturer guidelines. Total RNA purity and concentration was determined by NanoDrop.

cDNA synthesis was performed using SuperScript IV Reverse Transcriptase (Thermo Fisher) following manufacturer guidelines unless stated otherwise. Separate reactions were performed using 2 µM strand-specific reverse transcription (ssRT) primer (Table 1, PS-tag-RT or NS-tag-RT) or 50 µM oligo(dT)$_{20}$. A total mass of 500 ng RNA was added to each reaction, together with dNTP mix containing 10 mM of each nucleotide (Promega). Combined reaction mixtures were incubated at 50^{o}C for 10 minutes prior to inactivation at 85^{o}C for 10 minutes.

qPCR assays

cDNA synthesised from total RNA was diluted 100-fold in nuclease-free H_2O unless otherwise stated. qPCR was performed using GoTaq qPCR Master Mix (Promega). Reaction mixes were made up in 10 µL total volume containing 500 nM forward and reverse primers (Table 1, PS-qPCR-tag & PS-qPCR-R or NS-qPCR-tag & NS-qPCR-R or β-actin-F & β-actin-R) and 2 µL diluted cDNA. qPCR conditions followed the manufacturer recommended fast cycling program with polymerase activation (95^{o}C, 2 minutes) followed by 40 cycles (denaturation at 95^{o}C, 3 seconds and annealing/extension at 60^{o}C, 30 seconds) with melt curve analysis on a CFX Connect Real-Time PCR Detection System (Bio-Rad).

Nascent RNA labelling assays

Nascent RNA labelling assays were performed using the Click-iT™ Nascent RNA Capture Kit (Thermo Fisher). Replication assays were performed as described previously in 6-well plates, with addition of 0.2 mM 5-ethynyl uridine (5-EU) diluted from a 200 mM stock, to transfection complex mixtures immediately prior to addition to cells. Cells were harvested 6 hours post transfection through the addition of 1 mL TRI reagent (Sigma-Aldrich, Merck) directly to the monolayer, with collected samples stored at -20^{o}C until further processing. RNA was extracted as indicated above. Biotinylation of RNA by click reaction was performed following manufacturer guidelines using 10 µg total RNA and 0.5 mM biotin azide in a 50 µL reaction. RNA was precipitated in ammonium acetate overnight at -80^{o}C, pelleted, washed and solubilised in nuclease-free H_2O, as recommended. RNA was quantified by NanoDrop and 1 µg biotinylated RNA mixed with 12 µL Dynabeads® MyOne™ Streptavidin T1 magnetic beads. Following 30 minutes incubation at room temperature with agitation at 600 rpm, beads were

washed for 3 minutes at room temperature with agitation at 600 rpm 5x using washer buffer 1 and 5x

using wash buffer 2, with magnetic capture performed between washes. Bead mixtures were

transferred to a new tube after every three washes to minimise tube contaminant carry over.

Following the final wash, beads were resuspended in 36 μL wash buffer 2 and 12 μL slurry divided

between three separate tubes for separate reverse transcription of positive-strand genomes,

negative-strand genomes and total poly(A) RNA. Bead suspensions were heated at $70^{\circ}C$ for 5 minutes

prior to immediate addition of 17 μL master mix containing 15 μL NF-H_2O, 1 μL 10 mM dNTPs and 1

μL 2 μM ssRT primer (Table 1) or 50 μM Oligo(dT)$_{20}$. Mixes were cooled to room temperature with

agitation at 600 rpm for 10 minutes before addition of 8 μL 5x SuperScript IV reaction buffer, 2 μL 100

mM DTT and 1 μL SuperScript IV enzyme to each reaction. Samples were incubated at $50^{\circ}C$ for 1 hour

with agitation at 600 rpm. Reaction mixtures were heated at $85^{\circ}C$ for 10 minutes to inactivate reverse

transcription and release cDNA from beads. Tubes were pulse centrifuged to pellet liquid prior to bead

immobilisation using a magnetic rack and aspiration of cDNA. cDNA was diluted 100-fold for positive-

strand reactions and 10-fold for negative-strand reactions. Oligo(dT)$_{20}$ reactions were diluted

accordingly. qPCR reactions were performed as previously described above with quantification using

the ΔΔCq method. Samples in which no signal was detected were arbitrarily assigned a C_T value of 40

to permit quantification.

Results

Transfection with replicon RNA is not compatible with strand-specific discrimination by qPCR.

Reverse genetics is a powerful tool for investigating the molecular biology of many viruses. However,

a common drawback with many reverse genetics systems is the inefficient delivery of *in vitro*

transcribed viral genomes, which can overwhelm assays deigned to dissect individual steps of viral

lifecycles such as strand-specific genome synthesis. To overcome this restriction we have developed

assays involving nascent RNA labelling, reverse transcription and strand-specific qPCR to examine the

individual steps of negative- and/or positive-strand synthesis.

Our previously reported assays of replication utilised a FMDV replicon encoding the GFP reporter

protein in place of the viral structural proteins (60, 61). Transfection of *in vitro* transcribed replicon

RNA into cells allows measurement of a completed replication cycle via GFP expression prior to RNA

extraction, cDNA synthesis and measurement of strand-specific specific RNA replication by qPCR

(Figure 1A). Following transfection with *in vitro* transcribed pRep-ptGFP replicon RNA (WT or the

replication-deficient 3D^{GNN}) or yeast tRNA (as a control), cells were examined using an Incucyte S3

instrument to visualise phase contrast and green images (representative images Figure 1B) and the

number of ptGFP positive cells determined using pre-defined analysis parameters (Figure 1C).

Strand-specific primers with a non-viral tag sequence were designed for use during reverse

transcription reactions to generate unique 5' sequences providing specificity for each strand (Table 1)

similar to published protocols (55-58). In order to minimise the potential for cross-contamination,

primers were designed towards the 5' and 3' ends of the FMDV replicon genome for the positive-

strand and negative-strand, respectively (Figure 1D). Reverse transcription using oligo(dT)$_{20}$ was also

performed for each sample to permit normalisation between samples by amplifying β-actin as a stably

expressed reference gene. A primer complementary to the strand-specific non-viral tag and an

internal primer complementary to the viral genome were designed for the strand-specific qPCR. Initial

assays were performed comparing strand production between WT and 3D^{GNN} replicons. As

anticipated, only a ≈2 fold-difference was observed between WT and 3D^{GNN} for both strands (Figure

1E&F), suggesting that the ability to demonstrate nascent viral RNA production was masked by the

large quantity of input RNA necessary to efficiently initiate replication.

Removal of input replicon genomes by nascent RNA labelling.

We speculated that the inability of the strand-specific qPCR to distinguish between RNA isolated from

cells transfected with either replication-competent or -incompetent replicon constructs was due to

the large amount of input RNA necessary to initiate replication in the majority of the cells. To address

this, we developed a method to specifically label and isolate newly synthesised RNA. To facilitate

selection, transfected cells were incubated with an alkyne-modified uridine, 5-EU, from the time of

transfection so that only cellular RNAs and viral genomes synthesised during replication incorporated

the modified nucleotide (Figure 2A). Control assays indicated that addition of 0.2 mM 5-EU to the

culture medium did not inhibit replication (Figure 2B).

Following RNA extraction, strand-specific qPCR was performed to determine whether incorporation

of 5-EU had any adverse effects upon the assay. These assays showed that detection of positive-strand

and negative-strand genomes (Figure 2C&D, respectively) was comparable to that observed without

labelling (Figure 1E&F). Copper-catalysed click reactions were performed to covalently link azide-

modified biotin to the newly synthesised RNA. This acted as a bait for capture with streptavidin

magnetic beads and removal by washing of input replicon RNA. Following capture and on-bead reverse

transcription, analysis of the eluted cDNA confirmed that the method had efficiently removed replicon

input RNA, with the 3D^{GNN} replicon signal reduced to 0.03 and 0.02 of WT for positive-strand and

negative-strand, respectively (Figure 2E&F). This confirmed that labelling of newly synthesised RNA

with 5-EU prior to biotin modification by click reaction and bead capture provided a method suitable

for replicon-based investigation of differential viral strand synthesis.

Cre **is essential for negative-strand production.**

The *cre* is essential to replication of picornaviruses by acting as a template for uridylylation of VPg by

$3D^{pol}$. There is also evidence that the poly(A) tail of poliovirus may act as a template for uridylylation

in cell-free replication, albeit-it less efficiently (52-54). However, whether c*re* is essential for VPg

uridylylation to prime the negative-strand intermediate during replication in cells is unknown. The role

of the *cre* in strand synthesis was therefore investigated using replicons harbouring a complete *cre*

deletion (Δ*cre*) or with a single nucleotide substitution of the functional AAACA motif (termed A1G,

containing mutation of the first A nucleotide to G). Both of these mutations are known to prevent the

complete virus lifecycle (25). Loss of replication competency was confirmed here for both mutants by

replicon assay (Figure 3A). Strand-specific assays confirmed that the positive-strand could not be

synthesised by either mutant replicon (Figure 3B). Negative-strand production was also ablated,

showing that the c*re* is also essential during intracellular replication for production of the negative-

strand (Figure 3C).

Discussion

The ability to reliably differentiate positive-strand and negative-strand RNAs is important for

investigating details of the mechanism of picornavirus replication. This can be achieved using qPCR

techniques, but the selection of strand-specific primers can be challenging. This problem is

exacerbated when studying replicons, as reliable transfection requires a large quantity of input RNA.

Despite these drawbacks, replicons facilitate investigations of replication that would otherwise be

technically challenging, e.g. with non-recoverable or poorly replicating viruses, or to address issues

regarding containment and safety. We therefore designed a strand-specific qPCR assay to distinguish

synthesis of each strand separately and explored how mutation to the viral genome may restrict

synthesis.

Initial assays identified the large quantity of input RNA required to ensure transfection of the majority

of cells as a major restriction to demonstrating differences between replication-competent and -

defective replicons (Figure 1). In the case of the FMDV replicon, the ≈ 7 kb genome relates to

approximately 1.8×10^{12} genome copies/μg, and with up to 5 μg/well of replicon being transfected

into cells, carry-over of input RNA was unsurprising. RNA self-priming, a consequence of high

secondary structure at the 3' terminus of RNA and reverse transcriptases lacking RNase H activity, has

been reported previously as a source of contamination by complementary sequences in strand-

specific assays (57, 58). In addition, residual fragments of plasmid DNA used as template for *in vitro* transcription of the transfecting RNA could theoretically also be a source of contaminating complementary sequences. However, contamination by any interfering sequences is eliminated by the specific selection of newly synthesised RNA as described here, thus providing an assay that can reliably distinguish complementary strands.

The essential role of the *cre* in picornavirus replication is well documented and was also confirmed here in replicon assays (Figure 3A) (36, 52-54, 62). It is possible that the poly(A) tail could template uridylylation of VPg (i.e. in addition to and independently of the c*re*, as reported in cell-free assays) (52-54). However, it was not established whether the c*re* is required for negative-strand synthesis in an intracellular assay. We therefore investigated the role of two c*re* mutations (A1G and Δ*cre*) during strand synthesis and found that neither positive- nor negative-strand RNAs were transcribed. Given that negative-strand intermediates were not produced, it is unsurprising that positive-strand production was also restricted. Due to a lack of negative-strand RNA it could not be established in this assay whether synthesis of new positive-strand genomes requires the *cre*. However, given evidence from the literature of the importance to positive-strand RNA synthesis, these findings would suggest that the c*re* is essential for efficient priming of replication by 3D^{pol} in the infected cell.

In conclusion, we present a method using 5-EU labelling and purification of nascent RNA prior to strand-specific qPCR that can be applied to determine the role of viral elements in transcription of positive- and negative-strands using replicons containing specific mutations. Here, this was applied to the known c*re* mutations A1G and Δc*re* and showed that neither are capable of producing a negative-strand intermediate. We will continue to probe other mutants in order to discover the role of viral elements that control viral replication of positive- and negative-strands.

Funding information

This project was funded by the Biotechnology and Biological Sciences Research council UK (project grant reference BB/T015748/1).

Acknowledgements

We are grateful to group members for their helpful comments on the project.

Author contributions

282 The project was conceived by SJD, MRH, DJR and NJS. The experiments were carried out and the data

283 analysed by SJD, JCW undertook mutagenesis. The manuscript was drafted by SJD and was edited and

284 revised by all authors.

Conflicts of interest

288 The authors declare that there are no conflicts of interest.

References

1.   Paton DJ, Di Nardo A, Knowles NJ, Wadsworth J, Pituco EM, Cosivi O, et al. The history of foot-
and-mouth disease virus serotype C: the first known extinct serotype? Virus Evol. 2021;7(1):veab009.

2.   Jamal SM, Belsham GJ. Foot-and-mouth disease: past, present and future. Vet Res.
2013;44:116.

3.   Knight-Jones TJ, Rushton J. The economic impacts of foot and mouth disease - what are they,
how big are they and where do they occur? Prev Vet Med. 2013;112(3-4):161-73.

4.   Feng S, Patton M, Davis J. Market Impact of Foot-and-Mouth Disease Control Strategies: A UK
Case Study. Front Vet Sci. 2017;4:129.

5.   Thompson D, Muriel P, Russell D, Osborne P, Bromley A, Rowland M, et al. Economic costs of
the foot and mouth disease outbreak in the United Kingdom in 2001. Rev Sci Tech. 2002;21(3):675-
87.

6.   Mahapatra M, Parida S. Foot and mouth disease vaccine strain selection: current approaches
and future perspectives. Expert Rev Vaccines. 2018;17(7):577-91.

7.   Park JH. Requirements for improved vaccines against foot-and-mouth disease epidemics. Clin
Exp Vaccine Res. 2013;2(1):8-18.

8.   Stenfeldt C, Eschbaumer M, Rekant SI, Pacheco JM, Smoliga GR, Hartwig EJ, et al. The Foot-
and-Mouth Disease Carrier State Divergence in Cattle. J Virol. 2016;90(14):6344-64.

9.   Dawe PS, Sorensen K, Ferris NP, Barnett IT, Armstrong RM, Knowles NJ. Experimental
transmission of foot-and-mouth disease virus from carrier African buffalo (Syncerus caffer) to cattle
in Zimbabwe. Vet Rec. 1994;134(9):211-5.

10.   Dawe PS, Flanagan FO, Madekurozwa RL, Sorensen KJ, Anderson EC, Foggin CM, et al. Natural
transmission of foot-and-mouth disease virus from African buffalo (Syncerus caffer) to cattle in a
wildlife area of Zimbabwe. Vet Rec. 1994;134(10):230-2.

11.   Gao Y, Sun SQ, Guo HC. Biological function of Foot-and-mouth disease virus non-structural
proteins and non-coding elements. Virol J. 2016;13:107.

12.   Roberts PJ, Belsham GJ. Identification of critical amino acids within the foot-and-mouth
disease virus leader protein, a cysteine protease. Virology. 1995;213(1):140-6.

13.   Vakharia VN, Devaney MA, Moore DM, Dunn JJ, Grubman MJ. Proteolytic processing of foot-
and-mouth disease virus polyproteins expressed in a cell-free system from clone-derived transcripts.
J Virol. 1987;61(10):3199-207.

14.   Moffat K, Knox C, Howell G, Clark SJ, Yang H, Belsham GJ, et al. Inhibition of the secretory
pathway by foot-and-mouth disease virus 2BC protein is reproduced by coexpression of 2B with 2C,
and the site of inhibition is determined by the subcellular location of 2C. J Virol. 2007;81(3):1129-39.

15.   Yost SA, Marcotrigiano J. Viral precursor polyproteins: keys of regulation from replication to
maturation. Curr Opin Virol. 2013;3(2):137-42.

16.   Ferrer-Orta C, Arias A, Agudo R, Perez-Luque R, Escarmis C, Domingo E, et al. The structure of
a protein primer-polymerase complex in the initiation of genome replication. EMBO J. 2006;25(4):880-
8.

17.   Birtley JR, Knox SR, Jaulent AM, Brick P, Leatherbarrow RJ, Curry S. Crystal structure of foot-
and-mouth disease virus 3C protease. New insights into catalytic mechanism and cleavage specificity.
J Biol Chem. 2005;280(12):11520-7.

18.   Grubman MJ, Zellner M, Bablanian G, Mason PW, Piccone ME. Identification of the active-site
residues of the 3C proteinase of foot-and-mouth disease virus. Virology. 1995;213(2):581-9.

19.   Forss S, Schaller H. A tandem repeat gene in a picornavirus. Nucleic Acids Res.
1982;10(20):6441-50.

20.   King AM, Sangar DV, Harris TJ, Brown F. Heterogeneity of the genome-linked protein of foot-
and-mouth disease virus. J Virol. 1980;34(3):627-34.

21.   Gonzalez-Magaldi M, Martin-Acebes MA, Kremer L, Sobrino F. Membrane topology and
cellular dynamics of foot-and-mouth disease virus 3A protein. PLoS One. 2014;9(9):e106685.

22.   Ferrer-Orta C, Arias A, Perez-Luque R, Escarmis C, Domingo E, Verdaguer N. Structure of foot-
and-mouth disease virus RNA-dependent RNA polymerase and its complex with a template-primer
RNA. J Biol Chem. 2004;279(45):47212-21.

23.   Herod MR, Gold S, Lasecka-Dykes L, Wright C, Ward JC, McLean TC, et al. Genetic economy in
picornaviruses: Foot-and-mouth disease virus replication exploits alternative precursor cleavage
pathways. PLoS Pathog. 2017;13(10):e1006666.

24.   Ward JC, Lasecka-Dykes L, Neil C, Adeyemi OO, Gold S, McLean-Pell N, et al. The RNA
pseudoknots in foot-and-mouth disease virus are dispensable for genome replication, but essential
for the production of infectious virus. PLoS Pathog. 2022;18(6):e1010589.

25.   Mason PW, Bezborodova SV, Henry TM. Identification and characterization of a cis-acting
replication element (cre) adjacent to the internal ribosome entry site of foot-and-mouth disease virus.
J Virol. 2002;76(19):9686-94.

26.   Kanda T, Ozawa M, Tsukiyama-Kohara K. IRES-mediated translation of foot-and-mouth
disease virus (FMDV) in cultured cells derived from FMDV-susceptible and -insusceptible animals. BMC
Vet Res. 2016;12:66.

27.   Kloc A, Diaz-San Segundo F, Schafer EA, Rai DK, Kenney M, de Los Santos T, et al. Foot-and-
mouth disease virus 5'-terminal S fragment is required for replication and modulation of the innate
immune response in host cells. Virology. 2017;512:132-43.

28.   Carrillo C, Tulman ER, Delhon G, Lu Z, Carreno A, Vagnozzi A, et al. Comparative genomics of
foot-and-mouth disease virus. J Virol. 2005;79(10):6487-504.

29.   Lopez de Quinto S, Martinez-Salas E. Conserved structural motifs located in distal loops of
aphthovirus internal ribosome entry site domain 3 are required for internal initiation of translation. J
Virol. 1997;71(5):4171-5.

30. Lopez de Quinto S, Saiz M, de la Morena D, Sobrino F, Martinez-Salas E. IRES-driven translation
is stimulated separately by the FMDV 3'-NCR and poly(A) sequences. Nucleic Acids Res.
2002;30(20):4398-405.

31. Xiang W, Harris KS, Alexander L, Wimmer E. Interaction between the 5'-terminal cloverleaf
and 3AB/3CDpro of poliovirus is essential for RNA replication. J Virol. 1995;69(6):3658-67.

32. Gamarnik AV, Andino R. Interactions of viral protein 3CD and poly(rC) binding protein with the
5' untranslated region of the poliovirus genome. J Virol. 2000;74(5):2219-26.

33. Harris KS, Xiang W, Alexander L, Lane WS, Paul AV, Wimmer E. Interaction of poliovirus
polypeptide 3CDpro with the 5' and 3' termini of the poliovirus genome. Identification of viral and
cellular cofactors needed for efficient binding. J Biol Chem. 1994;269(43):27004-14.

34. Yang Y, Rijnbrand R, Watowich S, Lemon SM. Genetic evidence for an interaction between a
picornaviral cis-acting RNA replication element and 3CD protein. J Biol Chem. 2004;279(13):12659-67.

35. Yang F, Zhu Z, Cao W, Liu H, Wei T, Zheng M, et al. Genetic Determinants of Altered Virulence
of Type O Foot-and-Mouth Disease Virus. J Virol. 2020;94(7).

36. Kloc A, Rai DK, Rieder E. The Roles of Picornavirus Untranslated Regions in Infection and Innate
Immunity. Front Microbiol. 2018;9:485.

37. Zhu Z, Yang F, Cao W, Liu H, Zhang K, Tian H, et al. The Pseudoknot Region of the 5'
Untranslated Region Is a Determinant of Viral Tropism and Virulence of Foot-and-Mouth Disease Virus.
J Virol. 2019;93(8).

38. Mohapatra JK, Pawar SS, Tosh C, Subramaniam S, Palsamy R, Sanyal A, et al. Genetic
characterization of vaccine and field strains of serotype A foot-and-mouth disease virus from India.
Acta Virol. 2011;55(4):349-52.

39. Escarmis C, Dopazo J, Davila M, Palma EL, Domingo E. Large deletions in the 5'-untranslated
region of foot-and-mouth disease virus of serotype C. Virus Res. 1995;35(2):155-67.

40. Paul AV, Rieder E, Kim DW, van Boom JH, Wimmer E. Identification of an RNA hairpin in
poliovirus RNA that serves as the primary template in the in vitro uridylylation of VPg. J Virol.
2000;74(22):10359-70.

41. Nayak A, Goodfellow IG, Belsham GJ. Factors required for the Uridylylation of the foot-and-
mouth disease virus 3B1, 3B2, and 3B3 peptides by the RNA-dependent RNA polymerase (3Dpol) in
vitro. J Virol. 2005;79(12):7698-706.

42. Schein CH, Ye M, Paul AV, Oberste MS, Chapman N, van der Heden van Noort GJ, et al.
Sequence specificity for uridylylation of the viral peptide linked to the genome (VPg) of enteroviruses.
Virology. 2015;484:80-5.

43. Paul AV, Yin J, Mugavero J, Rieder E, Liu Y, Wimmer E. A "slide-back" mechanism for the
initiation of protein-primed RNA synthesis by the RNA polymerase of poliovirus. J Biol Chem.
2003;278(45):43951-60.

44. Paul AV, Peters J, Mugavero J, Yin J, van Boom JH, Wimmer E. Biochemical and genetic studies
of the VPg uridylylation reaction catalyzed by the RNA polymerase of poliovirus. J Virol.
2003;77(2):891-904.

45. Paul AV, van Boom JH, Filippov D, Wimmer E. Protein-primed RNA synthesis by purified
poliovirus RNA polymerase. Nature. 1998;393(6682):280-4.

46. Belsham GJ, Brangwyn JK. A region of the 5' noncoding region of foot-and-mouth disease virus
RNA directs efficient internal initiation of protein synthesis within cells: involvement with the role of
L protease in translational control. J Virol. 1990;64(11):5389-95.

47. Bienz K, Egger D, Pfister T, Troxler M. Structural and functional characterization of the
poliovirus replication complex. J Virol. 1992;66(5):2740-7.

48. Lundquist RE, Maizel JV, Jr. In vivo regulation of the poliovirus RNA polymerase. Virology.
1978;89(2):484-93.

49. Barton DJ, Flanegan JB. Synchronous replication of poliovirus RNA: initiation of negative-
strand RNA synthesis requires the guanidine-inhibited activity of protein 2C. J Virol. 1997;71(11):8482-
9.

50. Richards OC, Martin SC, Jense HG, Ehrenfeld E. Structure of poliovirus replicative intermediate
RNA. Electron microscope analysis of RNA cross-linked in vivo with psoralen derivative. J Mol Biol.
1984;173(3):325-40.

51. Triantafilou K, Vakakis E, Kar S, Richer E, Evans GL, Triantafilou M. Visualisation of direct
interaction of MDA5 and the dsRNA replicative intermediate form of positive strand RNA viruses. J Cell
Sci. 2012;125(Pt 20):4761-9.

52. Goodfellow IG, Polacek C, Andino R, Evans DJ. The poliovirus 2C cis-acting replication element-
mediated uridylylation of VPg is not required for synthesis of negative-sense genomes. J Gen Virol.
2003;84(Pt 9):2359-63.

53. Morasco BJ, Sharma N, Parilla J, Flanegan JB. Poliovirus cre(2C)-dependent synthesis of
VPgpUpU is required for positive- but not negative-strand RNA synthesis. J Virol. 2003;77(9):5136-44.

54. Murray KE, Barton DJ. Poliovirus CRE-dependent VPg uridylylation is required for positive-
strand RNA synthesis but not for negative-strand RNA synthesis. J Virol. 2003;77(8):4739-50.

55. Horsington J, Zhang Z. Analysis of foot-and-mouth disease virus replication using strand-
specific quantitative RT-PCR. J Virol Methods. 2007;144(1-2):149-55.

56. Plaskon NE, Adelman ZN, Myles KM. Accurate strand-specific quantification of viral RNA. PLoS
One. 2009;4(10):e7468.

57. Tuiskunen A, Leparc-Goffart I, Boubis L, Monteil V, Klingstrom J, Tolou HJ, et al. Self-priming
of reverse transcriptase impairs strand-specific detection of dengue virus RNA. J Gen Virol. 2010;91(Pt
4):1019-27.

58. Vashist S, Urena L, Goodfellow I. Development of a strand specific real-time RT-qPCR assay for
the detection and quantitation of murine norovirus RNA. J Virol Methods. 2012;184(1-2):69-76.

59. Muller M, Slivinski N, Todd E, Khalid H, Li R, Karwatka M, et al. Chikungunya virus requires
cellular chloride channels for efficient genome replication. PLoS Negl Trop Dis. 2019;13(9):e0007703.

60. Tulloch F, Pathania U, Luke GA, Nicholson J, Stonehouse NJ, Rowlands DJ, et al. FMDV replicons
encoding green fluorescent protein are replication competent. J Virol Methods. 2014;209:35-40.

61. Herod MR, Loundras EA, Ward JC, Tulloch F, Rowlands DJ, Stonehouse NJ. Employing
transposon mutagenesis to investigate foot-and-mouth disease virus replication. J Gen Virol.
2015;96(12):3507-18.

62. Steil BP, Barton DJ. Cis-active RNA elements (CREs) and picornavirus RNA replication. Virus
Res. 2009;139(2):240-52.

Figure legends

Figure 1: Transfection of FMDV replicon RNA is not compatible with distinguishing the synthesis of positive and negative strand molecules during replication. (A) Schematic overview of a replicon transfection assay with imaging prior to RNA extraction, cDNA synthesis and strand-specific qPCR. (B) Incucyte S3 imaging of BHK-21 cells transfected with WT and replication deficient 3D[GNN] (GNN) FMDV pRep-ptGFP RNA and yeast tRNA (as a control), with phase contrast and green visualisation over time. Scale bar = 400 µm. (C) Detection over time of ptGFP following transfection with FMDV replicon RNA. (D) Schematic representation of primer binding regions with P2 and P3 genomic regions of FMDV for the cDNA synthesis and qPCR detection of positive- and negative-strands. RNA extracted from transfected cells was reverse transcribed using strand-specific primers prior to qPCR to detect positive-strand (E) and negative-strand (F) expression. Analysis was performed using the ΔΔCq method relative to WT. Statistical analysis was performed by one-way ANOVA with ** *P*-value = *<0.005* and **** *P*-value = *<0.0001*. n = 3.

Figure 2: Removal of input replicon RNA by 5-EU labelling and RNA pulldown to select newly synthesised RNA. (A) Schematic representation of a 5-EU nascent RNA labelling assay. BHK cells were transfected with FMDV pRep-ptGFP RNA (WT and GNN), with addition of 0.2 mM 5-EU to label nascent RNA. An unlabelled control was also included (termed WT no EU). Cells were imaged using an Incucyte S3 instrument prior to harvesting of RNA 6 hours post transfection. 5-EU labelled RNA was biotinylated by click reaction and captured with streptavidin magnetic beads. Input RNA was removed by washing prior to on-bead cDNA synthesis and strand-specific qPCR. (B) ptGFP positive cells detected by Incucyte S3 imaging 6 hours post transfection to determine effect of 5-EU labelling on replication. Following RNA extraction, strand-specific qPCR to detect positive-strands (C) and negative-strands (D) prior to click reaction, was performed. Click reaction mediated biotinylation of nascent RNA, magnetic bead capture and on bead cDNA synthesis was performed prior to strand-specific assays to determine positive-strand (E) and negative-strand (F) expression. Analysis was performed using the ΔΔCq method relative to WT. Statistical analysis was performed by one-way ANOVA with ** *P*-value = *<0.005* and **** *P*-value = *<0.0001*. n = 3.

Figure 3: FMDV *cre* is essential to the production of negative-strand genomes. (A) BHK-21 cells were transfected with pRep-ptGFP constructs containing *cre* mutations A1G and Δ*cre* (alongside WT and GNN controls as described in Figure 1) with addition of 0.2 mM 5-EU to label nascent transcribed RNA. ptGFP positive cells were visualised 6 hours post-transfection using an Incucyte S3 instrument. Total RNA extracted 6 hours post transfection was biotinylated by click reaction, captured with streptavidin beads and reverse transcribed using strand-specific or oligo dT$_{(20)}$ as a primer. cDNA was used in

478   strand-specific qPCR assays to determine positive-strand (B) and negative-strand (C) expression.

479   Analysis was performed using the ΔΔCq method relative to WT. Statistical analysis was performed by

480   one-way ANOVA with **** *P*-value = *<0.0001*. n = 3.

481 ## Tables and figures

Table 1: Primers used in RT and qPCR reactions. Underlined are the non-viral tag sequences for both the RT and qPCR primers.

Target	Application	Name	Sequence (5'-3')
Positive-strand	RT	PS-tag-RT	<u>GCAGGAGCTAAGCGCTGG</u>GGTCGCGATGATGACTTTGC
	qPCR	PS-qPCR-TAG	<u>GCAGGAGCTAAGCGCTGG</u>
	qPCR	PS-qPCR-R	GATTTGGGCCAGAACCCTGA
Negative-strand	RT	NS-tag-RT	<u>TGACGTTGCGACGAGTTCG</u>CCAAAGACGGAGCTGACACT
	qPCR	NS-qPCR-TAG	<u>TGACGTTGCGACGAGTTCG</u>
	qPCR	NS-qPCR-R	GTCCACAATCAACCCCTCGT
β-actin	qPCR	β-actin-F	GTGCTATGTTGCCCTGGACT
	qPCR	β-actin-R	GCTCGTTGCCAATGGTGATG

Figure 1

Figure 2

Figure 3