Whole transcriptome profiling of placental pathobiology in SARS-CoV-2 pregnancies identifies placental dysfunction signatures.

Nataly Stylianou[1#], Ismail Sebina[2#], Nicholas Matigian[3], James Monkman[2], Hadeel Doehler[1], Joan Röhl[4], Mark Allenby[5], Andy Nam[6], Liuliu Pan[6], Anja Rockstroh[1], Habib Sadeghirad[2], Kimberly Chung[2], Thais Sobanski[1], Ken O'Byrne[1], Ana Clara Simoes Florido Almeida[7], Patricia Zadorosnei Rebutini[7], Cleber Machado-Souza[8], Emanuele Therezinha Schueda Stonoga[9], Majid E Warkiani[10], Carlos Salomon[11], Kirsty Short[12], Lana McClements[10], Lucia de Noronha[7], Ruby Huang[13], Gabrielle T. Belz[2], Fernando Souza-Fonseca-Guimaraes[2], Vicki Clifton[14], Arutha Kulasinghe[2*].

[1] Centre for Genomics and Personalised Health, School of Biomedical Sciences, Queensland University of Technology, Brisbane, QLD 4102, Australia

[2] Frazer Institute, Faculty of Medicine, The University of Queensland, Brisbane, QLD 4102, Australia.

[3] QCIF Bioinformatics, St Lucia, QLD 4102, Australia

[45] Faculty of Health Sciences and Medicine, Bond University, Queensland, Australia

[5] BioMimetic Systems Engineering Lab, School of Chemical Engineering, University of Queensland (UQ), St Lucia, QLD, Australia

[6] Nanostring Technologies, Inc, Seattle, WA, USA

[7] Postgraduate Program of Health Sciences, School of Medicine, Pontifícia Universidade Católica do Paraná´ -PUCPR, Curitiba, Brazil.

[8] Postgraduate Program in Biotechnology Applied in Health of Children and Adolescent, Instituto de Pesquisa Pelé Pequeno Príncipe, Faculdades Pequeno Príncipe, Curitiba, Brazil

[9] Department of Medical Pathology, Clinical Hospital, Universidade Federal do Paraná´ - UFPR, Curitiba, Brazil.

[10] *School of Life Sciences & Institute for Biomedical Materials and Devices, Faculty of*

Science, University of Technology Sydney, NSW, Australia.

[11] *Exosome Biology Laboratory, Centre for Clinical Diagnostics, University of Queensland*

Centre for Clinical Research, Royal Brisbane and Women's Hospital, Faculty of Medicine,

The University of Queensland, Brisbane, Australia.

[12] *School of Chemistry and Molecular Biosciences, Faculty of Science, The University of*

Queensland, St Lucia QLD 4067, Australia.

[13] *School of Medicine, College of Medicine, National Taiwan University, Taipei, Taiwan.*

[14] *Mater Medical Research Institute – University of Queensland, Brisbane, Australia.*

 # These authors contributed equally

*Correspondence: Dr Arutha Kulasinghe, Frazer Institute, Faculty of Medicine, The

University of Queensland. 37 Kent Street, Woolloongabba, Queensland 4012, Australia. E:

arutha.kulasinghe@uq.edu.au

Graphical abstract: In this study, using spatial digital profiling transcriptomic approaches,

we demonstrate that SARS-CoV-2 infection in pregnancy disrupts optimal placental function

by altering the genomic architecture of trophoblasts and villous core stromal cells.

ABSTRACT

44 **Objectives:** Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) virus

45 infection in pregnancy is associated with higher incidence of placental dysfunction, referred

46 to by a few studies as a "preeclampsia-like syndrome". However, the mechanisms

47 underpinning SARS-CoV-2-induced placental malfunction are still unclear. Here, we

investigated whether the transcriptional architecture of the placenta is altered in response to SARS-CoV-2 infection.

Methods: We utilized whole-transcriptome, digital spatial profiling, to examine gene expression patterns in placental tissues from participants who contracted SARS-CoV-2 in the third trimester of their pregnancy (n=7) and those collected prior to the start of the coronavirus disease 2019 (COVID-19) pandemic (n=9).

Results: Through comprehensive spatial transcriptomic analyses of the trophoblast and villous core stromal cell subpopulations in the placenta, we identified signatures associated with hypoxia and placental dysfunction during SARS-CoV-2 infection in pregnancy. Notably, genes associated with vasodilation (*NOS3*), oxidative stress (*GDF15*, *CRH*), and preeclampsia (*FLT1, EGFR, KISS1, PAPPA2),* were enriched with SARS-CoV-2. Pathways related to increased nutrient uptake, vascular tension, hypertension, and inflammation, were also enriched in SARS-CoV-2 samples compared to uninfected controls.

Conclusions: Our findings demonstrate the utility of spatially resolved transcriptomic analysis in defining the underlying pathogenic mechanisms of SARS-CoV-2 in pregnancy, particularly its role in placental dysfunction. Furthermore, this study highlights the significance of digital spatial profiling in mapping the intricate crosstalk between trophoblasts and villous core stromal cells, thus shedding light on pathways associated with placental dysfunction in pregnancies with SARS-CoV-2 infection.

KEYWORDS: Placental dysfunction, SARS-CoV-2, COVID-19, digital spatial profiling, gene expression profiling, trophoblasts, villous core stroma.

INTRODUCTION

Viral infections in pregnancy can disrupt placental function and predispose pregnancy complications, including late-onset preeclampsia, preterm birth, stillbirth, and intrauterine fetal demise[1-4]. Recent studies have revealed that pregnant women who contract Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2, which causes coronavirus disease 2019 [COVID-19]), can experience placental dysfunction and what has been referred to as "preeclampsia-like syndrome" [5-10]. Placental tissues from COVID-19 patients exhibit increased vasculopathy and inflammation, which are characteristic pathological features of preeclampsia [11]. Moreover, clinical manifestations observed in COVID-19 patients, such as COVID-19-associated hypoxia, hypertension, endothelial dysfunction, kidney disease, thrombocytopenia, and liver injury, overlap with those observed in preeclampsia[5, 12]. However, mechanisms through which SARS-CoV-2 infection predisposes pregnancies to these preeclampsia-like pathological features are largely unclear.

The placenta is vital for fetal development and growth throughout gestation as it is a functional interface between the mother and fetus[13]. This interface comprises various anatomically distinct sites, including the decidua basalis, where maternal immune cells and decidual stromal cells interact with fetal extravillous trophoblasts[14]. The maternal-fetal interface also consists of the placental intervillous space, where maternal immune cells interact with fetal syncytiotrophoblasts, and the boundary between the parietalis and the chorion laeve in the chorioamniotic membranes[14]. Other cell types within this interface, such as villous cytotrophoblasts, column cytotrophoblasts, fibroblasts, endothelial cells, and Hofbauer cells, contribute to nutrient and waste exchange, hormone production, protection from pathogens, and maternal immune responses essential for fetal development [13, 14].

Whether SARS-CoV-2 infection modifies the transcriptomic architecture and functional characteristics of different cell types within these distinct placental sites is still unclear.

In this study, we utilised digital spatial whole-transcriptomic analysis of human placental tissue to delineate molecular pathways associated with SARS-CoV-2 infection-induced placental pathology in pregnancy. Specifically, we focused our analysis on defining the distinct transcriptional profiles of trophoblasts and villous core stromal cell populations (the latter including endothelial, fibroblast, and immune cells), in the context of SARS-CoV-2 infection. We identified several pivotal pathways that underlie the development of a "preeclampsia-like syndrome" associated with SARS-CoV-2 infection in pregnancy.

RESULTS

Characterization of patient demographics and histopathology in collected placentae.

Tissue microarrays were constructed using placental cores that were collected immediately after birth from unvaccinated participants who had tested positive within 15 days prior to delivery (Alpha strain, April 2020, n=7), and unvaccinated participants who were negative for SARS-CoV-2 throughout their pregnancy (n=9; Table 1). There were no significant differences in placental weight, fetal weight, gestational age, comorbidities, or maternal age between the two groups (Table 1). Within the SARS-CoV-2 group, 3/7 newborns were born preterm, compared to 4/9 in the control group (Table 1).

No SARS-CoV-2 viral load was detected in the placental cores from the SARS-CoV-2 infected group through examination by RNAscope of the SARS-CoV-2 spike mRNA (data not shown). With the aid of a trained placental pathologist, an area featuring an anchoring

villus, and an area featuring a cluster of terminal villi, were designated as two areas of interest (AOI) within each placental core (Figure 1a). AOIs where immunofluorescently stained with PanCK to identify trophoblast populations, and vimentin to identify stromal populations (i.e., fibroblasts, endothelial cells). Transcriptional expression was collected separately for PanCK positive and separately for vimentin positive cells within each AOI using the Whole Transcriptome Atlas kit (Nanostring; Figure 1a). Subsequent cell deconvolution was performed to assess the purity of each collection (Figure 1b). As expected, transcriptional expression from PanCK positive regions within the AOIs had high proportion of trophoblast populations, compared to vimentin positive regions that had higher proportions of fibroblast and endothelial cells (Figure 1b-f). Due to the overlapping nature of cells, all samples captured a proportion of immune cell types (macrophages, monocytes, Hofbauer cells), except for the PanCK positive regions that displaying a proportion of granulocytes that was absent from the vimentin positive regions within the AOIs (Figure 1b, g-j). SARS-CoV-2 infection did not significantly alter the transcriptional proportion of any cell type assessed when compared to controls (Figure 1b-j). In subsequent analyses, the PanCK positive regions within the AOIs will be referred to as "Trophoblasts" and the vimentin positive regions will be referred to as "Villous Core Stroma" compartments, due to the predominant enriched cell type they represent.

SARS-CoV-2 infection related pathways enriched in the placenta despite absence of detectable viral particles.

Unsupervised clustering of the normalised gene counts by principal component analysis showed that SARS-CoV-2 samples clustered separately to control samples in dimension 1, and further by phenotype in dimension 2, supporting that infection with SARS-CoV-2 markedly alters the transcriptional profiles of the trophoblast and villous core stroma cell populations (Figure 2a-b). Notably, there was very high overlap of genes differentially

expressed between the anchoring and the terminal villi; for instance, trophoblasts at the anchoring villi had 1,791 differentially expressed genes versus 493 at the terminal villi, with 405 genes in common between them (82% overlap; Supplementary table 1, Figure 2c). Similarly, the villous core stroma cells at the anchoring villi had 1,139 differentially expressed genes versus 601 at the terminal villi, with 458 genes in common between them (76% overlap; Supplementary table 1, Figure 2c). As expected, there was minimal overlap in differential gene expression between the trophoblasts and villous core stroma compartments, which highlights their distinct cell phenotypes (Figure 2c).

Despite the SARS-CoV-2 samples showing undetectable SARS-CoV-2 by RNA-scope, transcriptional profiling showed positive enrichment of SARS-CoV-2 related pathways in the SARS-CoV-2 samples, such as "SARS_COV_2_INFECTION", and "SARS_COV_2_HOST_INTERACTIONS" from the Reactome database, as well as the Interferon Alpha Response pathway from the Hallmark database, which is a first-line immune response pathway that has been associated with SARS-CoV-2 infection (Figure 2d-f)[15] . These pathways were supported by increased expression of genes that have been associated with SARS-CoV-2, such as the inflammatory marker *IFI16*[16], disease progression marker *IFI27*[17], disease prognosticator *B2M*[18], activation of Janus Kinases (i.e., *JAK1*), and expression of *STAT3*[19] (Figure 2 g-j). Notably, gene expression for these markers was elevated predominantly in the villous core stroma cell compartment, presumably stemming from the immune subpopulation within the stroma. Indeed, specific analysis of the villus core stroma compartment revealed enrichment of several immune related pathways from the Hallmarks database such as IL6/JAK/STAT3 signalling, IL2/STAT5 signalling, TNF-alpha signalling, inflammatory response, and complement pathways (Figure 3a), supporting that the immune cells within the placental villi are actively responding to SARS-CoV-2 infection.

SARS-CoV-2 infection enriches hypoxia and placental dysfunction pathways.

Pathway enrichment analysis of the genes differentially expressed in response to SARS-CoV-2 infection, revealed pathways related to placental disfunction, in both the trophoblast and villous core stroma compartments. For instance, hypoxia and oxidative phosphorylation pathways were enriched in the villous core stroma (Figure 3a), both of which have been previously linked with placental dysfunction and increased incidence of developing preeclampsia[20]. Hypoxia triggers TGF-β signalling and angiogenesis [21], and both TGF- β signalling and angiogenesis related pathways were enriched in the villous core stroma in response to SARS-CoV-2 infection (Figure 3b). Furthermore, pathways related to haemorrhage were upregulated and pathways related to vascular tension, such as hyperaldosteronin/renin pathways, acetylcholine channels, and olfactory receptors, were downregulated in the villous core stroma (Figure 3b). Trophoblast cells exhibited enrichment of pathways related to nitric oxide production (Figure 3c), which is a potent vasodilator[22]. Conversely, pathways related to calcium import[23] and vasoconstriction were downregulated in trophoblasts (Figure 3c), supporting the notion that the placenta actively reduces vascular tension during SARS-CoV-2 infection. In parallel, trophoblasts showed an increase in cell-cell adherence, communication, and transmembrane amino acid transport, including MTORC1 signalling[24], suggesting that nutritional uptake to the foetus is enhanced in response to SARS-CoV-2 infection (Figure 3a and 3c). Further, pathways related to allograft rejection and MHC molecules were decreased, suggestive of a defensive mechanism by the trophoblast layer to protect gestation (Figure 3a and 3c).

Markers associated with preeclampsia are elevated with SARS-CoV-2

Placentae from the SARS-CoV-2 group showed several markers that have been previously associated with hypoxia and placental dysfunction. For instance, the hypoxia and preeclampsia associated markers Fms Related Receptor Tyrosine Kinase 1 (*FLT1*) , *FLT4*, epidermal growth factor receptor (*EGFR),* and pappalysin-2 (*PAPPA2*) were increased in

trophoblasts from the SARS-CoV-2 group (Figure 3e-i)[25-28]. Additionally, markers associated

with placental dysfunction and oxidative stress such as Nitric oxide synthase 3 (*NOS3*)[29],

corticotrophin-releasing hormone (*CRH*)[30], kisspeptin 1 (*KISS1*)[31], Growth Differentiation

Factor 15 (*GDF15*)[32], and tissue factor pathway inhibitor 2 (*TFPI-2*)[33], were also elevated in

the trophoblasts from the SARS-CoV-2 group (Figure 3e, 3j-k). Transforming growth factor

b1 (*TGFβ-1*), and the PAPP-A2 substrates *IGFBP4/5* (Figure 3d and 3l), were also elevated

in the villous core stroma, both markers associated with increased preeclampsia risk[21, 34].

Given that a number of these pathways and genes are associated with preeclampsia, as well as

several recent studies reporting SARS-CoV-2 in predisposing pregnant individuals to

preeclampsia[5-10], we next assessed the enrichment of a preeclampsia-specific gene set

generated from published patient cohorts[35]. The gene set was generated by Moslehi et al.,

where they found 419 genes to be common between four studies examining preeclampsia

versus healthy pregnancies[35]. These 419 genes are involved in pathways relevant to

preeclampsia, such as oxidative stress, hypoxia, and immune response[35-37]. In our data, this

preeclampsia signature was positively enriched in patient samples from the SARS-CoV-2

group (NES 3.49, FDR <0.001; Figure 3m), which aligns with the positive enrichment of

hypoxia, immune, and oxidative stress, related pathways we observed in our studies (Figure

3a).

DISCUSSION

Using digital spatial profiling, we quantified the expression of key markers within distinct

cellular compartments of the placenta, providing a detailed picture of the molecular changes

occurring in response to SARS-CoV-2 infection. Although our study is limited by its

relatively small cohort, cross-sectional nature, and low availability of clinical data, it offers

valuable insights into the interplay between trophoblasts and the cells within the villous core

stroma in the placenta and how this relationship is influenced by SARS-CoV-2 infection.

Close examination of the transcriptional alterations occurring in the placental trophoblast and

villous core stroma in response to SARS-CoV-2, revealed a notable number of genes that are

enriched in biological pathways previously associated with placental dysfunction.

Trophoblasts from the SARS-CoV-2-infected group had significantly higher levels of *NOS3*

compared to the control group (Figure 3j). The upregulation of NOS3 is associated with

increased endogenous production of the vasodilator nitric oxide, as a response to altered

vascular reactivity, endothelial dysfunction, and hypertension[22, 38]. Interestingly, NOS has

been previously found to be highly upregulated to supraphysiological levels in animal models

of infection-mediated inflammation during pregnancy, leading researchers to hypothesise that

increased NOS may play a role in placental inflammation[39-41]. In response to increased NOS

by the trophoblasts, the villous core stromal showed increased expression in biological

pathways related to systemic pressure and vasodilation. This included the downregulation of

olfactory receptors, acetylcholine channels, and hyperaldosteronin/renin pathways, alongside

upregulated hypoxia pathways, suggesting deregulation of the vascular tone and blood

pressure due to a hypoxic environment[42].

Additional transcriptional analysis of trophoblast and villous core stromal compartments from

SARS-CoV-2-infected samples identified several transcriptional variations that have been

previously associated with preeclampsia (Figure 3). Trophoblasts had higher expression of

EGFR, a marker that increases with hypoxia and is known to upregulate *FLT1*, where

excessive release of soluble FLT1 by syncytiotrophoblasts is a characteristic marker of late

onset preeclampsia[43]. There was prominent increase of *PAPPA2* in the trophoblasts, the

latter considered to become upregulated in response to hypoxia, and placental pathologies,

including preeclampsia[44]. Notably, the PAPPA2 substrates IGFBP4 and IGFBP5 were

concurrently upregulated in the villous core stromal compartment, whereby the interaction of

PAPPA2 with IGFBP4/5 increases levels of IGF2, which was also increased in the villous

core stromal cells in the SARS-CoV-2-infected group (Figure 3d)[34]. Additionally, the villous

core stromal compartment had decreased levels of Isthmin-2 (*ISM2*), a placental marker that

is downregulated with preeclampsia[45]. *GDF15*, *TFPI-2*, *KISS1,* and *CRH* genes were also

upregulated in SARS-CoV-2-infected trophoblasts, all previously associated with placental

oxidative stress, hypertension, and preeclampsia[34, 46-48].

 In conclusion, our data suggest that the placenta from pregnancies with SARS-CoV-2 adopts

a transcriptional profile aligning with placental dysfunction that has been observed in

pregnant participants who develop 'preeclampsia-like' syndrome. Using digital spatial

profiling, our studies showcased the crosstalk between the trophoblast and villous core

stromal cell populations, and how this is enriched with pathways associated with placental

dysfunction. Our findings set the foundation for a more comprehensive understanding of

placental dysfunction in pregnant individuals with SARS-CoV-2 infection and offer

important insights into the potential mechanisms through which SARS-CoV-2 may impact

pregnancy outcomes and fetal development.

METHODS

Study Design

The SARS-CoV-2 group (n=7) consisted of pregnant, unvaccinated participants, who were symptomatic with COVID-19 in their third trimester (confirmed by RT-qPCR from nasopharyngeal swabs). Placental tissue samples were collected at birth at the Hospital de Clínicas (HC) and Hospital Nossa Senhora das Graças, Brazil, with ethics approval from the National Commission for Research Ethics (CONEP) under approval number 30188020.7.1001.0020[49]. The control group was comprised of archived placentae from nine COVID19-negative people collected during delivery at the Complexo Hospital de Clínicas, Universidade Federal do Paraná, Curitiba, Brazil between 2016 and 2018. To account for maternal co-morbidities, maternal age, and gestational age, the control group was selected to match these clinical features as presented in the SARS-CoV-2 group. Participant cohort and their clinical characteristics are summarised in Table 1. Morphological analysis was performed in all placentas from SARS-CoV-2-infected and control groups using the Amsterdam Placental Workshop Group Consensus Statement[50]. Histological sections were systematically identified and evaluated by two experienced pathologists to obtain samples for tissue microarray (TMA) construction, as described in a previous work[49]. Two TMAs were prepared from the placental samples, following the workflow demonstrated in Figure 1.

RNAscope

A serial section from the TMAs (4 um) was incubated with RNAscope probes targeting SARS-CoV-2 spike mRNA (nCoV2019, #848568-C3, ACDBio, CA, USA), as per manufacturer's instructions for automation on Leica Bond RX. DNA was visualised with Syto13 (500 nM, #S7575, ThermoFisher Scientific, MA, USA), and SARS-CoV-2 spike mRNA with TSA Plus CY5 (1:1500, #NEL745001KT, Akoya Biosciences, MA, USA). Fluorescent images were acquired with NanoString GeoMX DSP at 20×.

Digital spatial profiling with Nanostring GeoMX platform

TMA slides were freshly sectioned (4 um thick serial sections) and prepared according to the Nanostring GeoMX Digital Spatial Profiler (DSP) slide preparation for RNA profiling (NanoString, WA, USA). Slides were hybridised with the NanoString Technologies Whole Transcriptome Atlas (WTA) barcoded probe set (~18,000 genes), followed by fluorescent staining with Pan-Cytokeratin (PanCK, clone AE-1/AE-3, AF488, Santa Cruz NBP2-33200AF488, [2 µg/mL], CA, USA) to identify trophoblasts, vimentin (VIM, clone E-5, AF594, Santa Cruz sc-373717, [1 µg/mL], CA, USA) to identify endothelial and mesenchymal stromal cells, and SYTO83 to identify nuclei. With the aid of a trained placental pathologist, an area featuring an anchoring villus, and an area featuring a cluster of terminal villi, were designated as two areas of interest (AOI) within each placental core. Oligonucleotides linked to hybridized mRNA targets were cleaved separately for PanCK positive regions within each AOI, and separately for vimentin positive regions. Cleaved oligonucleotides were collected for counting using Illumina i5 and i7 dual indexing as described previously[51, 52]. Paired-end sequencing (2 × 75) was performed using an Illumina NextSeq550 up to 400M total aligned reads. Fastq files were processed using the Nanostring DND system and uploaded to the GeoMX DSP system where raw counts were aligned with their respective AOIs.

Data normalisation, differential expression analysis, and pathway enrichment analysis

Raw data were normalised to the 134 negative probes in the Human Whole Transcriptome Atlas probe set followed by upper quantile normalisation using the R package RUVseq[53]. Transcriptional data from PanCK positive regions within each AOI were normalised separately to the vimentin positive regions. Differential gene expression analysis between SARS-CoV-2 positive and negative groups was performed separately for PanCK positive

regions within each AOI, and separately for vimentin positive regions, using the R package

limma[54]. Bayesian adjusted t-statistic method was used where foetal sex and TMA slide

number were considered as co-variants. A fold change of +/- 1.5 and *P*-value $\leq$ 0.05

(adjusted for a false discovery rate of 5%) was considered significant. Pathway enrichment

analysis was performed using the Gene Set Enrichment Analysis program (GSEA, v4.3.2,

Broad Institute, MA, USA) for biological pathways obtained from the Molecular Signatures

Database (MSigDB, Broad Institute, Human v2022.1, MA, USA). The preeclampsia gene-set

was obtained from Moslehi et al[35]. GSEA parameters: 1000 permutations, weighted analysis.

Gene set enrichment data were further clustered and visualised using the R package vissE

with the parameters: computeMsigOverlap (thresh = 0.25), findMsigClusters (alg =

cluster_walktrap, minSize =2)[55].

AUTHOR CONTRIBUTIONS

325 **Designing research study**: NS, JR, MA, LdN, AK. **Conducting experiments**: NS, JM, AN,

326 LP, PZR, CMS, ETSS, LdN

327 **Acquiring data**: JM, AN, LP. **Analyzing data**: NS, IS, NM, AR, LM, GTB, FSFG, VC, AK.

328 **Writing the manuscript**: all authors

ACKNOWLEGMENTS

331 This study was funded by the Queensland University of Technology ECR funds for AK, JR,

332 MA, NS. The following authors are supported by fellowships from the NHMRC (AK –

333 1157741, GB - 2008542 and 1135898), US DOD (NS – PC190533). The authors thank Fred

334 Hutch pathology (Miki Haraguchi & Stephanie Weaver) for histology assistance.

DATA AVAILABILITY

The data generated in this study are available in the Gene Expression Omnibus (GEO) under GSE223612.

CONFLICTS OF INTEREST

Andy Nam and Liuliu Pan are employed by Nanostring Technologies. Nicolas Matigian is employed by QCIF Bioinformatics.

REFERENCES

1.   Bordt EA, Shook LL, Atyeo C, *et al.* Maternal SARS-CoV-2 infection elicits sexually dimorphic placental immune responses. *Sci Transl Med* 2021; **13**: eabi7428.

2.   Edlow AG, Li JZ, Collier AY, *et al.* Assessment of Maternal and Neonatal SARS-CoV-2 Viral Load, Transplacental Antibody Transfer, and Placental Pathology in Pregnancies During the COVID-19 Pandemic. *JAMA Netw Open* 2020; **3**: e2030455.

3.   Schwartz DA, Mulkey SB, Roberts DJ. SARS-CoV-2 Placentitis, Stillbirth and Maternal COVID-19 Vaccination: Clinical-Pathological Correlations. *Am J Obstet Gynecol* 2022.

4.   Racicot K, Mor G. Risks associated with viral infections during pregnancy. *J Clin Invest* 2017; **127**: 1591-1599.

5.   Palomo M, Youssef L, Ramos A, *et al.* Differences and similarities in endothelial and angiogenic profiles of preeclampsia and COVID-19 in pregnancy. *Am J Obstet Gynecol* 2022; **227**: 277 e271-277 e216.

6.   Mendoza M, Garcia-Ruiz I, Maiz N, *et al.* Pre-eclampsia-like syndrome induced by

severe COVID-19: a prospective observational study. *Bjog* 2020; **127**: 1374-1380.

7.   Serrano B, Bonacina E, Garcia-Ruiz I, *et al.* Confirmation of preeclampsia-like

syndrome induced by severe COVID-19: an observational study. *Am J Obstet*

Gynecol MFM 2022; **5**: 100760.

8.   Naeh A, Berezowsky A, Yudin MH, Dhalla IA, Berger H. Preeclampsia-Like

Syndrome in a Pregnant Patient With Coronavirus Disease 2019 (COVID-19). *J*

Obstet Gynaecol Can 2022; **44**: 193-195.

9.   Celewicz A, Celewicz M, Michalczyk M, *et al.* SARS CoV-2 infection as a risk factor

of preeclampsia and pre-term birth. An interplay between viral infection, pregnancy-

specific immune shift and endothelial dysfunction may lead to negative pregnancy

outcomes. *Ann Med* 2023; **55**: 2197289.

10.   Gonzalez-Vanegas O, Martinez-Perez O. SARS-CoV-2 Infection and Preeclampsia-

How an Infection Can Help Us to Know More about an Obstetric Condition. *Viruses*

2023; **15**.

11.   Schwartz DA, Avvad-Portari E, Babál P, *et al.* Placental Tissue Destruction and

Insufficiency From COVID-19 Causes Stillbirth and Neonatal Death From Hypoxic-

Ischemic Injury. *Arch Pathol Lab Med* 2022; **146**: 660-676.

12.   Huang C, Wang Y, Li X, *et al.* Clinical features of patients infected with 2019 novel

coronavirus in Wuhan, China. *The Lancet* 2020; **395**: 497-506.

13.   Maltepe E, Fisher SJ. Placenta: the forgotten organ. *Annu Rev Cell Dev Biol* 2015; **31**:

523-552.

14.   Buchrieser J, Degrelle SA, Couderc T, *et al.* IFITM proteins inhibit placental

syncytiotrophoblast formation and promote fetal demise. *Science* 2019; **365**: 176-180.

15.   Wedenoja S, Yoshihara M, Teder H, *et al.* Fetal HLA-G mediated immune tolerance
and interferon response in preeclampsia. *EBioMedicine* 2020; **59**: 102872.

16.   Hamldar S, Kiani SJ, Khoshmirsafa M, *et al.* Expression profiling of inflammation-
related genes including IFI-16, NOTCH2, CXCL8, THBS1 in COVID-19 patients.
Biologicals 2022; **80**: 27-34.

17.   Kulasinghe A, Tan CW, Ribeiro Dos Santos Miggiolaro AF, *et al.* Profiling of lung
SARS-CoV-2 and influenza virus infection dissects virus-specific host responses and
gene signatures. *Eur Respir J* 2022; **59**.

18.   Conca W, Alabdely M, Albaiz F, *et al.* Serum beta2-microglobulin levels in
Coronavirus disease 2019 (Covid-19): Another prognosticator of disease severity?
PLoS One 2021; **16**: e0247758.

19.   Jain NK, Tailang M, Jain HK, *et al.* Therapeutic implications of current Janus kinase
inhibitors as anti-COVID agents: A review. *Front Pharmacol* 2023; **14**: 1135145.

20.   Hu XQ, Zhang L. Hypoxia and Mitochondrial Dysfunction in Pregnancy
Complications. *Antioxidants (Basel)* 2021; **10**.

21.   Rana S, Burke SD, Karumanchi SA. Imbalances in circulating angiogenic factors in
the pathophysiology of preeclampsia and related disorders. *Am J Obstet Gynecol*
2022; **226**: S1019-S1034.

22.   Mukosera GT, Clark TC, Ngo L, *et al.* Nitric oxide metabolism in the human placenta
during aberrant maternal inflammation. *J Physiol* 2020; **598**: 2223-2241.

23.   Van Hove CE, Van der Donckt C, Herman AG, Bult H, Fransen P. Vasodilator
efficacy of nitric oxide depends on mechanisms of intracellular calcium mobilization
in mouse aortic smooth muscle cells. *Br J Pharmacol* 2009; **158**: 920-930.

24.   Jansson T, Aye IL, Goberdhan DC. The emerging role of mTORC1 signaling in
placental nutrient-sensing. *Placenta* 2012; **33 Suppl 2**: e23-29.

25.   Hastie R, Brownfoot FC, Pritchard N, *et al.* EGFR (Epidermal Growth Factor Receptor) Signaling and the Mitochondria Regulate sFlt-1 (Soluble FMS-Like Tyrosine Kinase-1) Secretion. *Hypertension* 2019; **73**: 659-670.

26.   Kramer AW, Lamale-Smith LM, Winn VD. Differential expression of human placental PAPP-A2 over gestation and in preeclampsia. *Placenta* 2016; **37**: 19-25.

27.   Neuman RI, Alblas van der Meer MM, Nieboer D, *et al.* PAPP-A2 and Inhibin A as Novel Predictors for Pregnancy Complications in Women With Suspected or Confirmed Preeclampsia. *J Am Heart Assoc* 2020; **9**: e018219.

28.   Lamale-Smith LM, Gumina DL, Kramer AW, *et al.* Uteroplacental Ischemia Is Associated with Increased PAPP-A2. *Reprod Sci* 2020; **27**: 529-536.

29.   Guerby P, Tasta O, Swiader A, *et al.* Role of oxidative stress in the dysfunction of the placental endothelial nitric oxide synthase in preeclampsia. *Redox Biol* 2021; **40**: 101861.

30.   Karteris E, Vatish M, Hillhouse EW, Grammatopoulos DK. Preeclampsia is associated with impaired regulation of the placental nitric oxide-cyclic guanosine monophosphate pathway by corticotropin-releasing hormone (CRH) and CRH-related peptides. *J Clin Endocrinol Metab* 2005; **90**: 3680-3687.

31.   Ibanoglu MC, Oskovi-Kaplan ZA, Ozgu-Erdinc AS, Kara O, Sahin D. Comparison of the Kisspeptin levels in early onset preeclampsia and late-onset preeclampsia. *Arch Gynecol Obstet* 2022; **306**: 991-996.

32.   Cruickshank T, MacDonald TM, Walker SP, *et al.* Circulating Growth Differentiation Factor 15 Is Increased Preceding Preeclampsia Diagnosis: Implications as a Disease Biomarker. *J Am Heart Assoc* 2021; **10**: e020302.

33.   Kobayashi H, Matsubara S, Yoshimoto C, Shigetomi H, Imanaka S. Tissue Factor

Pathway Inhibitors as Potential Targets for Understanding the Pathophysiology of

Preeclampsia. *Biomedicines* 2023; **11**.

34.   Nishizawa H, Pryor-Koishi K, Suzuki M*, et al.* Increased levels of pregnancy-

associated plasma protein-A2 in the serum of pre-eclamptic patients. *Mol Hum*

Reprod 2008; **14**: 595-602.

35.   Moslehi R, Mills JL, Signore C, Kumar A, Ambroggio X, Dzutsev A. Integrative

transcriptome analysis reveals dysregulation of canonical cancer molecular pathways

in placenta leading to preeclampsia. *Sci Rep* 2013; **3**: 2407.

36.   Aouache R, Biquard L, Vaiman D, Miralles F. Oxidative Stress in Preeclampsia and

Placental Diseases. *Int J Mol Sci* 2018; **19**.

37.   Redman CW, Staff AC. Preeclampsia, biomarkers, syncytiotrophoblast stress, and

placental capacity. *Am J Obstet Gynecol* 2015; **213**: S9 e1, S9-11.

38.   Mistry HD, Czajka AN, Kurlak LO, Pipkin FB, Taggart MJ, Tribe RT. Differential

placental caveolin-1 gene expression in women with pre-eclampsia. *Archives of*

Disease in Childhood - Fetal and Neonatal Edition 2011; **96**: Fa125-Fa126.

39.   Nowicki B, Singhal J, Fang L, Nowicki S, Yallampalli C. Inverse relationship

between severity of experimental pyelonephritis and nitric oxide production in

C3H/HeJ mice. *Infect Immun* 1999; **67**: 2421-2427.

40.   Ogando DG, Paz D, Cella M, Franchi AM. The fundamental role of increased

production of nitric oxide in lipopolysaccharide-induced embryonic resorption in

mice. *Reproduction* 2003; **125**: 95-110.

41.   Kong L, Zhang Q, Chao J*, et al.* Polarization of macrophages induced by Toxoplasma

gondii and its impact on abnormal pregnancy in rats. *Acta Trop* 2015; **143**: 1-7.

42. Wareing M, Bai X, Seghier F, *et al.* Expression and function of potassium channels in the human placental vasculature. *Am J Physiol Regul Integr Comp Physiol* 2006; **291**: R437-446.

43. Roberts JM, Rich-Edwards JW, McElrath TF, Garmire L, Myatt L, Global Pregnancy C. Subtypes of Preeclampsia: Recognition and Determining Clinical Usefulness. *Hypertension* 2021; **77**: 1430-1441.

44. Wagner PK, Otomo A, Christians JK. Regulation of pregnancy-associated plasma protein A2 (PAPPA2) in a human placental trophoblast cell line (BeWo). *Reprod Biol Endocrinol* 2011; **9**: 48.

45. Martinez C, Gonzalez-Ramirez J, Marin ME, *et al.* Isthmin 2 is decreased in preeclampsia and highly expressed in choriocarcinoma. *Heliyon* 2020; **6**: e05096.

46. Sugulle M, Dechend R, Herse F, *et al.* Circulating and placental growth-differentiation factor 15 in preeclampsia and in pregnancy complicated by diabetes mellitus. *Hypertension* 2009; **54**: 106-112.

47. Hogg K, Blair JD, McFadden DE, von Dadelszen P, Robinson WP. Early onset pre-eclampsia is associated with altered DNA methylation of cortisol-signalling and steroidogenic genes in the placenta. *PLoS One* 2013; **8**: e62969.

48. Zheng L, Huang J, Su Y, Wang F, Kong H, Xin H. Overexpression of tissue factor pathway inhibitor 2 attenuates trophoblast proliferation and invasion in preeclampsia. *Hum Cell* 2020; **33**: 512-520.

49. Rebutini PZ, Zanchettin AC, Stonoga ETS, *et al.* Association Between COVID-19 Pregnant Women Symptoms Severity and Placental Morphologic Features. *Front Immunol* 2021; **12**: 685919.

50.   Khong TY, Mooney EE, Ariel I, *et al.* Sampling and Definitions of Placental Lesions:

Amsterdam Placental Workshop Group Consensus Statement. *Arch Pathol Lab Med*

2016; **140**: 698-713.

51.   Merritt CR, Ong GT, Church SE, *et al.* Multiplex digital spatial profiling of proteins

and RNA in fixed tissue. *Nat Biotechnol* 2020; **38**: 586-599.

52.   Jerby-Arnon L, Neftel C, Shore ME, *et al.* Opposing immune and genetic mechanisms

shape oncogenic programs in synovial sarcoma. *Nat Med* 2021; **27**: 289-300.

53.   Risso D, Ngai J, Speed TP, Dudoit S. Normalization of RNA-seq data using factor

analysis of control genes or samples. *Nat Biotechnol* 2014; **32**: 896-902.

54.   Ritchie ME, Phipson B, Wu D, *et al.* limma powers differential expression analyses

for RNA-sequencing and microarray studies. *Nucleic Acids Res* 2015; **43**: e47.

55.   Dharmesh D. Bhuva CWT, Ning Liu, Holly J. Whitfield, Nicholas Papachristos, Sam

Lee, Malvika Kharbanda, Ahmed Mohamed, Melissa J. Davis. vissE: A versatile tool

to identify and visualise higher-order molecular phenotypes from functional

enrichment analysis. *bioRxiv* 2022.

TABLES AND FIGURE LEGENDS

Patient De-id.	Group	Sample Code	Participant Age	Gestational Age	Comorbidities	SARS-CoV-2 symptoms/severity	Fetal sex	Fetal Outcome	Delivery Method	Fetal Weight (grams)	Placental Weight (grams)	Macroscopic observations
20-3594	SARS-CoV-2	2	25-30	30-35	Hypothyroidism and hypertensive disorder in pregnancy	+/+	Male	Preterm	na	2450	448	Infarcts and intervillous thrombosis (<5%)
20-3561	SARS-CoV-2	4	35-40	25-30	Hypothyroidism	+/+	Female	Preterm	C-section	na	245	-
20-3744	SARS-CoV-2	8	25-30	30-35	Gestational diabetes, bipolar disorder, hypothyroidism and syphilis (treated)	+/-	Female	Preterm	C-section	na	412	Infarcts (<5%)
20-5105	SARS-CoV-2	12	25-30	35-40	None	-	Female	Term	C-section	2960	462	-
20-3369	SARS-CoV-2	13	25-30	35-40	Gestational diabetes and hyperthyroidism	+/-	Female	Term	Assisted Vaginal	2600	358	Retroplacental and marginal hematoma, infarcts (<5%)
20-5869	SARS-CoV-2	18	25-30	35-40	None	+/-	Male	Term	C-section	2345	370	-
20-2916	SARS-CoV-2	22	20-25	35-40	None	-	Female	Term	Assisted Vaginal	3030	650	-
16-7859	CONTROL	1	20-25	30-35	Hypothyroidism		Male	Preterm	C-section	1180	270	Placental hypoplasia
18-13016	CONTROL	3	20-25	35-40	Hypothyroidism and hypertension		Female	Preterm	Assisted Vaginal	2223	498	-
16-8315	CONTROL	5	15-20	35-40	Obesity		Female	Term	Assisted Vaginal	3810	514	-
18-4906	CONTROL	9	20-25	25-30	None		Male	Preterm	Assisted Vaginal	1205	248	-
18-14057	CONTROL	10	40-45	30-35	Diabetes, hypertension, bipolar disorder		Male	Preterm	C-section	1650	243	Placental hypoplasia
16-7599	CONTROL	11	25-30	35-40	Gestational diabetes		Male	Term	C-section	3460	480	-
16-3340	CONTROL	15	35-40	35-40	None		Female	Term	C-section	3005	395	-
18-9951	CONTROL	16	20-25	35-40	None		Male	Term	C-section	3690	574	-
16-6144	CONTROL	19	25-30	35-40	None		Male	Term	C-section	3345	394	-

Table 1: Clinical information of the SARS-CoV-2 and control cohort

Figure 1: Study design and cell deconvolution. (a) 1. Placental cores collected at delivery from the SARS-CoV-2 (n=7) and control (n=9) groups were assembled into tissue microarray slides (TMAs). 2. TMAs were stained with fluorescent markers to differentiate cell types within anchoring (pink outline) and terminal villi (red outline). Barcodes were cleaved and collected from each region of interest by UV light. *3.* Cleaved barcodes were sequenced and counted using an Illumina® sequencer in preparation for transcriptomic analysis. Data were normalised before downstream differential expression analysis. **(b)** Transcriptional cell deconvolution map. **(c-j)** Box-plots of indicated cell type proportions from 1b. AV: anchoring villi, TV: terminal villi, SARS-CoV-2 group is n=7 and control group is n=9.

Figure 2: Enrichment of SARS-CoV-2 related pathways. **(a)** Principal component analysis of normalised gene counts from trophoblasts and villous core stromal compartments from SARS-CoV-2 (n=7) and control (n=9) groups at the anchoring or terminal villi (AV; TV). **(b)** Principal component dimensions. **(c)** Upset plot of differential gene expression in trophoblasts and villous core stromal compartments from the AV and TV in response to

SARS-CoV-2 infection. The bar charts on the left indicate the total number of differentially expressed genes for the indicated sample group and the bar charts on the top show the gene overlap for the comparisons indicated by the purple lines. Black dots denote differentially expressed genes that are unique for the indicated sample group. Fold change +/- 1.5, *P*-value <= 0.05, FDR < 0.05. **(d)** Enrichment of significant SARS-CoV-2 related pathways from the Reactome database, in the SARS-CoV-2 and control samples. Blue: significantly negatively enriched, red: significantly positively enriched, grey: not significant. The full list of enriched pathways from the Reactome database can be found in Supplementary table 2 and table 3. **(e)** Gene set enrichment analysis (GSEA) plot of the SARS_COV_2_INFECTION pathway from the Reactome database and **(f)** INTERFERON_ALPHA_RESPONSE from the Hallmark database, in SARS-CoV-2 samples versus controls. **(g-j)** Normalised expression counts of IFI27, B2M, JAK1, and STAT3 genes, in trophoblast (TB) or villous core stroma (VCS) compartments from the SARS-CoV-2 (n=7) and control samples (n=9). ** *P*-value < 0.01, *** *P*-value < 0.001.

Figure 3: Enrichment of placental dysfunction pathways with SARS-CoV-2. (a) Differentially enriched pathways form the Hallmarks database in trophoblasts (TB) and villous core stroma (VCS) compartments in SARS-CoV-2 (n=7) vs control group (n=9). Colour gradient refers to the normalised enrichment score. **(b)** Enriched gene ontology biological processes (GO-BP) pathway clusters from the Molecular Signatures Database (MSigDb) generated using vissE, in the VCS and **(c)** TB. Top row depicts upregulated pathways, bottom row depicts downregulated pathways. **(d)** Volcano plot of gene expression from VCS or **(e)** TB in response to SARS-CoV-2 infection. Fold change (FC) +/- 1.5, *P*-value <= 0.05, FDR < 0.05, the full list of differentially expressed genes can be found in Supplementary table 4. **(f-l)** Normalised expression counts of indicated genes in TB or VCS

537 compartments from the SARS-CoV-2 (n=7) and control samples (n=9). * P-value < 0.05, **

538 P-value < 0.01, *** P-value < 0.001, NS: not significant. **(m)** Gene set enrichment analysis

539 (GSEA) plot of the preeclampsia signature generated by Moslehi et al, in SARS-CoV-2

540 samples versus controls.

Figure 1: Study design and cell deconvolution. (a) 1. Placental cores collected at delivery from the SARS-CoV-2 (n=7) and control (n=9) groups were assembled into tissue microarray slides (TMAs). 2. TMAs were stained with fluorescent markers to differentiate cell types within anchoring (pink outline) and terminal villi (red outline). Barcodes were cleaved and collected from each region of interest by UV light. *3.* Cleaved barcodes were sequenced and counted using an Illumina® sequencer in preparation for transcriptomic analysis. Data were normalised before downstream differential expression analysis. **(b)** Transcriptional cell deconvolution map. **(c-j)** Box-plots of indicated cell type proportions from 1b. AV: anchoring villi, TV: terminal villi, SARS-CoV-2 group is n=7 and control group is n=9.

Figure 2: Enrichment of SARS-CoV-2 related pathways. **(a)** Principal component analysis of normalised gene counts from trophoblasts and villous core stromal compartments from SARS-CoV-2 (n=7) and control (n=9) groups at the anchoring or terminal villi (AV; TV). **(b)** Principal component dimensions. **(c)** Upset plot of differential gene expression in trophoblasts and villous core stromal compartments from the AV and TV in response to SARS-CoV-2 infection. The bar charts on the left indicate the total number of differentially expressed genes for the indicated sample group and the bar charts on the top show the gene overlap for the comparisons indicated by the purple lines. Black dots denote differentially expressed genes that are unique for the indicated sample group. Fold change +/- 1.5, *P*-value <= 0.05, FDR < 0.05. **(d)** Enrichment of significant SARS-CoV-2 related pathways from the Reactome database, in the SARS-CoV-2 and control samples. Blue: significantly negatively enriched, red: significantly positively enriched, grey: not significant. The full list of enriched pathways from the Reactome database can be found in Supplementary table 2 and table 3. **(e)** Gene set enrichment analysis (GSEA) plot of the SARS_COV_2_INFECTION pathway from the Reactome database and **(f)** INTERFERON_ALPHA_RESPONSE from the Hallmark database, in SARS-CoV-2 samples versus controls. **(g-j)** Normalised expression counts of IFI27, B2M, JAK1, and STAT3 genes, in trophoblast (TB) or villous core stroma (VCS) compartments from the SARS-CoV-2 (n=7) and control samples (n=9). ** *P*-value < 0.01, *** *P*-value < 0.001.

Figure 3: Enrichment of placental dysfunction pathways with SARS-CoV-2.

(a) Differentially enriched pathways form the Hallmarks database in trophoblasts (TB) and villous core stroma (VCS) compartments in SARS-CoV-2 (n=7) vs control group (n=9). Colour gradient refers to the normalised enrichment score. **(b)** Enriched gene ontology biological processes (GO-BP) pathway clusters from the Molecular Signatures Database (MSigDb) generated using vissE, in the VCS and **(c)** TB. Top row depicts upregulated pathways, bottom row depicts downregulated pathways. **(d)** Volcano plot of gene expression from VCS or **(e)** TB in response to SARS-CoV-2 infection. Fold change (FC) +/- 1.5, *P*-value <= 0.05, FDR < 0.05, the full list of differentially expressed genes can be found in Supplementary table 4. **(f-l)** Normalised expression counts of indicated genes in TB or VCS compartments from the SARS-CoV-2 (n=7) and control samples (n=9). * *P*-value < 0.05, ** *P*-value < 0.01, *** *P*-value < 0.001, NS: not significant. **(m)** Gene set enrichment analysis (GSEA) plot of the preeclampsia signature generated by Moslehi et al, in SARS-CoV-2 samples versus controls.

Patient De-id.	Group	Sample Code	Participant Age	Gestational Age	Comorbidities	SARS-CoV-2 symptoms/ severity	Fetal sex	Fetal Outcome	Delivery Method	Fetal Weight (grams)	Placental Weight (grams)	Macroscopic observations
20-3594	SARS-CoV-2	2	25-30	30-35	Hypothyroidism and hypertensive disorder in pregnancy	+/+	Male	Preterm	na	2450	448	Infarcts and intervillous thrombosis (<5%)
20-3561	SARS-CoV-2	4	35-40	25-30	Hypothyroidism	+/+	Female	Preterm	C-section	na	245	-
20-3744	SARS-CoV-2	8	25-30	30-35	Gestational diabetes, bipolar disorder, hypothyroidism and syphilis (treated)	+/-	Female	Preterm	C-section	na	412	Infarcts (<5%)
20-5105	SARS-CoV-2	12	25-30	35-40	None	-	Female	Term	C-section	2960	462	-
20-3369	SARS-CoV-2	13	25-30	35-40	Gestational diabetes and hyperthyroidism	+/-	Female	Term	Assisted Vaginal	2600	358	Retroplacental and marginal hematoma, infarcts (<5%)
20-5869	SARS-CoV-2	18	25-30	35-40	None	+/-	Male	Term	C-section	2345	370	-
20-2916	SARS-CoV-2	22	20-25	35-40	None	-	Female	Term	Assisted Vaginal	3030	650	-
16-7859	CONTROL	1	20-25	30-35	Hypothyroidism		Male	Preterm	C-section	1180	270	Placental hypoplasia
18-13016	CONTROL	3	20-25	35-40	Hypothyroidism and hypertension		Female	Preterm	Assisted Vaginal	2223	498	-
16-8315	CONTROL	5	15-20	35-40	Obesity		Female	Term	Assisted Vaginal	3810	514	-
18-4906	CONTROL	9	20-25	25-30	None		Male	Preterm	Assisted Vaginal	1205	248	-
18-14057	CONTROL	10	40-45	30-35	Diabetes, hypertension, bipolar disorder		Male	Preterm	C-section	1650	243	Placental hypoplasia
16-7599	CONTROL	11	25-30	35-40	Gestational diabetes		Male	Term	C-section	3460	480	-
16-3340	CONTROL	15	35-40	35-40	None		Female	Term	C-section	3005	395	-
18-9951	CONTROL	16	20-25	35-40	None		Male	Term	C-section	3690	574	-
16-6144	CONTROL	19	25-30	35-40	None		Male	Term	C-section	3345	394	-

Table 1: Clinical information of the SARS-CoV-2 and control cohort