1 **Lipopolysaccharide integrity primes bacterial sensitivity to a cell wall-degrading intermicrobial toxin**
2

3 Kristine L Trotta[1], Beth M Hayes[1], Johannes P Schneider[2], Jing Wang[2], Horia Todor[3], Patrick Rockefeller
4 Grimes[1], Ziyi Zhao[1], William L Hatleberg[4], Melanie R Silvis[3], Rachel Kim[1], Byoung Mo Koo[3], Marek Basler[2],
5 Seemay Chou[1]*
6
7 [1] Department of Biochemistry & Biophysics, University of California – San Francisco, San Francisco, CA, USA
8 [2] Focal Area Infection Biology, Biozentrum, University of Basel, Klingelbergstrasse 50/70, CH - 4056
9 Basel, Switzerland
10 [3] Department of Cell and Tissue Biology, University of California – San Francisco, San Francisco, CA, USA
11 [4] Independent Researcher, Pittsburgh, Pennsylvania, USA
12
13 *Corresponding author: Seemay Chou (seemaychou@gmail.com)

14
15 **ABSTRACT**
16

Gram-negative bacteria can antagonize neighboring microbes using a type VI secretion system (T6SS) to deliver
toxins that target different essential cellular features. Despite the conserved nature of these targets, T6SS
potency can vary across recipient species. To understand the molecular basis of intrinsic T6SS susceptibility, we
screened for essential *Escherichia coli* genes that affect its survival when antagonized by a cell wall-degrading
T6SS toxin from *Pseudomonas aeruginosa*, Tae1. We revealed genes associated with both the cell wall and a
separate layer of the cell envelope, surface lipopolysaccharide, that modulate Tae1 toxicity *in vivo*. Disruption of
lipopolysaccharide synthesis provided *Escherichia coli (Eco)* with novel resistance to Tae1, despite significant cell
wall degradation. These data suggest that Tae1 toxicity is determined not only by direct substrate damage, but
also by indirect cell envelope homeostasis activities. We also found that Tae1-resistant *Eco* exhibited reduced cell
wall synthesis and overall slowed growth, suggesting that reactive cell envelope maintenance pathways could
promote, not prevent, self-lysis. Together, our study highlights the consequences of co-regulating essential
pathways on recipient fitness during interbacterial competition, and how antibacterial toxins leverage cellular
vulnerabilities that are both direct and indirect to their specific targets *in vivo*.

INTRODUCTION

Many bacteria live in mixed-species microbial communities where they compete with each other for limited space and resources[1]. Intermicrobial competition is mediated by a diverse array of molecular strategies that can exclude or directly interfere with other microbes, both near and far[2]. Nearly 25% of Gram-negative bacteria encode a type VI secretion system (T6SS)[3], which antagonizes neighboring cells by injection of toxic protein effectors into a recipient cell[4–6]. The opportunistic human pathogen *Pseudomonas aeruginosa* (*Pae*) harbors an interbacterial T6SS (H1-T6SS)[7] that can kill the model bacterium *Escherichia coli* (*Eco*)[8–10]. Studies of H1-T6SS-mediated competition between these genetically tractable species have provided fundamental insights into the molecular basis of T6SS function and regulation.

Key to *Pae* H1-T6SS toxicity are its seven known effectors, each with a unique biochemical activity[6,11–15]. The T6S amidase effector 1 (Tae1) from *Pae* plays a dominant role in H1-T6SS-dependent killing of *Eco* by degrading peptidoglycan (PG), a structural component of the cell wall that is critical for managing cell shape and turgor[16,17]. Early efforts to understand Tae1 toxicity focused on its *in vitro* biochemical activity against PG, which offered key insights about how H1-T6SS targets select bacterial species. Tae1 specifically digests γ-D-glutamyl-meso-2,6-diaminopimelic acid (D-Glu-*m*DAP) peptide bonds, which are commonly found in PG from Gram-negative bacteria [8,18]. Tae1 toxicity is further restricted to non-kin cells through a *Pae* cognate immunity protein, T6S amidase immunity protein 1 (Tai1), which binds and inhibits Tae1 in kin cells[11,19,20].

However, biochemical specificity is not sufficient to explain the toxicity and organismal selectivity of T6SS effectors *in vivo*. Bacteria antagonized by T6SSs ('recipients') can actively regulate effector toxicity through adaptive stress responses. *Eco* upregulates its envelope stress responses Rcs and BaeSR after exposure to the *Vibrio cholerae* (V52) T6SS effectors TseH (a PG hydrolase)[21] and TseL (a lipase)[22], suggesting that *Eco* could counter cell envelope damage by re-enforcing its surface[23]. Similarly, *Bacillus subtilis* triggers protective sporulation in response to a *Pseudomonas chlororaphis* (PCL1606) T6SS effector, Tse1 (a muramidase)[24]. Additional recipient-cell coordinators of T6SS effector toxicity include reactive oxygen species[25] and glucose-dependent gene expression[26]. These studies demonstrate that T6SS effector toxicity *in vivo* may also depend on downstream adaptive features of recipient cells.

The cell wall is a complex and dynamic substrate that is actively regulated to protect the cell[27–32], yet *Eco* is highly susceptible to lysis by Tae1 *in vivo*. We therefore hypothesized that Tae1 activity promotes H1-T6SS-mediated lysis in *Eco* through a unique strategy to overcome neutralization by the recipient cell. In this study, we investigated the *Eco* cellular features that drive its intrinsic sensitivity to H1-T6SS and the Tae1 toxin during interbacterial competition with *Pae*. Many T6SS effectors target essential cell features, so we screened the entire complement of essential *Eco* genes (plus some conditionally essential PG genes) for Tae1 susceptibility determinants. This approach complements previous genetic screens for T6SS recipient fitness which focused on nonessential gene candidates[33,34]. While cell wall-related genes indeed impacted *Eco* susceptibility to Tae1, we also discovered a strong relationship between survival and another component of the cell envelope, lipopolysaccharide (LPS). Perturbation of LPS synthesis genes *msbA* and *lpxK* rendered *Eco* conditionally resistant to lysis by Tae1 from *Pae*. Our work revealed that LPS-related resistance was mediated through cell-biological processes that were independent of the biochemical Tae1–PG interaction. Our findings suggest that beyond biochemical specificity and adaptive stress responses lies a role for essential homeostatic processes in defining T6SS effector toxicity *in vivo*.

RESULTS

Adaptation of native T6SS competitions to study *Eco* susceptibility to Tae1

We developed an *in vivo* screen for genetic interactions between the cell wall-degrading H1-T6SS effector Tae1 from *Pae* and the model target bacterium *Eco*. Our screen had two fundamental design requirements: (1) the ability to distinguish between general (T6SS-dependent) and specific (Tae1-dependent) genetic interactions, and (2) the capacity to test a broad array of target cell features. We adapted an established interbacterial competition co-culture assay between H1-T6SS-active *Pae* and *Eco*, the outcome of which is sensitive to the specific contribution of Tae1[8]. In this assay *Eco* exhibits a greater fitness advantage when competed against *Pae* missing *tae1* (*Pae^Δtae1*) relative to an equivalent control strain (*Pae^WT*) (**Figure 1a**). We hypothesized that the *Pae*:*Eco* co-culture assay could be leveraged to quantitatively compare recipient cell fitness against both Tae1 (toxin-specific fitness) and the H1-T6SS (Tae1-independent fitness) in interbacterial competition.

To screen broadly for *Eco* determinants, we adopted an established *Eco* CRISPR interference (CRISPRi)
platform that generates hypomorphic mutants through intermediate gene expression knockdowns (KDs)[35]. In
contrast to knock-outs or transposon mutagenesis studies, CRISPRi is systematically amenable to essential
genes and thus provided an opportunity to make unique insights about genes that are typically challenging to
screen for. This includes many essential (or conditionally essential) genes related to peptidoglycan (PG)
metabolism, whose KDs
we predicted would impact
Tae1 toxicity. In this
CRISPRi system, inducible
sgRNA expression is
coupled with constitutive
dCas9 expression to
conditionally repress
transcription at specific loci
with and without induction
("induced" and "basal"
CRISPRi, respectively)
(**Supplemental Figure 1a**).
In total, our CRISPRi
collection was composed of
596 *Eco* strains with KDs
representing most cellular
functions as defined by the
NCBI clusters of
orthologous genes (COG)
system (**Supp. Fig. 1b**).
Our collection also included
50 negative control strains
with non-targeting sgRNAs,
including *rfp-KD*, to ensure
CRISPRi alone did not
impact inherent *Eco*
susceptibility to *Pae* (**Supp.**
Fig. 2a).
123
For the interbacterial
competition screen, we co-
cultured *Pae* with the
pooled *Eco* CRISPRi
collection to test
competitive fitness across
all KD strains in parallel
(**Fig. 1b**). To compare
Tae1-dependent and -
independent fitness
determinants, we
conducted screens against
H1-T6SS-active *Pae*
strains that either secrete
Tae1 (*Pae^WT^*; *ΔretSΔpppA*)
or are Tae1-deficient
(*Pae^Δtae1^*;
ΔretSΔpppAΔtae1). As
negative controls, we also
competed the *Eco*
collection against a
genetically H1-T6SS-
inactivated *Pae* strain
(*Pae^inactive^*;

Figure 1: Adaptation of native T6SS competitions to study *Eco* susceptibility to Tae1
a) Tae1 from *Pseudomonas aeruginosa* (*Pae*) degrades the *Escherichia coli* (*Eco*) cell wall to promote H1-T6SS-mediated lysis. *Left*: *Pae^WT^* (dark grey) outcompetes *Eco* (light grey) using H1-T6SS to deliver a cocktail of toxic effectors, including Tae1 (triangle) which degrades peptidoglycan (orange). *Center*: Tae1 hydrolyzes D-Glu-*m*DAP peptide bonds in the donor stem peptides of 4,3-crosslinked peptidoglycan. *Right*: *Pae^Δtae1^* is less effective at outcompeting *Eco* using H1-T6SS. **b) Method for a genetic screen to test *Eco* gene function toward fitness against Tae1 from *Pae*.** *Left*: *Pae* strains (dark grey) were engineered with modified H1-T6SS activities including: constitutively active *Pae^WT^* (*ΔretSΔpppA*), Tae1-deficient *Pae^Δtae1^* (*ΔretSΔpppAΔtae1*), and T6SS-inactive *Pae^inactive^* (*ΔretSΔpppAΔicmF*). Each *Pae* strain was mixed with a pool of *Eco* KD (knockdown) strains engineered to conditionally disrupt a single gene (CRISPRi induced vs. basal). *Center*: each *Pae* strain was cocultured with an *Eco* CRISPRi strain pool for 6 hours. The *Eco* CRISPRi strain pool was also grown for 6 hours without *Pae* (*Eco^ctrl^*) as a negative control. Genomic sgRNA sequences harvested from competitions were amplified into Illumina sequencing libraries. *Right*: sgRNA barcode abundances after 6 hours were used to calculate a normalized log$_2$ fold-change (L2FC) for each *Eco* KD strain under each condition. Above a -log10 *p*-value cutoff, a positive L2FC value indicates a KD strain which is resistant to a given condition relative to WT *Eco*; a negative L2FC value indicates a KD strain which is sensitive to a given condition relative to WT *Eco*. **c) Interbacterial competition and CRISPRi induction have distinct effects on the composition of the *Eco* CRISPRi strain library.** Principal component analysis of *Eco* library composition after competition against *Pae^WT^* (blue), *Pae^Δtae1^* (purple), or *Pae^inactive^* (green), with induced (solid circles) or basal (hollow circles) CRISPRi induction. Four biological replicates per condition.

ΔretSΔpppAΔicmF) and included a condition in which the collection was grown without *Pae* present (*Eco^ctrl*).
Experiments were performed under both induced and basal CRISPRi conditions to distinguish between general
Eco fitness changes and those due to transcriptional knockdown. We used high-throughput sequencing to
quantify KD strain abundance at the beginning and end of each six-hour competition. To understand the
contribution of each KD to *Eco* survival against *Pae* in the presence or absence of H1-T6SS or Tae1, we
calculated log$_2$ fold-change (L2FC) values for each KD strain after competition and normalized against abundance
after growth without competition (*Eco^ctrl*)[36,37]. Across four biological replicates per condition, L2FC values were
reproducible (**Supp. Fig. 3a**; median Pearson's *r* between all replicates = 0.91). L2FC was used as a proxy for
competitive fitness of KD strains across different competition conditions.
157
To determine if our screen was sensitive to the effects of Tae1, H1-T6SS, and CRISPRi, we conducted a principal
component analysis of L2FC values for each strain under every competition condition (**Fig. 1c**). We observed
clear separation of datasets by CRISPRi induction (induced versus basal) across the first principal component
(PC1; 85.41%), indicating that KD induction was a major contributor to the performance of the KD library in the
pooled screen. We also observed clustering of datasets according to *Pae* competitor (PC2; 2.66%). These results
indicate that each *Pae* competitor yielded a distinct effect on the fitness of the CRISPRi library and demonstrates
that our screen was sensitive to the presence (*Pae^WT*) or absence (*Pae^Δtae1*) of Tae1 delivery from H1-T6SS.
From these data we conclude that our screen successfully captured the unique impacts of CRISPRi, Tae1, and
H1-T6SS on pooled *Eco* CRISPRi libraries during interbacterial competition.
167
168 **CRISPRi reveals toxin-specific and non-specific determinants of *Eco* fitness against H1-T6SS**
169
To reveal specific *Eco* genes that shape intrinsic susceptibility to H1-T6SS-mediated antagonism, we identified
KD strains which were significantly depleted or enriched at least three-fold (L2FC<-1.585 for depletion or
L2FC>1.585 for enrichment, and -log10 p-adj <0.05) after competition against *Pae^WT, Pae^Δtae1, or Pae^inactive*. Our
goal was to prioritize KDs which had a unique effect on fitness against *Pae^WT* relative to conditions lacking Tae1.
With CRISPRi induced, we found a select cohort of KDs with significant loss of fitness (*n*=12) or gain of fitness
(*n*=11) against *Pae^WT* (**Fig. 2a**). We were surprised that some KDs caused resistance to Tae1 despite the
combined challenge of essential gene depletion and H1-T6SS antagonism.
177
Competition against *Pae^WT* with basal CRISPRi diminished the pool of significant candidate KDs (**Supp. 4a**),
reinforcing our observation that KD strains' fitness changes against *Pae* are dependent on CRISPRi induction.
Against *Pae^Δtae1* (CRISPRi induced), we observed seventeen KDs with significant fitness changes (**Fig. 2b**) which
were also CRISPRi-dependent (**Supp. 4b**). These KDs were mostly distinct from those that affected *Eco* fitness
against *Pae^WT* (**Fig. 2c**). These results indicate that the presence or absence of Tae1 had a unique effect on the
T6SS competition and thus had a distinct impact on KD fitness. Finally, we found few candidate KDs that affected

Figure 2: CRISPRi reveals toxin-specific and non-specific determinants of *Eco* fitness against H1-T6SS
a-b) CRISPRi knockdowns promote *Eco* survival against *Pae^WT* (a) and *Pae^Δtae1*(b). Volcano plots showing log$_2$-fold change (L2FC)
values for each KD strain after interbacterial competition (induced CRISPRi). Data shown: mean from four biological replicates. Statistical
test: Wald test. Vertical dotted lines indicate arbitrary cutoffs for L2FC at x =-1.58 and x=1.58 (absolute FC x=-3 or x= 3). Horizontal dotted
line indicates statistical significance cutoff for log$_{10}$ adjusted p-value (≤ 0.05). Orange points represent KDs with L2FC ≥ 1.58 or ≤ -1.58 and
log10-adj. *p*-value ≤0.05. Dark purple points represent non-targeting negative control KDs (*n*=50). Lavender points represent KDs that do
not meet cutoffs for L2FC or statistical test. **c) T6SS competitions identify CRISPRi strains with distinct fitness changes against T6SS
and Tae1.** Venn diagram of total KDs significantly enriched OR depleted after competition against *Pae^WT* (n=23), *Pae^Δtae1*(n=17), and
Pae^inactive(n=5).

fitness against *Pae^inactive* regardless
of CRISPRi induction condition
(**Supp. 4a-b**)**,** suggesting that most
significant phenotypes were H1-
T6SS-dependent, if not Tae1-
dependent. In fact, L2FC values in
Pae^inactive and *Eco^ctrl* datasets had
high correlation (**Supp. 4 c-d**,
median Pearson correlation *r*=
0.98), indicating that *Pae* is a
neutral co-culture partner with its
H1-T6SS inactivated.
With our interest in Tae1-specific
determinants, we focused our
attention on the 20 KDs which had
a unique effect on *Eco* fitness
against Tae1 (*Pae^WT* +CRISPRi
induced; **Table 1**). Most KDs in this
group targeted genes related to the
cell envelope (COG category M:
cell wall/membrane/envelope
biogenesis, *n*=13/20). Composed
of concentric layers of inner
membrane (IM), cell wall PG, outer
membrane (OM), and
lipopolysaccharide (LPS)[38](**Fig.**
3b), the cell envelope is a critical
structure for protecting *Eco* against
environmental stress. Tae1-
sensitized strains were dominated
by gene targets related to the
synthesis of PG (*murA, ftsI, murC,*
murI, mcrB, murJ). Given that Tae1
targets the cell wall, these results
support our initial hypothesis that
PG structural integrity or
composition are direct
determinants of Tae1 *susceptibility.*

KD target	pathway/process	Avg. L2FC (*Pae^WT*)	fitness against *Pae^WT*
murA	PG synthesis	-7.40	sensitive
ftsI	Cell division	-6.85	sensitive
accD	Lipid metabolism	-4.37	sensitive
lptC	LPS transport	-3.35	sensitive
murC	PG synthesis	-2.61	sensitive
bamA	OM protein assembly	-2.46	sensitive
murI	PG synthesis	-1.86	sensitive
mrcB	PG synthesis	-1.60	sensitive
murJ	PG transport	-1.59	sensitive
pssA	Lipid metabolism	1.77	resistant
hemE	Heme metabolism	1.79	resistant
msbA	LPS transport	1.84	resistant
waaA	LPS synthesis	1.91	resistant
lpxA	LPS synthesis	2.18	resistant
ffs	Membrane trafficking/ secretion	2.25	resistant
acpP	Lipid metabolism	2.25	resistant
ffh	Membrane trafficking/ secretion	2.30	resistant
kdsB	LPS synthesis	2.35	resistant
lpxK (lpxK_-1as)	LPS synthesis	2.39	resistant
lpxK (lpxK_32as)	LPS synthesis	2.69	resistant

Table 1: Cell envelope gene KDs develop strong fitness changes against Tae1 in competition. KDs that target PG synthesis can increase *Pae^WT* sensitivity, while targeting other cell envelope processes can result in sensitivity or resistance. Data shown: normalized L2FC values for all 20 KD strains with unique and significant fitness changes against *Pae^WT* (which secretes Tae1); average of four biological replicates.

KDs related to lipid membrane metabolism and transport offered either resistance (*pssA, acpP, ffs, ffh)* or
sensitivity (*accD, bamA*) to Tae1, indicating that cell envelope factors indirect to the effector-substrate interaction
could impact Tae1 toxicity. To our surprise, most KDs that rendered *Eco* resistant to *Pae^WT* were related to LPS
synthesis and transport. Tae1 is not known to directly interact with the IM, OM, or LPS as part of its molecular
mechanism but metabolic crosstalk does occur between the PG, LPS, and lipid biosynthesis pathways[31,39]. Thus,
our data raised the possibility that regulation of other cell envelope structures could also be implicated in
mediating cell wall attack.
229
230 ***msbA-KD* disrupts LPS biosynthesis and imparts Tae1 resistance**
231
To investigate the hypothesis that non-PG components of the cell envelope may also shape Tae1 toxicity, we
focused downstream studies on Tae1-resistant KDs related to the synthesis of LPS, an essential lipidated surface
sugar that offers protection and structure to the OM[40]. Candidate KDs targeted highly-conserved, essential genes
in Kdo_2-Lipid A synthesis and transport (*lpxA, lpxK, kdsB, waaA, msbA*) (**Fig. 3a**). Kdo_2-Lipid A synthesis is the
most-upstream arm of LPS biosynthesis with rate-limiting control over the entire pathway[41,42]. In our screen, the
strongest resistance phenotypes we observed were in KDs targeting *lpxK* (*lpxK_-1as* and *lpxK_32as*) (**Table 1**).
LpxK is a kinase that phosphorylates the Lipid-A intermediate tetraacyldisaccharide 1-phosphate to form Lipid
IV_A[43,44]. In *Eco*, *lpxK* is in an operon with *msbA* (**Fig. 3b**), which encodes the IM Kdo_2-Lipid A flippase MsbA[45,46]. A
KD of *msbA* (*msbA_40as*) also conferred resistance to *Pae^WT* in our screen(**Table 1**).
241

Figure 3: *msbA-KD* disrupts LPS biosynthesis and imparts Tae1 resistance
a) Tae1 resistance emerges in KDs that target the lipopolysaccharide (LPS) biosynthesis pathway. Schematic representation of the LPS biosynthesis pathway and its distribution across the *Eco* cell envelope. Genes with KDs that render *Eco* resistant to Pae^{WT} are involved in the biosynthesis of Kdo2-Lipid A (*lpxA, lpxK, kdsA, waaA, msbA*). Note that Tae1 (grey triangle) targets peptidoglycan (PG), which is physically separate from Kdo₂-Lipid A synthesis in the IM. **b) The Kdo₂-Lipid A biogenesis genes *msbA* and *lpxK* are integral members of the *ycaI-msbA-lpxK-ycaQ* operon in *Eco*.** *msbA* (orange) and *lpxK* (purple) are co-expressed at the transcriptional level. **c-e) *msbA-KD* loses sensitivity to Tae1 in interbacterial competition against *Pae* but *lpxK-KD* does not.** Interbacterial competitions between *Pae* (Pae^{WT}, $Pae^{\Delta tae1}$, $Pae^{inactive}$) and *rfp-KD* (c; grey), *msbA-KD* (d; orange), or *lpxK-KD* (e; purple). Data shown are average fold-change in *Eco* colony forming units (CFUs) after 6 hours of competition (geometric mean 3 biological replicates ± s.d). Statistical test: unpaired two-tailed *t*-test; *p*-value ≤0.05 displayed in bold font. **f-h) Kdo2-Lipid A mutants develop structural damage to membranes.** Cryo-EM tomographs of *rfp-KD* (f), *msbA-KD* (g), and *lpxK-KD* (h) with CRISPRi induced, highlighting cross-sections of the cell envelope (including IM and OM; black arrows). Deformed membranes (red arrows) and novel intracellular vesicles (blue arrows) are demarcated in (g) and (h). Scale bar: 100nm.

We first experimentally validated pooled screen results by individually testing *lpxK-KD* and *msbA-KD* fitness in binary competitions against *Pae*. We regenerated and validated KD strains for *lpxK* (*lpxK_-1as*; "*lpxK-KD*") and *msbA* ("*msbA-KD*") for use in these experiments (**Supp. 6**). Consistent with our screen, *msbA-KD* gained Tae1-specific resistance in H1-T6SS-mediated competitions (**Fig. 3d**), exhibiting loss of sensitivity to Pae^{WT} relative to $Pae^{\Delta tae1}$. In contrast, we could not validate Tae1 resistance for *lpxK-KD* (**Fig. 3e**). Like *rfp-KD* (**Fig. 3c**), *lpxK-KD* maintains sensitivity to Pae^{WT} relative to $Pae^{\Delta tae1}$. The gene expression of *msbA* and *lpxK* are co-dependent, so we were surprised that *msbA-KD* and *lpxK-KD* did not equally reproduce Tae1 resistance. However, CRISPRi-dependent phenotypes could be controlled by factors such as transcriptional polar effects or off-target CRISPRi effects. To address their phenotypic disparities, we quantified transcriptional KD efficacy and specificity for *lpxK-KD* and *msbA-KD* with qRT-PCR. For *msbA-KD* with CRISPRi induced, we found repression of *msbA* (29-fold), l*pxK* (15-fold), and *ycaQ* (3.6-fold) expression (**Supp. Fig. 6a**). Thus, owing to downstream polar effects, our *msbA-KD* strain is a KD of both LPS candidate genes, *msbA* and *lpxK*. Conversely, *lpxK-KD* only repressed *lpxK* (71-fold) and *ycaQ* (11-fold) (**Supp. Fig. 6b**), but not *msbA*. Therefore, *msbA-KD* and *lpxK-KD* yield distinct transcriptional consequences despite targeting the same operon using CRISPRi.

Next, we investigated phenotypic consequences of inducing CRISPRi in *msbA-KD* and *lpxK-KD* by comparing
their cellular morphologies with cryo-electron tomography. Disruption of *msbA* and *lpxK* typically leads to
structural deformation in the *Eco* cell envelope from aberrant accumulation of Kdo₂-Lipid A intermediates in the

IM[44,46,47]. Unlike *rfp-KD* negative control cells (**Fig. 3e**), *msbA-KD* cells developed irregular buckling in the IM and
OM (**Fig. 3f**, red arrows). We also observed vesicular or tubular membrane structures within the cytoplasm (**Fig.**
3f, blue arrows). Such structural abnormalities are consistent with physical crowding of Kdo_2-Lipid A intermediates
in the IM that are relieved by vesicular internalization. On the other hand, while *lpxK-KD* had a distended IM and
vesicles (**Fig. 3g**, red and blue arrows), the OM appeared smooth and regular. This phenotypic divergence points
to two distinct KD effects: defects in the IM (both *msbA-KD* and *lpxK-KD*) and defects in the OM (*msbA-KD* only).
Together with our transcriptional analyses, these results demonstrate that *msbA-KD* and *lpxK-KD* have unique
consequences for LPS integrity and Tae1 susceptibility despite targeting the same operon. We focused the
remainder of our study on the validated *msbA-KD* strain which damages the IM and OM.
310
311 **Resistance to Tae1 in *msbA-KD* is independent of cell wall hydrolysis**
312
Identifying *msbA* and *lpxK* as potential Tae1 resistance determinants provided us a chance to study mechanisms
by which LPS impacts
susceptibility to cell wall
damage. Such mechanisms
could span several scales
including: direct Tae1-PG
interactions (**Fig. 4**), cellular
responses to Tae1 hydrolysis
(**Fig. 5**), broad physiological
conditions that affect
mechanical lysis (**Fig. 6**), or
some combination of these. To
investigate, we used an
orthogonal *in vivo* assay to
directly test the effect of Tae1
activity in *msbA-KD* cells in the
absence of *Pae* and other co-
delivered H1-T6SS toxins. We
measured lysis for *rfp-KD* and
msbA-KD upon induction of
exogenous wild-type Tae1
(Tae1[WT]) expression in the cell
wall-containing periplasm[8,48]and
found that *msbA-KD* had
increased survival against
Tae1[WT] relative to *rfp-KD* (**Fig.**
4a). *Eco* resistance was
dependent on Tae1 activity, as
evidenced by loss of the *msbA-*
KD resistance phenotype with
catalytically-attenuated
Tae1[C30A] (**Fig. 4b**) and no-
enzyme (empty) (**Fig. 4c**)
controls. There were no major
differences in Tae1 expression
levels across conditions (**Supp.**
Fig. 7a-b), which ruled out the
possibility that fitness was tied
to toxin dose. Complementation
of *msbA* by overexpression
partially rescued Tae1[WT]
susceptibility in *msbA-KD*
(**Supp. Fig. 8a-c,g**), while *lpxK*
overexpression did not (**Supp.**
Fig. 8d-f,h). Given the
multigenic knockdown in *msbA-*

Figure 4: Resistance to Tae1 in *msbA-KD* is independent of cell wall hydrolysis
a-c) *msbA-KD* **populations have a Tae1-dependent growth advantage.** OD_{600} growth curves
of *msbA-KD* (orange) and *rfp-KD* (black) with CRISPRi induced, overexpressing
(a)*pBAD24::pelB-tae1[WT]* (Tae1[WT]), (b) *pBAD24::pelB-tae1[C30A]* (Tae1[C30A]), or (c) *pBAD24* (empty).
Data shown: average of 3 biological replicates ± s.d. Dotted vertical line indicates plasmid
induction timepoint (at OD_{600}=0.25). **d) The muropeptide composition of *msbA-KD* PG is**
identical to control *rfp-KD*. HPLC chromatograms of muropeptides purified from *msbA-KD*
(orange) and *rfp-KD* (grey) expressing *pBAD24* (empty). *Inset:* major muropeptide species in
Eco include tetrapeptide monomers (M4; r.t. ~10 minutes) and 4,3-crosslinked tetra-tetra dimers
(D44; r.t. ~15.5 minutes). Tae1 digests D44 peptides (black arrow). Data shown: representative
from 3 biological replicates. **e) Tae1[WT] digests PG from both *msbA-KD* and *rfp-KD* PG *in***
***vivo*.** HPLC chromatograms of muropeptides purified from *msbA-KD* (orange) and *rfp-KD* (grey)
expressing *pBAD24::pelB-tae1[WT]* (Tae1[WT]). Black arrow indicates D44 peptide partially digested
by Tae1. Data shown: representative from 3 biological replicates. **f) Tae1 is equally efficient at**
digesting PG in *msbA-KD* and *rfp-KD*. Percent loss of D44 peptide after 60 minutes of
periplasmic Tae1[WT] or Tae1[C30A] expression. Data shown: average of 3 biological replicates (±
s.d.). Statistical test: two-tailed unpaired *t*-test; *p*-value ≤0.05 displayed in bold font.

	Tae1[WT]	Tae1[C30A]	*p*-value (WT vs. C30A)
rfp-KD	-30.77% (±2.66%)	-4.74% (±5.83%)	*0.0022*
msbA-KD	-27.73% (±1.76%)	-3.01% (±1.96%)	*0.000084*
p-value (rfp-KD vs. msbA-KD)	0.1744	0.6511	

359 *lpxK-ycaQ* in *msbA-KD*, these data suggest that *msbA* is a partial determinant of Tae1 susceptibility in the strain.
360
Next, we tested whether *msbA-KD* directly impacts Tae1–PG physical interactions by triggering changes to the
chemical composition of *Eco* PG, which can occur downstream of OM stress[31]. PG remodeling could alter intrinsic
Tae1 susceptibility by changing the relative abundance of targetable peptides in the cell wall. We isolated and
characterized the composition of PG purified from *rfp-KD* and *msbA-KD* by HPLC muropeptide analysis. Both
strains had highly similar and stereotypical *Eco* muropeptide profiles (**Fig. 4d**). PG peptides containing the scissile
bond and structural context for Tae1 recognition (4,3-crosslinked dimers; D44)[8] were found at an approximate 1:1
ratio with another dominant species of muropeptide (tetrapeptide monomers; M4)[49]. Our results suggest that the

PG composition of *msbA-KD* is not modified downstream
of LPS damage, indicating that Tae1 resistance cannot be
explained by biochemical changes to the Tae1:PG
interaction.
372
We tested an alternative hypothesis that resistance may
derive from decreased efficiency in Tae1 hydrolysis. We
reasoned that structural deformations in the *msbA-KD* cell
envelope (**Fig. 3f**) could occlude or delay the accessibility
of PG to Tae1, thus slowing the kinetics of cell wall
degradation and cell lysis. To test this, we monitored the
relative degradation of D44 peptides after Tae1 induction
in *rfp-KD* and *msbA-KD* populations. Empty-vector and
Tae1[C30A] conditions were included as negative controls
(**Fig4d,f; Supp. Fig. 9a**). At 60 minutes of induction (just
prior to lysis in *rfp-KD* populations*)*, we found that D44
peptides were similarly hydrolyzed between strains, with a
32.58% loss in *rfp-KD* and 27.73% of in *msbA-KD* (**Fig.
4e-f**). Thus, Tae1 hydrolyzes *msbA-KD* PG as efficiently
as *rfp-KD* PG. Collectively, these data show that both cell
wall recognition and hydrolysis by Tae1 are unchanged in
msbA-KD, ruling out the possibility that direct changes to
PG are responsible for differential cellular lysis outcomes.
391

392 **PG synthesis is suppressed in *msbA-KD* but sensitive
393 to Tae1 activity**
394

Given that we did not find any effects on direct Tae1–cell
wall interactions in *msbA-KD*, we next explored indirect
resistance mechanisms. The PG sacculus is dynamically
synthesized, edited, and recycled *in vivo* to maintain
mechanical support to the cell during growth and
stress[27,50]. We hypothesized that Tae1 hydrolysis could
also impact PG synthesis activity in *Eco* by generating a
need to replace damaged PG with new substrate. The
ability to repair PG could thus be a valuable determinant
of Tae1 susceptibility. To determine if PG synthesis is
sensitive to Tae1 exposure, we measured the
incorporation of the fluorescent D-amino acid HADA into
rfp-KD cell walls both with and without exogenous Tae1
expression. When normalized against control cells
(*empty*), PG synthesis in *rfp-KD* cells increased by 22% in
response to Tae1[WT] and decreased by 6.5% in response
to Tae1[C30A] (**Fig. 5a; Table 2**). These data show that PG

**Figure 5: PG synthesis is suppressed in *msbA-KD* but
sensitive to Tae1 activity**
a-b) PG synthesis activity is sensitive to Tae1 overexpression.
Single-cell fluorescence intensity measurements for *rfp-KD* (a;
grey) or *msbA-KD* (b; orange) after incorporating the fluorescent D-
amino acid HADA into PG after 60 minutes of overexpressing
pBAD24::pelB-tae1[WT] (Tae1[WT]), *pBAD24::pelB-tae1[C30A]* (Tae1[C30A),
or *pBAD24* (empty), with CRISPRi induced. Data shown: 600 cells
(200 cells x 3 biological replicates), with average ± s.d. Statistical
test: unpaired two-tailed *t*-test; *p*-value ≤0.05 displayed in bold
font.

		% change (intra-strain)	% change (*rfp-KD* norm.)
rfp-KD	Tae1[WT]	22% (±3.6%)	
	Tae1[C30A]	-6.5% (±2.6%)	
	empty	0% (±1.6%)	
msbA-KD	Tae1[WT]	26.5% (±2.5%)	12% (±2.5%)
	Tae1[C30A]	2.82% (±3.2%)	-9% (±3.2%)
	empty	0% (±2.0%)	-11.5% (±2.0%)

**Table 2: PG synthesis activity is sensitive to CRISPRi
and Tae1 overexpression**. Descriptive statistics for
normalized percent change in HADA fluorescence in *rfp-KD* and
msbA-KD as related to **Fig.5** and **Supp. Fig. 10**. Data shown:
average of 600 single-cell measurements ±s.d.

synthesis is stimulated by Tae1 exposure, and this response is dependent on toxin activity.

PG synthesis is also coordinated to other essential processes in *Eco*, and sensitive to their genetic or chemical
perturbations[31,51]. We investigated if *msbA-KD* impacts the dynamic PG synthesis response to Tae1. Tae1[WT]
exposure yielded a 26.5% increase in PG activity in *msbA-KD*, and no significant change in activity with Tae1[C30A]
(**Fig. 5b; Table 2**). These results indicate that PG synthesis is still actively regulated in *msbA-KD* in accordance
with relative Tae1 activity. However, when normalized against baseline *rfp-KD* activity, all PG synthesis
measurements for *msbA-KD* were significantly diminished (**Supp. Fig. 10; Table 2**). This observation indicates
that PG synthesis activity is globally suppressed as a consequence of CRISPRi in *msbA*. Thus, we conclude that
PG dynamism in *Eco* is sensitive to Tae1 hydrolysis of PG, and that *msbA-KD* alters the global capacity for PG
synthesis activity without altering its sensitivity to Tae1. Furthermore, these data suggest a reactive crosstalk
between LPS and PG synthesis activities *in vivo*.

Blocks to growth and protein synthesis accompany Tae1 resistance in *msbA-KD*

Based on its responsiveness to Tae1 exposure, we might hypothesize that *Eco* stimulates PG synthesis to
attempt protection against lysis by Tae1. However, suppressed PG synthesis activity alongside tolerance to
wildtype-levels of PG damage in *msbA-KD* suggested that *msbA-KD* may survive lysis by Tae1 using an
additional strategy to support or even supersede PG integrity. *Eco* can resist lysis upon acute PG stress by
transiently arresting homeostatic
functions like cell division, DNA
replication, and protein synthesis
to prioritize stress responses to
critical damage[52–54]. A recent
study showed that a CRISPRi KD
in *lpxA*, the first enzyme in Lipid A
biosynthesis, triggered hallmark
signs of a dormancy stress
response called the stringent
response[55]. Thus, we
hypothesized that decreased PG
synthesis activity in *msbA-KD*
may be symptomatic of a general,
KD-dependent slow growth
phenotype which could protect
against Tae1 activity by passive
tolerance.

To observe the effects of Tae1
and CRISPRi on cellular growth
and lysis behaviors over time, we
performed timelapse microscopy
of *rfp-KD* and *msbA-KD* cells in
competition with *Pae*. Across all
Pae competitions, *msbA-KD* cells
grew slowly without dividing or
lysing (**Fig. 6a; Supp. Fig. 11a-
b**). By contrast, *rfp-KD* cells grew
and divided rapidly, but lysed
when in competition against *Pae*

Figure 6: Blocks to growth and protein synthesis accompany Tae1 resistance in *msbA-KD*

a) *msbA-KD* cells resist lysis from *Pae[WT]* while growing slowly without dividing. Representative frames from time-course imaging of *rfp-KD* (left column; grey) and *msbA-KD* (right column; grey) co-cultured with *Pae[WT]* (green), with CRISPRi induced. Green foci in *Pae[WT]* indicate aggregates of GFP-labelled ClpV, which signal a H1-T6SS firing event. Red arrow indicates lysed cell. Data shown are merged phase and fluorescent channels. Scale bar: 2µm. **b) Protein synthesis activity is attenuated in *msbA-KD*.** Single-cell fluorescence intensity measurements for *rfp-KD* (grey) or *msbA-KD* (orange) cells after incorporating fluorescently-labelled O-propargyl-puromycin (OPP) into nascent peptides during overexpression of *pBAD24::pelB-tae1[WT]* (Tae1[WT]), *pBAD24::pelB-tae1[C30A]* (Tae1[C30A]), or *pBAD24* (empty), with CRISPRi induced. All data normalized to average OPP signal in *rfp-KD* + empty. Data shown: 100 cells/condition, with average ± s.d. Statistical test: unpaired two-tailed *t*-test; *p*-value ≤0.05 displayed in bold font.

strains with active H1-T6SSs (*Pae[WT]*, *Pae[Δtae1]*) (**Fig. 6a; Supp. Fig. 11a-b**). These data demonstrate that stunted
cell growth and division are additional consequences of CRISPRi in *msbA-KD*. We orthogonally tested the effect
of *msbA-KD* on global cell physiology by measuring nascent protein synthesis activity in *msbA-KD* and *rfp-KD*.
Overall protein synthesis levels were significantly lower in *msbA-KD* relative to *rfp-KD* under all conditions tested
(**Fig. 6a**). From these data we conclude that *msbA-KD* cells exhibit broad changes in cellular physiology that may
underscore their unique ability to survive PG damage by Tae1.

We propose a model in which Tae1 susceptibility *in vivo* is determined at multiple levels of specificity in *Eco*: not
only at the level of local PG damage but also by crosstalk between essential cell envelope pathways and the
general growth state of the cell. As mediated through damage to LPS in *msbA-KD*, we posit that such crosstalk
between essential cell functions can be helpful for slowing reactivity and thus increasing tolerance to acute PG

stress. By the same token, the enmeshment of essential pathways may render fast-growing *Eco* vulnerable to
Tae1 by creating a sudden chain-reaction of imbalances in critical functions which the cell must also resolve
alongside the initial PG damage.
475
476 **DISCUSSION**
477
The species composition of mixed-microbial communities can be driven by competitive strategies that bacteria
use to antagonize their neighbors. However, our understanding of microbial weapons is primarily derived from *in*
vitro studies of their molecular mechanisms. In this study, we wanted to understand how Tae1, a PG-degrading
H1-T6SS effector toxin, specifically aided *Pae* in antagonizing *Eco in vivo*. By combining T6SS-mediated
competition with CRISPRi against essential *Eco* genes, our high-throughput genetic screen was poised to
uncover new molecular details about the interaction between Tae1 and essential functions in recipient cells.
Related studies have successfully identified roles for nonessential genes that contribute to recipient survival
against individual T6SS effectors[33,34]. Our study expands our understanding of intrinsic fitness against T6SS
effectors by demonstrating how essential, homeostatic cell activities can have both direct (PG) and indirect (LPS,
growth) impact on the effector-substrate interaction *in vivo*. We find that Tae1 toxicity is driven not only by its
ability to destroy PG but also by broader physiological and regulatory contexts.
489
Through the lens of LPS perturbation (*msbA-KD*), we discovered that slowing cell growth is associated with
resistance to Tae1-dependent lysis. The protective nature of abject dormancy has been demonstrated for survival
against other cell wall-degrading enzymes, lytic bacteriophages, and antibiotics[56–60]. However, previous work on
interbacterial competition has shown that fast growth protects recipient cells from T6SS by establishing stable
microcolonies more quickly than T6SS can kill the recipient cell type[61,62]. Our study suggests that slow recipient
growth could also offer a fitness advantage against lytic T6SS effectors. Similarly to how dead (unlysed) cells can
physically block T6SS-wielding competitors from progressing in space [63], slow-growing cells could also absorb
T6SS attacks to protect their kin in community settings. A compelling direction for future work could be to
determine if slowing cell growth by an orthogonal mechanism, such as a bonafide stringent response, is sufficient
to recapitulate resistance to Tae1 or other lytic T6SS effectors.
500
A surprising feature of lysis resistance in *msbA-KD* was its tolerance to PG damage by Tae1 alongside additional
damage to its IM and OM. Structural destabilization of the cell envelope commonly renders *Eco* hypersensitive to
lysis[64,65]. However, our observations suggest that integrity of individual envelope components is not always
sufficient to explain cell lysis. Indeed, PG and the OM can work together to bear cellular turgor pressure changes
by sharing the mechanical load across both surfaces[66]. The damaged OM observed in *msbA-KD* could therefore
maintain its turgor-bearing properties to protect cells against lysis when Tae1 hydrolyzes PG. Additionally, the
mechanical integrity of the cell envelope in *msbA-KD* may be fortified by covalently-bound Braun's lipoprotein or
changes to membrane composition which could increase cell envelope stiffness[67,68]. Another unique feature for
msbA-KD is that its LPS damage does not stimulate PG remodeling, unlike other depletion alleles for
LPS biosynthesis affecting transport to the OM[31]. We suggest that this indicates multiple nodes for co-regulation
between PG synthesis and LPS synthesis pathways with distinct phenotypic consequences. In line with this
hypothesis, our screen revealed opposing Tae1 sensitivity phenotypes for KDs of *lptC* (LPS transport to OM;
sensitive) and every other LPS hit from the screen (Lipid A-Kdo$_2$ synthesis/transport; resistant). This observation
invites deeper investigation into the potential for multiple types of LPS and PG crosstalk which may inform the
complex underpinnings of mechanical integrity within the bacterial cell envelope.
516
Another key insight from our study is that PG synthesis is stimulated in response to Tae1, indicative of an active
Eco counterresponse. However, wild-type levels of PG synthesis were coincident with, not counter to, lytic death.
Diminished PG synthesis activity in *msbA-KD* could therefore enable resistance by suppressing a toxic
dysregulation of homeostatic activities. We propose that Tae1 activity leads to *Eco* cell death, in part, by triggering
a futile cycle of Tae1 hydrolysis and PG synthesis that does not resolve in cell wall homeostasis. An exciting
prospect for future studies could involve determining the molecular mechanisms that control PG synthesis
stimulation after Tae1 hydrolysis, including whether Tae1 may also synergize or hijack specific endogenous cell
wall enzymes to amplify its damage to PG[69]. The dynamic regulation of PG features indirect to Tae1's peptide
target, such as the glycan backbone, interpeptide crosslinks (type and amount), and recycling could also intersect
with the toxin's acute function to affect its overall impact on the cell wall.
527
In conclusion, our work highlights how recipient susceptibility in interbacterial competition may be more complex
than direct -substrate interactions alone. Toxins with essential targets not only impact specific molecules but also
a dynamic network of interconnected pathways. T6SSs often encode multiple toxins that antagonize different

essential features[70], including components of the cell envelope and other metabolic pathways. We posit that
T6SSs deploy a cocktail of toxins that can act in coordination to disrupt the network beyond repair, or even
weaponize protective homeostatic mechanisms themselves. This study points to the importance of studying the
role of essential genes in the context of T6SS-mediated bacterial antagonism.

METHODS

Bacterial growth and selection

Escherichia coli strains were cultured in LB or LB-no salt (LBNS) at 37°C with orbital shaking. *Pseudomonas aeruginosa* strains were cultured in LB+ 0.01% Triton at 37°C with orbital shaking. Interbacterial competitions between *Eco* and *Pae*, and all *Eco* assays requiring solid growth, were conducted on LB+agar or LBNS+agar plates at 30°C. Where necessary, bacterial strains and plasmids were selected for growth using the following antibiotics: carbenicillin (Carb; 50 µg/ml) (Grainger), chloramphenicol (Chl; 25 µg/ml) (MP Biomedicals), gentamicin (Gent; 50 µg/ml)(Alfa Aesar), irgasan (Irg; 25 µg/ml) (Sigma-Aldrich), trimethoprim (Trm;15 µg/ml) (Sigma-Aldrich), or kanamycin (Kan; 50 µg/ml.) (VWR).

Eco CRISPRi library construction and use

The *Eco* CRISPRi collection was received in pooled format as a gift from the laboratory of Carol Gross (UCSF). CRISPRi strains were derived from K12 strain BW25113[71] and are each engineered with a chromosomal insertion of *dcas9* (constitutive expression) and a custom sgRNA sequence for inducible dCas9-mediated knockdown of a single gene-of-interest[35]. Transcriptional knockdown is induced with addition of 100µM IPTG ("induced") into growth media, though growth without inductant also results in a mild knockdown phenotype ("basal")[35]. Except where indicated, CRISPRi knockdown is induced in this study. CRISPRi strains *msbA-KD* and *lpxK-KD* were reconstructed from the parent strain for individual use in this study. Reconstructed strains were validated by Sanger sequencing (of the sgRNA and dCas9 chromosomal inserts), qRT-PCR (for knockdown efficiency), and Western blot (for dCas9 expression). See **Table S1** for strain descriptions and **Table S2** for primer sequences used for construction and validation.

Pae strain construction

Pae$^{\Delta tae1}$ (*ΔretSΔpppAΔtae1; clpV-GFP*) and *Pae*^{inactive} (*ΔretSΔpppAΔicmF; clpV-GFP*) strains were constructed from biparental mating of parent strain *Pae*^{WT} (B515: PAO1 *ΔretSΔpppA; clpV-GFP*)[72] with *Eco* SM10 λpir[73] bearing suicide vector pEXG2 cloned with homology to the gene(s) of interest and a spacer sequence for replacement. pEXG2 plasmids were cloned using splice-overlap extension[11]. After mating, transformants were isolated by negative selection on LB-agar + 5% sucrose and confirmed as scarless knockout mutants by colony PCR of the locus of interest. See **Table S1** for strain descriptions and **Table S2** for primer sequences used for construction and validation.

Pooled interbacterial competition screen

Competition assays were performed with overnight *Pae* cultures (*Pae*^{WT}, *Pae*$^{\Delta tae1}$, *Pae*^{inactive}) and pooled *Eco* CRISPRi libraries. Flash-frozen glycerol stocks of *Eco* pools were resuspended in LB, backdiluted to OD$_{600}$=0.25, and recovered for 90 minutes at 37°C with shaking. All cultures were washed twice with fresh LB, then OD$_{600}$-adjusted to 2.0 (for *Pae*) or 1.0 (for *Eco*) in either LB (basal CRISPRi) or LB+100µM IPTG (induced CRISPRi) . An aliquot of each CRISPRi pool was reserved by pelleting and flash-freezing for sequencing-based analysis of strain abundances in the starting population. Media-matched *Pae* and *Eco* were mixed at a 1:1 volumetric ratio, except for *Eco*^{ctrl} populations (for which *Eco* pools were not mixed with *Pae)*. Six, 10µl aliquots of coculture were applied to nitrocellulose membranes (0.2µm, GVS) atop LB-agar (basal CRISPRi) or LB-agar +100µM IPTG (induced CRISPRi) plates to match liquid media conditions. Covering the agar surface with nitrocellulose allows for nutrient transfer from the media to the bacteria, while aiding in bacterial recovery from the surface after competition. Cocultures were dried down to the membrane under flame-sterilization, then incubated at 30°C for 6h. Cocultures were removed from the plate by scalpel-excision of surrounding nitrocellulose and resuspended into 1ml fresh PBS by bead-beating for 45s on a tabletop vortex. The six aliquots per experiment were pooled, centrifuged (2min at 9000xG, RT), and PBS was decanted. Pellets were flash frozen in liquid nitrogen and stored at -80°C.

Sequencing library preparation

Genomic DNA was extracted from frozen bacterial pellets by phenol: chloroform extraction and RNase treatment[74], followed by quantification on a Nanodrop 2000 spectrophotometer (Thermo Scientific). PCR amplification was used to isolate *Eco* sgRNA sequences from mixed genomic DNA and to attach Illumina Truseq index adapters for high-throughput sequencing. Sequencing libraries were purified by gel electrophoresis on 8% TBE gels (Invitrogen Novex), stained with SYBR Gold (Invitrogen) to visualize library bands, and scalpel-excised (200-300bp region) under blue light imaging (Azure Biosystems c600). Excised libraries were gel-extracted and precipitated[75], then resuspended in nuclease-free distilled water (Invitrogen UltraPure). Library concentration was quantified on a Qubit 2.0 fluorimeter (Invitrogen) using the dsDNA high-sensitivity assay, and assayed for purity on a 2100 Bioanalyzer (Agilent) using the high-sensitivity DNA assay. Single-end sequencing was performed on an Illumina NextSeq 500 using a custom sequencing primer and a read length of 75bp. Multiplexed samples were

594   spiked with 5% PhiX Control v3 DNA (Illumina) to account for low diversity among sgRNA sequences. See **Table**
595   **S2** for custom primers used for library preparation and sequencing.
596
Sequencing data analysis
Raw FASTQ files were aligned to the library oligos and counted using ScreenProcessing
(https://github.com/mhorlbeck/ScreenProcessing). Counts were normalized to a total of 20,000,000 reads,
pseudocounts of 1 were added, and $\log_2$ fold change (L2FC) from t0 was calculated for each strain with at least
100 counts at t0. L2FC was further corrected by subtracting the median L2FC of the non-targeting control
sgRNAs from that sample[76]. The L2FC of each sgRNA were averaged across four biological replicates to
calculate the L2FC for that condition. Finally, to account for differences in the number of generations experienced
(growth) in each of the experimental conditions, L2FC values for the Pae^{WT}, $Pae^{\Delta tae1}$, $Pae^{inactive}$ experiments were
corrected by the coefficient of a robust (MM-type) intercept free linear regression between the experimental L2FC
values and the CRISPRi induction-matched (induced/basal) Eco^{ctrl} experiment. See Table S3 for correction
coefficients and corrected L2FC values. Differences between conditions were then calculated for each sgRNA as:
608
609   Diff = (L2FC [*condition*]) – (L2FC Eco^{ctrl})
610
611   Final Diff values are listed in **Table S4** and were used for all further analyses.
612
613   **COG analysis**
614   Gene ontology information was compiled from the NIH Database of Clusters of Orthologous Genes (COGs)
615   (https://www.ncbi.nlm.nih.gov/research/cog) and reported previously[35].
616
617   **Data availability and software**
618   Illumina sequencing data from this study is accessible at the NCBI Sequence Read Archive under accession
619   PRJNA917770. Principal component analysis was performed using R[77] and visualized using ggplot2[78]. All other
620   data visualizations were prepared using GraphPad Prism 9.4.1 (GraphPad Software, San Diego, California USA,
621   www.graphpad.com).
622
Pairwise Interbacterial T6SS competition assay
Competition assays were performed with overnight liquid cultures of *Pae* and *Eco* CRISPRi strains. *Eco* cultures
were backdiluted 1:4 in LB-no salt (LBNS; cite) + 100µM IPTG and grown for 1h at 37°C with shaking to pre-
induce CRISPRi before competition. Strains were washed and mixed in a 1:1 volumetric ratio of *Pae* (OD_{600}=2)
and *Eco* (OD_{600}=1) in LBNS+100µM IPTG. Three, 10µl aliquots of each liquid co-culture applied to nitrocellulose
membranes (0.2µm, GVS) atop LB-agar+100µM IPTG and dried down by flame-sterilization to encourage
interbacterial competition. Cocultures were incubated at 30°C for 6h. For initial *Eco* colony-forming unit
measurements ($CFU_{t=0h}$), 20µl of each liquid co-culture input was serially diluted (10-fold dilutions x 8) in a 96-well
plate (Corning) and plated onto LB-agar + Gent (*Eco*-selective). After the competition, coculture spots were
harvested from the plate by scalpel-excision of the surrounding nitrocellulose, and pooled by resuspension into
1ml fresh PBS by bead-beating for 45s on a tabletop vortex. Resuspensions were serially diluted (10x8) and
plated onto LB+Gent. All serial dilution plates were incubated overnight at 37°C. Dilution plates with
approximately 20-200 colonies-per-plate were counted for *Eco* CFU abundance ($CFU_{t=0h}$, $CFU_{t=6h}$). Fold-change
in *Eco* CFUs was determined by back-calculating CFUs per ml from dilution plates, and then calculating
$CFU_{t=6h}/CFU_{t=0h}$. Experiment was performed for three biological replicates. Statistical test: two-tailed unpaired t-
test.
639
qRT-PCR
Overnight cultures of *Eco* were washed and OD_{600}-corrected to 1.0 in LB or LBNS +/-100µl IPTG. Three, 10µl
aliquots of each culture were applied to nitrocellulose membranes (0.2µm, GVS) atop LB-agar+100µM IPTG or
LBNS-agar+100µM IPTG and dried down by flame-sterilization. After growing 6 hours at 30°C, the spots were
scalpel-excised, pooled, and resuspended into PBS by bead beating, then pelleted for RNA extraction. RNA was
extracted using TRIzol Reagent (Invitrogen) with Max Bacterial Enhancement Reagent (Invitrogen), followed by
treatment with Turbo DNA-free kit (Invitrogen) to remove contaminating DNA. After quantification by Nanodrop
(Thermo Scientific), total RNA was reverse transcribed into cDNA using qScript cDNA Supermix (QuantaBio). A
1:5 dilution of cDNA and custom primers were input into qPCR reactions with PowerUP SYBR Green Master Mix
(Applied Biosystems).qRT-PCR was performed using a QuantStudio 3 Real Time PCR system (ThermoFisher
Scientific) using cycling parameters as defined by the master mix instructions. Fold-change in transcript levels
was calculated using $\Delta\Delta C_t$ analysis, using *rpoD* as a control gene. Three biological and three technical replicates

were used per experiment. Statistical test: two-tailed unpaired t-test. Custom primers for qPCR of *Eco* genes can
be found in Table 3.
654
Cryo-ET imaging
Overnight cultures of *E.coli* strains were diluted in LB 1:100 and grown at 37°C. At OD_{600}=0.2, 150 µM IPTG was
added to the liquid culture to induce CRISPRi knockdown. Bacteria were grown for another 90 min and then flash-
frozen in liquid nitrogen. Cell cultures were mixed with 10 nm protein A gold at 20:1 ratio (Utrecht), then aliquots
of 3 µL mixtures were applied to glow-discharged R2/2, 200 mesh copper Quantifoil grids (Quantifoil Micro Tools).
The sample was blotted for 3 s at 20°C and at 80% humidity. The grids were plunge-frozen in liquid ethane using
Leica EM GP system (Leica Microsystems) and stored in liquid nitrogen. Cryo-ET was performed on a Talos
electron microscope equipped with a Ceta CCD camera (ThermoFisher). Images were taken at magnification
22,000x corresponding to a pixel size of 6.7 Å. Tilt series were collected using SerialEM[79] with a continuous tilt
scheme (−48° to 48°, every 3° increment). The defocus was set to -6 to -8 µm and the cumulative exposure per tilt
series was 150 e⁻/A². Tomograms were reconstructed with the IMOD software package[80].
666
Overexpression plasmid construction and use
Plasmids for periplasmic Tae1 overexpression in *Eco* were constructed using splice-overlap extension cloning of
tae1^WT and *tae1^C30A* coding sequences derived from *P.aeruginosa* (PAO1) into *pBAD24*[48,81]. A *pelB* leader
sequence was fused to *tae1* for localization to the periplasm. Expression from *pBAD24* plasmids transformed into
Eco was induced by addition of 0.125% arabinose (w/v) (Spectrum Chemical) into liquid LBNS media at early log
phase (OD_{600} ~0.25). Overexpression constructs for *msbA* and *lpxK* were constructed by cloning each full-length
gene from *Eco* into the NdeI/HindIII restriction sites of *pSCrhaB2*[82]. Overexpression from *pSCrhaB2* plasmids
transformed into *Eco* was induced by addition of 0.1% rhamnose (w/v) (Thermo Scientific) into liquid media. See
Table S2 for primer sequences used for cloning and PCR validation.
676
Tae1 overexpression lysis assay
Chemically competent *Eco* were transformed with Tae1 overexpression constructs (*pBAD24::tae1^WT*,
pBAD24::tae1^C30A, *pBAD24*) by standard 42°C heat-shock and a 45-minute recovery in LB at 37°C with shaking.
A transformant population was selected overnight in liquid LB+Carb; the more-traditional method of selecting on
solid media was skipped to discourage the formation of Tae1-resistant compensatory mutations. Overnight
transformant cultures were backdiluted to OD_{600}=0.1 in LBNS+Carb +/- 100µM IPTG, then incubated in a Synergy
H1 plate reader (BioTek) at 37°C with shaking (2 technical x 3 biological replicates). OD_{600} reads were taken
every five minutes to generate a growth curve. At OD_{600}=0.25 (early log-phase), Tae1 expression was induced
from *pBAD24* with the addition of 0.125% arabinose to each well, and grown for 500 minutes at 37°C with
shaking. Bacterial growth curves were normalized to blank growth curves (LBNS+Carb, no bacteria), and average
growth curves from all biological and technical replicates were plotted in Prism (GraphPad).
688
689 For *msbA* and *lpxK* complementation assays, *pSCrhaB2* plasmids were transformed alongside *pBAD24* plasmids,
690 and overnight selection was performed in liquid LB+Carb+Trm. The next day, cultures were washed and
691 backdiluted at OD_{600}=0.1 into LBNS+Carb+Trm+0.1% rhamnose. The experiment then proceeded in the plate
692 reader as described above.
693
Western blotting
dCas9 detection: Total protein was extracted from the organic layer of bacterial pellets treated with TRIzol
Reagent (prepared as described in **qRT-PCR**), according to manufacturer's protocol. Protein samples were
diluted to 1mg/ml in PBS + 1x Laemmli denaturing buffer, boiled for 10 minutes then centrifuged at 20,000xg at
RT for 2 minutes. Fifteen µl of supernatant was loaded onto an anyKD MiniPROTEAN gel (BioRad), alongside
ProteinPlus Ladder (BioRad). Gels were run according to manufacturer's protocol in 1x SDS-PAGE running buffer
to separate proteins. Protein was transferred to nitrocellulose (0.2µm; GVS) via semi-dry transfer with a TransBlot
Turbo transfer system (BioRad) and matching transfer buffer (BioRad) under the following conditions: 45 min @
15V, 2.5 Amp. Transfer was validated by Ponceau stain. Blots were blocked for one hour at RT with shaking in
3% milk+TBST. Primary antibody was applied: 1:1000 mouse anti-Cas9 (Abcam ab191468) in TBST, overnight,
at 4C with shaking. Blots were washed four times in TBST. Secondary antibody was applied: 1:5000 anti-mouse
HRP (Advansta R-05071-500) in TBST, for one hour at RT, with shaking. Blots were washed four times in TBST.
Blots were treated with Clarity ECL Western blotting substrate (BioRad) for chemiluminescent detection on an
Azure c400 imager. Visible light images were also taken to visualize protein ladder. Densitometry analysis was
performed in Fiji[83,84]. Statistical test: two-tailed unpaired t-test. Three biological replicates.
Tae1 detection: Chemically competent *Eco* cells were transformed with Tae1 overexpression constructs
(*pBAD24::tae1^WT*, *pBAD24::tae1^C30A*, *pBAD24*) by standard 42°C heat-shock and a 45-minute recovery in LB at

37°C with shaking. A transformant population was selected overnight in liquid LB+Carb. Cultures were
backdiluted to OD_{600}=0.1 in LBNS + Carb +100µM IPTG, then incubated in a Synergy H1 plate reader (BioTek) at
37°C with shaking (2 technical x 3 biological replicates). OD_{600} reads were taken every five minutes to track
population growth. At OD_{600}=0.25, Tae1 expression was induced with the addition of 0.125% arabinose to each
well. Bacteria were grown for 60 minutes with Tae1 induction, before technical replicates were harvested and
pooled. Samples were pelleted by centrifugation and media was decanted before cells were resuspended in PBS
+ 1x Laemmli denaturing buffer. Western blotting protocol then proceeded as above, excepting the use of a
custom rabbit anti-Tae1 primary antibody (1:2500 in TBST) (ThermoFisher) and anti-rabbit HRP secondary
antibody (1:5000 in TBST) (Advansta R-05072-500).
720
Muropeptide analysis
Chemically competent *Eco* cells were transformed with Tae1 overexpression constructs (*pBAD24::tae1^WT^*,
*pBAD24::tae1^C30A^*, *pBAD24*) by standard 42°C heat-shock and a 45-minute recovery in LB at 37°C with shaking.
A transformant population was selected overnight in liquid LB+Carb. Cultures were backdiluted to OD_{600}=0.1 in
LBNS+Carb +100µM IPTG, and grown with shaking. At early log phase (OD_{600}=0.25), 0.125% arabinose was
added to induce *pBAD24* expression. Cells were grown for 60 minutes, then harvested by centrifugation. For PG
purification, cells were boiled in 3% SDS to extract crude PG, then treated with Pronase E (100µg/ml in Tris-HCl
(pH 7.2) + 0.06% NaCl) (VWR Chemicals) for 2 hours at 60C to remove proteins covalently bound to PG.
Mutanolysin digestion (40µg/ml in Tris-HCl (pH 7.2) + 0.06% NaCl) was performed overnight at 37C to solubilize
PG into muropeptides for HPLC analysis. Samples were reduced with sodium borohydride (Fisher Chemical) then
pH-corrected to 3-4 using o-phosphoric acid(Fisher Chemical)[85]. Muropeptides were separated on a 1220 Infinity
II HPLC (Agilent) with UV-visible detection (λ=206nm). Muropeptide separation was achieved over 54 minutes at
0.5 ml/min using a Hypersil ODS C18 column (Thermo Scientific) and a gradient elution from 50mM sodium
phosphate + 0.04% NaN_3 (Buffer A) to 75mM sodium phosphate +15% methanol (Buffer B). Chromatograms
were integrated in ChemStation software (Agilent) to determine peak area, height, and elution time. Experimental
chromatograms were normalized against a chromatogram from a blank run (ddH_2O). Chromatograms were also
internally normalized against the most abundant M4 (monomer muropeptide) peak; this allowed for direct relative
comparisons of peak heights between samples.
739
To calculate the percent change in D44 (4,3-crosslinked dimer) peptides after Tae1 overexpression, the
normalized area under the curve (AUC) for D44 was divided by the total chromatogram area to calculate the
relative D44 peak area for each condition (AUC_{WT}, AUC_{C30A}, AUC_{EV}). Then, within a given strain,
(AUC_{WT}/AUC_{EV})*100 and (AUC_{C30A}/AUC_{EV})*100 were calculated to determine the percent of D44 peak area lost to
Tae1^WT^ or Tae1^C30A^ treatment, relative to EV treatment. Three biological replicates were performed per condition.
Statistical test: two-tailed unpaired t-test.
746
HADA incorporation imaging
Chemically competent cells were transformed with *pBAD24* constructs: (*pBAD24::tae1^WT^*, *pBAD24::tae1^C30A^*,or
pBAD24) and selected with Carb overnight in liquid LB. Transformant cultures were backdiluted to OD_{600}=0.1 in
1ml LBNS+Carb +100µM IPTG, and grown with shaking. At early log phase (OD_{600}=0.25), 0.125% arabinose
added to induce *pBAD24* expression. Cells were grown for 30 minutes, then 250µM HADA added to culture. Cells
were grown an additional 30 minutes, then collected by centrifugation and washed 3x with cold PBS + sodium
citrate (pH 3.0) to block hydrolysis of labelled septal PG[86]. Cells were fixed by treatment with 3% PFA for 15
minutes on ice. Fixed cells were washed 3x in cold PBS, then resuspended in PBS +20% DMSO. Fluorescence
imaging was performed on a Nikon Eclipse Ti2-E inverted microscope equipped with a 100x/1.40 oil-immersion
phase objective and an EMCCD camera (Prime 95B). Fluorescence (DAPI channel) and phase-contrast images
were captured using NIS-Elements AR Viewer 5.20. Images were analyzed for single-cell fluorescence intensity
using MicrobeJ for Fiji[84,87]. 200 cells/sample measured, 3 biological replicates. Statistical test: unpaired t-test.
759
Nascent protein synthesis imaging
Chemically competent cells were transformed with *pBAD24* constructs: (*pBAD24::tae1^WT^*, *pBAD24::tae1^C30A^*,or
pBAD24) and selected with Carb overnight in liquid LB. Cultures were diluted by 1:100 and grown in LBNS+ Carb+
100µM IPTG at 37 °C with shaking. At early log phase (~80 minutes) 0.125% arabinose was added to induce Tae1
expression. After 35 minutes, 13µM O-propargyl-puromycin (OPP) was added to cultures to label new peptide
synthesis before harvesting (Click-iT™ Plus OPP Alexa Fluor™ 488 Protein Synthesis Assay Kit, Invitrogen)[88]. After
labelling, cells were pelleted and fixed in 3.7% formaldehyde in PBS. Cells were permeabilized with 0.3% Triton X-
100 in PBS for 15 min, then labelled for imaging with Click-iT reaction cocktail for 20 min in the dark, washed then
resuspended in PBS. Fluorescence imaging was performed on a Nikon Eclipse Ti2-E inverted microscope equipped
with a 100x/1.40 oil-immersion objective and an EMCCD camera (Prime 95B). The 488-nm laser illumination

fluorescence and phase-contrast images were captured using NIS-Elements AR Viewer 5.20 and analyzed using MicrobeJ software for Fiji[84,87].

Time-lapse imaging of T6SS competitions

Competition microscopy experiments were performed with overnight liquid cultures of *Pae* (LB) and *Eco* CRISPRi strains (LB+Gent+Cam). Cultures were diluted 1:50 in fresh medium and grown for 2h. *Pae* cells were diluted again 1:50 in fresh medium (LB) and grown at 37°C to OD 1.2 – 1.5 (~1 hour). Similarly, *E. coli* strains were diluted 1:100 in fresh medium (LB+150µM IPTG) supplemented with antibiotics (Gent / Cam) and grown at 37°C to OD 1.2 – 1.5 (~1 hour). Then, cultures were washed with LB, resuspended in LB + 150µM IPTG and mixed 2:1 (*Pae:Eco*). 1 µl of the mixed cells was spotted on an agarose pad containing propidium iodide and imaged for 2h at 37°C. A Nikon Ti-E inverted motorized microscope with Perfect Focus System and Plan Apo 1003 Oil Ph3 DM (NA 1.4) objective lens was used to acquired images. If not indicated otherwise, time-lapse series of competitions were acquired at 10 s acquisition frame rate during 120 min. SPECTRA X light engine (Lumencore), ET-GFP (Chroma #49002) and ET-mCherry (Chroma #49008) filter sets were used to excite and filter fluorescence. VisiView software (Visitron Systems, Germany) was used to record images with a sCMOS camera pco.edge 4.2 (PCO, Germany) (pixel size 65 nm). The power output of the SPECTRA X light engine was set to 20% for all excitation wavelengths. GFP, phase-contrast and RFP / propidium iodide (PI) images were acquired with 50-100 ms exposure time. Temperature and humidity were set to 37°C, 95% respectively, using an Okolab T-unit objective heating collar as well as a climate chamber (Okolab). Fiji was used for imaging processing[84]. Acquired time-lapse series were drift-corrected using a custom StackReg based software [89,90].

SUPPLEMENTAL INFORMATION

Table S1. Bacterial strains and plasmids used in this study
Table S2. Primer sequences
Table S3. Corrected L2FC values from screen
Table S4. Final Diff values from screen

ACKNOWLEDGEMENTS

We are grateful to all members of the Chou and Basler labs for their support throughout this project (with special thanks to Atanas Radkov, Krisna Van Dyke, Sebastian Flores, and Eleanor Wang). We thank Carol Gross (UCSF) and members of her lab (Jason Peters, Marco Jost, John Hawkins) for their assistance in adapting their CRISPRi system for our project. We thank Michelle Tan, Rene Sit, and Norma Neff (Chan-Zuckerberg Biohub) for assistance with high-throughput sequencing. We thank Naomi Ziv (UCSF), and DeLaine Larsen and Kari Harrington (UCSF Nikon Imaging Center) for assistance with fluorescence microscopy. We are grateful to KC Huang (Stanford University), Waldemar Vollmer (Newcastle University), and Alessandra Polissi (University of Milan) for fruitful conversations regarding the complex biology of the bacterial cell envelope. We thank Sandra Catania and Lauren Trotta for their generous feedback toward data analysis and manuscript preparation.

This work was funded by: NIH NIAID award T32AI060535 (KLT), a UCSF Moritz-Heyman Discovery Fellowship (KLT), the Swiss National Science Foundation (grant BSSGI0_155778) (MB), a European Research Council consolidator grant (865105 -"AimingT6SS") (MB), the Chan-Zuckerberg Biohub (SC), and the Pew Biomedical Scholars Program (SC).

AUTHOR CONTRIBUTIONS

Kristine L Trotta (Conceptualization, Methodology, Research, Data analysis, Data visualization, Writing, Reviewing)
Beth M Hayes (Conceptualization, Research, Data analysis, Reviewing)
Johannes P Schneider (Methodology, Research, Data analysis, Reviewing)
Jing Wang (Methodology, Research, Data analysis, Reviewing)
Horia Todor (Data analysis, Data visualization, Reviewing)
Patrick Rockefeller Grimes (Research, Reviewing)
Ziyi Zhao (Methodology, Research, Data analysis, Reviewing)
William L Hatleberg (Data visualization, Reviewing)
Melanie R Silvis (Conceptualization, Methodology, Reviewing)
Rachel Kim (Methodology, Reviewing)
Byoung-Mo Koo (Methodology, Reviewing)
Marek Basler (Project administration, Reviewing)
Seemay Chou (Project administration, Funding, Conceptualization, Writing, Reviewing)

830 **COMPETING INTERESTS**

831 The authors declare no competing interests. Seemay Chou is the president and CEO of Arcadia Science.

REFERENCES

(1) Ghoul, M.; Mitri, S. The Ecology and Evolution of Microbial Competition. *Trends Microbiol* **2016**, *24* (10), 833–845. https://doi.org/10.1016/j.tim.2016.06.011.

(2) Granato, E. T.; Meiller-Legrand, T. A.; Foster, K. R. The Evolution and Ecology of Bacterial Warfare. *Current Biology* **2019**, *29* (11), R521–R537. https://doi.org/10.1016/j.cub.2019.04.024.

(3) Boyer, F.; Fichant, G.; Berthod, J.; Vandenbrouck, Y.; Attree, I. Dissecting the Bacterial Type VI Secretion System by a Genome Wide in Silico Analysis: What Can Be Learned from Available Microbial Genomic Resources? *BMC Genomics* **2009**, *10*, 104. https://doi.org/10.1186/1471-2164-10-104.

(4) Basler, M. Type VI Secretion System: Secretion by a Contractile Nanomachine. *Philos Trans R Soc Lond B Biol Sci* **2015**, *370* (1679), 20150021. https://doi.org/10.1098/rstb.2015.0021.

(5) Pukatzki, S.; Ma, A. T.; Sturtevant, D.; Krastins, B.; Sarracino, D.; Nelson, W. C.; Heidelberg, J. F.; Mekalanos, J. J. Identification of a Conserved Bacterial Protein Secretion System in Vibrio Cholerae Using the Dictyostelium Host Model System. *Proc Natl Acad Sci U S A* **2006**, *103* (5), 1528–1533. https://doi.org/10.1073/pnas.0510322103.

(6) Hood, R. D.; Singh, P.; Hsu, F.; Güvener, T.; Carl, M. A.; Trinidad, R. R. S.; Silverman, J. M.; Ohlson, B. B.; Hicks, K. G.; Plemel, R. L.; Li, M.; Schwarz, S.; Wang, W. Y.; Merz, A. J.; Goodlett, D. R.; Mougous, J. D. A Type VI Secretion System of Pseudomonas Aeruginosa Targets a Toxin to Bacteria. *Cell Host Microbe* **2010**, *7* (1), 25–37. https://doi.org/10.1016/j.chom.2009.12.007.

(7) Mougous, J. D.; Cuff, M. E.; Raunser, S.; Shen, A.; Zhou, M.; Gifford, C. A.; Goodman, A. L.; Joachimiak, G.; Ordoñez, C. L.; Lory, S.; Walz, T.; Joachimiak, A.; Mekalanos, J. J. A Virulence Locus of Pseudomonas Aeruginosa Encodes a Protein Secretion Apparatus. *Science* **2006**, *312* (5779), 1526–1530. https://doi.org/10.1126/science.1128393.

(8) Chou, S.; Bui, N. K.; Russell, A. B.; Lexa, K. W.; Gardiner, T. E.; LeRoux, M.; Vollmer, W.; Mougous, J. D. Structure of a Peptidoglycan Amidase Effector Targeted to Gram-Negative Bacteria by the Type VI Secretion System. *Cell Reports* **2012**, *1* (6), 656–664. https://doi.org/10.1016/j.celrep.2012.05.016.

(9) Ho, B. T.; Basler, M.; Mekalanos, J. J. Type 6 Secretion System-Mediated Immunity to Type 4 Secretion System-Mediated Horizontal Gene Transfer. *Science* **2013**, *342* (6155), 250–253. https://doi.org/10.1126/science.1243745.

(10) LeRoux, M.; Kirkpatrick, R. L.; Montauti, E. I.; Tran, B. Q.; Peterson, S. B.; Harding, B. N.; Whitney, J. C.; Russell, A. B.; Traxler, B.; Goo, Y. A.; Goodlett, D. R.; Wiggins, P. A.; Mougous, J. D. Kin Cell Lysis Is a Danger Signal That Activates Antibacterial Pathways of Pseudomonas Aeruginosa. *eLife 4*. https://doi.org/10.7554/eLife.05701.

(11) Russell, A. B.; Hood, R. D.; Bui, N. K.; LeRoux, M.; Vollmer, W.; Mougous, J. D. Type VI Secretion Delivers Bacteriolytic Effectors to Target Cells. *Nature* **2011**, *475* (7356), 343–347. https://doi.org/10.1038/nature10244.

(12) Whitney, J. C.; Chou, S.; Russell, A. B.; Biboy, J.; Gardiner, T. E.; Ferrin, M. A.; Brittnacher, M.; Vollmer, W.; Mougous, J. D. Identification, Structure, and Function of a Novel Type VI Secretion Peptidoglycan Glycoside Hydrolase Effector-Immunity Pair *. *Journal of Biological Chemistry* **2013**, *288* (37), 26616–26624. https://doi.org/10.1074/jbc.M113.488320.

(13) Whitney, J. C.; Quentin, D.; Sawai, S.; LeRoux, M.; Harding, B. N.; Ledvina, H. E.; Tran, B. Q.; Robinson, H.; Goo, Y. A.; Goodlett, D. R.; Raunser, S.; Mougous, J. D. An Interbacterial NAD(P)+ Glycohydrolase Toxin Requires Elongation Factor Tu for Delivery to Target Cells. *Cell* **2015**, *163* (3), 607–619. https://doi.org/10.1016/j.cell.2015.09.027.

(14) LaCourse, K. D.; Peterson, S. B.; Kulasekara, H. D.; Radey, M. C.; Kim, J.; Mougous, J. D. Conditional Toxicity and Synergy Drive Diversity among Antibacterial Effectors. *Nat Microbiol* **2018**, *3* (4), 440–446. https://doi.org/10.1038/s41564-018-0113-y.

(15) Pissaridou, P.; Allsopp, L. P.; Wettstadt, S.; Howard, S. A.; Mavridou, D. A. I.; Filloux, A. The Pseudomonas Aeruginosa T6SS-VgrG1b Spike Is Topped by a PAAR Protein Eliciting DNA Damage to Bacterial Competitors. *Proc Natl Acad Sci U S A* **2018**, *115* (49), 12519–12524. https://doi.org/10.1073/pnas.1814181115.

(16) Höltje, J.-V. Growth of the Stress-Bearing and Shape-Maintaining Murein Sacculus of Escherichia Coli. *Microbiol Mol Biol Rev* **1998**, *62* (1), 181–203.

(17) Pazos, M.; Peters, K. Peptidoglycan. In *Bacterial Cell Walls and Membranes*; Kuhn, A., Ed.; Subcellular Biochemistry; Springer International Publishing: Cham, 2019; pp 127–168. https://doi.org/10.1007/978-3-030-18768-2_5.

(18) Russell, A. B.; Singh, P.; Brittnacher, M.; Bui, N. K.; Hood, R. D.; Carl, M. A.; Agnello, D. M.; Schwarz, S.; Goodlett, D. R.; Vollmer, W.; Mougous, J. D. A Widespread Bacterial Type VI Secretion Effector Superfamily Identified Using a Heuristic Approach. *Cell Host & Microbe* **2012**, *11* (5), 538–549. https://doi.org/10.1016/j.chom.2012.04.007.

(19) Shang, G.; Liu, X.; Lu, D.; Zhang, J.; Li, N.; Zhu, C.; Liu, S.; Yu, Q.; Zhao, Y.; Zhang, H.; Hu, J.; Cang, H.; Xu, S.; Gu, L. Structural Insight into How *Pseudomonas Aeruginosa* Peptidoglycanhydrolase Tse1 and Its Immunity Protein Tsi1 Function. *Biochemical Journal* **2012**, *448* (2), 201–211. https://doi.org/10.1042/BJ20120668.

(20) Ding, J.; Wang, W.; Feng, H.; Zhang, Y.; Wang, D.-C. Structural Insights into the Pseudomonas Aeruginosa Type VI Virulence Effector Tse1 Bacteriolysis and Self-Protection Mechanisms. *J Biol Chem* **2012**, *287* (32), 26911–26920. https://doi.org/10.1074/jbc.M112.368043.

(21) Hersch, S. J.; Watanabe, N.; Stietz, M. S.; Manera, K.; Kamal, F.; Burkinshaw, B.; Lam, L.; Pun, A.; Li, M.; Savchenko, A.; Dong, T. G. Envelope Stress Responses Defend against Type Six Secretion System Attacks Independently of Immunity Proteins. *Nat Microbiol* **2020**, *5* (5), 706–714. https://doi.org/10.1038/s41564-020-0672-6.

(22) Kamal, F.; Liang, X.; Manera, K.; Pei, T.-T.; Kim, H.; Lam, L. G.; Pun, A.; Hersch, S. J.; Dong, T. G. Differential Cellular Response to Translocated Toxic Effectors and Physical Penetration by the Type VI Secretion System. *Cell Reports* **2020**, *31* (11), 107766. https://doi.org/10.1016/j.celrep.2020.107766.

(23) Toska, J.; Ho, B. T.; Mekalanos, J. J. Exopolysaccharide Protects Vibrio Cholerae from Exogenous Attacks by the Type 6 Secretion System. *Proceedings of the National Academy of Sciences* **2018**, *115* (31), 7997–8002. https://doi.org/10.1073/pnas.1808469115.

(24) Pérez-Lorente, A. I.; Molina-Santiago, C.; de Vicente, A.; Romero, D. *Sporulation Activated via σ^W Protects* Bacillus *from a Tse1 Peptidoglycan Hydrolase T6SS Effector*; preprint; Microbiology, 2022. https://doi.org/10.1101/2022.02.23.481616.

(25) Dong, T. G.; Dong, S.; Catalano, C.; Moore, R.; Liang, X.; Mekalanos, J. J. Generation of Reactive Oxygen Species by Lethal Attacks from Competing Microbes. *Proc Natl Acad Sci U S A* **2015**, *112* (7), 2181–2186. https://doi.org/10.1073/pnas.1425007112.

(26) Crisan, C. V.; Nichols, H. L.; Wiesenfeld, S.; Steinbach, G.; Yunker, P. J.; Hammer, B. K. Glucose Confers Protection to Escherichia Coli against Contact Killing by Vibrio Cholerae. *Sci Rep* **2021**, *11*, 2935. https://doi.org/10.1038/s41598-021-81813-4.

(27) Typas, A.; Banzhaf, M.; Gross, C. A.; Vollmer, W. From the Regulation of Peptidoglycan Synthesis to Bacterial Growth and Morphology. *Nat Rev Microbiol* **2011**, *10* (2), 123–136. https://doi.org/10.1038/nrmicro2677.

(28) Yang, D. C.; Tan, K.; Joachimiak, A.; Bernhardt, T. G. A Conformational Switch Controls Cell Wall-Remodelling Enzymes Required for Bacterial Cell Division. *Mol Microbiol* **2012**, *85* (4), 768–781. https://doi.org/10.1111/j.1365-2958.2012.08138.x.

(29) Delhaye, A.; Collet, J.-F.; Laloux, G. Fine-Tuning of the Cpx Envelope Stress Response Is Required for Cell Wall Homeostasis in *Escherichia Coli*. *mBio* **2016**, *7* (1), e00047-16. https://doi.org/10.1128/mBio.00047-16.

(30) Peters, K.; Kannan, S.; Rao, V. A.; Biboy, J.; Vollmer, D.; Erickson, S. W.; Lewis, R. J.; Young, K. D.; Vollmer, W. The Redundancy of Peptidoglycan Carboxypeptidases Ensures Robust Cell Shape Maintenance in Escherichia Coli. *mBio* **2016**, *7* (3), e00819-16. https://doi.org/10.1128/mBio.00819-16.

(31) Morè, N.; Martorana, A. M.; Biboy, J.; Otten, C.; Winkle, M.; Serrano, C. K. G.; Montón Silva, A.; Atkinson, L.; Yau, H.; Breukink, E.; den Blaauwen, T.; Vollmer, W.; Polissi, A. Peptidoglycan Remodeling Enables Escherichia Coli To Survive Severe Outer Membrane Assembly Defect. *mBio* **2019**, *10* (1), e02729-18. https://doi.org/10.1128/mBio.02729-18.

(32) Mueller, E. A.; Egan, A. J.; Breukink, E.; Vollmer, W.; Levin, P. A. Plasticity of Escherichia Coli Cell Wall Metabolism Promotes Fitness and Antibiotic Resistance across Environmental Conditions. *Elife* **2019**, *8*, e40754. https://doi.org/10.7554/eLife.40754.

(33) Lin, H.-H.; Yu, M.; Sriramoju, M. K.; Hsu, S.-T. D.; Liu, C.-T.; Lai, E.-M. A High-Throughput Interbacterial Competition Screen Identifies ClpAP in Enhancing Recipient Susceptibility to Type VI Secretion System-Mediated Attack by Agrobacterium Tumefaciens. *Front Microbiol* **2020**, *10*, 3077. https://doi.org/10.3389/fmicb.2019.03077.

(34) Hersch, S. J.; Sejuty, R. T.; Manera, K.; Dong, T. G. *High Throughput Identification of Genes Conferring Resistance or Sensitivity to Toxic Effectors Delivered by the Type VI Secretion System*; preprint; Microbiology, 2021. https://doi.org/10.1101/2021.10.06.463450.

(35) Silvis, M. R.; Rajendram, M.; Shi, H.; Osadnik, H.; Gray, A. N.; Cesar, S.; Peters, J. M.; Hearne, C. C.; Kumar, P.; Todor, H.; Huang, K. C.; Gross, C. A. Morphological and Transcriptional Responses to CRISPRi Knockdown of Essential Genes in Escherichia Coli. *mBio* **2021**, *12* (5), e02561-21. https://doi.org/10.1128/mBio.02561-21.

(36) Kampmann, M.; Bassik, M. C.; Weissman, J. S. Integrated Platform for Genome-Wide Screening and Construction of High-Density Genetic Interaction Maps in Mammalian Cells. *PNAS* **2013**, *110* (25), E2317–E2326. https://doi.org/10.1073/pnas.1307002110.

(37) Horlbeck, M. A.; Gilbert, L. A.; Villalta, J. E.; Adamson, B.; Pak, R. A.; Chen, Y.; Fields, A. P.; Park, C. Y.;
Corn, J. E.; Kampmann, M.; Weissman, J. S. Compact and Highly Active Next-Generation Libraries for
CRISPR-Mediated Gene Repression and Activation. *Elife* **2016**, *5*. https://doi.org/10.7554/eLife.19760.
(38) Silhavy, T. J.; Kahne, D.; Walker, S. The Bacterial Cell Envelope. *Cold Spring Harb Perspect Biol* **2010**, *2* (5),
a000414. https://doi.org/10.1101/cshperspect.a000414.
(39) Klein, G.; Raina, S. Regulated Control of the Assembly and Diversity of LPS by Noncoding SRNAs. *Biomed*
Res Int **2015**, *2015*. https://doi.org/10.1155/2015/153561.
(40) Bertani, B.; Ruiz, N. Function and Biogenesis of Lipopolysaccharides. *EcoSal Plus* **2018**, *8* (1).
https://doi.org/10.1128/ecosalplus.ESP-0001-2018.
(41) Klein, G.; Raina, S. Regulated Assembly of LPS, Its Structural Alterations and Cellular Response to LPS
Defects. *Int J Mol Sci* **2019**, *20* (2), 356. https://doi.org/10.3390/ijms20020356.
(42) Guest, R. L.; Rutherford, S. T.; Silhavy, T. J. Border Control: Regulating LPS Biogenesis. *Trends in*
Microbiology **2021**, *29* (4), 334–345. https://doi.org/10.1016/j.tim.2020.09.008.
(43) Garrett, T. A.; Kadrmas, J. L.; Raetz, C. R. Identification of the Gene Encoding the Escherichia Coli Lipid A 4'-
Kinase. *J Biol Chem* **1997**, *272* (35), 21855–21864. https://doi.org/10.1074/jbc.272.35.21855.
(44) Garrett, T. A.; Que, N. L. S.; Raetz, C. R. H. Accumulation of a Lipid A Precursor Lacking the 4'-Phosphate
Following Inactivation of the Escherichia Coli LpxKGene *. *Journal of Biological Chemistry* **1998**, *273* (20),
12457–12465. https://doi.org/10.1074/jbc.273.20.12457.
(45) Karow, M.; Georgopoulos, C. The Essential Escherichia Coli MsbA Gene, a Multicopy Suppressor of Null
Mutations in the HtrB Gene, Is Related to the Universally Conserved Family of ATP-Dependent
Translocators. *Mol Microbiol* **1993**, *7* (1), 69–79. https://doi.org/10.1111/j.1365-2958.1993.tb01098.x.
(46) Polissi, A.; Georgopoulos, C. Mutational Analysis and Properties of the MsbA Gene of Escherichia Coli,
Coding for an Essential ABC Family Transporter. *Molecular Microbiology* **1996**, *20* (6), 1221–1233.
https://doi.org/10.1111/j.1365-2958.1996.tb02642.x.
(47) Alexander, M. K.; Miu, A.; Oh, A.; Reichelt, M.; Ho, H.; Chalouni, C.; Labadie, S.; Wang, L.; Liang, J.;
Nickerson, N. N.; Hu, H.; Yu, L.; Du, M.; Yan, D.; Park, S.; Kim, J.; Xu, M.; Sellers, B. D.; Purkey, H. E.;
Skelton, N. J.; Koehler, M. F. T.; Payandeh, J.; Verma, V.; Xu, Y.; Koth, C. M.; Nishiyama, M. Disrupting
Gram-Negative Bacterial Outer Membrane Biosynthesis through Inhibition of the Lipopolysaccharide
Transporter MsbA. *Antimicrob Agents Chemother* **2018**, *62* (11), e01142-18.
https://doi.org/10.1128/AAC.01142-18.
(48) Radkov, A.; Sapiro, A. L.; Flores, S.; Henderson, C.; Saunders, H.; Kim, R.; Massa, S.; Thompson, S.;
Mateusiak, C.; Biboy, J.; Zhao, Z.; Starita, L. M.; Hatleberg, W. L.; Vollmer, W.; Russell, A. B.; Simorre, J.-
P.; Anthony-Cahill, S.; Brzovic, P.; Hayes, B.; Chou, S. Antibacterial Potency of Type VI Amidase Effector
Toxins Is Dependent on Substrate Topology and Cellular Context. *eLife* **2022**, *11*, e79796.
https://doi.org/10.7554/eLife.79796.
(49) Desmarais, S. M.; De Pedro, M. A.; Cava, F.; Huang, K. C. Peptidoglycan at Its Peaks: How
Chromatographic Analyses Can Reveal Bacterial Cell Wall Structure and Assembly. *Molecular Microbiology*
2013, *89* (1), 1–13. https://doi.org/10.1111/mmi.12266.
(50) Egan, A. J. F.; Errington, J.; Vollmer, W. Regulation of Peptidoglycan Synthesis and Remodelling. *Nat Rev*
Microbiol **2020**, *18* (8), 446–460. https://doi.org/10.1038/s41579-020-0366-3.
(51) Egan, A. J. F.; Cleverley, R. M.; Peters, K.; Lewis, R. J.; Vollmer, W. Regulation of Bacterial Cell Wall Growth.
FEBS J **2017**, *284* (6), 851–867. https://doi.org/10.1111/febs.13959.
(52) Scherrer, R.; Moyed, H. S. Conditional Impairment of Cell Division and Altered Lethality in HipA Mutants of
Escherichia Coli K-12. *J Bacteriol* **1988**, *170* (8), 3321–3326.
(53) Kwan, B. W.; Valenta, J. A.; Benedik, M. J.; Wood, T. K. Arrested Protein Synthesis Increases Persister-Like
Cell Formation. *Antimicrob Agents Chemother* **2013**, *57* (3), 1468–1473. https://doi.org/10.1128/AAC.02135-
12.
(54) Liu, S.; Wu, N.; Zhang, S.; Yuan, Y.; Zhang, W.; Zhang, Y. Variable Persister Gene Interactions with
(p)PpGpp for Persister Formation in Escherichia Coli. *Frontiers in Microbiology* **2017**, *8*.
(55) Roghanian, M.; Semsey, S.; Løbner-Olesen, A.; Jalalvand, F. (P)PpGpp-Mediated Stress Response Induced
by Defects in Outer Membrane Biogenesis and ATP Production Promotes Survival in Escherichia Coli. *Sci*
Rep **2019**, *9* (1), 2934. https://doi.org/10.1038/s41598-019-39371-3.
(56) Łoś, M.; Golec, P.; Łoś, J. M.; Węglewska-Jurkiewicz, A.; Czyż, A.; Węgrzyn, A.; Węgrzyn, G.; Neubauer, P.
Effective Inhibition of Lytic Development of Bacteriophages λ, P1 and T4 by Starvation of Their Host,
Escherichia Coli. *BMC Biotechnology* **2007**, *7* (1), 13. https://doi.org/10.1186/1472-6750-7-13.
(57) Pearl, S.; Gabay, C.; Kishony, R.; Oppenheim, A.; Balaban, N. Q. Nongenetic Individuality in the Host–Phage
Interaction. *PLoS Biol* **2008**, *6* (5), e120. https://doi.org/10.1371/journal.pbio.0060120.

(58) Meeske, A. J.; Nakandakari-Higa, S.; Marraffini, L. A. Cas13-Induced Cellular Dormancy Prevents the Rise of CRISPR-Resistant Bacteriophage. *Nature* **2019**, *570* (7760), 241–245. https://doi.org/10.1038/s41586-019-1257-5.

(59) Wiuff, C.; Zappala, R. M.; Regoes, R. R.; Garner, K. N.; Baquero, F.; Levin, B. R. Phenotypic Tolerance: Antibiotic Enrichment of Noninherited Resistance in Bacterial Populations. *Antimicrob Agents Chemother* **2005**, *49* (4), 1483–1494. https://doi.org/10.1128/AAC.49.4.1483-1494.2005.

(60) Lee, A. J.; Wang, S.; Meredith, H. R.; Zhuang, B.; Dai, Z.; You, L. Robust, Linear Correlations between Growth Rates and β-Lactam–Mediated Lysis Rates. *Proceedings of the National Academy of Sciences* **2018**, *115* (16), 4069–4074. https://doi.org/10.1073/pnas.1719504115.

(61) Wilmoth, J. L.; Doak, P. W.; Timm, A.; Halsted, M.; Anderson, J. D.; Ginovart, M.; Prats, C.; Portell, X.; Retterer, S. T.; Fuentes-Cabrera, M. A Microfluidics and Agent-Based Modeling Framework for Investigating Spatial Organization in Bacterial Colonies: The Case of Pseudomonas Aeruginosa and H1-Type VI Secretion Interactions. *Front Microbiol* **2018**, *9*, 33. https://doi.org/10.3389/fmicb.2018.00033.

(62) Borenstein, D. B.; Ringel, P.; Basler, M.; Wingreen, N. S. Established Microbial Colonies Can Survive Type VI Secretion Assault. *PLOS Computational Biology* **2015**, *11* (10), e1004520. https://doi.org/10.1371/journal.pcbi.1004520.

(63) Smith, W. P. J.; Vettiger, A.; Winter, J.; Ryser, T.; Comstock, L. E.; Basler, M.; Foster, K. R. The Evolution of the Type VI Secretion System as a Disintegration Weapon. *PLOS Biology* **2020**, *18* (5), e3000720. https://doi.org/10.1371/journal.pbio.3000720.

(64) Nichols, R. J.; Sen, S.; Choo, Y. J.; Beltrao, P.; Zietek, M.; Chaba, R.; Lee, S.; Kazmierczak, K. M.; Lee, K. J.; Wong, A.; Shales, M.; Lovett, S.; Winkler, M. E.; Krogan, N. J.; Typas, A.; Gross, C. A. Phenotypic Landscape of a Bacterial Cell. *Cell* **2011**, *144* (1), 143–156. https://doi.org/10.1016/j.cell.2010.11.052.

(65) Paradis-Bleau, C.; Kritikos, G.; Orlova, K.; Typas, A.; Bernhardt, T. G. A Genome-Wide Screen for Bacterial Envelope Biogenesis Mutants Identifies a Novel Factor Involved in Cell Wall Precursor Metabolism. *PLOS Genetics* **2014**, *10* (1), e1004056. https://doi.org/10.1371/journal.pgen.1004056.

(66) Rojas, E. R.; Billings, G.; Odermatt, P. D.; Auer, G. K.; Zhu, L.; Miguel, A.; Chang, F.; Weibel, D. B.; Theriot, J. A.; Huang, K. C. The Outer Membrane Is an Essential Load-Bearing Element in Gram-Negative Bacteria. *Nature* **2018**, *559* (7715), 617–621. https://doi.org/10.1038/s41586-018-0344-3.

(67) Mathelié-Guinlet, M.; Asmar, A. T.; Collet, J.-F.; Dufrêne, Y. F. Lipoprotein Lpp Regulates the Mechanical Properties of the E. Coli Cell Envelope. *Nat Commun* **2020**, *11*, 1789. https://doi.org/10.1038/s41467-020-15489-1.

(68) Effects of Membrane Lipid Composition and Antibacterial Drugs on the Rigidity of Escherichia Coli: Different Contributions of Various Bacterial Substructures. https://doi.org/10.1002/sca.21243.

(69) Singh, S. K.; SaiSree, L.; Amrutha, R. N.; Reddy, M. Three Redundant Murein Endopeptidases Catalyse an Essential Cleavage Step in Peptidoglycan Synthesis of Escherichia Coli K12. *Molecular Microbiology* **2012**, *86* (5), 1036–1051. https://doi.org/10.1111/mmi.12058.

(70) Jurėnas, D.; Journet, L. Activity, Delivery, and Diversity of Type VI Secretion Effectors. *Molecular Microbiology* **2021**, *115* (3), 383–394. https://doi.org/10.1111/mmi.14648.

(71) Baba, T.; Ara, T.; Hasegawa, M.; Takai, Y.; Okumura, Y.; Baba, M.; Datsenko, K. A.; Tomita, M.; Wanner, B. L.; Mori, H. Construction of Escherichia Coli K-12 in-Frame, Single-Gene Knockout Mutants: The Keio Collection. *Molecular Systems Biology* **2006**, *2* (1), 2006.0008. https://doi.org/10.1038/msb4100050.

(72) Vettiger, A.; Basler, M. Type VI Secretion System Substrates Are Transferred and Reused among Sister Cells. *Cell* **2016**, *167* (1), 99-110.e12. https://doi.org/10.1016/j.cell.2016.08.023.

(73) Simon, R.; Priefer, U.; Pühler, A. A Broad Host Range Mobilization System for In Vivo Genetic Engineering: Transposon Mutagenesis in Gram Negative Bacteria. *Nat Biotechnol* **1983**, *1* (9), 784–791. https://doi.org/10.1038/nbt1183-784.

(74) He, F. E. Coli Genomic DNA Extraction. *Bio-101* **2011**, *e97*. https://doi.org/10.21769/BioProtoc.97.

(75) Churchman, L. S.; Weissman, J. S. Nascent Transcript Sequencing Visualizes Transcription at Nucleotide Resolution. *Nature* **2011**, *469* (7330), 368–373. https://doi.org/10.1038/nature09652.

(76) Hawkins, J. S.; Silvis, M. R.; Koo, B.-M.; Peters, J. M.; Osadnik, H.; Jost, M.; Hearne, C. C.; Weissman, J. S.; Todor, H.; Gross, C. A. Mismatch-CRISPRi Reveals the Co-Varying Expression-Fitness Relationships of Essential Genes in Escherichia Coli and Bacillus Subtilis. *Cell Syst* **2020**, *11* (5), 523-535.e9. https://doi.org/10.1016/j.cels.2020.09.009.

(77) R: A Language and Environment for Statistical Computing. https://www.R-project.org/.

(78) Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*; Springer-Verlag New York.

(79) Mastronarde, D. N. Automated Electron Microscope Tomography Using Robust Prediction of Specimen Movements. *Journal of Structural Biology* **2005**, *152* (1), 36–51. https://doi.org/10.1016/j.jsb.2005.07.007.

(80) Kremer, J. R.; Mastronarde, D. N.; McIntosh, J. R. Computer Visualization of Three-Dimensional Image Data Using IMOD. *J Struct Biol* **1996**, *116* (1), 71–76. https://doi.org/10.1006/jsbi.1996.0013.

(81) Guzman, L. M.; Belin, D.; Carson, M. J.; Beckwith, J. Tight Regulation, Modulation, and High-Level
Expression by Vectors Containing the Arabinose PBAD Promoter. *Journal of Bacteriology* **1995**, *177* (14),
4121–4130. https://doi.org/10.1128/jb.177.14.4121-4130.1995.
(82) Cardona, S. T.; Valvano, M. A. An Expression Vector Containing a Rhamnose-Inducible Promoter Provides
Tightly Regulated Gene Expression in Burkholderia Cenocepacia. *Plasmid* **2005**, *54* (3), 219–228.
https://doi.org/10.1016/j.plasmid.2005.03.004.
(83) Stael, S.; Miller, L. P.; Fernández-Fernández, Á. D.; Van Breusegem, F. Detection of Damage-Activated
Metacaspase ActivityActivitiesby Western Blot in Plants. In *Plant Proteases and Plant Cell Death: Methods
and Protocols*; Klemenčič, M., Stael, S., Huesgen, P. F., Eds.; Methods in Molecular Biology; Springer US:
New York, NY, 2022; pp 127–137. https://doi.org/10.1007/978-1-0716-2079-3_11.
(84) Schindelin, J.; Arganda-Carreras, I.; Frise, E.; Kaynig, V.; Longair, M.; Pietzsch, T.; Preibisch, S.; Rueden, C.;
Saalfeld, S.; Schmid, B.; Tinevez, J.-Y.; White, D. J.; Hartenstein, V.; Eliceiri, K.; Tomancak, P.; Cardona, A.
Fiji: An Open-Source Platform for Biological-Image Analysis. *Nat Methods* **2012**, *9* (7), 676–682.
https://doi.org/10.1038/nmeth.2019.
(85) Desmarais, S. M.; Cava, F.; de Pedro, M. A.; Huang, K. C. Isolation and Preparation of Bacterial Cell Walls
for Compositional Analysis by Ultra Performance Liquid Chromatography. *J Vis Exp* **2014**, No. 83, 51183.
https://doi.org/10.3791/51183.
(86) Peters, K.; Pazos, M.; VanNieuwenhze, M. S.; Vollmer, W. Optimized Protocol for the Incorporation of FDAA
(HADA Labeling) for in Situ Labeling of Peptidoglycan. *Bio Protoc* **2019**, *9* (15), e3316.
https://doi.org/10.21769/BioProtoc.3316.
(87) Ducret, A.; Quardokus, E. M.; Brun, Y. V. MicrobeJ, a Tool for High Throughput Bacterial Cell Detection and
Quantitative Analysis. *Nat Microbiol* **2016**, *1* (7), 1–7. https://doi.org/10.1038/nmicrobiol.2016.77.
(88) Diez, S.; Ryu, J.; Caban, K.; Gonzalez, R. L.; Dworkin, J. The Alarmones (p)PpGpp Directly Regulate
Translation Initiation during Entry into Quiescence. *Proceedings of the National Academy of Sciences* **2020**,
117 (27), 15565–15572. https://doi.org/10.1073/pnas.1920013117.
(89) Ringel, P. D.; Hu, D.; Basler, M. The Role of Type VI Secretion System Effectors in Target Cell Lysis and
Subsequent Horizontal Gene Transfer. *Cell Rep* **2017**, *21* (13), 3927–3940.
https://doi.org/10.1016/j.celrep.2017.12.020.
(90) Thevenaz, P.; Ruttimann, U. E.; Unser, M. A Pyramid Approach to Subpixel Registration Based on Intensity.
IEEE Transactions on Image Processing **1998**, *7* (1), 27–41. https://doi.org/10.1109/83.650848.

SUPPLEMENTAL FIGURES

 Supplement 1: CRISPRi conditionally knocks down transcription across hundreds of *Eco* gene targets
 a) CRISPRi induction produces mild transcriptional knockdown of endogenous *rfp* (11.7-fold decrease) in *Eco*. qRT-PCR
 measurement of relative *rfp* RNA expression in *Eco* strain SC363 after 6 hours of growth on solid LB media with basal or induced CRISPRi.
 Data shown: 3 biological replicates with mean ± s.d. Statistical test: unpaired two-tailed *t*-test. **b) CRISPRi targets *Eco* genes that**
 collectively represent 21 clusters of orthogonal genes (COGs). CRISPRi target genes (*n*=596) were binned by their NCBI COG functional
 assignment. The relative representation of each COG in the strain collection is displayed as a percent of all COGs. Some genes are
 represented by multiple COGs, resulting in a greater number of COGs (*n*=624) than target genes. Non-targeting negative controls ("control",
 n=50) genes without COG assignments ("none", *n*=34), and genes coding for non-coding RNAs ("ncRNA", *n*=7) were also binned.

1107
Supplement 2: Non-targeting CRISPRi induction has little effect on *Eco* fitness in T6SS competition
a) CRISPRi induction does not disrupt T6SS- and Tae1-dependent targeting of *Eco* by *Pae*. Interbacterial competition between *Pae*
(*Pae^WT^*, *Pae^Δtae1^*, *Pae^inactive^*) and an *Eco* negative-control KD strain (*rfp-KD*), with induced or basal CRISPRi. Data shown: mean fold-change (±
geometric s.d.) of *rfp-KD* colony forming units (CFUs) after six hours of competition against *Pae*. Statistical test: unpaired two-tailed *t*-test; *p*-
value ≤0.05 displayed in bold font.

Supplement 3: CRISPRi library fitness in T6SS screen is reproducible across biological replicates
a) CRISPRi library fitness in T6SS screen is reproducible across biological replicates. Replica plots showing the uncorrected L2FC
values for each *Eco* CRISPRi strain after competition against *Pae*^WT^, *Pae*^Δtae1^, *Pae*^inactive^, for four biological replicates. For each plot, replicate 1
is compared to replicate 2 (grey), replicate 3 (red), or replicate 4 (blue). Median Pearson's *r* between all replicates = 0.91.
1117

1118
Supplement 4: Pooled T6SS competitions with basal CRISPRi attenuate significant fitness phenotypes
a-b) Basal CRISPRi attenuates *Eco* fitness phenotypes against *Pae^WT* (a) and *Pae^Δtae1*(b). Volcano plots showing log$_2$-fold change (L2FC)
values for each KD strain after interbacterial competition (basal CRISPRi). Data shown: mean from four biological replicates. Statistical test:
Wald test. Vertical dotted lines indicate arbitrary cutoffs for L2FC at x =-1.58 and x=1.58 (absolute FC x=-3 or x= 3). Horizontal dotted line
indicates statistical significance cutoff for log$_{10}$ adjusted p-value (≤ 0.05). Orange points represent KDs with L2FC ≥ 1.58 or ≤ -1.58 and log10-
adj. *p*-value ≤0.05. Dark purple points represent non-targeting negative control KDs (*n*=50). Lavender points represent KDs that do not meet
cutoffs for L2FC or statistical test.

Supplement 5: *Pae^inactive* is a neutral co-culture partner for *Eco*

a-b) Competition against *Pae^inactive* reveals few *Eco* fitness determinants. Volcano plots showing log$_2$-fold change (L2FC) values for each KD strain after interbacterial competition with induced (a) or basal (b) CRISPRi. Data shown: mean from four biological replicates. Statistical test: Wald test. Vertical dotted lines indicate arbitrary cutoffs for L2FC at x =-1.58 and x=1.58 (absolute FC x=-3 or x= 3). Horizontal dotted line indicates statistical significance cutoff for log$_{10}$ adjusted p-value (≤ 0.05). Orange points represent KDs with L2FC ≥ 1.58 or ≤ -1.58 and log10-adj. *p*-value ≤0.05. Dark purple points represent non-targeting negative control KDs (*n*=50). Lavender points represent KDs that do not meet cutoffs for L2FC or statistical test. **c-d) KD strain abundance is highly similar after competition with *Pae^inactive* and after growth without competition (*Eco^ctrl*).** Scatter plots comparing mean L2FC for each *Eco* KD strain after competition with *Pae^inactive* or *Eco^ctrl* treatment, with basal (c) or induced (d) CRISPRi. Median Pearson correlation *r*= 0.98.

1137
Supplement 6: *lpxK-KD* and *msbA-KD* modulate target gene expression and show polar effects
a-b) Transcriptional knockdowns in *msbA* and *lpxK* have off-target polar effects on transcription in their operon. qRT-PCR analysis of
transcriptional fold-change in *ycaI-msbA-lpxK-ycaQ* in *msbA-KD* (a) and in *lpxK-KD* (b) after growth for 6 hours with induced CRISPRi,
normalized to expression in *rfp-KD*. Data shown are geometric average of 3 biological replicates ± s.d. Statistical test: unpaired two-tailed *t*-
test; *p*-value ≤0.05 displayed in bold font. **c-d) *msbA-KD* and *lpxK-KD* express a catalytically dead Cas9 (dCas9) enzyme for CRISPRi-
mediated transcriptional knockdown.** Western blot analysis of dCas9 protein expression (160 kDa) from *rfp-KD, msbA-KD* (c), and *lpxK-KD*
(d). Three independent biological replicates shown.
1145

1146
1147 **Supplement 7: Tae1 protein expression is unaffected in _msbA-KD_**
1148 **a-b) Bulk Tae1 protein expression is similar between _msbA-KD_ and _rfp-KD_**. Western blot analysis of periplasmic Tae1 protein (17kDa)
1149 from (a) _pBAD24::pelB-tae1^WT_ (Tae1^WT) or (b) _pBAD24::pelB-tae1^C30A_ (Tae1^C30A) in _rfp-KD_ and _msbA-KD_ (with induced CRISPRi). Protein
1150 expression of RNA polymerase (β subunit) (150kDa) is used as an internal loading control.

Supplement 8: Plasmid-borne overexpression of _msbA_ partially rescues Tae1 sensitivity in _msbA-KD_

a-c) Plasmid-borne _msbA_ overexpression partially rescues _msbA-KD_ resistance to lysis by Tae1. OD_{600} growth curves of _msbA-KD_ with induced CRISPRi, overexpressing _pSCrhaB2::msbA_ (orange) or _pSCrhaB2_ (empty) (grey) alongside (a)_pBAD24::pelB-tae1^{WT}_ (Tae1^{WT}), (b) _pBAD24::pelB-tae1^{C30A}_ (Tae1^{C30A}), or (c) _pBAD24_ (empty). Data shown: average of 3 biological replicates ± s.d. Dotted vertical line indicates _pBAD24_ induction timepoint (at OD_{600}=0.25) (0.125% arabinose w/v). **d-f) Plasmid-borne _lpxK_ overexpression enhances _msbA-KD_ resistance to lysis by Tae1.** OD_{600} growth curves of _msbA-KD_ with CRISPRi induced, overexpressing _pSCrhaB2::lpxK_ (purple) or _pSCrhaB2_ (empty) (grey) alongside (d)_pBAD24::pelB-tae1^{WT}_ (Tae1^{WT}), (e) _pBAD24::pelB-tae1^{C30A}_ (Tae1^{C30A}), or (f) _pBAD24_ (empty). Data shown: average of 3 biological replicates ± s.d. Dotted vertical line indicates _pBAD24_ induction timepoint (at OD_{600}=0.25) (0.125% arabinose w/v). **g-h) _pSCrhaB2_ vectors selectively rescue transcription of their target gene by overexpression.** qRT-PCR analysis of transcriptional fold-change in (g)_msbA_ or (h)_lpxK_ expression with constitutive rhamnose induction of _pSCrhaB2::msbA_ (orange), _pSCrhaB2::lpxK_ (purple), or (c)_pSCrhaB2_ (empty; grey) in _msbA-KD_ with induced CRISPRi. Expression normalized against basal _msbA_ expression in _rfp-KD_ + _pSCrhaB2_ (empty). Data shown: geometric average of 3 biological replicates ± s.d. Statistical test: unpaired two-tailed _t_-test; _p_-value ≤0.05 displayed in bold font.

1165
1166 **Supplement 9: Tae1^{C30A} hydrolyzes D44 muropeptides in *rfp-KD* and *msbA-KD***
1167 **a)Tae1^{C30A} overexpression yields minor digestion of D44 muropeptides**. HPLC chromatograms of muropeptides purified from *msbA-KD*
1168 (orange) and *rfp-KD* (grey) expressing *pBAD24::pelB-tae1^{C30A}* (Tae1^{C30A}). Black arrow indicates D44 peptide partially digested by Tae1^{C30A}.
1169 Data shown: representative from 3 biological replicates.
1170

1171
Supplement 10: PG synthesis activity is _msbA-KD_ is suppressed across all conditions
a) PG synthesis activity in _msbA-KD_ is attenuated under all tested conditions. Single-cell fluorescence intensity measurements for _rfp-_
_KD_ (grey) or _msbA-KD_ (orange) incorporating the fluorescent D-amino acid HADA into PG after 60 minutes of overexpressing _pBAD24::pelB-_
tae1^{WT}(Tae1^{WT}), _pBAD24::pelB-tae1_^{C30A} (Tae1^{C30A}), or _pBAD24_ (empty), with CRISPRi induced. All data normalized to average HADA signal in
rfp-KD + empty. Data shown: 600 cells (200 cells x 3 biological replicates), with average ± s.d. Statistical test: unpaired two-tailed _t_-test; _p_-
value ≤0.05 displayed in bold font.

1178
1179 **Supplement 11: _msbA-KD_ growth defects are independent of _Pae_ T6SS activity**
1180 _a-b) **msbA-KD** cells maintain growth defects regardless of **Pae** competitor_. Representative frames from time-course imaging of _rfp-KD_
1181 (left column; grey cells) and _msbA-KD_ (right column; grey cells) co-cultured with _Pae_ [Δtae1](a) or _Pae_ [inactive] (b) (green cells), and with induced
1182 CRISPRi. Green foci in _Pae_[WT] indicate aggregates of GFP-labelled ClpV, which signal H1-T6SS firing events. Red arrow indicates lysed cell.
1183 Data shown are merged phase and fluorescent channels. Scale bar: 2μm.