Shift from a Simplified to Complex Gut Microbiota Reduces Adenoma Burden in a Preclinical Rat Model of Colon Cancer

Susheel Bhanu Busi[1], Daniel Davis[2], Jacob Moskowitz[3], and James Amos-Landgraf[3,#].

[1]Department of Molecular Microbiology and Immunology, University of Missouri School of Medicine, One Hospital Drive, MA204, Columbia, MO 65212

[2]Animal Modeling Core, Office of Research, University of Missouri, 1600 Rollins Street, Columbia, MO 65211

[3]Department of Veterinary Pathobiology, University of Missouri School of Veterinary Medicine, 1600 Rollins Road, Columbia, MO 65211

Running title: Complex Gut Microbiota reduces adenomas in Pirc rats

[#]Corresponding author: Dr. James Amos-Landgraf, amoslandgrafj@missouri.edu

Abstract

Specific bacterial taxa in the gastrointestinal tract have been strongly associated with cases of colorectal cancer (CRC) cancer in familial adenomatous polyposis and spontaneous disease cases in humans. This has been recapitulated in animal models of CRC with positive correlations with many commensals and pathogens. However, many of these studies are performed either in germ-free animals or employ an antibiotic regime, overlooking the complex interactions of the commensals within the colon. To simplify the challenges associated with the complexity of the microbiota in the GI tract we established the Pirc rat model of colon cancer on an Altered Schaedler Flora (ASF) gut microbiota (GM), maintained in a barrier room. To elucidate the role of the simplified (ASF) and conventional GMs on disease susceptibility, We conventionalized ASF Pirc littermates. We found that the conventionalized F1 rats had increased microbial diversity and decreased colonic adenoma multiplicity. Our findings show that the complexity and the interactions of the GM community and not a *Firmicutes* to *Bacteroidetes* ratio are an important factor affecting disease susceptibility.

Introduction

Colorectal cancer (CRC) models, including both mice and rats, have been used to understanding the etiology of human diseases for decades [1,2]. The ideal model should recapitulate the phenotype observed in humans, but also elucidate contributing factors such as the host microbiota and its relationship to the mechanisms of the disease. Recent evidence suggests that the gut microbiome, i.e. the collection of microorganisms in the large intestine plays an important role in the etiology of the disease [3,4]. Several studies have tried to elucidate the mechanisms by which specific bacteria

contribute to disease susceptibility by various methods including the utilization of germ-free [5,6] or

monocolonized animals [7,8], or the use of antibiotics to eliminate endogenous gut microbiota (GM)

populations [3]. These majority of studies use the $Apc^{+/Min}$ mouse model that develop the majority

of their tumors in the small intestine unlike human disease. Since the GM population has been

shown to be different in the small intestine compared to the colon the translatability of these studies

may be limited.

The Pirc (F344/NTac-$Apc^{+/am1137}$) rat model of human colon cancer demonstrates a more

consistent colonic tumor phenotype compared to the $Apc^{+/Min}$ mice and has been shown to have an

altered phenotype with altered gut microbiota [2,9,10]. To model more closely, the large number of

endogenous commensals found in human CRC patients, we previously showed that the

endogenous GM could be modulated through complex microbiota targeted rederivation [2,9–11].

Determining the mechanisms and most importantly the interactions between commensals still pose

challenges, considering the multiple permutations and combinations with the taxa found in the

model.

In order to tackle the challenge of complexity, we established the Pirc rat on an Altered Schaedler

Flora (ASF) gut microbiota [12]. The ASF model includes an eight-community gut microbiome

derived from mice, was developed in 1970 by R.P. Orcutt [13]. Considering the highly diverse

microbiome in animal models, the ASF community allows for in-depth identification of host-

microbe interactions, including their spatial arrangment [12]. Instituting Pirc rats on a minimal GM

profile could potentially serve as a model for understanding mechanisms and interactions of

specific bacteria, in the context of a well-defined, yet complex gut microbiome profile. Using

CRASF (Charles-River ASF) rats as surrogates, F1-Pirc rats were established, and at weaning,

littermates were transferred from a barrier room to a conventional status room in the animal

facility. We hypothesized that transferring the Pirc rats to a conventional room compared to the

cleaner, barrier room would increase the colonic tumor burden at sacrifice. Contrary to our

hypothesis, we found that the animals maintained in the barrier (clean) room had significantly

more colonic adenomas. This is the first time Pirc rats have been established on an Altered

Schaedler Flora gut microbiota, but more importantly, suggest an even more central role for the

gut microbiota in modulating the colon tumor phenotype of animal models for studying human

diseases.

Results

Nominal taxa incursion in the Charles River Altered Schaedler Flora (CRASF)

In order to establish F344/NTac-$Apc^{+/am1137}$ rats onto a CRASF gut microbiota (GM), we first had

to ensure that the simplified GM profile could be maintained in our facility. We housed four female

and four male LEW/Crl ASF (CRASF) rats in a barrier room setting with individually ventilated

racks in microisolator cages. Fecal samples collected prior to arrival at the facility and upon

housing for 3 months at the facility, showed minimal addition of species to the GM profile. Over

time, the LEW/Crl ASF animals acquired *Lachnospiraceae UCG-001, Lachnospiraceae UCG-*

006, Anaerotruncus, [Eubacterium], Enterococcus and Staphylococcus (Fig.1a). Several cross-

fostering experiments were needed to obtain a Pirc rat on a CRASF GM background. Out of 4 that

were successfully cross-fostered, we only had one Pirc male rat. The F344/NTac-$Apc^{+/am1137}$ rat

that was so cross-fostered onto the CRASF surrogates, at 1 month of age, showed a stable GM

similar to that of the CRASF rats (Fig.1a). The ZymoBIOMICS mock microbial community

standards simultaneously only acquired *Enterobacteriaceae,* potentially via the sample processing

or sequencing or the bioinformatics analysis and annotation pipeline. Interestingly, the incursion

of six taxa into CRASF rats led to significant differences when visualized using a Principal

Coordinate analysis (PCoA) to understand the similarities between samples pre- and post- arrival,

using the bray-curtis distance matrix (Fig.1b). The majority of the GM was maintained stably after

housing the CRASF animals in a barrier setting for 3 months. To determine if the OTUs were

acquired as a means of shipping to our facility, sequencing was performed on the bedding and gel-

paks that the animals arrived with and found that four of the OTUs were possibly assimilated

through the gel-paks, with *Muribaculaceae* making a significant contribution to the overall GM

profile (Fig.1c).

Simplified gut microbiota increases susceptibility to colonic adenomas

LEWF344F1-*Apc^{+/am1137}* CRASF rats obtained via the breeding set up were used to understand

how the complexity of the gut microbiota (GM) may modulate disease susceptibility to adenomas

in the rat model of colon cancer. At weaning, F1 Pirc littermates were separated into two separate

rooms of the animal facility; a barrier room, where all cage changes were performed in a

biocontainment hood, and a conventional room (Supplementary Fig.1). We found that the animals

housed in the conventional setting had significantly fewer colonic adenomas than those housed

under barrier conditions (Fig.2a). This differential tumor abundance was found in both male and

female F1-Pirc rats. Interestingly, male rats from the conventional room had significantly more

small intestinal tumors compared to the barrier rats, while female rats showed a similar trend (Fig.2b).

Altered Schaedler Flora alters the colonic adenoma phenotype and the physiology of the gastrointestinal tract

Animals housed in the barrier room post-weaning demonstrated an increase in the number of proximal adenomas compared to conventionalized CRASF Pirc rats (Fig.3a). Most of these adenomas were 1 mm or smaller in diameter, however the rats with conventional GM did not show a similar phenotype (Fig.3b). Only one of twelve F1 Pirc rats separated at weaning and housed under a conventional settings had adenomas in the proximal colon that was slightly larger than 1 mm (Table.1). We also found that the overall number of small adenomas was significantly higher in the barrier room animals, irrespective of sex (Fig.3c), and the adenomas larger than 1 mm did not show any significant differences between the barrier and conventional rats' (Fig.3d).

Furthermore, we sacrificed a cohort of F1 Pirc ASF animals at weaning and found no differences in their cecal size (Fig.3e). However, sacrifice after housing under barrier or conventional settings for 4 months, revealed considerable differences in cecum size. We found that rats maintained in the barrier room had ceca that were nearly 2-fold larger compared to conventionally-housed rats (Fig.3f). These results suggest that the lack of taxa from the conventional GM and/or their interactions with the Altered Schaedler Flora (ASF) in barrier rats is capable of modifying the physiology and the phenotype of the F1 Pirc rats.

Conventional housing affects the GM architecture at 4 months of age

Considering the husbandry, handling, cecal and tumor multiplicity differences between the barrier and conventional rooms, we used 16S rRNA gene sequencing to determine the GM architecture in the F1-Pirc rats. At weaning, we found that the GM of rats at the time of separation into barrier and conventional rooms were similar to each other as indicated by bar graph (Fig.4a) and the Principal Coordinate analysis in Fig.4b (using Bray-Curtis distance matrix) and the overall richness determined by the number of OTUs observed in the samples (Fig.4c). They also resembled the GM profile of the parents, except for the conspicuous decrease in the relative abundance of genus *Mucispirillum* (Fig.4a).

At sacrifice (4 months of age), considerable differences were observed in the overall Genus' profile of the GM between the barrier and conventionally-housed F1-Pirc rats (Fig.5a). At the genus level (Supplementary Fig.2A), several taxa including *Parabacteroides, ASF356, Blautia, and Mucispirillum* were elevated in the barrier F1-Pirc rats. In the conventionalized rats there was an observed increase in the relative abundance of over 50 taxa, the top 35 are depicted in the heatmap (Supplementary Fig.2a). The overall GM profile composition differences are visualized using a Principal Coordinate analysis (Fig.5b). The most separation was observed along PCo1, suggesting that the room differences contribute to the majority of the variability in the GM architecture. There were also significant increases in the richness and diversity indices such as Chao1 and Shannon (Fig.5c-e). These results suggest that the contribution of the room differences, such as husbandry, handling, and exposure to conventional animals have a crucial effect on the acquired taxa. *Firmicutes* and *Tenericutes* were increased in the conventionally-housed rats, whereas

Bacteroidetes was decreased. This also led to a significant shift in the *Firmicutes:Bacteroidetes*

ratio between the two groups (Fig.5f).

We found significant correlations between certain taxa from the barrier (Fig.6a and Supplementary

Fig.3a) room at weaning with the colonic tumor burden including the small adenomas. In these

F1-Pirc rats, decrease in *Erysipelotrichaceae* and the genus *Parabacteroides* were associated with

an increase in the colonic tumor count, whereas order *Peptostreptococcaceae* was found to show

a positive correlation with tumor burden. Other taxa such as *Ruminococcaceae* and

Lachnospiraceae showed similar correlations. Similarly, *Bacteroides, Peptococcus, Clostridiales,*

Peptococcaceae and *Candidatus saccharomonas* showed significantly positive correlations.

Correlation analysis with the colonic tumor burden of the conventionally raised F1-Pirc animals

(Fig.6b and Supplementary Fig.3b) showed that family *Prevotellaceae* at weaning was negatively

correlated with tumor burden. *Clostridum Family XIII, Lachnospiraceae, Ruminococcus UCG-*

006, UCG-008, UCG-010 and *Clostridiales*, on the other hand were positively correlated with

tumor increase, suggesting a causative role for the taxa from these operational taxonomic units.

Simultaneously, we found significant negative correlations between *Parabacteroides* and specific

OTUs from the barrier room at 4 months of age. These OTUs included *Ruminococcus NK4A136,*

Roseburia, Lachnospiraceae, Instestimonas and Oscillibacter. Other OTUs including

Parabacteroides and *Prevotellaceae* had positive and negative correlations respectively with the

other commensals such as *ASF356, Mucispirillum, Lachnospiraceae* and *Ruminococcus UCG*

taxa.

We used PICRUSt (Phylogenetic Investigation of Communities by Reconstruction of Unobserved

States) and the HMP Unified Metabolic Analysis Network (HUMAnN) to understand the

functional capacity of bacterial taxa in the fecal samples collected at 4 months of age

(Supplementary Fig.2b). We found that the barrier room F1 Pirc rats had a substantial increase in

the Spermidine-putrescine transport system and the succinate dehydrogenase pathways.

Conversely, the predicted functional capacity of the conventional room rats showed increased

abundance of pathways related to bacterial cell doubling time including Kreb's cycle, increased

amino acid biosynthesis (methionine and leucine), the iron transport system and increased sulfate

reduction to hydrogen sulfide.

180 Barrier and conventional room diets have distinct GM populations

Targeting the 16S rRNA gene, we sequenced the feed from the barrier and conventional rooms

where we found that the GM profile of the feed from the two rooms did not differ significantly

(Supplementary Fig.4a). Feed from the barrier and conventional rooms demonstrated similar

abundances of order *Streptophyta* and *Zea luxurians*; genetic content likely derived from plant

material used in the preparative process for feed (Supplementary Fig.5a). Though the community

profile appeared similar, the relative abundances of *Lactobacillus, Leuconostoc, Sphingomonas*

and *Fusobacterium* were significantly increased in the barrier room feed compared to the

conventional diet (Supplementary Fig.5c). More importantly, to delineate between residual genetic

content after autoclaving and potential taxa that may colonize the rats in the conventional room

we cultured the feed from both rooms. Under anaerobic conditions, we cultured the feed overnight

at 37 ˚C using brain heart infusion media. We observed several taxa in the feed from the

conventional room grew abundantly, whereas the barrier room chow had minimally detectable

levels of taxa such as *Clostridium* and *Bacillus* (Supplementary Fig.4b). This was also observed

in the rarefaction curves (Supplementary Fig.5b) when sampling the observed species in each of

the samples. We found that the cultures with the conventional feed had several species that were

identifiable compared to both the uncultured conventional and barrier feed, including the cultured

barrier feed. This suggested that bacteria from the feed could potentially have colonized the

gastrointestinal (GI) tract of the F1 Pirc rats that were housed under conventional conditions,

possibly altering the phenotype. In fact, we found that operational taxonomic units (OTUs) found

in the feed such as *Lachospiraceae, RF39, Ruminiclostridium, Oscillibacter, Ruminicoccaceae*

and several genus' of the *Muribaculaceae* order were found in the conventionally raised F1-Pirc

animals, but were undetectable in the barrier room rats (Supplementary Fig.5a).

Discussion

The human colon is host to approximately 10^{14} bacteria alone, aside from viruses and fungi, which

together form the gut microbiota. The interaction between the host and the endogenous GM is

highly varied and complex which may be a crucial part of disease susceptibility. However,

modelling the interactions of the GM in a complex setting is challenging. Therefore, we generated

F344/NTac-*Apc*$^{+/am1137}$ (Pirc) rats and fostered them onto a Charles River Altered Schaedler Flora

(CRASF) gut microbiota profile. We were able to stably maintain the ASF GM with only the

acquisition of a few OTUs such as *Anaerotruncus, and Staphylococcus* etc. More importantly, the

F1-Pirc (LEWF344F1-*Apc*$^{+/am1137}$ CRASF) animals generated resembled the CRASF parents at

weaning except genus *Mucispirillum* which was decreased in the offspring compared to the

breeders. This taxon is difficult to culture *in vitro* compared to other ASF taxa, however, it is still

not known whether it is inhibited by the presence of other OTUs usually observed in altered

Schaedler flora colonies [14]. Based on reports of the CRASF populations by Matthias *et al* [15] we

find a majority of the original CRASF populations within our colony.

Dove et al [16] showed that germ-free model of the Apc mice developed two-fold fewer adenomas

than the conventional. In rats, Reddy et al [17] also showed that germ-free animals, though treated

with chemical carcinogens, had fewer tumors than conventional rats. Similarly, previous studies

by Zackular *et al* [3], where the same model was treated with antibiotics showed a signficiant

reduction in both small intestinal and colonic tumors. We hypothesized based on the phenotype of

colon cancer in germ-free or antibiotic-treated animal models, that the F1-Pirc CRASF rats in the

barrier room would have fewer colonic adenomas. Contrary to our hypothesis, animals maintained

under barrier conditions had an increased tumor burden, including significantly increased number

of smaller adenomas especially in the proximal section of the colon. In the original report of the

Pirc rat [2,3], microadenomas required histopathological confirmation and were recorded as being

smaller than 0.5 mm in diameter. However, in our case the differences between the barrier and

conventional rooms were grossly apparent, including an increase in the number of smaller

adenomas when compared to previous reports of conventional and germ-free rats. This forms an

interesting hypothesis as to why and which OTUs of the GM may be initiating the shift from a

distal colonic to a proximal. This disparity will be addressed in future studies. It must be noted

though that excluding the smaller adenoma numbers, there was a trend towards increased tumors

in the conventional animals as originally hypothesized. This posits for future studies where F1-

Pirc rats would be aged longer than 4 months to understand if the observed small adenomas may

develop into adenomas larger than 1 mm in diameter. Another observation from our study was the

alteration of the colonic tumor phenotype observed in the F1-Pirc rats from the barrier room.

Typically, Pirc rats demonstrate a colonic phenotype where the adenomas develop in the middle

and distal portion of the colon with few in the proximal regions, as we observed in the F1-Pirc

animals from the conventional room. However, F1-Pirc animals from the barrier room had several

adenomas in the proximal section of the colon with few or none in the middle and distal regions.

Similar to previous reports of germ free animals, these animals had enlarged ceca compared to

conventional F1-Pirc rats [18][19,20]. Zackular *et al.* showed in an AOM/DSS-treated mouse model that

a decrease in the overall GM population through the administration of antibiotics, led to a

significant decrease in tumor burden. Another study similarly demonstrated that transferring

tumor-associated microbiota into germ-free mice increased the tumor burden of the mice,

otherwise significantly reduced when mice were maintained germ-free [21]. Based on these reports,

our findings of animals maintained in a barrier room having significantly elevated adenomas is

intriguing. Although studies have shown that bacteria are needed for a phenotype to be manifested

in animals [22,23], our results suggest that a consortium of taxa may influence disease. Also, our

observations that a simple, yet diverse consortium of taxa within the gut microbiome, still

maintains a large cecal size as compared to conventionally-housed rats is notable. However, in

previous reports by [19,23,24], the cecal size is reported as a measure of percent body weight. Future

studies will analyze the differences in cecal sizes compared to other germ-free or mono-colonized

models. In 1962, Skelly et al [25] reported that certain Clostridium species restored cecal size to what

is observed in normal conventional settings. Alternatively, Loesche in 1969 reported that B.fragilis

caused an increase in cecal size [24,25]. We observed both Clostridium and Bacteroides species within

our microbiome analyses, and will need additional studies including germ-free, ASF, and

conventional Pirc rats to identify the source of this variation.

The barrier room was maintained with irradiated chow, paper chip bedding, autoclaved water and

animals were always handled in a biosafety cabinet. We housed the conventional room rats with

non-irradiated feed, non-autoclaved bedding and used animal handling techniques that did not

require aseptic methods. We hypothesized that this would alter the existing CRASF microbiota to

a more complex GM. We used 16S rRNA sequencing to determine if the GM, known to be

modulated by husbandry factors [2626–28] was the crucial modulator of the phenotype observed in our

study at 3-months after introduction into the conventional facility, we found the conventional rats

had acquired OTUs including *Prevotellaceae, Ruminococcaceae, Muribaculaceae, Parasutterella*

and *Desulfovibrionaceae*. *Prevotellaceae* and *Desulfovibrionaceae* have been reported to be

associated with healthy patients or a decreased tumor burden in colon cancer studies [3,10,29]. On the

other hand, *Blautia, Enterococcus,* and some *Lachnospiraceae* taxa found in the barrier room F1-

Pirc rats have been associated with an increased tumor susceptibility [30–33]. This was equally evident

from the correlations where *Peptococcaceae, Clostridiales, Lachnospiraceae,* previously reported

to be associated with an increased tumor burden were elevated and positively correlated with the

tumor burden in the barrier rats [34,35]. Correlation analysis also found that certain OTUs introduced

into the conventional rats had a negative association with *Parabacteroides*, potentially suggesting

that these OTUs inhibit the proliferation or take over the niche occupied by the latter, i.e.

competitive interactions [36]. In the barrier room rats we also found predicted functional pathways

such as succinate dehydrogenase and spermidine-putrescine transport system to be elevated. Host

succinate dehydrogenase mutations are very commonly found in colon cancer [37,38]. This raises the

possibility of a breakdown of the host dehydrogenases, thereby leading to an increase in the

bacterial dehydrogenase expression to counteract the toxic effect of succinate. Alternatively, many

rumen bacteria are known to produce succinate which in turn has been identified as a biomarker

for colon cancer via mass spectrometry [39]. This suggests elevated levels of succinate, reportedly

an onco-metabolite [40] could be promoting tumorigenesis in the barrier room rats via inhibition of

PHD (prolyl hydroxylase domain-containing) enzymes [41] via activation of hypoxia-induced factor

alpha (HIF-α). Succinate quantitation via metabolomics and PHD enzyme activity will however

need to be validated in future studies to determine the mechanisms contributing to increased

succinate levels. Similarly, polyamines such as spermidine and putrescine have been reported to

be biomarkers for colorectal cancer in human patients [42]. In 1988, Upp *et al.* analyzed the

polyamine levels including spermidine and putrescine in colon cancer patients and found that they

may be used to identify at-risk patients of the disease [42,43]. More recently, it was identified that GI

bacteria such as *Bacteroides fragilis* upregulates spermine oxidase which induces production of

spermidine, hydrogen peroxide and aldehydes [42–44], potentially causing DNA damage. Another

thought-provoking observation in our study is the presence of OTUs found in the diet that were

detected in the barrier and conventional room fecal samples from F1-Pirc rats. Although, the

barrier room rats were not over-ridden by the taxa found via 16S sequencing, this was not true for

the conventionally raised rats. We found that the conventionally housed F1-Pirc rats had

significant amounts of bacterial taxa that were also detected in the diet, and that were anaerobically

cultivable. This suggests that the non-irradiated diet, may be one source of the variation, although

it is also possible that the rare OTUs picked up are nonviable residual DNA from dead bacteria or

spores. More importantly, this source, potentially led to a significant shift in the phenotype, i.e.

number of adenomas.

Colorectal cancer (CRC) animal models have been extensively used to study and understand the

etiology of the disease including initiation, development and factors affecting susceptibility.

Despite the development of the $Apc^{+/am1137}$ rat, the $Apc^{+/Min}$ mouse model of colon cancer is still

largely used for various studies owing to cost and the ease of genetic manipulation techniques.

However, the Pirc ($Apc^{+/am1137}$) rat with a colonic phenotype has created a potentially more

translatable alternative to the mouse when studying colon cancer. While not the primary focus of

this study, we realize that it is not inconceivable that there may be inherent differences between

the gut microbiome of wild type and CRASF-Pirc rats and that genetics play a crucial role in the

composition of the GM. However, with studies recently reporting evidence of the role of the gut

microbiota in diseases susceptibility including colon cancer [3,4,6,33], the importance of

reproducibility in disease models is critical. Several reports have identified *Fusobacterium*, in

particular *Fusobacterium nucleatum,* as a significant modifier of disease burden [45,46]. This bacteria

along with Enterotoxigenic *Bacteroides fragilis* (ETBF) has often been associated with increased

tumor burden and/or carcinoma samples in human patients [4,7]. However, it should be noted that

most of these studies do not take into account the constant interactions and synergistic nature of

the commensals within the GI tract. GM populations are a constant source of nutrients and

metabolites, which are contingent on the action of one bacterium on the by-products of the

replicative processes of another. To model and establish a simplified GM profile to study the role

of specific bacteria and their interactions with the host and other commensals, we established

Apc$^{+/am1137}$ rats on a CRASF (Charles River Altered Schaedler Flora) gut microbiome profile. The

observance of increased tumor number in a limited GM microbiome provides a platform for

probiotic experimentation. It can also allow for more refined metabolite profiling and longitudinal

assessment in changes in metabolic processing. Utilizing a simplified GM profile for

understanding the pathophysiology of colon cancer, may provide insights into the interactions

between commensals and with the host, including the mechanisms by which specific taxa promote

or prevent adenomagenesis.

Methods

Animal Care and Use

All procedures were performed according to the guidelines regulated by the Guide for the Use and Care of Laboratory Animals, the Public Health Service Policy on Humane Care and Use of Laboratory Animals, and the Guidelines for the Welfare of Animals in Experimental Neoplasia and were approved by the University of Missouri Institutional Animal Care and Use Committee.

Charles River Altered Schaedler Flora (CRASF) rats and cross-fostering

7 week old Lewis rats with a limited Altered Schaedler Flora (n = 4 males, and 4 females) were purchased from Charles River Laboratories Inc. Laboratories (Wilmington, MA). The animals were shipped overnight in a sterile double-enclosed isolator cage with sterile bedding and Hydrogel® gel packs (Portland, ME) to the Discovery Ridge animal facility at University of Missouri. Fecal samples were collected prior to shipping and upon arrival at the facility for 16S rRNA sequencing. Simultaneously, bedding and gel-pak samples that the animals were shipped with were also collected for sequencing. The animals were housed in a barrier room on ventilated racks (Thoren, Hazleton, PA) in micro-isolator cages with autoclaved paper chip bedding (Shepherd Specialty Paper, Milford, NJ) and were fed irradiated 5053 PicoLab Mouse Diet 20 (LabDiet, St. Louis, MO), and water (reverse osmosis purified and sulfuric acid (pH 2.5-2.8) acidified followed by autoclaving), and allowed to acclimatize for a week, after which they were setup into breeder pairs. Timed matings for fostering were set up with our F344/NTac $Apc^{+/am1137}$ (generation, N=28) conventional rat colony.

Female F344/NTac rats were checked for plugs, and on day 21 post observation of plugs, a

Caesarean was performed. The uterus was tied-off at both ends prior to surgical resection and then

transferred in a sterile petri dish with betadine solution to the barrier room. In a biosafety hood,

the uterus was opened with a pair of sterile scissors and the pups were physically manipulated after

removing the amniotic sac and warmed under a heat lamp. Only CRASF breeders with pups on

the ground within 36 hours were used as surrogates for fostering the F344/NTac-$Apc^{+/am1137}$ pups.

Half the litter and bedding was removed from the CRASF surrogate, and mixed with the to-be

fostered pups thoroughly, before placing the F344/NTac-$Apc^{+/am1137}$ fostered pups along with a

few of the CRASF pups with the surrogate mom. At 12 days of age, all pups including the fostered

ones were clipped for genotyping.

Genotyping and animal identification

Pups were ear-punched prior to weaning at 12 days of age using sterile technique. DNA was

extracted using the "HotSHOT" genomic DNA preparation method previously outlined [47]. DNA

was used for genotyping using a high resolution melt (HRM) analysis as described previously [10,47].

Experimental design, animal husbandry (breeding) and barrier room housing

F1-Pirc rats were generated by crossing one founder male, F344/NTac-$Apc^{+/am1137}$ CRASF Pirc rat

established via cross-fostering, with wild-type female LEW/Crl ASF rats. The rats were housed

on ventilated racks (Thoren, Hazleton, PA) in micro-isolator cages as described above. Animal

handling required complete personal protective equipment (PPE) including face masks, hair nets

and TyVek sterile sleeves (Cat.No.17988110, Fisher Scientific, USA) and the use of 10% bleach

spray to treat surfaces including gloves and sleaves. Prior to breeding fecal samples were collected

from both the breeders using aseptic methods. LEWF344F1-$Apc^{+/am1137}$ (F1 generation) ASF pups

were generated and genotyped at 12 days of age.

Conventional room housing

375   At weaning, F1 Pirc rats were co-housed in the conventional room with F344/NTac animals from

376   the holding colony with an endogenous complex GM when available, in microisolator cages on

377   ventilated racks with nonsterile paper chip bedding. Cage changes for conventional rats were done

378   on open benches. Rats in the conventional room were fed non-irradiated 5008 Lab diet and had *ad*

379   *libitum* access to acidified (sulfuric acid, pH 2.5-2.8), autoclaved water.

Fecal sample collection

381   Fecal samples were collected from the pups at weaning, and monthly thereafter starting until

382   sacrifice at 4-months of age. Briefly, fecal samples were collected by placing the animal in a clean,

383   sterile cage without bedding. Freshly evacuated feces were speared with a sterile toothpick or

384   forceps and placed into a sterile Eppendorf tube. All samples were stored at -80 ˚C until further

385   processing.

Fecal DNA extraction, 16S library preparation, sequencing and analysis

Fecal samples were pared down to 65 mg using a sterile blade and then extracted using methods

described previously [10]. Amplification and sequencing of the V4 hypervariable region of the 16s

rDNA was performed at the University of Missouri Metagenomics center and DNA core facility

(Columbia, MO) and the results annotated using the SILVA 16S database [10,48]. Samples with a

read count below 15,000 were removed from the analysis due to insufficient rarefaction. The

average read counts for all samples was 57,863. Microbial Community DNA Standards from

ZymoBIOMICS [TM] were used to account for any errors via extraction and sequencing processes.

All OTUs with a relative abundance below 0.001% were excluded from analysis. Principal

Coordinate analyses were performed in PAST (PAleontological STatisitcs, version 3.2) [49].

PERMANOVA with default permutations (N=9999) was used to determine significant differences

between groups when performing PCoA analyses using the module embedded into PAST3.2.

Simultaneously, a scree plot was generated using the chemometrics.R script under the

m*etaboanalyst* package to identify which principal coordinates to plot for the figures. Heatmaps

were generated using the plotHeatMap function from the same package along with the hclust

function from the *stat* package. For the heatmaps, Euclidean distance was used as the similarity

measure, while Ward's clustering algorithm accounting for average linkage was used to create the

Dendrogram. Correlation analyses testing the relationship of OTUs' relative abundance with tumor

burden was assessed using the *corrgram* package in R (version 3.4.1), assessing the top 50 OTUs

based on the individual relative abundance. PICRUSt, HUMAnN and LEfSE analysis was

performed after re-annotating (closed-reference) the 16S rDNA gene sequences against the

Greengenes (May, 2013) database as described previously [49,50].

Anaerobic culturing of the lab diet feed and DNA extraction

3 samples of 0.5g of feed from the barrier and conventional rooms was introduced anaerobically

into an autoclaved serum vial, closed with a sterile rubber stopper and an aluminum crimp seal.

Oxygen was purged from vials and 5ml of sterile brain heart infusion (BD Difco, ThermoFisher

Scientific, USA) media was added using a syringe. The inoculum and media was then incubated

anaerobically overnight at 37 ˚C in a 5% CO_2 incubator. After incubation, the contents of the vial

were used for DNA extraction using previously established methods including manual DNA

precipitation and the DNEasy kit (Qiagen, USA) [51].

Tumor counts

All animals were humanely euthanized with CO_2, administration and necropsied at 16 weeks of

age. The small intestine and colon from the rats were placed on to bibulous paper and then splayed

opened longitudinally. Tissues were then fixed overnight in Carnoys fixative (30% Choloroform,

10% glacial acetic acid and 60% absolute ethanol), and were replaced with 70% ethanol for long

term storage until adenoma counting was performed.

Statistical analyses and figures

423 Statistical analyses and graphing for figures (except Fig.1) were prepared through GraphPad Prism

424 version 7 for Windows (GraphPad Software, La Jolla, CA). *p*-values were set to identify

425 significance at a value less than 0.05, unless otherwise indicated.

Ethics Approval and Consent to Participate

428 The current study was conducted in accordance with the guidelines set forth by the Guide for the

429 Use and Care of Laboratory Animals and the Public Health Service Policy on Humane Care and

430 Use of Laboratory Animals. All studies and protocols were approved by the University of Missouri

431 Institutional Animal Care and Use Committees.

Availability of Data and Material

434 The sequences and raw 16S rDNA read files generated and analyzed during the current study are

435 available at the NCBI GenBank database under BioProject ID# PRJNA490292.

Acknowledgements

The authors wish to acknowledge Giedre Turner, Becky Dorfmeyer and the MU Metagenomics

Center (MUMC) for their assistance with 16S rRNA gene sequencing; Brittany Lister and Office

of Animal Resources staff for assistance with animal husbandry; Charles River Laboratories Inc.

for assistance making the CRASF animals available and with sample collection prior to shipping.

This research was funded by the University of Missouri System Research Board grant. The funders

had no role in study design, data collection and analysis, decision to publish, or preparation of the

manuscript.

Author Contributions

447   Conceived and designed the experiments: S.B.B., J.M.A.L., D.D. Performed the experiments:

448   S.B.B. J.M.A.L., J.M. and D.D. Analyzed the data: S.B.B., D.D. and J.M.A.L, Manuscript

449   preparation: S.B.B., J.M.A.L. All authors contributed to the review of the manuscript.

Competing Interests

452   The authors declare no competing interests.

References

1.   Taketo, M. M. Colon Cancer and Polyposis Models. *Genetically Engineered Mice*

Handbook 233–251 Preprint at https://doi.org/10.1201/9781420039078.ch18 (2005).

2.   Amos-Landgraf, J. M. *et al.* A target-selected Apc-mutant rat kindred enhances the

modeling of familial human colon cancer. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 4036–4041

(2007).

3. Zackular, J. P., Baxter, N. T., Iverson, K. D. & Sadler, W. D. The gut microbiome

modulates colon tumorigenesis. *MBio* (2013).

4. Allen, J. & Sears, C. L. Impact of the gut microbiome on the genome and epigenome of

colon epithelial cells: contributions to colorectal cancer development. *Genome Med.* **11**, 11

(2019).

5. Slowicka, K. *et al.* Zeb2 drives invasive and microbiota-dependent colon carcinoma. *Nat*

Cancer **1**, 620–634 (2020).

6. Drewes, J. L. *et al.* Human Colon Cancer-Derived Clostridioides difficile Strains Drive

Colonic Tumorigenesis in Mice. *Cancer Discov.* **12**, 1873–1885 (2022).

7. DeStefano Shields, C. E. *et al.* Bacterial-Driven Inflammation and Mutant Expression

Combine to Promote Murine Colon Tumorigenesis That Is Sensitive to Immune Checkpoint

Therapy. *Cancer Discov.* **11**, 1792–1807 (2021).

8. Sears, C. L. The who, where and how of fusobacteria and colon cancer. *eLife* vol. 7 Preprint

at https://doi.org/10.7554/elife.28434 (2018).

9. Vitali, F. *et al.* Intestinal microbiota profiles in a genetic model of colon tumorigenesis

correlates with colon cancer biomarkers. *Sci. Rep.* **12**, 1432 (2022).

10. Ericsson, A. C. *et al.* Differential susceptibility to colorectal cancer due to naturally

occurring gut microbiota. *Oncotarget* **6**, 33689–33704 (2015).

11. Moskowitz, J. E. *et al.* Integration of genomics, metagenomics, and metabolomics to

identify interplay between susceptibility alleles and microbiota in adenoma initiation. *BMC*

Cancer **20**, 600 (2020).

12. Brand, M. W. *et al.* The Altered Schaedler Flora: Continued Applications of a Defined

Murine Microbial Community. *ILAR J.* **56**, 169 (2015).

13. Dewhirst, F. E. *et al.* Phylogeny of the Defined Murine Microbiota: Altered Schaedler

Flora. *Applied and Environmental Microbiology* vol. 65 3287–3292 Preprint at

https://doi.org/10.1128/aem.65.8.3287-3292.1999 (1999).

14. Biggs, M. B. *et al.* Systems-level metabolism of the altered Schaedler flora, a complete gut

microbiota. *ISME J.* **11**, 426–438 (2017).

15. Stehr, M. *et al.* Charles River altered Schaedler flora (CRASF) remained stable for four

years in a mouse colony housed in individually ventilated cages. *Lab. Anim.* **43**, (2009).

16. Taketo, M. M. Apc gene knockout mice, a model for human colorectal cancer. *Drug*

Metabolism and Pharmacokinetics vol. 12 145–145 Preprint at

https://doi.org/10.2133/dmpk.12.supplement_145 (1997).

17. Reddy, B. S., Narisawa, T. & Weisburger, J. H. Colon carcinogenesis in germ-free rats with

intrarectal 1,2-dimethylhydrazine and subcutaneous azoxymethane. *Cancer Res.* **36**, 2874–

2876 (1976).

18. Lynn, M. A., Ryan, F. J., Tee, Y. C. & Lynn, D. J. Protocol to colonize gnotobiotic mice in

early life and assess the impact on early life immune programming. *STAR Protoc* **3**, 101914

(2022).

19. Thompson, G. R. & Trexler, P. C. Gastrointestinal structure and function in germ-free or

gnotobiotic animals. *Gut* vol. 12 230–235 Preprint at https://doi.org/10.1136/gut.12.3.230

(1971).

20. Savage, D. C. & McAllister, J. S. Cecal enlargement and microbial flora in suckling mice

given antibacterial drugs. *Infect. Immun.* **3**, 342–349 (1971).

21. Zackular, J. P., Baxter, N. T., Chen, G. Y. & Schloss, P. D. Manipulation of the Gut

Microbiota Reveals Role in Colon Tumorigenesis. *mSphere* **1**, (2016).

22. Knights, D., Lassen, K. G. & Xavier, R. J. Advances in inflammatory bowel disease

pathogenesis: linking host genetics and the microbiome. *Gut* **62**, 1505–1510 (2013).

23. Holmes, E., Li, J. V., Marchesi, J. R. & Nicholson, J. K. Gut microbiota composition and

activity in relation to host metabolic phenotype and disease risk. *Cell Metab.* **16**, 559–564

(2012).

24. Loesche, W. J. Effect of bacterial contamination on cecal size and cecal contents of

gnotobiotic rodents. *J. Bacteriol.* **99**, 520–526 (1969).

25. Skelly, B. J., Trexler, P. C. & Tanami, J. Effect of a Clostridium species upon cecal size of

gnotobiotic mice. *Proc. Soc. Exp. Biol. Med.* **100**, 455–458 (1962).

26. Franklin, C. L. & Ericsson, A. C. Microbiota and reproducibility of rodent models. *Lab*

Anim. (2017).

27. Ericsson, A. C. *et al.* The influence of caging, bedding, and diet on the composition of the

microbiota in different regions of the mouse gut. *Scientific Reports* vol. 8 Preprint at

https://doi.org/10.1038/s41598-018-21986-7 (2018).

28. Rausch, P. *et al.* Analysis of factors contributing to variation in the C57BL/6J fecal

microbiota across German animal facilities. *Int. J. Med. Microbiol.* **306**, 343–355 (2016).

29. Sobhani, I. *et al.* Microbial dysbiosis in colorectal cancer (CRC) patients. *PLoS One* **6**,

e16393 (2011).

30. Arthur, J. C. *et al.* Intestinal inflammation targets cancer-inducing activity of the

microbiota. *Science* **338**, 120–123 (2012).

31. Wang, T. *et al.* Structural segregation of gut microbiota between colorectal cancer patients

and healthy volunteers. *The ISME Journal* vol. 6 320–329 Preprint at

https://doi.org/10.1038/ismej.2011.109 (2012).

32.   Weir, T. L. *et al.* Stool microbiome and metabolome differences between colorectal cancer

patients and healthy adults. *PLoS One* **8**, e70803 (2013).

33.   Baxter, N. T., Zackular, J. P., Chen, G. Y. & Schloss, P. D. Structure of the gut microbiome

following colonization with human feces determines colonic tumor burden. *Microbiome* **2**,

20 (2014).

34.   Reddy, B. S., Weisburger, J. H. & Wynder, E. L. Effects of high risk and low risk diets for

colon carcinogenesis on fecal microflora and steroids in man. *J. Nutr.* **105**, 878–884 (1975).

35.   Leung, A., Tsoi, H. & Yu, J. *Fusobacterium*and*Escherichia*: models of colorectal cancer

driven by microbiota and the utility of microbiota in colorectal cancer screening. *Expert*

Review of Gastroenterology & Hepatology vol. 9 651–657 Preprint at

https://doi.org/10.1586/17474124.2015.1001745 (2015).

36.   Kamada, N., Chen, G. Y., Inohara, N. & Núñez, G. Control of pathogens and pathobionts

by the gut microbiota. *Nat. Immunol.* **14**, 685–690 (2013).

37.   Kitazawa, S. *et al.* Succinate dehydrogenase B-deficient cancer cells are highly sensitive to

bromodomain and extra-terminal inhibitors. *Oncotarget* **8**, 28922–28938 (2017).

38.   Zhang, D. *et al.* Reduced succinate dehydrogenase B expression is associated with growth

and de-differentiation of colorectal cancer cells. *Tumor Biology* vol. 34 2337–2347 Preprint

at https://doi.org/10.1007/s13277-013-0781-4 (2013).

39.   Deng, L. *et al.* Development and Validation of a High-Throughput Mass Spectrometry

Based Urine Metabolomic Test for the Detection of Colonic Adenomatous Polyps.

Metabolites vol. 7 32 Preprint at https://doi.org/10.3390/metabo7030032 (2017).

40.   Wishart, D., Mandal, R., Stanislaus, A. & Ramirez-Gaona, M. Cancer Metabolomics and

the Human Metabolome Database. *Metabolites* vol. 6 10 Preprint at

https://doi.org/10.3390/metabo6010010 (2016).

41. Boroughs, L. K. & DeBerardinis, R. J. Metabolic pathways promoting cancer cell survival

and growth. *Nature Cell Biology* vol. 17 351–359 Preprint at

https://doi.org/10.1038/ncb3124 (2015).

42. Löser, C., Fölsch, U. R., Paprotny, C. & Creutzfeldt, W. Polyamines in colorectal cancer.

Evaluation of polyamine concentrations in the colon tissue, serum, and urine of 50 patients

with colorectal cancer. *Cancer* vol. 65 958–966 Preprint at https://doi.org/10.1002/1097-

0142(19900215)65:4<958::aid-cncr2820650423>3.0.co;2-z (1990).

43. Upp, J. R., Jr *et al.* Polyamine levels and gastrin receptors in colon cancers. *Ann. Surg.* **207**,

662–669 (1988).

44. Goodwin, A. C. *et al.* Polyamine catabolism contributes to enterotoxigenic *Bacteroides*

fragilis -induced colon tumorigenesis. *Proceedings of the National Academy of Sciences*

vol. 108 15354–15359 Preprint at https://doi.org/10.1073/pnas.1010203108 (2011).

45. Wu, N. *et al.* Fusobacterium nucleatum promotes colon cancer progression by changing the

mucosal microbiota and colon transcriptome in a mouse model. *World J. Gastroenterol.* **28**,

1981–1995 (2022).

46. Garrett, W. S. The gut microbiota and colon cancer. *Science* **364**, 1133–1135 (2019).

47. Truett, G. E. *et al.* Preparation of PCR-quality mouse genomic DNA with hot sodium

hydroxide and tris (HotSHOT). *Biotechniques* **29**, 52, 54 (2000).

48. Pfeiffer, S. *et al.* Improved group-specific primers based on the full SILVA 16S rRNA gene

reference database. *Environmental Microbiology* vol. 16 2389–2407 Preprint at

https://doi.org/10.1111/1462-2920.12350 (2014).

49. Hammer, Ø. & Harper, D. A. T. *Paleontological Data Analysis*. (John Wiley & Sons,

2008).

50. Montonye, D. R. *et al.* Acclimation and Institutionalization of the Mouse Microbiota

Following Transportation. *Front. Microbiol.* **9**, 1085 (2018).

51. Hart, M. L., Meyer, A., Johnson, P. J. & Ericsson, A. C. Comparative Evaluation of DNA

Extraction Methods from Feces of Multiple Host Species for Downstream Next-Generation

Sequencing. *PLoS One* **10**, e0143334 (2015).

Tables

Table 1: Altered Schaedler Flora alters the colonic adenoma phenotype and the physiology

of the gastrointestinal tract

Number of adenomas found in the small intestine (SI) and colon. Colonic adenomas smaller than

1 mm of diameter, in addition to the colonic adenomas are also provided. The last column indicates

the presence or absence of small, proximal colonic adenomas which are not typically seen in the

Pirc model of colon cancer.

Animal #	Sex	Room	SI adenomas	Colonic adenomas (diameter more than 1mm)	Small colonic adenomas (diameter less than 1mm)	Proximal small adenomas

1852	F	Barrier	5	3	9	Yes
1853	F	Barrier	3	4	0	No
1882	F	Barrier	3	2	14	Yes
1902	F	Barrier	8	4	5	No
1920	F	Barrier	2	1	5	Yes
1863	M	Barrier	12	6	7	Yes
1883	M	Barrier	11	2	20	Yes
1884	M	Barrier	14	1	28	Yes
1886	M	Barrier	10	1	28	Yes
1908	M	Barrier	21	8	4	No
1926	M	Barrier	11	2	13	Yes
1940	M	Barrier	11	2	8	Yes
1943	M	Barrier	14	2	7	Yes
1860	F	Conventional	9	3	0	No
1879	F	Conventional	6	2	2	No
1880	F	Conventional	5	6	2	No
1922	F	Conventional	4	3	1	No
1934	F	Conventional	5	3	0	No
1865	M	Conventional	23	5	4	Yes
1887	M	Conventional	17	3	3	No
1890	M	Conventional	18	6	6	No
1891	M	Conventional	14	4	5	No
1927	M	Conventional	20	6	0	No

| 1930 | M | Conventional | 24 | 7 | 0 | No |
| 1938 | M | Conventional | 16 | 5 | 0 | No |

Figure Legends

Figure 1. 16S sequencing analysis of fecal microbiota in CRASF rats pre and post shipping

(a) Relative abundance (percentages) of each operational taxonomic unit (OTU) at the Genus level is shown for the ASF rats purchased from Charles River Laboratories, before shipping and 3-months post arrival at the Discovery Ridge animal facility. Also shown is the GM profile of the F344/CRASF-*Apc*$^{+/am1137}$ (JA1047.D4) that was fostered onto a CRASF dam. Bar graphs depicting the 16S sequencing data for the ZymoBIOMICS^{TM} microbial community DNA standard is shown on the left that were used as processing and sequencing controls. *OTUs picked up after arrival and housing for 3 months. Lewis rats 1-4 were females, while rats 5-7 were males. (b) Principal Coordinate Analysis (PCoA) for the 16S rRNA sequencing data shows that Pre and Post samples (black, filled circles) of the CRASF rats are significantly different (PERMANOVA, F=6.272 and p=0.0001). The fostered *Apc*$^{+/am1137}$ rat is shown as the purple filled triangle. (c) Bar graphs representing each OTU as a single color show the relative abundance of taxa detected in the bedding and the gel-paks via 16S rRNA gene sequencing.

Figure 2. Colonic and small intestinal adenoma multiplicity of barrier and conventional rats at 4 months of age

Colonic (a) and small intestinal (b) adenoma multiplicity for male and female F1 Pirc rats from the barrier and conventional rooms is shown with adenoma counts on the y-axis and the groups on

609   the x-axis. Significance was assessed by a Student's t-test, with a p-value less than 0.05 was

610   observed. Error bars indicate standard error of the mean ($\pm$SEM).

Figure 3. Effect of Altered Schaedler Flora on the colonic adenoma phenotype and the

physiology of the gastrointestinal tract

(a) Representative proximal colon section of the (n=13) F1 Pirc rats from the barrier room. Arrows

indicate small adenomas, less than 1 mm in diameter. Scale bar = 1mm. Depicted small adenoma

sizes: 1 = 0.363mm, 2 = 0.858mm, 3 = 0.875 mm, 4 = 0.993mm, 5 = 0.969mm, and 6 = 0.378mm.

(b) Representative proximal colonic region for (n=12) conventionally-housed rats. Images were

captured on a Leica M165FC microscope with 1X magnification and a 40X objective. (c) Number

of small adenomas determined in males and females respectively in the barrier and conventional

rooms. (d) Adenoma multiplicity differences in males and females respectively were determined

by excluding the number of small adenomas seen in the F1-Pirc rats. Significance was assessed by

a Student's t-test, with a p-value less than 0.05 was observed as significant. Error bars indicate

standard error of the mean ($\pm$SEM). (e) Representative images of the cecum at weaning, from the

barrier and conventional rooms. (f) Barrier and conventional room ceca obtained at sacrifice

(representative images), indicating the difference in size between the housing conditions. Images

were captured using a Nikon D5200. A ruler is shown for comparison between groups.

Figure 4. 16S sequencing analysis of fecal microbiota in F1-Pirc rats at weaning

(a) Gut microbiota profiles of the dams and sire along with the barrier and conventionally raised

F1-Pirc rats at weaning are displayed as a bar graph depicting the relative abundance of each OTU

in percentages. Each color represents a single OTU. (b) Principal Coordinate Analysis using a

Bray-Curtis distance matrix depicts the overall similarity or dissimilarity within the groups: barrier

(brown, filled circle), conventional (blue, filled square), dams (black, filled triangle), and sire

(green, filled triangle). PERMANOVA was used for significance testing; F=1.112 and p=0.3172.

(c) The total number of OTUs observed, i.e. richness of the groups is depicted with the groups

along the x-axis and the number of OTUs along the y-axis. No significant differences were found

(ANOVA, Tukey's post hoc, $p<0.05$)

Figure 5. Effect of conventional housing on the GM at 4 months of age

(a) Bar graphs depicting the Phyla and Genus at 4-months of age from the barrier and conventional

F1-Pirc rats demonstrate the individual OTUs as a different color. (b) Genus level OTUs were used

to visualize the similarities/dissimilarities between each samples and the groups at 4-months of

age using a PCoA. PERMANOVA was used to determine significant differences with a p-value

less than 0.05. Based on the Genus data with a cutoff of 0.001% (accounting for sequencing error

rates), the richness (c), and diversity indices – Chao1 (d) and Shannon (d) were measured from the

raw read counts after normalizing the sequences to 21,639 per sample. (f) The

Firmicutes:Bacteroidetes ratio of the two housing strategies is depicted. Significance assessed by

$p<0.05$ was determined using a Student's t-test.

Figure 6. Correlation analysis of OTUs from barrier and conventional rooms with colonic

tumor count at 4 months of age

Correlation analyses was performed using *Corrgram* R package, to determine positive or negative

correlations with individual taxa at weaning in the barrier (a) and conventional (b) housing

conditions. Correlations with a significant p-value of less than 0.05 are depicted by filled circles

or diamonds. Empty cells indicate no significant correlations. Positive correlations with $r^2>0.75$

are shown as blue diamonds and as blue circles for $r^2 < 0.75$. Negative correlations are shown as

red diamonds ($r^2 > 0.75$), and red circles for < 0.75.