Wolbachia is a nutritional symbiont in Drosophila melanogaster

Amelia RI Lindsey[1*], Audrey J Parish[2], Irene LG Newton[2], Jason M Tennessen[2], Megan W

Jones[1], Nicole Stark[2]

[*]To whom correspondence should be addressed (alindsey@umn.edu)

[1]Department of Entomology, University of Minnesota, St. Paul, Minnesota, 55108

[2]Department of Biology, Indiana University, Bloomington, Indiana, 47405

ABSTRACT

The intracellular bacterium *Wolbachia* is a common symbiont of many arthropods and nematodes,

well studied for its impacts on host reproductive biology. However, its broad success as a vertically

transmitted infection cannot be attributed to manipulations of host reproduction alone. Using the

Drosophila melanogaster model and their natively associated *Wolbachia* strain "*w*Mel"*,* we show

that *Wolbachia* infection supports fly development and buffers against nutritional stress.

Wolbachia infection across several fly genotypes and a range of nutrient conditions resulted in

reduced pupal mortality, increased adult emergence, and larger size. We determined that the

exogenous supplementation of pyrimidines rescued these phenotypes in the *Wolbachia*-free, flies

suggesting that *Wolbachia* plays a role in providing this metabolite that is normally limiting for fly

growth. Additionally, *Wolbachia* was sensitive to host pyrimidine metabolism: *Wolbachia* titers

increased upon transgenic knockdown of the *Drosophila de novo* pyrimidine synthesis pathway

but not knockdown of the *de novo* purine synthesis pathway. We propose that *Wolbachia* acts as

a nutritional symbiont to supplement fly development and enhance host fitness*.*

SIGNIFICANCE STATEMENT

Wolbachia is a bacterial symbiont of arthropods and nematodes, well described for its

manipulations of arthropod reproduction. However, many have theorized there must be more to

this symbiosis, even in well-studied *Wolbachia*-host relationships such with *Drosophila*.

Reproductive impacts alone cannot explain the success and ubiquity of this bacterium. Here, we

use *Drosophila melanogaster* and their native *Wolbachia* infections to show that *Wolbachia*

supports fly development and significantly buffers flies against nutritional stress. Additionally, we

show that this advantage is likely mediated by pyrimidines. These developmental advantages

might help explain the ubiquity of *Wolbachia* infections.

INTRODUCTION

Maternally transmitted microbes have evolved numerous ways to alter host biology, ultimately facilitating their own transmission. Strategies include supplementation of the host's diet via nutrient provisioning, protection against parasites and infections, and even direct manipulation of host reproduction (1). Many insects, other terrestrial arthropods, and some nematode species are infected with bacteria in the genus *Wolbachia* (Alphaproteobacteria: Rickettsiales), a maternally transmitted infection long appreciated for its direct impacts on host reproduction and sex ratios (1-10). Importantly, *Wolbachia* can spread through populations in the absence of such reproductive manipulations (11). And, increasingly, other *Wolbachia*-mediated benefits to the host are being uncovered (12). For example, a suite of *Wolbachia* strains have been characterized for their ability to protect host insects against secondary infections, especially viruses (13). Certain strains of *Wolbachia* are obligate beneficial nutritional symbionts, such as the bed bug-infecting *Wolbachia* that produce B-vitamins to support these hematophagous insects (14). In other insects, facultative associations with *Wolbachia* have been implicated in the balance of cholesterol, iron, or B-vitamins (15-18).

We previously identified that *Wolbachia* infection results in a suite of changes to the expression of *Drosophila melanogaster* nucleotide metabolism pathways (19). Additionally, comparative genomic data indicate that the biosynthetic pathways for the *de novo* synthesis of purines and pyrimidines have been conserved across *Wolbachia*, pointing towards their broad importance (20). Furthermore, *Wolbachia* encode for an abundance of amino acid importers, which would provide necessary precursors for *Wolbachia*'s *de novo* synthesis of nucleotides (3, 20). The conservation of these metabolic pathways might simply represent essential functions for *Wolbachia*'s own metabolism, or they might also be important in the context of interaction with the host. We hypothesized that *Wolbachia* plays a supporting role in host nutrition, and this might be mediated by nucleotides. We show that *Wolbachia* infection is beneficial for flies, especially

under nutrient limited conditions, and that *Wolbachia* is responsive to changes in the expression

of fly metabolic pathways.

METHODS

<u>Fly husbandry</u>

Drosophila melanogaster stocks with or without their native *Wolbachia* strain "*w*Mel" were

maintained on standard Bloomington cornmeal-agar medium at 25 °C on a 24-hour, 12:12

light:dark cycle under density-controlled conditions and 50% relative humidity. *Wolbachia*

colonization status was confirmed with PCR assays using *Wolbachia*-specific 16S primers

WspecF and WspecR (21). Genotypes used in nutritional assays (below) included: DGRP-320

(RRID:BDSC_29654), a *Wolbachia*-infected isogenic wild-type strain with genome sequence

available which we refer to as "wild type" below (22), and a *Wolbachia*-infected white line, w^{145}

(RRID:BDSC_145). *Wolbachia*-cleared counterpart stocks were generated with antibiotics via

three generations of tetracycline treatment (20 μg/mL in the fly food for three generations),

followed by re-inoculation of the gut microbiome by transfer to bottles that previously harbored

male flies from the original stock that had fed and defecated on the media for one week (23).

Stocks were allowed to recover from any potential transgenerational effects of the antibiotic

treatment for at least an additional ten generations prior to use in any experiment and re-screened

for *Wolbachia* infection immediately prior to all experiments. All other fly stocks (e.g., RNAi lines),

are detailed in Supplementary Table S1.

<u>Nutritional stress experiments</u>

To generate poor media, Nutri-fly® Bloomington Formulation (Genesee Scientific 66-121) was

prepared according to manufacturer's instructions and diluted to indicated percentages while

maintaining the full concentrations of agar (5.3 g/L) and propionic acid (4.8 ml/L). Five ml of media

was aliquoted into each vial. Flies were placed in mating cages on grape agar plates streaked

with yeast paste and acclimated for 24 hours, prior to initiating 4-hour egg lays. Upon hatching, <

4-hour old L1 larvae were transferred to media vials (n=20 per vial), and development was scored

every 24 hours. Pupation was defined by eversion of the anterior spiracles (24). Each vial was

tracked until all pupae eclosed, or until three consecutive days without additional adult

emergence.

Gnotobiotic assays

To test if the gut microbiome was playing a role in *Wolbachia*-mediated developmental

phenotypes, we reared wild type flies with and without *Wolbachia* on 100% and 25% strength

media under gnotobiotic conditions (i.e., without the gut or food microbiome). 10 mL of media

(prepared according to protocols in "nutritional stress experiments") was aliquoted into

polypropylene wide vials, capped with a Cellulose Acetate Flug® (Flystuff 49-101), and

autoclaved on liquid setting for 20 minutes to ensure sterility. Embryos were collected on grape

agar with yeast paste as described previously and transferred into 70-micron mesh cell strainers

(Falcon 352350). Embryos in the cell strainers were washed with *Drosophila* embryo wash

solution (7% NaCl, 0.5% Triton X-100) to remove food debris. After an initial wash, embryos were

immersed in a 10% bleach solution for 3 minutes, with gentle mixing every 30 seconds. The

embryos were then washed with sterile *Drosophila* embryo wash solution to remove excess

bleach. Embryos were subjected to a final rinse with sterile phosphate buffered saline (PBS) and

transferred to a sterile agar culture plate (2% agar in deionized water) stained with blue gel food

coloring to facilitate counting. A flame-sterilized probe was used to transfer 20 embryos into each

vial of sterile media.

Nutritional supplementation experiments

To test the impact of specific nutrients, we prepared 12.5% strength media as above, and

selectively added in protein or nucleotides: the precursors and end points of *de novo* nucleotide

biosynthesis. Treatments included casein (for bulk addition of amino acids at 64.1 g/L; Sigma

C7078), inosine (0.74 g/L; Sigma I4125), uridine (0.67 g/L; Sigma U3003), or both inosine and

uridine at the aforementioned concentrations. The concentrations of additives were calculated as

87.5% of a previously defined holidic diet (25), to adjust for the 12.5% BDSC media strength.

Developmental assays were carried out as described above.

<u>Pupal size measurements</u>

After selected developmental assays, pupal casings and dead pupae were removed from vials

using a wet paint brush and transferred to glass slides. Pupae were imaged with brightfield

microscopy on an ECHO Revolve at 4X. Images were manually annotated in Echo Labs Pro

software to measure sizes. Pupal length was defined as the distance between the base of the

posterior spiracles to the midway point between the anterior spiracles. Width measurements were

taken at the widest part of the pupa. The volume of each pupa was calculated assuming a prolate

spheroid shape [$V = (4/3) \pi (width/2)^2 (length/2)$] (Supplemental Figure S1) (26, 27).

<u>Knockdown of fly nucleotide metabolism genes</u>

Wolbachia-infected Transgenic RNAi Project ("TRiP") stocks were obtained from the Bloomington

Drosophila Stock Center to knock down expression of genes in the purine and pyrimidine

nucleotide biosynthesis pathways (Supplemental File S1). Unmated females carrying a UAS-

gene-specific short hairpin were crossed to males with an Act5C-Gal4 driver (RRID:BDSC_3954:

y^1w^*; P{w^{+mC}=Act5C-GAL4}17bFO1/TM6B, Tb^1). Unmated, F1 female progeny were sorted as

adults into those with the short hairpin and Gal4-driver or those with the short hairpin and the

TM6B balancer. At three days old, the sorted flies were flash frozen in liquid nitrogen and then

stored at -80 °C for later processing. To determine the effect of the TM6B balancer on *Wolbachia*

titer, we crossed *Wolbachia*-infected, Act5C-Gal4 driver females to *Wolbachia*-uninfected UAS-

anti-Ppyr\LUC males (RRID:BDSC_31603: $y^1 v^1$; P{$y^{+t7.7} v^{+t1.8}$=TRiP.JF01355}attP2). Unmated,

F1 female progeny were sorted and stored as described for the nucleotide biosynthesis gene

knockdowns. To determine spatial differences in *Wolbachia* titer, we dissected ovaries from two-

day old *r-l* knockdown and sibling control flies. We separated ovaries from living, anesthetized

flies in sterile PBS. Ovaries and carcasses from three flies were pooled for each biological

replicate and flash frozen, followed by storage at -80 °C for later processing.

Real-time quantitative RT-PCR analyses of target gene expression

Flies were homogenized in TRIzol reagent (Invitrogen), and total RNA was extracted following

manufacturer's instructions. RNA extractions were DNase treated (RQ1 RNase-free DNase, New

England Biolabs) according to the manufacturer's instructions. cDNA was synthesized using

MMuLV Reverse Transcriptase (New England Biolabs) with random hexamer primers (Integrated

DNA Technologies). Quantitative RT-PCR reactions were performed with SensiFAST SYBR Hi-

ROX kit (Bioline) and gene-specific primers (Supplemental File S2). All samples were run in

technical duplicate alongside negative controls on an Applied Bioscience StepOnePlus qPCR

machine (Life Technologies). Gene expression was normalized to endogenous 18S rRNA

expression using the Livak method (28).

Real-time quantitative PCR analyses of *Wolbachia* titer

DNA was extracted from individual flies using the Quick-DNA/RNA Pathogen Miniprep kit (Zymo

Research) according to the manufacturer's protocol. *ftsZ* primers were used to quantify *Wolbachia*

genome copy numbers, which were normalized to host genome copies via 18S quantification

using the Livak method (28). Reactions were performed with SensiFAST SYBR Hi-ROX kit

(Bioline), and all samples and negative controls were run in technical duplicate on an Applied

Bioscience StepOnePlus qPCR machine (Life Technologies).

Nucleotide metabolism biosynthesis and gene expression

Nucleotide biosynthesis pathway information was downloaded from BioCyc by querying the

"*Wolbachia* endosymbiont of *Drosophila melanogaster w*Mel, version 26.5" and "*Drosophila*

melanogaster, version 26.5" databases (29). Enzymes for each step were cross-referenced with

KEGG Pathways, release 105.0 (30). To assess expression of *Wolbachia* metabolic pathways,

we analyzed the *Wolbachia* transcriptomes previously extracted from the MODENCODE dataset

(31, 32). We leveraged the previously published FPKM values for each *Wolbachia* gene. For

simplicity of visualization, mean FPKM is displayed for developmental time points represented by

multiple libraries. Fly gene expression for *Wolbachia*-infected and *Wolbachia* uninfected adult

females was previously reported (19).

<u>Statistics and Data Visualization</u>

Statistics and data visualization were carried out in R version 3.5.0 (33). Significant differences in

development were assessed with generalized linear mixed-effects models (package: 'lme4',

function 'glmer' (34)) including the proportion of flies that reached a given stage (pupa or adult)

as a binomial response, *Wolbachia* presence, day of development, media, and the interaction of

the three as fixed effects, and vial as a random effect to account for repeated measures.

Significant differences in pupal volume of flies reared on 12.5% media was assessed with a two-

way ANOVA (function 'aov') including *Wolbachia*, mortality status, and their interaction as fixed

effects. Significant differences in sizes of pupae reared on supplemented media was first

assessed with a two-way ANOVA (function 'aov') including *Wolbachia*, media, and their

interaction as fixed effects. Paired comparisons between *Wolbachia*-infected and uninfected flies

were then assessed with Wilcoxon rank sum tests (function 'wilcox.test'). Gene expression and

Wolbachia titers from knockdown experiments were assessed with two-way ANOVAs (function

'aov') including delta delta Ct as the response, and target locus, genotype (knockdown versus

sibling), and their interaction as fixed effects. *Wolbachia* titers in dissected flies were also

assessed with ANOVA, here with genotype, tissue, and their interaction as fixed effects. Pairwise

comparisons were performed with t-tests.

RESULTS

Wolbachia is beneficial under nutrient limited conditions

To assess the impact of *Wolbachia* infection on fly development, we reared flies on 100% strength

and 25% strength media and quantified pupation and adult emergence. We found a significant

interaction between *Wolbachia*, media strength, and time that impacted fly pupation ($F_{1,1063}$ =

24.7147, p < 0.0001) and adult emergence ($F_{1,1063}$ = 22.2897, p < 0.0001). Additionally, we found

a significant interaction between the presence of *Wolbachia* and the media strength (wild type

pupae: $F_{1,1063}$ = 0.6517, p = 0.0063; wild type adults: $F_{1,1063}$ = 0.1746, p = 0.0062; w^{145} pupae: $F_{1,711}$

= 8.8129, p = 0.0009; w^{145} adults: $F_{1,711}$ = 19.0998, p < 0.0001), along with a significant impact of

Wolbachia infection alone on fly pupation and adult emergence (wild type pupae: $F_{1,1063}$ = 0.2874,

p < 0.0001; wild type adults: $F_{1,1063}$ = 0.5375, p < 0.0001; w^{145} pupae: $F_{1,711}$ = 10.5483, p < 0.0001;

w^{145} adults: $F_{1,711}$ = 7.0718, p < 0.0001). These effects are driven by the nutrient-limited conditions

where *Wolbachia*-infected flies developed faster (1-5 days depending on genotype, Figures 1B

and 1E) and a larger percentage of flies reached adulthood. For example, while the wild type flies

with and without *Wolbachia* reared on 100% strength media reached adulthood in 82% and 81%

of cases, on the 25% strength media these were reduced to 75% and 66% respectively. The

Wolbachia-mediated advantage was more prominent for a second fly genetic background, w^{145}.

Even on 100% strength media, the *Wolbachia*-free w^{145} flies experienced a 5% reduction in adult

emergence relative to *Wolbachia*-infected flies (Figure 1D). When reared on 25% strength media,

the *Wolbachia*-infected w^{145} flies were more than twice as likely to reach adulthood (70% versus

31%, Figure 1E). Across these assays, we found delays in entering pupation are approximately

equal to the delay in adult emergence: *i.e.,* the time spent in metamorphosis did not change as a

factor of *Wolbachia* infection.

Given that the effect size for the *Wolbachia* advantage was genotype dependent and given the

more subtle impact of *Wolbachia* in the wild-type flies, we wondered if an even weaker media

would result in a stronger advantage for the *Wolbachia*-infected flies. Conversely, *Wolbachia*

could be a burden under more significant nutritional stress. Indeed, when flies were raised on

12.5% strength media, we again saw significant effects of the interaction between *Wolbachia* and

time on the rate and success of pupation and adult emergence (pupae: $F_{1,347}$ = 77.135, p < 0.0001;

adults: $F_{1,347}$ = 58.9296, p < 0.0001). However, *Wolbachia*-infected flies now developed slower

than their uninfected counterparts (6-7 days delayed), but they pupated at higher rates (68%

versus 61%) and were more likely to reach adulthood (54% versus 40%, Figure 1C). Furthermore,

although *Wolbachia*-infected fly development was slower, it resulted in them attaining, on

average, a 23% larger size at pupation ($F_{1,191}$ = 32.076, p < 0.0001, Figure 1F). Additionally,

Wolbachia-infected flies experienced lower levels of pupal mortality (35% versus 19%), perhaps

related to the finding that flies that died during pupation were significantly smaller ($F_{1,191}$ = 4.625,

p = 0.0328, Figure 1F).

To determine if the developmental advantage was due to direct effects of *Wolbachia* or indirect

effects via *Wolbachia*-mediated impacts on the gut microbiome, we performed the same

developmental assay under gnotobiotic conditions (i.e., without a gut or food microbiome). Under

gnotobiotic conditions, we saw a strong interactive effect of *Wolbachia* and time that resulted in

significantly more, and faster, pupation and adult emergence for infected flies (Figure 1G; pupae:

$F_{1,1920}$ = 153.3796, p < 0.0001; adults: $F_{1,1920}$ = 129.6080, p < 0.0001). Even on the 100% strength

media, only 43% of *Wolbachia*-free flies pupated, compared to 61% of *Wolbachia*-infected flies.

Again, *Wolbachia*-free flies had higher levels of pupal mortality: 21% of those that pupated did

not emerge as adults, as compared to 13% of the *Wolbachia*-infected pupae. While removing the

microbiota negatively impacted fly developmental timing as expected, this effect was exacerbated

in the *Wolbachia*-free flies (Figures 1A and 1G). As compared to the conventionally-reared flies

on 100% strength media (Figure 1A), gnotobiotic *Wolbachia*-infected flies experienced a two-day

developmental delay, and *Wolbachia*-free flies were delayed three days (Figure 1G). The

combination of gnotobiotic conditions and 25% strength media resulted in especially impaired fly

development: less than 20% of flies reached adulthood (Figure 1H). While there were no

significant differences in the proportion of flies that pupated or emerged as adults between

Wolbachia-infected and uninfected flies in these high-mortality conditions (pupae: p = 0.9547;

adults: p = 0.6210), it is notable that *Wolbachia*-infected flies on average started emerging five

days ahead of the *Wolbachia*-free flies (Figure 1H). Finally, we measured the puparia of the

gnotobiotic flies reared on 100% strength media (Figure 1I) and found that *Wolbachia* infection

resulted in, on average, 10% larger pupal volumes ($F_{1,200}$ = 11.69, p = 0.0008, Figure 1I), again

indicating a significant developmental advantage.

Figure 1. *Wolbachia* infection is beneficial under nutrient limited conditions. Flies (wild type

and *w*[145], with and without *Wolbachia*) were reared on different concentrations of media to test

the impact of *Wolbachia* infection. Biological replicates included 20 larvae per each of 10 vials.

Dotted and solid lines indicate cumulative pupation and adult emergence, respectively, with

shaded regions defining standard error. All *Wolbachia*-infected treatments are in red, and

Wolbachia-uninfected in grey. **(A)** Conventionally reared wild type flies on 100%, **(B)** 25%, and

(C) 12.5% strength media. **(D)** Conventionally reared w^{145} flies on 100% and **(E)** 25% strength

media. **(F)** Pupal volumes of wild type flies with and without *Wolbachia*, derived from the 12.5%

media-reared flies in (C). **(G)** Gnotobiotic wild type flies on 100% and **(H)** 25% strength media. **(I)**

Pupal volumes of the gnotobiotic-reared flies on 100% strength media derived from (G). Orange

datapoints (H-I) indicate pupae that did not eclose into adults. *** $p < 0.001$

<u>Dynamics of the redundant *de novo* nucleotide biosynthesis pathways</u>

We next considered the role that *Wolbachia* may be playing in supporting fly development during

these nutritionally limited conditions. We previously found that nucleotide metabolism is a

significant point of interaction for flies and *Wolbachia* (19), pathways known to significantly impact

fly growth and development (35). Importantly, the pathways for the *de novo* synthesis of purines

and pyrimidines are conserved across all *Wolbachia*, and they are redundantly encoded by the

fly (Figure 2A-B). Additionally, we previously showed that *Wolbachia* infection results in

downregulation of the fly's pyrimidine synthesis enzymes *rudimentary* (-0.273 Fold Change (FC)

compared to uninfected flies), *Dhod* (-0.164 FC), and *rudimentary-like* (-0.301 FC) (19). Strikingly,

the repressor *su(r)* is upregulated (+0.505 FC)(Figure 2C). In contrast, purine metabolism overall

was not obviously distinct between *Wolbachia*-infected and uninfected flies: some enzymes were

up- or down- regulated but not the entire pathway (Figure 2C).

Given the developmental effects of *Wolbachia* infection, we hypothesized that *Wolbachia*

nucleotide metabolic pathways would be more highly expressed during growth stages (*i.e.,* pre-

metamorphosis). Indeed, from embryogenesis through the third larval instar (L3), most of

Wolbachia's nucleotide metabolism genes were upregulated. At the onset of pupation (the white

pre-pupae "WPP" stage), *Wolbachia*'s metabolic gene expression was sharply and broadly

downregulated (Figure 2D). These data indicate that *Wolbachia* may be supplementing host

nutrition during critical growth stages, and either limiting negative impacts of infection during

metamorphosis and reproductive stages or shifting roles to better interface with host reproduction

and ensure transmission.

Figure 2. Flies and *Wolbachia* redundantly encode for *de novo* nucleotide biosynthesis.

(A) Metabolic pathway for *de novo* pyrimidine biosynthesis in *Wolbachia* (left side, purple bacterial

schematic) and *Drosophila melanogaster* (right side, fly schematic). **(B)** Metabolic pathway for *de*

novo purine biosynthesis in *Wolbachia* (left side or bottom, purple bacterial schematic) and

Drosophila melanogaster (right side or top, fly schematic). **(C)** Expression of *Drosophila*

melanogaster nucleotide biosynthesis loci in *Wolbachia*-infected female flies relative to

Wolbachia-uninfected controls, derived from published expression data (19). **(D)** Expression of

Wolbachia nucleotide biosynthesis loci across fly development. *Wolbachia* transcriptomes were

generated as part of the MODENCODE project (31, 32). Data are row normalized for inference of

per-gene transcriptional changes across development. Fly images are from BioRender.com.

Abbreviations: UMP: uridine monophosphate; IMP: inosine monophosphate. Fly gene alternate

names: Ade2: Pfas; Ade3: Gart; Ade5: Paics.

<u>Pyrimidines contribute to the *Wolbachia* infection advantage</u>

We reasoned that deficiencies in the diet supplemented by *Wolbachia* could be supplemented by

exogenous nutrient addition to the fly food. Therefore, flies with and without *Wolbachia* would

have the same growth parameters if this important nutrient was added. Specifically, we focused

on the precursors (amino acids) and end-products (uridine and inosine) of the *de novo* nucleotide

synthesis pathways, which we supplemented back into the 12.5% media. The developmental

assays on supplemented media were repeated with the wild type stocks, avoiding w^{145} because

the *white* transporter is known to affect nucleotide metabolism (36). We again saw the *Wolbachia*-

mediated developmental advantage in the 12.5% strength media (pupae: $F_{1,235}$ = 13.9505, p =

0.0002; adults: $F_{1,235}$ = 12.7568, p = 0.0004). Additionally, we saw a subtle but significant positive

impact of *Wolbachia* in the 100% strength media as measure by adult emergence (pupae: $F_{1,235}$

= 1.3002, p = 0.2538; adults: $F_{1,235}$ = 28.0455, p < 0.0001), due to 6% pupal mortality in the

Wolbachia-free flies that the *Wolbachia*-infected flies did not experience (Figure 3A). While the

addition of casein benefited both *Wolbachia*-infected and uninfected flies, *Wolbachia*-free flies

pupated at significantly increased rates relative to *Wolbachia*-infected flies ($F_{1,235}$ = 9.8699, p =

0.0017). Again, pupal mortality was quite high in the *Wolbachia*-free flies (36% as compared to

9% with *Wolbachia*), resulting in significantly fewer numbers of adults ($F_{1,235}$ = 6.9988, p = 0.0082).

The addition of uridine (product of *de novo* pyrimidine synthesis) specifically benefited *Wolbachia*-

free flies, but not the flies that had *Wolbachia* infections. Indeed, uridine supplementation resulted

in a rescue phenotype where equivalent numbers of *Wolbachia*-infected and uninfected flies

emerged as adults (p = 0.9281). Importantly, emergence of *Wolbachia*-free flies supplemented

with uridine was not significantly different than the *Wolbachia*-infected flies on the standard 12.5%

strength media (p = 0.9097) (Figure 3A). In fact, *Wolbachia*-free flies supplemented with uridine

developed slightly faster than their infected counterparts (pupae: $F_{1,235}$ = 2.2401, p = 0.1348;

adults: $F_{1,235}$ = 10.5842, p = 0.0011). The addition of inosine (product of *de novo* purine synthesis)

was toxic and reduced pupation and emergence for all flies. As such there were no significant

differences between *Wolbachia*-infected and uninfected flies in these high mortality (*i.e.*, zero-

inflated) conditions (pupae: $F_{1,235}$ = 0.9217, p = 0.335; adults: $F_{1,235}$ = 0.3519, p = 0.553).

Simultaneously adding uridine and inosine resulted in more intermediate, albeit still quite lethal

developmental phenotypes with no significant impact of *Wolbachia* ($F_{1,235}$ = 1.3471, p = 0.2459;

adults: $F_{1,235}$ = 1.0400, p = 0.307).

Congruent with the previous developmental assays (Figure 1), *Wolbachia* infection had a

significant impact on pupal size, both due to interacting effects with the media ($F_{5,470}$ = 2.55, p =

0.0272), and *Wolbachia* alone ($F_{1,470}$ = 37.31, p < 0.0001) (Figure 3B). Furthermore, *Wolbachia*-

infected flies were significantly larger when reared on 100% (p = 0.0401), 12.5% (p = 0.0316),

and casein-enriched 12.5% media (p < 0.0001). However, the addition of uridine to the 12.5%

media resulted in similarly sized flies regardless of *Wolbachia* infection (p = 0.3685). In summary,

Wolbachia-free flies were rescued to the same size and level of adult emergence as *Wolbachia*-

infected counterparts in uridine (i.e., pyrimidine) supplemented media.

Figure 3. Uridine rescues *Wolbachia*-free flies. Wild type flies were reared on 100% media, 12.5% media, or 12.5% media containing either casein, uridine, inosine, or both inosine and uridine. The media designations in the middle correspond to the figures above and below. Biological replicates included 20 larvae per each of five vials. **(A)** Dotted and solid lines indicate cumulative pupation and adult emergence, respectively, with shaded regions defining standard error. *Wolbachia*-infected treatments are in red and *Wolbachia*-uninfected are in grey. **(B)** Pupae from (A) were removed post-developmental assay, measured, and their volume was calculated. Orange datapoints indicate pupae that did not eclose into adults. Significance annotations for

pupal sizes comparisons are the same regardless of whether flies that died during pupation were

included. ***p < 0.001; **p < 0.01; *p < 0.05; ns = not significant.

Wolbachia infection is responsive to fly pyrimidine biosynthesis gene expression

To determine whether *Wolbachia* infection was sensitive to changes in host metabolism, we

knocked down individual genes within the fly's *de novo* nucleotide synthesis pathways (Figure

2A-B) using *Wolbachia*-infected RNAi fly stocks (Figure 4A). Short-hairpin RNA (shRNA)

expression was driven in the entire body of the fly by an Act5C-Gal4 driver. We compared gene

expression and *Wolbachia* titer in knockdown flies to sibling controls (flies not expressing a gene-

targeting shRNA, and instead with the TM6B balancer). We successfully knocked down three

genes in the pyrimidine synthesis pathway (*su(r)*, *Dhod*, *rudimentary-like*) and four genes in the

purine synthesis pathway (*prat2*, *ade2*, *ade5*, *CG11089*) (Figure 4A, Supplemental Figure S2).

Several gene knockdowns were lethal to fly development, including *rudimentary* in the de novo

pyrimidine biosynthesis pathway and *CG2246*, *CG6767*, *Prat*, and *AdSL* in the de novo purine

biosynthesis pathway. Broad-spectrum knockdown of *de novo* purine synthesis had no significant

impacts on *Wolbachia* titer (gene*knockdown: $F_{3,31}$ = 0.652, p = 0.588; gene: $F_{3,31}$ = 0.644, p =

0.593; knockdown: $F_{1,31}$ = 0.593, p = 0.447; Supplemental Figure S2). In contrast, knockdown of

the *de novo* pyrimidine synthesis pathway resulted in significant increases in *Wolbachia* titer ($F_{1,38}$

= 10.3916, p = 0.0026; Figure 4B). This pattern was specifically driven by the enzymatic loci

(*Dhod*: p = 0.034 and *rudimentary-like* (*r-l*)*:* p = 0.013). In contrast, knockdown of the negative

regulator *sur(r)* had no effect on *Wolbachia* titer (p = 0.530), perhaps indicating that *Wolbachia*

titers are responsive to low fly pyrimidine synthesis activity, but not elevated levels. To account

for any effects of the TM6B balancer on measurements of *Wolbachia* abundance, we crossed the

Act5C-Gal4 driver to a RNAi line expressing shRNA against a foreign gene: *Photinus pyralis*

luciferase (UAS-anti-Ppyr\LUC). We found no significant changes in *Wolbachia* titer, or in

nucleotide gene expression that could be driving the results in Figure 4B (Supplemental Figure

S3). We then asked if the change in *Wolbachia* titer was tissue-specific by measuring the relative

Wolbachia titers in ovaries and the remaining carcasses after knockdown of *r-l* (Figure 4C). There

was no significant interaction of knockdown and tissue ($F_{1,32}$ = 0.20, p = 0.658) or of tissue alone

($F_{1,32}$ = 0.20, p = 0.658) on *Wolbachia* titers. We again saw a significant effect of *r-l* knockdown

on *Wolbachia* titers ($F_{1,32}$ = 13.07, p = 0.001) both in ovaries (p = 0.007) and in carcasses (p =

0.0127) (Figure 4C). Finally, given these interactions with pyrimidine synthesis, we used flies with

a homozygous hypomorphic allele *Dhod²* (RRID:BDSC_1936, Supplemental Table S1) to assess

the importance of *Wolbachia* in a pyrimidine synthesis-impaired genetic background. This stock

is naturally infected with *Wolbachia*, but removal of the symbiont lead to significant delays and

reductions in pupation and adult emergence (Figure 4D; pupae: $F_{1,247}$ = 11.0058, p = 0.0009;

adults: $F_{1,247}$ = 12.0550, p = 0.0057).

Figure 4. *Wolbachia* titers are responsive to fly pyrimidine synthesis gene expression.

Expression of fly genes in the *de novo* pyrimidine pathway was constitutively knocked down using

standard transgenic approaches. **(A)** *Drosophila* gene knockdown was verified with qRT-PCR

leveraging gene specific primers and normalization to 18S expression. **(B)** Relative *Wolbachia*

titers in whole adult females and **(C)** dissected ovaries and carcasses were quantified with qPCR

using *Wolbachia*-specific primers and normalization to fly genome copy number via 18S

amplification. Legend: *** $p <0.001$; ** $p < 0.01$; * $p < 0.05$; ns p = not significant. **(D)** Development

of conventionally reared *Wolbachia*-infected and -uninfected $Dhod^2$ p^2 flies on 100% strength

media. Biological replicates included 20 larvae per each of ten vials. Dotted and solid lines indicate

cumulative pupation and adult emergence, respectively, with shaded regions defining standard

error. *Wolbachia*-infected flies are in red and *Wolbachia*-uninfected are in grey.

DISCUSSION

Wolbachia has long been appreciated for its diverse impacts on a range of ecdysozoan hosts.

There has been increasing speculation as to the role *Wolbachia* plays in host metabolism more

broadly, often times in the context of specific metabolites (12, 19, 20). For example, *Wolbachia*-

infection contributes to B-vitamin supplementation in a planthopper (18), bed bugs (14), and likely

some species of solitary bees (37). Iron has been the subject of a number of investigations into

Wolbachia-host interactions (15). In *Drosophila melanogaster, Wolbachia* infection buffers

against stresses associated with low or high concentrations of iron in the diet (38). *Wolbachia*

infection is additionally associated with changes in host ferritin expression and iron homeostasis

in other insects such as *Drosophila simulans* and the parasitoid wasp *Asobara tabida* (16). Given

the range of metabolites and physiological processes that have been implicated in *Wolbachia*-

host interactions, we focused initial experiments on using a more holistic approach to

understanding fly development in the context of diet and *Wolbachia* infection.

Here, we show that *Wolbachia* significantly impacts *Drosophila melanogaster* development,

resulting in larger flies more likely to reach adulthood. The strength of this phenotype varied by

fly genotype and the overall concentration of nutrition in the food: *Wolbachia* infection was highly

beneficial under nutrient limited conditions. Removal of the extracellular microbial community

indicated that this advantage was directly mediated by *Wolbachia*. In fact, *Wolbachia* significantly

buffered flies against the stresses normally associated with removal of the microbiome (39, 40).

Furthermore, *Wolbachia* titers were responsive to changes in the expression of fly pyrimidine

biosynthesis, and the supplementation of pyrimidines rescued adult emergence of the *Wolbachia*-

uninfected flies. We propose that pyrimidines are a significant component of this developmental

advantage provided by *Wolbachia*. Indeed, the patterns of *Wolbachia* gene expression seem to

be attuned to fly development which could be a strategy for supporting the exponential growth

that occurs during larval stages and mitigating fitness impacts of the infection during reproductive

stages. Importantly, while the transcription of *Wolbachia*'s nucleotide metabolism pathways are

quite high pre-pupation, the flies have comparatively low *Wolbachia* titers during that time (31).

Genomic data suggest that *Wolbachia* imports amino acids from the host: while these may be

used as precursors for nucleotide biosynthesis, the depletion of free amino acids could induce

host stress responses (13, 41). Changes in *Wolbachia* titer and physiology across development

and under different nutritional scenarios likely reflect such balancing acts.

While *Wolbachia* infections have received considerable attention for their impacts on

reproduction, we know much less about how *Wolbachia* interact with their host during juvenile

stages. Importantly, many insects spend the majority of their lifetime as juveniles. Juvenile insect

metabolism is typically quite distinct from adults, and this time period has profound impacts on

metamorphosis, adult physiology, and fitness (42). Critically, all insect growth occurs during the

larval period: in *Drosophila melanogaster,* larvae increase in body mass by ~200-fold over the

course of 4 days (42). We found that *Wolbachia*-free flies had higher levels of pupal mortality,

especially during nutritional stress, suggesting insufficient metabolic reserves. This effect was

particularly pronounced when flies were reared on an excess of protein (Figure 3). Importantly,

amino acids are a key nutritional cue for determining the timing of pupation (i.e., signaling critical

weight) due to a direct and indirect effects on the timing and intensity of ecdysone pulses (43-46).

An influx of amino acids without other essential nutrients needed for growth might impact

nutritional signaling and thus the timing of critical weight, perhaps triggering flies to pupate early

without sufficient reserves for metamorphosis. Given *Wolbachia* likely imports amino acids, this

may reduce the concentrations of free amino acids and delay signaling critical weight, allowing

flies to accumulate more metabolic reserves before pupation.

During the ~200-fold increase in larval body mass, there is a concomitant 25-fold increase in DNA

and RNA content that creates a huge demand for pyrimidines, well exceeding the fly's

endogenous biosynthetic capacity (47). Considering that the rate of *Drosophila* development is

highly sensitive to *de novo* pyrimidine synthesis (47), bacterially-derived nucleotides could give

the host a significant advantage. Pyrimidines are synthesized by a series of metabolic reactions

that use ribose sugar and the amino acids glutamine and aspartate to generate uridine-5'-

monophosphate, UMP (48): the precursor for uracil, cytosine, and thymine. This pathway is

conserved in both *Drosophila* and *Wolbachia*, suggesting that infected flies generate pyrimidines

in multiple compartments: the host cytoplasm/mitochondria and *Wolbachia*. *Wolbachia*'s

production of pyrimidines could also represent far more than simply additional biosynthesis, as

the enzyme dihydroorotate dehydrogenase (*Dhod*) is located in the inner mitochondrial

membrane where it couples pyrimidine synthesis to electron transport (49, 50). As such,

provisioning of *Wolbachia*-generated pyrimidines could reduce the biosynthetic burden normally

imposed upon mitochondria. Additionally, fly enzymes used for pyrimidine synthesis, CAD and

UMP synthase, are negatively regulated by UMP (51). Therefore, *Wolbachia* could simultaneously

impact free amino acid pools, supplement pyrimidines, reduce mitochondrial burdens, and

spatially segregate components of a normally inhibitory interaction.

While we found that pyrimidines seemed to be important for the *Drosophila-Wolbachia* symbiosis,

we did not see any changes in *Wolbachia* titer due to suppression of purine synthesis. In fact, the

addition of inosine to media negatively impacted flies (Figure 3). Purine nucleotides act as

signaling molecules that impact a range of functions. For example, activation of the adenosine

receptor globally suppresses fly metabolism (52), which can cause insecticidal effects (53).

Another possibility is misincorporation of inosine into RNAs. In mammalian cell lines, this can

happen when inosine is in excess, ultimately impacting translation (54). While altering the

expression of fly purine metabolic gene expression did not impact the relative abundance pf

Wolbachia in our experiments, we cannot not rule out the potential for other changes in *Wolbachia*

gene expression or physiology. Indeed, we previously found an interactive effect of *Wolbachia*

infection and purine gene expression on virus replication (19), highlighting both the importance of

purines to the symbiosis and the complexity of the metabolic landscape.

The role of *Wolbachia* as a nutritional symbiont raises numerous questions about the cell biology

of *Wolbachia* and how the relationship with the host is regulated. Is *Wolbachia* specifically

providing pyrimidines (*e.g.,* via nucleotide transporters)? Or are flies farming and consuming their

Wolbachia infections (*e.g.,* via autophagy) and pyrimidines so happen to be abundant in

Wolbachia cells and also a limiting metabolite for the fly? As is the case for many symbioses, it is

not always clear if a change in the symbiont (*e.g.,* gene expression or titer), is due to the microbe

detecting and responding to the host's status, or the host releasing control of the microbe and

allowing a change in microbial physiology. *Wolbachia* do encode for a number of proteins that

would allow them to detect their environment and regulate gene expression accordingly (55).

However, each *Wolbachia* is enclosed by a host-derived vesicle (56, 57), so perhaps the host

tightly regulates *Wolbachia*'s "experience".

While beneficial nutritional symbioses in insects are not rare, most of the systems for which this

is described are of a much more binary nature. Insects that exclusively feed on unbalanced diets

(e.g., blood, plant sap) have obligate microbial symbionts to synthesize metabolites that the host

cannot generate *de novo*: the partners complement each other (58, 59). Here, both flies and

Wolbachia have their own complete pathways for nucleotide biosynthesis, which perhaps has led

us to overlook the relevance of this biosynthetic redundancy. Our finding that *Wolbachia*

significantly impacts the developmental timing, size, and susceptibility to nutritional stress in a

well-studied model such as *Drosophila melanogaster* raises the question of the broader relevance

of *Wolbachia* infections to insect metabolism and development.

DECLARATIONS

<u>Acknowledgements</u>

This work was supported by UMN AGREETT startup funds to ARIL. ILGN in supported by the

National Institute of Allergy and Infectious Diseases of the National Institutes of Health under R01

AI144430. JMT is supported by the National Institute of General Medical Sciences of the National

Institutes of Health under a R35 Maximizing Investigators' Research Award (MIRA;

1R35GM119557). The authors thank Richard Hardy for support and feedback. Research was

further made possible by Flybase (NIH 5U41HG000739), and stocks obtained from the

Bloomington *Drosophila* Stock Center (NIH P40OD018537).

<u>Conflicts of interest</u>

The authors declare that they have no competing interests.

<u>Data Availability</u>

Supplemental Table S1. Fly stocks

Supplemental Table S2. Primers

Supplemental Figure S1. Pupal measurement.

Supplemental Figure S2. Purine knockdowns.

Supplemental Figure S3. Knockdown controls.

REFERENCES

1.   R. Kaur *et al.*, Living in the endosymbiotic world of *Wolbachia*: A centennial review. *Cell*

Host & Microbe (2021).

2.   O. Duron *et al.*, The diversity of reproductive parasites among arthropods: *Wolbachia* do

not walk alone. *BMC Biol.* **6**, 1-12 (2008).

3.   M. Wu *et al.*, Phylogenomics of the reproductive parasite *Wolbachia* pipientis *w*Mel: A

streamlined genome overrun by mobile genetic elements. *PLoS Biol.* **2**, 327-341 (2004).

4.   J. H. Werren, W. Zhang, L. R. Guo, Evolution and phylogeny of *Wolbachia* - reproductive

parasites of arthropods. *Proc. R. Soc. Lond. B* **261**, 55-63 (1995).

5.   N. Bensaadi-Merchermek, J.-C. Salvado, C. Cagnon, S. Karama, C. Mouchès,

Characterization of the unlinked 16S rDNA and 23S-5S rRNA operon of *Wolbachia*

pipientis, a prokaryotic parasite of insect gonads. *Gene* **165**, 81-86 (1995).

6.   R. Zug, P. Hammerstein, Still a host of hosts for *Wolbachia*: Analysis of recent data

suggests that 40% of terrestrial arthropod species are infected. *PLoS One* **7** (2012).

7.   K. Hilgenboecker, P. Hammerstein, P. Schlattmann, A. Telschow, J. H. Werren, How

many species are infected with *Wolbachia*? - A statistical analysis of current data. *FEMS*

Microbiol. Lett. **281**, 215-220 (2008).

8.   F. Rousset, D. Bouchon, B. Pintureau, P. Juchault, M. Solignac, *Wolbachia*

endosymbionts responsible for various alterations of sexuality in arthropods. *Proc. R. Soc.*

Lond. B **250**, 91-98 (1992).

9.   K. Fenn, M. Blaxter, *Wolbachia* genomes: revealing the biology of parasitism and

mutualism. *Trends Parasitol.* **22**, 60-65 (2006).

10. R. Stouthamer, J. A. J. Breeuwer, G. D. D. Hurst, *Wolbachia pipientis*: Microbial

manipulator of arthropod reproduction. *Annu. Rev. Microbiol.* **53**, 71-102 (1999).

11. C. A. Hamm *et al.*, *Wolbachia* do not live by reproductive manipulation alone: infection

polymorphism in *Drosophila suzukii* and *D. subpulchrella*. *Mol. Ecol.* **23**, 4871-4885

(2014).

12. R. Zug, P. Hammerstein, Bad guys turned nice? A critical assessment of *Wolbachia*

mutualisms in arthropod hosts. *Biological Reviews* **90**, 89-111 (2015).

13. A. R. I. Lindsey, T. Bhattacharya, I. L. G. Newton, R. W. Hardy, Conflict in the intracellular

lives of endosymbionts and viruses: A mechanistic look at *Wolbachia*-mediated pathogen-

blocking. *Viruses* **10**, 141 (2018).

14. T. Hosokawa, R. Koga, Y. Kikuchi, X. Y. Meng, T. Fukatsu, *Wolbachia* as a bacteriocyte-

associated nutritional mutualist. *Proc. Natl. Acad. Sci.* **107**, 769-774 (2010).

15. A. C. Gill, A. C. Darby, B. L. Makepeace, Iron necessity: the secret of *Wolbachia*'s

success? *PLoS Negl. Trop. Dis.* **8**, e3224 (2014).

16. N. Kremer *et al.*, *Wolbachia* interferes with ferritin expression and iron metabolism in

insects. *PLoS Path.* **5**, e1000630 (2009).

17. V. Geoghegan *et al.*, Perturbed cholesterol and vesicular trafficking associated with

dengue blocking in *Wolbachia*-infected *Aedes aegypti* cells. *Nat. Commun.* **8**, 526 (2017).

18. J.-F. Ju *et al.*, *Wolbachia* supplement biotin and riboflavin to enhance reproduction in

planthoppers. *The ISME journal* **14**, 676-687 (2020).

19.   A. R. Lindsey, T. Bhattacharya, R. W. Hardy, I. L. Newton, *Wolbachia* and virus alter the

host transcriptome at the interface of nucleotide metabolism pathways. *mBio* **12** (2021).

20.   I. L. Newton, D. W. Rice, The Jekyll and Hyde symbiont: could *Wolbachia* be a nutritional

mutualist? *J. Bacteriol.* **202** (2020).

21.   J. H. Werren, D. M. Windsor, *Wolbachia* infection frequencies in insects: Evidence of a

global equilibrium? *Proc. R. Soc. Lond. B* **267**, 1277-1285 (2000).

22.   W. Huang *et al.*, Natural variation in genome architecture among 205 *Drosophila*

melanogaster Genetic Reference Panel lines. *Genome Res.* **24**, 1193-1208 (2014).

23.   M. W. Jones, L. C. Fricke, C. J. Thorpe, L. O. V. Esch, A. R. I. Lindsey, Infection dynamics

of cotransmitted reproductive symbionts are mediated by sex, tissue, and development.

Appl. Environ. Microbiol. **88**, e00529-00522 (2022).

24.   S. Bainbridge, M. Bownes, Staging the metamorphosis of *Drosophila melanogaster*.

Journal of Embryology and Experimental Morphology **66**, 57-80 (1981).

25.   T. Reis, Effects of synthetic diets enriched in specific nutrients on *Drosophila*

development, body fat, and lifespan. *PLoS One* **11**, e0146758 (2016).

26.   F. Demontis, N. Perrimon, Integration of Insulin receptor/Foxo signaling and dMyc activity

during muscle growth regulates body size in *Drosophila*. (2009).

27.   L. Moss-Taylor, A. Upadhyay, X. Pan, M.-J. Kim, M. B. O'Connor, Body size and tissue-

scaling is regulated by motoneuron-derived activinß in *Drosophila melanogaster*. *Genetics*

213, 1447-1464 (2019).

28.   K. J. Livak, T. D. Schmittgen, Analysis of relative gene expression data using real-time

quantitative PCR and the 2− ΔΔCT method. *methods* **25**, 402-408 (2001).

29.   R. Caspi *et al.*, The MetaCyc database of metabolic pathways and enzymes and the

BioCyc collection of pathway/genome databases. *Nucleic Acids Res.* **44**, D471-D480

(2016).

30.   M. Kanehisa, S. Goto, KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids*

Res. **28**, 27-30 (2000).

31.   F. Gutzwiller *et al.*, Dynamics of *Wolbachia pipientis* gene expression across the

Drosophila melanogaster life cycle. *G3: Genes| Genomes| Genetics* **5**, 2843-2856 (2015).

32.   B. R. Graveley *et al.*, The developmental transcriptome of *Drosophila melanogaster*.

Nature **471**, 473-479 (2011).

33.   R Core Team (2014) R: A language and environment for statistical computing. (URL

http://www.R-project.org/, R Foundation for Statistical Computing, Vienna, Austria).

34.   D. Bates, M. Maechler, B. Bolker, S. Walker, lme4: Linear mixed-effects models using

Eigen and S4. *R package version* **1** (2014).

35.   D. Falk, R. Bellamy, M. El Kouni, F. Naguib, A genetic and dietary study of the physiology

of pyrimidine synthesis in *Drosophila melanogaster*. *J. Insect Physiol.* **26**, 735-740 (1980).

36.   A. Sasaki, T. Nishimura, T. Takano, S. Naito, S. K. Yoo, *white* regulates proliferative

homeostasis of intestinal stem cells during ageing in *Drosophila*. *Nature Metabolism* **3**,

546-557 (2021).

37.   M. Gerth, C. Bleidorn, Comparative genomics provides a timeframe for *Wolbachia*

evolution and exposes a recent biotin synthesis operon transfer. *Nat. Micro.* **2**, 16241

(2016).

38.   J. C. Brownlie *et al.*, Evidence for metabolic provisioning by a common invertebrate

endosymbiont, *Wolbachia pipientis*, during periods of nutritional stress. *PLoS Path.* **5**,

e1000368 (2009).

39.   N. A. Broderick, B. Lemaitre, Gut-associated microbes of *Drosophila melanogaster*. *Gut*

Microbes **3**, 307-321 (2012).

40.   D. N. Lesperance, N. A. Broderick, Microbiomes as modulators of *Drosophila*

melanogaster homeostasis and disease. *Current Opinion in Insect Science* **39**, 84-90

(2020).

41.   H. P. Harding *et al.*, An integrated stress response regulates amino acid metabolism and
resistance to oxidative stress. *Mol. Cell* **11**, 619-633 (2003).

42.   J. M. Tennessen, C. S. Thummel, Coordinating growth and maturation—insights from
Drosophila. Curr. Biol. **21**, R750-R757 (2011).

43.   S. Layalle, N. Arquier, P. Léopold, The TOR pathway couples nutrition and developmental
timing in *Drosophila. Dev. Cell* **15**, 568-577 (2008).

44.   C. Géminard, E. J. Rulifson, P. Léopold, Remote control of insulin secretion by fat cells in
Drosophila. *Cell Metabolism* **10**, 199-207 (2009).

45.   J. Colombani *et al.*, A nutrient sensor mechanism controls *Drosophila* growth. *Cell* **114**,
739-749 (2003).

46.   M. J. Texada, T. Koyama, K. Rewitz, Regulation of body size and growth control. *Genetics*
216, 269-313 (2020).

47.   D. Falk, R. Bellamy, M. el Kouni, N. F, A genetic and dietary study of the physiology of
pyrimidine synthesis in *Drosophila melanogaster. J. Insect Physiol.* **26**, 735-740 (1980).

48.   D. R. Evans, H. I. Guy, Mammalian pyrimidine biosynthesis: fresh insights into an ancient
pathway. *J. Biol. Chem.* **279**, 33035-33038 (2004).

49.   J.-J. Chen, M. E. Jones, The cellular location of dihydroorotate dehydrogenase: relation
to de novo biosynthesis of pyrimidines. *Archives of Biochemistry and Biophysics* **176**, 82-
90 (1976).

50.   J. M. Rawls, Genetic complementation and enzyme correlates at the locus encoding the
last two steps of de novo pyrimidine biosynthesis in *Drosophila melanogaster. Molecular*
and General Genetics MGG **184**, 174-179 (1981).

51.   M. Löffler, E. A. Carrey, W. Knecht, The pathway to pyrimidines: The essential focus on
dihydroorotate dehydrogenase, the mitochondrial enzyme coupled to the respiratory
chain. *Nucleosides Nucleotides Nucl. Acids* **39**, 1281-1305 (2020).

52.   A. Bajgar, T. Dolezal, Extracellular adenosine modulates host-pathogen interactions

through regulation of systemic metabolism during immune response in Drosophila. *PLoS*

Path. **14**, e1007022 (2018).

53.   M. Fang, Y. Chai, G. Chen, H. Wang, B. Huang, N6-(2-hydroxyethyl)-adenosine exhibits

insecticidal activity against Plutella xylostella via adenosine receptors. *PLoS One* **11**,

e0162859 (2016).

54.   J. H. Schroader *et al.*, Disease-associated inosine misincorporation into RNA hinders

translation. *Nucleic Acids Res.* **50**, 9306-9318 (2022).

55.   A. R. I. Lindsey, Sensing, Signaling, and Secretion: A review and analysis of systems for

regulating host interaction in *Wolbachia*. *Genes* **11**, 813 (2020).

56.   K.-O. Cho, G.-W. Kim, O.-K. Lee, *Wolbachia* bacteria reside in host Golgi-related vesicles

whose position is regulated by polarity proteins. *PLoS One* **6**, e22703 (2011).

57.   P. M. White *et al.*, Reliance of *Wolbachia* on high rates of host proteolysis revealed by a

genome-wide RNAi screen of *Drosophila* cells. *Genetics* **205**, 1473-1488 (2017).

58.   N. A. Moran, G. R. Plague, J. P. Sandström, J. L. Wilcox, A genomic perspective on

nutrient provisioning by bacterial symbionts of insects. *Proc. Natl. Acad. Sci.* **100**, 14543-

14548 (2003).

59.   N. A. Moran, Symbiosis. *Curr. Biol.* **16**, R866-871 (2006).

681   **Supplemental Tables and Figures**

683   **Table S1. *Drosophila melanogaster* stocks used in this study.**

BDSC RRID	Complete Genotype	*Wolbachia* infection
41694	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS02259}attP2/TM3, Sb[1]	positive
60086	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC05080}attP40	positive

43296	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS02669}attP40	positive
51492	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC03244}attP2	positive
36686	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS01574}attP2	positive
62241	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC05248}attP40	positive
34347	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS01336}attP2	positive
53332	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC03561}attP40	positive
60019	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC05011}attP40	positive
53339	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC03568}attP40	positive
51801	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.HMC03359}attP40	positive
55183	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC03871}attP40	positive
3954	y[1] w[*]; P{w[+mC]=Act5C-GAL4}17bFO1/TM6B, Tb[1]	positive
31603	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF01355}attP2	negative
1936	Dhod[2] p[p]	positive

686 **Table S2. qPCR primers used in this study.**

Target	FlyBase ID	Forward Primer Sequence (5'-3')	Reverse Primer Sequence (5'-3')
ftsZ	n/a	TTTTGTTGTCGCAAATACCG	CCATTCCTGCTGTGATGAAA
18S	FBgn0085802	CGAAAGTTAGAGGTTCGAAGGCGA	CCGTGTTGAGTCAAATTAAGCCGC
Prat2	FBgn0041194	AATAGTGACCAGTTTGGGCAAG	CCCAGGTTACCCTTTAACTTCC
ade2	FBgn0000052	CAGCCAGGTTTGTTCCTCTG	TAGCTGCTCGTCAAAGCTCG
ade5	FBgn0020513	CCCGAGGGCTACAGGTTCT	AACTTGGCGGAGACAATTTGC
CG11089	FBgn0039241	CTAGCTTGCGTGGCGCTG	CTGGATAAGATCGAAGCCCTGC
su(r)	FBgn0086450	CATCAAGCACACCACATTGTCG	GTGGGACAGGACTTTTGGCA
Dhod	FBgn0000447	TACGGCATACAAGAACCAGGA	TCATCATGGTACTGTGAAACCG
r-l	FBgn0003257	TGTGATGCAAACCGTATCCGA	GTCCTCGACAATCAGACAGGT

Figure S1. Pupal measurements. Pupal volume was calculated based on length and width (red

lines) and assuming a prolate spheroid shape [V = (4/3) π (width/2)^2 (length/2)].

Figure S2. *Wolbachia* titer is not responsive to changes in de novo purine biosynthesis

gene expression. (A) qRT-PCR validation of knockdown across four purine loci. Legend: *** p

<0.001; ** p < 0.01; * p < 0.05; ns p = not significant. **(B)** Knockdown of purine synthesis had no

significant impacts on *Wolbachia* titer (gene*knockdown: $F_{3,31}$ = 0.652, p = 0.588; gene: $F_{3,31}$ =

0.644, p = 0.593; knockdown: $F_{1,31}$ = 0.593, p = 0.447).

Figure S3. Validation of knockdown controls. Impact of balancer chromosomes on (A) purine

gene expression, (B) pyrimidine gene expression, and (C) *Wolbachia* titer. **(A)** The TM6B

balancer had no significant impacts on purine gene expression (gene*balancer: $F_{3,24}$ = 1.311, p =

0.2939; gene: $F_{3,24}$ = 1.311, p = 0.2939; knockdown: $F_{1,24}$ = 4.146, p = 0.0529). Importantly, while

the mean expression of the purine loci may have been marginally higher in the presence of the

balancer chromosome, this means that we would underestimate the effects of knockdown due to

an even larger difference between the sibling and knockdown flies (see main text). **(B)** The TM6B

balancer had no significant impacts on pyrimidine gene expression (gene*balancer: $F_{2,18}$ = 0.227,

p = 0.799; gene: $F_{2,18}$ = 0.227, p = 0.799; knockdown: $F_{1,18}$ = 1.180, p = 0.292). **(C)** There was no

significant difference in *Wolbachia* titer due to the balancer chromosome (t-test, t = 1.8404, df =

8, p = 0.1153). Again, the trend towards marginally higher *Wolbachia* titer in the presence of the

balancer only means that we would underestimate the effects of knockdown.