Wolbachia is a nutritional symbiont

Amelia RI Lindsey[1*], Audrey J Parish[2], Irene LG Newton[2], Jason M Tennessen[2], Megan W

Jones[1], Nicole Stark[2]

[*]To whom correspondence should be addressed (alindsey@umn.edu)

[1]Department of Entomology, University of Minnesota, St. Paul, Minnesota, 55108

[2]Department of Biology, Indiana University, Bloomington, Indiana, 47405

ABSTRACT

Approximately a quarter of eukaryotes are infected with the bacterium *Wolbachia*. Its broad

success as a vertically transmitted infection has been historically difficult to explain given the

myriad of parasitic impacts characterized across *Wolbachia*'s host range. Using the *Drosophila*

model and their natively associated *Wolbachia,* we show that *Wolbachia* infection supports fly

development and buffers against nutritional stress. *Wolbachia* infection across several fly

genotypes and a range of nutrient conditions resulted in reduced pupal mortality, increased adult

emergence, and larger size. We determined that the exogenous supplementation of pyrimidines

rescued these phenotypes in the *Wolbachia*-free, flies suggesting that *Wolbachia* plays a role in

providing this metabolite that is normally limiting for insect growth. Additionally, *Wolbachia* was

sensitive to host pyrimidine metabolism: *Wolbachia* titers increased upon transgenic knockdown

of the *Drosophila de novo* pyrimidine synthesis pathway but not knockdown of the *de novo* purine

synthesis pathway. We propose that *Wolbachia* acts as a nutritional symbiont to supplement

insect development and increase host fitness: a selective advantage that could contribute to its

high frequency in nature.

SIGNIFICANCE STATEMENT

Wolbachia is a bacterial symbiont of ~1/4 of all eukaryotic species on earth, often deemed a

"parasite" due to selfish manipulations of arthropod reproduction. However, many have theorized

there must be more to this symbiosis: parasitic and reproductive impacts alone cannot explain

the success and ubiquity of this bacterium. Here, we use *Drosophila* and their native *Wolbachia*

infections to show that *Wolbachia* supports fly development and significantly buffers flies against

nutritional stress. Additionally, we show that this advantage is likely mediated by pyrimidines: a

biosynthetic pathway that all *Wolbachia* lineages encode for. These developmental advantages

might help explain the ubiquity of *Wolbachia* infections.

INTRODUCTION

Maternally transmitted microbes have evolved numerous ways to manipulate host physiology

ultimately facilitating their own success. Strategies include supplementation of the host's diet via

nutrient provisioning, protection against parasites and infections, and even direct manipulation of

the host's reproduction (1). A high percentage of arthropod species are infected with bacteria in

the genus *Wolbachia* (Alphaproteobacteria: Rickettsiales), a maternally transmitted infection long

considered to be a "reproductive parasite" (2-8). This classification was due to the seemingly

selfish ways in which *Wolbachia* alters host reproductive outcomes ultimately facilitating its own

spread through the population (1, 9, 10). For example, the conversion to asexual reproduction or

the induction of sperm-egg incompatibilities (so-called "cytoplasmic incompatibility") by *Wolbachia*

can result in a rapid sweep of *Wolbachia*-infected matrilines across a population (11, 12).

However, the line between "parasite" and "mutualist" is becoming increasingly blurred as

Wolbachia-mediated benefits for the host are uncovered (13). For example, certain strains of

Wolbachia are clear nutritional mutualists, such as the bed bug-infecting *Wolbachia* that produce

B vitamins to support these obligately hematophagous insects (14). Many other *Wolbachia* strains

have been characterized for their ability to protect host insects against secondary infections,

especially viruses (15). Indeed, evolutionary theory predicts that vertical transmission will select

for mutualism, a process that we have been fortunate to watch happen in real time in some

Wolbachia-host associations (16). Additionally, there are data indicating that *Wolbachia* can

spread through a population even in the absence of so-called "parasitic" reproductive

manipulations (17).

We previously identified that *Wolbachia* infection results in a suite of changes to the expression

of fly nucleotide metabolism pathways (18). Additionally, comparative genomic data indicate that

the biosynthetic pathways for the *de novo* synthesis of purines and pyrimidines have been

conserved across *Wolbachia*, pointing towards their broad importance (19). Furthermore,

Wolbachia encode for an abundance of amino acid importers, which would provide necessary

precursors for *Wolbachia*'s *de novo* synthesis of nucleotides (3, 19). We hypothesized that

Wolbachia plays a supporting role in host nutrition. We show that *Wolbachia* infection is beneficial

for flies, especially under nutrient limited conditions, and that *Wolbachia* is responsive to changes

in the expression of fly metabolic pathways. These data support the idea that *Wolbachia* may

broadly act as a nutritional symbiont, potentially underlying its evolutionary success as a common

infection in the most abundant group of eukaryotes on earth.

METHODS

<u>Fly husbandry</u>

Fly stocks were maintained on standard Bloomington cornmeal-agar medium at 25 °C on a 24-

hour, 12:12 light:dark cycle under density-controlled conditions and 50% relative humidity.

Wolbachia colonization status was confirmed with PCR assays using *Wolbachia*-specific 16S

primers WspecF and WspecR (20). Genotypes used in nutritional assays (below) included:

DGRP-320 (RRID:BDSC_29654), a *Wolbachia*-infected isogenic wild-type strain with genome

sequence available which we refer to as "wild type" below (21), and a *Wolbachia*-infected white

line, w^{145} (RRID:BDSC_145). *Wolbachia*-cleared counterpart stocks were generated with

antibiotics via three generations of tetracycline treatment (20 µg/mL in the fly food for three

generations), followed by re-inoculation of the gut microbiome by transfer to bottles that previously

harbored male flies from the original stock that had fed and defecated on the media for one week

(22). Stocks were allowed to recover from any potential transgenerational effects of the antibiotic

treatment for at least an additional ten generations prior to use in any experiment. All other fly

stocks (e.g., RNAi lines), are detailed in Supplementary Table S1.

Nutritional stress experiments

To generate poor media, Nutri-fly® Bloomington Formulation (Genesee Scientific 66-121) was

prepared according to manufacturer's instructions and diluted to indicated percentages while

maintaining the full concentrations of agar (5.3 g/L) and propionic acid (4.8 ml/L). Five ml of media

was aliquoted into each vial. Flies were placed in mating cages on grape agar plates streaked

with yeast paste and acclimated for 24 hours, prior to initiating 4-hour egg lays. Upon hatching, <

4-hour old L1 larvae were transferred to media vials (n=20 per vial), and development was scored

every 24 hours. Pupation was defined by eversion of the anterior spiracles (23). Each vial was

tracked until all pupae eclosed, or until three consecutive days without additional adult

emergence.

Gnotobiotic assays

To test if the gut microbiome was playing a role in *Wolbachia*-mediated developmental

phenotypes, we reared wild type flies with and without *Wolbachia* on 100% and 25% strength

media under gnotobiotic conditions (i.e., without the gut or food microbiome). 10 mL of media

(prepared according to protocols in "nutritional stress experiments") was aliquoted into

polypropylene wide vials, capped with a Cellulose Acetate Flug® (Flystuff 49-101), and

autoclaved on liquid setting for 20 minutes to ensure sterility. Embryos were collected on grape

agar with yeast paste as described previously and transferred into 70-micron mesh cell strainers

(Falcon 352350). Embryos in the cell strainers were washed with *Drosophila* embryo wash

solution (7% NaCl, 0.5% Triton X-100) to remove food debris. After an initial wash, embryos were

immersed in a 10% bleach solution for 3 minutes, with gentle mixing every 30 seconds. The

embryos were then washed with sterile *Drosophila* embryo wash solution to remove excess

bleach. Embryos were subjected to a final rinse with sterile phosphate buffered saline (PBS) and

transferred to a sterile agar culture plate (2% agar in deionized water) stained with blue gel food

coloring to facilitate counting. A flame-sterilized probe was used to transfer 20 embryos into each

vial of sterile media.

<u>Nutritional supplementation experiments</u>

To test the impact of specific nutrients, we prepared 12.5% strength media as above, and

selectively added in protein or nucleotides. Treatments included casein (64.1 g/L; Sigma C7078),

inosine (0.74 g/L; Sigma I4125), uridine (0.67 g/L; Sigma U3003), or both inosine and uridine at

the aforementioned concentrations. The concentrations of additives were calculated as 87.5% of

a previously defined holidic diet (24), to adjust for the 12.5% BDSC media strength.

Developmental assays were carried out as described above.

<u>Pupal size measurements</u>

After selected developmental assays, pupal casings and dead pupae were removed from vials

using a wet paint brush and transferred to glass slides. Pupae were imaged with brightfield

microscopy on an ECHO Revolve at 4X. Images were manually annotated in Echo Labs Pro

software to measure sizes. Pupal length was defined as the distance between the base of the

posterior spiracles to the midway point between the anterior spiracles. Width measurements were

taken at the widest part of the pupa. The volume of each pupa was calculated assuming a prolate

spheroid shape $[V = (4/3) \pi (width/2)^2 (length/2)]$ (Supplemental Figure S1) (25, 26).

<u>Knockdown of fly nucleotide metabolism genes</u>

Wolbachia-infected TRiP stocks were obtained from the Bloomington *Drosophila* Stock Center to

knock down expression of genes in the purine and pyrimidine nucleotide biosynthesis pathways

(Supplemental File S1). Unmated females carrying a UAS-gene-specific short hairpin were

crossed to males with an Act5C-Gal4 driver (RRID:BDSC_3954: y^1w^*; P{w[+mC]=Act5C-

GAL4}17bFO1/TM6B, Tb[1]). Unmated, F1 female progeny were sorted as adults into those with

the short hairpin and Gal4-driver or those with the short hairpin and the TM6B balancer. At three

days old, the sorted flies were flash frozen in liquid nitrogen and then stored at -80 °C for later

processing. To determine the effect of the TM6B balancer on *Wolbachia* titer, we crossed

Wolbachia-infected, Act5C-Gal4 driver females to *Wolbachia*-uninfected UAS-anti-Ppyr\LUC

males (RRID:BDSC_31603: y^1 v^1; P{$y^{+t7.7}$ $v^{+t1.8}$=TRiP.JF01355}attP2). Unmated, F1 female

progeny were sorted and stored as described for the nucleotide biosynthesis gene knockdowns.

To determine spatial differences in *Wolbachia* titer, we dissected ovaries from two-day old *r-l*

knockdown and sibling control flies. We separated ovaries from living, anesthetized flies in sterile

PBS. Ovaries and carcasses from three flies were pooled for each biological replicate and flash

frozen, followed by storage at -80 °C for later processing.

<u>Real-time quantitative RT-PCR analyses of target gene expression</u>

Flies were homogenized in TRIzol reagent (Invitrogen), and total RNA was extracted following

manufacturer's instructions. RNA extractions were DNase treated (RQ1 RNase-free DNase, New

England Biolabs) according to the manufacturer's instructions. cDNA was synthesized using

MMuLV Reverse Transcriptase (New England Biolabs) with random hexamer primers (Integrated

DNA Technologies). Quantitative RT-PCR reactions were performed with SensiFAST SYBR Hi-

ROX kit (Bioline) and gene-specific primers (Supplemental File S2). All samples were run in

technical duplicate alongside negative controls on an Applied Bioscience StepOnePlus qPCR

machine (Life Technologies). Gene expression was normalized to endogenous 18S rRNA

expression using the Livak method (27).

<u>Real-time quantitative PCR analyses of *Wolbachia* titer</u>

DNA was extracted from individual flies using the Quick-DNA/RNA Pathogen Miniprep kit (Zymo

Research) according to the manufacturer's protocol. *ftsZ* primers were used to quantify *Wolbachia*

genome copy numbers, which were normalized to host genome copies via 18S quantification

using the Livak method (27). Reactions were performed with SensiFAST SYBR Hi-ROX kit

(Bioline), and all samples and negative controls were run in technical duplicate on an Applied

Bioscience StepOnePlus qPCR machine (Life Technologies).

<u>Nucleotide metabolism biosynthesis and gene expression</u>

Nucleotide biosynthesis pathway information was downloaded from BioCyc by querying the

"*Wolbachia* endosymbiont of *Drosophila melanogaster w*Mel, version 26.5" and "*Drosophila*

melanogaster, version 26.5" databases (28). Enzymes for each step were cross-referenced with

KEGG Pathways, release 105.0 (29). To assess expression of *Wolbachia* metabolic pathways,

we analyzed the *Wolbachia* transcriptomes previously extracted from the MODENCODE dataset

(30, 31). We leveraged the previously published FPKM values for each *Wolbachia* gene. For

simplicity of visualization, mean FPKM is displayed for developmental time points represented by

multiple libraries. Fly gene expression for *Wolbachia*-infected and *Wolbachia* uninfected adult

females was previously reported (18).

<u>Statistics and Data Visualization</u>

Statistics and data visualization were carried out in R version 3.5.0 (32). Significant differences in

development were assessed with generalized linear mixed-effects models (package: 'lme4',

function 'glmer' (33)) including the proportion of flies that reached a given stage (pupa or adult)

as a binomial response, *Wolbachia* presence, day of development, media, and the interaction of

the three as fixed effects, and vial as a random effect to account for repeated measures.

Significant differences in pupal volume of flies reared on 12.5% media was assessed with a two-

way ANOVA (function 'aov') including *Wolbachia*, mortality status, and their interaction as fixed

effects. Significant differences in sizes of pupae reared on supplemented media was first

assessed with a two-way ANOVA (function 'aov') including *Wolbachia*, media, and their

interaction as fixed effects. Paired comparisons between *Wolbachia*-infected and uninfected flies

were then assessed with Wilcoxon rank sum tests (function 'wilcox.test'). Gene expression and

Wolbachia titers from knockdown experiments were assessed with two-way ANOVAs (function

'aov') including delta delta Ct as the response, and target locus, genotype (knockdown versus

sibling), and their interaction as fixed effects. *Wolbachia* titers in dissected flies were also

assessed with ANOVA, here with genotype, tissue, and their interaction as fixed effects. Pairwise

comparisons were performed with t-tests.

RESULTS

<u>*Wolbachia* is beneficial under nutrient limited conditions</u>

To assess the impact of *Wolbachia* infection on fly development, we reared flies on 100% strength

and 25% strength media and quantified pupation and adult emergence. We found a significant

interaction between *Wolbachia*, media strength, and time that impacted fly pupation ($F_{1,1063}$ =

24.7147, $p < 0.0001$) and adult emergence ($F_{1,1063}$ = 22.2897, $p < 0.0001$). Additionally, we found

a significant interaction between the presence of *Wolbachia* and the media strength (wild type

pupae: $F_{1,1063}$ = 0.6517, $p = 0.0063$; wild type adults: $F_{1,1063}$ = 0.1746, $p = 0.0062$; w^{145} pupae: $F_{1,711}$

= 8.8129, $p = 0.0009$; w^{145} adults: $F_{1,711}$ = 19.0998, $p < 0.0001$), along with a significant impact of

Wolbachia infection alone on fly pupation and adult emergence (wild type pupae: $F_{1,1063}$ = 0.2874,

$p < 0.0001$; wild type adults: $F_{1,1063}$ = 0.5375, $p < 0.0001$; w^{145} pupae: $F_{1,711}$ = 10.5483, $p < 0.0001$;

w^{145} adults: $F_{1,711}$ = 7.0718, $p < 0.0001$). These effects are driven by the nutrient-limited conditions

where *Wolbachia*-infected flies developed faster (1-5 days depending on genotype, Figures 1B

and 1E) and a larger percentage of flies reached adulthood. For example, while the wild type flies

with and without *Wolbachia* reared on 100% strength media reached adulthood in 82% and 81%

of cases, on the 25% strength media these were reduced to 75% and 66% respectively. The

Wolbachia-mediated advantage was more prominent for a second fly genetic background, w^{145}.

Even on 100% strength media, the *Wolbachia*-free w^{145} flies experienced a 5% reduction in adult

emergence relative to *Wolbachia*-infected flies (Figure 1D). When reared on 25% strength media,

the *Wolbachia*-infected w^{145} flies were more than twice as likely to reach adulthood (70% versus

31%, Figure 1E). Across these assays, we found delays in entering pupation are approximately

equal to the delay in adult emergence: *i.e.,* the time spent in metamorphosis did not change as a

factor of *Wolbachia* infection.

Given that the effect size for the *Wolbachia* advantage was genotype dependent and given the

more subtle impact of *Wolbachia* in the wild-type flies, we wondered if an even weaker media

would result in a stronger advantage for the *Wolbachia*-infected flies. Conversely, *Wolbachia*

could be a burden under more significant nutritional stress. Indeed, when flies were raised on

12.5% strength media, we again saw significant effects of the interaction between *Wolbachia* and

time on the rate and success of pupation and adult emergence (pupae: $F_{1,347}$ = 77.135, p < 0.0001;

adults: $F_{1,347}$ = 58.9296, p < 0.0001). However, *Wolbachia*-infected flies now developed slower

than their uninfected counterparts (6-7 days delayed), but they pupated at higher rates (68%

versus 61%) and were more likely to reach adulthood (54% versus 40%, Figure 1C). Furthermore,

although *Wolbachia*-infected fly development was slower, it resulted in them attaining, on

average, a 23% larger size at pupation ($F_{1,191}$ = 32.076, p < 0.0001, Figure 1F). Additionally,

Wolbachia-infected flies experienced lower levels of pupal mortality (35% versus 19%), perhaps

related to the finding that flies that died during pupation were significantly smaller ($F_{1,191}$ = 4.625,

p = 0.0328, Figure 1F).

To determine if the developmental advantage was due to direct effects of *Wolbachia* or indirect

effects via *Wolbachia*-mediated impacts on the gut microbiome, we performed the same

developmental assay under gnotobiotic conditions (i.e., without a gut or food microbiome). Under

gnotobiotic conditions, we saw a strong interactive effect of *Wolbachia* and time that resulted in

significantly more, and faster, pupation and adult emergence for infected flies (Figure 1G; pupae:

$F_{1,1920}$ = 153.3796, p < 0.0001; adults: $F_{1,1920}$ = 129.6080, p < 0.0001). Even on the 100% strength

media, only 43% of *Wolbachia*-free flies pupated, compared to 61% of *Wolbachia*-infected flies.

Again, *Wolbachia*-free flies had higher levels of pupal mortality: 21% of those that pupated did

not emerge as adults, as compared to 13% of the *Wolbachia*-infected pupae. While removing the

microbiota negatively impacted fly developmental timing as expected, this effect was exacerbated

in the *Wolbachia*-free flies (Figures 1A and 1G). As compared to the conventionally-reared flies

on 100% strength media (Figure 1A), gnotobiotic *Wolbachia*-infected flies experienced a two-day

developmental delay, and *Wolbachia*-free flies were delayed three days (Figure 1G). The

combination of gnotobiotic conditions and 25% strength media resulted in especially impaired fly

development: less than 20% of flies reached adulthood (Figure 1H). While there were no

significant differences in the proportion of flies that pupated or emerged as adults between

Wolbachia-infected and uninfected flies in these high-mortality conditions (pupae: p = 0.9547;

adults: p = 0.6210), it is notable that *Wolbachia*-infected flies on average started emerging five

days ahead of the *Wolbachia*-free flies (Figure 1H).

Figure 1. *Wolbachia* infection is beneficial under nutrient limited conditions. Flies (wild type

and *w*[145], with and without *Wolbachia*) were reared on different concentrations of media to test

the impact of *Wolbachia* infection. Biological replicates included 20 larvae per each of 10 vials.

Dotted and solid lines indicate cumulative pupation and adult emergence, respectively, with

shaded regions defining standard error. All *Wolbachia*-infected treatments are in red, and

Wolbachia-uninfected in grey. **(A)** Conventionally reared wild type flies on 100%, **(B)** 25%, and

(C) 12.5% strength media. **(D)** Conventionally reared w^{145} flies on 100% and **(E)** 25% strength

media. **(F)** Pupal volumes of wild type flies with and without *Wolbachia*, derived from the 12.5%

media-reared flies in (C). Orange datapoints indicate pupae that did not eclose into adults. *** p

< 0.001 **(G)** Gnotobiotic wild type flies on 100% and **(H)** 25% strength media.

<u>Dynamics of the redundant *de novo* nucleotide biosynthesis pathways</u>

We next considered the role that *Wolbachia* may be playing in supporting fly development during

these nutritionally limited conditions. We previously found that nucleotide metabolism is a

significant point of interaction for flies and *Wolbachia* (18), pathways known to significantly impact

fly growth and development (34). Importantly, the pathways for the *de novo* synthesis of purines

and pyrimidines are conserved across all *Wolbachia*, and they are redundantly encoded by the

fly (Figure 2A-B). Additionally, we previously showed that *Wolbachia* infection results in

downregulation of the fly's pyrimidine synthesis enzymes *rudimentary* (-0.273 Fold Change

compared to uninfected flies), *Dhod* (-0.164 FC), and *rudimentary-like* (-0.301 FC) (18). Strikingly,

the repressor *su(r)* is upregulated (+0.505 FC)(Figure 2C). In contrast, purine metabolism overall

was not obviously distinct between *Wolbachia*-infected and uninfected flies: some enzymes were

up- or down- regulated but not the entire pathway (Figure 2C).

Given the developmental effects of *Wolbachia* infection, we hypothesized that *Wolbachia*

nucleotide metabolic pathways would be more highly expressed during growth stages (*i.e.,* pre-

metamorphosis). Indeed, from embryogenesis through the third larval instar (L3), most of

*Wolbachia'*s nucleotide metabolism genes were upregulated. At the onset of pupation (the white

pre-pupae "WPP" stage), *Wolbachia*'s metabolic gene expression was sharply and broadly

downregulated (Figure 2D). These data indicate that *Wolbachia* may be supplementing host nutrition during critical growth stages, and either limiting negative impacts of infection during metamorphosis and reproductive stages or shifting roles to better interface with host reproduction and ensure transmission.

Figure 2. Flies and *Wolbachia* redundantly encode for *de novo* nucleotide biosynthesis. **(A)** Metabolic pathway for *de novo* pyrimidine biosynthesis. **(B)** Metabolic pathway for *de novo* purine biosynthesis. **(C)** Expression of *Drosophila melanogaster* nucleotide biosynthesis loci in *Wolbachia*-infected female flies relative to *Wolbachia*-uninfected controls, modified from (18). **(D)** Expression of *Wolbachia* nucleotide biosynthesis loci across fly development. *Wolbachia* transcriptomes were generated as part of the MODENCODE project (30, 31). Data are row normalized for inference of per-gene transcriptional changes across development. Fly images are

from BioRender.com. Abbreviations: UMP: uridine monophosphate; IMP: inosine

monophosphate. Fly gene alternate names: Ade2: Pfas; Ade3: Gart; Ade5: Paics.

<u>Pyrimidines contribute to the *Wolbachia* infection advantage</u>

We reasoned that deficiencies in the diet supplemented by *Wolbachia* could be supplemented by

exogenous nutrient addition to the fly food. Therefore, flies with and without *Wolbachia* would

have the same growth parameters if this important nutrient was added. Specifically, we focused

on the precursors and end-products of the *de novo* nucleotide synthesis pathways, which we

supplemented back into the 12.5% media. The developmental assays on supplemented media

were repeated with the wild type stocks, avoiding w^{145} because the *white* transporter is known to

affect nucleotide metabolism (35). We again saw the *Wolbachia*-mediated developmental

advantage in the 12.5% strength media (pupae: $F_{1,235}$ = 13.9505, p = 0.0002; adults: $F_{1,235}$ =

12.7568, p = 0.0004). Additionally, we saw a subtle but significant positive impact of *Wolbachia*

in the 100% strength media as measure by adult emergence (pupae: $F_{1,235}$ = 1.3002, p = 0.2538;

adults: $F_{1,235}$ = 28.0455, p < 0.0001), due to 6% pupal mortality in the *Wolbachia*-free flies that the

Wolbachia-infected flies did not experience (Figure 3A). While the addition of casein benefited

both *Wolbachia*-infected and uninfected flies, *Wolbachia*-free flies pupated at significantly

increased rates relative to *Wolbachia*-infected flies ($F_{1,235}$ = 9.8699, p = 0.0017). Again, pupal

mortality was quite high in the *Wolbachia*-free flies (36% as compared to 9% with *Wolbachia*),

resulting in significantly fewer numbers of adults ($F_{1,235}$ = 6.9988, p = 0.0082). The addition of

uridine (product of *de novo* pyrimidine synthesis) specifically benefited *Wolbachia*-free flies, but

not the flies that had *Wolbachia* infections. Indeed, uridine supplementation resulted in a rescue

phenotype where equivalent numbers of *Wolbachia*-infected and uninfected flies emerged as

adults (p = 0.9281), almost identical to the rates of adult emergence for *Wolbachia*-infected flies

on the standard 12.5% strength media (Figure 3A). In fact, *Wolbachia*-free flies supplemented

with uridine developed slightly faster than their infected counterparts (pupae: $F_{1,235}$ = 2.2401, p =

0.1348; adults: $F_{1,235}$ = 10.5842, p = 0.0011). The addition of inosine (product of *de novo* purine

synthesis) was toxic and reduced pupation and emergence for all flies. As such there were no

significant differences between *Wolbachia*-infected and uninfected flies in these high mortality

(*i.e.*, zero-inflated) conditions (pupae: $F_{1,235}$ = 0.9217, p = 0.335; adults: $F_{1,235}$ = 0.3519, p = 0.553).

Simultaneously adding uridine and inosine resulted in more intermediate, albeit still quite lethal

developmental phenotypes with no significant impact of *Wolbachia* ($F_{1,235}$ = 1.3471, p = 0.2459;

adults: $F_{1,235}$ = 1.0400, p = 0.307).

Congruent with the previous developmental assays (Figure 1), *Wolbachia* infection had a

significant impact on pupal size, both due to interacting effects with the media ($F_{5,470}$ = 2.55, p =

0.0272), and *Wolbachia* alone ($F_{1,470}$ = 37.31, p < 0.0001) (Figure 3B). Furthermore, *Wolbachia*-

infected flies were significantly larger when reared on 100% (p = 0.0401), 12.5% (p = 0.0316),

and casein-enriched 12.5% media (p < 0.0001). However, the addition of uridine to the 12.5%

media resulted in similarly sized flies regardless of *Wolbachia* infection (p = 0.3685). In summary,

Wolbachia-free flies were rescued to the same size and level of adult emergence as *Wolbachia*-

infected counterparts in uridine (i.e., pyrimidine) supplemented media.

Figure 3. Uridine rescues *Wolbachia*-free flies. Wild type flies were reared on 100% media, 12.5% media, or 12.5% media containing either casein, uridine, inosine, or both inosine and uridine. The media designations in the middle correspond to the figures above and below. Biological replicates included 20 larvae per each of five vials. **(A)** Dotted and solid lines indicate cumulative pupation and adult emergence, respectively, with shaded regions defining standard error. *Wolbachia*-infected treatments are in red and *Wolbachia*-uninfected are in grey. **(B)** Pupae from (A) were removed post-developmental assay, measured, and their volume was calculated. Orange datapoints indicate pupae that did not eclose into adults. Significance annotations for

pupal sizes comparisons are the same regardless of whether flies that died during pupation were

included. ***p < 0.001; **p < 0.01; *p < 0.05; ns = not significant.

Wolbachia infection is responsive to pyrimidine biosynthesis gene expression

To determine whether _Wolbachia_ infection was sensitive to changes in host metabolism, we

knocked down individual genes within the fly's _de novo_ nucleotide synthesis pathways (Figure

2A-B) using _Wolbachia_-infected TRiP RNAi fly stocks (Figure 4A). Short-hairpin RNA (shRNA)

expression was driven in the entire body of the fly by an Act5C-Gal4 driver. We compared gene

expression and _Wolbachia_ titer in knockdown flies to sibling controls (flies not expressing a gene-

targeting shRNA, and instead with the TM6B balancer). We successfully knocked down three

genes in the pyrimidine synthesis pathway (_su(r)_, _Dhod_, _rudimentary-like_) and four genes in the

purine synthesis pathway (_prat2_, _ade2_, _ade5_, _CG11089_) (Figure 4A, Supplemental Figure S2).

Several gene knockdowns were lethal to fly development, including _rudimentary_ in the de novo

pyrimidine biosynthesis pathway and _CG2246_, _CG6767_, _Prat_, and _AdSL_ in the de novo purine

biosynthesis pathway. Broad-spectrum knockdown of _de novo_ purine synthesis had no significant

impacts on _Wolbachia_ titer (gene*knockdown: $F_{3,31}$ = 0.652, p = 0.588; gene: $F_{3,31}$ = 0.644, p =

0.593; knockdown: $F_{1,31}$ = 0.593, p = 0.447; Supplemental Figure S2). In contrast, knockdown of

the _de novo_ pyrimidine synthesis pathway resulted in significant increases in _Wolbachia_ titer ($F_{1,38}$

= 10.3916, p = 0.0026; Figure 4B). This pattern was specifically driven by the enzymatic loci

(_Dhod_: p = 0.034 and _rudimentary-like_ (_r-l_)_:_ p = 0.013). In contrast, knockdown of the negative

regulator _sur(r)_ had no effect on _Wolbachia_ titer (p = 0.530), perhaps indicating that _Wolbachia_

titers are responsive to low fly pyrimidine synthesis activity, but not elevated levels. To account

for any effects of the TM6B balancer on measurements of _Wolbachia_ abundance, we crossed the

Act5C-Gal4 driver to a TRiP line expressing shRNA against a foreign gene: _Photinus pyralis_

luciferase (UAS-anti-Ppyr\LUC). We found no significant changes in _Wolbachia_ titer, or in

nucleotide gene expression that could be driving the results in Figure 4B (Supplemental Figure

S3). We then asked if the change in *Wolbachia* titer was tissue-specific by measuring the relative

Wolbachia titers in ovaries and the remaining carcasses after knockdown of *r-l* (Figure 4C). There

was no significant interaction of knockdown and tissue ($F_{1,32}$ = 0.20, p = 0.658) or of tissue alone

($F_{1,32}$ = 0.20, p = 0.658) on *Wolbachia* titers. We again saw a significant effect of *r-l* knockdown

on *Wolbachia* titers ($F_{1,32}$ = 13.07, p = 0.001) both in ovaries (p = 0.007) and in carcasses (p =

0.0127) (Figure 4C). Finally, given these interactions with pyrimidine synthesis, we used flies with

a homozygous hypomorphic allele *Dhod^2* (RRID:BDSC_1936, Supplemental Table S1) to assess

the importance of *Wolbachia* in a pyrimidine synthesis-impaired genetic background. This stock

is naturally infected with *Wolbachia*, but removal of the symbiont lead to significant delays and

reductions in pupation and adult emergence (Figure 4D; pupae: $F_{1,247}$ = 11.0058, p = 0.0009;

adults: $F_{1,247}$ = 12.0550, p = 0.0057).

Figure 4. *Wolbachia* titers are responsive to fly pyrimidine synthesis gene expression.

Expression of fly genes in the *de novo* pyrimidine pathway was constitutively knocked down using

standard transgenic approaches. **(A)** *Drosophila* gene knockdown was verified with qRT-PCR

leveraging gene specific primers and normalization to 18S expression. **(B)** *Wolbachia* titers in

whole adult females and **(C)** dissected ovaries and carcasses were quantified with qPCR using

Wolbachia-specific primers and normalization to fly genome copy number via 18S amplification.

Legend: *** p <0.001; ** p < 0.01; * p < 0.05; ns p = not significant. **(D)** Development of

conventionally reared *Wolbachia*-infected and -uninfected *Dhod^2 p^2* flies on 100% strength media.

Biological replicates included 20 larvae per each of ten vials. Dotted and solid lines indicate

cumulative pupation and adult emergence, respectively, with shaded regions defining standard

error. *Wolbachia*-infected flies are in red and *Wolbachia*-uninfected are in grey.

DISCUSSION

Wolbachia has long been appreciated for its diverse impacts on a range of ecdysozoan hosts,

and there has been increasing speculation as to the role *Wolbachia* might play in host metabolism

(13, 18, 19). Here, we show that *Wolbachia* significantly impacts fly development, resulting in

larger flies more likely to reach adulthood. The strength of this phenotype varied by fly genotype

and the makeup of the food: *Wolbachia* infection was highly beneficial under nutrient limited

conditions. Removal of the extracellular microbial community indicated that this advantage was

directly mediated by *Wolbachia*. In fact, *Wolbachia* significantly buffered flies against the stresses

normally associated with removal of the microbiome (36, 37). Furthermore, *Wolbachia* titers were

responsive to changes in the expression of fly pyrimidine biosynthesis, and the supplementation

of pyrimidines rescued adult emergence of the *Wolbachia*-uninfected flies. We propose that

pyrimidines are a significant component of the advantage provided by *Wolbachia* infection.

Indeed, the patterns of *Wolbachia* gene expression seem to be attuned to fly development which

could be a strategy for supporting the exponential growth that occurs during larval stages, and

mitigating fitness impacts of the infection during reproductive stages. Importantly, while the

transcription of *Wolbachia*'s nucleotide metabolism pathways are quite high pre-pupation, the flies

have comparatively low *Wolbachia* titers during that time (30). Genomic data suggest that

Wolbachia imports amino acids from the host: while these may be used as precursors for

nucleotide biosynthesis, the depletion of free amino acids could induce host stress responses (15,

38). Changes in *Wolbachia* titer and physiology across development and under different

nutritional scenarios likely reflect such balancing acts.

While *Wolbachia* infections have received considerable attention for their impacts on

reproduction, we know much less about how *Wolbachia* interact with their host during juvenile

stages. Importantly, many insects spend the majority of their lifetime as juveniles. Juvenile insect

metabolism is typically quite distinct from adults, and this time period has profound impacts on

metamorphosis, adult physiology, and fitness (39). Critically, all insect growth occurs during the

larval period: in *Drosophila,* larvae increase in body mass by ~200-fold over the course of 4 days

(39). We found that *Wolbachia*-free flies had higher levels of pupal mortality, especially during

nutritional stress, suggesting insufficient metabolic reserves. This effect was particularly

pronounced when flies were reared on an excess of protein (Figure 3). Importantly, amino acids

are a key nutritional cue for determining the timing of pupation (i.e., signaling critical weight) due

to a direct and indirect effects on the timing and intensity of ecdysone pulses (40-43). An influx of

amino acids without other essential nutrients needed for growth might impact nutritional signaling

and thus the timing of critical weight, perhaps triggering flies to pupate early without sufficient

reserves for metamorphosis. Given *Wolbachia* likely imports amino acids, this may reduce the

concentrations of free amino acids and delay signaling critical weight, allowing flies to accumulate

more metabolic reserves before pupation.

During the ~200-fold increase in larval body mass, there is a concomitant 25-fold increase in DNA

and RNA content that creates a huge demand for pyrimidines, well exceeding the fly's

endogenous biosynthetic capacity (44). Considering that the rate of *Drosophila* development is

highly sensitive to *de novo* pyrimidine synthesis (44), bacterially-derived nucleotides could give

the host a significant advantage. Pyrimidines are synthesized by a series of metabolic reactions

that use ribose sugar and the amino acids glutamine and aspartate to generate uridine-5'-

monophosphate, UMP (45): the precursor for uracil, cytosine, and thymine. This pathway is

conserved in both *Drosophila* and *Wolbachia*, suggesting that infected flies generate pyrimidines

in multiple compartments: the host cytoplasm/mitochondria and *Wolbachia*. *Wolbachia*'s

production of pyrimidines could also represent far more than simply additional biosynthesis, as

the enzyme dihydroorotate dehydrogenase (*Dhod*) is located in the inner mitochondrial

membrane where it couples pyrimidine synthesis to electron transport (46, 47). As such,

provisioning of *Wolbachia*-generated pyrimidines could reduce the biosynthetic burden normally

imposed upon mitochondria. Additionally, fly enzymes used for pyrimidine synthesis, CAD and

UMP synthase, are negatively regulated by UMP (48). Therefore, *Wolbachia* could simultaneously

impact free amino acid pools, supplement pyrimidines, reduce mitochondrial burdens, and

spatially segregate components of a normally inhibitory interaction.

While we found that pyrimidines seemed to be important for the fly-*Wolbachia* symbiosis, we did

not see any changes in *Wolbachia* titer due to repression of purine synthesis. In fact, the addition

of inosine to media negatively impacted flies (Figure 3). Purine nucleotides act as signaling

molecules that impact a range of functions. For example, activation of the adenosine receptor

globally suppresses fly metabolism (49), which can cause insecticidal effects (50). Another

possibility is misincorporation of inosine into RNAs. In mammalian cell lines, this can happen

when inosine is in excess, ultimately impacting translation (51). While altering the expression of

fly purine metabolic gene expression did not impact *Wolbachia* titers in our experiments, we

cannot not rule out the potential for other changes in *Wolbachia* gene expression or physiology.

Indeed, we previously found an interactive effect of *Wolbachia* infection and purine gene

expression on virus replication (18), highlighting both the importance of purines to the symbiosis

and the complexity of the metabolic landscape.

The role of *Wolbachia* as a nutritional symbiont raises numerous questions about the cell biology

of *Wolbachia* and how the relationship with the host is regulated. Is *Wolbachia* specifically

providing pyrimidines (e.g., via nucleotide transporters)? Or are flies farming and consuming their

Wolbachia infections (e.g., autophagy) and pyrimidines so happen to be abundant in *Wolbachia*

cells and also a limiting metabolite for the fly? As is the case for many symbioses, it is not always

clear if a change in the symbiont (e.g., gene expression or titer), is due to the microbe detecting

and responding to the host's status, or the host releasing control of the microbe and allowing a

change in microbial physiology. *Wolbachia* do encode for a number of proteins that would allow

them to detect their environment and regulate gene expression accordingly (52). However, each

Wolbachia is enclosed by a host-derived vesicle (53, 54), so perhaps the host tightly regulates

Wolbachia's "experience".

While nutritional mutualisms in insects are not rare, most of the systems for which this is described

are of a much more binary nature. Insects that exclusively feed on unbalanced diets (e.g., blood,

plant sap) have obligate microbial symbionts to synthesize metabolites that the host cannot

generate *de novo*: the partners complement each other (55, 56). Here, both flies and *Wolbachia*

have their own complete pathways for nucleotide biosynthesis, which perhaps has led us to

overlook the relevance of this biosynthetic redundancy. We propose that the metabolic and

developmental impacts of *Wolbachia* are adaptive and might explain the broad success and

spread of *Wolbachia* even in the absence of direct reproductive manipulations.

DECLARATIONS

Acknowledgements

This work was supported by UMN AGREETT startup funds to ARIL. ILGN in supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under R01 AI144430. JMT is supported by the National Institute of General Medical Sciences of the National Institutes of Health under a R35 Maximizing Investigators' Research Award (MIRA; 1R35GM119557). The authors thank Richard Hardy for support and feedback. Research was further made possible by Flybase (NIH 5U41HG000739), and stocks obtained from the Bloomington *Drosophila* Stock Center (NIH P40OD018537).

Conflicts of interest

The authors declare that they have no competing interests.

Data Availability

Supplemental Table S1. Fly stocks

Supplemental Table S2. Primers

Supplemental Figure S1. Pupal measurement.

Supplemental Figure S2. Purine knockdowns.

Supplemental Figure S3. Knockdown controls.

REFERENCES

1.   R. Kaur *et al.*, Living in the endosymbiotic world of *Wolbachia*: A centennial review. *Cell Host & Microbe* (2021).

2.   O. Duron *et al.*, The diversity of reproductive parasites among arthropods: *Wolbachia* do not walk alone. *BMC Biol.* **6**, 1-12 (2008).

3.   M. Wu *et al.*, Phylogenomics of the reproductive parasite *Wolbachia* pipientis *w*Mel: A

streamlined genome overrun by mobile genetic elements. *PLoS Biol.* **2**, 327-341 (2004).

4.   J. H. Werren, W. Zhang, L. R. Guo, Evolution and phylogeny of *Wolbachia* - reproductive

parasites of arthropods. *Proc. R. Soc. Lond. B* **261**, 55-63 (1995).

5.   N. Bensaadi-Merchermek, J.-C. Salvado, C. Cagnon, S. Karama, C. Mouchès,

Characterization of the unlinked 16S rDNA and 23S-5S rRNA operon of *Wolbachia*

pipientis, a prokaryotic parasite of insect gonads. *Gene* **165**, 81-86 (1995).

6.   R. Zug, P. Hammerstein, Still a host of hosts for *Wolbachia*: Analysis of recent data

suggests that 40% of terrestrial arthropod species are infected. *PLoS One* **7** (2012).

7.   K. Hilgenboecker, P. Hammerstein, P. Schlattmann, A. Telschow, J. H. Werren, How

many species are infected with *Wolbachia*? - A statistical analysis of current data. *FEMS*

Microbiol. Lett. **281**, 215-220 (2008).

8.   F. Rousset, D. Bouchon, B. Pintureau, P. Juchault, M. Solignac, *Wolbachia*

endosymbionts responsible for various alterations of sexuality in arthropods. *Proc. R. Soc.*

Lond. B **250**, 91-98 (1992).

9.   K. Fenn, M. Blaxter, *Wolbachia* genomes: revealing the biology of parasitism and

mutualism. *Trends Parasitol.* **22**, 60-65 (2006).

10.   R. Stouthamer, J. A. J. Breeuwer, G. D. D. Hurst, *Wolbachia pipientis*: Microbial

manipulator of arthropod reproduction. *Annu. Rev. Microbiol.* **53**, 71-102 (1999).

11.   M. Turelli, A. A. Hoffmann, Rapid spread of an inherited incompatibility factor in California

Drosophila. *Nature* **353**, 440-442 (1991).

12.   R. Stouthamer, J. E. Russell, F. Vavre, L. Nunney, Intragenomic conflict in populations

infected by Parthenogenesis Inducing *Wolbachia* ends with irreversible loss of sexual

reproduction. *BMC Evol. Biol.* **10**, 12 (2010).

13.   R. Zug, P. Hammerstein, Bad guys turned nice? A critical assessment of *Wolbachia*

mutualisms in arthropod hosts. *Biological Reviews* **90**, 89-111 (2015).

14.   T. Hosokawa, R. Koga, Y. Kikuchi, X. Y. Meng, T. Fukatsu, *Wolbachia* as a bacteriocyte-
559   associated nutritional mutualist. *Proc. Natl. Acad. Sci.* **107**, 769-774 (2010).

560   15.   A. R. I. Lindsey, T. Bhattacharya, I. L. G. Newton, R. W. Hardy, Conflict in the intracellular
561   lives of endosymbionts and viruses: A mechanistic look at *Wolbachia*-mediated pathogen-
562   blocking. *Viruses* **10**, 141 (2018).

563   16.   A. R. Weeks, M. Turelli, W. R. Harcombe, K. T. Reynolds, A. A. Hoffmann, From parasite
564   to mutualist: Rapid evolution of *Wolbachia* in natural populations of *Drosophila*. *PLoS Biol.*
565   **5**, e114 (2007).

566   17.   C. A. Hamm *et al.*, *Wolbachia* do not live by reproductive manipulation alone: infection
567   polymorphism in *Drosophila suzukii* and *D. subpulchrella*. *Mol. Ecol.* **23**, 4871-4885
568   (2014).

569   18.   A. R. Lindsey, T. Bhattacharya, R. W. Hardy, I. L. Newton, *Wolbachia* and virus alter the
570   host transcriptome at the interface of nucleotide metabolism pathways. *mBio* **12** (2021).

571   19.   I. L. Newton, D. W. Rice, The Jekyll and Hyde symbiont: could *Wolbachia* be a nutritional
572   mutualist? *J. Bacteriol.* **202** (2020).

573   20.   J. H. Werren, D. M. Windsor, *Wolbachia* infection frequencies in insects: Evidence of a
574   global equilibrium? *Proc. R. Soc. Lond. B* **267**, 1277-1285 (2000).

575   21.   W. Huang *et al.*, Natural variation in genome architecture among 205 *Drosophila*
576   *melanogaster* Genetic Reference Panel lines. *Genome Res.* **24**, 1193-1208 (2014).

577   22.   M. W. Jones, L. C. Fricke, C. J. Thorpe, L. O. V. Esch, A. R. I. Lindsey, Infection dynamics
578   of cotransmitted reproductive symbionts are mediated by sex, tissue, and development.
579   *Appl. Environ. Microbiol.* **88**, e00529-00522 (2022).

580   23.   S. Bainbridge, M. Bownes, Staging the metamorphosis of *Drosophila melanogaster*.
581   *Journal of Embryology and Experimental Morphology* **66**, 57-80 (1981).

582   24.   T. Reis, Effects of synthetic diets enriched in specific nutrients on *Drosophila*
583   development, body fat, and lifespan. *PLoS One* **11**, e0146758 (2016).

25.   F. Demontis, N. Perrimon, Integration of Insulin receptor/Foxo signaling and dMyc activity
during muscle growth regulates body size in *Drosophila*. (2009).

26.   L. Moss-Taylor, A. Upadhyay, X. Pan, M.-J. Kim, M. B. O'Connor, Body size and tissue-
scaling is regulated by motoneuron-derived activinß in *Drosophila melanogaster*. *Genetics*
213, 1447-1464 (2019).

27.   K. J. Livak, T. D. Schmittgen, Analysis of relative gene expression data using real-time
quantitative PCR and the 2− ΔΔCT method. *methods* **25**, 402-408 (2001).

28.   R. Caspi *et al.*, The MetaCyc database of metabolic pathways and enzymes and the
BioCyc collection of pathway/genome databases. *Nucleic Acids Res.* **44**, D471-D480
(2016).

29.   M. Kanehisa, S. Goto, KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids*
Res. **28**, 27-30 (2000).

30.   F. Gutzwiller *et al.*, Dynamics of *Wolbachia pipientis* gene expression across the
Drosophila melanogaster life cycle. *G3: Genes| Genomes| Genetics* **5**, 2843-2856 (2015).

31.   B. R. Graveley *et al.*, The developmental transcriptome of *Drosophila melanogaster*.
Nature **471**, 473-479 (2011).

32.   R Core Team (2014) R: A language and environment for statistical computing. (URL
http://www.R-project.org/, R Foundation for Statistical Computing, Vienna, Austria).

33.   D. Bates, M. Maechler, B. Bolker, S. Walker, lme4: Linear mixed-effects models using
Eigen and S4. *R package version* **1** (2014).

34.   D. Falk, R. Bellamy, M. El Kouni, F. Naguib, A genetic and dietary study of the physiology
of pyrimidine synthesis in *Drosophila melanogaster*. *J. Insect Physiol.* **26**, 735-740 (1980).

35.   A. Sasaki, T. Nishimura, T. Takano, S. Naito, S. K. Yoo, *white* regulates proliferative
homeostasis of intestinal stem cells during ageing in *Drosophila*. *Nature Metabolism* **3**,
546-557 (2021).

36.   N. A. Broderick, B. Lemaitre, Gut-associated microbes of *Drosophila melanogaster*. *Gut*
Microbes **3**, 307-321 (2012).

37.   D. N. Lesperance, N. A. Broderick, Microbiomes as modulators of *Drosophila*
melanogaster homeostasis and disease. *Current Opinion in Insect Science* **39**, 84-90
(2020).

38.   H. P. Harding *et al.*, An integrated stress response regulates amino acid metabolism and
resistance to oxidative stress. *Mol. Cell* **11**, 619-633 (2003).

39.   J. M. Tennessen, C. S. Thummel, Coordinating growth and maturation—insights from
Drosophila. *Curr. Biol.* **21**, R750-R757 (2011).

40.   S. Layalle, N. Arquier, P. Léopold, The TOR pathway couples nutrition and developmental
timing in *Drosophila*. *Dev. Cell* **15**, 568-577 (2008).

41.   C. Géminard, E. J. Rulifson, P. Léopold, Remote control of insulin secretion by fat cells in
Drosophila. *Cell Metabolism* **10**, 199-207 (2009).

42.   J. Colombani *et al.*, A nutrient sensor mechanism controls *Drosophila* growth. *Cell* **114**,
739-749 (2003).

43.   M. J. Texada, T. Koyama, K. Rewitz, Regulation of body size and growth control. *Genetics*
216, 269-313 (2020).

44.   D. Falk, R. Bellamy, M. el Kouni, N. F, A genetic and dietary study of the physiology of
pyrimidine synthesis in *Drosophila melanogaster*. *J. Insect Physiol.* **26**, 735-740 (1980).

45.   D. R. Evans, H. I. Guy, Mammalian pyrimidine biosynthesis: fresh insights into an ancient
pathway. *J. Biol. Chem.* **279**, 33035-33038 (2004).

46.   J.-J. Chen, M. E. Jones, The cellular location of dihydroorotate dehydrogenase: relation
to de novo biosynthesis of pyrimidines. *Archives of Biochemistry and Biophysics* **176**, 82-
90 (1976).

47.   J. M. Rawls, Genetic complementation and enzyme correlates at the locus encoding the

last two steps of de novo pyrimidine biosynthesis in *Drosophila melanogaster*. *Molecular*

and General Genetics MGG **184**, 174-179 (1981).

48.   M. Löffler, E. A. Carrey, W. Knecht, The pathway to pyrimidines: The essential focus on

dihydroorotate dehydrogenase, the mitochondrial enzyme coupled to the respiratory

chain. *Nucleosides Nucleotides Nucl. Acids* **39**, 1281-1305 (2020).

49.   A. Bajgar, T. Dolezal, Extracellular adenosine modulates host-pathogen interactions

through regulation of systemic metabolism during immune response in Drosophila. *PLoS*

Path. **14**, e1007022 (2018).

50.   M. Fang, Y. Chai, G. Chen, H. Wang, B. Huang, N6-(2-hydroxyethyl)-adenosine exhibits

insecticidal activity against Plutella xylostella via adenosine receptors. *PLoS One* **11**,

e0162859 (2016).

51.   J. H. Schroader *et al.*, Disease-associated inosine misincorporation into RNA hinders

translation. *Nucleic Acids Res.* **50**, 9306-9318 (2022).

52.   A. R. I. Lindsey, Sensing, Signaling, and Secretion: A review and analysis of systems for

regulating host interaction in *Wolbachia*. *Genes* **11**, 813 (2020).

53.   K.-O. Cho, G.-W. Kim, O.-K. Lee, *Wolbachia* bacteria reside in host Golgi-related vesicles

whose position is regulated by polarity proteins. *PLoS One* **6**, e22703 (2011).

54.   P. M. White *et al.*, Reliance of *Wolbachia* on high rates of host proteolysis revealed by a

genome-wide RNAi screen of *Drosophila* cells. *Genetics* **205**, 1473-1488 (2017).

55.   N. A. Moran, G. R. Plague, J. P. Sandström, J. L. Wilcox, A genomic perspective on

nutrient provisioning by bacterial symbionts of insects. *Proc. Natl. Acad. Sci.* **100**, 14543-

14548 (2003).

56.   N. A. Moran, Symbiosis. *Curr. Biol.* **16**, R866-871 (2006).

 Supplemental Tables and Figures

661 **Table S1. *Drosophila melanogaster* stocks used in this study.**

BDSC RRID	Complete Genotype	*Wolbachia* infection
41694	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS02259}attP2/TM3, Sb[1]	positive
60086	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC05080}attP40	positive
43296	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS02669}attP40	positive
51492	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC03244}attP2	positive
36686	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS01574}attP2	positive
62241	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC05248}attP40	positive
34347	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMS01336}attP2	positive
53332	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC03561}attP40	positive
60019	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC05011}attP40	positive
53339	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC03568}attP40	positive
51801	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.HMC03359}attP40	positive
55183	y[1] sc[*] v[1] sev[21]; P{y[+t7.7] v[+t1.8]=TRiP.HMC03871}attP40	positive
3954	y[1] w[*]; P{w[+mC]=Act5C-GAL4}17bFO1/TM6B, Tb[1]	positive
31603	y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF01355}attP2	negative
1936	Dhod[2] p[p]	positive

664 **Table S2. qPCR primers used in this study.**

Target	FlyBase ID	Forward Primer Sequence (5'-3')	Reverse Primer Sequence (5'-3')
ftsZ	n/a	TTTTGTTGTCGCAAATACCG	CCATTCCTGCTGTGATGAAA
18S	FBgn0085802	CGAAAGTTAGAGGTTCGAAGGCGA	CCGTGTTGAGTCAAATTAAGCCGC
Prat2	FBgn0041194	AATAGTGACCAGTTTGGGCAAG	CCCAGGTTACCCTTTAACTTCC
ade2	FBgn0000052	CAGCCAGGTTTGTTCCTCTG	TAGCTGCTCGTCAAAGCTCG
ade5	FBgn0020513	CCCGAGGGCTACAGGTTCT	AACTTGGCGGAGACAATTTGC
CG11089	FBgn0039241	CTAGCTTGCGTGGCGCTG	CTGGATAAGATCGAAGCCCTGC
su(r)	FBgn0086450	CATCAAGCACACCACATTGTCG	GTGGGACAGGACTTTTGGCA
Dhod	FBgn0000447	TACGGCATACAAGAACCAGGA	TCATCATGGTACTGTGAAACCG
r-l	FBgn0003257	TGTGATGCAAACCGTATCCGA	GTCCTCGACAATCAGACAGGT

Figure S1. Pupal measurements. Pupal volume was calculated based on length and width (red

lines) and assuming a prolate spheroid shape [$V = (4/3) \pi (width/2)^2 (length/2)$].

Figure S2. *Wolbachia* titer is not responsive to changes in de novo purine biosynthesis

gene expression. (A) qRT-PCR validation of knockdown across four purine loci. Legend: *** p

<0.001; ** p < 0.01; * p < 0.05; ns p = not significant. **(B)** Knockdown of purine synthesis had no

significant impacts on *Wolbachia* titer (gene*knockdown: $F_{3,31} = 0.652$, p = 0.588; gene: $F_{3,31} =$

0.644, p = 0.593; knockdown: $F_{1,31} = 0.593$, p = 0.447).

Figure S3. Validation of knockdown controls. Impact of balancer chromosomes on (A) purine

gene expression, (B) pyrimidine gene expression, and (C) *Wolbachia* titer. **(A)** The TM6B

balancer had no significant impacts on purine gene expression (gene*balancer: $F_{3,24}$ = 1.311, p =

0.2939; gene: $F_{3,24}$ = 1.311, p = 0.2939; knockdown: $F_{1,24}$ = 4.146, p = 0.0529). Importantly, while

the mean expression of the purine loci may have been marginally higher in the presence of the

balancer chromosome, this means that we would underestimate the effects of knockdown due to

an even larger difference between the sibling and knockdown flies (see main text). **(B)** The TM6B

balancer had no significant impacts on pyrimidine gene expression (gene*balancer: $F_{2,18}$ = 0.227,

p = 0.799; gene: $F_{2,18}$ = 0.227, p = 0.799; knockdown: $F_{1,18}$ = 1.180, p = 0.292). **(C)** There was no

significant difference in *Wolbachia* titer due to the balancer chromosome (t-test, t = 1.8404, df =

8, p = 0.1153). Again, the trend towards marginally higher *Wolbachia* titer in the presence of the

balancer only means that we would underestimate the effects of knockdown.