Broadly Neutralizing Antibody Epitopes on HIV-1 Particles are exposed after Virus

Interaction with Host Cells

Priyanka Gadam Rao[a], Gregory S. Lambert[a], and Chitra Upadhyay[a]*

[a]Division of Infectious Disease, Department of Medicine, Icahn School of Medicine at

Mount Sinai, New York, New York, USA

Running Head: Unmasking epitopes for HIV-1 Neutralization

* Address correspondence to chitra.upadhyay@mssm.edu

Abstract Word Count: 250

Importance word count: 136

Text word count: 7397

Keywords: Human immunodeficiency virus; neutralization; antibodies; broadly

neutralizing antibodies (bNAbs); non-neutralizing antibodies (nNAbs); envelope

conformations, glycosylation, lectins, glycans.

Abstract

The envelope glycoproteins (Env) on HIV-1 virions are the sole target of broadly

neutralizing antibodies (bNAb) and the focus of vaccines. However, many cross-reactive

conserved epitopes are often occluded on virus particles, contributing to the evasion of

humoral immunity. This study aimed to identify the Env epitopes that are

exposed/occluded on HIV-1 particles and to investigate the mechanisms contributing to

their masking. Using a flow cytometry-based assay, three HIV-1 isolates, and a panel of

antibodies, we show that only select epitopes including V2i, gp120-g41 interface, and

gp41-MPER are accessible on HIV-1 particles, while V3, V2q, and select CD4bs epitopes

are masked. These epitopes become accessible after allosteric conformational changes are

induced by pre-binding of select Abs, prompting us to test if similar conformational

changes are required for these Abs to exhibit their neutralization capability. We tested HIV-

1 neutralization where virus-mAb mix was pre-incubated/not pre-incubated for one hour

prior to adding the target cells. Similar levels of neutralization were observed under both

assay conditions, suggesting that the interaction between virus and target cells sensitizes

the virions for neutralization via bNAbs. We further show that lectin-glycan interactions

can also expose these epitopes. However, this effect is dependent on the lectin specificity.

Given that, bNAbs are the ideal for providing sterilizing immunity and are the goal of

current HIV-1 vaccine efforts, these data offer insight on how HIV-1 may occlude these

vulnerable epitopes from the host immune response. In addition, the findings can guide the

formulation of effective antibody combinations for therapeutic use.

Importance

The human immunodeficiency virus (HIV-1) envelope (Env) glycoprotein mediates viral

entry, and is the sole target of neutralizing antibodies. Our data suggest that antibody

epitopes including V2q (e.g., PG9, PGT145), CD4bs (e.g., VRC01, 3BNC117) and V3

(2219, 2557) are masked on HIV-1 particles. The PG9 and 2219 epitopes became

accessible for binding after conformational unmasking was induced by pre-binding of

select mAbs. Attempts to understand the masking mechanism led to the revelation that

interaction between virus and host cells is needed to sensitize the virions for neutralization

by broadly neutralizing antibodies (bNAbs). These data provide insight on how bNAbs

may gain access to these occluded epitopes to exert their neutralization effects and block

HIV-1 infection. These findings have important implications for the way we evaluate the

neutralizing efficacy of antibodies and can potentially guide vaccine design.

1. Introduction

The envelope glycoprotein (Env) of human immunodeficiency virus type-1 (HIV-1) is

the virus attachment protein that interacts with the host-cell receptor CD4 and chemokine

receptors CCR5 or CXCR4 to initiate infection (1). HIV-1 displays an average of 7-14 Env

trimers per particle. However, this number varies among isolates (2). Env is synthesized as

a gp160 precursor protein in the endoplasmic reticulum (ER), where signal peptide

cleavage, folding, addition of high-mannose glycans, and trimerization in association with

molecular chaperones takes place (3-6). Once the nascent polypeptide attains its native

folding state, the Env egresses from the ER and translocates to the Golgi apparatus (3, 7-

13). In the Golgi, gp160 is cleaved by host furin-like proteases to generate a transmembrane

gp41 subunit and a non-covalently associated surface gp120 subunit. Three gp120-gp41

heterodimers assemble to form the trimeric functional Env spikes that are then directed to

the plasma membrane for incorporation into virions (3, 11-13). The gp120 subunit

comprises five variable (V1-V5) and five conserved (C1-C5) regions. The V1V2 from each

of the protomers join at the top to form the trimer apex, an immunogenic, structurally

conserved region targeted by some of the most potent HIV-1 bNAbs (14-18). The gp41

subunit comprises the fusion machinery, mediating entry of the virus into the target cell by

facilitating membrane fusion (19, 20). While trimeric Env is expected to be the most

abundant form of Env present on virions, non-functional Env such as dimers, monomers,

and gp41 stumps may also be present (21-23). These nonfunctional Env are

immunodominant and act as decoys to divert the immune response away from vulnerable

conserved epitopes resulting in higher titers of non-neutralizing antibody responses (23,

24), further adding a layer of diversity to the already complex Env protein.

HIV-1 Env is the only viral protein on the surface of HIV-1 particles, and is therefore

a principal target for HIV-1 broadly neutralizing antibodies (bNAbs) that neutralize the

virus and prevent infection of host cells. Passive transfer of various bNAbs has been shown

to protect non-human primates and humanized mice from viral challenge (25, 26). Thus,

bNAbs are being investigated for both therapeutic use and to guide the design of Env-based

immunogens (27, 28). Major efforts are being invested in improving HIV-1 antigen design

and strategies that can stimulate production of antibodies with similarly broad and

neutralizing responses. Several approaches using the trimer-mimic and virus-like particles

have been explored, but eliciting bNAbs via vaccination has proven extremely difficult.

Despite large research efforts, immunogens capable of eliciting bNAbs do not exist yet.

HIV-1 has evolved multiple mechanisms such as high antigenic diversity, conformational

flexibility, and a dense coat of N-linked glycans that masks the conserved epitopes on Env

(29). This masking of conserved epitopes is one of several factors that has impeded the

development of an Env-derived immunogen capable of eliciting bNAbs against HIV-1.

Currently, five regions have been identified on the HIV-1 Env that are the focus of bNAb

elicitation. These include the V2 apex, the V3-N332 supersite, the membrane proximal

external region (MPER), the CD4 binding site (CD4bs), and the gp120-gp41 interface (30-

32). Thus, identifying the Env epitopes that are occluded on virions and understanding the

mechanisms that govern their unmasking is crucial to inform design and development of a

protective HIV-1 vaccine.

Several monoclonal antibodies (mAbs) have been isolated from HIV-1 infected

individuals and vaccinees from clinical trials that have helped to identify and map the

vulnerable Env epitopes. These mAbs are categorized into distinct types based on their

neutralization breadth and potency. The highly potent neutralizing antibodies that can

neutralize broad arrays of HIV-1 isolates are referred as bNAbs. The weak- or non-

neutralizing antibodies (wNAbs, nNAbs) bind to the Env in a manner that is not efficient

at blocking virus infection. The bNAbs known to date have been isolated from HIV-1

infected individuals, and elicitation of these kind of mAbs via vaccination has not yet been

achieved. In contrast, nNAbs or wNAbs, including those that target the V1V2 and V3

regions, are readily elicited via various Env vaccines as well as during HIV-1 infection

(33-42). In fact, the early antibody responses are typically directed against the gp41 domain

of Env and later evolves to include gp120 (43). Only after several years of infection do a

fraction of infected individual develop bNAbs (44). The HIV-1 antibodies are further

categorized based on the distinct Env region that they target, such as V1V2, V3, CD4bs,

gp41-MPER, and the gp120-gp41 interface (45). While most mAbs are conformation-

dependent, some also recognize linear epitopes. For example, the V1V2 mAbs are

categorized into three types (V2i, V2p and V2q) based on their binding mode (46, 47). V2i-

specific mAbs, (e.g., 697, 830A, and 2158) recognize V1V2 when its V2C region is in a

β-strand configuration, as determined by X ray crystallography; the epitope region of these

V2i mAbs is discontinuous, highly conformational, and overlaps the α4β7 integrin-binding

motif (46). The V2p mAbs (eg., CH58 and CH59) were isolated from RV144 vaccinees

and target a linear peptide region on V2; these mAbs recognize V2 when the V2C strand

region is in an α-helix and extended coil conformation (48-50). The epitope region

recognized by V2q mAbs (e.g., PG9, PG16, and PGT145) includes two N-linked glycans

(N156 and N160) and consists of the V2C in its β-strand configuration (15, 51-53); only

V2q mAbs are bNAbs. Another key distinction is whether the mAbs recognize Env in the

context of functional trimers for example, PG9 and PG16 (21). The conformational

diversity of Env, and the presence of functional and non-functional Envs on virions, can

significantly influence both antibody efficacy and their elicitation.

In this study, we used a panel of Abs specific to different regions of Env to identify

the overall antigenic makeup of Env on virus particles. We established a flow cytometry-

based assay to evaluate the neutralizing and non-neutralizing epitopes that are masked or

accessible on virions, and to identify the mechanisms that can unmask the occluded

epitopes. We show that virus interaction with host cells induces changes in Env, exposing

the epitopes and allowing antibody binding and virus neutralization. We further show that

a virus-mAb pre-incubation step, widely used in neutralization assays in the field, does not

appear to be required for neutralization by these bNAbs. Similar exposure of epitopes can

also be achieved by lectin and Env glycan interactions. However, this activity is dependent

on the lectin specificity. These findings have important implications for the way we

evaluate the neutralizing efficacy of antibodies, and provides important clues for improving

vaccine design and the formulation of effective antibody combinations for therapeutic use.

2. Materials and Methods

Cell lines: HEK293T/17 cells were obtained from the American Type Culture Collection

(ATCC, Manassas, VA). The following reagent was obtained through the NIH HIV

Reagent Program, Division of AIDS, NIAID, NIH: TZM-bl Cells, ARP-8129, contributed

by Dr. John C. Kappes, Dr. Xiaoyun Wu, and Tranzyme Inc. (54). For all experiments,

HEK293T/17 cells (293T) were used to produce infectious HIV-1 viruses and the TZM.bl

cell line was used to assay virus infectivity. TZM.bl cell line is derived from HeLa cells

and is genetically modified to express high levels of CD4, CCR5 and CXCR4 and contains

reporter cassettes of luciferase and β-galactosidase that are each expressed from an HIV-1

LTR. The 293T and TZM.bl cell lines were routinely sub-cultured every 3 to 4 days by

trypsinization and were maintained in Dulbecco's Modified Eagle's Medium (DMEM)

supplemented with 10% heat-inactivated fetal bovine serum (FBS), HEPES pH 7.4 (10

mM), L-glutamine (2 mM), penicillin (100 U/ml), and streptomycin (100 μg/ml) at 37°C

in a humidified atmosphere with 5% CO_2.

Plasmids: A full-length transmitted/founder (T/F) infectious molecular clone (IMC) of

pREJO.c/2864 (REJO, ARP-11746) was obtained through the NIH HIV Reagent Program,

Division of AIDS, NIAID, NIH, contributed by Dr. John Kappes and Dr. Christina

Ochsenbauer (55). REJO is a tier 2, clade B, T/F isolate. IMCs of SF162 (chronic; clade

B, chronic, tier 1B) and CMU06 (acute; clade AE, acute, tier 2) were generated by cloning

the Env into a pNL4.3 backbone to construct pNL-CMU06 and pNL-SF162, respectively

(56). The human CD4 expression plasmid was kindly provided by Prof. F. Kirchhoff (57).

Antibodies and lectins: The following antibody reagents used in this study were obtained

through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: Anti-HIV-1

gp120 monoclonal PG9, PG16, PGT145, PGT121, PGT128, from IAVI (58); anti-HIV-1

gp120 monoclonal CH59 from Drs. Barton F. Haynes and Hua-Xin Liao (49); anti-HIV-1

gp120 monoclonal VRC01 from Dr. John Mascola (59); anti-HIV-1 gp120 Monoclonal

b12 from Dr. Dennis Burton and Carlos Barbas (60); anti-HIV-1 gp120 monoclonal

3BNC117 from Dr. Michel C. Nussenzweig (61); anti-HIV-1 gp120 monoclonal 17b from

Dr. James E. Robinson (62); anti-HIV-1 gp41-gp120 monoclonal 35O22, from Drs. Jinghe

Huang and Mark Connors (63); anti-HIV-1 gp41-gp120 monoclonal PGT151 from Dr.

Dennis Burton; anti-HIV-1 gp41 monoclonal 2F5 and 4E10 from Polymun Scientific (64).

The V2i and V3 mAbs were obtained from the laboratory of Dr. Susan Zolla-Pazner (65-

72). An irrelevant anti-anthrax mAb 3685 (73) was used as a negative control. The lectin

Concanavalin A (ConA) was purchased from Vector Laboratories. The lectin Griffithsin

was obtained through NIH HIV Reagent Program, Division of AIDS, NIAID, NIH,

contributed by Drs. Barry O'Keefe and James McMahon.

Virus production and purification: Infectious viruses were generated by transfecting

293T cells with pREJO, pNL-SF162 and pNL-CMU06 plasmids using jetPEI transfection

reagent (Polyplus, New York, NY) (74). Supernatants were harvested after 48 hours and

clarified by centrifugation and 0.45μm filtration. Virus infectivity was assessed on TZM.bl

cells as described (56, 74). Briefly, serial two-fold dilutions of virus stock in 10% DMEM

were incubated with TZM.bl cells (in duplicates for each dilution) in half-area 96-well

plates in the presence of DEAE-dextran (12.5 μg/ml) for 48 hours at 37°C. Virus infectivity

was measured by β-galactosidase activity (Promega, Madison, WI). Virus stocks were

concentrated (20X) by ultracentrifugation over 20% (w/v) sucrose in 1X phosphate

buffered saline (PBS) at 28,000 RPM for 2 hours in an SW-28 swinging bucket rotor

(Sorvall, Thermofisher Scientific). Supernatants were decanted and pellets dried briefly

before resuspension in PBS. Inactivation of virions was carried out using Aldrithiol-2 (AT-

2) (21, 75). Briefly, 125 ul of sucrose-purified virus was incubated with 0.5 mM AT-2 in

DMSO for 2 hours at 37°C, followed by centrifugation at 13,000 rpm for 2 hours. The

supernatant was discarded, and the pellet re-suspended in 125 ul PBS. Inactivation was

confirmed by measuring infectivity in TZM.bl cells and Env content was checked by

Western blotting.

Western blotting: To quantify and monitor the expression of Env in each virus preparation

Western blot analyses were performed. The sucrose-purified virus particles were lysed,

resolved by SDS-PAGE on 4–20% tris-glycine gels (Bio-Rad, Hercules, CA), and blotted

onto membranes, which were then probed with antibodies. A cocktail of anti-human anti-

gp120 MAbs (anti-V3: 391, 694, 2219, 2558; anti-C2: 841, 1006; anti-C5: 450, 670, 722;

1μg/ml each) was used to detect Env. MAb 91-5D (1μg/ml) was used to detect Gag p24.

Membranes were developed with Clarity Western ECL Substrate (Bio-Rad, Hercules, CA)

and imaged by a ChemiDoc Imaging System (Bio-Rad, Hercules, CA). Purified

recombinant gp120 and p24 proteins were also loaded at a known concentration as controls

and quantification standards. Band intensities were quantified using the Image Lab

Software Version 5.0 (Bio-Rad, Hercules, CA).

Coupling of fluorescent beads to virions: The coupling strategy used in this study is a

standard method widely used in the Luminex based multiplex immunoassays and is

published previously (76-79). We and others have used this protocol (xMAP Technology,

Luminex Corp) and carboxylated beads to assess the Fc-mediated ADCP function of anti-

HIV-1 and anti-COVID-19 antibodies (80-82). Here we applied this approach for coupling

the virus to the beads. Sucrose-purified, inactivated virions were covalently coupled to

carboxylate-modified microspheres (1.0 µm) using a two-step carbodiimide reaction with

the xMAP Ab Coupling (AbC) Kit according to manufacturers' instructions (Luminex,

Austin, TX). Carboxylated beads purchased from Thermo Fisher (cat# F8823), were

coupled to 125 µl of 20X concentrated virus preparations (~36.4x10^9 beads per reaction).

Briefly, the stock microspheres were vortexed and sonicated to resuspend and disperse the

microspheres and 12 µl was transferred to a tube containing 1200 ul of 1% BSA/1X PBS

(per virus). The microspheres were washed twice with 500 µl of activation buffer followed

by vortexing and sonication after each step. The microspheres were activated with 250 µL

of activation buffer, 50 µL of 50 mg/ml Sulfo-NHS (N-hydroxysulfosuccinimide), 50 µL

of 40 mg/mL ethyl dimethylaminopropyl carbodiimide hydrochloride (EDC) and

incubated for 20 min at room temperature with end-to-end rotation. The microspheres were

washed three times in activation buffer, then incubated with AT-2 inactivated virus in

activation buffer for 2 hours at room temperature. We typically used a volume of 20X

concentrated virus that equals to ~175 ng total. The microspheres were subsequently

washed and resuspended in 1.2 mL of 0.1% BSA/PBS and stored at 4°C until ready to use.

Virus-associated Env binding assay (VAEBA): The coupled microspheres were

dispensed into 96-well plate (10 µl/well) and blocked with 100 ul of 3% BSA for 30 mins

at 4°C. Plates were centrifuged at 2000 x g for 10 minutes and the supernatant was

discarded. The microspheres were incubated with serially diluted mAbs for 30 min at 37°C

followed by addition of anti-human-biotin (1:500 in 0.1% BSA/PBS) and incubated in the

dark for 30 mins. Plates were washed 3 times with PBS and incubated with Streptavidin-

PE (1:1000 in 0.1% BSA/PBS) for 30 min followed by resuspension in 200µl of 0.5%

paraformaldehyde. Plates were washed 3 times with PBS after each step. Analysis was

done with Attune NxT flow cytometer, and $\geq$30,000 events were collected in the

phycoerythrin (PE)+ gate. Data analysis was carried out using FCS-Express software as

follows: FITC positive microspheres were selected from a plot of forward-area vs. FITC

(FSC-A/FITC-A) from which doublets were excluded in a forward scatter height vs

forward scatter area plot (FSC-H/FSC-A). Geometric mean fluorescent intensity (MFI)

values of PE+ beads representing anti-Env-stained virus particles coupled to FITC

microspheres, were determined and plotted. Background MFI, as determined from

microspheres stained without primary antibodies was subtracted from all Env-mAb pairs.

Area under the titration curves was also calculated for select data sets. Unless otherwise

noted in the figure legends the mAbs were tested at the following concentrations: 697,

830A, 2158, 3685 at 100 µg/ml; 2219, 2557 at 50 µg/ml; PG9, PG16, PGT145, CH59,

VRC01, 3BNC117 at 25 µg/ml; 4E10 at 20 µg/ml; PGT151 at 10 µg/ml; NIH45-46 at 5

µg/ml.

For time-dependent binding studies, the assay was performed as above using biotinylated

mAbs that were incubated with the virus-coupled beads for different times (0, 15, 30, 45,

60 and 75 min) followed by Streptavidin-PE. Data was acquired by Attune NxT flow

cytometer, as above.

For conformational epitope exposure in response to mAbs, virus coupled microspheres

were pre-incubated with titrated mAbs for 30 min at 37°C. Plates were centrifuged and

further incubated with biotinylated mAbs (PG9 or 2219) for 30 min at 37°C followed by

Streptavidin-PE. Data was acquired by Attune NxT flow cytometer, as above.

For virus-binding assays in the presence of 293T cells +/- human CD4, we transfected 293T

cells with plasmids expressing full-length functional human CD4 (57), as below. Cells

were harvested 24 hours post-transfection using trypsin-free cell-dissociation buffer,

stained with Live/Dead fixable Aqua stain and labelled with Vybrant DiD cell labelling

solution (Thermo Fisher Scientific) for 30 minutes at 4°C. Cells were distributed in 96 well

V-bottom plates followed by addition of virus-coupled microspheres and serially diluted

antibodies, and incubated for 30 minutes at 4°C. Binding was detected using biotinylated

anti-human IgG and Streptavidin-PE. Data was acquired by Attune NxT flow cytometer,

as above. Analysis was carried out using FCS-Express software using two gating strategies

as follows: 293T cells and virus-coupled microspheres were selected from a plot of

forward-area vs. side scatter-area (FSC-A/SSC-A). Gating strategy 1: DiD labelled cells

were gated for live cells by selecting for Aqua-negative gating, followed by gating on cells

that are positive for FITC (virus-coupled FITC microspheres) and geometric mean

fluorescent intensity (MFI) of PE+ events, representing anti-Env-stained beads, were

quantified. Gating strategy 2: FITC positive virus-coupled microspheres were gated for

DiD positive cells, live cells were selected by Aqua-negative gating, and geometric mean

fluorescent intensity (MFI) of PE+ beads/cells, representing anti-Env-stained beads, were

quantified.

For conformational epitope exposure in response to lectin ConA, virus coupled

microspheres were pre-incubated with titrated amounts of ConA for 30 min at 37°C. Plates

were centrifuged and further incubated with mAbs (830A, PG9, 2219, NIH 45-46, VRC01,

and negative control 3685) at one dilution for 30 min at 37°C followed by Streptavidin-

PE. Data was acquired by Attune NxT flow cytometer, as above.

Cell-associated Env binding assay. An assay to detect antibody binding to cell surface

expressed Env was performed as described (56, 83, 84). Briefly, monolayers of 293T cells

(4×10^6) seeded in 100-mm tissue culture dishes were transfected with 20 μg of gp160

expression plasmid using jetPEI (DNA:jetPEI ratio of 1:3) following manufacturer's

instructions (Polyplus, New York, NY). Transfected cells were incubated for 24 hours at

37°C, washed with PBS, detached with trypsin-free cell-dissociation buffer, and

resuspended in PBS containing 2% BSA. Cells were stained with Live/Dead fixable Aqua

stain and distributed into 96-well V-bottom tissue culture plates (5×10^4/well) for individual

staining reactions. Cells were incubated with mAbs at concentrations detailed in figure

legends. For detection of mAb binding, biotinylated goat anti-Human IgG Fc (1:500)

followed by Streptavidin-PE (1:1000) was used. The cells were washed 3X with PBS-B

(PBS plus 1% BSA) after each step and all incubation steps were performed on ice for 30

min. Cells were analyzed with a BD Fortessa flow cytometer, and >30,000 events were

collected in the PE+ gate. Analysis was carried out using FCS-Express software as follows:

293T cells were selected from a plot of forward-area vs. side scatter-area (FSC-A/SSC-A)

from which doublets were excluded in a forward scatter height vs forward scatter area plot

(FSC-H/FSC-A). Live cells were selected by Aqua-negative gating, and geometric mean

fluorescent intensity (MFI) of PE+ cells, representing anti-Env-stained cells, was

quantified. Background MFI, as determined from cells stained without primary antibodies,

was subtracted from all Env-mAb pairs.

Lectin Capture Enzyme-Linked ImmunoSorbent Assay (ELISA): The relative binding

of mAbs to Env on virions was also measured by a lectin capture ELISA. Briefly, half-area

high-binding ELISA plates (Costar) were coated with five-fold titrated amounts of ConA

lectin (50 – 0.08 μg/ml in PBS) for 1 hour at 37°C. Sucrose pelleted REJO virus was AT-2

inactivated as above and added to the ConA coated plate. The plate was incubated for 1

hour at 37°C, washed, and blocked with 5% non-fat dry milk in whey solution for 1 hour

at room temperature or overnight at 4°C. Antibodies were incubated with captured virus at

a single dilution (830A, 2219, PG9, 3685 at 50 μg/ml; VRC01 at 25 μg/ml and NIH45-46

at 2.5 μg/ml), followed by biotinylated anti-human IgG and Streptavidin-HRP. The bound

mAbs were detected with SuperSignal ELISA Pico Chemiluminescent Substrate (Thermo

Fisher Scientific) and relative luminescence units (RLUs) were measured using a BioTek

Cytation luminometer. For the lectin Griffithsin (GRFT), plates were coated with GRFT at

50 μg/ml, incubated with virus particles, and probed with five-fold serially diluted mAbs

(830A, 2219, PG9, 3685 at 50 μg/ml; VRC01 at 25 μg/ml and NIH45-46 at 2.5 μg/ml).

The bound mAbs were detected as above. The plates were washed 3X with PBS-T (PBS

plus 0.05% Tween 20) after each step.

Neutralization assay. Virus neutralization was measured using a β-galactosidase-based

assay (Promega) with TZM.bl target cells (85). Serial dilutions of mAbs were added to the

virus in half-area 96-well plates (Costar, Corning, NY) and incubated for designated time

periods (0 min and 60 min) at 37°C. TZM.bl cells were then added along with DEAE-

dextran (12.5 μg/ml; Sigma). After a 48-hour incubation for each assay condition, at 37°C

and in a 5% CO2 incubator, the β-galactosidase activity was measured. Each condition was

tested in duplicate. Assay controls included replicate wells of TZM.bl cells alone (cell

control) and TZM.bl cells with virus alone (virus control). The highest antibody

concentrations tested were based on known neutralization titers and varied per mAb. PG9

and 2219 were analyzed in the range of 50 μg/ml to 0.78 μg/ml, PGT145 was analyzed in

the range of 5 μg/ml to 0.001 μg/ml, and NIH45-46 was analyzed in the range of 2.5 μg/ml

to 0.004 μg/ml, while 830A and 3685 were tested in the range of 100 μg/ml to 0.006 μg/ml.

Percent neutralization was determined on the basis of virus control under the specific assay

conditions. The virus input corresponded to titrated stocks yielding 150,000 to 200,000

relative luminescence units (RLU).

Statistical analysis: All statistical analyses were performed with GraphPad Prism 9.4.1

(GraphPad, San Diego, CA). ANOVA and Mann-Whitney t-tests were performed as

appropriate and are mentioned in the figure legends.

Results

Detection of REJO Env on HIV-1 particles: To analyze the representation of Env

epitopes on virus particles, we established a flow cytometry-based assay. The assay

developed is based on (a) coupling the virions to fluorescent microspheres; (b) staining

virions immobilized on microspheres with antibodies targeting different epitopes; (c)

staining the resulting complex with biotinylated anti-human IgG and streptavidin-PE; and

(d) detecting Env-Ab interaction via flow cytometry (Fig. 1A).

Initially, we used REJO—a T/F, clade B, tier 2, HIV-1 isolate—and select

monoclonal antibodies (mAbs) that recognize the conformational CD4bs (NIH45-46) and

linear V3 (2219) epitopes on Env. A non-HIV-1 mAb 3685 was also used as a negative

control. To optimize the amount of sucrose-pelleted virus required to couple with the

microspheres, we tested four different amounts (25, 50, 100 and 125 μl corresponding to

~35, 70, 142 and 177 ng total Env, respectively) of 20X concentrated virus preparations.

The Env content was measured by Western blot (Fig. S1) as in (56, 74).

At 125 μl of virus, the binding of positive control mAb NIH45-46 was significantly

enhanced while the negative control mAb 3685 background binding levels were low (Fig.

1B). Thus, based on the fold AUC change over the non-HIV-1 mAb 3685, we elected to

use 125 μl of 20X concentrated virus preparation, equivalent to 177 ng Env. At this amount,

the negative control mAb stained the virions minimally (1.7%) (Fig. 1C). The mAb NIH45-

46 bound 84% of REJO virions, while only 1.8% of particles bound to V3 mAb 2219 (Fig.

1C). Next, we used the assay to analyze the Env epitopes present on the REJO virus

particles using the extended mAb panel (Table 1) that covers most of the major Env

domains.

The REJO virus-bound microspheres were treated with titrated amounts of mAbs and the

assay was conducted as outlined in the Methods section. All three V2i mAbs (697, 830A,

and 2158) bound to Env on virions, with mAb 830A exhibiting the strongest binding (Fig.

2A-C, S2). In contrast, binding of trimer-preferring V2q bNAbs (PG9, PG16, and PGT145)

and peptide recognizing V2p mAbs (CH58 and CH59) was minimal and similar to the

negative control mAb 3685 (Fig. 2A-C). Binding of V3 specific mAbs 2219 and 2557 also

was similar to 3685 (neg control mAb) (Fig. 2). Recognition of CD4bs epitopes by most

mAbs, except NIH45-46, was also negligible. The gp120-gp41 interface (PGT151) and

gp41 MPER mAbs (4E10) displayed efficient binding. Higher binding of MPER specific

4E10 mAb may be either due to lipid cross-reactivity, increased exposure of the gp41 base,

or presence of gp41 stumps (21, 86, 87). Notably, PGT151 which is known to be trimer-

specific and binds only to properly formed, cleaved trimers, had comparable binding to the

V2i mAb 830A (88).

To compare the relative levels of each epitope exposed on virus particles, we

calculated AUC values from the titration curves. Figure 2B represents the overall

presentation of different epitopes on the virions as measured by mAb binding. A similar

pattern was also observed when geometric mean fluorescent intensities (MFI) of mAbs

tested at the highest dilution were plotted (Figure 2C).

Pattern of Env epitope exposure on virions from different clades, tiers, or infection

stage: We next tested two other HIV-1 isolates: SF162 (chronic; clade B, tier 1) and

CMU06 (acute; clade AE, tier 2) using the same panel of mAbs. In case of SF162 virions,

the V2i mAbs 697 and 830A showed greater levels of binding than 2158 (Fig 3, S3). As

seen with REJO, V2q, V2p, and V3 mAbs showed negligible binding to SF162 virions and

were similar to negative control mAb 3685 (Fig 3, S3). All CD4bs mAbs tested showed

binding to SF162 virions that was above background level. In case of CMU06, a similar

pattern was observed with mAbs 830A, PGT151, and 4E10 showing highest binding levels,

while little to no binding was seen with the mAbs targeting the V2q and V3 epitopes (Fig

4, S4). Negligible binding was seen with the negative control mAb 3685. Overall, the

binding strength of mAbs to the SF162 Env on virions was higher compared to CMU06

and REJO. The epitopes recognized by mAbs 830A, PGT151, and 4E10 were consistently

accessible on all three viruses tested in this study. Differences in the recognition of the

CD4bs mAbs among the three isolates were also evident - mAbs 3BNC117 and VRC01

showed better binding to SF162 than the other two viruses. The CD4bs mAb NIH45-46 is

a more potent clonal variant of VRC01 with high sequence and structural similarities to

VRC01, but has a distinct mode of binding to gp120 (89). The potency of the CD4bs mAbs

tested here is in the order NIH45-46 > 3BNC117 > VRC01 (90). Differences in binding,

in that order, are evident in the binding data shown in Fig 2-4.

Binding profile of mAbs to Env expressed on the cell surface is different from Env on

virions: We next sought to compare the binding of mAbs to virions with binding to

transiently expressed Env on cells. We transfected 293T cells with gp160 expression

plasmids and probed Env expressed on the cell surface with the mAbs in Table 1. A

different profile was observed when staining of Env on virus particles was compared to

Env expressed on transfected cells (Fig. 5, S5, S6, and S7). All mAbs tested bound to Env

expressed on cells, while this was not the case with HIV-1 particles (Fig. 2-4). The V2q,

V2p, and V3 mAbs which showed negligible binding to Env on virions displayed efficient

binding to REJO and CMU06 Env expressed on 293T cell surface. Thus, Env recognized

by V2q mAbs are efficiently expressed however, the mechanism of Env incorporation may

affect its conformation and expression at the virion surface. In addition, Env on producer

cells may also be populated by uncleaved Env monomers at the cell surface that are not

incorporated into virions (91), which may also account for the observed differences in

binding. The CD4bs mAbs also displayed greater binding to cell-surface expressed Env

compared to Env on virus particles. Env from all three isolates also bound efficiently to

PGT151, which binds only to properly formed, cleaved trimers (92). Thus, the binding

profile of mAbs to Env differs based on the location of the Env (cell surface versus virus

surface).

Prolonging the mAb-virion interaction time does not unmask the occluded epitopes:

Binding of antibodies to Env is essential in order for them to neutralize the virus. In a

typical standard neutralization assay, virus and mAbs are pre-incubated for 1 hour at 37°C

prior to adding the target cells. In our Env binding assay, mAbs-virus were incubated for

only 30 min at 37°C, prompting us to test if increasing the mAb-virus interaction time

would facilitate binding, especially for the mAbs (V2q, V3, and CD4bs) that showed little

to no binding after 30 minutes.

To test this idea, microspheres coupled with REJO virus were incubated with select

mAbs: V2i (830), V2q (PG9), V3 (2219), CD4bs (NIH45-46), or control mAb (3685) for

various periods of time (0 to 75 min). The binding of mAbs to the Env on REJO virus was

measured by flow cytometry (Fig. 6, S8). Little binding to REJO Env was observed for all

mAbs when no incubation (0 min) was allowed, but a significant linear increase in binding

of 830A and NIH45-46 mAbs was detected over time (Fig. 6, S8). The V2q mAb PG9 and

V3 mAb 2219 showed no increase in binding to REJO Env at any time point, indicating

that epitopes of these mAbs remained occluded on the virus even after 75 min of incubation

(Fig. 6). As expected, no binding was detected for the negative control mAb 3685 at any

time point.

Masked Env epitopes can be exposed by pre-binding of other mAbs. We wondered if

microsphere-virion coupling method might be contributing to the inability to detect binding

of mAbs to these epitopes. If that stands correct then binding of V2q mAb PG9 or V3 mAb

2219 will remain low on virions even under conditions that are known to conformationally

unmask these epitopes (93, 94). We tested this idea by pre-treating the virus coupled

microspheres with select mAbs followed by detecting the binding of V2q (PG9) and V3

(2219) mAbs. We incubated the microsphere-coated virions first with mAbs 697, 830A,

2158, 447, or 2219 to induce structural changes in Env. After washing away the unbound

mAb, we probed for V2q epitope exposure using biotinylated PG9 mAb. Non-biotinylated

PG9 and non-HIV-1 mAb 3685 were used as controls. As seen in Fig. 7 and S9, binding

of all three V2i mAbs (697, 830A, and 2158) presumably induced an allosteric

conformational change that exposed the PG9 binding epitopes. This concurs with the

published study showing that V2i 830A and 2158 increase the binding of PG9 on A244

gp120 (94). In contrast, the V3 mAbs (447 and 2219) did not alter PG9 binding and were

comparable to the negative control mAb 3685. Since PG9 epitopes are also occluded on

virions, negligible binding was seen in this control as well. Thus, these data further support

the notion that PG9 epitopes are conformationally masked on virions. We next evaluated

if V3 epitopes can be similarly exposed. The V3 loops are tucked beneath V1V2, thus Abs

that induce any movement in the V1V2 loops should expose the V3 epitopes. We made use

of the following CD4bs mAbs: the bNAb NIH45-46 that binds strongly to Env on virions,

the non-neutralizing mAb 654 that is known to expose V3 loops, and the V2i mAb 830A

(93, 94). Non-biotinylated 2219 and 3685 were used as controls. Binding of 830A and 654

exposed the V3 epitopes, allowing the binding of biotinylated 2219 mAb. In contrast,

NIH45-46, despite binding well to REJO virions, did not induce changes in Env that might

expose the 2219 epitopes (Fig 7, S9). The 2219 epitopes are not available on virions in the

absence of allosteric alterations; thus, no binding was detected in this control.

Neutralization of REJO and CMU06 viruses by bNAbs: The findings from above

suggest that allosteric changes, but not time, can allow exposure of the epitopes targeted

by the V2q mAb PG9. However, given the reported neutralization breadth and potency of

PG9, the little to no binding of PG9 to the Env on virion surface was perplexing (58, 59).

This led us to speculate that perhaps the interaction of virus with the target cells induces

similar changes in Env that allows bNAbs such as PG9 to bind to their epitopes. In such a

case, we would expect that neutralization of HIV-1 with PG9 should be similar in the

presence or absence of a virus-mAb pre-incubation step.

Indeed, as seen in Fig. 8, V2q bNAbs PG9 and PGT145 were able to neutralize REJO

virus with equal efficacy independent of a pre-incubation step. Among the CD4bs mAbs,

neutralization by 3BNC117 and NIH45-46 was also similar at both t = 0 and 60 minutes,

while VRC01 had slightly better neutralization at t =60 min (AUC 111) compared to t = 0

(AUC 85); however, the difference was not statistically significant. These data suggest that

interaction of virus with target cells appears to induce localized antigenic changes in Env

on virions, allowing for bNAbs to latch onto their epitopes. This is the first time, to the best

of our knowledge that the V2q and CD4bs mAbs tested here are shown to require pre-

interaction of virus with target cells to exhibit their neutralization effect. Interestingly,

mAbs such as 830A and 4E10 show strong binding to their respective epitopes on virions.

However, these mAbs fail to neutralize the REJO virus. One possible explanation for this

observation may be that the 830A and 4E10 epitopes that are available for binding on

virions are associated with non-functional Env. Similar results were also observed when

neutralization of CMU06 virus was compared at $t = 0$ vs 60 min (Fig 9). Collectively, these

data highlight the importance of gaining a better understanding of the multiple mechanisms

that are utilized by HIV-1 to avoid interaction with host-mounted Abs and maintain

infectivity. Also, these findings encourage the idea to re-evaluate the assay formats that are

widely used in the field.

Influence of virus-cell interaction on Env epitope exposure: To further verify that virus

interaction with host cells can indeed improve the exposure of Env epitopes on virions, we

measured the binding of select mAbs to virion coupled fluorescent microspheres in the

presence of host cells. The TZM.bl cells used for neutralization assays are HIV-1

permissive reporter HeLa cells modified to express the receptor (human CD4) and co-

receptors (human CCR5 and CXCR4), allowing the virus to initiate infection in these cells

(85). In addition, the surface of every eukaryotic cell is known to be covered by a dense

and diverse array of cell surface glycans, the "glycocalyx" (95). Studies have shown that

host cell glycocalyx can capture HIV-1 via specific, non-electrostatic oligomannose-

GlcNAc glycan-glycan interactions to support viral entry (96, 97). To test the idea that

host-cell interaction exposes Env epitopes, we used 293T cells that are non-permissive to

HIV-1 entry for this assay. We transfected 293T cells with a plasmid expressing human

CD4 (57, 98), to study the effect of surface CD4 on Env epitope exposure. In addition, we

used un-transfected 293T cells, to study how glycan-glycan interaction (i.e, host

glycocalyx and Env glycan interactions) may affect epitope exposure on virions. As seen

in Fig 10 and S10, differences in binding of mAbs are evident, particularly for PG9 and

2219 that is relatively higher in the presence of 293T cells, compared to 293T-hCD4 cells.

In our previously published study, we have shown that lectin interaction with HIV-

1 Env glycans can influence the exposure of V2i and V3 epitopes (99). To test if lectins

can also expose the V2q epitopes on virions, we used Concanavalin A (ConA), which is a

carbohydrate-binding protein extracted from jack-bean (*Canavalia ensiformis*) seeds and

binds to the mannose residues of various glycoproteins (100, 101). ConA binds specifically

to the α-mannose/α-glucose moieties on N-linked oligomannose and complex-type glycans

and is not known to bind O-glycans on glycoproteins (102). We incubated the microsphere-

coupled virions first with titrated amounts of ConA followed by select mAbs (830A, PG9,

2219, NIH45-46, VRC01, and 3685) at a single concentration. As seen in Fig 11 and S11,

binding of all anti-HIV-1 mAbs including PG9, VRC01, and 2219 was detected in the

presence of ConA. We further verified these findings using an alternative ELISA-based

approach where we do not couple the virus particles to microspheres. We used ConA to

capture the virus particles and detected the binding of different mAbs to captured virions

by ELISA. As seen in Fig 12A, virions captured by ConA also displayed improved

exposure of the PG9 and VRC01 epitopes, similar to the pattern observed in the flow

cytometry based virus-binding assay (Fig 11). In contrast, another lectin, Griffithsin

(GRFT), which is specific for α1–2 mannose on oligomannose glycans, exposed 2219 and

VRC01 but not PG9 epitopes (fig 12B). Thus, these data further support the notion that

HIV-1 masks the vulnerable epitopes on its Env and different factors (e.g., glycan-glycan

and lectin-glycan interactions) may modulate the antibody accessibility of these epitopes

on the virus particles.

4. Discussion

This study focused on investigating the antigenic landscape of HIV-1 Env on the

infectious virus particles that are encountered by the host immune system. We developed

a flow cytometry-based bead assay to quantitate the relative levels of Env epitopes that are

exposed on HIV-1 particles. Although bead-based assays have been reported before, our

approach varies from previous assays in that it alleviates the requirement of producing

fluorescent virus particles (103). In addition, other assays make use of two anti-HIV-1 Abs,

one Ab to capture the virus particles and other Ab to detect binding (104). As shown in this

study, application of two Abs may skew the results and fail to differentiate between the

actual mAb binding and binding due to the conformational changes in Env influenced by

the capture Ab. Furthermore, the assay described in this study can be easily translated to

study other viruses (e.g., SARS-CoV-2, Influenza etc.,) and antigens in different formats

(e.g. gp120, gp140 etc).

Findings from this study show that epitopes targeted by mAbs like 830A (V2i),

PGT151 (gp120-gp41 interface) and 4E10 (gp41) are present at relatively higher levels on

the surface of REJO, CMU06 and SF162 viruses, suggesting that elements that regulate

Env epitope abundance are likely to be conserved among isolates. In contrast, the V2q and

V3 epitopes are inaccessible on the virus and remained so even when longer virion-mAb

interaction times were allowed. The V2q and V3 epitopes become available for binding by

mAbs only when conformational changes were induced. This is expected for V3 epitopes,

as V3 loops are suggested to be tucked underneath the V1V2 loops (105) and

conformational changes such as movement or displacement of V1V2 can release the V3

loop allowing for its recognition by V3 targeting mAbs (106). Computational studies have

also shown that the V3 loop can flicker in and out of its tucked position without disrupting

the V1V2 organization at the apex (105, 107). However, these observations came from the

simulation studies using the X-ray structure of the JR-FL SOSIP.559 trimer (PDB

accession no. 5FYK), not virion associated Env, and therefore may or may not be

applicable to all Env trimers. Even slight opening of the trimer apex can expose the highly

immunogenic V3 loop (24). In such a case, we would expect to observe some level of

binding to V3 mAbs. However, the V3 mAbs (2219, 2557) tested here did not show any

binding even when the REJO virus and mAb interaction time were extended up to 75 min.

The V3 epitopes also remained occluded in SF162 which is highly sensitive to

neutralization by these V3 mAbs (71). The SF162 Env has historically been presumed to

be in an open conformation (108-110), but based on the sensitivity of SF162 to

neutralization by a family of mAbs that recognizes an SF162 type-specific quaternary

epitope that bridges the V2 and V3 domains (111-114), it has been suggested that this Env

actually exists predominantly in the closed conformation (115). This may explain why we

observed masked V3 epitopes on SF162 in our assay. These data support the idea that V3

exposure varies among different HIV-1 isolates and was not applicable to the viruses tested

here. Native-like Env trimers are the leading platform for HIV-1 vaccine design and several

modifications have been made to reduce the Env metastability and exposure of the V3-loop

(116-118). Understanding how the viruses retain the V3 in a tucked-in position may help

these efforts.

The trimer apex epitopes on virions were inaccessible to the V2q mAbs that target

these epitopes. Neutralization of HIV-1, or any other viral pathogen, minimally requires an

initial encounter with an Ab. The PG9, PG16, and PGT145 mAbs can efficiently neutralize

REJO virus (74). Thus, limited accessibility of these epitopes on virions was unexpected,

particularly since V3 is shielded only when the native Env trimer is in the closed pre-fusion

conformation: a conformation that is required for the binding of V2q mAbs (105, 119). It

is known that vulnerable epitopes on HIV-1 particles are masked either conformationally

or via the glycan shield (45, 120, 121). It is also known that due to this conformational

masking, accessibility of epitopes to antibodies can be limited (122). The V2q mAbs are

some of the most broad and potent bNAbs, making the V2q epitope region a major site of

vulnerability (16). Thus, the finding that these functional and vulnerable bNAb Env

epitopes are occluded on virions is not surprising. Considering that induction of bNAbs,

including V2q type Abs, by immunization is a major goal of vaccine design, the data

presented here helps us to better understand the strategies that can be employed to improve

the presentation of these epitopes.

Our previously published study has shown that different mechanisms are involved in

occluding the V2i and V3 epitopes (47). While V3 epitopes become accessible after

engagement of CD4 receptor, this is not the case for V2i epitopes. Again, V2i epitopes

such as those targeted by the 830A mAb are abundantly available on the virions, but both

REJO and CMU06 viruses are resistant to neutralization by 830A when tested under

standard 1 hour incubation assay conditions (47, 56, 74). Both viruses become

neutralization sensitive if the virus-mAb incubation is allowed to continue to 24 hours prior

to adding the target TZM.bl cells (56, 74). Thus, prolonged incubation time allows for

conformational unmasking or allows Env breathing that may be transiently exposing the

V2i and V3 epitopes. Allosteric changes induced by binding of V2i mAbs, but not

prolonged interaction time with virus, allowed PG9 to bind to V2q epitopes on REJO virus.

As both V2i and V2q mAbs bind V1V2 when V2C is in a β-strand conformation (46, 123),

the mechanism by which the binding of V2i helped expose V2q remains unknown.

Likewise, we observed no significant differences in neutralization of REJO virus by the

V2q and CD4bs mAbs, when the neutralization assay omitted the commonly adopted one-

hour pre-incubation step (85, 124-130). These data suggest that HIV-1 interaction with

target cells causes changes that allow these bNAbs to bind to their target epitopes and

neutralize virions. Glycan-glycan interactions are emerging as new class of high-affinity

biomolecular interaction (131), and are shown to support the binding of SARS-CoV-2 to

the host cell membrane (131-133). Similarly, it has been shown that glycan-glycan

interaction between the HIV-1 Env N-linked glycans and the N-acetylglucosamine

(GlcNAc) on the host cell glycocalyx can support virus adhesion to cells (96). Here we

show that binding of PG9 to Env on REJO virions is detected in presence of 293T cells

while negligible binding is observed when cells are not present, supporting that initial

interaction with host cells may influence the Env organization exposing these epitopes. In

addition, lectin-glycan interactions are also shown to impact virus transmission, virus

attachment (134-136), and virus neutralization (99). Both ConA and GRFT lectins tested

here can interact with Env to improve the exposure of the occluded epitopes. However, this

effect is also dependent on the specificity of the lectin. Nonetheless, the evidence presented

regarding how the occluded bNAb epitopes on virions may be unmasked, allowing the

mAbs to bind to their epitopes and block infection, highlights an unexpected role for Env-

cell interaction and a new perspective on the conformational dynamics of Env.

In summary, we show that only select Env epitopes are exposed on the HIV-1 surface,

with V2i epitopes being the most abundant and potentially providing an explanation as to

why Abs against these immunogenic epitopes are more readily elicited during infection

(65, 66, 69, 70, 137-141). Pre-existing V2i antibodies can form immune complexes with

HIV-1 Env and skew antibody elicitation to a more cross-reactive immune response (93,

94), suggesting a possible role of these Abs in influencing the quality of humoral immune

response. We also show a previously unknown fact that interaction with host cells is

required for bNAbs to be able to access their epitopes on virus particles. It will also be

interesting to know if other viruses, for example SARS-CoV-2 and Influenza, have similar

requirements concerning the interaction with host cells in order for the virus-specific

neutralizing Abs to exert their effect. A greater understanding of epitope exposure on

individual virions upon interaction with host cells would contribute greatly to

understanding how humoral responses recognize different Env domains and how it may

influence virus infection and antibody responses.

Acknowledgments:

The authors would like to thank Dr. Susan Zolla-Pazner and Ms. Xiaomei Liu for providing

HIV-1-specific mAbs.

Figure Legends

Figure. 1. Virus-binding assay for detection of antibody binding to Env expressed on

virions. (A) Schematic of the assay; (B) Optimization of virus amount to be coupled to the

microspheres; (C) Representative dot plots and histogram showing the gating strategy for

analysis of antibodies binding to virus particle by flow cytometry. The mAbs were tested

at following concentrations: NIH45-46 at 5 µg/ml; 2219 and 3685 at 50 µg/ml. 1°, primary;

2° secondary.

Figure 2. Binding of mAbs to REJO Env on virions. (A) Microspheres coupled to REJO

virus were reacted with serially diluted mAbs targeting different Env epitopes. (B) Area

under the curve (AUC) were calculated from the curves in A and plotted. (C) Geometric

mean fluorescent intensity (MFI) of each mAb tested at a single dilution are plotted. Virus

coupled microspheres stained with biotin and PE alone were used to set the background

MFI and subtracted.

Figure 3. Binding of mAbs to SF162 Env on virions. (A) Microspheres coupled to SF162

virus were reacted with serially diluted mAbs targeting different Env epitopes. Virus

coupled microspheres stained with biotin and PE alone were used to set the background

MFI and subtracted. (B) Area under the curve (AUC) were calculated from the curves in

A and plotted. (C) Geometric mean fluorescent intensity (MFI) of each mAb tested at a

single dilution are plotted.

Figure 4. Binding of mAbs to CMU06 Env on virions. (A) Microspheres coupled to

CMU06 virus were reacted with mAbs targeting different Env epitopes. Geometric mean

fluorescent intensity (MFI) of each mAb tested at a single dilution are plotted. Virus

coupled microspheres stained with biotin and PE alone were used to set the background

MFI and subtracted.

Figure 5. Binding of mAbs to trimeric Env expressed on cell surface. 293T cells were

transiently transfected with plasmid encoding for full length gp160 of (A) REJO, (B)

CMU06 and (C) SF162. Cells were analyzed for binding of different mAbs 24-hours post-

transfection. Geometric mean fluorescence intensity (MFI) and SD from duplicates or

triplicates in one experiment are shown. Transfected cells stained with biotin and PE alone

were used to set the background MFI and subtracted. The mAbs were tested at following

concentrations: 697, 830A, 2158, 3685 at 100 μg/ml; 2219, 2557 at 50 μg/ml; PG9, PG16,

PGT145, PGDM1400, CH59, VRC01, 3BNC117, b12 at 25 μg/ml; 4E10, 2F5 at 20 μg/ml;

PGT151 at 10 μg/ml; NIH45-46 at 5 μg/ml.

Figure 6. Time-dependent increase of Env binding. Microspheres coupled to REJO

virus particles were treated with mAbs at 37°C for various time from 0 to 75 min. Binding

was detected as in Figure 2 and geometric mean fluorescence intensity (MFI) are shown.

Virus coupled microspheres stained with biotin and PE alone were used to set the

background MFI and subtracted. The mAbs were tested at following concentrations: 830A,

3685 at 100 μg/ml; 2219 at 50 μg/ml; PG9 at 25 μg/ml; NIH45-46 at 2.5 μg/ml. ****, p=

<0.0001; **, p= 0.0026 vs t = 0 min by ANOVA.

Figure 7. Changes in reactivity of PG9 and 2219 mAbs. Microspheres coupled with

REJO virus were incubated with titrated amounts of antibodies followed by biotinylated

V2q mAb PG9 (25 μg/ml) or biotinylated V3 mAb 2219 (50 μg/ml). Binding was detected

with streptavidin-PE and geometric mean fluorescent intensity (MFI) are plotted. Virus

coupled microspheres stained with biotin and PE alone were used to set the background

MFI and subtracted. **** p< 0.0001 by ANOVA.

Figure 8. Neutralization of REJO virus. Virus neutralization was measured after REJO

virus was incubated with serially diluted mAbs for the designated period of time at 37°C

prior to the addition of TZM-bl target cells. Virus infectivity was assessed 48 h later based

on β-galactosidase activity. (A) Neutralization curves are shown. Means and standard

errors calculated from two different experiments (each in duplicate) are shown. Statistical

analyses were performed on the neutralization curves reaching ≥50%. Comparison is made

between neutralization curves at T = 0 vs T = 60 minutes. P= ns, not significant by

nonparametric Mann-Whitney t-test. (B) Area under the neutralization curves were

calculated and plotted as bar graph. Statistical analysis was performed on AUC graph by

ANOVA (P = 0.99; ns, not significant).

Figure 9. Neutralization of CMU06 virus. Virus neutralization was measured after

CMU06 virus was incubated with serially diluted mAbs for the designated period of time

at 37°C prior to the addition of TZM-bl target cells. Virus infectivity was assessed 48 h

later based on β-galactosidase activity. (A) Neutralization curves are shown. Means and

standard errors calculated from two different experiments (each in duplicate) are shown.

Statistical analyses were performed on the neutralization curves reaching ≥50%.

Comparison is made between neutralization curves at T = 0 vs T = 60 minutes. P= ns, not

significant by nonparametric Mann-Whitney t-test. (B) Area under the neutralization

curves were calculated and plotted as bar graph. Statistical analysis was performed on AUC

graph by ANOVA (P = 0.99; ns, not significant).

Figure 10. Influence of virus-cell interaction on Env epitope exposure. Binding of

mAbs to virus-coupled microspheres were measured in the presence of 293T cells by flow

cytometry. Cells were transfected or not transfected with human CD4 plasmid for 24 h and

treated with virus-coupled microspheres and V2i, V3, V2q, CD4bs or control mAbs at 4°C.

MAb binding was detected with biotin-conjugated anti-human IgG and PE-Streptavidin.

Percent PE positive microspheres from one experiment are shown. Gating strategy used

for data analysis is shown in Fig S10. MAb titration curves and bar graphs showing

geometric mean fluorescent intensity (MFI) of each mAb tested at a highest dilution are

plotted. Gray dotted line represents MFI of negative control mAb 3685.

Figure 11. Effect of lectin Concanavalin A on exposure of Env epitopes. Microspheres

coupled with REJO virus were incubated with five-fold titrated amounts of lectin

Concanavalin A (ConA) (50 - 0.08 μg/ml) followed by mAbs at single dilutions (830A,

2219, PG9, 3685 at 50 μg/ml; VRC01 at 25 μg/ml and NIH45-46 at 2.5 μg/ml). (50 μg/ml).

Binding was detected with streptavidin-PE. Geometric mean fluorescent intensity (MFI)

and fold change in binding relative to no ConA are shown. Virus coupled microspheres

stained with biotin and PE alone were used to set the background MFI and subtracted. ****

p= 0.0002, * p= 0.039 by ANOVA relative to negative control mAb 3685.

Figure 12. Effect of Lectins on Env epitope exposure in ELISA. Sucrose pelleted REJO

virus was captured by (A) five-fold titrated lectin ConA (50 - 0.08 μg/ml) and reacted with

different mAbs at one dilution (830A, 2219, PG9, 3685 at 50 μg/ml; VRC01 at 25 μg/ml

and NIH45-46 at 2.5 μg/ml). (B) Sucrose pelleted REJO virus was captured by lectin

Griffithsin (50 μg/ml) and reacted with five-fold titrated mAbs (830A, 2219, PG9, 3685 at

50 μg/ml; VRC01 at 25 μg/ml and NIH45-46 at 2.5 μg/ml). AUC values were calculated

from binding curves. AUC changes relative to the respective WT are shown. Statistical

analysis was performed on binding curves by 2-way ANOVA (**** $p < 0.0001$) relative

to negative control mAb 3685.

Supplementary Figure 1. Measurement of Env in virus preparations by Western blot. (A)

Viruses produced in 293T cells, were concentrated (20X) by sucrose pelleting. Four µl of

each 20X concentrated virus particle prep were lysed and analyzed by SDS-PAGE (4–

20%) and Western blot. An anti-gp120 MAb cocktail (V3: 391/95-D, 694/98-D, 2219,

2558; C2: 847-D, 1006-30D; C5: 450-D, 670-D) was used to quantitate the levels of Env

associated with virions. REJO gp120 protein loaded at different concentrations was used

as a standard. The band density of the REJO gp120 protein was used to generate the linear

curve (B) and to calculate the amount of Env in each virus preparation.

Supplementary Figure 2. Histogram plots related to Fig 2 showing reactivity of serially

diluted mAbs to REJO virus particles coupled to fluorescent microspheres. Virus coupled

microspheres stained with biotin and PE alone were used to set the background staining

(gray). Experiment was repeated to probe for binding of CD4-IgG, 17b and negative

control mAb 3685 (Right panel)

Supplementary Figure 3. Histogram plots related to Fig 3 showing reactivity of serially

diluted mAbs to SF162 virus particles coupled to fluorescent microspheres. Virus coupled

microspheres stained with biotin and PE alone were used to set the background staining

(gray).

Supplementary Figure 4. Representative histogram plots related to Fig 4 showing

reactivity of mAbs to CMU06 virus particles coupled to fluorescent microspheres. Binding

was detected with streptavidin-PE and fluorescent intensity (MFI) are shown. Virus

coupled microspheres stained with biotin and PE alone were used to set the background

staining (gray).

Supplementary Figure 5. Histogram plots related to Fig 5A showing reactivity of mAbs

to REJO Env expressed on cell surface. Replicates from one representative experiment are

shown. Binding was detected with streptavidin-PE. Transfected cells stained with biotin

and PE alone were used to set the background MFI (gray). MAb 3685 was used as negative

control.

Supplementary Figure 6. Histogram plots related to Fig 5B showing reactivity of mAbs

to CMU06 Env expressed on cell surface. Triplicates are shown. Binding was detected with

streptavidin-PE. Transfected cells stained with biotin and PE alone were used to set the

background MFI (gray). MAb 3685 was used as negative control.

Supplementary Figure 7. Histogram plots related to Fig 5C showing reactivity of mAbs

to SF162 Env expressed on cell surface. Triplicates for all antibodies and duplicates for

CD4-IgG and PGT151 are shown. Binding was detected with streptavidin-PE. Transfected

cells stained with biotin and PE alone were used to set the background MFI (gray). MAb

3685 was used as negative control.

Supplementary Figure 8. Representative histogram plots related to Fig 6 showing time-

dependent changes in reactivity of mAbs. Microspheres coupled to REJO virus particles

were treated with mAbs at 37°C for various time from 0 to 75 min. Binding was detected

with streptavidin-PE.

Supplementary Figure 9. Representative histogram plots related to Fig 7 showing

changes in reactivity of PG9 and 2219 mAbs by different antibodies. Microspheres coupled

with REJO virus were incubated with five-fold titrated antibodies followed by biotinylated

V2q mAb PG9 (25 μg/ml). Binding was detected with streptavidin-PE.

Supplementary Figure 10. Gating strategy (A) **and histograms related to Fig 10.** Plots showing binding of mAbs to virions in presence of (B) 293T cells transfected to express hCD4 and (C) un-transfected 293T cells. (D) Histograms showing cell surface expression of human CD4 293T cells. Cells expressing hCD4 are shown in blue and un-transfected cells are shown in gray.

Supplementary Figure 11. Representative histogram plots related to Fig 11 showing changes in reactivity of mAbs by Concanavalin A (ConA).

Funding: "This research was funded by NIH R01 grant (AI140909 to C.U.).

Institutional Review Board Statement: "Not applicable".

Informed Consent Statement: "Not applicable".

Data Availability Statement: All available data are presented in the article.

Conflicts of Interest: "The authors declare no conflict of interest."

References

1.   Wyatt R, Sodroski J. 1998. The HIV-1 envelope glycoproteins: fusogens, antigens,
and immunogens. Science 280:1884-8.

2.   Brandenberg OF, Magnus C, Rusert P, Regoes RR, Trkola A. 2015. Different
infectivity of HIV-1 strains is linked to number of envelope trimers required for
entry. PLoS Pathog 11:e1004595.

3.   Bosch V, Pawlita M. 1990. Mutational analysis of the human immunodeficiency
virus type 1 env gene product proteolytic cleavage site. Journal of Virology
64:2337-2344.

4.   Fennie C, Lasky LA. 1989. Model for intracellular folding of the human
immunodeficiency virus type 1 gp120. Journal of Virology 63:639-646.

5.   Li Y, Luo L, Thomas DY, Kang CY. 2000. The HIV-1 Env protein signal sequence
retards its cleavage and down-regulates the glycoprotein folding. Virology
272:417-28.

6.   Willey RL, Bonifacino JS, Potts BJ, Martin MA, Klausner RD. 1988. Biosynthesis,
cleavage, and degradation of the human immunodeficiency virus 1 envelope
glycoprotein gp160. Proc Natl Acad Sci U S A 85:9580-4.

7.   Martoglio B, Dobberstein B. 1998. Signal sequences: more than just greasy
peptides. Trends Cell Biol 8:410-5.

8.   Land A, Zonneveld D, Braakman I. 2003. Folding of HIV-1 envelope glycoprotein
involves extensive isomerization of disulfide bonds and conformation-dependent
leader peptide cleavage. FASEB J 17:1058-67.

9.   Li Y, Bergeron JJ, Luo L, Ou WJ, Thomas DY, Kang CY. 1996. Effects of
inefficient cleavage of the signal sequence of HIV-1 gp 120 on its association with
calnexin, folding, and intracellular transport. Proc Natl Acad Sci U S A 93:9606-
11.

10.   Li Y, Luo L, Rasool N, Kang CY. 1993. Glycosylation is necessary for the correct
804   folding of human immunodeficiency virus gp120 in CD4 binding. J Virol 67:584-
805   8.

806   11.   Decroly E, Vandenbranden M, Ruysschaert JM, Cogniaux J, Jacob GS, Howard
807   SC, Marshall G, Kompelli A, Basak A, Jean F, Lazure C, Benjannet S, Chrétien M,
808   Day R, Seidah NG. 1994. The convertases furin and PC1 can both cleave the human
809   immunodeficiency virus (HIV)-1 envelope glycoprotein gp160 into gp120 (HIV-I
810   SU) and gp41 (HIV-I TM). Journal of Biological Chemistry 269:12240-12247.

811   12.   Kantanen ML, Leinikki P, Kuismanen E. 1995. Endoproteolytic cleavage of HIV-
812   1 gp160 envelope precursor occurs after exit from the trans-Golgi network (TGN).
813   Archives of Virology 140:1441-1449.

814   13.   Pfeiffer T, Zentgraf H, Freyaldenhoven B, Bosch V. 1997. Transfer of endoplasmic
815   reticulum and Golgi retention signals to human immunodeficiency virus type 1 gp
816   160 inhibits intracellular transport and proteolytic processing of viral glycoprotein
817   but does not influence the cellular site of virus particle budding. Journal of General
818   Virology 78:1745-1753.

819   14.   Doria-Rose NA, Schramm CA, Gorman J, Moore PL, Bhiman JN, DeKosky BJ,
820   Ernandes MJ, Georgiev IS, Kim HJ, Pancera M, Staupe RP, Altae-Tran HR, Bailer
821   RT, Crooks ET, Cupo A, Druz A, Garrett NJ, Hoi KH, Kong R, Louder MK, Longo

NS, McKee K, Nonyane M, O'Dell S, Roark RS, Rudicell RS, Schmidt SD,
Sheward DJ, Soto C, Wibmer CK, Yang Y, Zhang Z, Program NCS, Mullikin JC,
Binley JM, Sanders RW, Wilson IA, Moore JP, Ward AB, Georgiou G, Williamson
C, Abdool Karim SS, Morris L, Kwong PD, Shapiro L, Mascola JR. 2014.
Developmental pathway for potent V1V2-directed HIV-neutralizing antibodies.
Nature 509:55-62.

15.   McLellan JS, Pancera M, Carrico C, Gorman J, Julien JP, Khayat R, Louder R,
Pejchal R, Sastry M, Dai K, O'Dell S, Patel N, Shahzad-ul-Hussan S, Yang Y,
Zhang B, Zhou T, Zhu J, Boyington JC, Chuang GY, Diwanji D, Georgiev I, Kwon
YD, Lee D, Louder MK, Moquin S, Schmidt SD, Yang ZY, Bonsignori M, Crump
JA, Kapiga SH, Sam NE, Haynes BF, Burton DR, Koff WC, Walker LM, Phogat
S, Wyatt R, Orwenyo J, Wang LX, Arthos J, Bewley CA, Mascola JR, Nabel GJ,
Schief WR, Ward AB, Wilson IA, Kwong PD. 2011. Structure of HIV-1 gp120
V1/V2 domain with broadly neutralizing antibody PG9. Nature 480:336-43.

16.   Walker LM, Phogat SK, Chan-Hui PY, Wagner D, Phung P, Goss JL, Wrin T,
Simek MD, Fling S, Mitcham JL, Lehrman JK, Priddy FH, Olsen OA, Frey SM,
Hammond PW, Kaminsky S, Zamb T, Moyle M, Koff WC, Poignard P, Burton
DR. 2009. Broad and potent neutralizing antibodies from an African donor reveal
a new HIV-1 vaccine target. Science 326:285-9.

17.   Walker LM, Huber M, Doores KJ, Falkowska E, Pejchal R, Julien JP, Wang SK,
Ramos A, Chan-Hui PY, Moyle M, Mitcham JL, Hammond PW, Olsen OA, Phung
P, Fling S, Wong CH, Phogat S, Wrin T, Simek MD, Koff WC, Wilson IA, Burton
DR, Poignard P. 2011. Broad neutralization coverage of HIV by multiple highly
potent antibodies. Nature 477:466-70.

18.   Sok D, van Gils MJ, Pauthner M, Julien J-P, Saye-Francisco KL, Hsueh J, Briney
B, Lee JH, Le KM, Lee PS. 2014. Recombinant HIV envelope trimer selects for
quaternary-dependent antibodies targeting the trimer apex. Proceedings of the
National Academy of Sciences 111:17624-17629.

19.   Gallo SA, Finnegan CM, Viard M, Raviv Y, Dimitrov A, Rawat SS, Puri A, Durell
S, Blumenthal R. 2003. The HIV Env-mediated fusion reaction. Biochim Biophys
Acta 1614:36-50.

20.   Eckert DM, Kim PS. 2001. Mechanisms of viral membrane fusion and its
inhibition. Annu Rev Biochem 70:777-810.

21.   Stieh DJ, King DF, Klein K, Aldon Y, McKay PF, Shattock RJ. 2015. Discrete
partitioning of HIV-1 Env forms revealed by viral capture. Retrovirology 12:81.

22.   Crooks ET, Tong T, Osawa K, Binley JM. 2011. Enzyme digests eliminate
nonfunctional Env from HIV-1 particle surfaces, leaving native Env trimers intact
and viral infectivity unaffected. J Virol 85:5825-39.

23.   Moore PL, Crooks ET, Porter L, Zhu P, Cayanan CS, Grise H, Corcoran P, Zwick
MB, Franti M, Morris L, Roux KH, Burton DR, Binley JM. 2006. Nature of
nonfunctional envelope proteins on the surface of human immunodeficiency virus
type 1. J Virol 80:2515-28.

24.   Sanders RW, Derking R, Cupo A, Julien JP, Yasmeen A, de Val N, Kim HJ,
Blattner C, de la Pena AT, Korzun J, Golabek M, de Los Reyes K, Ketas TJ, van
Gils MJ, King CR, Wilson IA, Ward AB, Klasse PJ, Moore JP. 2013. A next-
generation cleaved, soluble HIV-1 Env trimer, BG505 SOSIP.664 gp140, expresses

multiple epitopes for broadly neutralizing but not non-neutralizing antibodies.
PLoS Pathog 9:e1003618.
25.   Pegu A, Hessell AJ, Mascola JR, Haigwood NL. 2017. Use of broadly neutralizing
antibodies for HIV-1 prevention. Immunol Rev 275:296-312.
26.   Phelps M, Balazs AB. 2021. Contribution to HIV Prevention and Treatment by
Antibody-Mediated Effector Function and Advances in Broadly Neutralizing
Antibody Delivery by Vectored Immunoprophylaxis. Front Immunol 12:734304.
27.   Stephenson KE, Barouch DH. 2016. Broadly Neutralizing Antibodies for HIV
Eradication. Curr HIV/AIDS Rep 13:31-7.
28.   Sanders RW, Moore JP. 2017. Native-like Env trimers as a platform for HIV-1
vaccine design. Immunol Rev 275:161-182.
29.   Wei X, Decker JM, Wang S, Hui H, Kappes JC, Wu X, Salazar-Gonzalez JF,
Salazar MG, Kilby JM, Saag MS, Komarova NL, Nowak MA, Hahn BH, Kwong
PD, Shaw GM. 2003. Antibody neutralization and escape by HIV-1. Nature
422:307-12.
30.   Burton DR, Hangartner L. 2016. Broadly Neutralizing Antibodies to HIV and Their
Role in Vaccine Design. Annu Rev Immunol 34:635-59.
31.   Kwong PD, Mascola JR. 2012. Human antibodies that neutralize HIV-1:
identification, structures, and B cell ontogenies. Immunity 37:412-25.
32.   West AP, Jr., Scharf L, Scheid JF, Klein F, Bjorkman PJ, Nussenzweig MC. 2014.
Structural insights on the role of antibodies in HIV-1 vaccine and therapy. Cell
156:633-48.
33.   Zolla-Pazner S, Edlefsen PT, Rolland M, Kong XP, deCamp A, Gottardo R,
Williams C, Tovanabutra S, Sharpe-Cohen S, Mullins JI, deSouza MS, Karasavvas
N, Nitayaphan S, Rerks-Ngarm S, Pitisuttihum P, Kaewkungwal J, O'Connell RJ,
Robb ML, Michael NL, Kim JH, Gilbert P. 2014. Vaccine-induced Human
Antibodies Specific for the Third Variable Region of HIV-1 gp120 Impose Immune
Pressure on Infecting Viruses. EBioMedicine 1:37-45.
34.   Zolla-Pazner S, Cohen S, Pinter A, Krachmarov C, Wrin T, Wang S, Lu S. 2009.
Cross-clade neutralizing antibodies against HIV-1 induced in rabbits by focusing
the immune response on a neutralizing epitope. Virology 392:82-93.
35.   Visciano ML, Tuen M, Gorny MK, Hioe CE. 2008. In vivo alteration of humoral
responses to HIV-1 envelope glycoprotein gp120 by antibodies to the CD4-binding
site of gp120. Virology 372:409-420.
36.   Lynch RM, Rong R, Boliar S, Sethi A, Li B, Mulenga J, Allen S, Robinson JE,
Gnanakaran S, Derdeyn CA. 2011. The B cell response is redundant and highly
focused on V1V2 during early subtype C infection in a Zambian seroconverter.
Journal of Virology 85:905-915.
37.   Kumar R, Tuen M, Liu J, Nadas A, Pan R, Kong X, Hioe CE. 2013. Elicitation of
broadly reactive antibodies against glycan-modulated neutralizing V3 epitopes of
HIV-1 by immune complex vaccines. Vaccine 31:5413-21.
38.   Hessell AJ, McBurney S, Pandey S, Sutton W, Liu L, Li L, Totrov M, Zolla-Pazner
S, Haigwood NL, Gorny MK. 2016. Induction of neutralizing antibodies in rhesus
macaques using V3 mimotope peptides. Vaccine 34:2713-2721.
39.   Hessell AJ, Malherbe DC, Pissani F, McBurney S, Krebs SJ, Gomes M, Pandey S,
Sutton WF, Burwitz BJ, Gray M, Robins H, Park BS, Sacha JB, La Branche CC,

Fuller DH, Montefiori DC, Stamatatos L, Sather DN, Haigwood NL. 2016.
Achieving potent autologous neutralizing antibody responses against tier 2 HIV-1
viruses by strategic selection of envelope immunogens. Journal of Immunology
196:3064-3078.
40. Davis KL, Gray ES, Moore PL, Decker JM, Salomon A, Montefiori DC, Graham
BS, Keefer MC, Pinter A, Morris L, Hahn BH, Shaw GM. 2009. High titer HIV-1
V3-specific antibodies with broad reactivity but low neutralizing potency in acute
infection and following vaccination. Virology 387:414-426.
41. Davis KL, Bibollet-Ruche F, Li H, Decker JM, Kutsch O, Morris L, Salomon A,
Pinter A, Hoxie JA, Hahn BH, Kwong PD, Shaw GM. 2009. Human
immunodeficiency virus type 2 (HIV-2)/HIV-1 envelope chimeras detect high titers
of broadly reactive HIV-1 V3-specific antibodies in human plasma. Journal of
Virology 83:1240-1259.
42. Carbonetti S, Oliver BG, Glenn J, Stamatatos L, Sather DN. 2014. Soluble HIV-1
envelope immunogens derived from an elite neutralizer elicit cross-reactive V1V2
antibodies and low potency neutralizing antibodies. PLoS ONE 9.
43. Tomaras GD, Yates NL, Liu P, Qin L, Fouda GG, Chavez LL, Decamp AC, Parks
RJ, Ashley VC, Lucas JT, Cohen M, Eron J, Hicks CB, Liao HX, Self SG, Landucci
G, Forthal DN, Weinhold KJ, Keele BF, Hahn BH, Greenberg ML, Morris L,
Karim SS, Blattner WA, Montefiori DC, Shaw GM, Perelson AS, Haynes BF.
2008. Initial B-cell responses to transmitted human immunodeficiency virus type
1: virion-binding immunoglobulin M (IgM) and IgG antibodies followed by plasma
anti-gp41 antibodies with ineffective control of initial viremia. J Virol 82:12449-
63.
44. Klein F, Mouquet H, Dosenovic P, Scheid JF, Scharf L, Nussenzweig MC. 2013.
Antibodies in HIV-1 vaccine development and therapy. Science 341:1199-204.
45. Parker Miller E, Finkelstein MT, Erdman MC, Seth PC, Fera D. 2021. A Structural
Update of Neutralizing Epitopes on the HIV Envelope, a Moving Target. Viruses
13.
46. Pan R, Gorny MK, Zolla-Pazner S, Kong XP. 2015. The V1V2 Region of HIV-1
gp120 Forms a Five-Stranded Beta Barrel. J Virol 89:8003-10.
47. Upadhyay C, Mayr LM, Zhang J, Kumar R, Gorny MK, Nadas A, Zolla-Pazner S,
Hioe CE. 2014. Distinct mechanisms regulate exposure of neutralizing epitopes in
the V2 and V3 loops of HIV-1 envelope. J Virol 88:12853-65.
48. Aiyegbo MS, Shmelkov E, Dominguez L, Goger M, Battacharya S, deCamp AC,
Gilbert PB, Berman PW, Cardozo T. 2017. Peptide Targeted by Human Antibodies
Associated with HIV Vaccine-Associated Protection Assumes a Dynamic alpha-
Helical Structure. PLoS One 12:e0170530.
49. Liao HX, Bonsignori M, Alam SM, McLellan JS, Tomaras GD, Moody MA,
Kozink DM, Hwang KK, Chen X, Tsao CY, Liu P, Lu X, Parks RJ, Montefiori DC,
Ferrari G, Pollara J, Rao M, Peachman KK, Santra S, Letvin NL, Karasavvas N,
Yang ZY, Dai K, Pancera M, Gorman J, Wiehe K, Nicely NI, Rerks-Ngarm S,
Nitayaphan S, Kaewkungwal J, Pitisuttithum P, Tartaglia J, Sinangil F, Kim JH,
Michael NL, Kepler TB, Kwong PD, Mascola JR, Nabel GJ, Pinter A, Zolla-Pazner
S, Haynes BF. 2013. Vaccine induction of antibodies against a structurally

heterogeneous site of immune pressure within HIV-1 envelope protein variable
regions 1 and 2. Immunity 38:176-86.

50. Wibmer CK, Richardson SI, Yolitz J, Cicala C, Arthos J, Moore PL, Morris L.
2018. Common helical V1V2 conformations of HIV-1 Envelope expose the
alpha4beta7 binding site on intact virions. Nat Commun 9:4489.

51. Pancera M, Zhou T, Druz A, Georgiev IS, Soto C, Gorman J, Huang J, Acharya P,
Chuang GY, Ofek G, Stewart-Jones GB, Stuckey J, Bailer RT, Joyce MG, Louder
MK, Tumba N, Yang Y, Zhang B, Cohen MS, Haynes BF, Mascola JR, Morris L,
Munro JB, Blanchard SC, Mothes W, Connors M, Kwong PD. 2014. Structure and
immune recognition of trimeric pre-fusion HIV-1 Env. Nature 514:455-61.

52. Lee JH, Andrabi R, Su CY, Yasmeen A, Julien JP, Kong L, Wu NC, McBride R,
Sok D, Pauthner M, Cottrell CA, Nieusma T, Blattner C, Paulson JC, Klasse PJ,
Wilson IA, Burton DR, Ward AB. 2017. A Broadly Neutralizing Antibody Targets
the Dynamic HIV Envelope Trimer Apex via a Long, Rigidified, and Anionic beta-
Hairpin Structure. Immunity 46:690-702.

53. Sok D, van Gils MJ, Pauthner M, Julien JP, Saye-Francisco KL, Hsueh J, Briney
B, Lee JH, Le KM, Lee PS, Hua Y, Seaman MS, Moore JP, Ward AB, Wilson IA,
Sanders RW, Burton DR. 2014. Recombinant HIV envelope trimer selects for
quaternary-dependent antibodies targeting the trimer apex. Proc Natl Acad Sci U S
A 111:17624-9.

54. Platt EJ, Wehrly K, Kuhmann SE, Chesebro B, Kabat D. 1998. Effects of CCR5
and CD4 cell surface concentrations on infections by macrophagetropic isolates of
human immunodeficiency virus type 1. J Virol 72:2855-64.

55. Ochsenbauer C, Edmonds TG, Ding H, Keele BF, Decker J, Salazar MG, Salazar-
Gonzalez JF, Shattock R, Haynes BF, Shaw GM, Hahn BH, Kappes JC. 2012.
Generation of transmitted/founder HIV-1 infectious molecular clones and
characterization of their replication capacity in CD4 T lymphocytes and monocyte-
derived macrophages. J Virol 86:2715-28.

56. Upadhyay C, Feyznezhad R, Cao L, Chan KW, Liu K, Yang W, Zhang H, Yolitz
J, Arthos J, Nadas A, Kong XP, Zolla-Pazner S, Hioe CE. 2020. Signal peptide of
HIV-1 envelope modulates glycosylation impacting exposure of V1V2 and other
epitopes. PLoS Pathog 16:e1009185.

57. Kmiec D, Iyer SS, Sturzel CM, Sauter D, Hahn BH, Kirchhoff F. 2016. Vpu-
Mediated Counteraction of Tetherin Is a Major Determinant of HIV-1 Interferon
Resistance. MBio 7.

58. Walker LM, Phogat SK, Chan-Hui PY, Wagner D, Phung P, Goss JL, Wrin T,
Simek MD, Fling S, Mitcham JL, Lehrman JK, Priddy FH, Olsen OA, Frey SM,
Hammond PW, Protocol GPI, Kaminsky S, Zamb T, Moyle M, Koff WC, Poignard
P, Burton DR. 2009. Broad and potent neutralizing antibodies from an African
donor reveal a new HIV-1 vaccine target. Science 326:285-9.

59. Wu X, Yang ZY, Li Y, Hogerkorp CM, Schief WR, Seaman MS, Zhou T, Schmidt
SD, Wu L, Xu L, Longo NS, McKee K, O'Dell S, Louder MK, Wycuff DL, Feng
Y, Nason M, Doria-Rose N, Connors M, Kwong PD, Roederer M, Wyatt RT, Nabel
GJ, Mascola JR. 2010. Rational design of envelope identifies broadly neutralizing
human monoclonal antibodies to HIV-1. Science 329:856-61.

60.   Barbas CF, 3rd, Bjorling E, Chiodi F, Dunlop N, Cababa D, Jones TM, Zebedee
SL, Persson MA, Nara PL, Norrby E, et al. 1992. Recombinant human Fab
fragments neutralize human type 1 immunodeficiency virus in vitro. Proc Natl
Acad Sci U S A 89:9339-43.
61.   Scheid JF, Mouquet H, Ueberheide B, Diskin R, Klein F, Oliveira TY, Pietzsch J,
Fenyo D, Abadir A, Velinzon K, Hurley A, Myung S, Boulad F, Poignard P, Burton
DR, Pereyra F, Ho DD, Walker BD, Seaman MS, Bjorkman PJ, Chait BT,
Nussenzweig MC. 2011. Sequence and structural convergence of broad and potent
HIV antibodies that mimic CD4 binding. Science 333:1633-7.
62.   Thali M, Moore JP, Furman C, Charles M, Ho DD, Robinson J, Sodroski J. 1993.
Characterization of conserved human immunodeficiency virus type 1 gp120
neutralization epitopes exposed upon gp120-CD4 binding. J Virol 67:3978-88.
63.   Buchacher A, Predl R, Strutzenberger K, Steinfellner W, Trkola A, Purtscher M,
Gruber G, Tauer C, Steindl F, Jungbauer A, et al. 1994. Generation of human
monoclonal antibodies against HIV-1 proteins; electrofusion and Epstein-Barr
virus transformation for peripheral blood lymphocyte immortalization. AIDS Res
Hum Retroviruses 10:359-69.
64.   Stiegler G, Kunert R, Purtscher M, Wolbank S, Voglauer R, Steindl F, Katinger H.
2001. A potent cross-clade neutralizing human monoclonal antibody against a
novel epitope on gp41 of human immunodeficiency virus type 1. AIDS Res Hum
Retroviruses 17:1757-65.
65.   Gorny MK, Moore JP, Conley AJ, Karwowska S, Sodroski J, Williams C, Burda
S, Boots LJ, Zolla-Pazner S. 1994. Human anti-V2 monoclonal antibody that
neutralizes primary but not laboratory isolates of human immunodeficiency virus
type 1. J Virol 68:8312-20.
66.   Gorny MK, Pan R, Williams C, Wang XH, Volsky B, O'Neal T, Spurrier B,
Sampson JM, Li L, Seaman MS, Kong XP, Zolla-Pazner S. 2012. Functional and
immunochemical cross-reactivity of V2-specific monoclonal antibodies from HIV-
1-infected individuals. Virology 427:198-207.
67.   Nyambi PN, Mbah HA, Burda S, Williams C, Gorny MK, Nadas A, Zolla-Pazner
S. 2000. Conserved and exposed epitopes on intact, native, primary human
immunodeficiency virus type 1 virions of group M. J Virol 74:7096-107.
68.   Pinter A, Honnen WJ, He Y, Gorny MK, Zolla-Pazner S, Kayman SC. 2004. The
V1/V2 domain of gp120 is a global regulator of the sensitivity of primary human
immunodeficiency virus type 1 isolates to neutralization by antibodies commonly
induced upon infection. J Virol 78:5205-15.
69.   Gorny MK, Conley AJ, Karwowska S, Buchbinder A, Xu JY, Emini EA, Koenig
S, Zolla-Pazner S. 1992. Neutralization of diverse human immunodeficiency virus
type 1 variants by an anti-V3 human monoclonal antibody. J Virol 66:7538-42.
70.   Gorny MK, Williams C, Volsky B, Revesz K, Wang XH, Burda S, Kimura T,
Konings FA, Nadas A, Anyangwe CA, Nyambi P, Krachmarov C, Pinter A, Zolla-
Pazner S. 2006. Cross-clade neutralizing activity of human anti-V3 monoclonal
antibodies derived from the cells of individuals infected with non-B clades of
human immunodeficiency virus type 1. J Virol 80:6865-72.

71.   Hioe CE, Wrin T, Seaman MS, Yu X, Wood B, Self S, Williams C, Gorny MK,
1049   Zolla-Pazner S. 2010. Anti-V3 monoclonal antibodies display broad neutralizing
1050   activities against multiple HIV-1 subtypes. PLoS One 5:e10254.
1051   72.   Gigler A, Dorsch S, Hemauer A, Williams C, Kim S, Young NS, Zolla-Pazner S,
1052   Wolf H, Gorny MK, Modrow S. 1999. Generation of neutralizing human
1053   monoclonal antibodies against parvovirus B19 proteins. J Virol 73:1974-9.
1054   73.   Bradley KA, Mogridge J, Mourez M, Collier RJ, Young JA. 2001. Identification of
1055   the cellular receptor for anthrax toxin. Nature 414:225-9.
1056   74.   Upadhyay C, Feyznezhad R, Yang W, Zhang H, Zolla-Pazner S, Hioe CE. 2018.
1057   Alterations of HIV-1 envelope phenotype and antibody-mediated neutralization by
1058   signal peptide mutations. PLoS Pathog 14:e1006812.
1059   75.   Rossio JL, Esser MT, Suryanarayana K, Schneider DK, Bess JW, Jr., Vasquez GM,
1060   Wiltrout TA, Chertova E, Grimes MK, Sattentau Q, Arthur LO, Henderson LE,
1061   Lifson JD. 1998. Inactivation of human immunodeficiency virus type 1 infectivity
1062   with preservation of conformational and functional integrity of virion surface
1063   proteins. J Virol 72:7992-8001.
1064   76.   Weiss S, Itri V, Pan R, Jiang X, Luo CC, Morris L, Malherbe DC, Barnette P,
1065   Alexander J, Kong XP, Haigwood NL, Hessell AJ, Duerr R, Zolla-Pazner S. 2022.
1066   Differential V2-directed antibody responses in non-human primates infected with
1067   SHIVs or immunized with diverse HIV vaccines. Nat Commun 13:903.
1068   77.   Haynes BF, Gilbert PB, McElrath MJ, Zolla-Pazner S, Tomaras GD, Alam SM,
1069   Evans DT, Montefiori DC, Karnasuta C, Sutthent R, Liao HX, DeVico AL, Lewis
1070   GK, Williams C, Pinter A, Fong Y, Janes H, DeCamp A, Huang Y, Rao M, Billings
1071   E, Karasavvas N, Robb ML, Ngauy V, de Souza MS, Paris R, Ferrari G, Bailer RT,
1072   Soderberg KA, Andrews C, Berman PW, Frahm N, De Rosa SC, Alpert MD, Yates
1073   NL, Shen X, Koup RA, Pitisuttithum P, Kaewkungwal J, Nitayaphan S, Rerks-
1074   Ngarm S, Michael NL, Kim JH. 2012. Immune-correlates analysis of an HIV-1
1075   vaccine efficacy trial. N Engl J Med 366:1275-86.
1076   78.   Mkhize NN, Durgiah R, Ashley V, Archary D, Garrett NJ, Karim QA, Karim SS,
1077   Moore PL, Yates N, Passmore JA, Tomaras GD, Morris L. 2016. Broadly
1078   neutralizing antibody specificities detected in the genital tract of HIV-1 infected
1079   women. AIDS 30:1005-14.
1080   79.   Yates NL, deCamp AC, Korber BT, Liao HX, Irene C, Pinter A, Peacock J, Harris
1081   LJ, Sawant S, Hraber P, Shen X, Rerks-Ngarm S, Pitisuttithum P, Nitayapan S,
1082   Berman PW, Robb ML, Pantaleo G, Zolla-Pazner S, Haynes BF, Alam SM,
1083   Montefiori DC, Tomaras GD. 2018. HIV-1 Envelope Glycoproteins from Diverse
1084   Clades Differentiate Antibody Responses and Durability among Vaccinees. J Virol
1085   92.
1086   80.   Klingler J, Lambert GS, Itri V, Liu S, Bandres JC, Enyindah-Asonye G, Liu X,
1087   Simon V, Gleason CR, Kleiner G, Chiu HP, Hung CT, Kowdle S, Amanat F, Lee
1088   B, Zolla-Pazner S, Upadhyay C, Hioe CE. 2021. Detection of Antibody Responses
1089   Against SARS-CoV-2 in Plasma and Saliva From Vaccinated and Infected
1090   Individuals. Front Immunol 12:759688.
1091   81.   Klingler J, Lambert GS, Bandres JC, Emami-Gorizi R, Nadas A, Oguntuyo KY,
1092   Amanat F, Bermudez-Gonzalez MC, Gleason C, Kleiner G, Simon V, Lee B, Zolla-

Pazner S, Upadhyay C, Hioe CE. 2022. Immune profiles to distinguish hospitalized
versus ambulatory COVID-19 cases in older patients. iScience 25:105608.

82.   Butler AL, Fallon JK, Alter G. 2019. A Sample-Sparing Multiplexed ADCP Assay.
Front Immunol 10:1851.

83.   Altman JB, Liu X, Itri V, Zolla-Pazner S, Powell RLR. 2018. Optimized protocol
for detection of native, full-length HIV-1 envelope on the surface of transfected
cells. Health Sci Rep 1:e74.

84.   Ivan B, Sun Z, Subbaraman H, Friedrich N, Trkola A. 2019. CD4 occupancy
triggers sequential pre-fusion conformational states of the HIV-1 envelope trimer
with relevance for broadly neutralizing antibody activity. PLoS Biol 17:e3000114.

85.   Montefiori DC. 2005. Evaluating neutralizing antibodies against HIV, SIV, and
SHIV in luciferase reporter gene assays. Curr Protoc Immunol Chapter 12:Unit 12
11.

86.   Matyas GR, Beck Z, Karasavvas N, Alving CR. 2009. Lipid binding properties of
4E10, 2F5, and WR304 monoclonal antibodies that neutralize HIV-1. Biochim
Biophys Acta 1788:660-5.

87.   Alam SM, McAdams M, Boren D, Rak M, Scearce RM, Gao F, Camacho ZT,
Gewirth D, Kelsoe G, Chen P, Haynes BF. 2007. The role of antibody
polyspecificity and lipid reactivity in binding of broadly neutralizing anti-HIV-1
envelope human monoclonal antibodies 2F5 and 4E10 to glycoprotein 41
membrane proximal envelope epitopes. J Immunol 178:4424-35.

88.   Falkowska E, Le KM, Ramos A, Doores KJ, Lee JH, Blattner C, Ramirez A,
Derking R, van Gils MJ, Liang CH, McBride R, von Bredow B, Shivatare SS, Wu
CY, Chan-Hui PY, Liu Y, Feizi T, Zwick MB, Koff WC, Seaman MS, Swiderek
K, Moore JP, Evans D, Paulson JC, Wong CH, Ward AB, Wilson IA, Sanders RW,
Poignard P, Burton DR. 2014. Broadly neutralizing HIV antibodies define a glycan-
dependent epitope on the prefusion conformation of gp41 on cleaved envelope
trimers. Immunity 40:657-68.

89.   Diskin R, Scheid JF, Marcovecchio PM, West Jr AP, Klein F, Gao H,
Gnanapragasam PNP, Abadir A, Seaman MS, Nussenzweig MC, Bjorkman PJ.
2011. Increasing the potency and breadth of an HIV antibody by using structure-
based rational design. Science 334:1289-1293.

90.   Scheid JF, Mouquet H, Ueberheide B, Diskin R, Klein F, Oliveira TYK, Pietzsch
J, Fenyo D, Abadir A, Velinzon K, Hurley A, Myung S, Boulad F, Poignard P,
Burton DR, Pereyra F, Ho DD, Walker BD, Seaman MS, Bjorkman PJ, Chait BT,
Nussenzweig MC. 2011. Sequence and Structural Convergence of Broad and
Potent HIV Antibodies That Mimic CD4 Binding. Science 333:1633-1637.

91.   Moulard M, Hallenberger S, Garten W, Klenk HD. 1999. Processing and routage
of HIV glycoproteins by furin to the cell surface. Virus Res 60:55-65.

92.   Blattner C, Lee JH, Sliepen K, Derking R, Falkowska E, de la Pena AT, Cupo A,
Julien JP, van Gils M, Lee PS, Peng W, Paulson JC, Poignard P, Burton DR, Moore
JP, Sanders RW, Wilson IA, Ward AB. 2014. Structural delineation of a quaternary,
cleavage-dependent epitope at the gp41-gp120 interface on intact HIV-1 Env
trimers. Immunity 40:669-80.

93.   Hioe CE, Kumar R, Upadhyay C, Jan M, Fox A, Itri V, Peachman KK, Rao M, Liu
L, Lo NC, Tuen M, Jiang X, Kong XP, Zolla-Pazner S. 2018. Modulation of

1139    Antibody Responses to the V1V2 and V3 Regions of HIV-1 Envelope by Immune
1140    Complex Vaccines. Front Immunol 9:2441.

1141 94. Tang AF, Enyindah-Asonye G, Hioe CE. 2021. Immune Complex Vaccine
1142    Strategies to Combat HIV-1 and Other Infectious Diseases. Vaccines (Basel) 9.

1143 95. Varki A. 2011. Evolutionary forces shaping the Golgi glycosylation machinery:
1144    why cell surface glycans are universal to living cells. Cold Spring Harb Perspect
1145    Biol 3.

1146 96. Spillings BL, Day CJ, Garcia-Minambres A, Aggarwal A, Condon ND, Haselhorst
1147    T, Purcell DFJ, Turville SG, Stow JL, Jennings MP, Mak J. 2022. Host glycocalyx
1148    captures HIV proximal to the cell surface via oligomannose-GlcNAc glycan-glycan
1149    interactions to support viral entry. Cell Rep 38:110296.

1150 97. Buckley J, Battaglia G. 2022. The Role of Glycocalyx Steric Effects on Viral
1151    Endocytosis*. bioRxiv doi:10.1101/2022.02.12.480189:2022.02.12.480189.

1152 98. Hotter D, Kirchhoff F, Sauter D. 2013. HIV-1 Vpu does not degrade interferon
1153    regulatory factor 3. J Virol 87:7160-5.

1154 99. Jan M, Upadhyay C, Sharma A, Hioe CE, Arora SK. 2017. Short Communication:
1155    Manalpha1-2Man-Binding Anti-HIV Lectins Enhance the Exposure of V2i and V3
1156    Crown Neutralization Epitopes on the V1/V2 and V3 Hypervariable Loops of HIV-
1157    1 Envelope. AIDS Res Hum Retroviruses 33:941-945.

1158 100. Cavada BS, Pinto-Junior VR, Osterne VJS, Nascimento KS. 2018. ConA-Like
1159    Lectins: High Similarity Proteins as Models to Study Structure/Biological
1160    Activities Relationships. Int J Mol Sci 20.

1161 101. Jang H, Lee DH, Kang HG, Lee SJ. 2020. Concanavalin A targeting N-linked
1162    glycans in spike proteins influence viral interactions. Dalton Trans 49:13538-
1163    13543.

1164 102. Cummings RD, Etzler ME. 2009. Antibodies and Lectins in Glycan Analysis. *In*
1165    Varki A, Cummings RD, Esko JD, Freeze HH, Stanley P, Bertozzi CR, Hart GW,
1166    Etzler ME (ed), Essentials of Glycobiology, 2nd ed, Cold Spring Harbor (NY).

1167 103. Staropoli I, Dufloo J, Ducher A, Commere PH, Sartori-Rupp A, Novault S, Bruel
1168    T, Lorin V, Mouquet H, Schwartz O, Casartelli N. 2020. Flow Cytometry Analysis
1169    of HIV-1 Env Conformations at the Surface of Infected Cells and Virions: Role of
1170    Nef, CD4, and SERINC5. J Virol 94.

1171 104. Arakelyan A, Fitzgerald W, King DF, Rogers P, Cheeseman HM, Grivel JC,
1172    Shattock RJ, Margolis L. 2017. Flow virometry analysis of envelope glycoprotein
1173    conformations on individual HIV virions. Sci Rep 7:948.

1174 105. Zolla-Pazner S, Cohen SS, Boyd D, Kong XP, Seaman M, Nussenzweig M, Klein
1175    F, Overbaugh J, Totrov M. 2016. Structure/Function Studies Involving the V3
1176    Region of the HIV-1 Envelope Delineate Multiple Factors That Affect
1177    Neutralization Sensitivity. J Virol 90:636-49.

1178 106. Wang H, Barnes CO, Yang Z, Nussenzweig MC, Bjorkman PJ. 2018. Partially
1179    Open HIV-1 Envelope Structures Exhibit Conformational Changes Relevant for
1180    Coreceptor Binding and Fusion. Cell Host Microbe 24:579-592 e4.

1181 107. Powell RLR, Totrov M, Itri V, Liu X, Fox A, Zolla-Pazner S. 2017. Plasticity and
1182    Epitope Exposure of the HIV-1 Envelope Trimer. J Virol doi:10.1128/JVI.00410-
1183    17.

108. Stamatatos L, Wiskerchen M, Cheng-Mayer C. 1998. Effect of major deletions in
the V1 and V2 loops of a macrophage-tropic HIV type 1 isolate on viral envelope
structure, cell entry, and replication. AIDS Res Hum Retroviruses 14:1129-39.
109. Saunders CJ, McCaffrey RA, Zharkikh I, Kraft Z, Malenbaum SE, Burke B, Cheng-
Mayer C, Stamatatos L. 2005. The V1, V2, and V3 regions of the human
immunodeficiency virus type 1 envelope differentially affect the viral phenotype in
an isolate-dependent manner. J Virol 79:9069-80.
110. Brandenberg OF, Rusert P, Magnus C, Weber J, Boni J, Gunthard HF, Regoes RR,
Trkola A. 2014. Partial rescue of V1V2 mutant infectivity by HIV-1 cell-cell
transmission supports the domain's exceptional capacity for sequence variation.
Retrovirology 11:75.
111. Krachmarov C, Lai Z, Honnen WJ, Salomon A, Gorny MK, Zolla-Pazner S,
Robinson J, Pinter A. 2011. Characterization of structural features and diversity of
variable-region determinants of related quaternary epitopes recognized by human
and rhesus macaque monoclonal antibodies possessing unusually potent
neutralizing activities. J Virol 85:10730-40.
112. Robinson JE, Franco K, Elliott DH, Maher MJ, Reyna A, Montefiori DC, Zolla-
Pazner S, Gorny MK, Kraft Z, Stamatatos L. 2010. Quaternary epitope specificities
of anti-HIV-1 neutralizing antibodies generated in rhesus macaques infected by the
simian/human immunodeficiency virus SHIVSF162P4. J Virol 84:3443-53.
113. Honnen WJ, Krachmarov C, Kayman SC, Gorny MK, Zolla-Pazner S, Pinter A.
2007. Type-specific epitopes targeted by monoclonal antibodies with exceptionally
potent neutralizing activities for selected strains of human immunodeficiency virus
type 1 map to a common region of the V2 domain of gp120 and differ only at single
positions from the clade B consensus sequence. J Virol 81:1424-32.
114. Gorny MK, Stamatatos L, Volsky B, Revesz K, Williams C, Wang XH, Cohen S,
Staudinger R, Zolla-Pazner S. 2005. Identification of a new quaternary neutralizing
epitope on human immunodeficiency virus type 1 virus particles. J Virol 79:5232-
7.
115. Qualls ZM, Choudhary A, Honnen W, Prattipati R, Robinson JE, Pinter A. 2018.
Identification of Novel Structural Determinants in MW965 Env That Regulate the
Neutralization Phenotype and Conformational Masking Potential of Primary HIV-
1 Isolates. J Virol 92.
116. Kulp DW, Steichen JM, Pauthner M, Hu X, Schiffner T, Liguori A, Cottrell CA,
Havenar-Daughton C, Ozorowski G, Georgeson E, Kalyuzhniy O, Willis JR,
Kubitz M, Adachi Y, Reiss SM, Shin M, de Val N, Ward AB, Crotty S, Burton DR,
Schief WR. 2017. Structure-based design of native-like HIV-1 envelope trimers to
silence non-neutralizing epitopes and eliminate CD4 binding. Nat Commun 8:1655.
117. Chuang GY, Geng H, Pancera M, Xu K, Cheng C, Acharya P, Chambers M, Druz
A, Tsybovsky Y, Wanninger TG, Yang Y, Doria-Rose NA, Georgiev IS, Gorman
J, Joyce MG, O'Dell S, Zhou T, McDermott AB, Mascola JR, Kwong PD. 2017.
Structure-Based Design of a Soluble Prefusion-Closed HIV-1 Env Trimer with
Reduced CD4 Affinity and Improved Immunogenicity. J Virol 91.
118. Kwon YD, Pancera M, Acharya P, Georgiev IS, Crooks ET, Gorman J, Joyce MG,
Guttman M, Ma X, Narpala S, Soto C, Terry DS, Yang Y, Zhou T, Ahlsen G, Bailer
RT, Chambers M, Chuang GY, Doria-Rose NA, Druz A, Hallen MA, Harned A,

Kirys T, Louder MK, O'Dell S, Ofek G, Osawa K, Prabhakaran M, Sastry M, Stewart-Jones GB, Stuckey J, Thomas PV, Tittley T, Williams C, Zhang B, Zhao H, Zhou Z, Donald BR, Lee LK, Zolla-Pazner S, Baxa U, Schon A, Freire E, Shapiro L, Lee KK, Arthos J, Munro JB, Blanchard SC, Mothes W, Binley JM, et al. 2015. Crystal structure, conformational fixation and entry-related interactions of mature ligand-free HIV-1 Env. Nat Struct Mol Biol 22:522-31.

119. Friedrich N, Stiegeler E, Glogl M, Lemmin T, Hansen S, Kadelka C, Wu Y, Ernst P, Maliqi L, Foulkes C, Morin M, Eroglu M, Liechti T, Ivan B, Reinberg T, Schaefer JV, Karakus U, Ursprung S, Mann A, Rusert P, Kouyos RD, Robinson JA, Gunthard HF, Pluckthun A, Trkola A. 2021. Distinct conformations of the HIV-1 V3 loop crown are targetable for broad neutralization. Nat Commun 12:6705.

120. Kwong PD, Doyle ML, Casper DJ, Cicala C, Leavitt SA, Majeed S, Steenbeke TD, Venturi M, Chaiken I, Fung M, Katinger H, Parren PWIH, Robinson J, Van Ryk D, Wang L, Burton DR, Freire E, Wyatt R, Sodroski J, Hendrickson WA, Arthos J. 2002. HIV-1 evades antibody-mediated neutralization through conformational masking of receptor-binding sites. Nature 420:678-682.

121. Lewis GK, Finzi A, DeVico AL, Pazgier M. 2015. Conformational Masking and Receptor-Dependent Unmasking of Highly Conserved Env Epitopes Recognized by Non-Neutralizing Antibodies That Mediate Potent ADCC against HIV-1. Viruses 7:5115-32.

122. Kwong PD, Doyle ML, Casper DJ, Cicala C, Leavitt SA, Majeed S, Steenbeke TD, Venturi M, Chaiken I, Fung M, Katinger H, Parren PW, Robinson J, Van Ryk D, Wang L, Burton DR, Freire E, Wyatt R, Sodroski J, Hendrickson WA, Arthos J. 2002. HIV-1 evades antibody-mediated neutralization through conformational masking of receptor-binding sites. Nature 420:678-82.

123. Lai JI, Eszterhas SK, Brooks SA, Guo C, Zolla-Pazner S, Seaman MS, Bailey-Kellogg C, Griswold KE, Ackerman ME. 2020. Induction of cross-reactive HIV-1 specific antibody responses by engineered V1V2 immunogens with reduced conformational plasticity. Vaccine 38:3436-3446.

124. Kubota-Koketsu R, Yunoki M, Okuno Y, Ikuta K. 2021. Virus Neutralization by Human Intravenous Immunoglobulin Against Influenza Virus Subtypes A/H5 and A/H7. Biologics 15:87-94.

125. Kasel JA, Couch RB, Gerin JL, Schulman JL. 1973. Effect of influenza anti-neuraminidase antibody on virus neutralization. Infect Immun 8:130-1.

126. Valenzuela Nieto G, Jara R, Watterson D, Modhiran N, Amarilla AA, Himelreichs J, Khromykh AA, Salinas-Rebolledo C, Pinto T, Cheuquemilla Y, Margolles Y, Lopez Gonzalez Del Rey N, Miranda-Chacon Z, Cuevas A, Berking A, Deride C, Gonzalez-Moraga S, Mancilla H, Maturana D, Langer A, Toledo JP, Muller A, Uberti B, Krall P, Ehrenfeld P, Blesa J, Chana-Cuevas P, Rehren G, Schwefel D, Fernandez LA, Rojas-Fernandez A. 2021. Potent neutralization of clinical isolates of SARS-CoV-2 D614 and G614 variants by a monomeric, sub-nanomolar affinity nanobody. Sci Rep 11:3318.

127. Hu J, Peng P, Wang K, Fang L, Luo FY, Jin AS, Liu BZ, Tang N, Huang AL. 2021. Emerging SARS-CoV-2 variants reduce neutralization sensitivity to convalescent sera and monoclonal antibodies. Cell Mol Immunol 18:1061-1063.

128. Garcia-Beltran WF, Lam EC, St Denis K, Nitido AD, Garcia ZH, Hauser BM,
Feldman J, Pavlovic MN, Gregory DJ, Poznansky MC, Sigal A, Schmidt AG,
Iafrate AJ, Naranbhai V, Balazs AB. 2021. Multiple SARS-CoV-2 variants escape
neutralization by vaccine-induced humoral immunity. Cell 184:2372-2383 e9.
129. Diamond M, Chen R, Xie X, Case J, Zhang X, VanBlargan L, Liu Y, Liu J, Errico
J, Winkler E, Suryadevara N, Tahan S, Turner J, Kim W, Schmitz A, Thapa M,
Wang D, Boon A, Pinto D, Presti R, O'Halloran J, Kim A, Deepak P, Fremont D,
Corti D, Virgin H, Crowe J, Droit L, Ellebedy A, Shi PY, Gilchuk P. 2021. SARS-
CoV-2 variants show resistance to neutralization by many monoclonal and serum-
derived polyclonal antibodies. Res Sq doi:10.21203/rs.3.rs-228079/v1.
130. Cheng CC, Platen L, Christa C, Tellenbach M, Kappler V, Bester R, Liao BH,
Holzmann-Littig C, Werz M, Schonhals E, Platen E, Eggerer P, Treguer L, Kuchle
C, Schmaderer C, Heemann U, Renders L, Protzer U, Braunisch MC. 2022.
Improved SARS-CoV-2 Neutralization of Delta and Omicron BA.1 Variants of
Concern after Fourth Vaccination in Hemodialysis Patients. Vaccines (Basel) 10.
131. Poole J, Day CJ, von Itzstein M, Paton JC, Jennings MP. 2018. Glycointeractions
in bacterial pathogenesis. Nat Rev Microbiol 16:440-452.
132. Sheng Y, Vinjamuri A, Alvarez MRS, Xie Y, McGrath M, Chen S, Barboza M,
Frieman M, Lebrilla CB. 2022. Host Cell Glycocalyx Remodeling Reveals SARS-
CoV-2 Spike Protein Glycomic Binding Sites. Front Mol Biosci 9:799703.
133. Clausen TM, Sandoval DR, Spliid CB, Pihl J, Perrett HR, Painter CD, Narayanan
A, Majowicz SA, Kwong EM, McVicar RN, Thacker BE, Glass CA, Yang Z,
Torres JL, Golden GJ, Bartels PL, Porell RN, Garretson AF, Laubach L, Feldman
J, Yin X, Pu Y, Hauser BM, Caradonna TM, Kellman BP, Martino C, Gordts P,
Chanda SK, Schmidt AG, Godula K, Leibel SL, Jose J, Corbett KD, Ward AB,
Carlin AF, Esko JD. 2020. SARS-CoV-2 Infection Depends on Cellular Heparan
Sulfate and ACE2. Cell 183:1043-1057 e15.
134. Ugolini S, Mondor I, Sattentau QJ. 1999. HIV-1 attachment: another look. Trends
Microbiol 7:144-9.
135. Ng PS, Bohm R, Hartley-Tassell LE, Steen JA, Wang H, Lukowski SW, Hawthorne
PL, Trezise AE, Coloe PJ, Grimmond SM, Haselhorst T, von Itzstein M, Paton
AW, Paton JC, Jennings MP. 2014. Ferrets exclusively synthesize Neu5Ac and
express naturally humanized influenza A virus receptors. Nat Commun 5:5750.
136. Bobardt MD, Saphire AC, Hung HC, Yu X, Van der Schueren B, Zhang Z, David
G, Gallay PA. 2003. Syndecan captures, protects, and transmits HIV to T
lymphocytes. Immunity 18:27-39.
137. Li L, Wang XH, Williams C, Volsky B, Steczko O, Seaman MS, Luthra K, Nyambi
P, Nadas A, Giudicelli V, Lefranc MP, Zolla-Pazner S, Gorny MK. 2015. A broad
range of mutations in HIV-1 neutralizing human monoclonal antibodies specific for
V2, V3, and the CD4 binding site. Mol Immunol 66:364-374.
138. Moore JP, Sattentau QJ, Yoshiyama H, Thali M, Charles M, Sullivan N, Poon SW,
Fung MS, Traincard F, Pinkus M, et al. 1993. Probing the structure of the V2
domain of human immunodeficiency virus type 1 surface glycoprotein gp120 with
a panel of eight monoclonal antibodies: human immune response to the V1 and V2
domains. J Virol 67:6136-51.

139.   McKeating JA, Shotton C, Cordell J, Graham S, Balfe P, Sullivan N, Charles M,
1321   Page M, Bolmstedt A, Olofsson S, et al. 1993. Characterization of neutralizing
1322   monoclonal antibodies to linear and conformation-dependent epitopes within the
1323   first and second variable domains of human immunodeficiency virus type 1 gp120.
1324   J Virol 67:4932-44.
1325   140.   Tuen M, Bimela JS, Banin AN, Ding S, Harkins GW, Weiss S, Itri V, Durham AR,
1326   Porcella SF, Soni S, Mayr L, Meli J, Torimiro JN, Tongo M, Wang X, Kong XP,
1327   Nadas A, Kaufmann DE, Brumme ZL, Nanfack AJ, Quinn TC, Zolla-Pazner S,
1328   Redd AD, Finzi A, Gorny MK, Nyambi PN, Duerr R. 2019. Immune Correlates of
1329   Disease Progression in Linked HIV-1 Infection. Front Immunol 10:1062.
1330   141.   Liu L, Li L, Nanfack A, Mayr LM, Soni S, Kohutnicki A, Agyingi L, Wang XH,
1331   Tuen M, Shao Y, Totrov M, Zolla-Pazner S, Kong XP, Duerr R, Gorny MK. 2019.
1332   Anti-V2 antibody deficiency in individuals infected with HIV-1 in Cameroon.
1333   Virology 529:57-64.
1334

Figure. 1. Virus-binding assay for detection of antibody binding to Env expressed on virions. (A) Schematic of the assay; (B) Optimization of virus amount to be coupled to the microspheres; (C) Representative dot plots and histogram showing the gating strategy for analysis of antibodies binding to virus particle by flow cytometry. The mAbs were tested at following concentrations: NIH45-46 at 5 μg/ml; 2219 and 3685 at 50 μg/ml. 1°, primary; 2° secondary.

Figure 2. Binding of mAbs to REJO Env on virions. (A) Microspheres coupled to REJO virus were reacted with serially diluted mAbs targeting different Env epitopes. (B) Area under the curve (AUC) were calculated from the curves in A and plotted. (C) Geometric mean fluorescent intensity (MFI) of each mAb tested at a single dilution are plotted. Virus coupled microspheres stained with biotin and PE alone were used to set the background MFI and subtracted.

Figure 3. Binding of mAbs to SF162 Env on virions. (A) Microspheres coupled to SF162 virus were reacted with serially diluted mAbs targeting different Env epitopes. Virus coupled microspheres stained with biotin and PE alone were used to set the background MFI and subtracted. (B) Area under the curve (AUC) were calculated from the curves in A and plotted. (C) Geometric mean fluorescent intensity (MFI) of each mAb tested at a single dilution are plotted.

Figure 4. Binding of mAbs to CMU06 Env on virions. (A) Microspheres coupled to CMU06 virus were reacted with mAbs targeting different Env epitopes. Geometric mean fluorescent intensity (MFI) of each mAb tested at a single dilution are plotted. Virus coupled microspheres stained with biotin and PE alone were used to set the background MFI and subtracted.

Figure 5. Binding of mAbs to trimeric Env expressed on cell surface. 293T cells were transiently transfected with plasmid encoding for full length gp160 of (A) REJO, (B) CMU06 and (C) SF162. Cells were analyzed for binding of different mAbs 24-hours post-transfection. Geometric mean fluorescence intensity (MFI) and SD from duplicates or triplicates in one experiment are shown. Transfected cells stained with biotin and PE alone were used to set the background MFI and subtracted. The mAbs were tested at following concentrations: 697, 830A, 2158, 3685 at 100 µg/ml; 2219, 2557 at 50 µg/ml; PG9, PG16, PGT145, PGDM1400, CH59, VRC01, 3BNC117, b12 at 25 µg/ml; 4E10, 2F5 at 20 µg/ml; PGT151 at 10 µg/ml; NIH45-46 at 5 µg/ml.

Figure 6. Time-dependent increase of Env binding. Microspheres coupled to REJO virus particles were treated with mAbs at 37°C for various time from 0 to 75 min. Binding was detected as in Figure 2 and geometric mean fluorescence intensity (MFI) are shown. Virus coupled microspheres stained with biotin and PE alone were used to set the background MFI and subtracted. The mAbs were tested at following concentrations: 830A, 3685 at 100 µg/ml; 2219 at 50 µg/ml; PG9 at 25 µg/ml; NIH45-46 at 2.5 µg/ml. ****, p= <0.0001; **, p= 0.0026 vs t = 0 min by ANOVA.

Figure 7. Changes in reactivity of PG9 and 2219 mAbs. Microspheres coupled with REJO virus were incubated with titrated amounts of antibodies followed by biotinylated V2q mAb PG9 (25 µg/ml) or biotinylated V3 mAb 2219 (50 µg/ml). Binding was detected with streptavidin-PE and geometric mean fluorescent intensity (MFI) are plotted. Virus coupled microspheres stained with biotin and PE alone were used to set the background MFI and subtracted. **** $p < 0.0001$ by ANOVA.

Figure 8. Neutralization of REJO virus. Virus neutralization was measured after REJO virus was incubated with serially diluted mAbs for the designated period of time at 37°C prior to the addition of TZM-bl target cells. Virus infectivity was assessed 48 h later based on β-galactosidase activity. (A) Neutralization curves are shown. Means and standard errors calculated from two different experiments (each in duplicate) are shown. Statistical analyses were performed on the neutralization curves reaching ≥50%. Comparison is made between neutralization curves at T = 0 vs T = 60 minutes. P= ns, not significant by nonparametric Mann-Whitney t-test. (B) Area under the neutralization curves were calculated and plotted as bar graph. Statistical analysis was performed on AUC graph by ANOVA (P = 0.99; ns, not significant).

Figure 9. Neutralization of CMU06 virus. Virus neutralization was measured after CMU06 virus was incubated with serially diluted mAbs for the designated period of time at 37°C prior to the addition of TZM-bl target cells. Virus infectivity was assessed 48 h later based on β-galactosidase activity. (A) Neutralization curves are shown. Means and standard errors calculated from two different experiments (each in duplicate) are shown. Statistical analyses were performed on the neutralization curves reaching ≥50%. Comparison is made between neutralization curves at T = 0 vs T = 60 minutes. P= ns, not significant by nonparametric Mann-Whitney t-test. (B) Area under the neutralization curves were calculated and plotted as bar graph. Statistical analysis was performed on AUC graph by ANOVA (P = 0.99; ns, not significant).

Figure 10. Influence of virus-cell interaction on Env epitope exposure. Binding of mAbs to virus-coupled microspheres were measured in the presence of 293T cells by flow cytometry. Cells were transfected or not transfected with human CD4 plasmid for 24 h and treated with virus-coupled microspheres and V2i, V3, V2q, CD4bs or control mAbs at 4°C. MAb binding was detected with biotin-conjugated anti-human IgG and PE-Streptavidin. Percent PE positive microspheres from one experiment are shown. Gating strategy used for data analysis is shown in Fig S10. mAb titration curves and bar graphs showing geometric mean fluorescent intensity (MFI) of each mAb tested at a highest dilution are plotted. Gray dotted line represents MFI of negative control mAb 3685.

Figure 11. Effect of lectin Concanavalin A on exposure of Env epitopes. Microspheres coupled with REJO virus were incubated with five-fold titrated amounts of lectin Concanavalin A (ConA) (50 - 0.08 µg/ml) followed by mAbs at single dilutions (830A, 2219, PG9, 3685 at 50 µg/ml; VRC01 at 25 µg/ml and NIH45-46 at 2.5 µg/ml). (50 µg/ml). Binding was detected with streptavidin-PE. Geometric mean fluorescent intensity (MFI) and fold change in binding relative to no ConA are shown. Virus coupled microspheres stained with biotin and PE alone were used to set the background MFI and subtracted. **** p= 0.0002, * p= 0.039 by ANOVA relative to negative control mAb 3685.

Figure 12. Effect of Lectins on Env epitope exposure in ELISA. Sucrose pelleted REJO virus was captured by (A) five-fold titrated lectin ConA (50 - 0.08 µg/ml) and reacted with different mAbs at one dilution (830A, 2219, PG9, 3685 at 50 µg/ml; VRC01 at 25 µg/ml and NIH45-46 at 2.5 µg/ml). (B) Sucrose pelleted REJO virus was captured by lectin Griffithsin (50 µg/ml) and reacted with five-fold titrated mAbs (830A, 2219, PG9, 3685 at 50 µg/ml; VRC01 at 25 µg/ml and NIH45-46 at 2.5 µg/ml). AUC values were calculated from binding curves. AUC changes relative to the respective WT are shown. Statistical analysis was performed on binding curves by 2-way ANOVA (**** $p < 0.0001$) relative to negative control mAb 3685.

Table 1. Binding specificity of the monoclonal antibodies (mAbs) used in the study.

Antibody name	Epitope/ binding site	Structural preference
697 830A 2158	gp120 V1V2 (V2i)	Conformational
CH58 CH59	gp120 V1V2 (V2p)	Linear
PG9 PG16 PGT145	gp120 V1V2 (V2q)	Conformational
2219 2557	gp120 V3	Linear
NIH45-46 VRC01 b12 3BNC117 CD4-IgG	gp120 CD4 binding site (CD4BS)	Conformational
17b	CD4 induced epitopes (CD4i)	
PGT151	gp120-gp41 interface	
VRC34	gp41 fusion peptide/gp120 glycan	
4E10 2F5	gp41 MPER	Linear
3685	anti-anthrax	Non-binding