Effects of early life stress paired with adolescent alcohol consumption on two-bottle choice alcohol drinking behaviors in mice

Thomas W. Perry, Harrison M. Carvour, Amanda N. Reichert, Elizabeth A. Sneddon, Charlotte A.E.G. Roemer, Ying Ying Gao, Kristen M. Schuh, Natalie A. Shand, Jennifer J. Quinn, & Anna K. Radke

Department of Psychology and Center for Neuroscience and Behavior, Miami University, Oxford, OH, USA

Correspondence to: Anna K. Radke, Ph.D., 90 N Patterson Ave, Oxford, OH, USA 45056

email: aradke@miamioh.edu

phone: 513-529-6941

fax: 513-529-2420

Funding sources: This work was supported by NIH grants R21 AA028602 (AKR and JJQ) and F99 NS118727 (EAS), the Psychology Department and the Office of Research for Undergraduates at Miami University, and a Psi Chi Graduate Research Award (ANR).

Conflict of interests: The authors declare no conflicting interests in this work.

Abstract

Background: In humans, early life stress (ELS) is associated with an increased risk for developing both alcohol use disorder (AUD) and post-traumatic stress disorder (PTSD). We previously used an infant footshock model that produces stress-enhanced fear learning (SEFL) in rats and mice and increases aversion-resistant alcohol drinking in rats to explore this shared predisposition. The goal of the current study was to extend this model of comorbid PTSD and AUD to male and female C57BL/6J mice.

Methods: Acute ELS was induced using 15 footshocks on postnatal day (PND) 17. In adulthood, alcohol drinking behavior was tested in one of three two-bottle choice drinking paradigms. In continuous access, mice were given 24 h access to 5% or 10% ethanol and water for five consecutive drinking sessions each. In limited access drinking in the dark, mice were given 2 h of access to 15% ethanol and water across 15 sessions 3 h into the dark cycle. In intermittent access, mice were presented with 20% ethanol and water Monday, Wednesday, and Friday, for four consecutive weeks. In a fifth week of intermittent access drinking, increasing concentrations of quinine (10 mg/L, 100 mg/L, and 200 mg/L) were added to the ethanol to test aversion-resistant drinking. Intermittent access drinking was tested with and without a period of adolescent drinking (PND 35).

Results: Infant footshock did not alter drinking in the continuous or limited access tasks. Adult consumption and preference were lower in the intermittent access task when adolescent drinking was included and there were ELS-induced differences in consumption of quinine-adulterated ethanol in females.

Conclusions: Our results demonstrate that infant footshock followed by a period of adolescent drinking is a viable model of comorbid PTSD and AUD in rats and mice.

Introduction

Alcohol use disorder (AUD) is a complex psychiatric disorder influenced by both genetic and environmental factors (Mayfield et al., 2008; Tawa et al., 2016; Young-Wolff et al., 2011). Stress is one environmental variable that is strongly associated with alcohol abuse and the development of AUD (Guinle and Sinha, 2020; Sinha, 2001). Among individuals with PTSD, AUD is reported to be up to three times more prevalent than it is in the general public (Gilpin and Weiner, 2017; Kessler et al., 1995; Smith and Cottler, 2018). This suggests that PTSD and AUD may be products of common alterations brought about by stress. Early life stress (ELS) in humans is associated with an increased risk of alcohol dependence, early onset alcohol use, and PTSD-AUD comorbidity (Enoch, 2011; Lee et al., 2018; Pechtel and Pizzagalli, 2011). These associations indicate that in early stages of development the brain may be particularly susceptible to stress-induced changes (Lupien et al., 2009).

Consistent with clinical literature, in preclinical models, chronic ELS in the form of isolation or maternal separation generally increases levels of alcohol consumption later in adulthood (Becker et al., 2011; Cruz et al., 2008; McCool and Chappell, 2009). Additionally, a number of studies have found that ELS increases aversion-resistant drinking when alcohol is paired with an aversive stimulus such as quinine (Bertagna et al., 2021; Radke et al., 2020; Shaw et al., 2020). This drinking despite a negative consequence has been used to model compulsive-like drinking, a central characteristic of AUD (Hopf et al., 2010; Hopf and Lesscher, 2014; Radke et al., 2017; Sneddon et al., 2019). Sex has been reported to play a role in the relationship between ELS and alcohol-drinking behaviors; however, the exact influence of sex remains unclear (Bertagna et al., 2021; Radke et al., 2020; Shaw et al., 2020).

In a recent study using Long-Evans rats, we found that acute ELS in the form of repeated footshock on postnatal day (PND) 17 increased aversion-resistant drinking in an intermittent access alcohol drinking paradigm (Radke et al., 2020). This form of acute infantile trauma is commonly used to sensitize fear learning, a phenomenon known as stress-enhanced fear learning (SEFL), as a preclinical model for PTSD and PTSD susceptibility (e.g., Poulos et al., 2014; Quinn et al., 2014; Minshall et al., in press). The common influence of infant footshock on disordered drinking behavior and fear learning, but

not other forms of hippocampal-dependent learning or cognitive flexibility (Sneddon et al., 2021),

suggests that an acute infant footshock paradigm may be a viable stress model for investigating AUD-

PTSD comorbidity, as well as the shared neural mechanisms underlying both disorders.

To extend the utility of this preclinical model, the current study explored how acute ELS in the

form of infant footshock influences consumption of alcohol alone and when paired with an aversive

outcome in C57BL/6J mice. We used an infant stress protocol (15 footshocks on PND 17) that we have

previously found increases fear learning and resistance to extinction in mice (Sneddon et al., 2021). To

comprehensively characterize the effects of infant footshock on drinking, adult mice were exposed to

alcohol in one of three two-bottle choice drinking tasks: limited access drinking in the dark, continuous

access, or intermittent access. Based on previous findings in rats that had undergone PND 17 footshock

and alcohol drinking (Radke et al., 2020), we ran a second intermittent access task where mice were

presented with alcohol during adolescence (PND 35-55) to measure the necessity of alcohol exposure

during that developmental stage on adult outcomes. To model aversion-resistant drinking, alcohol was

adulterated with increasing concentrations of quinine, an aversive bittering agent, in the drinking in the

dark and intermittent access paradigms. Additionally, considering published findings (Hennessy et al.,

2009) and data from our own labs (Reichert et al., unpublished data) showing that exposure to

conspecifics following trauma, otherwise known as social buffering, can modulate the effect of trauma,

we investigated whether social buffering could alter the effects of infantile trauma on intermittent access

drinking. Finally, due to robust evidence of enhanced vulnerability to both stress-related disorders and

addictive behaviors in females (Bangasser and Valentino, 2014; Minshall et al., 2024; Quigley et al.,

2021; Radke et al., 2021b), we examined sex as a potential moderator of the effects of infant footshock on

alcohol drinking in all experiments.

Materials and Methods

Subjects

C57BL/6J mice were generated from breeding pairs from The Jackson Laboratory (Bar Harbor,

ME, USA). Mice were provided food (Rodent Diet 5001, Cincinnati Lab Supply, Cincinnati, OH) and

reverse osmosis (RO) water *ad libitum*. Mice were on a 12:12 light/dark cycle. Other than PND 17

footshock and adolescent alcohol exposure, all behavioral tests were conducted during adulthood (PND

60+). During infancy, all mice were briefly anesthetized with isoflurane and given an ear snip for

identification purposes. Prior to home cage drinking, mice were group housed (2-4 mice/cage). One week

before the first drinking session mice were individually housed in standard shoe box Udel polysulfone

rectangular mouse cages (18.4 cm x 29.2 cm x 12.7 cm) outfitted with 2-bottle cage tops. All animals

were cared for in accordance with the guidelines set by the National Institute for Health. All procedures

were approved by the Institutional Animal Care and Use Committee (IACUC) at Miami University.

Early life stress exposure

Mice were placed in a novel MED-Associates conditioning chamber (Context A) on PND 17

(after Quinn et al., 2014). Context A was brightly lit, contained a uniform grid floor, was scented with

vanilla, and was cleaned with odorless 5% sodium hydroxide. Mice received either 15 footshocks (1 mA,

1 sec) or no footshocks during a 60-min session beginning at 60 sec following placement in the chamber.

Shocks were delivered according to a variable inter-shock interval of 120-240 sec. Progressive scan video

cameras containing visible light filters (VID-CAMMONO-2A; Med Associates, Inc.) monitored mice

throughout the session.

For the continuous access and drinking in the dark experiments, mice were returned to a cage

with their littermates in a brightly lit room separate from the conditioning chambers following shock

exposure. Mice were run in squads of 4 and the entire litter was returned to the home cage 3 h after the

final squad's session.

For the first intermittent access experiment (Experiment 3), we discovered an issue with the

shock delivery boxes that resulted in animals receiving a variable number of shocks on PND 17. After

determining that 6 or more footshocks produced the SEFL effect (i.e., significantly higher fear learning in

adult mice exposed to shock vs. no shock) (Reichert et al., unpublished data), we excluded mice that

received fewer than 6 footshocks from the drinking analyses. In this experiment, we also examined the

effects of social buffering on SEFL (Reichert et al., unpublished data) and alcohol drinking in adulthood.

Immediately following the ELS exposure session, mice were placed into their assigned social buffering

condition for 90 minutes. During this time, pups in the isolation condition were housed individually while

pups in the littermates condition were housed as a group. All buffering took place in polysulfone holding

cages (19 x 29.2 x 12.7 cm) that are similar to the housing cages but without bedding. All holding cages

were in a room with overhead lighting and were placed on top of Sunbeam-brand heated mats, half on the

pad and half off. For the second intermittent access experiment (Experiment 4), all mice were in the

isolation condition following footshock.

Fear Conditioning and Memory Test

Before the commencement of drinking, on PND 60, mice that took part in intermittent access

drinking (Experiments 3 and 4) were fear conditioned in a novel context. This context consisted of four

identical conditioning chambers (32.4 x 25.4 x 21.6 cm; Med-Associates, Inc., Georgia, VT) housed

inside sound-attenuating cabinets. The chambers were distinct from the PND 17 footshock and isolation

contexts in several ways: the room was dark except lit by red lamps, the room smelled of vinegar, there

were black Plexiglass triangular inserts within the chamber, and the floor had staggered stainless steel

rods (18 rods, two rows, .05 cm vertically apart). Mice were placed into the fear conditioning chamber

and received one footshock (0.25 mA, 1s) 180 seconds following placement within the chamber. Thirty

seconds following footshock, mice were returned to the home cage. For Experiment 3, mice were

administered a fear retention test session in this context on PND 90 to test incubation of fear (data not

shown).

Two-bottle choice ethanol drinking

For all experiments, mice were given access to RO drinking water and ethanol in RO water (v/v)

in their home cage. Mice were weighed at the beginning of every session and solutions were made fresh

for each drinking session. Bottles were alternated each session to equate side biases. Two "dummy" cages

were outfitted with bottles to account for spillage and evaporation.

Experiment 1: Continuous access ethanol drinking

Forty mice (20 male and 20 female) were given access to 5% or 10% ethanol (**Figure 1A**). Each

ethanol concentration was provided for 5 consecutive, 24-h continuous drinking sessions and bottle

weights were recorded every 24 h.

Experiment 2: Limited access drinking in the dark

Fifty-five mice (28 male and 27 female) were presented with 15% ethanol following a limited

access, two-bottle choice protocol for 2 h a day, five day/week (Monday-Friday) (**Figure 2A**). Bottles

were placed on the home cage 3 h after lights out and bottle weights were recorded before and after each

session.

Experiment 3: Intermittent access ethanol drinking in adulthood

Sixty-six mice (37 male and 24 female) were presented with 20% ethanol following a two-bottle

choice, intermittent access protocol for 5 weeks (**Figure 3A**). Ethanol-drinking sessions took place on

Monday, Wednesday, and Friday. On intervening sessions (Tuesday, Thursday, Saturday, and Sunday),

mice had access to two bottles of drinking water only. Bottle weights were recorded at the beginning of

each ethanol-drinking session, after 30 min, and after 24 h. After 4 weeks (12 sessions) of drinking,

quinine hemisulfate was added to the ethanol bottle in increasing concentrations of 10 mg/L, 100 mg/L,

and 200 mg/L (equivalent to 25, 250, and 500 µM).

Experiment 4: Intermittent access ethanol drinking in adolescence and adulthood

Following ELS on PND 17, 40 mice (21 male and 27 female) underwent a 3-week intermittent

access procedure where they received 20% ethanol and RO water on Mondays, Wednesdays, and Fridays

between PND 35-55 (**Figure 4A**). On intervening days, mice received two bottles of RO water. Mice

were fear conditioned on PND 60 after completing adolescent intermittent access (data not shown). After

fear conditioning, mice started adult intermittent access for 5 weeks (17 male and 24 female) (**Figure

5A**). Mice received 20% ethanol and RO water every other day. On intervening days, mice received RO

water. During week 5, escalating concentrations of quinine hemisulfate (10 mg/L, 100 mg/L, and 200

mg/L) were mixed with 20% ethanol on Monday, Wednesday, and Friday.

Following the fifth week, mice were given only RO water for a week, before undergoing a

quinine sensitivity test as a control. For this test, mice were given two bottles of RO water on Monday to

establish a baseline. After establishing this baseline, mice were given escalating concentrations of quinine

(10 mg/L, 100 mg/L, and 200 mg/L) in one of the RO water bottles on Wednesday, Friday, and the

following Monday.

Data Analysis

Bottle weights were converted to grams (g) of ethanol or milliliters (mL) of water for each

mouse. Consumption was calculated as = *(Initial Bottle Weight – Final Bottle Weight) – Average Weight*

Loss of Dummy Bottles. Preference was calculated as = *(Volume of Ethanol [Volume of Ethanol + Volume*

of Water]) x 100. Consumption and preference were averaged across mice for each group and analyzed

using three-way repeated measures (RM) ANOVA with infant footshock, sex, and concentration as

factors. All results were first analyzed with session as a factor, but as no significant interaction effects

were found, data were averaged across sessions for final analysis and visualization. In the adult

intermittent access experiment, social buffering condition (i.e., isolated vs. littermates) was included as a

factor, but the data were ultimately collapsed across as there were no significant effects. The Greenhouse-

Geisser correction was applied when the assumption of sphericity was violated ($\varepsilon < 0.75$). Follow-up

Holm-Sidak's tests were used as appropriate to assess differences between groups. Aversion-resistant

drinking was defined *a priori* as a significant reduction from baseline (= 0 g/L) consumption and was

assessed in all groups with a Dunnett's test to correct for multiple comparisons.

Results

Experiment 1: Continuous access ethanol drinking is not altered by infant footshock exposure.

Sex, but not stress, impacted ethanol consumption in the continuous access drinking test, with

females consuming more than males (**Figure 1B**). A three-way ANOVA revealed no effect of infant

footshock on average ethanol consumption across the two concentrations of ethanol presented ($F_{(1, 36)} =$

0.085, $p = 0.773$). There was, however, a significant main effect of sex on ethanol consumption ($F_{(1, 36)} =$

21.300, p < 0.001) and a main effect of ethanol concentration on consumption ($F_{(1, 36)}$ = 10.550, p =

0.003). An interaction between ethanol concentration, sex, and infant footshock neared significance ($F_{(1,}$

$_{36)}$ = 3.943, p = 0.055). There were no other significant interactions between the factors (all p > 0.409).

Holm Sidak's multiple comparison tests comparing consumption of the 5% and 10% concentrations for

each group, demonstrated a greater consumption of 5% vs. 10% ethanol in 15-shock females only (p =

0.008).

Similar to consumption, preference for ethanol vs. water was higher in females than males but

was not affected by stress (**Figure 1C**). A three-way ANOVA revealed no significant effect of infant

footshock ($F_{(1, 36)}$ = 0.009, p = 0.925) or concentration ($F_{(1, 36)}$ = 3.125, p = 0.086) on ethanol preference.

There was, however, a significant main effect of sex on ethanol preference ($F_{(1, 36)}$ = 11.220, p = 0.002).

There was also a significant interaction between ethanol concentration, sex, and infant footshock ($F_{(1, 36)}$ =

5.182, p = 0.029). There were no other significant interactions between the factors (all p > 0.487). Holm

Sidak's multiple comparison tests comparing preference for ethanol across concentrations in each group

found that 15-shock females had significantly higher preference for 5% vs. 10% ethanol (p = 0.047).

Experiment 2: Limited access drinking in the dark is not altered by infant footshock exposure.

When assessing limited access consumption, there were no effects of stress, but females

consumed more than males (**Figure 2B**). A two-way ANOVA revealed no significant effect of infant

footshock on ethanol consumption ($F_{(1, 51)}$ = 0.245, p = 0.623). Analysis, however, indicated a significant

main effect of sex ($F_{(1, 51)}$ = 22.100, p < 0.001). The interaction between the factors was not significant

($F_{(1, 51)}$ = 0.328, p = 0.569). For ethanol preference, there were no main effects of sex ($F_{(1, 51)}$ = 0.165, p =

0.687) or stress ($F_{(1, 51)}$ = 1.637, p = 0.207) and the interaction was not significant ($F_{(1, 51)}$ = 0.012, p =

0.912) (**Figure 2C**).

Experiment 3: Intermittent access ethanol drinking in adulthood is not altered by infant footshock

exposure.

We next assessed whether infant footshock influenced adult drinking using an intermittent access

drinking task. There were no effects of social buffering observed on any measure, therefore data from

mice housed alone vs. with littermates following the ELS session on PND 17 were collapsed for

visualization and analysis. Additionally, mice in this experiment received a variable number of footshocks

(between 6 and 15) due to an equipment malfunction (average number of shocks males = 11.56 ± 0.72,

females = 10.40 ± 0.90) but still displayed significant increases in fear conditioning (data not shown).

Ethanol consumption averaged across the first twelve sessions of intermittent access drinking was

higher in females but not affected by stress (**Figure 3B**). A two-way ANOVA revealed no significant

effect of infant footshock on average consumption of ethanol ($F_{(1, 57)}$ = 0.057, p = 0.812). There was,

however, a significant main effect of sex on average ethanol consumption ($F_{(1, 57)}$ = 7.490, p = 0.008). No

significant interaction effect was indicated ($F_{(1, 57)}$ = 2.289, p = 0.136). Ethanol preference was also

similar across groups, regardless of sex or stress exposure (**Figure 3C**). A two-way ANOVA revealed no

significant effect of infant footshock ($F_{(1, 57)}$ = 0.133, p = 0.717), or sex ($F_{(1, 57)}$ = 0.148, p = 0.702) on

average preference for ethanol vs. water. No significant interaction effect was indicated, ($F_{(1, 57)}$ = 2.365, p

= 0.130). To determine whether the absence of a stress effect was due to an insufficient number of shocks

in some animals, we correlated the number of shocks received on PND 17 with average consumption

(males: r = 0.070, p = 0.796; females: r = -0.227, p = 0.528) and preference (males: r = -0.385, p = 0.131;

females: r = -0.096, p = 0.791). These correlations were not significant, further supporting the idea that

infant footshock did not influence consumption or preference in this experiment.

For aversion-resistant drinking on sessions 12-15, females continued to consume more than

males, and quinine reduced consumption equally in all groups (**Figure 3D**). A three-way ANOVA

revealed no significant effect of infant footshock on consumption of quinine adulterated ethanol across

sessions 12-15. ($F_{(1, 57)}$ = 0.403, p = 0.528). There were, however, significant main effects of sex ($F_{(1, 57)}$ =

14.590, p < 0.001) and quinine concentration ($F_{(3, 171)}$ = 56.800, p < 0.001) on consumption. No significant

interaction effects were indicated (all p > 0.257). To assess aversion resistant consumption, two-way

ANOVAs followed by Dunnett's multiple comparisons tests were performed for each sex. In males, there

was a significant main effect of quinine concentration ($F_{(3, 105)}$ = 54.770, p < 0.0001) but no main effect of

stress ($F_{(1, 35)}$ = 2.006, p = 0.166) or interaction between the two factors ($F_{(3, 105)}$ = 0.026, p = 0.994).

Compared to baseline, consumption was decreased in no-shock and 15-shock males at the 100 and 200

g/L concentrations (all $p < 0.0001$), but not at a quinine concentration of 10 g/L ($p = 0.770$ and 0.754). In

females, there was a significant main effect of quinine concentration ($F_{(3, 66)} = 14.390$, $p < 0.0001$).

Neither the main effect of stress nor interaction reached the threshold for significance. Aversion

resistance was observed at the 10 g/L concentration in no-shock ($p = 0.802$) and 15-shock ($p = 0.992$)

females while consumption was reduced at the 100 and 200 g/L concentrations in both groups (all $p <$

0.05).

Experiment 4: Intermittent access ethanol drinking is reduced following exposure to infant footshock and

adolescent drinking.

Because our prior results demonstrating an effect of infant footshock on drinking in rats included

an adolescent drinking period (Radke et al., 2020), we tested the necessity of this exposure in a second

cohort of mice that consumed ethanol from PND 35-55. This experiment revealed that infant footshock

influences adult consumption when preceded by an adolescent drinking period.

Infant footshock did not affect average ethanol consumption across the nine adolescent drinking

sessions (**Figure 4B**). A two-way ANOVA revealed no significant main effects of sex ($F_{(1, 44)} = 1.040$, $p =$

0.313), shock ($F_{(1, 44)} = 1.223$, $p = 0.275$), or an interaction between sex and shock ($F_{(1, 44)} = 0.816$, $p =$

0.371). For average ethanol preference across these sessions, a two-way ANOVA revealed there was a

significant interaction between sex and shock ($F_{(1, 44)} = 5.126$, $p < 0.05$) (**Figure 4C**). However, there

were no significant main effects of sex ($F_{1, 44} = 0.170$, $p = 0.682$) or shock ($F_{(1, 44)} = 0.571$, $p = 0.454$) and

follow-up Holm-Sidak's tests comparing preference between shocked groups in males and females were

not significant ($p > 0.108$).

In mice that consumed ethanol during adolescence, infant footshock affected average adult

ethanol consumption across twelve sessions (**Figure 5B**). A two-way ANOVA revealed significant main

effects of sex ($F_{(1, 37)} = 10.870$, $p < 0.01$) and shock ($F_{(1, 37)} = 12.250$, $p < 0.01$). However, there was no

significant interaction between sex and shock ($F_{(1, 37)} = 0.0143$, $p = 0.906$), demonstrating that infant

footshock exposure produced lower levels of adult alcohol consumption in mice that drank alcohol as

adolescents. For average ethanol preference across twelve adult drinking sessions, a two-way ANOVA

revealed a significant main effect of shock ($F_{(1, 36)}$ = 7.006, p < 0.05), but not sex ($F_{(1, 36)}$ = 0.037, p =

0.8498) (**Figure 5C**). Additionally, there was no significant interaction between sex and shock ($F_{(1, 36)}$ =

0.0006, p = 0.981).

Consumption of quinine-adulterated ethanol was affected by infant footshock in mice that

consumed ethanol during adolescence (**Figure 5D**). A three-way ANOVA revealed significant main

effects of quinine concentration ($F_{(2.724, 98.05)}$ = 81.760, p < 0.0001), sex ($F_{(1, 36)}$ = 14.530, p < 0.001), and

shock ($F_{(1, 36)}$ = 17.810, p < 0.001). There was also a significant interaction between quinine

concentration, sex, and shock ($F_{(3, 108)}$ = 3.045, p < 0.05). However, there were no other significant

interaction effects (all p > 0.3111). To assess aversion-resistant ethanol consumption, two-way ANOVAs

followed by Dunnett's multiple comparisons tests were performed for each sex. In males, there was a

significant main effect of quinine concentration ($F_{(2.096, 29.34)}$ = 25.330, p < 0.0001) and shock ($F_{(1, 14)}$ =

10.550, p < 0.01), but no interaction between quinine concentration and shock ($F_{(3, 42)}$ = 1.908, p = 0.143).

Compared to baseline, consumption was significantly decreased in both no shock and 15 shock males at

the 100 and 200 mg/L concentrations (all < 0.01), but not at 10 mg/L (p = 0.807 and 0.211). In females,

there was a significant main effect of quinine concentration ($F_{(3, 66)}$ = 67.770, p < 0.0001) and shock ($F_{(1,}$

$_{22)}$ = 6.873, p < 0.05), but no interaction between quinine concentration and shock ($F_{(3, 66)}$ = 0.822, p =

0.486). Compared to baseline, consumption was significantly decreased in both no-shock and 15-shock

females at the 100 and 200 mg/L concentrations (all p < 0.0001). However, the 15-shock females

demonstrated aversion resistance, as they did not significantly decrease consumption compared to

baseline at 10 mg/L (p = 0.405), while the no-shock females did decrease (p < 0.05).

As a control, we examined sensitivity to quinine by administering quinine in RO drinking water

one week after the cessation of ethanol drinking. This experiment revealed no effects of stress or sex on

quinine sensitivity in mice exposed to infant footshock and adolescent ethanol (**Table 1**). A three-way

RM ANOVA uncovered a main effect of concentration ($F_{(3, 108)}$ = 21.710, p < 0.0001) and a main effect of

sex ($F_{(1, 36)}$ = 6.899, p < 0.05), the latter of which agrees with our prior demonstrations of greater water

consumption in female vs. male mice (Sneddon et al., 2023, 2022). There were no other significant main

effects of interactions (all $p > 0.320$). To determine which concentrations of quinine produced aversion

resistance, follow up two-way ANOVAs and Dunnett's tests were performed. In males, there was a

significant main effect of concentration ($F_{(3, 42)} = 14.850$, $p < 0.0001$) but not stress. Consumption of water

adulterated with 10 mg/L quinine was not reduced in either group (no shock: $p = 0.585$, 15 shock: $p =$

0.975) while consumption at the 100 mg/L (no shock: $p = 0.007$, 15 shock: $p = 0.047$) and 200 mg/L (no

shock: $p < 0.0001$, 15 shock: $p = 0.010$) concentrations was reduced in all animals. Results in females

were similar, with a main effect of concentration ($F_{(3, 66)} = 13.980$, $p < 0.0001$) but not stress.

Consumption was significantly reduced from baseline at the 100 mg/L (no shock: $p = 0.0003$, 15 shock: p

$= 0.003$) and 200 mg/L (no shock: $p = 0.002$, 15 shock: $p = 0.003$) concentrations in both groups, but not

the 10 mg/L concentration.

Discussion

The central finding of this study is that acute exposure to footshock on PND 17, using a protocol

that enhances adult fear learning (Sneddon et al., 2021), affects alcohol drinking behavior in C57BL/6J

mice when followed by a period of adolescent intake. Mice exposed to ELS followed by adolescent

alcohol had lower alcohol preference and consumption as adults than no-shock mice. In addition, infant

stress affected consumption of quinine-adulterated alcohol only in mice that underwent adolescent

drinking. In these mice, shocked females did not avoid the lowest quinine concentration while no-shock

females did, although we interpret these results with caution as discussed below. There was also a clear

effect of sex on drinking behaviors, such that females consumed more alcohol than males when

accounting for body weight. However, preference for alcohol vs. water was only higher in females under

continuous access conditions.

In a prior study using Long-Evans rats, which investigated alcohol consumption in an intermittent

access paradigm following an adolescent drinking period, we demonstrated that PND 17 footshock

increased aversion-resistant drinking in male and female subjects but did not alter consumption of

unadulterated alcohol (Radke et al., 2020). The current studies were designed to examine whether these

effects could be replicated in mice and with other drinking protocols. After failing to find effects of infant

footshock on adult consumption using continuous access, limited access, and intermittent access, we

tested the necessity of the adolescent drinking period. Our results revealed an effect of infant footshock

on alcohol consumption in mice that consumed alcohol in adolescence, which aligns with the original

protocol tested in rats (Radke et al., 2020). Thus, we conclude that the infant SEFL protocol can influence

adult alcohol drinking behaviors across species and sexes when combined with adolescent alcohol

drinking, although the manifestation of the behavioral change differs in rats and mice. For example, infant

footshock had no effect on levels of alcohol consumption and preference in rats but decreased both

measures of drinking in mice. Further, infant footshock increased the concentration of quinine in alcohol

tolerated by rats of both sexes, while the results in mice were less robust. Female mice in the 15-shock

group were resistant to 10 mg/L quinine compared to their no-shock counterparts, but this effect should

be interpreted with caution considering that mice also did not decrease consumption when this

concentration of quinine was added to water. Although we have demonstrated that 10 mg/L quinine in

water is avoided by rats and mice previously (Carvour et al., 2024; Radke et al., 2020), since it was not

avoided in this experiment we cannot definitively say that 15-shock females were resistant to aversion.

Because we saw increased aversion-resistance when using the same protocol in rats, we expected

the infant footshock protocol to similarly enhance vulnerability to alcohol drinking behaviors in mice.

Thus, the observation of decreased consumption and preference was surprising and more difficult to

reconcile with the hypothesis that ELS enhances vulnerability to alcohol. One possibility is that infant

footshock followed by adolescent alcohol consumption produces an anhedonic affective state in adults,

leading to reductions in reward-seeking (Hu et al., 2020). This hypothesis could be tested by assessing

operant responding for other drugs and natural rewards, intracranial self-stimulation, or social interaction.

Another consideration is that adolescent drinking often causes an increase in adult alcohol consumption

(Spear, 2016) that may have been prevented in shocked mice. Because Experiments 3 and 4 were

conducted separately, we cannot statistically compare the results. However, there is a notable trend

toward increased consumption in the no-shock animals between the two experiments (Experiment 3

average consumption male = 12.66 ± 0.89, female = 17.43 ± 1.45; Experiment 4 average consumption

male = 16.61 ± 1.50, female = 21.59 ± 1.14) that was not seen in the shocked mice. In either case, our

results suggest that ELS and adolescent alcohol consumption engage overlapping mechanisms to produce

alterations in adult behavior.

In preclinical studies, stress has been found to have wide-ranging effects on alcohol drinking

(e.g., Becker et al., 2011; Radke et al., 2014). At least one prior review article estimated that acute stress

increased drinking in fewer than 50% of studies (Becker et al., 2011). The limited and/or variable effects

of acute stress exposure on drinking are in contrast with chronic stress exposure, which increase drinking

behaviors much more consistently, especially when administered in the early stages of development

(Becker et al., 2011; Radke et al., 2014). Further, adolescent alcohol exposure has previously been shown

to render adult rats more susceptible to the effects of stressors, including footshock (Füllgrabe et al.,

2007; Siegmund et al., 2005). Thus, when considered alongside the literature, the current studies highlight

the variability inherent to acute stress effects on alcohol drinking behaviors and the importance of other

developmental experiences, be it stress or drug exposure, to adult outcomes. It is important to consider

that consumption of alcohol could be described as a stressor in and of itself. Alcohol exposure increased

activity in the hypothalamic-pituitary-adrenal (HPA) axis, and withdrawal from alcohol increased levels

of corticotropin-releasing factor (Olive et al., 2002; Zorrilla et al., 2001). The intermittent access drinking

procedure also requires animals to be individually housed, which is another significant source of stress

(Butler et al., 2016; Walker et al., 2019). Alcohol drinking in adolescence may consequently reveal the

effects of infant footshock on adult drinking by acting as a secondary stressor that potentiates the effects

of ELS. Further studies will be necessary to determine whether alcohol exposure alone, without the stress

of individual housing, can replicate the effects seen here and in our rat studies.

Our final experiment incorporated additional stressors beyond adolescent alcohol exposure. Mice

in this study were isolated from their dam and littermates following footshock and were fear-conditioned

during adulthood. Based on the results of Experiment 3, and our studies in rats, isolation from littermates

does not appear to be a necessary condition to produce effects on drinking, although it is unclear at

present whether isolation from the dam contributes to the ELS effects. Adult fear conditioning was not

incorporated in the drinking in the dark or continuous access experiments, which could possibly drive the

absence of a drinking effect in those studies. However, the lack of an effect in Experiment 3, which did

include adult fear conditioning, makes it more likely that adolescent drinking is the key factor. These

factors could be tested more systematically in the future to determine the number of stressful experiences

necessary to produce an effect on alcohol drinking behaviors.

It is important to consider why the effects of infant stress followed by adolescent alcohol drinking

were model-species dependent. There are several notable differences between Long-Evans rats and

C57BL/6J mice. Long-Evans rats are an outbred line, and thus in a given set of subjects there is increased

genetic diversity compared to the inbred C57BL/6J line. The effect of stress on alcohol consumption has

been found to vary between mouse strains (Radke et al., 2014). Furthermore, PTSD-AUD comorbidity is

not ubiquitous in individuals who have experienced trauma (Gilpin and Weiner, 2017; Young-Wolff et

al., 2011). Thus, the effect of early life trauma is likely dependent on genetic predispositions (Koenen et

al., 2009; Young-Wolff et al., 2011). Another notable difference between C57BL/6J mice and Long-

Evans rats is that C57BL/6J mice are bred to be high sucrose-preferring mice and are known to consume

more alcohol than other strains of mice. In contrast, Long-Evans rats have no predisposition to

maladaptive alcohol drinking. Thus, consumption may not be high enough in Long-Evans rats to observe

reductions in drinking like those seen in mice.

Our results show a clear effect of sex on alcohol consumption, with females consuming more

alcohol than males, which is consistent with a large body of previous literature (Finn, 2020; Radke et al.,

2021a, 2021b). Further, males and females displayed similar preference for alcohol vs. water in the

limited access and intermittent access experiments. This finding is consistent with previous studies of sex

differences in alcohol drinking behaviors in our lab that have found divergences in the effects of

biological sex on alcohol consumption vs. preference (Sneddon et al., 2022, 2019), and may give insight

into the origin of sex differences in alcohol drinking. Sex differences in alcohol preference have been

found to differ based on alcohol concentration (Radke et al., 2021b; Sneddon et al., 2022), but it is

unclear if the lack of an effect on preference seen here resulted from the different concentrations of

alcohol used in the three experiments or from the intermittent access vs. continuous access conditions. It

is also interesting that female rats are more sensitive to the effects of infant footshock on adult fear

behaviors (Minshall et al., 2024) than males and that shocked females demonstrated greater aversion-

resistance in the studies reported here. Thus, the model developed here may capture the increased

vulnerability for comorbid PTSD-AUD observed in female patients (Breslau et al., 1997; Gavranidou and

Rosner, 2003).

In summary, our study found no effect of infant footshock on alcohol drinking behaviors in mice

unless a period of adolescent alcohol drinking was included. A major goal of this line of work is to

establish a viable model of comorbid PTSD and AUD in male and female rodents. Infant footshock is

appealing for its relatively selective effects on PTSD-relevant behaviors, including fear learning and

anxiety behaviors (Poulos et al., 2014; Quinn et al., 2014) as well as resistance to fear extinction

(Sneddon et al., 2021). This model is also unique in that stress is administered prior to weaning and

before animals develop the ability to form associative memories, unlike other models in which stress

exposure occurs during adolescence (Shaw et al., 2020; Skelly et al., 2015). Importantly, the current

model has also been validated in male and female animals. It should be noted that contradictory results

and poor replicability are common in studies investigating the relationship between stress and alcohol

exposure. Thus, in order to understand the intricate differences in the relationship between stressors and

alcohol consumption, future studies should continue to investigate the specific conditions necessary to

increase alcohol drinking following ELS and include measurements of physical indicators of the stress

response such as corticosterone levels. Such continued development and refinement of mouse models of

comorbid PTSD and AUD will be beneficial to uncovering the shared neural mechanisms of these two

debilitating disorders.

Acknowledgements

The authors are grateful to Caroline Scribner, Brandon Arnold, and Annemarie Thomas for assistance

with behavioral experiments.

Author contributions: Thomas Perry: Formal Analysis, Investigation, Visualization, Writing – Original

Draft. Harrison Carvour: Formal Analysis, Investigation, Visualization, Writing - Original Draft. Amanda

Reichert: Formal Analysis, Investigation, Writing – Original Draft. Elizabeth Sneddon: Formal Analysis,

Investigation, Writing – Original Draft. Charlotte Roemer: Formal Analysis, Investigation. Ying Ying

Gao: Formal Analysis, Investigation. Kristen Schuh: Formal Analysis, Investigation. Natalie Shand:

Investigation, Writing – Original Draft. Jennifer Quinn: Conceptualization, Funding Acquisition,

Supervision, Writing - Review and Editing. Anna Radke: Conceptualization, Funding Acquisition,

Supervision, Visualization, Writing - Review and Editing.

References

Bangasser DA, Valentino RJ (2014) Sex differences in stress-related psychiatric disorders:

neurobiological perspectives. Front Neuroendocrinol 35:303–319.

Becker HC, Lopez MF, Doremus-Fitzwater TL (2011) Effects of stress on alcohol drinking: a review of

animal studies. Psychopharmacology 218:131–156.

Bertagna NB, Favoretto CA, Rodolpho BT, Palombo P, Yokoyama TS, Righi T, Loss CM, Leão RM,

Miguel TT, Cruz FC (2021) Maternal separation stress affects voluntary ethanol intake in a sex

dependent manner. Front Physiol 12:775404.

Breslau N, Davis GC, Andreski P, Peterson EL, Schultz LR (1997) Sex differences in posttraumatic stress

disorder. Arch Gen Psychiatry 54:1044–1048.

Butler TR, Karkhanis AN, Jones SR, Weiner JL (2016) Adolescent Social Isolation as a Model of

Heightened Vulnerability to Comorbid Alcoholism and Anxiety Disorders. Alcohol Clin Exp Res

40:1202–1214.

Carvour HM, Roemer CA, Underwood DP, Padilla ES, Sandoval O, Robertson M, Miller M, Parsadanyan

N, Perry TW, Radke AK (2024) Mu-opioid receptor knockout on Foxp2-expressing neurons reduces

aversion-resistant alcohol drinking. bioRxivorg 2023.11.29.569252.

Cruz FC, Quadros IM, Planeta C da S, Miczek KA (2008) Maternal separation stress in male mice: long-

term increases in alcohol intake. Psychopharmacology 201:459–468.

Enoch M-A (2011) The role of early life stress as a predictor for alcohol and drug dependence.

Psychopharmacology 214:17–31.

Finn DA (2020) The Endocrine System and Alcohol Drinking in Females. Alcohol Res 40:02.

Füllgrabe MW, Vengeliene V, Spanagel R (2007) Influence of age at drinking onset on the alcohol

deprivation effect and stress-induced drinking in female rats. Pharmacol Biochem Behav 86:320–

326.

Gavranidou M, Rosner R (2003) The weaker sex? Gender and post-traumatic stress disorder. Depress

Anxiety 17:130–139.

Gilpin NW, Weiner JL (2017) Neurobiology of comorbid post-traumatic stress disorder and alcohol-use

474   disorder. Genes Brain Behav 16:15–43.

475   Guinle MIB, Sinha R (2020) The Role of Stress, Trauma, and Negative Affect in Alcohol Misuse and

476   Alcohol Use Disorder in Women. Alcohol Res 40:05.

477   Hennessy MB, Kaiser S, Sachser N (2009) Social buffering of the stress response: diversity, mechanisms,

478   and functions. Front Neuroendocrinol 30:470–482.

479   Hopf FW, Chang S-J, Sparta DR, Bowers MS, Bonci A (2010) Motivation for alcohol becomes resistant

480   to quinine adulteration after 3 to 4 months of intermittent alcohol self-administration. Alcohol Clin

481   Exp Res 34:1565–1573.

482   Hopf FW, Lesscher HMB (2014) Rodent models for compulsive alcohol intake. Alcohol 48:253–264.

483   Hu P, Maita I, Phan ML, Gu E, Kwok C, Dieterich A, Gergues MM, Yohn CN, Wang Y, Zhou J-N, Qi X-

484   R, Swaab DF, Pang ZP, Lucassen PJ, Roepke TA, Samuels BA (2020) Early-life stress alters

485   affective behaviors in adult mice through persistent activation of CRH-BDNF signaling in the oval

486   bed nucleus of the stria terminalis. Transl Psychiatry 10:396.

487   Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB (1995) Posttraumatic stress disorder in the

488   National Comorbidity Survey. Arch Gen Psychiatry 52:1048–1060.

489   Koenen KC, Amstadter AB, Nugent NR (2009) Gene-environment interaction in posttraumatic stress

490   disorder: an update. J Trauma Stress 22:416–426.

491   Lee RS, Oswald LM, Wand GS (2018) Early life stress as a predictor of co-occurring alcohol use disorder

492   and post-traumatic stress disorder. Alcohol Res 39:147–159.

493   Lupien SJ, McEwen BS, Gunnar MR, Heim C (2009) Effects of stress throughout the lifespan on the

494   brain, behaviour and cognition. Nat Rev Neurosci 10:434–445.

495   Mayfield RD, Harris RA, Schuckit MA (2008) Genetic factors influencing alcohol dependence: Genetic

496   factors and alcohol dependence. Br J Pharmacol 154:275–287.

McCool BA, Chappell AM (2009) Early social isolation in male Long-Evans rats alters both appetitive

and consummatory behaviors expressed during operant ethanol self-administration. Alcohol Clin

Exp Res 33:273–282.

Minshall BL, Peguero AZ, Scheive KM, Wasylyshyn CF, Claflin DI, Quinn JJ. (in press). Acute

early life stress alters threat processing in adult rats. Behavioral Neuroscience.

Minshall BL, Skipper RA, Riddle CA, Wasylyshyn CF, Claflin DI, Quinn JJ (2024) Sex differences in

acute early life stress-enhanced fear learning in adult rats. Dev Psychobiol 66:e22511.

Olive MF, Koenig HN, Nannini MA, Hodge CW (2002) Elevated extracellular CRF levels in the bed

nucleus of the stria terminalis during ethanol withdrawal and reduction by subsequent ethanol intake.

Pharmacol Biochem Behav 72:213–220.

Pechtel P, Pizzagalli DA (2011) Effects of early life stress on cognitive and affective function: an

integrated review of human literature. Psychopharmacology 214:55–70.

Poulos AM, Reger M, Mehta N, Zhuravka I, Sterlace SS, Gannam C, Hovda DA, Giza CC, Fanselow MS

(2014) Amnesia for early life stress does not preclude the adult development of posttraumatic stress

disorder symptoms in rats. Biol Psychiatry 76:306–314.

Quigley JA, Logsdon MK, Turner CA, Gonzalez I, Leonardo N, Becker JB (2021) Sex differences in

vulnerability to addiction. Neuropharmacology 108491.

Quinn JJ, Skipper RA, Claflin DI (2014) Infant stress exposure produces persistent enhancement of fear

learning across development. Dev Psychobiol 56:1008–1016.

Radke AK, Held IT, Sneddon EA, Riddle CA, Quinn JJ (2020) Additive influences of acute early life

stress and sex on vulnerability for aversion-resistant alcohol drinking. Addict Biol 25:e12829.

Radke AK, Jury NJ, Kocharian A, Marcinkiewcz CA, Lowery-Gionta EG, Pleil KE, McElligott ZA,

McKlveen JM, Kash TL, Holmes A (2017) Chronic EtOH effects on putative measures of

compulsive behavior in mice. Addict Biol 22:423–434.

Radke AK, Pickens CL, Holmes A (2014) The effects of stress on measures of alcohol drinking in rodents

In: Neurobiology of Alcohol Dependence (Noronha AB, Cui C, Harris RA, Crabbe JC ed), pp 97–

110. Oxford: Academic Press.

Radke AK, Sneddon EA, Frasier RM, Hopf FW (2021a) Recent Perspectives on Sex Differences in

Compulsion-Like and Binge Alcohol Drinking. Int J Mol Sci 22:3788.

Radke AK, Sneddon EA, Monroe SC (2021b) Studying Sex Differences in Rodent Models of Addictive

Behavior. Curr Protoc 1:e119.

Shaw GA, Bent MAM, Council KR, Pais AC, Amstadter A, Wolstenholme JT, Miles MF, Neigh GN

(2020) Chronic repeated predatory stress induces resistance to quinine adulteration of ethanol in

male mice. Behav Brain Res 382:112500.

Siegmund S, Vengeliene V, Singer MV, Spanagel R (2005) Influence of age at drinking onset on long-

term ethanol self-administration with deprivation and stress phases. Alcohol Clin Exp Res 29:1139–

1145.

Sinha R (2001) How does stress increase risk of drug abuse and relapse? Psychopharmacology 158:343–

359.

Skelly MJ, Chappell AE, Carter E, Weiner JL (2015) Adolescent social isolation increases anxiety-like

behavior and ethanol intake and impairs fear extinction in adulthood: Possible role of disrupted

noradrenergic signaling. Neuropharmacology 97:149–159.

Smith NDL, Cottler LB (2018) The epidemiology of post-traumatic stress disorder and alcohol use

disorder. Alcohol Res 39:113–120.

Sneddon EA, Masters BM, Shi H, Radke AK (2023) Removal of the ovaries suppresses ethanol drinking

and promotes aversion-resistance in C57BL/6J female mice. Psychopharmacology 240:2607–2616.

Sneddon EA, Rasizer LN, Cavalco NG, Jaymes AH, Ostlie NJ, Minshall BL, Masters BM, Hughes MR,

Hrncir H, Arnold AP, Radke AK (2022) Gonadal hormones and sex chromosome complement

differentially contribute to ethanol intake, preference, and relapse-like behaviour in four core

genotypes mice. Addict Biol 27:e13222.

Sneddon EA, Riddle CA, Schuh KM, Quinn JJ, Radke AK (2021) Selective enhancement of fear learning

and resistance to extinction in a mouse model of acute early life trauma. Learn Mem 28:12–16.

Sneddon EA, White RD, Radke AK (2019) Sex Differences in Binge-Like and Aversion-Resistant

Alcohol Drinking in C57BL/6J Mice. Alcohol Clin Exp Res 43:243–249.

Spear LP (2016) Consequences of adolescent use of alcohol and other drugs: Studies using rodent

552   models. Neurosci Biobehav Rev 70:228–243.

553   Tawa EA, Hall SD, Lohoff FW (2016) Overview of the genetics of Alcohol Use Disorder. Alcohol

554   Alcohol 51:507–514.

555   Walker DM, Cunningham AM, Gregory JK, Nestler EJ (2019) Long-term behavioral effects of post-

556   weaning social isolation in males and females. Front Behav Neurosci 13:66.

557   Young-Wolff KC, Enoch M-A, Prescott CA (2011) The influence of gene-environment interactions on

558   alcohol consumption and alcohol use disorders: a comprehensive review. Clin Psychol Rev 31:800–

559   816.

560   Zorrilla EP, Valdez GR, Weiss F (2001) Changes in levels of regional CRF-like-immunoreactivity and

561   plasma corticosterone during protracted drug withdrawal in dependent rats. Psychopharmacology

562   (Berl) 158:374–381.

Figure Legends

Figure 1. Sex, but not infant footshock, influences ethanol consumption and preference in a

continuous access drinking task. (**A**) Mice (n = 10/group) drank 5% and 10% ethanol for five 24-h

sessions each. (**B**) Infant footshock did not impact ethanol consumption (g/kg). Females consumed

significantly more ethanol than males, **p < 0.001 main effect of sex. Female mice that received

footshock consumed more 5% ethanol than 10% ethanol, *p = 0.008 Holm-Sidak's test. (**C**) Infant

footshock did not impact preference for ethanol vs. water. Ethanol preference was higher in females vs.

males, **p = 0.002 main effect of sex. Ethanol preference was higher in female mice that received

footshock at the 5% vs. 10% ethanol concentration, *p = 0.047 Holm-Sidak's test.

Figure 2. Sex but not infant footshock influences ethanol consumption in a limited access drinking

task. (**A**) Mice (n = 13-14/group) drank 15% ethanol for ten 2-h sessions. (**B**) There was a significant

influence of sex on ethanol consumption (g/kg), *p < 0.001 main effect of sex. (**C**) Neither sex nor stress

altered preference for ethanol vs. water.

Figure 3. Sex but not infant footshock influences ethanol consumption in an intermittent access

drinking task. (**A**) Mice n = 10-21/group) drank 20% ethanol for twelve 24-h sessions. Quinine was

added to the ethanol solution on sessions 13-15 in increasing concentrations (10, 100, and 200 mg/L). (**B**)

Females consumed more ethanol than males, **p < 0.001 main effect of sex. Infant footshock did not

impact ethanol consumption. (**C**) Neither sex nor stress altered preference for ethanol vs. water. (**D**)

Females consumed more ethanol mixed with quinine than males, **p < 0.001 main effect of sex. Quinine

decreased ethanol consumption regardless of sex or footshock exposure, **p < 0.001 main effect of

quinine concentration. Consumption was reduced at the 100 and 200 mg/L concentrations, but not 10

mg/L, in all groups, * p < 0.05, **P < 0.01 Dunnett's test.

Figure 4. Infant footshock and sex do not influence ethanol consumption in adolescent mice in an

intermittent access drinking task. (**A**) Mice (n = 7-14/group) drank 20% ethanol for nine 24-h sessions.

(**B**) Infant footshock did not impact ethanol consumption at adolescence. (**C**) Infant footshock did not

impact ethanol preference at adolescence.

Figure 5. Infant footshock paired with adolescent alcohol exposure influences ethanol consumption

and aversion-resistant drinking. (**A**) Mice (n = 6-13/group) drank 20% ethanol for twelve 24-h

sessions. Quinine was added to the ethanol solution on sessions 13-15 in increasing concentrations (10,

100, and 200 mg/L). (**B**) 15 shock mice had significantly lower ethanol consumption compared to no

shock animals, $**p < 0.01$ main effect of shock. Female mice drank significantly more ethanol than their

male counterparts, $**p < 0.01$ main effect of sex. (**C**) 15 shock mice had significantly lower ethanol

preference compared to no shock animals, $*p < 0.05$ main effect of shock. (**D**) 15 shock mice had

significantly lower ethanol consumption compared to no shock animals, $**p < 0.01$ main effect of shock.

Female mice drank significantly more ethanol than their male counterparts, $**p < 0.01$ main effect of sex.

Quinine decreased ethanol consumption regardless of sex or footshock exposure, $**p < 0.01$ main effect

of quinine concentration. Consumption was reduced at the 100 and 200 mg/L concentrations for all

groups, except for no shock females, who also decreased consumption compared to baseline at 10 mg/L,

$*p < 0.05$, $**p < 0.01$ Dunnett's test.

Tables

Sex*	ELS	Concentration (mg/L)**	Consumption (mL/kg)
Male	no shock	0	88.56 ± 10.81
		10	72.71 ± 13.79
		100	**40.88 ± 20.20**
		200	**3.95 ± 0.81**
	15 shock	0	68.10 ± 12.89
		10	61.63 ± 11.77
		100	**20.67 ± 15.26**
		200	**8.69 ± 4.50**
Female	no shock	0	127.22 ± 6.80
		10	97.33 ± 19.72
		100	**34.77 ± 19.68**
		200	**47.45 ± 19.33**
	15 shock	0	128.00 ± 9.59
		10	112.86 ± 28.20
		100	**43.80 ± 25.39**
		200	**44.55 ± 20.18**

Table 1. Consumption of quinine in water over 24 h. Sensitivity to quinine was not altered by sex or ELS exposure. * $p < 0.05$, ** $p < 0.01$ significant main effect. Values in bold typeface are significantly different from baseline ($p < 0.05$, Dunnett's test).

A. Continuous access two-bottle choice

5 sessions — *5% EtOH vs. H2O*

5 sessions — *10% EtOH vs. H2O*

B. Consumption (g/kg in 24h) vs. EtOH Concentration (5%, 10%), Male and Female, no shock / 15 shock. Male ** Female; *

C. Preference (%) vs. EtOH Concentration (5%, 10%), Male and Female, no shock / 15 shock. Male ** Female; *

A.

Limited access two-bottle choice "Drinking in the dark"

10 sessions

15% EtOH vs. H2O

A.

Intermittent access two-bottle choice

12 sessions | 3 sessions

20% EtOH vs. H2O | 20% EtOH + quinine vs. H2O

B.

Consumption (g/kg in 24 h)

Male | Female
Sex****

no shock
15 shock

C.

Preference (%)

Male | Female
Sex

no shock
15 shock

D.

Male ⊢——— ** ———⊣ Female

Consumption (g/kg in 24 h)

no shock
15 shock

Quinine Concentration (mg/L)**

A.

Intermittent access two-bottle choice

B.

Consumption (g/kg in 24 h)

- no shock
- 15 shock

Sex — Male, Female

C.

Preference (%)

- no shock
- 15 shock

Sex — Male, Female

A.

Intermittent access two-bottle choice

Adolescent intermittent access

12 sessions

20% EtOH vs. H2O

3 sessions

20% EtOH + quinine vs. H2O

B.

no shock
15 shock **

C.

no shock
15 shock **

D.