The modular biochemical reaction network structure of cellular translation

Bruno Cuevas-Zuviría [a,b] , Evrim Fer [a,c], Zachary R. Adam [a,d] and Betül Kaçar [a,*]

[a] Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA
[b] Centro de Biotecnología y Genómica de Plantas, Universidad Politécnica de Madrid, Madrid, Spain
[c] Microbiology Doctoral Training Program, University of Wisconsin-Madison, Madison, WI, USA
[d] Department of Geosciences, University of Wisconsin-Madison, Madison, WI, USA

*To whom correspondence should be addressed: bkacar@wisc.edu

Abstract

Translation is an essential attribute of all living cells. At the heart of cellular operation, it is a chemical information decoding process that begins with an input string of nucleotides and ends with the synthesis of a specific output string of peptides. The translation process is interconnected with gene expression, physiological regulation, transcription, and responses to signaling molecules, among other cellular functions. Foundational efforts have uncovered a wealth of knowledge about the mechanistic functions of and many interactions between components of translation, but the broader biochemical connections between translation, metabolism and polymer biosynthesis that enable translation to occur have not been comprehensively mapped. Here we present a multilayer graph of biochemical reactions describing the translation, polymer biosynthesis and metabolism networks of an *Escherichia coli* cell. Intriguingly, the compounds that compose these three layers are distinctly aggregated into three modes regardless of their layer categorization. Multimodal mass distributions are well-known in ecosystems, but this is the first such distribution reported at the biochemical level. The degree distributions of the translation and metabolic networks are each likely to be heavy-tailed, but the polymer biosynthesis network is not. A multimodal mass-degree distribution indicates that the translation and metabolism networks are each distinct, adaptive biochemical modules, and that the gaps between the modes reflect evolved responses to the functional use of metabolite, polypeptide and polynucleotide compounds. The chemical reaction network of cellular translation opens new avenues for exploring complex adaptive phenomena such as percolation and phase changes in biochemical contexts.

Introduction

Cellular life is a network of molecular components that are dynamic and adaptive. Life can only
exist as a gradient of free energy, and metabolic networks can account for the uptake and
excretion of substrates alongside the synthesis of key biomolecules[1]. Metabolic networks trace
the routes that energy travels through cells, but genetic sequences and their associated
enzymatic polymers[2–4] specify which pathways will be constructed, governed, and maintained in
any given cell. While a great deal of current research is focused on filling specific knowledge gaps
at the biochemical level, studies of the overall cellular network can provide insights into how the
details of biochemistry lead to the emergence of life's foundational properties.

At the cellular level, analysis of metabolic networks has inspired decades of research into
biochemical complexity[5–9]. These analyses have drawn connections between network attributes
such as a heavy-tailed degree distributions and general complex behaviors such as resilience,
adaptability, and modularity. Following initial studies of metabolic networks, there have been
numerous descriptions of heavy-tailed networks at other biological levels such as ecological
interactions, including theories of generative mechanisms that are relevant to characterizing the
prebiotic world[10–14]. These generative mechanisms lead to continuous, heavy-tailed distributions
consistent with a power law as an asymptotic sampling limit is approached; discontinuous or
multimodal distributions are not explicitly prescribed by these physical models.

While the cellular level is usually the fundamental starting point to describe life properties, life
manifests at different nested scales (*e.g.*, cell, organism, ecosystem). At the largest biological
levels of populations and ecologies, systems show emergent traits as *multimodality* or even
discontinuities in their properties. One of the most studied properties showing discontinuities is
size: depending on the ecosystem, organisms spanning certain size ranges will not exist. Examples
are found in arboreal forest[15], bird[16], fish and plankton[17], and mammal[18] groupings that can
exhibit multimodal or discontinuous size distributions. Ecological multimodal aggregations arise
from a confluence of dynamic processes consisting of an overlapping combination of both
positive feedback (*e.g.*, overabundant energy resources available for consumption at one size or
temporal scale but not at others) and negative feedback (*e.g.*, predation pressures such as grazing
that can make survival at specific scales extremely challenging, or few and highly variable
available resources)[19]. Discontinuities and modalities in biology also appear to be idiosyncratic;
gaps or troughs in one ecosystem do not necessarily map to those found in others, and persistent
gaps can nevertheless vary and shift over time[20]. It is not clear whether these network-level
attributes of biology ought to extend downward to the biomolecular scale.

Considering these scale-dependent aspects of biological networks, a compelling possibility is
which (if any) topological features associated with complex high-level biological systems can be
found at the cellular level. We hypothesized that these features, if they show up within a cellular
system, necessitate an explicit description of the processes that shape chemical hierarchical
structure. In ecosystems, these processes include mapping trophic relationships and sources of
energy input. Therefore, we address this question at the cellular level by building a translation
reaction network in *Escherichia coli (E. coli)*, which we have combined with its metabolic and
biosynthetic networks (**Figure 1**). Translation is a molecular information decoding process that
begins with a ribonucleotide sequence as input and a peptide sequence as output, with
remarkable robustness, accuracy and adaptability[21–27]. The processes that allow translation to
occur are carried out through the coordinated actions of numerous ribonucleic and protein
polymers, which we refer to as the translation machinery (TM). The decoding process that
converts triplets of RNA monomers into the attachment of a specific amino acid residue to a
growing peptide sequence is remarkable in that it is the earliest (and possibly only) emergent
system that processes discretized information and manifests semantic relationships entirely in
chemical terms. The chemosynthetic nature of the TM (*i.e.*, information processing via chemical
reactions, as compared to electronic signals in the brain or in digital computers) enables its
description as a chemical reaction network, which is amenable to systems-level analyses as has
been done for metabolic networks[5,28].
91
Here, we present the translation reaction network that includes the core processes of translation
(initiation, elongation, termination), detailed assembly steps required to replicate the ribosome,
and other well-documented features of the TM that maintain translation functionality (**Figure
2A**). To connect the translation and metabolic networks, we constructed a polymer biosynthesis
network that explicitly links enzymatic components catalyzing the metabolic reactions with the
mRNA reading and peptide synthesis steps of translation (**Figure 3A**). Comprehensive mapping
of the connections between translation and metabolism is necessary to investigate context-
dependent feedbacks between the cellular environment, substrate/nutrient availability, and
ribosomal protein activity.
101

Methods

Generation of the translation and polymer biosynthesis reaction databases

The purposes of this study are (i) to describe translation as a chemosynthetic process, (ii) to
integrate this process with a chemical reaction network describing metabolism; (iii) to map
network-level patterns between object mass and object connectivity across different chemical
domains; and (iv) to assess whether network-level patterns observed at the highest levels of the
biotic hierarchy are present at its lowest levels of biochemistry. These purposes circumscribe the

scope of the resulting network's component description. Objects and reactions that describe the
chemical components of the cellular translation and metabolic systems will be transcribed and
included in the network. Those reactions that do not alter the explicit biochemical components
of an *E. coli* cell (*i.e.,* gene regulatory interactions, sequence-specific motifs and their mutual
interactions, isomerization reactions, *etc*.) may be otherwise important, but they fall outside the
scope of this particular chemosynthetic study and will be excluded. To be sufficient for inclusion
in the network, an object and its associated reactions must have been reported in peer-reviewed
literature and be based on observable, experimental data; inferences based only on modeling
alone are not sufficient.

To serve this purpose, the level of description for translation includes a chemical accounting of
ribosome assembly and replication, as well as adjacent active site (A, P and E) interactions, since
each of these interactions has downstream effects on the synthesis of biopolymers that enable
autocatalytic replication of the entire system (**Figure 2B**). The chemical reaction network is
intended to focus on the cellular chemosynthetic process, and does not explicitly include
sequence-specific or site-specific (physiological or anatomical) information of the cell.

A list of *E. coli* translation components was obtained from the EcoCyc database[29] (GO: 006412).
Supplementary File 1 summarizes the components of translation used in this network, apart
from the ions and metabolites (*e.g.* GTPs) required to operate. The reactions for tRNA
aminoacylation by tRNA synthetases, rRNA maturation and modifications, ribosomal protein
modifications and maturations and peptidyl tRNA hydrolysis were obtained from the EcoCyc[29]
database and were also confirmed by literature synthesis (**Supplementary File 1**). The reactions
for ribosomal subunit assembly, initiation, elongation and termination steps and stress response
were built for *E. coli* (**Supplementary File 2**). The generation of translation reactions focused on
five processes: a) elongation, which is largely conserved across all organisms; b) initiation, where
some variations have been described between and within domains of life; c) termination; d)
translation machinery biosynthesis, which are the assembly steps for replicating a ribosome; and
e) the response to amino acid starvation. The reactions were manually annotated. To describe
the elongation process in detail, the reaction model accounts for each of the three ribosome
active sites. This includes a large number of reactions describing each of the distinct combinations
of amino acids at each position. The metabolic and translation component masses were retrieved
from the Kyoto Encyclopedia of Genes and Genomes (KEGG) Application Programming Interface
(API). The TM layer reaction network file is attached as **Supplementary File 3**.

The polymer biosynthesis layer was constructed to link the metabolic and translation layers
together by writing non-stoichiometric reactions that approximate the assembly of amino acids

into the assembled enzymatic polypeptides. Each reaction within the metabolic layer associated
with a specific enzyme resulted in a new reaction placed within the polymer biosynthesis layer
network. For reactions that involved multiple protein components (*i.e.*, the formation of
multimer heterocomplexes), the logical relationships of the protein components that enabled
specific reactions were automatically read and parsed to generate protein-protein interaction
maps within the molecular biosynthesis layer. The polymer biosynthesis layer is attached as
Supplementary File 6.

The translation process is carried out as a repetition of steps for each of the proteins. Given that
we are not considering sequence-specific processes or relationships, we must employ a
placeholder "peptide" node that is used in chemosynthesis steps carried out by translation prior
to completion of enzyme synthesis. It denotes any incomplete polypeptide sequence that is in
the process of being assembled through the translation process. It does not possess any specific
chemical function or identity that must be subsequently tracked in other portions of the chemical
reaction network.

Metabolic reaction database
The metabolic layer was obtained from Feist *et al.*[30], who reconstructed *E. coli* metabolism
(**Supplementary File 4**). We linked these reactions to their corresponding enzyme(s). We
obtained chemical information about each of these enzymes through KEGG. The metabolic layer
is attached as **Supplementary File 5.** We also considered metabolic models of alternative strains
of *E. coli* obtained from BiGG[31]. The subtle differences between the strains do not lead to any
significant difference in the degree distributions (see **Figure S1**), so we only consider one strain
in this article.

Chemical reaction network integration
All chemical reactions from each of the three layers (translation, polymer biosynthesis and
metabolism) were uniformly processed into an aggregate network data file for graphing and
analysis (**Figures 1 and 3A**). The aggregate network data file forms a bipartite graph with one
class of nodes representing chemical compounds, and another class of nodes representing logical
chemical relationships (*i.e.*, chemosynthesis reactions and biosynthesis assembly processes) that
describe generating biomolecules from underlying substrates. The integrated network is
composed of 1433 reactions for translation, 1455 reactions for polymer biosynthesis and 2085
reactions for cellular metabolism, each with varying proportions of intra- and internetwork
connectivity. The complete network is provided in **Supplementary File 7.**

Network visualization, processing, and analyses

We define the degree of a compound based on the number of different reactions involving it
(whether as reactant, product, transporter or catalyst). We have built upon prior network
analysis efforts[5] to explicitly consider enzymes and macromolecular assemblies in addition to
metabolites. The complementary cumulative distribution function (CCDF), a binned list of the
number of compounds with a degree value in a given network, was analyzed using the power-
law Python library[32] to determine the likeliest statistical model that fits the distribution. We
performed discrete and continuous fits using models of exponential and power-law distributions,
and we compared the log-likelihood ratios among them to determine if a given degree
distribution is likelier to be homogeneous or heterogeneous, respectively[33].
193
Network visualizations with many pieces of information tend to become cluttered, which can
obscure observational insights. Techniques that reduce a network's information density without
altering its fundamental relationships are useful for recovering these insights. We simplified
network visualizations by grouping sets of reactions into combined nodes (*node-contracting*)
according to their *module* labels. These labels were either provided by the original metabolic data
without further modification[30](*e.g.,* transport, pyruvate synthesis, *etc.*), or manually assigned by
the authors for the translation network (*e.g.,* elongation, initiation, *etc.*). The term 'module'
denotes a subset of nodes in a network that are densely connected to each other, while being
sparsely connected or entirely disconnected from nodes in other modules[34,35]. Compounds that
participate only in reactions within a single module are grouped within combined nodes with
their module labels, while those compounds that interact across different modules are conserved
and graphed as individual nodes. The resulting contracted network is still a bipartite network
with modules and reactions connected by compounds. We contracted a network formed by the
translation and metabolism layers, and we perform an analysis to find out which nodes are more
relevant in the traversal of this contracted network using the betweenness centrality metric. This
metric consists of measuring the number of times that a node appears in the shortest path
between two different nodes[36]:
211

212
$$BC(n) = \sum_{s \neq n \neq t} \frac{\sigma_{st}(n)}{\sigma_{st}}$$

213
214 Where σ_{st} is the number of paths joining the nodes s and t, and $\sigma_{st}(n)$ is the number of those
215 paths that pass through the node n. Reaction directionality is accounted for when using this
216 method. A different contraction approach is taken in Figure 3.B, where we collapse all the
217 reactions belonging to the same layer, and we represent in the edges the number of compounds
218 that participate within each layer or that connect the different layers.
219

220 The KEGG database provided the molecular structures and sequences required to compute the
221 molecular weight of each of the chemical compounds (proteins, RNA, metabolites, *etc.*). In the
222 case of complexes, we add the masses of their components but we disregard their stoichiometry
223 of assembly due to a lack of experimental information.
224
Large networks are highly dimensional objects where similarity relationships are difficult to
assess. Network embeddings allow the mathematical representation of network entities to ease
their comparison[37]. The network embeddings are generated using the node2vec algorithm[38] as
implemented in PyTorch Geometric[39]. We use standard parameters to generate the embeddings.
The resulting 128 dimensions of each of these embeddings were converted into 2D embeddings
by using the t-distributed Stochastic Neighborhood Embedding (t-SNE) technique. Visualization
of the integrated network and its individual layers was carried out in Gephi 0.9.2[40]. The code
employed to perform this analysis together with the data are provided in the article GitHub
repository https://github.com/kacarlab/Translation-Network.
234

Results

236
Translation is a process described in textbooks in great detail, but there are still many unknowns
about its function and modes of integration with other cellular systems. By reviewing the
literature, we compiled the reactions of translation and generated a bipartite graph represented
in **Figure 2B**. This TM network includes the core of translation processes (initiation, elongation
and termination) together with the assembly and recycling of the ribosome, and the response to
amino acid starvation. A feature of this network is the high level of redundancy of some of the
reactions, given that translation is based on parallel processes involving different tRNAs.
244
All chemical reactions for translation and polymer biosynthesis were then integrated with that
for metabolism. The chemical object names in each network were screened for consistent
annotation, their molecular masses and object classifications were cross-referenced or inferred
(*i.e.*, proteins, RNAs, codons, amino acids, energy molecules, and relevant ions). A full map of the
integrated network, with node sizes weighted by node degree and reactions color-coded by
network layer category is displayed in **Figure 3A**. The overall network contains more than 8000
nodes and 24000 links. When nodes are placed using a graph-layout algorithm (Force-Atlas 2 in
this case), each layer clusters as a separated group of nodes that are mostly distinct from one
another. A breakdown of the numbers of objects connected within and between the different
layers is displayed through the use of numbered arrows in **Figure 3B**.
255

To understand how their components are connected within and outside each network, we
applied Node2Vec non-supervised learning to generate multi-dimensional representations
(embeddings) as vectors for each node in the network that reflect its neighborhood of
connections. We then reduced these multi-dimensional vectors to two dimensions using a t-
Distributed Stochastic Neighbor Embedding (t-SNE) dimensionality reduction technique. The
resulting embeddings provide a representation of which nodes share a common neighborhood
by placing them closer in a unitless 2D plane. The translation and the metabolic layers represent
two different domains that can be easily distinguished (**Figure S2**), while the biosynthesis nodes
appear scattered across these two different domains without any distinct clustering. Within the
translation network, we find two large clusters of nodes and many smaller groups. Within these
smaller groups there are nodes representing each of the amino acid processing cycles and the
tRNA-synthetase process. In contrast, we do not find any cluster in the metabolic network
containing distinct components of the TM.

Despite the large number of new connections mapped across the integrated network, the
sparsity between translation and metabolic layers may be explained in part by the relatively few
compounds joining them directly together (25 and 27 to and from each layer, **Figure 3B**), which
is limited to the molecules providing peptide bond formation energy and substrate molecules for
RNA polymers. It seems likely that the reactions that form connections across the metabolism
and translation layers might have a central role keeping the overall network together. We
contracted reactions and compounds involved in identified modules (*e.g.,* cofactor synthesis,
elongation, *etc.*), and color-coded the resulting network using the betweenness centrality for
each node of the contracted network in **Figure S3**. In this case, this metric represents how often
a node is visited when traversing across the different layers of the integrated translation-
metabolic network. The network depiction of these data shows a distinct separation between
the translation and metabolism layers, with connections across the separation maintained
through modules such as tRNA charging providing the raw materials for polypeptide elongation.
Together with transport modules that provide the bulk of metabolic precursors for heterotrophic
E. coli metabolism, tRNA charging is one of the most central modules of the cell.

An important attribute of large networks is their degree distribution: the probability of finding
nodes with a given number of connections. In bipartite reaction networks, such as the integrated
network assembled here, the degree of each compound represents the number of reactions in
which it participates. Networks associated with complex systems usually show heterogeneous
degree distributions, where a few nodes show a remarkable number of connections, while
random networks are characterized by having more homogeneous degree distributions peaking
around a mean number of connections[14]. We use the log-ratios of different statistical fits to
determine the most likely degree distribution (exponential for homogeneous networks, power-

law for heterogeneous networks). For objects that compose the metabolism and translation
layers, their degree distributions follow statistically probable heavy-tailed distributions (**Figure**
4B, **Table 1**), which is not the case of the polymer biosynthesis layer degree distribution (p>0.05).
The only object in the polymer biosynthesis network that is heavily connected to other
compounds is a generic node 'peptide' (**Figure 3A**), which represents intermediate peptide
sequences that for various reasons have not completed the termination step, and thus have no
distinct chemosynthetic or functional impact on the network. The overall degree distribution of
the full network follows a heavy-tailed degree distribution with an exponent around 2.0 (**Figure**
4A, Table 1).

Table 1. A tabulated comparison of power law and exponential distributions (log-likelihood
ratios), significance estimates (p-value), minimal degree value at which we start fitting the
distribution (k_{min}), and best-fit exponent values (γ) for the complementary cumulative distribution
functions of the translation, biosynthesis and metabolism networks. See Clauset *et al.*[33]. For a
more detailed discussion of the fitting methods.

309
310

Domain	Fit type					
	Continuous			Discrete		
	γ	k_{min}	p-value	γ	k_{min}	p-value
All layers	2.118	3	$2.4 \cdot 10^{-17}$	1.929	3	$4.5 \cdot 10^{-13}$
Biosynthesis	3.470	3	$9.9 \cdot 10^{-2}$	2.703	3	$9.1 \cdot 10^{-1}$
Metabolism	2.290	6	$9.0 \cdot 10^{-8}$	2.229	3	$3.0 \cdot 10^{-7}$
Translation	3.526	40	$3.2 \cdot 10^{-14}$	3.448	40	$1.2 \cdot 10^{-13}$

311
312
While the degree of a compound might reflect its relevance in shaping the chemical network, its
mass correlates with the amount of energy required for its synthesis; larger molecules require
more energy to synthesize than smaller ones[41]. Our objective is to assess the relationships, if any,
between the connection degrees and the masses of the compounds in the network. A multimodal
distribution can be discerned in the panels of **Figure 4C** (categorized by chemical compound type)
and **Figure 4D** (categorized by network layer). Mass distribution concentrations (modes) across
all network layers occur at around 240, 40,000 and 2,000,000 Daltons (Da). The central regions
of these modes contain both the highest concentrations of compounds and the most highly
connected compounds from the different layers. There are bins in troughs between the modes
without reported molecules, which indicates that the modal mass distribution may also be
discontinuous. All modes contain some objects from the TM network; enzymatic polymers that
enable metabolism are almost entirely confined to the middle mode; and metabolic compounds
compose the two smallest modes. The structural composition of the chemical compounds (small
molecules, polypeptides, and polynucleotides) is the likeliest determinant of the mode regardless

of the functions they perform or the layer they are assigned to within this network (**Figures 4C**
and 4D).
329

Discussion

Bioinformatic databases are constructed and annotated in such a way as to enable the high-
throughput reconstruction of many biosynthetic steps of metabolism and gene regulation[42], and
also to reconcile these steps with sequences found within an organism's genome in an organized
pipeline (gene → protein → function[43]). Prior network descriptions and analyses of translation
have focused on its dynamic attributes afforded by its autocatalytic motifs[44], genetic regulation
of metabolic processes[45], the links between TM function and organism-level cellular dynamics[46]
and a mapping of genotype to phenotype useful for interpreting a wealth of -omics cellular
data[47]. The chemosynthetic function of the TM and the topological attributes of the overall
network did not figure significantly into these prior efforts.
340

The integration of translation and metabolism reactions for *E. coli* describes a naturally occurring
chemical network that is hierarchical and spans at least six orders of magnitude of compound
mass. Hierarchy is most clearly resolved when degree per compound is plotted against compound
mass (**Figure 4C and 4D, upper panels**). By inference, the energy costs associated with
synthesizing and replicating the largest TM components outstrip those of enzymes and
metabolites by at least 1-2 orders of magnitude. The energy cost per molecular component
demonstrates the hierarchical importance of the TM in specifying chemical functions made
possible by biopolymers within the cell.
349

As cellular operation requires a hierarchy of different mass entities, it also involves modules with
different connectivity patterns. The embedding analysis depicted in **Figure S3** reveals that the
metabolic and polymer biosynthesis reactions do not cluster apart from each other in any
significant way, but the TM clusters in a distinct region of the graph. The metabolic and TM layers
exhibit degree distributions that are likely heterogeneous, but the biopolymer layer does not.
Taken together, this would indicate that protein biosynthesis is mostly shaped by the same
structural constraints of the intracellular metabolism module and is not likely to exhibit
independent (*i.e.*, intra-layer) attributes associated with complexity.
358

359 **Continuous, multimodal or discontinuous distributions**
360

Based on prior observations of biological networks across the hierarchy of life, we articulated
three distinct possibilities for the distribution of compounds in the integrated translation network

(see **Figure S4**). Each of these possibilities has distinct, predicted implications that may be
compared to the observed mass distribution. In a parallel to 'trophic hierarchy', one hypothesized
driver of hierarchical connection in cellular biochemistry is the ability to distribute energy and
maximize power in a dynamic way across independently functioning layers. For this possibility,
each distinct molecular layer is self-organizing and both feeds upon and into the others. The
predicted pattern is that each layer contains distinct connectivity peaks, and the overall mass
distribution is continuous. In an 'energy availability' distribution, the hierarchical structures of
polymer biosynthesis and translation are built directly atop, and extensions of, energy pathways
enabled by metabolism. The predicted pattern is that though the translation and polymer
biosynthesis layers of the network may contain 'heavy tails', the bulk of compounds in each layer
all coincide with a peak generated by metabolic compounds. In a distribution shaped by
'molecular distinction', the association of compounds with a given biofunctional layer is less
significant than the compounds' chemical attributes in the cellular environment. The predicted
pattern for this distribution is that multimodality or discontinuity caused by distinct molecular
properties would lead to sorting at the network level. Each of these predicted distributions may
carry basic implications about the primordial relationship(s) between emergent metabolic,
enzymatic and translation functions, and about identifying measurable network attributes that
fundamentally distinguish abiotic and biotic systems. We predicted that energy availability would
be the dominant organizing driver of cellular biochemistry, given that energy transduction is
indispensable to the overall function of translation.

The observed multimodal (and perhaps discontinuous) mass-degree distribution of the network
seems most consistent with the 'distinct molecules' prediction. The network's multimodal mass-
degree distribution is the first such distribution reported for biological systems at the cellular
level. The multimodal mass distribution of the integrated chemical reaction network is
independently supported by chemical assays of living *E. coli* cells. Non-targeted metabolomic
assays of small molecular mass compounds using mass spectrometry (TOF-MS) and different
preparatory analytic weighting techniques consistently show a peak in metabolite compounds at
approximately 150-350 Da, and a tapering of compounds approaching ~2000 Da, even after
accounting for decreased detector sensitivity with increasing mass[48,49]. This approximately aligns
with the centroid and distribution range of the smallest mass mode. For polypeptides, proteomic
assays conducted across taxa as diverse as *E. coli, S. cerevisiae* and *H. sapiens* consistently show
a broad correlation between translated genomic length and *in vivo* enzyme length[50]. For *E. coli*,
this specifically includes a median peak length of about 209 amino acids, which corresponds to a
peak in the mass distribution of polypeptides of approximately 28,000 Da if the average amino
acid mass is assumed to be 136 Da (the average molar mass of all 20 essential amino acids). This
approximately aligns with the centroid of the second mass mode. A whole-cell assay of all *E. coli*
enzymes, regardless of composition and function, show that highly connected objects central to

both metabolism and translation (*e.g.*, the ribosome, TCA cycle components, glycolysis and
fermentation components, *etc.*) compose comparable protein mass fractions under a variety of
growing conditions[51]. The ribosome itself is consistently composed of 2:1 mass fractions of RNA
to protein[52]. In sum, independent raw cell molecular assays indicate sharply discretized
compound mass modes that are attributable to the chemical properties of RNA polymers,
proteinaceous polypeptides, and base metabolites, regardless of their cellular functions they
carry out or the modules to which they may be assigned. Molecules that fall within the troughs
between the modes are undoubtedly present in cells in some form, but would not seem to play
essential roles in generating an *E. coli*'s chemosynthetic pathways, cellular physiology, or
adaptive responses to stimuli. Multimodal distributions present a unique challenge to theoretical
studies of network generation, and may require elaboration using multiplexes that can account
for multimodality[53].
413
414 **Causes and evolutionary implications of biochemical modularity**
415
An assessment of the possible drivers of multimodality and discontinuity in the chemical reaction
networks of bacteria may open new ways of characterizing the evolvability of cellular systems,
and for distinguishing them from complex abiotic systems. Different network analysis techniques
(**Figures 3A, S2 and S3**) and raw plots of mass distribution modes (**Figures 4C and 4D**) trace
modularized systems of translation and metabolism that, by virtue of their chemical isolation
from one another, are each able to independently adapt to conditions that the cell may
encounter[48]. Modules can exhibit hierarchical properties and function as key units of adaptive
evolution because organismal fitness depends on their performance[7,54–57].
424
The *E. coli* metabolic network is only able to exist via the functions afforded by polymeric
peptides. Accounting for these polymeric peptides in the chemical reaction network introduces
the first modal trough (**Figure 4C**) between the largest metabolites and the smallest polypeptides.
This trough is also present in objects that compose the translation network that are found across
this range of masses (**Figure 4D**). From a purely physical perspective, there are no *a priori* reasons
for a discontinuity of mass size between these compounds. A discontinuity may be
understandable, though, in biofunctional terms if resources for synthesis are limited[41]. The
synthesis of an extremely large metabolite necessitates the precise placement of assorted atoms
and molecular groups, connected with different bond types. If there is a single placement error,
or inadequate feedstock of starting materials, the resulting compound is unlikely to be functional.
An expenditure of chemical resources to produce increasingly larger (and increasingly error-
prone) metabolites has adverse effects on the persistence of the overall cellular system. The
synthesis of a linear oligopeptide of comparable mass, by contrast, can make use of repetitive
interconnections of a much smaller number of object types (*i.e.*, amino acids), many of which

also have similar chemical properties. A reduction of the number of building block types and of
the kinds of bonds through which they are connected can increase production fidelity (and
resultant functionality) of peptidic complexes compared to massive metabolites. The polypeptide
assembly process cannot scale downward indefinitely, however, since there are natural limits to
how small a chemically functional polypeptide may be. Oligopeptides must attain some minimal
length to form structural folds, or to form a pocket in which specialized cofactor chemistry can
occur that is isolated from the cellular environment[57]. The tradeoffs between synthesizing
extremely large metabolites or extremely small polypeptides to ensure overall functionality can
introduce a natural trough in mass distribution between these two chemical regimes that is
attributable to both the constraints brought about by the chemical properties of their different
building blocks and of the emergent functions they enable.
450
The trough between the second (polypeptide) and third (polynucleotide assemblies) modes
appears to arise for entirely different reasons that are distinct to the TM. As with the first mode
gap, there are no obvious physical or chemical reasons why there should not be an overlap
between large polypeptide and small ribonucleotide polymer compounds, and the observed
trough is counter to our predictions. The ribosome, however, has evolved to localize many of the
TM's protein-protein, protein-ribozyme, and ribosomal interactions on itself, thereby spatially
isolating the functions of translation from the surrounding metabolic and physiological activities
in the cell[55]. These functions are further isolated in time and space by regulatory mechanisms,
such as via binding with the properly formed initiation factors[58–61], by corresponding activated
tRNA complexes[62] and through spatial heterogeneity arising from complex physiological
relationships with nucleoids[63]. The TM network description includes a rich array of intraribosomal
subunit interactions, which we record as interactions involving an object with the mass of the
ribosome. This creates a sharp, isolated peak in the distribution centered at the ribosome's mass.
The ready clustering of the TM in the network topology (**Figures 3A**) and embedded vector and
dimensionality reduction analysis (**Figure S2**) depict and corroborate the biochemical facets of
translation's modularity, as do *in vitro* experimental systems that employ cell-free protein
synthesis[64,65].
468
There are numerous examples of systems that may be characterized as complex self-organizing,
adaptive phenomena, but which lack the evolvability, emergent novelty and agency of living
systems. Far-from-equilibrium physical systems such as galactic, stellar, and planetary
structures[66], earthquakes[67], solar flares [68] and sandpile or avalanche dynamics[69] all exhibit
continuous scaling across many orders of magnitude, but few or no reported discontinuities.
Discrete size classes observed in biology, on the other hand, are evidence of multiple dynamic
regimes[70] in which signaling, and the perturbative effects of novelty can propagate and be
controlled in directions both up and down scales of a hierarchy[71].

A multimodal biochemical distribution may be interpreted as indicating an evolved response of
modularization to significant constraints placed on a complex adaptive chemical reaction
network[34]. In biochemical systems, energy flows through the entire hierarchy mostly in the
direction from metabolites up to the translation components, but modal gaps and troughs arise
due to a complicated reconciliation process between the basic building blocks that compose large
objects in the system, energy availability for synthesis, and optimization of functional traits that
large objects can exhibit[72]. In conjunction with a heavy-tailed degree distribution, a multimodal
mass distribution may be one of a select suite of indicators diagnostic of biological dynamics that
can be observed across the entirety of life's various spatiotemporal scales.

Implications for prebiotic chemistry

Mapping the full extent of chemosynthetic relationships between translation and metabolism
(**Figures 3A** and **3B**) graphically captures long-recognized aspects of the hierarchical relationship
between these two cellular functions. Translation of a transcribed mRNA directs the specificity
of polymer-catalyzed metabolic chemistry, and this chemistry in turn supplies the required
energy and materials for ribosomal replication and operation. At the same time, though, the
existence of multiple distinct mass discontinuities that correlate more with differences in
chemical composition (**Figure 4D**) than with cellular function (**Figure 4C**) opens new possibilities
about the underlying dynamic circumstances in which translation arose. Chemical modes and
discontinuities may be so fundamental as to have been contemporaneous with (or have
preceded) the origins of cellular life. As an extension, it is possible that some information-
processing attributes of the TM module are primordially rooted in complex feedbacks between
ribonucleotides and metabolites, intra-network dynamics of the TM, or physicochemical
reinforcement of modularization (such as spatiotemporal isolation) afforded by discrete
compound class modes, as much as from the emergence of the ribosome's large and small
subunit mechanical activities.

Generating hierarchically connected networks across so many orders of magnitude, with
modules that each contain at least one autocatalytic cycle[44,73–75] presents a significant challenge
to the field of prebiotic chemistry[76–78]. If the biochemical modality and modularity observed in
this network are genuinely universal attributes of life, they may arise due to general physical
phenomena such as percolation[79]. The observed coupling between translation, protein
biosynthesis and metabolic networks presents a new model system in which the network
phenomenon of percolation may be studied in a chemical context[80] as an exemplary emergent
system in which multiple networks with distinct topologies coexist and are connected via
common elements[81] and discontinuities stem from mechanisms that lead to network

(dis)assortativity[82,83]. Theoretical analyses of such conditions show a possible connection
between the number of distinct modes and the possibility of multiple discontinuous phase
transitions enabled by multiplex networks[84]. Phase transitions have long been recognized as
potentially important behavioral features that are correlated with complex adaptive systems[85].

The relationships between complex adaptive systems and information processing attributes
afforded by network topology are directly relevant to developing new theories of the origin of
translation. The decoding process that links nucleotide sequence identity to peptide sequence
functionality is highly optimized for error tolerance through a property known as code
degeneracy, which can be described in informatic and computational terms without explicit
reference to the underlying biochemistry. It is unclear how an inherently informatic, network-
level property such as code degeneracy would arise on the basis of tracing the individual origins
of the ribosome's chemical components and interaction partners in isolation from this wider
network[86]. Recent discoveries have shown that 'biological' attributes such as self-reproduction,
adaptation and evolvability (which together characterize much of biological agency) are
intrinsically linked to one another through coupling of Hopf and Turing instabilities within a small
network of reactions with a cubic autocatalytic motif[87]. These attributes are observable even in
the absence of the intricacies of macromolecular mechanisms or a diversified biochemical
metabolic network. Further studies of coupled metabolic and translation chemical reaction
network properties may help to develop new hypotheses about the origin of translation as an
information-processing modular network, rather than about how translation can arise due to the
chemical relationships between amino acids and nucleotides or through the specific biochemical
activities of TM components. Specifically, one may test whether the modularization of
information processing centered on the ribosome in modern cells can be decentralized across a
primordial TM network.

Conclusions

Life exists, in part, by cohesively channeling energy from abundant, small-mass objects into the
synthesis of a specific array of large-mass objects. These larger objects afford functions and
activities that smaller objects, for various reasons, cannot perform. The inherent tension
between utilizing available energy in an efficient way and ensuring diverse functions and
activities will be reliably carried out can lead to multimodal mass distributions. Multimodality in
this sense may be diagnostically biological, as an evolved trait that is exhibited by adaptive, self-
organizing and self-assembling populations that can exhibit and propagate novel responses to
stimuli and which are subject to natural selection. It is a property that has been observed in many
different ecological systems. Our study of a combined network of metabolism and translation
shows that mass multimodality can arise in biochemistry, with distinct drivers for different

troughs between modes, even before physiological or gene regulatory mechanisms of cellular
operation are considered. This extends multimodality down to the lowest levels at which
biological population dynamics can exist: the chemistry within individual bacterial cells.
Translation stands apart in the cell as a functional module that is as much informatic as it is
chemosynthetic. Its modularity is achieved in part by concentrating many enzyme and ribozyme
component interactions on the ribosomal complex, which has the effect of chemically isolating
the translation function from much of the chemical activity going on elsewhere in the cell.
Looking forward, it may be possible to use an integrated metabolism and translation network to
study the emergence of the translation function as a network-level module in a way that is
abstracted from the mechanistic biochemical activities carried out by the ribosome and its
interaction partners.

564　Acknowledgements

566　This work is supported by the Wisconsin Alumni Research Foundation, the John Templeton
567　Foundation Grant #61926 and the Universidad Politécnica de Madrid (UPM) Margarita Salas
568　Fellowship, founded by the "Unión Europea-NextGeneration EU" (code UP2021-035).

570　Conflict of Interest

572　The authors declare no conflicts of interest

Code availability

The code necessary to reproduce this study can be found in the article Github repository: https://github.com/kacarlab/Translation-Network

Data availability

The data of this study can be found in the article Github repository: https://github.com/kacarlab/Translation-Network

Author contributions

B.C.Z., E.F., Z.R.A. and B.K. conceptualized and designed the study. B.C.Z. and E. F. performed the experiment and data analysis with input from Z.R.A. and B.K. B.C.Z. and B.K. wrote the paper with input from E.F. and Z.R.A. B.K. supervised the project. All authors read and approved the manuscript.

Figures and Tables

Figure 1. Schematic representation of the integrated cellular chemosynthesis network, composed of translation, polymer biosynthesis and metabolism layers. We built a translation layer (1) that includes all the relevant steps of that process such as initiation, peptidic bond formation, translocation, and termination. Translation generates peptides that appear in a biosynthesis layer (A). The biosynthesis layer describes interactions between polymers and the formation of reported hetero-complex multimers (2) to carry out metabolic functions. Proteins

601   in the biosynthesis layer act as enzymes or transporters enabling reactions (B) that generate
602   energy and building blocks of translation such as ATP, GTP and charged tRNA within the
603   metabolism layer (3). Finally, the free energy and building blocks of metabolism are employed as
604   substrate inputs for translation (C).

Figure 2. Overview of the translation portions of the integrated chemosynthesis network. . **A.** A simplified depiction of the 'canonical' translation process encoded within the translation network. **B.** A detailed visualization of the complete translation chemical reaction layer using Gephi 0.9.2 and a Force Atlas 2 layout[88] (settings: scaling=5.0, gravity=1.0, tolerance=1.0,

611   approximation=1.2). **C.** A depiction of the 30S and 50S component assembly steps that enable
612   ribosomal replication to occur.
613
614

Figure 3. Detailed and simplified visualizations of all reactions and compounds that compose the integrated chemosynthesis network. **A.** Full map with Translation (T) reactions in orange, Polymer Biosynthesis (B) reactions in green, and Metabolic (M) reactions in blue. Compounds (in white) are scaled in size according to their number of connections within the network. Network depicted with Gephi version 0.9.2 using a ForceAtlas2 algorithm[88] (settings: scaling=5.0, gravity=1.0, tolerance=1.0, approximation=1.2). **B.** A simplified depiction of the number of reaction connections between (directed arrows) and within (circular arrows) each of the layers.

Figure 4. Statistical analysis of the degree and molecular weight data of the integrated
chemosynthesis network and each of its layers. **A.** Complementary cumulative density function
(CCDF) of node connection plotted versus bins of node connection degree (k) and distribution

model comparisons for the complete chemosynthesis network; heterogeneous (exponential)

model shown in dashed line, and heterogeneous (power law) model shown in solid line. **B.** CCDF

plotted versus node connection degree (k) and distribution model comparisons for the

Translation (top), Biosynthesis (middle) and Metabolism (bottom) layers; heterogeneous

(exponential) model shown in dashed line, and heterogeneous (power law) model shown in solid

line, as in Panel A. **C.** Raw scatter data of the connection degree (k) and molecular weight (Da) of

all chemical objects in the network (upper panel) and same data depicted as a histogram of

molecular weight bins (middle panel), with histogram bin contributions color coded by molecule

type (bottom panel). **D.** Raw scatter data of the connection degree (k) and molecular weight (Da)

of all chemical objects in the network (upper panel) and same data depicted as a histogram of

molecular weight bins (middle panel), with histogram bin contributions color coded by layer

(bottom panel).

Supplementary Figures

642
643 **Figure S1**. Degree complementary cumulative distribution function (CCDF) of five different
644 strains.

645
Figure S2. Scatter plot of the t-distributed stochastic neighbor embedding (t-SNE) dimensional
reduction of node2vec embeddings of reactions from the complete integrated network.
Reactions within the different layers are color coded with all chemical compounds depicted in
gray and layer reactions from Translation in orange, Biosynthesis in green and Metabolism in
blue. Reactions from the translation and metabolism layers group distinctly from one another
with little overlap. Reactions from the biosynthesis layer do not group together in any distinct
way, but they broadly plot in the same areas that contain metabolic reactions.
653

Figure S3. Simplified view of the integrated network depicting the connections between
translation and metabolism modules. A module is a collection of reactions and compounds that
only connect to other reactions and compounds within the same assignment category.
Compounds that occur in more than one module are conserved (*i.e.*, not reduced to a module).
Module categories for translation were assigned during layer construction using labels as
shown in Figure 2B, while those for metabolism were taken without modification or alteration
from the source document from Feist et al [30]. Modules are coded from white (less central) to
purple (more central) using the betweenness centrality metric; the more visited a node is in a
path between two random nodes, the higher its betweenness-centrality.

668
669
670 **Figure S4**. Depiction of three different expected patterns of compound size distributions (top
671 panels) and their respective associated characteristics (bottom panels).
672
673
674

References

References

1. Oberhardt, M. A., Palsson, B. Ø. & Papin, J. A. Applications of genome-scale metabolic
reconstructions. *Mol Syst Biol* **5**, 320–320 (2009).
2. Lewis, M. A Tale of Two Repressors. *J Mol Biol* **409**, 14–27 (2011).
3. Chubukov, V., Gerosa, L., Kochanowski, K. & Sauer, U. Coordination of microbial metabolism.
Nat Rev Microbiol **12**, 327–340 (2014).
4. Chubukov, V., Zuleta, I. A. & Li, H. Regulatory architecture determines optimal regulation of
gene expression in metabolic pathways. *Proc National Acad Sci* **109**, 5127–5132 (2012).
5. Jeong, H., Tombor, B., Albert, R., Oltvai, Z. N. & Barabási, A.-L. The large-scale organization of
metabolic networks. *Nature* **407**, 651–654 (2000).
6. Barabási, A.-L., Albert, R. & Jeong, H. Mean-field theory for scale-free random networks. *Phys*
Statistical Mech Appl **272**, 173–187 (1999).
7. Ravasz, E., Somera, A. L., Mongru, D. A., Oltvai, Z. N. & Barabási, A.-L. Hierarchical
Organization of Modularity in Metabolic Networks. *Science* **297**, 1551–1555 (2002).
8. Newman, M. E. J. Analysis of weighted networks. *Phys Rev E* **70**, 056131 (2004).
9. Ma, W., Trusina, A., El-Samad, H., Lim, W. A. & Tang, C. Defining Network Topologies that Can
Achieve Biochemical Adaptation. *Cell* **138**, 760–773 (2009).
10. Nghe, P. *et al.* Prebiotic network evolution: six key parameters. *Mol Biosyst* **11**, 3206–3217
(2015).
11. Wagner, A. & Fell, D. A. The small world inside large metabolic networks. *Proc Royal Soc*
Lond Ser B Biological Sci **268**, 1803–1810 (2001).
12. Newman, M. E. J. The Structure and Function of Complex Networks. *Siam Rev* **45**, 167–256
(2003).
13. Shenhav, B., Solomon, A., Lancet, D. & Kafri, R. Transactions on Computational Systems
Biology I. in 14–27 (2005). doi:10.1007/978-3-540-32126-2_2.
14. Albert, R. & Barabási, A.-L. Statistical mechanics of complex networks. *Rev. Mod. Phys.* **74**,
47–97 (2002).
15. Holling, C. S. Cross-Scale Morphology, Geometry, and Dynamics of Ecosystems. *Ecol Monogr*
62, 447–502 (1992).
16. Fischer, J., Lindenmayer, D. B. & Montague-Drake, R. The role of landscape texture in
conservation biogeography: a case study on birds in south-eastern Australia. *Divers Distrib* **14**,
38–46 (2008).
17. Havlicek, T. D. & Carpenter, S. R. Pelagic species size distributions in lakes: Are they
discontinuous? *Limnol Oceanogr* **46**, 1021–1033 (2001).
18. Lambert, T. D., Malcolm, J. R. & Zimmerman, B. L. Amazonian small mammal abundances in
relation to habitat structure and resource abundance. *J Mammal* **87**, 766–776 (2006).
19. Stow, C., Allen, C. & Garmestani, A. Evaluating Discontinuities in Complex Systems: Toward
Quantitative Measures of Resilience. *Ecol Soc* **12**, (2007).
20. Nash, K. L. *et al.* Discontinuities, cross-scale patterns, and the organization of ecosystems.
Ecology **95**, 654–667 (2014).
21. Novozhilov, A. S., Wolf, Y. I. & Koonin, E. V. Evolution of the genetic code: partial
optimization of a random code for robustness to translation error in a rugged fitness landscape.
Biol Direct **2**, 24 (2007).

22. Koonin, E. V. & Novozhilov, A. S. Origin and Evolution of the Universal Genetic Code. *Annu*
Rev Genet **51**, 1–18 (2016).
23. Zaher, H. S. & Green, R. Quality control by the ribosome following peptide bond formation.
Nature **457**, 161–166 (2009).
24. Steinchen, W., Zegarra, V. & Bange, G. (p)ppGpp: Magic Modulators of Bacterial Physiology
and Metabolism. *Front Microbiol* **11**, 2072 (2020).
25. Prossliner, T., Gerdes, K., Sørensen, M. A. & Winther, K. S. Hibernation factors directly block
ribonucleases from entering the ribosome in response to starvation. *Nucleic Acids Res* **49**,
gkab017- (2021).
26. Starosta, A. L., Lassak, J., Jung, K. & Wilson, D. N. The bacterial translation stress response.
Fems Microbiol Rev **38**, 1172–1201 (2014).
27. Irving, S. E., Choudhury, N. R. & Corrigan, R. M. The stringent response and physiological
roles of (pp)pGpp in bacteria. *Nat Rev Microbiol* **19**, 256–271 (2021).
28. Fuente, I. M. D. la, Cortes, J. M., Pelta, D. A. & Veguillas, J. Attractor Metabolic Networks.
Plos One **8**, e58284 (2013).
29. Keseler, I. M. *et al.* The EcoCyc Database in 2021. *Front Microbiol* **12**, 711077 (2021).
30. Feist, A. M. *et al.* A genome-scale metabolic reconstruction for Escherichia coli K-12
MG1655 that accounts for 1260 ORFs and thermodynamic information. *Mol Syst Biol* **3**, 121
(2007).
31. King, Z. A. *et al.* BiGG Models: A platform for integrating, standardizing and sharing genome-
scale models. *Nucleic Acids Res.* **44**, D515–D522 (2016).
32. Alstott, J., Bullmore, E. & Plenz, D. powerlaw: A Python Package for Analysis of Heavy-Tailed
Distributions. *Plos One* **9**, e85777 (2014).
33. Clauset, A., Shalizi, C. R. & Newman, M. E. J. Power-Law Distributions in Empirical Data.
SIAM Rev. **51**, 661–703 (2009).
34. Hintze, A. & Adami, C. Evolution of Complex Modular Biological Networks. *Plos Comput Biol*
4, e23 (2008).
35. Srivastava, A., Kumar, S. & Ramaswamy, R. Two-layer modular analysis of gene and protein
networks in breast cancer. *Bmc Syst Biol* **8**, 81 (2014).
36. Girvan, M. & Newman, M. E. J. Community structure in social and biological networks. *Proc.*
Natl. Acad. Sci. **99**, 7821–7826 (2002).
37. Nelson, W. *et al.* To Embed or Not: Network Embedding as a Paradigm in Computational
Biology. *Frontiers Genetics* **10**, 381 (2019).
38. Grover, A. & Leskovec, J. node2vec: Scalable Feature Learning for Networks. *Arxiv* (2016).
39. Fey, M. & Lenssen, J. E. Fast Graph Representation Learning with PyTorch Geometric. *Arxiv*
(2019) doi:10.48550/arxiv.1903.02428.
40. Bastian, M., Heymann, S. & Jacomy, M. Gephi: An Open Source Software for Exploring and
Manipulating Networks. *Proc Int Aaai Conf Web Soc Media* **3**, 361–362 (2009).
41. Hu, X.-P., Dourado, H., Schubert, P. & Lercher, M. J. The protein translation machinery is
expressed for maximal efficiency in Escherichia coli. *Nat Commun* **11**, 5260 (2020).
42. Bar-Joseph, Z. *et al.* Computational discovery of gene modules and regulatory networks. *Nat*
Biotechnol **21**, 1337–1342 (2003).
43. Becker, S. A. & Palsson, B. O. Context-Specific Metabolic Networks Are Consistent with
Experiments. *Plos Comput Biol* **4**, e1000082 (2008).

44. Roy, A., Goberman, D. & Pugatch, R. A unifying autocatalytic network-based framework for
bacterial growth laws. *P Natl Acad Sci Usa* **118**, e2107829118 (2021).

45. Grimbs, A., Klosik, D. F., Bornholdt, S. & Hütt, M.-T. A system-wide network reconstruction
of gene regulation and metabolism in Escherichia coli. *Plos Comput Biol* **15**, e1006962 (2019).

46. Thiele, I. *et al.* Multiscale Modeling of Metabolism and Macromolecular Synthesis in E. coli
and Its Application to the Evolution of Codon Usage. *Plos One* **7**, e45635 (2012).

47. Thiele, I., Jamshidi, N., Fleming, R. M. T. & Palsson, B. Ø. Genome-Scale Reconstruction of
Escherichia coli's Transcriptional and Translational Machinery: A Knowledge Base, Its
Mathematical Formulation, and Its Functional Characterization. *Plos Comput Biol* **5**, e1000312
(2009).

48. Fuhrer, T., Heer, D., Begemann, B. & Zamboni, N. High-Throughput, Accurate Mass
Metabolome Profiling of Cellular Extracts by Flow Injection–Time-of-Flight Mass Spectrometry.
Anal Chem **83**, 7074–7080 (2011).

49. Dwivedi, P. *et al.* Metabolic profiling of Escherichia coli by ion mobility-mass spectrometry
with MALDI ion source. *J. Mass Spectrom.* **45**, 1383–1393 (2010).

50. Geiger, T., Wehner, A., Schaab, C., Cox, J. & Mann, M. Comparative Proteomic Analysis of
Eleven Common Cell Lines Reveals Ubiquitous but Varying Expression of Most Proteins*. *Mol
Cell Proteomics* **11**, M111.014050 (2012).

51. Mori, M. *et al.* From coarse to fine: the absolute Escherichia coli proteome under diverse
growth conditions. *Mol Syst Biol* **17**, e9536 (2021).

52. Kostinski, S. & Reuveni, S. Ribosome Composition Maximizes Cellular Growth Rates in E. coli.
Phys Rev Lett **125**, 028103 (2020).

53. Bianconi, G. Statistical mechanics of multiplex networks: Entropy and overlap. *Phys Rev E*
87, 062806 (2013).

54. Venkataram, S., Monasky, R., Sikaroodi, S. H., Kryazhimskiy, S. & Kacar, B. Evolutionary
stalling and a limit on the power of natural selection to improve a cellular module. *Proc
National Acad Sci* **117**, 18582–18590 (2020).

55. Hartwell, L. H., Hopfield, J. J., Leibler, S. & Murray, A. W. From molecular to modular cell
biology. *Nature* **402**, C47–C52 (1999).

56. Qi, Y. & Ge, H. Modularity and Dynamics of Cellular Networks. *Plos Comput Biol* **2**, e174
(2006).

57. Goldman, A. D. & Kacar, B. Cofactors are Remnants of Life's Origin and Early Evolution. *J Mol
Evol* **89**, 127–133 (2021).

58. Simonetti, A. *et al.* A structural view of translation initiation in bacteria. *Cell Mol Life Sci* **66**,
423 (2008).

59. Sharma, I. M. & Woodson, S. A. RbfA and IF3 couple ribosome biogenesis and translation
initiation to increase stress tolerance. *Nucleic Acids Res* **48**, 359–372 (2019).

60. Milón, P. & Rodnina, M. V. Kinetic control of translation initiation in bacteria. *Crit Rev
Biochem Mol* **47**, 334–348 (2012).

61. Benelli, D. & Londei, P. Begin at the beginning: evolution of translational initiation. *Res
Microbiol* **160**, 493–501 (2009).

62. Schuette, J. *et al.* GTPase activation of elongation factor EF-Tu by the ribosome during
decoding. *Embo J* **28**, 755–765 (2009).

63. Bakshi, S., Choi, H. & Weisshaar, J. C. The spatial biology of transcription and translation in
rapidly growing Escherichia coli. *Front Microbiol* **6**, 636 (2015).
64. Hartman, M. C. T., Josephson, K., Lin, C.-W. & Szostak, J. W. An Expanded Set of Amino Acid
Analogs for the Ribosomal Translation of Unnatural Peptides. *Plos One* **2**, e972 (2007).
65. Niederholtmeyer, H., Stepanova, V. & Maerkl, S. J. Implementation of cell-free biological
networks at steady state. *Proc National Acad Sci* **110**, 15985–15990 (2013).
66. Pérez-Mercader, J. Astrobiology, The Quest for the Conditions of Life. 337–360 (2002)
doi:10.1007/978-3-642-59381-9_22.
67. Gutenberg, B. & Richter, C. F. Frequency of earthquakes in California*. *B Seismol Soc Am* **34**,
185–188 (1944).
68. Krucker, S. & Benz, A. O. Energy Distribution of Heating Processes in the Quiet Solar Corona.
Astrophysical J Lett **501**, L213–L216 (1998).
69. Bak, P., Tang, C. & Wiesenfeld, K. Self-organized criticality. *Phys Rev A* **38**, 364–374 (1988).
70. Garmestani, A., Allen, C. & Gunderson, L. Panarchy: Discontinuities Reveal Similarities in the
Dynamic System Structure of Ecological and Social Systems. *Ecol Soc* **14**, (2009).
71. Holling, C. S. & Gunderson, L. H. Resilience and adaptive cycles. *In: Panarchy: Understanding*
Transformations in Human and Natural Systems, 25-62 (2002).
72. Shekhtman, L. M., Shai, S. & Havlin, S. Resilience of networks formed of interdependent
modular networks. *New J Phys* **17**, 123007 (2015).
73. Zubarev, D. Y., Rappoport, D. & Aspuru-Guzik, A. Uncertainty of Prebiotic Scenarios: The
Case of the Non-Enzymatic Reverse Tricarboxylic Acid Cycle. *Sci Rep-uk* **5**, 8009 (2015).
74. Braakman, R. & Smith, E. The compositional and evolutionary logic of metabolism. *Phys Biol*
10, 011001 (2013).
75. Preiner, M. *et al.* Catalysts, autocatalysis and the origin of metabolism. *Interface Focus* **9**,
20190072 (2019).
76. Smith, J. I., Steel, M. & Hordijk, W. Autocatalytic sets in a partitioned biochemical network. *J*
Syst Chem **5**, 2 (2014).
77. Orgel, L. In the beginning. *Nature* **439**, 915–915 (2006).
78. Shapiro, R. Small Molecule Interactions were Central to the Origin of Life. *Q Rev Biology* **81**,
105–126 (2006).
79. Sornette, D. Critical Phenomena in Natural Sciences, Chaos, Fractals, Self Organization and
Disorder: Concepts and Tools. *Springer Series Syne* 239–256 (2000) doi:10.1007/978-3-662-
04174-1_12.
80. Schwartz, N., Cohen, R., ben-Avraham, D., Barabási, A.-L. & Havlin, S. Percolation in directed
scale-free networks. *Phys Rev E* **66**, 015104 (2002).
81. Leicht, E. A. & D'Souza, R. M. Percolation on interacting networks. *Arxiv* (2009)
doi:10.48550/arxiv.0907.0894.
82. Newman, M. E. J. & Girvan, M. Finding and evaluating community structure in networks.
Phys Rev E **69**, 026113 (2004).
83. Newman, M. E. J. Mixing patterns in networks. *Phys Rev E* **67**, 026126 (2003).
84. Kryven, I. & Bianconi, G. Enhancing the robustness of a multiplex network leads to multiple
discontinuous percolation transitions. *Phys Rev E* **100**, 020301 (2019).
85. Paperin, G., Green, D. G. & Sadedin, S. Dual-phase evolution in complex adaptive systems. *J*
Roy Soc Interface **8**, 609–629 (2011).

86. Gonzalez, D. L., Giannerini, S. & Rosa, R. On the origin of degeneracy in the genetic code.
Interface Focus **9**, 20190038 (2019).
87. Muñuzuri, A. P. & Pérez-Mercader, J. Unified representation of Life's basic properties by a 3-
species Stochastic Cubic Autocatalytic Reaction-Diffusion system of equations. *Phys Life Rev* **41**,
64–83 (2022).
88. Jacomy, M., Venturini, T., Heymann, S. & Bastian, M. ForceAtlas2, a Continuous Graph
Layout Algorithm for Handy Network Visualization Designed for the Gephi Software. *Plos One* **9**,
e98679 (2014).