Hippocampal subfield volumes contribute to working memory interference control in aging: Evidence from longitudinal associations over 5 years.

Andersson, P[1], Samrani, G[2,3,4,5], Andersson, M[3,4.], and Persson J[1,2].

[1]Center for Life-span Developmental Research (LEADER), School of Law, Psychology, and Social Work, Örebro University, Sweden [2]Aging Research Center (ARC), Karolinska Institute and Stockholm University, Sweden; [3]Department of Radiation Sciences, Umeå University, Sweden; [4]Department of Integrative Medical Biology, Umeå University, Umeå, Sweden; [3]Umeå Center for Functional Brain Imaging, Umeå University, Sweden;

*Corresponding author: Pernilla Andersson, Center for Life-span Developmental Research (LEADER), School of Behavioral, Social and Legal sciences, Örebro University Fakultetsgatan 1, 702 81, Örebro, Sweden E-mail: pernilla.andersson@oru.se

ABSTRACT

In memory, familiar but no longer relevant information may disrupt encoding and retrieval of to-be-learned information. While it has been demonstrated that the ability to resolve proactive interference (PI) in working memory (WM) is reduced in aging, the neuroanatomical components of this decline have yet to be determined. Hippocampal (HC) involvement in age-related decline in control of PI is currently not known. In particular, the association between HC subfield volumes and control of PI in WM has not been examined previously. The associations between gray matter subfield volume and PI in WM across the adult life span was examined using cross-sectional (N = 280) and longitudinal (N = 157) data. Three main finding were obtained. First, cross-sectional analyses showed that HC volume was positively related to PI, but this effect was seen only in older adults. Additional analyses of HC subfield volume showed that in older adults this association was significant only in the subiculum. Second, 5-year longitudinal analyses across all participants revealed that reduced volume in the subiculum was related to impaired control of PI. Age-stratified analyses showed that this association was most pronounced in older adults. Third, we found that in older adults the effect of age on PI was mediated by GM volume in the HC. The current results show that HC volume is associated with the ability to control PI in WM, and that these associations are modulated by age.

INTRODUCTION

The constant accumulation of memory representations places high demands on a cognitive system that is efficient in controlling the content of memory in order to prevent interference from goal-irrelevant information. Proactive interference (PI) arises from existing memory representations conflicting with new target information. While PI has most commonly been investigated within the domain of episodic memory (Keppel, 1968; Roediger & McDermott, 1995), there is now much evidence that PI plays a critical role also in working memory (WM; Bunting, 2006; Emery, Hale, & Myerson, 2008). PI is also closely linked with performance levels on WM span tests (Kane & Engle, 2000; Lilienthal, 2017; Lustig, May, & Hasher, 2001; May, Hasher, & Kane, 1999).

While there is much evidence that WM declines with increasing age, evidence for age-related impairments in inhibition and interference control has been less consistent (Rey-Mermet & Gade, 2017; Verhaeghen, 2011). In general, it seems that in tasks in which conflicting information can be handled within the focus of attention show less of a decline compared to tasks in which the conflict arises from familiar, but no longer relevant information outside the focus of attention (Lustig & Jantz, 2014). One possibility is that age-related cognitive decline is mediated by deficits in interference control or other executive functions, with corresponding consequences on other aspects of cognition (Hasher & Zacks, 1988; McGabe, Roediger, McDaniel, Balota, & Hambrick, 2010). Indeed, an age-related decline in PI was recently demonstrated using a population-based life-span sample of adults (Samrani & Persson, 2021), and in this particular study PI was also found to predict mean level performance in other cognitive abilities.

Age-related impairments in interference control are most likely related to less efficient brain function, and changes in brain morphology and neuromodulation that occur with increasing age. Much of the work on the neural bases of interference control in WM has focused

on the prefrontal cortex. There is a general consensus that the left inferior frontal gyrus, together with the anterior cingulate cortex, and the insula are key regions involved in successful resolution of PI in WM (Badre & Wagner, 2005; Burgess & Braver, 2010; Henson, Shallice, Josephs, & Dolan, 2002; Jonides & Nee, 2006; Marklund & Persson, 2012; Nelson, Reuter-Lorenz, Persson, Sylvester, & Jonides, 2009; Persson, Larsson, & Reuter-Lorenz, 2013; Samrani, Bäckman, & Persson, 2019). However, despite evidence suggesting that the hippocampus (HC) is critical for interference control (Öztekin, Curtis, & McElree, 2008), little is known on how these regions contribute to resolving PI in WM. This is particularly important in studies on aging given the well documented effects of aging on HC gray matter volume (Gorbach et al., 2017; Persson et al., 2012; Raz, Rodrigue, Head, Kennedy, & Acker, 2004; Raz et al., 2003).

While the HC has been the focus of research of long-term memory functions for decades, its role in WM is less understood. There are indications that the HC may contribute to WM processes; activation in the HC has been demonstrated during fMRI WM tasks (Axmacher, Schmitz, Weinreic, Elger, & Fell, 2008; D E Nee, Jonides, & Berman, 2007), and patients with HC lesions are impaired in WM tasks (Cashdollar et al., 2009; Hannula, Tranel, & Cohen, 2006; Nichols, Kao, Verfaellie, & Gabrieli, 2006). There is also some recent functional imaging evidence showing that the HC is involved in successful resolution of PI (Öztekin et al., 2008). In particular, the HC may play a key role in distinguishing older from newer information in long term memory (Barredo, Öztekin, & Badre, 2015; Caplan, McIntosh, & De Rosa, 2007; Yassa & Stark, 2011), and WM (Leszczynski, 2011; Nauer, Whiteman, Dunne, Stern, & Schon, 2015; Ranganath & D'Esposito, 2001; Öztekin, Davachi, & McElree, 2010) thereby reducing PI. The HC could also be involved in resolving PI by contributing context or source information that can help distinguish familiar but no longer goal-relevant items from goal-relevant target items. This key function may be critical for both long-term memory and WM.

The HC might also be implicated in interference control through its association with pattern separation given that one critical feature of interference control is to separate between existing memory representations. Pattern separation is the ability to form distinct, non-overlapping representations from similar or overlapping information. It is often dissociated from pattern completion, which is an ability critical for reinstating information from a unique memory representation. Both these processes are central to how the brain processes information in an efficient manner. Pattern separation has been shown to rely on the dentate gyrus (DG) and Cornu Ammonis (CA3) regions in both animal work (Leutgeb, Leutgeb, Moser, & Moser, 2007; Rolls, 2016) and human studies (Azab, Stark, & Stark, 2014; Berron et al., 2016; Stevenson et al., 2016; Yassa & Stark, 2011). Recent studies have shown that also the subiculum plays a role in pattern separation (Bouyeure et al., 2021; Molitor, Sherrill, Morton, Miller, & Preston, 2021; Nash, Hodges, Muncy, & Kirwan, 2021; Potvin, Doré, & Goulet, 2009). The CA1 region of the HC is typically engaged during pattern completion processes (Bakker, Brock, Kirwan, Miller, & Stark, 2008). Also, recent studies have shown that pattern separation abilities decline with age (Stark, Yassa, Lacy, & Stark, 2013; Toner, Pirogovsky, Kirwan, & Gilbert, 2009), further suggesting that age-related hippocampal structural and functional changes could underlie impairments in pattern separation that may also affect interference control.

In contrast to episodic memory, the neuroanatomical underpinnings of interference control have received much less attention despite its likely role in age-related cognitive impairments. To the best of our knowledge, only one study to date has examined the cerebral morphological properties underlying interference control across the adult life span using both cross-sectional and longitudinal data (Samrani et al., 2019), and this particular study focused on frontal regions only. We focus here on the HC because of their known role in memory functions. In the current study, we use data from a population-based cohort study to examine cross-sectional (N = 280) and longitudinal (N = 157) associations between HC subfield

gray matter, and PI in WM across the adult life span. We hypothesize that larger gray matter volume in the HC should be related to better interference control (i.e., less PI). Since it has been demonstrated that brain function and volume might be differentially linked to cognition in younger and older adults (Koen & Rugg, 2019; Park et al., 2004; Rieckmann, Johnson, Sperling, Buckner, & Hedden, 2018; Van Petten, 2004), and that brain volume – cognition interactions may be stronger in older adults (Burzynska et al., 2012) we performed age-stratified analyses within groups on younger and older adults, as well as across the whole sample. We also obtained HC subfield volumes to test predictions regarding potential involvement of pattern-completion and pattern-separation processes in controlling familiar goal-irrelevant information in WM. We believe that this study takes critical steps to advance our understanding of hippocampal contributions to interference control across the adult life span.

MATERIAL AND METHODS

Participants

Participants were recruited from *The Betula prospective cohort study: Memory, health, and aging* (Nilsson et al., 1997), a deeply phenotyped longitudinal cohort. Participants were included from samples for which MRI measures were collected in 2008-2010 and 2013-2014. The age of the participants at baseline ranged from 25 to 90 years (mean = 56.9, standard deviation [SD] = 12.1; 49% female), and mean education level was 13 years (SD = 4.0 years). Individuals with clinical dementia and other neurological disorders at baseline were excluded. Participants were screened with the Mini-Mental State Examination (MMSE; Folstein, Folstein, & McHugh, 1975), and those scoring 25 and above were included. Participants with extremely low performance on the n-back task (proportion hits minus proportion false alarms < .1), indicating a very low adherence to task instructions, were not included. Six participants who did not pass the inclusion criteria for baseline testing but returned for the second time point and passed the performance-based criteria were included at follow-up testing. Thus, the total sample

consisted of 280 participants for cross-sectional analyses at baseline, and 157 participants for longitudinal analyses. Based on previous analyses conducted on this sample, this sample size has sufficient power. For a drop-out analysis, see Nyberg et al., (2019). To retain the diversity of the sample, exclusions were not made for: diabetes, hypertension, mild depressive symptoms, and other moderately severe medical conditions, which are common among the elderly. The Betula study was approved by the Regional Ethical Review Board in Umeå, and written consent was obtained from every participant. We investigated whether age modulated the relationship between brain volume and PI by dividing participants into two groups: one consisting of younger and middle-aged adults (25-65 years at baseline), and one consisting of older adults (70-90 years at baseline). Demographic information and cognitive performance can be found in Table 1.

Cognitive measures

PI was measured using a verbal 2-back WM task which included familiar lure items (Gray, Chabris, & Braver, 2003; Marklund & Persson, 2012; D E Nee et al., 2007) occurring either 1 or 2 trial(s) after the target position. The task included 40 trials; 21 non-familiar trials, 9 target trials, 8 3-back lures, and 2 4-back lures. *Target trials* matched the same stimuli as presented two trials earlier and required a 'Yes' response. *Lure trials* consisted of stimuli already presented 3- or 4 trials earlier and required a 'No' response. *New trials* were non-familiar trials that had never been presented and required a 'No' response. Stimuli and trial conditions were presented in the same fixed order for all participants. Stimuli consisted of Swedish nouns and were presented one at a time for 2.5 s, with an inter trial interval of 2 s. For each presented word, participants were instructed to press the "m" key on a standard Swedish keyboard, which corresponds to 'Yes' ("Yes, the word I now see has been shown two words ago") and the "x" key for 'No' ("No, the word I now see has not been shown two words ago"). Participants were instructed to answer as quickly and accurately as possible.

PI scores were calculated by combining the relative proportional difference in RT and accuracy between non-familiar trials (new trials) and 3- and 4-back familiar trials (lure trials). The PI scores based on accuracy and RTs were positively correlated, both at baseline (r = .295, P < 0.0005) and follow-up (r = .372. P < 0.0005), indicating shared variance between the two measures of PI. A relative difference score should represent a more salient and individual measure of executive control that considers baseline individual differences in the variables in question. Median RTs were used to reduce the influence of extreme values. Accuracy data for the n-back task is also reported under section 3.1.

MRI data acquisition and analyses

The MRI data was collected using a 3T GE scanner, equipped with a 32-channel head coil. The same scanner was used for baseline and follow-up data collection. T1-weighted images were acquired with a 3D fast spoiled gradient echo sequence (180 slices with a 1 mm thickness; TR: 8.2 ms, TE: 3.2 ms, flip angle: 12 degrees, field of view: 25 x 25 cm).

To extract HC volumes, T1-weighted images were processed using the Freesurfer image analysis suite. Hippocampal subfield segmentation was performed using the longitudinal stream (Reuter et al., 2012) of the Hippocampal Subfield module of FreeSurfer v.6.0 (https://surfer.nmr.mgh.harvard.edu/fswiki/HippocampalSubfields). The technical details of these procedures are described in prior publications (Dale, Fischl, & Sereno, 1999; Fischl et al., 2002; Fischl et al., 2004; Jovicich et al., 2006). Automated cortical and subcortical parcellation tools in the FreeSurfer software were used for volumetric segmentation, cortical surface reconstruction, and parcellation to quantify the brain volumes of interest. Cortical reconstructions and volumetric segmentations were performed on all images by executing a semi-automatic processing step (recon-all) within this software (Dale et al., 1999; Fischl et al., 2002). Identification of hippocampal subfields using FreeSurfer 6.0 should be considered

probabilistic, since it uses prior knowledge from ex vivo brains scanned at 7 T MRI combined with the available contrast information from MR images (Iglesias et al., 2016).

Scanner stability

From baseline to follow-up data collection, changes were implemented to the scanner software, and it is critical to check that these did not affect image quality. A quality assurance program based on Friedman and Glover (2006) was therefore run on a weekly basis on the scanner. For the structural data, the same T1-weighted fast spoiled gradient echo protocol as in the study was used to obtain volume data for the GE phantom. A threshold well above the noise level and the selected voxels were used to calculate the volume of the phantom. The relative volume change between the beginning of the quality assurance measurements and the time for T6 data collection was 0.45%, which is small compared to the expected average volume change in cortical regions in healthy elders (Fjell et al., 2009). Furthermore, the positive change in the measured volume of the phantom suggests that any significant volumetric decline observed in the study might be slightly underestimated.

Definition of cognitive and volumetric change

Change in proactive interference was estimated by dividing the PI-score from the second time point by the PI-score from the first time point. Similarly, we measured change in brain volumes by dividing the values from the second time point by the values from the first time point. The relative change from time points 1 to 2 takes into account the brain size for each individual, thus controlling for brain size without including TIV as a covariate in the analysis. Significant negative correlations between baseline level and rate of change in proactive interference were observed in both age cohorts (Pearson's r were -0.574 for younger adults and -0.704 for older adults). Individuals with low initial performance tended to improve or decline less rapidly and individuals starting with high levels of performance seemed to display more decline.

Selection of regions of interest

In order to reduce the number of comparisons, and because we did not have any specific hypotheses regarding laterality, we averaged volumes of the left and right hemisphere. For HC subfield analyses, we included the combined left and right CA1, dentate gyrus (DG), CA3, and subiculum subfields. We also performed analyses using total HC volume.

Statistical analyses

In the current study, relationships between cross-sectional estimates of brain volume and PI, as well as brain markers of change and change in PI were assessed using ordinary and partial correlation coefficients. Associations were controlled for age and education. Volumetric analyses were additionally controlled for total intracranial volume (TIV). Longitudinal analyses were additionally controlled for baseline level of PI. Partial eta squared (η_p^2) was used to measure effect size. Statistical analyses were performed using SPSS software ver. 26.0 (IBM, Armonk, NY, USA). Analyses were first performed across the whole age span, but since the relationship between local brain volume and cognition might change with increasing age (Burzynska et al., 2012; Razlighi et al., 2017), we also tested for associations within each of the three age groups. Multivariate outliers were identified using Mahalanobis distance at a $P < .001$.

In addition to analyses across the whole sample, we also tested whether there are age differences in the GM volume – PI associations. To do this, and similar to other studies (e.g. Andersson, Li, & Persson, 2022; Razlighi et al., 2017), we stratified participants into one group of younger (age at T5: 25-65), and one group of older (age at T5: 70-85) adults.

Moderation and mediation analyses were performed using the PROCESS macro (Hayes, 2013) implemented in SPSS. Using moderation analyses, we examined whether age was a significant moderator of the relationship between HC volume and PI. Mediation analyses were carried out to test whether the relation between the predictor (age) and the outcome (PI) was mediated – in total or in part – by a mediator variable (HC volume). For both moderation

and mediation analyses, a bootstrapping resampling strategy was implemented using 5000 bootstrap samples. For the mediation analysis, path *a* describes the direct effect of age on HC volume, path *b* represents the direct effect of HC volume on PI, and path *c* represents the direct effect between age and PI. Finally, path *c′* indicates the total effect of age on PI when HC volume is included in the model (Figure 3). Bias-corrected 95% confidence interval (CI) was computed to evaluate the contribution of the mediator (indirect effect, path a × b). CI reached significance when the interval range did not include zero. In order to examine the hypothesis that mediation would only be present in older adults, separate mediation analyses were carried out in younger and older adults, as well as in the whole sample. Standardized coefficients are reported for all mediation analyses.

RESULTS

Older age was associated with higher levels of proactive interference

Accuracy for non-familiar lures (98%) was significantly higher than for both familiar lures (60%; $F(279) = 575$, $p < .001$, $\eta_p^2 = .673$) and targets (77%; $F(279) = 404$, $p < .001$; $\eta_p^2 = .591$) and is on par with other studies using the modified version of the n-back task to induce familiarity based PI (Loosli, Rahm, Unterrainer, Weiller, & Kaller, 2014; Samrani, Bäckman, & Persson, 2017; Samrani & Persson, 2021) in older adults. Similarly, RT for non-familiar lures (1033 ms) was faster than for both familiar lures (1434 ms; $F(279) = 462$, $p < .001$, $\eta_p^2 = .629$) and targets (1144 ms; $F(279) = 37.7$, $p < .001$, $\eta_p^2 = .119$).

Moreover, older individuals had a lower ability to control interference as indicated by the positive correlation between age and PI. This association was found both at baseline ($r = .338$, $P < .0001$) and follow-up ($r = .432$, $P < .0001$). Change in PI over 5 years was, however, not significant ($P > .1$).

Cross-sectional and longitudinal effects of age on hippocampal brain volume

Linear regression analyses showed that age was a significant predictor of total HC volume both at baseline (r^2 = .210, β = -.458, P < .0001) and follow-up (r^2 = .281, β = -.549, P < .0001) after controlling for TIV. Similarly, analyses of HC sub-field volumes showed that age was a significant predictor of CA1 volume (baseline: r^2 = .1, β = -.313, P < .001; follow-up: r^2 = .112, β = -.351, P < .001), CA3 volume (baseline: r^2 = .067, β = -.229, P < .001; follow-up: r^2 = .085, β = -.298, P < .001), DG volume (baseline: r^2 = .180, β = -.298, P < .001; follow-up: r^2 = .181, β = -.467, P < .001) and subiculum volume (baseline: r^2 = .180, β = -.429, P < .001; follow-up: r^2 = .203, β = -.466, P < .001).

Cross-sectional associations between hippocampus volume and proactive interference in working memory

Regarding the relationship between local brain volume and proactive interference across all participants, the relationship at baseline showed a non-significant positive relationship (r = .065, P = .298). None of the associations with HC subfield volume were significant (all Ps > .1).

When analyses were performed for each age group separately, we found that for younger adults, PI was not associated with total HC volume, (r = -.082, P = .341). However, for older adults, we found a significant positive correlation with PI (r = .259, P = .003; Figure 1). A moderation analysis showed that age significantly moderated the association between HC volume and PI (F = 6.32, P = .0125, LLCI: .069, ULCI: .569). Analyses on HC subfields showed that the association between HC subfield volume and PI was significant for older adults in the subiculum (r = .191, P = .028) and in the CA1 (r = .178, P = .041). No other correlations were significant (all p*s* > .05).

Longitudinal associations between hippocampus volume and proactive interference in working memory

The correlation between 5-year change in HC volume and PI across the whole sample was not significant (r = -.037, P = .65). However, analyses on sub-field volume showed a significant correlation between change in subiculum volume and change in PI (r = -.213, P = .009). All other correlations were non-significant (P > .1). Age-stratified analyses for the whole HC were non-significant for both younger and older adults (P > .1), but analyses on sub-field volumes showed that in the group of older adults, change in HC sub-field volume was negatively associated with change in PI (Figure 2; CA1: r = -.297, P = .018; Subiculum: r = -.368, P = .003; DG: -.251, P = .047). Moderation analyses showed that age was not a significant moderator for the relationship between change in HC sub-field volume and change in PI. However, for the subiculum, results from the Johnson-Neyman procedure showed that this relationship was significant in participants above 58 years of age.

Mediation analyses

Single mediation analysis across all participants showed that HC volume was not a significant mediator of the effect of age on PI (estimated effect: -.035; lower level (LL) CI: -.006, higher level (HL) CI: .0019). Direct effects between age and HC volume (b = -.489, LLCI: -.044, HLCI: -.028, P < .001) and age and PI (b = .373, LLCI .0146, HLCI: .0305, P < .001) were significant, but the effect between HC volume and PI was not significant (b = .072, LLCI: -.049, HLCI: .165, P = .285). Age-stratified mediation analysis showed that for younger adults HC volume was not a significant mediator of the effect of age on PI (estimated effect: .012; LLCI: -.001, HLCI: .004). The direct effect between age and PI was significant (b = .273, LLCI: .008, HLCI: .032, P = .001), but the direct effects between and age and HC volume (b = -.123, LLCI -.021, HLCI: .003, P = .153) and HC volume and PI (b = -.099, LLCI: -.267, HLCI: .066, P = .235) were non-significant. For older adults, however, HC volume was a significant mediator of the effect of age on PI (estimated effect: -.137; LLCI: -.039, HLCI: -.008). All direct effects were significant (age-HC volume: b = -.478, LLCI: -.126, HLCI: -.064, P < .001;

age-PI: b = .286, LLCI: .0142, HLCI: .076, P = .045; HC volume-PI: b = .287 LLCI: .072, HLCI: .383, P = .045).

DISCUSSION

There is much evidence from patient and functional neuroimaging studies implicating HC as critical for efficient resolution of PI in WM. The current findings confirm the importance of the HC in interference control and suggest that this region is differentially involved in PI resolution in younger and older adults. Across all participants, HC volume was not significantly associated with PI. However, age-stratified as well as moderation analyses showed that in older, but not younger adults, larger HC volume was related to higher levels of PI. Further analyses of HC sub-field volumes revealed that larger subiculum and CA1 volume was related to higher levels of PI in older adults. Longitudinal analyses showed that in older adults, 5-year decrease in subiculum, CA1, and DG volume was related to decreased control of PI over 5 years. Finally, age-stratified mediation analyses revealed that, in older adults, whole HC volume significantly mediated the age – PI relationship. These results provide important insights into the neural architecture underlying interference control in WM.

Based on previous findings of a negative association between frontal cortex volume and PI (Samrani et al., 2019), along with data from fMRI of a link between HC activation and PI (D E Nee et al., 2007; Samrani & Persson, 2022; Öztekin et al., 2008), we expected that larger HC volume would be linked to less PI. While no significant correlations were observed across all participants, in older adults larger HC volume was associated with more, not less interference. Although speculative, the observed age-related impairment in interference control suggests that older adults may be less efficient in using recollection of contextual (temporal) information to resolve interference from familiar items. Instead, and in contrast to younger adults, larger HC volume in older adults may result in stronger memory

representations for all trials, and PI is likely to be affected by the underlying strength of the WM representation. This may increase the chance of familiar lure trials to exert a more interfering effect by inappropriate overgeneralization across contexts on the basis of familiarity alone.

Another possible, and yet largely unexplored, explanation for the link between HC volume and interference control is related to the role of this regions in pattern separation and completion. Without efficient pattern separation, memory representations can easily interfere with one another, resulting in PI from source confusion. Our results show that older individuals with both larger total HC volume and larger CA1 and subiculum volume are more affected by PI. One possibility is that larger volume in these regions is associated with a higher reliance on pattern completion during retrieval operations of WM information, and that this may result in increased rather than decreased PI. Indeed, theoretical models of hippocampal function predict that there should be a bias towards pattern completion in older adults (Wilson, Gallagher, Eichenbaum, & Tanila, 2006), and this view is also supported by empirical work (Vieweg, Stangl, Howard, & Wolbers, 2015; Yassa, Mattfeld, Stark, & Stark, 2011).

While the proposed idea which states that stronger WM representations may elicit more, and not less, interference is a sensible one (Szmalec, Verbruggen, Vandierendonck, & Kemps, 2011), it is not supported by recent finding showing that repeated encoding of items in WM resulted in less PI along with increased HC activation (Samrani & Persson, 2022). Additionally, studies that include individuals with memory impairments typically show more PI in these participants. For example, increased sensitivity to PI have been found in amnestic mild cognitive impairment (Ebert & Andersson, 2009; Loewenstein, Acevedo, Czaja, & Duara, 2004), Alzheimer´s disease (Crocco, Curiel, Acevedo, Czaja, & Loewenstein), and in patients with hippocampal lesions (Winocur, Moscovitch, & Bruni, 1996).

Longitudinal analyses of change across 5 years demonstrated that decreased HC volume was associated with decreased control of PI across all participants, but only in the subiculum. Age-stratified analyses found significant, negative, associations between change in CA1, subiculum and DG subfield volumes and change in PI in the older group. These findings corroborate previous cross-sectional findings that demonstrate a relationship between hippocampal function and PI in long-term memory (Blumenfeld & Ranganath, 2007; Wanjia, Favila, Kim, Molitor, & Kuhl, 2021). Moreover, it was recently demonstrated that 4 weeks of mindfulness training both improved the ability to control interference in WM and also resulted in an increase in HC volume (Greenberg et al., 2018). Importantly, the authors also found that mindfulness-related increase in HC volume was related to reduced PI.

Longitudinal subfield analyses did not show a separation between subfields implicated in pattern separation and pattern completion processes. Rather, the results show significant associations between increased PI and reduced volume in subfields implicated both in pattern separation and subfields implicated in pattern completion. While the literature at large agrees that the DG and CA3 subfields are differentially involved in pattern separation processes (Azab, Stark, & Stark, 2014; Berron et al., 2016; Stevenson et al., 2016; Yassa, Stark, et al., 2011) and the CA1 subfield is differentially involved in pattern completion processes (Bakker et al., 2008), some divergent results have been reported. For example, the CA3 subfield have been implicated in pattern completion in episodic memory (Guzowski, Knierim, & Moser, 2004; Rolls, 2016).

While the subiculum has received less attention than hippocampal subfields commonly considered part of the hippocampus proper, recent studies suggest that this region may play an important role in memory. While most reports suggest that the region contributes primarily to pattern separation (Potvin et al., 2009), observations of subiculum being involved in pattern completion have been reported as well (Bakker et al., 2008). As such, the specific

involvement of the subiculum in pattern separation and pattern completion processes is still unknown.

Interestingly, in a recent study, Hartopp and colleagues (2019) found significant associations between subiculum volume and fornix microstructure in a group of 31 healthy, older adults. Furthermore, this association was found to significantly contribute to episodic memory performance, suggesting that age-related alterations in subiculum structure could be driving fornix microstructural degradation. Similar investigations of the relationship between changes in fornix microstructure and volumetric changes in the subiculum subfield has, to our knowledge, not been conducted in relation to working memory. However, a recent study investigating correlations between white matter integrity and control of PI in working memory found that reduced fornix FA was associated with reduced ability to control PI across five years (Andersson et al., 2022). Follow-up analyses combining volumetric and microstructural measures could shed light on whether this fornix – subiculum relationship also contribute to the ability to control PI in working memory.

It is important to emphasize that we do not imply that the HC is uniquely involved in interference control. Rather, we believe that this region belongs to a network of regions, including the inferior frontal gyrus, anterior cingulate cortex, and insula that are all involved in control of PI. For example, animal tract-tracing studies have demonstrated direct connection between the HC and prefrontal cortex (Pandya, Van Hoesen, & Mesulam, 1981; Suzuki & Amaral, 1994) and anterior cingulate cortex (Morris, Pandya, & Petrides). Moreover, numerous human diffusion-tensor imaging (Draganski et al., 2008) and functional connectivity (D E Nee & Jonides, 2014) studies provide evidence for HC connectivity with the prefrontal cortex.

The current study has some notable limitations that needs to be mentioned. First, our longitudinal analyses contained only data from two measurement points. Therefore, we were not able to examine long-term trajectories of change, and this design also did not permit

independent estimations of retest effects, which are known to influence longitudinal data. Future studies that include three or more time points could provide data on long-term trajectories of change that we are unable to show in the current study design and may also provide more accurate estimation of retest effects. Second, the relatively low number of participants that remained for follow-up testing may have resulted in low statistical power in detecting longitudinal effects. This also restricted our age-differential analyses to include only cross-sectional data since the within-group sample size was too small for longitudinal analyses.

Taken together, our results support the idea that the hippocampus play a noticeable role in controlling PI in WM. Additionally, we demonstrate that the HC is differentially involved in PI resolution in younger and older adults. Somewhat unexpectedly, cross-sectional analyses across all participants showed that individuals with larger HC volume were more affected by PI, although age-specific analyses showed that this association was most pronounced in older adults. In addition, HC subfield analyses showed that larger subiculum volume was related to more PI across the whole sample, but this effect was largely driven by older adults. Speculatively, these results might indicate an age-related bias to pattern completion or familiarity-based recollection. Moreover, we provide new evidence that reduced HC subfield volumes over 5-years was related to concurrent increase in PI. We believe that these results provide new and important insights into the neural architecture underlying interference control in WM.

DATA AVAILABILITY

The data that support the findings of this study are available from the corresponding author upon reasonable request.

AUTHOR CONTRIBUTION

P.A: Writing Review & Editing, Writing – Original Draft Preparation,

G.S: Writing – Original Draft Preparation, Formal Analysis

M.A:

J.P: Writing – Original Draft Preparation, Writing Review & Editing, Conceptualization, Formal Analysis,

ACKNOWLEDGEMENTS

We acknowledge the contribution by the staff in the Betula project and all participants. The Freesurfer analyses were performed on resources provided by the Swedish National Infrastructure for Computing (SNIC) at HPC2N in Umeå.

FUNDING

The Betula Study was supported by the Bank of Sweden Tercentenary Foundation (Grant Nos. 1988-0082:17 and J2001-0682), Swedish Council for Planning and Coordination of Research (Grant Nos. D1988-0092, D1989-0115, D1990-0074, D1991-0258, D1992-0143, D1997-0756, D1997-1841, D1999-0739, and B1999-474), Swedish Council for Research in the Humanities and Social Sciences (Grant No. F377/1988-2000), the Swedish Council for Social Research (Grant Nos. 1988-1990: 88-0082 and 311/1991-2000), and the Swedish Research Council (Grant Nos. 345-2003-3883 and 315-2004-6977). The present research was additionally supported by a grant from the Swedish Research Council (421-2013-1039) to J.P.

CONFLICT OF INTEREST

None

REFERENCES

Andersson, P., Li, X., & Persson, J. (2022). The association between control of interference
and white-matter integrity: A cross-sectional and longitudinal investigation.
Neurobiology of Aging, 114, 49-60.
https://doi.org/10.1016/j.neurobiolaging.2022.03.002

Axmacher, N., Schmitz, D. P., Weinreic, I., Elger, C. E., & Fell, J. (2008). Interaction of
Working Memory and Long-Term Memory in the Medial Temporal Lobe. *Cerebral
Cortex, 18*, 2868-2878. https://doi.org/10.1093/cercor/bhn045

Azab, M., Stark, S. M., & Stark, C. E. L. (2014). Contributions of Human Hippocampal
Subfields to Spatial and Temporal Pattern Separation. *Hippocampus, 24*(3), 293-302.
https://doi.org/10.1002/hipo.22223

Badre, D., & Wagner, A. D. (2005). Frontal lobe mechanisms that resolve proactive
interference. *Cerebral Cortex, 15*(12), 2003-2012.
https://doi.org/10.1093/cercor/bhi075

Bakker, A., Brock, C., Kirwan, B., Miller, M., & Stark, C. E. L. (2008). Pattern Separation in
the Human Hippocampal CA3 and Dentate Gyrus. *Science, 319*(5870), 1640–1642.
https://doi.org/10.1126/science.1152882

Barredo, J., Öztekin, I., & Badre, D. (2015). Ventral fronto-temporal pathway supporting
cognitive control of episodic memory retrieval. *Cerebral Cortex, 25*(4), 1004–1019.
https://doi.org/10.1093/cercor/bht291

Berron, D., Schütze, H., Maass, A., Cardenas-Blanco, A., Kuijf, H., Kumaran, D., & Düzel,
E. (2016). Strong Evidence for Pattern Separation in Human Dentate Gyrus. *Journal
of Neuroscience, 36*, 7569-7579. https://doi.org/10.1523/JNEUROSCI.0518-16.2016

Blumenfeld, R. S., & Ranganath, C. (2007). Prefrontal cortex and long-term memory encoding: an integrative review of findings from neuropsychology and neuroimaging., *13*(3), 280–291. https://doi.org/10.1177/1073858407299290

Bouyeure, A., Patil, S., Mauconduit, F., Poiret, C., Isai, D., & Noulhiane, M. (2021). Hippocampal subfield volumes and memory discrimination in the developing brain *31*(11), 1202-1214. https://doi.org/10.1002/hipo.23385

Bunting, M. (2006). Proactive interference and item similarily in working memory. . *Journal of Experimental Psychology: Learning, Memory, and Cognition., 32*(2), 183-196. https://doi.org/10.1037/0278-7393.32.2.183

Burgess, G. C., & Braver, T. S. (2010). Neural mechanisms of interference control in working memory: effects of interference expectancy and fluid intelligence. *PLoS ONE, 5*(9). https://doi.org/10.1371/journal.pone.0012861

Burzynska, A. Z., Nagel, I. E., Preuschhof, C., Gluth, S., Bäckman, L., Li, S. C., . . . Heekeren, H. R. (2012). Cortical thickness is linked to executive functioning in adulthood and aging. *Human Brain Mapping, 33*, 1607–1620. https://doi.org/10.1002/hbm.21311

Caplan, J. B., McIntosh, A. R., & De Rosa, E. (2007). Two distinct functional networks for successful resolution of proactive interference. *Cerebral Cortex, 17*(7), 1650–1663. https://doi.org/10.1093/cercor/bhl076

Cashdollar, N., Malecki, U., Rugg-Gunn, F. J., Duncan, J. S., Lavie, N., & Düzel, E. (2009). Hippocampus-dependent and -independent theta-networks of active maintenance. *Proceedings of the National Academy of Sciences in the U S A, 106*(48), 20493-20498. https://doi.org/10.1073/pnas.0904823106

Crocco, E., Curiel, R. E., Acevedo, A., Czaja, S. J., & Loewenstein, D. A. An evaluation of deficits in semantic cueing and proactive and retroactive interference as early features

of Alzheimer's disease *American Journal of Geriatric Psychiatry, 22*(9), 889-897.

https://doi.org/10.1016/j.jagp.2013.01.066

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis. I.

Segmentation and surface reconstruction. *NeuroImage, 9*, 179-194.

https://doi.org/10.1006/nimg.1998.0395

Draganski, B., Kherif, F., Klöppel, S., Cook, P. A., Alexander, D. C., Parker, G. J., . . .

Frackowiak, R. S. (2008). Evidence for segregated and integrative connectivity

patterns in the human Basal Ganglia. *Journal of Neuroscience, 28*(28), 7143-7152.

https://doi.org/10.1523/JNEUROSCI.1486-08.2008

Ebert, P. L., & Andersson, N. D. (2009). Proactive and retroactive interference in young

adults, healthy older adults, and older adults with amnestic mild cognitive impairment

Journal of the International Neuropsychological Society, 15, 83 – 93.

https://doi.org/10.1017/S1355617708090115

Emery, L., Hale, S., & Myerson, J. (2008). Age differences in proactive interference, working

memory, and abstract reasoning. *Psychology and Aging, 23*(3), 634-645.

https://doi.org/10.1037/a0012577

Fischl, B., Salat, D. H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., . . . Dale, A. M.

(2002). Whole brain segmentation: automated labeling of neuroanatomical structures

in the human brain. *Neuron, 33*, 341-355. https://doi.org/10.1016/s0896-

6273(02)00569-x

Fischl, B., van der Kouwe, A., Destrieux, C., Halgren, E., Ségonne, F., Salat, D., . . . Dale, A.

M. (2004). Automatically parcellating the human cerebral cortex. *Cerebral Cortex,

14*(1), 11-22. https://doi.org/10.1093/cercor/bhg087

Fjell, A. M., Walhovd, K. B., Fennema-Notestine, C., McEvoy, L. K., Hagler, D. J., Holland,

D., . . . Dale, A. M. (2009). One-year brain atrophy evident in healthy aging. *Journal*

of Neuroscience, 29(48), 15223-15231. https://doi.org/10.1523/JNEUROSCI.3252-09.2009

Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). Mini-mental state: A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research., 12*(3), 189-198. https://doi.org/10.1016/0022-3956(75)90026-6

Friedman, L., & Glover, G. H. (2006). Report on a multicenter fMRI quality assurance protocol. *Journal of Magnetic Resonance Imaging, 22*(6), 827-839. https://doi.org/10.1002/jmri.20583

Gorbach, T., Pudas, S., Lundquist, A., Orädd, G., Josefsson, M., Salami, A., . . . Nyberg, L. (2017). Longitudinal association between hippocampus atrophy and episodic-memory decline. *Neurobiology of Aging, 51*, 167-176. https://doi.org/10.1016/j.neurobiolaging.2016.12.002

Gray, J. R., Chabris, C. F., & Braver, T. S. (2003). Neural mechanisms of general fluid intelligence. *Nature Neuoscience, 6*(3), 316-322. https://doi.org/10.1038/nn1014

Greenberg, J., Romero, V. L., Elkin-Frankston, S., Bezdek, M. A., Schumacher, E. H., & Lazar, S. W. (2018). Reduced interference in working memory following mindfulness training is associated with increases in hippocampal volume. *Brain Imaging and Behavior, 13*(2), 366-376. http://doi.org/10.1007/s11682-018-9858-4

Guzowski, J. F., Knierim, J. J., & Moser, E. I. (2004). Ensemble Dynamics of Hippocampal Regions CA3 and CA1,. *Neuron, 44*(4), 581-584. https://doi.org/10.1016/j.neuron.2004.11.003

Hannula, D. E., Tranel, D., & Cohen, N. J. (2006). The long and the short of it: relational memory impairments in amnesia, even at short lags. *Journal of Neuroscience, 26*(32), 8352-8359. https://doi.org/10.1523/JNEUROSCI.5222-05.2006

Hartopp, N., Wright, P., Ray, N. J., Evans, T. E., Metzler-Baddeley, C., Aggleton, J. P., & O'Sullivan, M. J. (2019). A Key Role for Subiculum-Fornix Connectivity in Recollection in Older Age. *Frontiers in Systems Neuroscience, 12*(70). https://doi.org/10.3389/fnsys.2018.00070

Hasher, L., & Zacks, R. T. (1988). Working memory, comprehension, and aging: A review and a new view. . In G. H. Bower (Ed.), *The Psychology of Learning and Motivation* (Vol. 22, pp. 193-225). New York: Academic Press. https://doi.org/10.1016/S0079-7421(08)60041-9

Hayes, A. F. (2013). *Introduction to Mediation, Moderation, and Conditional Process Analysis: A Regression Based Approach*: Guildford Press.

Henson, R. N. A., Shallice, T., Josephs, O., & Dolan, N. J. (2002). Functional magnetic resonance imaging of proactive interference during spoken cued recall. *NeuroImage, 17*, 543-558.

Iglesias, J. E., Van Leemput, K., Augustinack, J., Insausti, R., Fischl, B., & Reuter, M. (2016). Bayesian longitudinal segmentation of hippocampal substructures in brain MRI using subject-specific atlases. *NeuroImage, 141*, 542-555. https://doi.org/10.1016/j.neuroimage.2016.07.020

Jonides, J., & Nee, D. E. (2006). Brain mechanisms of proactive interference in working memory. *Neuroscience, 139*(1), 181-193. https://doi.org/10.1016/j.neuroscience.2005.06.042

Jovicich, J., Czanner, S., Greve, D., Haley, E., van der Kouwe, A., Gollub, R., . . . Dale, A. M. (2006). Reliability in multi-site structural MRI studies: effects of gradient non-linearity correction on phantom and human data. *NeuroImage, 30*, 436-443. https://doi.org/10.1016/j.neuroimage.2005.09.046

Kane, M. J., & Engle, R. W. (2000). Working-memory capacity, proactive interference, and divided attention: limits on long-term memory retrieval. *Journal of Experimental Psychology: Learning, Memory and Cognition, 26*, 336–358. https://doi.org/10.1037//0278-7393.26.2.336

Keppel, G. (1968). Retroactive and proactive inhibition. . In T. R. Dixon & D. L. Horton (Eds.), *Verbal behavior and general behavior theory* (pp. 172-213). Englewood Cliffs, NJ: Prentice Hall.

Koen, J. D., & Rugg, M. D. (2019). Neural Dedifferentiation in the Aging Brain. *Trends in Cognitive Sciences, 23*(7), 547-559. https://doi.org/10.1016/j.tics.2019.04.012

Leszczynski, M. (2011). How does hippocampus contribute to working memory processing? *Frontiers in human neuroscience, 5*(1-2). https://doi.org/10.3389/fnhum.2011.00168

Leutgeb, J. K., Leutgeb, S., Moser, M.-B., & Moser, E. I. (2007). Pattern Separation in the Dentate Gyrus and CA3 of the Hippocampus. *Science, 315*(5814), 961-966. https://doi.org/10.1126/science.1135801

Lilienthal, L. (2017). Individual differences in proactive interference in verbal and visuospatial working memory. *Memory, 25*(8), 1110-1116. https://doi.org/10.1080/09658211.2016.1268633

Loewenstein, D. A., Acevedo, A., Czaja, S. J., & Duara, R. (2004). Cognitive rehabilitation of mildly impaired Alzheimer disease patients on cholinesterase inhibitors. *American Journal of Geriatric Psychiatry, 12*(4), 395-402. https://doi.org/10.1176/appi.ajgp.12.4.395

Loosli, S. V., Rahm, B., Unterrainer, J. M., Weiller, C., & Kaller, C. P. (2014). Developmental change in proactive interference across the life span: evidence from two working memory tasks. *Developmental Psychology, 50*(4), 1060-1072. https://doi.org/10.1037/a0035231

Lustig, C., & Jantz, T. (2014). Questions of age difference in interference control: When and how, not if? *Brain Research, 1612*, 59-69. https://doi.org/10.1016/j.brainres.2014.10.024

Lustig, C., May, C. P., & Hasher, L. (2001). Working memory span and the role of proactive interference. *Journal of Experimental Psychology: General, 130*(2), 199-207. https://doi.org/10.1037/0096-3445.130.2.199

Marklund, P., & Persson, J. (2012). Context-dependent switching between proactive and reactive working memory control mechanisms in the right inferior frontal gyrus. *NeuroImage, 63*(3), 1552-1560. https://doi.org/10.1016/j.neuroimage.2012.08.016

May, C. P., Hasher, L., & Kane, M. J. (1999). The role of interference in memory span. . *Memory and Cognition, 27*(5), 759-767. https://doi.org/10.3758/BF03198529

McGabe, D. P., Roediger, H. L. I., McDaniel, M. A., Balota, D. A., & Hambrick, D. Z. (2010). The relationship between working memory capacity and executive functioning: Evidence for a common attention contruct. *Neuropsychology, 24*(2), 222-243. https://doi.org/10.1037/a0017619

Molitor, R. J., Sherrill, K. R., Morton, N. W., Miller, A. M., & Preston, A. R. (2021). Memory Reactivation during Learning Simultaneously Promotes Dentate Gyrus/CA2,3 Pattern Differentiation and CA1 Memory Integration *Journal of Neuroscience, 41*(4), 726-738. https://doi.org/10.1523/JNEUROSCI.0394-20.2020

Morris, R., Pandya, D. N., & Petrides, M. Fiber system linking the mid-dorsolateral frontal cortex with the retrosplenial/presubicular region in the rhesus monkey. *Journal of Comparative Neurology, 407*, 183–192. https://doi.org/10.1002/(sici)1096-9861(19990503)407:2<183::aid-cne3>3.0.co;2-n

Nash, M. I., Hodges, C. B., Muncy, N. M., & Kirwan, C. B. (2021). Pattern separation beyond the hippocampus: A high-resolution whole-brain investigation of mnemonic

discrimination in healthy adults *Hippocampus, 31*(4), 408-421.

https://doi.org/10.1002/hipo.23299

Nauer, R. K., Whiteman, A. S., Dunne, M. F., Stern, C. E., & Schon, K. (2015). Hippocampal

subfield and medial temporal cortical persistent activity during working memory

reflects ongoing encoding. . *Frontiers in Systems Neuroscience, 9*(30).

https://doi.org/10.3389/fnsys.2015.00030

Nee, D. E., & Jonides, J. (2014). Frontal–Medial Temporal Interactions Mediate Transitions

among Representational States in Short-Term Memory. *Journal of Neuroscience,*

34(23), 7964-7975. https://doi.org/10.1523/JNEUROSCI.0130-14.2014

Nee, D. E., Jonides, J., & Berman, M. G. (2007). Neural mechanisms of proactive

interference-resolution. *NeuroImage, 38*(4), 740-751.

https://doi.org/10.1016/j.neuroimage.2007.07.066

Nelson, J. K., Reuter-Lorenz, P. A., Persson, J., Sylvester, C. Y., & Jonides, J. (2009).

Mapping interference resolution across task domains: a shared control process in left

inferior frontal gyrus. *Brain Research, 1256*, 92-100.

https://doi.org/10.1016/j.brainres.2008.12.001

Nichols, E. A., Kao, Y. C., Verfaellie, M., & Gabrieli, J. D. E. (2006). Working memory and

long-term memory for faces: Evidence from fMRI and global amnesia for involvement

of the medial temporal lobes. *Hippocampus, 16*(7), 604-616.

https://doi.org/10.1002/hipo.20190

Nilsson, L. G., Bäckman, L., Erngrund, K., Nyberg, L., Adolfsson, R., Bucht, G., . . .

Winblad, G. (1997). The Betula prospective cohort study: Memory, health and aging.

Aging Neuropsychology and Cognition, 4(1), 1-32.

https://doi.org/10.1080/13825589708256633

Nyberg, L., Andersson, M., Lundquist, A., Salami, A., & Wåhlin, A. (2019). Frontal Contribution to Hippocampal Hyperactivity During Memory Encoding in Aging. *Frontiers in Molecular Neuroscience*. https://doi.org/10.3389/fnmol.2019.00229

Pandya, D. N., Van Hoesen, G. W., & Mesulam, M. M. (1981). Efferent connections of the cingulate gyrus in the rhesus monkey. *Experimental Brain Research, 42*, 319–330. https://doi.org/10.1007/BF00237497

Park, D. C., Polk, T. A., Park, R., Minear, M., Savage, A., & Smith, M. R. (2004). Aging reduces neural specialization in ventral visual cortex. *Proceedings of the National Academy of Sciences in the U S A, 101*(35), 13091-13095. https://doi.org/10.1073/pnas.0405148101

Persson, J., Larsson, A., & Reuter-Lorenz, P. A. (2013). Imaging Fatigue of Interference Control Reveals the Neural Basis of Executive Resource Depletion. *Journal of Cognitive Neuroscience, 25*(3), 338-351. https://doi.org/10.1162/jocn_a_00321

Persson, J., Pudas, S., Lind, J., Kauppi, K., Nilsson, L., & Nyberg, L. (2012). Longitudinal Structure-Function Correlates in Elderly Reveal MTL Dysfunction with Cognitive Decline. *Cerebral Cortex, 22*(10), 2297-2304. https://doi.org/10.1093/cercor/bhr306

Potvin, O., Doré, F. Y., & Goulet, S. (2009). Lesions of the dorsal subiculum and the dorsal hippocampus impaired pattern separation in a task using distinct and overlapping visual stimuli. *Neurobiology of Learning and Memory, 91*(3), 287–297. https://doi.org/10.1016/j.nlm.2008.10.003

Ranganath, C., & D'Esposito, M. (2001). Medial temporal lobe activity associated with active maintenance of novel information. *Neuron, 31*(5), 865-873. https://doi.org/10.1016/s0896-6273(01)00411-1

Raz, N., Rodrigue, K. M., Head, D., Kennedy, K. M., & Acker, J. D. (2004). Differential aging of the medial temporal lobe: A study of a five-year change. *Neurology, 62*, 433-438. https://doi.org/10.1212/01.wnl.0000106466.09835.46

Raz, N., Rodrigue, K. M., Kennedy, K. M., Head, D., Gunning-Dixon, F., & Acker, J. D. (2003). Differential aging of the human striatum: longitudinal evidence. *American Journal of Neuroradiology, 24*(9), 1849-1856. PMID: 14561615

Razlighi, Q., Oh, H., Habeck, C., O'Shea, D., Gazes, E., Eich, T., . . . Stern, Y. (2017). Dynamic Patterns of Brain Structure–Behavior Correlation Across the Lifespan. *Cerebral Cortex, 27*(7), 3586–3599. https://doi.org/10.1093/cercor/bhw179

Rey-Mermet, A., & Gade, M. (2017). Inhibition in aging: What is preserved? What declines? A meta-analysis. *Psychonomic Bulletin and Review*. https://doi.org/10.3758/s13423-017-1384-7

Rieckmann, A., Johnson, K. A., Sperling, R. A., Buckner, R. L., & Hedden, T. (2018). Dedifferentiation of caudate functional connectivity and striatal dopamine transporter density predict memory change in normal aging. *Proceedings of the National Academy of Sciences in the U S A, 115*(40), 10160-10165. https://doi.org/10.1073/pnas.1804641115

Roediger, H. L., & McDermott, K. B. (1995). Creating false memories: Remembering words not presented in lists. . *Journal of Experimental Psychology: Learning, Memory, and Cognition., 21*, 803-814. https://doi.org/10.1037/0278-7393.21.4.803

Rolls, E. T. (2016). Pattern separation, completion, and categorisation in the hippocampus and neocortex. *Neurobiology of Learning and Memory, 129*, 4-28. https://doi.org/10.1016/j.nlm.2015.07.008

Samrani, G., Bäckman, L., & Persson, J. (2017). Age-differences in the temporal properties of proactive interference in working memory. *Psychology and Aging, 32*(8), 722-731. https://doi.org/10.1037/pag0000204

Samrani, G., Bäckman, L., & Persson, J. (2019). Interference Control in Working Memory Is Associated with Ventrolateral Prefrontal Cortex Volume. *Journal of Cognitive Neuroscience, 31*(10), 1491-1505. https://doi.org/10.1162/jocn_a_01430

Samrani, G., & Persson, J. (2021). Proactive interference in working memory is related to adult age and cognitive factors: cross-sectional and longitudinal evidence from the Betula study. *Neuropsychology, Development and Cognition. Section B Aging, Neuropsychology and Cognition, 28*(1), 108-127. https://doi.org/10.1080/13825585.2020.1713982

Samrani, G., & Persson, J. (2022). Automatic and effortful control of interference in working memory can be distinguished by unique behavioral and functional brain representations. *NeuroImage, 253*(119098). https://doi.org/10.1016/j.neuroimage.2022.119098

Stark, S. M., Yassa, M. A., Lacy, J. W., & Stark, C. E. L. (2013). A task to assess behavioral pattern separation (BPS) in humans: Data from healthy aging and mild cognitive impairment. *Neuopsychologia, 51*(12), 2442-2449. https://doi.org/10.1016/j.neuropsychologia.2012.12.014.

Stevenson, B., Vieweg, P., Gao, F., Gilboa, A., Wolbers, T., Black, S. E., & Rosenbaum, S. (2016). The Human Dentate Gyrus Plays a Necessary Role in Discriminating New Memories. *Current Biology, 26*(19), 2629-2634. https://doi.org/10.1016/j.cub.2016.07.081

Suzuki, W. A., & Amaral, D. G. (1994). Perirhinal and parahippocampal cortices of the macaque monkey: cortical afferents. *Journal of Comparative Neurology, 350*, 497–533. https://doi.org/10.1002/cne.903500402

Szmalec, A., Verbruggen, F., Vandierendonck, A., & Kemps, E. (2011). Control of interference during working memory updating. *Journal of Experimental Psychology: Human Perception and Performance, 37*(1), 137–151. https://doi.org/10.1037/a0020365

Toner, C. K., Pirogovsky, E., Kirwan, C. B., & Gilbert, P. E. (2009). Visual object pattern separation deficits in nondemented older adults. *Learning & memory, 16*(5), 338–342. https://doi.org/10.1101/lm.1315109

Van Petten, C. (2004). Relationship between hippocampal volume and memory ability in healthy individuals across the lifespan: Review and meta-analysis. *Neuropsychologia, 42*, 1394-1413. https://doi.org/10.1016/j.neuropsychologia.2004.04.006

Verhaeghen, P. (2011). Aging and Executive Control: Reports of a Demise Greatly Exaggerated. *Current Directions in Psychological Science, 20*(3), 174-180. https://doi.org/10.1177/0963721411408772

Vieweg, P., Stangl, M., Howard, L. R., & Wolbers, T. (2015). Changes in pattern completion-a key mechanism to explain age-related recognition memory deficits? *Cortex, 64*, 343-351. https://doi.org/10.1016/j.cortex.2014.12.007

Wanjia, G., Favila, S. E., Kim, G., Molitor, R. J., & Kuhl, B. A. (2021). Abrupt hippocampal remapping signals resolution of memory interference. *Nature Communications, 12*(4816). https://doi.org/10.1038/s41467-021-25126-0

Wilson, I. A., Gallagher, M., Eichenbaum, H., & Tanila, H. (2006). Neurocognitive aging: Prior memories hinder new hippocampal encoding. *Trends in Neurosciences, 29*, 662–670. https://doi.org/10.1016/j.tins.2006.10.002

Winocur, G., Moscovitch, M., & Bruni, J. (1996). Heightened interference on implicit, but not explicit, tests of negative transfer: Evidence from patients with unilateral temporal lobe lesions and normal old people. *Brain and Cognition, 30*(1), 44–58. https://doi.org/10.1006/brcg.1996.0004

Yassa, M. A., Mattfeld, A. T., Stark, S. M., & Stark, C. E. (2011). Age-related memory deficits linked to circuit-specific disruptions in the hippocampus. *Proceedings of the National Academy of Sciences in the U S A, 108*(21), 8873-8868. https://doi.org/10.1073/pnas.1101567108

Yassa, M. A., & Stark, C. E. L. (2011). Pattern separation in the hippocampus. *Trends in Neurosciences, 34*(10), 515-525. https://doi.org/10.1016/j.tins.2011.06.006

Öztekin, I., Curtis, C. E., & McElree, B. (2008). The medial temporal lobe and inferior prefrontal cortex jointly support interference resolution in working memory. *Journal of Cognitive Neuroscience, 21*(10), 1967-1979. https://doi.org/10.1162/jocn.2008.21146

Öztekin, I., Davachi, L., & McElree, B. (2010). Are representations in working memory distinct from representations in long-term memory? Neural evidence in support of a single store. . *Psychological Science, 21*(8), 1123–1133. https://doi.org/10.1177/0956797610376651

Table 1. Demographic and cognitive measurements

	Younger adults	Older adults	
Baseline			P-value
N	139	141	
Age (range)	50.1 (25-60)	71(65-90)	<.001
Sex (F/M)	73/66	75/66	.91
Education (years); (SD)	14.4 (3.1)	12 (4.5)	<.001
Vocabulary (SD)	24.1 (3.4)	23.8 (3.7)	.42
MMSE (SD)	28.5 (1.3)	28.2 (1.3)	.11
2-back target accuracy (SD)	79 (14)	75 (16)	<.001
Blood pressure (systolic) (SD)	129.5 (17)	143.4 (16.6)	<.001
Blood pressure (diastolic) (SD)	79.9 (10.4)	81.1 (8.3)	.28
Follow-up			
N	90	67	
Age (range)	55.8 (30-65)	74.2 (70-90)	<.001
Sex (F/M)	41/49	31/33	.75
Education (years); (SD)	14.3 (3.3)	12.7 (4.7)	.01
Vocabulary (SD)	24.4 (3)	24.3 (3.4)	.92
MMSE (SD)	28.4 (1.2)	28.3 (1.4)	.59
2-back target accuracy (SD)	73.3 (20.1)	54.6 (19.1)	<.001
Blood pressure (systolic) (SD)	133.8 (14.2)	143.9 (17.7)	<.001
Blood pressure (diastolic) (SD)	80.7 (7.8)	80.9 (8.8)	.89

MMSE = Mini-mental state examination; SD = Standard deviation

Figure 1. Correlations between relative interference score and HC volume at baseline.

Standardized residuals are corrected for age, education, and TIV.

Figure 2. (Top) Correlations between relative interference score and HC volume at baseline for younger and older adults. Standardized residuals are corrected for age, education, and TIV. (Bottom) Correlations between relative interference score and HC sub-field volumes (CA1 and CA3/DG) at baseline for younger and older adults. Standardized residuals are corrected for age, education, and TIV.

All participants

Younger adults

Older adults

Figure 3. Mediation analysis investigating the three-way association between age, HC volume and PI. Note: "a" indicates the effect of age on HC volume, "b" indicates the effect HC volume on PI, , "c" indicates the direct effect of age on PI, and "c'" indicates the total effect (direct and indirect) of age on PI.