1

Evolution of phenotypic plasticity during environmental fluctuations

Running head: Evolution of phenotypic plasticity

Zuzana Sekajova[1], Erlend I. F. Fossen[1,2], Elena Rosa[1], Irja I. Ratikainen[3], Manon Tourniaire-Blum[1], Elisabeth Bolund[1] and Martin I. Lind[1,4] [*]

[1] Animal Ecology, Department of Ecology and Genetics, Uppsala

University, Uppsala, Sweden

[2] Oslo Centre for Biostatistics and Epidemiology, Institute of Basic Medical Sciences,

University of Oslo, Oslo, Norway

[3] Department of Biology, Centre for Biodiversity Dynamics, Norwegian University of

Science and Technology, Trondheim, Norway

[4] Department of Environmental and Biosciences, Halmstad University, Halmstad, Sweden

[*] Corresponding author

Corresponding author

Martin Lind

Animal Ecology, Dept. of Ecology and Genetics

Uppsala University

Norbyvägen 18D

752 36 Uppsala

Sweden

martin.lind@ebc.uu.se

Author contributions

ZS, MIL, IIR and EB designed the experiment, ZS, ER and MT-B performed the phenotypic

assays with the aid of MIL. EIFF and ZS analysed the data, with the aid of EB. ZS, MIL and

EIFF drafted the manuscript. All authors contributed to the revision of the manuscript.

Acknowledgements

We thank Alexei Maklakov for access to the experimental evolution lines. This work was

financed by the Swedish Research Council (Vetenskapsrådet, grant no. 2016-05195 and

2020-04388) and the Carl Tryggers Foundation (grant no. CTS 17:285) to MIL.

Conflict of Interest statement

The authors report no conflict of interest.

Data Availability Statement

Upon acceptance, the data will be archived at *Figshare*

Abstract

Evolution in fluctuating environments is predicted to disfavor specialization and instead

select for alternative strategies, such as phenotypic plasticity or possibly bet-hedging,

depending on the accuracy of environmental cues and type of fluctuations. While these two

alternatives are often contrasted in theoretical studies, their evolution are seldom studied

together in empirical work.

We used experimental evolution in the nematode worm *Caenorhabditis*

remanei to simultaneously study the evolution of plasticity and bet-hedging in environments

differing only in their temperature variability. We exposed worms for 30 generations to either

fluctuating or slowly increasing temperature, these two environments had the same average

temperature over evolutionary time. After experimental evolution, we scored size at sexual

maturity and fitness in full siblings reared in two different temperatures, optimal 20°C and

mildly stressful 25°C.

Experimental evolution in the fluctuating environment resulted in the evolution of

increased body size plasticity but not increased bet-hedging, compared to evolution in the

slowly changing environment. Plasticity followed the temperature size rule as size decreased

with increasing temperature and this plastic response was adaptive. In addition, we

documented substantial standing genetic variation in body size, which represents a potential

for further evolutionary change.

Introduction

Natural environments are generally not stable, but fluctuate on both long and short

timescales, and a developing organism needs to take this environmental variation into

account. While the parental environment can be a reliable cue if the environmental

fluctuations are slow and predictable (Lachmann & Jablonka, 1996; Kuijper & Hoyle, 2015;

Leimar & McNamara, 2015; Uller *et al.*, 2015) and can result in the evolution of anticipatory

parental effects (Dey *et al.*, 2016; Lind *et al.*, 2020), faster and random fluctuations select

against inheritance of the parental phenotype. Instead, phenotypic plasticity or bet-hedging is

predicted to be adaptive (Moran, 1992; Simons, 2011). If the developmental environment

provides reliable cues for the optimal offspring phenotype, theory predict the evolution of

adaptive phenotypic plasticity (Moran, 1992; Gavrilets & Scheiner, 1993; Simons, 2011)

which is the ability of a genotype to produce different phenotypes depending on

environmental conditions (West-Eberhard, 2003). If plasticity is present, individuals will

canalize fitness between environments by adjusting their phenotype according to the

environmental conditions.

Alternatively, individuals may express bet-hedging, which acts to reduce variation in

fitness (especially to avoid very low fitness values in certain environmental states) at the cost

of lowered arithmetic mean fitness, often by producing offspring with a range of phenotypes

(diversified bet-hedging), where some of the offspring always matches the environment and

is successful (Philippi & Seger, 1989). In contrast to phenotypic plasticity, bet-hedging is

generally seen as favored when environmental cues have low predictability (Cohen, 1966;

Slatkin, 1974; Meyers & Bull, 2002; Kussell & Leibler, 2005; Wolf *et al.*, 2005), or when

there is low correlation between the environment of development and selection (Tufto, 2015).

While both phenotypic plasticity and bet-hedging can be adaptive in variable environments (Simons, 2011; Furness *et al.*, 2015), most empirical studies of evolution in variable environments focus on the evolution of plasticity. While phenotypic plasticity is common and well documented (DeWitt & Scheiner, 2004), very few studies have investigated the evolution of plasticity. These studies generally follow two lines, either they investigate whether increased plasticity evolves in more variable environments (Moran, 1992; Tufto, 2015), or whether increased plasticity evolves as a (possibly transient) step in adaptation to a novel but stable environment (Lande, 2009; Chevin *et al.*, 2010; Levis & Pfennig, 2020). Studies focusing on the role of environmental heterogeneity have found stronger phenotypic plasticity in natural populations (Lind & Johansson, 2007; Lind *et al.*, 2011) or species (Hollander, 2008) originating from more variable environments. Moreover, two recent experimental evolution studies in microalgae have shown that variable environments can select for increased phenotypic plasticity (Schaum *et al.*, 2022), but that very unpredictable environments also can select against plasticity (Leung *et al.*, 2020). One possible explanation for selection against plasticity during rapid fluctuations is that it may take time for the plastic phenotype to be induced (Burton *et al.*, 2022; Dupont *et al.*, 2024). Studies that instead focus on the evolution of plasticity during adaptation to new stable environments have found that plasticity can rapidly evolve in a new environment (Sikkink *et al.*, 2014b; Corl *et al.*, 2018), but also that maladaptive plasticity can play a major role (Ghalambor *et al.*, 2015; Campbell-Staton *et al.*, 2021).

In contrast to phenotypic plasticity, empirical studies documenting bet-hedging are rare (Simons, 2011), and the few examples suggest that it is more likely to evolve if the environments differ dramatically in fitness (Bull, 1987), such as delayed germination in desert plants (Philippi, 1993; Clauss & Venable, 2000; Venable, 2007), diapause in killifish (Furness *et al.*, 2015) or experimental evolution in fluctuating environments with large fitness

differences (Beaumont *et al.*, 2009; Graham *et al.*, 2014). However, bet-hedging has been

suggested as a mechanism explaining the strain-specific variance in reproduction between

natural isolates of the nematode *Caenorhabditis elegans* (Diaz & Viney, 2014).

One environmental factor that is known to result in evolutionary adaptations

(Berteaux *et al.*, 2004; Geerts *et al.*, 2015), but also alternative strategies, is temperature. Not

only is temperature gradually increasing due to the ongoing climate change (Berteaux *et al.*,

2004), but climate change also results in increased temperature variability (Easterling *et al.*,

2000; Palmer & Räisänen, 2002; Van Aalst, 2006) potentially favoring evolution of increased

plasticity or possibly bet-hedging. Indeed, most documented responses to climate change in

natural populations are caused by pre-existing plasticity, while genetic adaptation seems rare

(Merilä & Hendry, 2014).

Among traits showing plastic responses to temperature, body size is of immense

importance to reproductive fitness. For females, a large body generally translates into

increased egg production, and also males often benefit from large size (Hedrick & Temeles,

1989; Andersson, 1994). Therefore, perhaps surprisingly, in warm environments organisms

generally increase growth rate, accelerate maturation but mature at a smaller size, which is

called the temperature-size rule (Ray, 1960; Atkinson, 1994; Verberk *et al.*, 2021). This has

sometimes been argued to be a passive by-product of other temperature-dependent processes

(Atkinson, 1994; van der Have & de Jong, 1996; Forster *et al.*, 2011). However, small size

may actually be actively regulated and beneficial in warm environments (Fryxell *et al.*, 2020;

Verberk *et al.*, 2021) for example by being advantageous for thermoregulation (Partridge &

Coyne, 1997), or allowing better regulation of oxygen demand and supply ratio (Walczyńska

et al., 2015; Verberk *et al.*, 2021).

We set out to investigate whether exposure to fluctuating temperature results in

evolution of increased phenotypic plasticity compared to evolution in an environment with

slowly increased temperature and whether any plasticity in body size is adaptive, using

experimental evolution in the nematode *Caenorhabditis remanei*. *C. remanei* has a fast

generation time and, since it is dioecious, harbors substantial standing genetic variation

which makes it ideal for experimental evolution studies (Teotónio *et al.*, 2017). In addition,

C. remanei has been shown to respond to manipulations in temperature (Sikkink *et al.*,

2014a; Lind *et al.*, 2020), has the ability to respond plastically to new environmental

conditions (Lind *et al.*, 2020), and the plasticity to withstand heat-shock can itself evolve

(Sikkink *et al.*, 2014b). Body size of *C. remanei* is under directional selection towards an

optimum larger than the mean size under standard temperature conditions (Stångberg *et al.*,

2020), and pharmacologically lowered body size results in lower female reproduction (Lind

et al., 2016). It's close relative *C. elegans* also follow the temperature-size rule (Kammenga

et al., 2007). As an alternative to phenotypic plasticity, we also investigated if populations

evolving in variable environments had evolved increased diversifying bet-hedging.

We used previously established experimental evolution populations of *C. remanei*

(described in Lind *et al.*, 2020). During experimental evolution, replicate populations were

exposed to 30 generations in one of two regimes; *Fast temperature cycles* or *Increased*

warming. The *Fast temperature cycles* regime were switched between two temperatures

(20°C, 25°C) every second generation, and although the fluctuations are deterministic, it

represents an uncorrelated (and therefore unpredictable) fluctuating environment each

generation, as the next generation will either be in the same or in a different temperature. The

evolution in this environment was compared to the *Increased warming* regime, where worms

were exposed to experimental evolution in a gradually increasing temperature which slowly

raised from 20°C to 25°C over 30 generations, and which served as a control. Importantly,

these two regimes had the same average temperature (22.5°C) over evolutionary time, and

only differed in the rate and predictability of environmental change. While no theoretical

model to our knowledge have investigated evolution of plasticity in fluctuating versus slowly

changing environments, it corresponds to different degrees of fluctuation, which is well

explored theoretically (Moran, 1992; Gavrilets & Scheiner, 1993; Tufto, 2015). After the 30

generations of experimental evolution, we reared full-siblings in either standard 20°C, or

warm 25°C, and scored them for reproduction and body size.

We predict that worms evolving in fluctuating temperature every 2nd generation

would evolve increased phenotypic plasticity (relative to the *Increased warming* regime),

since the timescale of this environmental variation is well within the parameter space where

plasticity (and bet-hedging) is favored (Tufto, 2015). We predict that evolution of phenotypic

plasticity is more likely than the evolution of bet hedging, since the differences in fitness

between the two temperatures is likely to be relatively small (Lind *et al.*, 2020). If increased

phenotypic plasticity has evolved, we predict that the *Fast temperature cycle* populations

would show increased size difference between temperatures, but not increased phenotypic

variance within one temperature. If instead increased diversifying bet-hedging had evolved,

we predict that the *Fast temperature cycle* populations will show (1) increased within-family

variance within each temperature, and (2) decreased heritability of size, because of an

increased environmental component of phenotypic variance (Tufto, 2015). We also predict

that any plasticity in body size will follow the temperature-size rule, and that this plasticity is

adaptive.

Methods

Experimental evolution

For the experimental evolution we used *C. remanei* nematode worms, strain SP8 which has been lab adapted for 15 generations at 20°C and subsequently exposed to 30 generations of experimental evolution in two regimes (*Increased warming* and *Fast temperature cycles*). The experimental evolution has been previously described in detail in Lind *et al.*, (2020b). Briefly, in the *Increased warming* experimental evolution regime, the temperature gradually raised from 20°C to 25°C, which is a novel and mildly stressful temperature. This gradual change over 30 generations represent an increase of 0.1°C every 2.13 days and results in a correlated parental and offspring environment. In the second regime, *Fast temperature cycles*, the temperature fluctuated every second generation between 20°C and 25°C, resulting in 14 temperature shifts but no exposure to the intermediate temperatures. The environmental change is deterministic (every second generation) but since parents and offspring would end up in either the same or in different temperature, it represents uncorrelated parental and offspring environment. The generation time in *C. remanei* is temperature dependent; 4 days long in 20°C and 3.4 days long in 25°C. Despite these differences, the average temperature and the total chronological time of experimental evolution were identical for both regimes, at 22.5°C and 110 days respectively.

Each evolutionary regime consisted of six replicate populations. The populations were maintained on 92 mm Petri dishes poured with NGM agar in climate chambers set to 60% relative humidity. In order to prevent bacterial and fungal contamination, the agar and bacterial LB contained the antibiotics streptomycin and kanamycin and the fungicide nystatin. The plates were seeded with 2 ml of an antibiotic-resistant OP50-1 (pUC4K) strain of *E. coli* (Stiernagle, 2006) that served as a source of food. Every 1-2 days, a piece of agar containing approximately 150 worms of mixed ages was cut and transferred to a new plate

containing fresh bacteria. This resulted in populations with overlapping generations that were

maintained in a constant exponential growth phase. After the experimental evolution,

populations were expanded for two generations and frozen in -80°C for later revival and

subsequent phenotypic assays.

Experimental set-up

Each replicate population of each of the two selection regimes was run in a separate block

resulting in 12 experimental blocks in total. For logistic reasons we focus on females, since

they are responsible for population growth rate and their fitness is straightforward to

measure.

Briefly, populations were revived from freezing and exposed to 25°C for 3

generations, to avoid any maternal effects associated with freezing. The third generation of

worms were split into eight families, each family consisting of one male and one female

worm. From each family, we randomly picked eight offspring females (full siblings) and

placed four females in 20°C and four in 25°C. Since our focus was evolution in females, their

fitness was assessed by mating them with standardized males from the ancestral line. Females

were kept together with two males (in case one of the males would be infertile/escape from

the plate). The ancestral line was, in contrast to selection regimes, maintained for three

generations in 20°C and subsequently split into 20°C and 25°C together with studied females.

For the detailed description of the experimental set up, see supplementary figure 1. In total,

we used 48 families per evolutionary regimes, and for each family we set up 4 offspring in

each temperature.

Phenotypic assays

Daily reproduction

Female and male worms were transferred to a new plate every 24 hours. The old plate was kept for two days to allow eggs to hatch and afterwards the number of viable offspring was counted to determine age-specific reproduction and calculate individual fitness. In the case of the males dying or escaping from the plate, they were replaced by a new male of the same age from the ancestral line. The female worm was discarded after dying, or after three consecutive days of zero reproduction.

Body size

The body size of the worm changes with time, reaching a peak before it declines (Lind *et al.*, 2016), and the age at maximum body size is temperature dependent. Worms in 20°C reach their peak size at day 4 of adulthood (Lind *et al.*, 2016). The peak size in 25°C is on day 2 of adulthood, which was determined during pilot assays. Photographs of worms were taken during their peak size using a Lumenera Infinity2-5C digital microscope camera mounted on a Leica M165C stereomicroscope. Body size was measured from the photographs using *ImageJ 1.46r* (https://imagej.nih.gov/ij/) as total cross-section area.

Statistical analyses

All statistical analyses were conducted in R 3.6.1 (R Core Team, 2015).

Individual fitness

We used the age-specific reproduction data to calculate rate-sensitive individual fitness λ_{ind} for each individual (Brommer *et al.*, 2002), which is analogous to the intrinsic rate of population growth (Stearns, 1992). Individual fitness was calculated by constructing a population projection matrix for each individual, and then calculating the dominant

eigenvalue of this matrix, following McGraw & Caswell, (1996). Since we kept the

population size and age structure constant during experimental evolution, individual fitness is

maximized during evolution and is therefore the most appropriate fitness measure for this

study (Mylius & Diekmann, 1995).

Thermal reaction norms

To test whether the degree of phenotypic plasticity has evolved, we used linear mixed-effect

models to separately estimate the thermal reaction norms of body size and individual fitness,

using the package *lme4* (Bates *et al.*, 2015) in R. The models included either body size (area)

or individual fitness (λ_{ind}) as response variables. The full model included three fixed effects:

mean-standardized temperature as a covariate, the experimental evolution regime as a

categorical factor, and an interaction between temperature and evolution regime. We expect

this interaction to be significant if the degree of plasticity has evolved during experimental

evolution. Experimental line and dam identity were included as random effects. Significance

of the fixed effects was evaluated using Wald χ^2 tests. Pseudo-R^2 values were calculated as

the squared correlation coefficient between fitted values from the model and observed values.

Selection

To test if temperature responses in size are adaptive, we estimated the selection on body size

and compared it to the observed temperature response. Selection on body size (area) was

estimated using mixed-effect models in R with the package *lme4* and individual fitness (λ_{ind})

as the response variable. The full model included the following fixed effects: area, area^2,

temperature (as a categorical factor with 2 levels), experimental evolution regime and all

interactions between these variables except for interactions involving both area and area^2

together. Experimental line was included as random effect, assumed to only affect the

variance of the intercept. Significance of fixed effects was evaluated using Wald χ^2 tests.

Pseudo-R^2 values were calculated as the squared correlation coefficient between fitted values

from the model and observed values. The optimal size (i.e. the area that maximizes fitness)

was calculated as: $-b/(2 \times c)$, where b = the slope of the regression (i.e. the linear selection

gradient) and c = the squared term of the regression (i.e. the quadratic selection gradient).

Confidence intervals of the temperature-specific optimal sizes were generated by

bootstrapping, implemented in the *boot* package using 10 000 bootstrap replicates.

Within family coefficient of variance

To test whether the degree of diversifying bet-hedging has evolved, we tested whether the

experimental evolution regimes differed in the mean within family variance within

temperatures. For each family, we used the trait values of the offspring (within a temperature)

to calculate the within family variance. To account for differences in the means of the traits,

we used within family means (i.e. the mean trait value of the family's offspring) to mean-

standardize the variance by calculating the within family coefficient of variance (CV): CV =

σ/μ, where σ = within family standard deviation, and μ = within family mean. An ANOVA

was used within each temperature to test if the evolution regimes differed in their mean

within family CV.

Since it is more difficult to detect differences in variances than differences in

means, we also performed power calculations on our ability to detect whether within-family

CVs differ between the selective regimes. Balanced one-way ANOVA power calculations

were performed to estimate the effect sizes possible to detect with power ranging from 0.70

to 0.95. Effect sizes, η^2, were obtained for our sample size of N = 48 per selection regime and

a significance level of 0.05. η^2 is calculated as the sum of squared explained by the treatment

(here: selection regime) divided by the total sum of squares, and has an equivalent

interpretation as an R^2.

Genetic variance and correlations

For body size (cross-section area), genetic variance and genetic correlations across

temperature were estimated using animal models using the package *MCMCglmm* (Hadfield,

2010) in R. Univariate models were used to estimate genetic variance, whereas bivariate

models were used to estimate genetic correlations, both using Gaussian family for trait

distribution. An inverse Wishart prior with parameters V = 1 and nu = 0.02 were used in both

univariate and bivariate models. Pedigree data linking offspring to parents, based on full-sib

relationships, was included in the models. Convergence of the models were ensured by

evaluating diagnostic plots of posterior distributions, using the convergence diagnostic half-

width test by Heidelberger and Welch (1983), and by ensuring that the autocorrelation

between MCMC samples was close to zero.

For univariate models, body area was used as the response variable. Temperature (as a

categorical factor with 2 levels), experimental evolution regime, and an interaction between

the two, were included as fixed effects. Genetic variance (V_G), variance due to differences

between experimental lines, and residual variances were estimated separately as random

effects in the full model for each temperature-by-evolution regime combination. The full

model ran for 4.2×10^6 MCMC iterations, 0.2×10^6 samples were discarded as burnin, and the

thinning interval was 4000, resulting in a sample size of 1000 MCMC-samples. Reduced

models, subset by temperature-by-evolution regime combination, were used to assess the

statistical significance of V_G, by comparing the deviance information criterion (DIC) of

models with versus without genetic variance included.

Broad sense heritability ($H^2 = V_G/V_P$, where V_P = total phenotypic variance after

accounting for variance due to experimental line effects) and broad sense evolvability (I^2 =

V_G/mean^2, Hansen *et al.*, 2003, 2011) were used to estimate the population's evolutionary

potential of body size. This was estimated for each temperature-by-evolution regime

combination. Evolvability measures the expected percentage change in a trait per generation

under unit strength of selection. Compared to heritability, evolvability is independent from

the environmental variance and represents a measure of the evolutionary potential that is

comparable across traits, populations and species when applied to traits with a natural zero

and which are strictly positive (Hansen *et al.*, 2011).

Genetic correlations of body size were estimated using bivariate animal models in

MCMCglmm. The data was split in two subsets based on the experimental evolution regimes,

resulting in genetic correlations being estimated separately for the two evolution regimes.

Body area was the response variable and was treated as two traits in the models, as area at

20°C and area at 25°C. Random effects in the full models included genetic covariance

between the two temperatures, whereas V_G, variance due to differences between experimental

lines, and residual variances were estimated separately for each temperature. The full models

ran for 2.05×10^6 MCMC iterations, 0.05×10^6 samples were discarded as burnin, and the

thinning interval was 2000, resulting in a sample size of 1000 MCMC-samples. Reduced

models without genetic covariance were used to access the statistical significance of the

genetic covariance, by comparing the DIC of models with versus without genetic covariance

included. The genetic correlation of body size across temperatures was calculated by dividing

the genetic covariance by the product of the genetic standard deviation of the two

temperatures. This was done on the posterior distributions, in order to carry the error

forwards in the analyses.

To compare posterior distributions of H^2, I^2 and genetic correlations across

temperatures and selection regimes, we calculated, within each MCMC sample, the pairwise

differences in these measures and checked if the posterior distributions of these differences

had a 95% credibility interval that included zero. Pairwise comparisons of distributions were

only performed between evolution regimes within temperature, and between temperatures

within evolution regimes.

Results

<u>Thermal reaction norms</u>

Size

Size decreased significantly with increasing temperature (Wald $\chi^2 = 309.93$, df = 1, p <

0.001, Fig. 1A). There was a significant interaction between temperature and evolution

regime, where *Fast temperature cycles* had a steeper slope, meaning that it had evolved

increased plasticity in size (Wald $\chi^2 = 5.82$, df = 1, p = 0.016). However, the intercepts

(representing size at the mean temperature) were not significantly different between evolution

regime (Wald $\chi^2 = 0.09$, df = 1, p = 0.769). The models pseudo $R^2 = 0.51$. Variance

components: $V_{dam} = 25.03$, $V_{Line} = 20.92$, and $V_{residual} = 108.19$.

Individual fitness (λ_{ind})

The mean total reproduction decreased with temperature (mean $\pm$ SE: 20°C, 780 $\pm$ 17; 25°C,

672 $\pm$ 17; p < 0.001 for the difference between the temperatures). However, the individual

fitness (λ_{ind}) increased significantly with increasing temperature (Wald $\chi^2 = 3670$, df = 1, p <

0.001, Fig. 1B). The evolution regimes did not differ significantly in intercepts (Wald χ^2 =

0.02, df = 1, p = 0.887), nor was there a significant interaction between temperature and

evolution regime (Wald χ^2 = 0.97, df = 1, p = 0.324). The best fit models pseudo R^2 = 0.86.

Variance components: V_{dam} = 0.05, V_{Line} = 0.01, and $V_{residual}$ = 0.21.

Figure 1. Thermal reaction norms showing means ± SE. (A) regression lines (with mean-standardized

temperature): Fast; Area = 80.87 ± 2.08 – 3.14 ± 0.22 × Temperature. Increase; Area = 81.74 ± 2.08 – 2.38 ±

0.22 × Temperature. (B) regression lines (with mean-standardized temperature): Fast; λ_{ind} = 4.85 ± 0.06 + 0.43 ±

0.01 × Temperature. Increase; λ_{ind} = 4.86 ± 0.06 + 0.41 ± 0.01 × Temperature.

<u>Selection</u>

There was significant linear and quadratic selection on body size (linear slope: Wald $\chi^2 =$

59.4, df = 1, p<0.001. Quadratic term: Wald $\chi^2 = 40.3$, df = 1, p < 0.001). The selection

differed significantly between temperatures (Fig. 2), given by a significant overall

temperature effect (Wald $\chi^2 = 2992$, df = 1, p < 0.001) and significant interaction effects

between temperature and size (linear slope: Wald $\chi^2 = 13.3$, df = 1, p < 0.001. Quadratic

term: Wald $\chi^2 = 12.8$, df = 1, p < 0.001). Maximum individual fitness (i.e. the optimal size) is

predicted to be 93.73 µm at 20°C [95% bootstrap CI: 87.61, 112.98], and 84.19 µm at 25°C

[95% bootstrap CI: 79.81, 92.61]. Selection was however not significantly different between

the experimental evolution regimes (p > 0.22 for main effect and interactions between

evolution regime and temperature or body size). No 3-way interaction between temperature,

body size and evolution regime were significant (p > 0.18). The best fit model's pseudo $R^2 =$

0.85. Variance components: $V_{Line} = 0.016$, and $V_{residual} = 0.204$.

Figure 2. Selection on body size (area). Experimental evolution regimes were not statistically different, but are

shown with separate lines. Overall regression line for 20°C: $\lambda_{ind} = 1.596 \pm 0.818 + 0.048 \pm 0.020 \times$ Area $-$

$2.57 \times 10^{-4} \pm 1.16 \times 10^{-4} \times$ Area^2. Overall regression line for 25°C: $\lambda_{ind} = 0.138 \pm 0.614 + 0.142 \pm 0.017 \times$ Area $-$

$8.43 \times 10^{-4} \pm 1.20 \times 10^{-4} \times$ Area^2. The mean size per temperature (m_{20}, and m_{25}) and optimal size (O_{20}, and O_{25}) are

shown for 20°C and 25°C respectively. Individual fitness is higher in 25°C due to decreased development time,

even if total reproduction is lower.

<u>Within family CV</u>

The evolution regimes did not differ significantly in within family CV of body size or of

individual fitness at either temperature (Table 1). Moreover, the distributions of within family

CV overlapped considerably between evolution regimes (Fig. 3). Power calculations showed

that we had a power of 90% to detect effects where the evolutionary regimes explained at

least 10% of the variation in within family CV (supplementary figure 2).

Table 1. Within family coefficient of variance (CV). Size (area) measured in μm^2, fitness as individual lambda

(λ_{ind})

Trait	Temp. (°C)	Experimental evol. regime	Within family CV (mean±SE)	Difference between evolution regimes	
				F (ndf=1, ddf=96)	p
Area	20	Fast temp. cycles	0.092 ± 0.009	3.799	0.054
		Increased warming	0.115 ± 0.009		
	25	Fast temp. cycles	0.100 ± 0.011	0.014	0.905
		Increased warming	0.102 ± 0.011		
Fitness	20	Fast temp. cycles	0.049 ± 0.005	0.266	0.608
		Increased warming	0.053 ± 0.005		
	25	Fast temp. cycles	0.087 ± 0.011	0.338	0.563
		Increased warming	0.079 ± 0.011		

Figure 3. Distribution of within family coefficient of variance (CV) for two traits at two temperatures. The within family CV was estimated within temperature for each family by estimating the standard deviation (σ) and the mean (μ) of the family's offspring trait values, where CV = σ/μ. The density is the number of families.

Genetic variance and correlations

There was overall significant genetic variance for body size (measured as area) for the 4 combinations of temperature and evolution regime (models with genetic variance were at least 7 DIC lower compared to models without genetic variance, Table 2). There was also significant genetic covariance between temperatures for both evolution regimes (*fast temp. cycle*: model with covariance included was 2.04 DIC lower than model without covariance; *increased warming*: model with covariance was 2.79 DIC lower than model without covariance). However, the pairwise comparisons of the posterior distributions of heritability, evolvability and genetic correlations were not significantly different between the 4 different combinations of temperature and evolution regimes (all 95% credibility intervals of the pairwise differences contained zero).

Table 2. Heritability (H^2), I^2 (Evolvability, in percentage) and genetic correlations for body size (area).

Experimental evol. regime	Temp. (°C)	Posterior mode H^2 (95% CI)	Posterior mode I^2 (95% CI)	Posterior mode genetic correlation across temperatures (95% CI)
Fast temp. cycles	20	0.59 (0.44, 0.70)	1.13 (0.56, 2.01)	0.20 (-0.10, 0.38)
	25	0.56 (0.42, 0.66)	1.30 (0.53, 2.67)	
Increased warming	20	0.50 (0.37, 0.65)	1.08 (0.57, 2.42)	0.20 (-0.12, 0.41)
	25	0.49 (0.42, 0.67)	1.54 (0.69, 2.97)	

Discussion

We found that evolution in an environment that changed in temperature every 2nd generation (*Fast temperature cycles* regime) resulted in the evolution of increased phenotypic plasticity in body size. In contrast, we did not find any evidence of increased diversifying bet hedging in this evolutionary regime, since there was neither increased phenotypic variance within families nor reduced heritability.

Evolution in variable environments with no environmental correlations across generations is predicted to result in increased importance of either phenotypic plasticity or bet-hedging (Tufto, 2015). While phenotypic plasticity should be favored when the environment contains predictable cues for development, bet-hedging should be favored instead in less predictable environments (Botero *et al.*, 2015; Tufto, 2015). Moreover, the timescale of environmental variation relative to the generation time is also important, and when modeled by Tufto, (2015), environmental changes every 2nd generation is identified as

476   the intersection between the evolution of bet-hedging, reversible plasticity or developmental

477   plasticity. Since the *Fast temperature cycle* regime experienced fluctuations every 2^{nd}

478   generation, they are ideally suited for investigating the evolution of plasticity and bet-hedging

479   in adult peak body size, an irreversible plastic trait closely connected to fitness.

480   We found evolution of increased phenotypic plasticity in the *Fast temperature cycle*

regime, manifested as a larger size difference between 20°C and 25°C (steeper reaction

norm). Evolution of increased phenotypic plasticity in more variable environments is

predicted by theory (Moran, 1992; Gavrilets & Scheiner, 1993; Kuijper & Hoyle, 2015), and

indeed studies using natural populations (Lind & Johansson, 2007; Lind *et al.*, 2011) or

species (Hollander, 2008) have found a positive correlation between the degree of

environmental heterogeneity and increased plasticity. Our study, using experimental

evolution, supports these results and pinpoint large temporal variation between temperature

extremes as the causative selection force underlying evolution of increased plasticity. This

has only been showed once before, in a recent experimental evolution study of the microalgae

Thalassiosira pseudonana, where populations evolving under fast fluctuations evolve

increased plasticity in photosynthesis (Schaum *et al.*, 2022). Our results also align with the

recent finding that laboratory-adapted populations of Zebra fish (*Danio rerio*), evolving in

very stable environments, have reduced physiological plasticity compared to their wild-

caught counterparts (Morgan *et al.*, 2022). Together, these studies demonstrate the

importance of environmental heterogeneity for the evolution of plasticity. It should however

be noted that very fast or unpredictable environmental change can make it impossible to

predict the environment, and in these circumstances plasticity may be selected against (Tufto,

2015), which was recently demonstrated using experimental evolution in the microalgae

Dunaliella salina (Leung *et al.*, 2020). Increased environmental variation is however not the

only factor that can influence the evolution of plasticity, but plasticity may also evolve when

a population is exposed to a novel (but stable) environment (Lande, 2009; Chevin *et al.*,

2010; Corl *et al.*, 2018). Evolution of increased plasticity has been shown for *C. remanei*

evolving in very heat-stressed environments (36.8°C), which demonstrates that evolution of

plasticity also can be a way to survive novel environment (Sikkink *et al.*, 2014b), a

phenomenon called plasticity-led evolution (Levis & Pfennig, 2020).

In contrast, we did not find any evolution of increased diversifying bet-hedging.

While empirical evidence for diversifying bet-hedging is much rarer than for phenotypic

plasticity, it is also much harder to detect since it is not trait means but trait variances that

needs to be measured. We have the power to detect differences in within-family CV where

the selection regimes explain at least 10% of the variation, which corresponds to a large

effect (Cohen, 1988). Still, there are a number of examples of bet-hedging, mainly regarding

delayed germination in desert plants (Philippi, 1993; Clauss & Venable, 2000; Venable,

2007), but also diapause in fish from ephemeral pools (Furness *et al.*, 2015). In addition,

diversifying bet-hedging has also evolved as a result of experimental evolution in

unpredictable environments in bacteria (Beaumont *et al.*, 2009) and fungi (Graham *et al.*,

2014). However, as predicted by Bull, (1987), a common factor in these examples is the very

strong fitness differences between environments, where one environmental state (for example

dry conditions) results in very low fitness. This contrasts to most examples of phenotypic

plasticity, where reproduction is possible to achieve in all environments, even if they are not

suitable without plastic adjustment of the phenotype. It has however been found that natural

isolates of *C. elegans* differ in their variance of total reproduction, which has been suggested

to result from a bet-hedging strategy where lines with more phenotypic variance may

originate from more variable environments (Diaz & Viney, 2014). We did not find any

difference in reproductive variance between our selection regimes, so further studies are

needed to understand why natural *C. elegans* isolates differ in their phenotypic variance of

reproduction.

When exposed to increasing temperatures, organisms generally show phenotypic

plasticity in size, and develop faster to mature smaller. Although there are exceptions,

especially in terrestrial arthropods (Horne *et al.*, 2015), this relationship is so general that it is

termed the temperature-size rule (Atkinson, 1994). Therefore, it is not surprising that

plasticity in size was present in both evolutionary regimes, and like *C. elegans* (Kammenga *et*

al., 2007), we find that *C. remanei* follows the temperature-size rule. Importantly however,

the regimes differed in the degree of size plasticity.

Whether the temperature-size rule (Atkinson, 1994) reflects an adaptive or non-

adaptive response to temperature is not resolved. Arguments for it being non-adaptive center

around constraints related to passive by-products of other temperature dependent processes

(Atkinson, 1994; van der Have & de Jong, 1996; Forster *et al.*, 2011). For example, small

body size can be a result of reduced growth rates which can be selected for as a by-product of

increased reproductive investment in warmer temperatures (Heino & Kaitala, 1999;

Walczyńska *et al.*, 2015), or, alternatively, as a trade-off between growth rate and resistance

to oxidative stress (Kim *et al.*, 2011) which increases due to increased metabolism in higher

temperatures (Birnie-Gauvin *et al.*, 2017).

However, small size in warm environments can also be adaptive. For example Fryxell

et al., (2020) showed that natural selection in warm temperatures favors smaller size in

mosquitofish *Gambusia affinis* and a similar result has been reported in snails (Arendt, 2015).

A possible advantage of small size in high temperatures can be a regulation of oxygen

demand and supply ratio (Walczyńska *et al.*, 2015). In addition, as a body composed of small

cells is more efficient in oxygen diffusion (Subczynski *et al.*, 1989; Verberk *et al.*, 2021)

there will be a particularly strong selection pressure on organisms such as *Caenorhabditis*

nematodes, which have a fixed number of cells and thus the cell size determines the final

body size.

To assess whether temperature-induced plasticity in size is adaptive in *C. remanei*, we

compared individual fitness of different-sized individuals in both temperatures (Figure 2).

We found evidence of directional selection on increased size in both temperatures, but also

significant stabilizing selection within each temperature. Stabilizing selection implies that the

fitness optimum in both 20°C and 25°C was present in individuals within the data size-range

(as opposed to at extreme phenotypes). If small size in warm temperatures were maladaptive,

we would expect the largest individuals to have the highest fitness. In contrast, we found that

individuals both smaller and larger than the optimum size had decreased fitness. This

optimum size in the warm temperature was also substantially smaller than the optimum size

at the normal temperature, thus the plastic response to decrease size as a response to warm

temperature must be considered adaptive in *C. remanei*.

Interestingly, because most individuals raised in 25°C exhibit smaller size than would

be optimal (Figure 2.; mean size is smaller than optimal size), we can consider this

temperature plasticity to be a hyperplastic response, a special case of plasticity when plastic

response overshoots the optimum and brings individuals to the other side of the new fitness

peak (King & Hadfield, 2019). Since plasticity nevertheless increases fitness (compared to a

hypothetical non-plastic genotype), this hyperplasticity should still be considered adaptive. In

addition, we also found linear selection for large size in 20°C with individuals raised in 20°C

also having slightly smaller size than would be optimal. A possible explanation is a sexual

conflict between male and female worms, as males' optimal size is smaller than females'

optimal size in *C. remanei* (Stångberg *et al.*, 2020) so that males drag females from their

phenotypic optimum.

In contrast to size, we didn't find any difference in individual fitness between the

regimes. While warm temperature caused a drop in total reproduction in both regimes,

individuals raised in 25°C had significantly higher rate-sensitive individual fitness, which is a

consequence of the temperature-induced alteration of the reproductive schedule, including a

faster development time (Sekajova *et al.*, 2022).

Previous experimental evolution and artificial selection studies in the SP8 line of *C.*

remanei, which was our founder population, have documented fast evolutionary responses to

selection in life history traits, suggesting a large amount of standing genetic variation (Chen

& Maklakov, 2012; Zwoinska *et al.*, 2016; Lind *et al.*, 2020). Accordingly, we found

substantial genetic variation for size, for all treatment × temperature combinations, which not

only allowed the lines to respond to selection, but also represents a potential for further

evolution. Since we used full-sibs, our estimates of genetic variance could potentially be

inflated by the presence of dominance variance and epistatic interactions. Epistatic

interactions are present for body size in the sister species *C. elegans* (Noble *et al.*, 2017;

Maulana *et al.*, 2022), but while most genetic variance for this trait was found to be additive

and with similar additive heritability to our estimate using an experimental evolution design

(Noble *et al.*, 2017), assessments of narrow-sense heritability using recombinant inbreed lines

found that additive effects played a smaller role for body-size (Maulana *et al.*, 2022).

Therefore, we assume that our broad sense heritability overestimates the additive genetic

effect, but to an unknown degree. Moreover, we didn't observe any significant differences in

broad-sense heritability between the lines, which further support our evidence of no evolution

of diversified bet-hedging, which comes with the prediction of lowered heritability in traits

(Tufto, 2015), nor did we observe any genetic correlations between trait values in the two

temperatures, suggesting that traits can largely evolve independently in each environment.

Conclusion

To summarize, we found that 30 generations of experimental evolution in a heterogeneous environment (*Fast temperature cycles)* resulted in the evolution of increased phenotypic plasticity, compared to evolution in a slowly changing environment (*Increased warming)*. We showed that plasticity followed the temperature size rule and was adaptive. In addition, substantial amount of standing genetic variation found in the line represents a potential for further evolutionary change.

References

Andersson, M. (1994) *Sexual selection*. Monographs in Behavior and Ecology. Princeton University Press, Princeton, New Jersey.

Arendt, J. (2015) Why get big in the cold? Size–fecundity relationships explain the temperature-size rule in a pulmonate snail (Physa). *Journal of Evolutionary Biology*, **28**, 169–178.

Atkinson, D. (1994) Temperature and organism size: a biological law for ectotherms? *Advances in ecological research*, **25**, 1–58.

Bates, D., Mächler, M., Bolker, B.M. & Walker, S.C. (2015) Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, **67**, 1–48.

Beaumont, H.J.E., Gallie, J., Kost, C., Ferguson, G.C. & Rainey, P.B. (2009) Experimental evolution of bet hedging. *Nature*, **462**, 90–93.

Berteaux, D., Réale, D., McAdam, A.G. & Boutin, S. (2004) Keeping pace with fast climate change: can arctic life count on evolution? *Integrative and Comparative Biology*, **44**, 140–151.

Birnie-Gauvin, K., Costantini, D., Cooke, S.J. & Willmore, W.G. (2017) A comparative and evolutionary approach to oxidative stress in fish: A review. *Fish and Fisheries*, **18**, 928–942.

Botero, C.A., Weissing, F.J., Wright, J. & Rubenstein, D.R. (2015) Evolutionary tipping

points in the capacity to adapt to environmental change. *Proceedings of the National*

Academy of Sciences, **112**, 184–189.

Brommer, J.E., Merilä, J. & Kokko, H. (2002) Reproductive timing and individual fitness.

Ecology Letters, **5**, 802–810.

Bull, J.J. (1987) Evolution of phenotypic variance. *Evolution*, **41**, 303–315.

Burton, T., Ratikainen, I.I. & Einum, S. (2022) Environmental change and the rate of

phenotypic plasticity. *Global Change Biology*, **28**, 5337–5345.

Campbell-Staton, S.C., Velotta, J.P. & Winchell, K.M. (2021) Selection on adaptive and

maladaptive gene expression plasticity during thermal adaptation to urban heat islands.

Nature Communications, **12**, 6195.

Chen, H. & Maklakov, A.A. (2012) Longer life span evolves under high rates of condition-

dependent mortality. *Current Biology*, **22**, 2140–2143.

Chevin, L.-M., Lande, R. & Mace, G.M. (2010) Adaptation, plasticity, and extinction in a

changing environment: towards a predictive theory. *PLoS Biol*, **8**, e1000357.

Clauss, M.J. & Venable, D.L. (2000) Seed germination in desert annuals: an empirical test of

adaptive bet hedging. *The American Naturalist*, **155**, 168–186.

Cohen, D. (1966) Optimizing reproduction in a randomly varying environment. *Journal of*

Theoretical Biology, **12**, 119–129.

Cohen, J. (1988) *Statistical power analysis for the behavioral sciences*. 2nd edition.

Routledge, Hillsdale, N.J.

Corl, A., Bi, K., Luke, C., Challa, A.S., Stern, A.J., Sinervo, B., *et al.* (2018) The genetic

basis of adaptation following plastic changes in coloration in a novel environment. *Current*

Biology, **28**, 2970-2977.e7.

DeWitt, T.J. & Scheiner, S.M. (2004) *Phenotypic plasticity: functional and conceptual*

approaches. Oxford University Press, New York, NY, USA.

Dey, S., Proulx, S.R. & Teotónio, H. (2016) Adaptation to temporally fluctuating environments by the evolution of maternal effects. *PLoS Biology*, **14**, e1002388.

Diaz, S.A. & Viney, M. (2014) Genotypic-specific variance in *Caenorhabditis elegans* lifetime fecundity. *Ecology and Evolution*, **4**, 2058–2069.

Dupont, L., Thierry, M., Zinger, L., Legrand, D. & Jacob, S. (2024) Beyond reaction norms: the temporal dynamics of phenotypic plasticity. *Trends in Ecology & Evolution*, **39**, 41–51.

Easterling, D.R., Meehl, G.A., Parmesan, C., Changnon, S.A., Karl, T.R. & Mearns, L.O. (2000) Climate extremes: observations, modeling, and impacts. *Science*, **289**, 2068–2074.

Forster, J., Hirst, A.G. & Woodward, G. (2011) Growth and development rates have different thermal responses. *The American Naturalist*, **178**, 668–678.

Fryxell, D.C., Hoover, A.N., Alvarez, D.A., Arnesen, F.J., Benavente, J.N., Moffett, E.R., *et al.* (2020) Recent warming reduces the reproductive advantage of large size and contributes to evolutionary downsizing in nature. *Proceedings of the Royal Society B: Biological Sciences*, **287**, 20200608.

Furness, A.I., Lee, K. & Reznick, D.N. (2015) Adaptation in a variable environment: Phenotypic plasticity and bet-hedging during egg diapause and hatching in an annual killifish. *Evolution*, **69**, 1461–1475.

Gavrilets, S. & Scheiner, S.M. (1993) The genetics of phenotypic plasticity. V. Evolution of reaction norm shape. *Journal of Evolutionary Biology*, **6**, 31–48.

Geerts, A.N., Vanoverbeke, J., Vanschoenwinkel, B., Van Doorslaer, W., Feuchtmayr, H., Atkinson, D., *et al.* (2015) Rapid evolution of thermal tolerance in the water flea *Daphnia*. *Nature Climate Change*, **5**, 665–668.

Ghalambor, C.K., Hoke, K.L., Ruell, E.W., Fischer, E.K., Reznick, D.N. & Hughes, K.A. (2015) Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in nature. *Nature*, **525**, 372–375.

Graham, J.K., Smith, M.L. & Simons, A.M. (2014) Experimental evolution of bet hedging under manipulated environmental uncertainty in *Neurospora crassa*. *Proceedings of the Royal Society of London B: Biological Sciences*, **281**, 20140706.

Hadfield, J. (2010) MCMC methods for multi-response generalized linear mixed models: the
MCMCglmm R package. *Journal of Statistical Software*, **33**, 1–22.

Hansen, T.F., Pélabon, C., Armbruster, W.S. & Carlson, M.L. (2003) Evolvability and
genetic constraint in *Dalechampia* blossoms: components of variance and measures of
evolvability. *Journal of Evolutionary Biology*, **16**, 754–766.

Hansen, T.F., Pélabon, C. & Houle, D. (2011) Heritability is not evolvability. *Evolutionary*
Biology, **38**, 258–277.

Have, T.M. van der & Jong, G. de. (1996) Adult size in ectotherms: temperature effects on
growth and differentiation. *Journal of Theoretical Biology*, **183**, 329–340.

Hedrick, A.V. & Temeles, E.J. (1989) The evolution of sexual dimorphism in animals:
Hypotheses and tests. *Trends in Ecology & Evolution*, **4**, 136–138.

Heino, M. & Kaitala, V. (1999) Evolution of resource allocation between growth and
reproduction in animals with indeterminate growth. *Journal of Evolutionary Biology*, **12**,
423–429.

Hollander, J. (2008) Testing the grain-size model for the evolution of phenotypic plasticity.
Evolution, **62**, 1381–1389.

Horne, C.R., Hirst, Andrew.G. & Atkinson, D. (2015) Temperature-size responses match
latitudinal-size clines in arthropods, revealing critical differences between aquatic and
terrestrial species. *Ecology Letters*, **18**, 327–335.

Kammenga, J.E., Doroszuk, A., Riksen, J.A.G., Hazendonk, E., Spiridon, L., Petrescu, A.-J.,
et al. (2007) A *Caenorhabditis elegans* wild type defies the temperature–size rule owing to a
single nucleotide polymorphism in tra-3. *PLoS Genet*, **3**, e34.

Kim, S.-Y., Noguera, J.C., Morales, J. & Velando, A. (2011) Quantitative genetic evidence
for trade-off between growth and resistance to oxidative stress in a wild bird. *Evolutionary*
Ecology, **25**, 461–472.

King, J.G. & Hadfield, J.D. (2019) The evolution of phenotypic plasticity when environments
fluctuate in time and space. *Evolution Letters*, **3**, 15–27.

Kuijper, B. & Hoyle, R.B. (2015) When to rely on maternal effects and when on phenotypic

plasticity? *Evolution*, **69**, 950–968.

Kussell, E. & Leibler, S. (2005) Phenotypic diversity, population growth, and information in

fluctuating environments. *Science*, **309**, 2075–2078.

Lachmann, M. & Jablonka, E. (1996) The inheritance of phenotypes: an adaptation to

fluctuating environments. *Journal of Theoretical Biology*, **181**, 1–9.

Lande, R. (2009) Adaptation to an extraordinary environment by evolution of phenotypic

plasticity and genetic assimilation. *Journal of Evolutionary Biology*, **22**, 1435–1446.

Leimar, O. & McNamara, J.M. (2015) The evolution of transgenerational integration of

information in heterogeneous environments. *The American Naturalist*, **185**, E55–E69.

Leung, C., Rescan, M., Grulois, D. & Chevin, L.-M. (2020) Reduced phenotypic plasticity

evolves in less predictable environments. *Ecology Letters*, **23**, 1664–1672.

Levis, N.A. & Pfennig, D.W. (2020) Plasticity-led evolution: A survey of developmental

mechanisms and empirical tests. *Evolution & Development*, **22**, 71–87.

Lind, M.I., Ingvarsson, P.K., Johansson, H., Hall, D. & Johansson, F. (2011) Gene flow and

selection on phenotypic plasticity in an island system of *Rana temporaria*. *Evolution*, **65**,

684–697.

Lind, M.I. & Johansson, F. (2007) The degree of phenotypic plasticity is correlated with the

spatial environmental heterogeneity experienced by island populations of *Rana temporaria*.

Journal of Evolutionary Biology, **20**, 1288–1297.

Lind, M.I., Zwoinska, M.K., Andersson, J., Carlsson, H., Krieg, T., Larva, T., *et al.* (2020)

Environmental variation mediates the evolution of anticipatory parental effects. *Evolution*

Letters, **4**, 371–381.

Lind, M.I., Zwoinska, M.K., Meurling, S., Carlsson, H. & Maklakov, A.A. (2016) Sex-

specific trade-offs with growth and fitness following lifespan extension by rapamycin in an

outcrossing nematode, *Caenorhabditis remanei*. *Journals of Gerontology Series A:*

Biological Sciences and Medical Sciences, **71**, 882–890.

Maulana, M.I., Riksen, J.A.G., Snoek, B.L., Kammenga, J.E. & Sterken, M.G. (2022) The

genetic architecture underlying body-size traits plasticity over different temperatures and

developmental stages in *Caenorhabditis elegans*. *Heredity*, **128**, 313–324.

McGraw, J.B. & Caswell, H. (1996) Estimation of individual fitness from life-history data.

The American Naturalist, **147**, 47–64.

Merilä, J. & Hendry, A.P. (2014) Climate change, adaptation, and phenotypic plasticity: the

problem and the evidence. *Evolutionary Applications*, **7**, 1–14.

Meyers, L.A. & Bull, J.J. (2002) Fighting change with change: adaptive variation in an

uncertain world. *Trends in Ecology & Evolution*, **17**, 551–557.

Moran, N.A. (1992) The evolutionary maintenance of alternative phenotypes. *American

Naturalist*, **139**, 971–989.

Morgan, R., Andreassen, A.H., Åsheim, E.R., Finnøen, M.H., Dresler, G., Brembu, T., *et al.*

(2022) Reduced physiological plasticity in a fish adapted to stable temperatures. *Proceedings

of the National Academy of Sciences*, **119**, e2201919119.

Mylius, S.D. & Diekmann, O. (1995) On evolutionarily stable life histories, optimization and

the need to be specific about density dependence. *Oikos*, **74**, 218–224.

Noble, L.M., Chelo, I., Guzella, T., Afonso, B., Riccardi, D.D., Ammerman, P., *et al.* (2017)

Polygenicity and epistasis underlie fitness-proximal traits in the *Caenorhabditis elegans*

multiparental experimental evolution (CeMEE) panel. *Genetics*, genetics.300406.2017.

Palmer, T.N. & Räisänen, J. (2002) Quantifying the risk of extreme seasonal precipitation

events in a changing climate. *Nature*, **415**, 512–514.

Partridge, L. & Coyne, J.A. (1997) Bergmann's rule in ectotherms: is it adaptive? *Evolution*,

51, 632–635.

Philippi, T. (1993) Bet-hedging germination of desert annuals: variation among populations

and maternal effects in *Lepidium lasiocarpum*. *The American Naturalist*, **142**, 488–507.

Philippi, T. & Seger, J. (1989) Hedging one's evolutionary bets, revisited. *Trends in Ecology

& Evolution*, **4**, 41–44.

R Core Team. (2015) R: A language and environment for statistical computing.

Ray, C. (1960) The application of Bergmann's and Allen's rules to the poikilotherms.
Journal of Morphology, **106**, 85–108.

Schaum, C.-E., Buckling, A., Smirnoff, N. & Yvon-Durocher, G. (2022) Evolution of
thermal tolerance and phenotypic plasticity under rapid and slow temperature fluctuations.
Proceedings of the Royal Society B: Biological Sciences, **289**, 20220834.

Sekajova, Z., Rosa, E., Spagopoulou, F., Zervakis, P.-I. & Lind, M.I. (2022) Temperature-
induced compensatory growth in the nematode *Caenorhabditis elegans* is regulated by a
thermosensitive TRP channel and influences reproductive rate. *Functional Ecology*, **36**,
2176–2187.

Sikkink, K.L., Ituarte, C.M., Reynolds, R.M., Cresko, W.A. & Phillips, P.C. (2014a) The
transgenerational effects of heat stress in the nematode *Caenorhabditis remanei* are negative
and rapidly eliminated under direct selection for increased stress resistance in larvae.
Genomics, Experimental evolution and the use of genomics, **104**, 438–446.

Sikkink, K.L., Reynolds, R.M., Ituarte, C.M., Cresko, W.A. & Phillips, P.C. (2014b) Rapid
evolution of phenotypic plasticity and shifting thresholds of genetic assimilation in the
nematode *Caenorhabditis remanei. G3: Genes | Genomes | Genetics*, **4**, 1103–1112.

Simons, A.M. (2011) Modes of response to environmental change and the elusive empirical
evidence for bet hedging. *Proceedings of the Royal Society of London B: Biological Sciences*,
278, 1601–1609.

Slatkin, M. (1974) Hedging one's evolutionary bets. *Nature*, **250**, 704–705.

Stångberg, J., Immonen, E., Moreno, P.P. & Bolund, E. (2020) Experimentally induced
intrasexual mating competition and sex-specific evolution in female and male nematodes.
Journal of Evolutionary Biology, **33**, 1677–1688.

Stearns, S.C. (1992) *The evolution of life histories*. Oxford University Press, New York, NY,
USA.

Stiernagle, T. (2006) Maintenance of *C. elegans*. *WormBook: the online review of C. elegans*
biology.

Subczynski, W.K., Hyde, J.S. & Kusumi, A. (1989) Oxygen permeability of
phosphatidylcholine--cholesterol membranes. *Proceedings of the National Academy of*
Sciences, **86**, 4474–4478.

Teotónio, H., Estes, S., Phillips, P.C. & Baer, C.F. (2017) Experimental evolution with
Caenorhabditis nematodes. *Genetics*, **206**, 691–716.

Tufto, J. (2015) Genetic evolution, plasticity, and bet-hedging as adaptive responses to
temporally autocorrelated fluctuating selection: A quantitative genetic model. *Evolution*, **69**,
2034–2049.

Uller, T., English, S. & Pen, I. (2015) When is incomplete epigenetic resetting in germ cells
favoured by natural selection? *Proc. R. Soc. B*, **282**, 20150682.

Van Aalst, M.K. (2006) The impacts of climate change on the risk of natural disasters.
Disasters, **30**, 5–18.

Venable, D.L. (2007) Bet hedging in a guild of desert annuals. *Ecology*, **88**, 1086–1090.

Verberk, W.C.E.P., Atkinson, D., Hoefnagel, K.N., Hirst, A.G., Horne, C.R. & Siepel, H.
(2021) Shrinking body sizes in response to warming: explanations for the temperature–size
rule with special emphasis on the role of oxygen. *Biological Reviews*, **96**, 247–268.

Walczyńska, A., Labecka, A.M., Sobczyk, M., Czarnoleski, M. & Kozłowski, J. (2015) The
temperature–size rule in *Lecane inermis* (Rotifera) is adaptive and driven by nuclei size
adjustment to temperature and oxygen combinations. *Journal of Thermal Biology*, **54**, 78–85.

West-Eberhard, M.J. (2003) *Developmental plasticity and evolution*. Oxford University Press
Inc, USA.

Wolf, D.M., Vazirani, V.V. & Arkin, A.P. (2005) Diversity in times of adversity:
probabilistic strategies in microbial survival games. *Journal of Theoretical Biology*, **234**,
227–253.

Zwoinska, M.K., Lind, M.I., Cortazar-Chinarro, M., Ramsden, M. & Maklakov, A.A. (2016)
Selection on learning performance results in the correlated evolution of sexual dimorphism in
life-history. *Evolution*, **70**, 342–357.