1

Circadian regulation of Homologous Recombination by Cryptochrome1-mediated

dampening of DNA end resection

One sentence summary: CCAR2-dependent inhibition of DNA end resection and

homologous recombination is controlled by the intrinsic cellular circadian clock protein

CRY1

Amador Romero-Franco[1,2], Cintia Checa-Rodríguez[1,2]#[†], Sonia Jimeno[1,2†], Maikel

Castellano-Pozo[1,2†], Paula Aguilera[2], Hector Miras[3], Amadeo Wals[3], Silvia Jimeno-

González[1,2], Andres Joaquin Lopez-Contreras[2], and Pablo Huertas[1, 2,]*

[1] Facultad de Biología, Universidad de Sevilla, Sevilla, 41080, Spain

[2] Centro Andaluz de Biología Molecular y Medicina Regenerativa-CABIMER,

Universidad de Sevilla-CSIC-Universidad Pablo de Olavide**,** Sevilla, 41092, Spain

[3] Servicios de Oncología Radioterápica y de Radiofísica, Hospital Universitario Virgen

Macarena, Sevilla, Spain

Present address: Instituto de Biomedicina de Sevilla-IBiS, Hospital Universitario

Virgen del Rocío/CSIC/Universidad de Sevilla, Sevilla, 41013, Spain

*To whom correspondence should be addressed. Tel: +34 954 467 667; Fax: +34 954

461 664; Email: phuertas@us.es

† These three authors contributed equally to this work

Lead Contact: Pablo Huertas

Keywords: DNA end resection/DNA repair/Homologous recombination/Circadian

clock/CCAR2/CRY1

Abstract

Genomic stability maintenance requires the repair of DNA breaks in the most accurate

fashion. So, an exquisite regulatory network controls the choice between different repair

mechanisms to maximize genome integrity. This relies mostly at the level of DNA end

resection, the initial steps of the homologous recombination. On the other hand, most

cellular activities follow a 24 h oscillation known as the circadian cycle. Here we

explore the regulation of the choice between different DNA break repair pathways

along the circadian cycle. We observed that in human cells DNA resection shows a

circadian oscillation, with a peak at early morning followed by a progressive reduction

until late afternoon. Such regulation depends on the circadian clock core component

CRY1, which modulates the anti-resection activity of CCAR2 to limit CtIP at nightfall.

Additionally, such regulation requires DNA-PK-mediated phosphorylation of CRY1.

Finally, this circadian regulation impacts cancer progression and response to radiation

therapy of specific tumours.

INTRODUCTION

DNA double-strand break (DSB) repair is a critical aspect of cellular biology

that has a great impact in different human diseases such as cancer[1]. DSBs can be

channelled through different repair pathways, with distinctive outcomes. Such pathways

belong, mostly, to two different families: homology-driven repair (HDR), that exploits

the presence of intact copies of the broken sequences as a template; and non-

homologous end-joining (NHEJ), the simple ligation of two DNA ends with little or no

processing[2–4]. Both can result in either faithful or mistaken repair, thus the choice

between them is tightly regulated. The best characterized regulatory step for DSB repair

pathway choice is the activation of DNA end resection, a degradation of one strand of

the broken DNA to create long tails of 3' protruding ssDNA[5,6]. Resection regulation

relies mostly on the control of the core resection factor CtIP trough specific interactions

and/or post-translational modifications[7,8]. Once CtIP is activated, resection ensues and

HDR prevails. If CtIP is blocked, resection is hampered and HDR is blocked. CCAR2

functions as a CtIP antagonist in resection regulation[9], acting as a downstream effector

of the Shieldin complex[10]. Both proteins interact constitutively, and only when CCAR2

is evicted from damaged chromatin, CtIP is unleashed, thus favouring resection[9].

Circadian rhythms are a series of periodic oscillations that follow the day/night

cycle[11]. Its presence is conserved throughout the whole photosensitive spectrum of

living organisms[12]. The circadian clock works as a built-in 24-hour oscillator that acts in

a two-tier fashion: an intrinsic cellular circadian circuitry with a roughly 24 h period

that can be synchronized in response to the external light, a process known as

photoentrainment[11]. In mammals, at the molecular level, the transcription factors

BMAL1 and CLOCK stimulate the expression of the genes that code for the Period

(PER1-3) and Cryptochrome (CRY1-2) families during the activation phase that, in

humans, coincides with daylight hours. Then, PER and CRY heterodimers translocate

into the nucleus where CRY negatively affect the transcriptional activity of

BMAL1/CLOCK, mainly during night hours[13]. Thus, a recurrent oscillation is created

by sequential transcriptional activation/repression waves. Additional accessory loops

have been characterized in the circadian clock, but the heterodimers BMAL1/CLOCK

and PER/CRY build the core cellular oscillator[14]. Importantly, the robustness of the

circadian cycle varies among tissues, but it is also affected by the age of the individuals

and even shows sex-dimorphic characteristics[15]. For simplicity, in this manuscript we

will refer to the terms "sunrise"/"dawn"/"early morning" or "sunset"/"dusk"/"late

afternoon" that correspond to an ideal circadian cycle of 12 h day followed by a 12 h night, i.e. roughly around 6-8 am or 6-8 pm, respectively. However, it is important to note that during the year, and especially closer to the poles, the real "sunrise"/"sunset" will take place at very different times.

Here we show that DNA end resection and HDR, are regulated in a circadian fashion. Indeed, resection activity peaks at "dawn" and slowly decreases during the day, followed by an increase during the night. Such regulation relies specifically and directly on the circadian factor CRY1. Mechanistically, CRY1 is recruited to sites of DSBs and reinforces the inhibitory effect of CCAR2 over CtIP by potentiating CCAR2 retention on damaged chromatin. Furthermore, DNA-PK finetunes this circadian effect. So, when CRY1 is recruited to damaged chromatin is phosphorylated by DNA-PK, increasing its retention at DSBs. This locks CCAR2 there, curbing CtIP activity. This negative modulation maximizes around nightfall in human cells, when CRY1 levels peak, whereas the absence of this protein at "dawn" breaks this regulatory circuit, hence licencing DNA end processing. Finally, this fine regulation of end resection affects the ability of cells to deal with DNA damaging agents in cellular models and mice Xenografts, but also in cancer clinical setups.

RESULTS

DNA end resection shows a circadian synchrony

As mentioned, all animal cells have a built-in circadian oscillator. Although *in vitro* grown cell cultures cannot be photoentrained, the addition of different drugs can reset them[16,17]. We synchronized U2OS cells with either Dexamethasone (DEX) or Forskolin (FSK) for two hours, resetting the internal clock to the equivalent of "early morning" (Fig.1A), i.e when CRY levels are lower. Then, we irradiated them to induce DSBs at fixed time points after release, that represent different times of the day (Fig. 1A). A control treated with the vehicle, i.e. an asynchronous population, was prepared in parallel. Circadian synchronization was followed by the levels of CRY1 and CRY2, that peak at "late afternoon" (Suppl. Fig. 1A for DEX). Indeed, such peak was due to circadian synchronization, as it was absent in the control experiment treated with the vehicle alone (Suppl. Fig. 1B). Then, we analysed DNA end resection by checking the formation of RPA foci by immunofluorescence upon 10 Gy of irradiation. U2OS cells synchronized with DEX (blue line) showed a marked oscillation in resection proficiency, clearly distinct to the observed in the control treated with the vehicle EtOH (red line Fig.

1B; see Suppl. Fig. 1C for representative images). Whereas at noon and midnight the number of cells actively resecting was like the control, a marked reduction was readily observed at "late afternoon". Moreover, at "early morning" resection was slightly elevated when compared with the asynchronous culture. Similar results were observed by adding FSK when compared with the mock treatment (Fig. 1C, Suppl. Fig. 1D). Although these results agreed with a circadian synchronization of DNA end resection, it has been described that during the first 24h after release from DEX there are some effects that might be mediated by immediate-early/early gene expression and not due to a true circadian effect. To clarify this issue, we repeated the RPA foci formation experiment upon DEX synchronization and release at different time points during 48 h. As shown in Fig. 1D (see Suppl. Fig. 1E for representative images and Suppl. Table 1 for statistical analysis), the same pattern was observed both during the first and second days upon release, strengthening the idea that resection is more proficient during the times that correspond to "early morning" and reduced at the time that correspond to "late afternoon". Indeed, as expected resection patterns behave opposite to the accumulation of the circadian protein CRY1, measured by immunofluorescence (Fig. 1E, Suppl. Table 1, Suppl. Fig. 1F) or western blot (Suppl. Fig. 1G).

Furthermore, this oscillation was also evident in the primary human cell line RPE1 (Fig. 1F, Suppl. Fig. 1H) and in mouse embryonic fibroblasts (MEFs; Fig. 1G, Suppl. Fig. 1I). From this point onward, only a single asynchronous sample was taken and represented as a dashed line. Cell cycle is a major regulator of DNA end resection[5–7,18] and both the circadian and the cell cycles are known to crosstalk[19]. However, our results could not be attributed to changes in cell cycle, as it was not affected by circadian synchronization (Suppl. Fig. 1J-L). To determine if the circadian rhythm controlled only the initiation of DNA end resection or if, instead, affected also the processivity of the process, we measured the actual length of resected DNA tracts at different positions on the circadian cycle (Fig. 1H). At "early morning", in agreement with a more permissive environment for resection, the length of DNA resected was longer than the asynchronous control. On the contrary, at "late afternoon" the extension of the resected DNA greatly diminished. Resection is controlled by multiple signals[5,6], chiefly the antagonistic relationship between the pro-resection BRCA1 and the anti-resection complex Shieldin[20–22]. So, we then analysed the recruitment of BRCA1 and the Shieldin subunit RIF1 1h after the irradiation at different times of the circadian cycle. In agreement with our previous results, BRCA1 was more avidly recruited when

DNA damage occurs at "early morning" whereas its localization was limited at "late afternoon" (Fig. 1I; Suppl. Fig 1M). RIF1 showed the opposite recruitment pattern (Fig. 1J; Suppl. Fig 1N). Finally, breaks induced at "dusk" were less likely to engage the recombinase RAD51 than those created at "dawn" (Fig. 1K; Suppl. Fig 1O). Thus, we conclude that the circadian cycle in humans creates a more permissive environment for resection early in the morning, but it is partially dampened at "late afternoon".

DNA end resection reacts to CRY1 levels

We took advantage of our previously published results in which we tested the effect of depletion of all human genes in the balance between HDR and NHEJ[23]. Interestingly, from the core and accessory circadian regulators, only CRY1 depletion skewed the balance between DSB repair pathways in a way that was compatible with our observations. Thus, we tested if resection was reacting directly to CRY1 levels. In agreement with a resection increase at "early morning", depletion of CRY1 stimulates the process and such effect was complemented by ectopic CRY1 expression (Fig. 2A; Suppl. Fig. 2A for depletion levels). Similar results were observed in RPE1 and MEFs (Suppl. Fig. 2B-C) and upon a CRISPR-mediated knockout of CRY1 in RPE1 cells (Suppl. Fig 2D). Strikingly, it is known that CRY1 loss does not abolish the circadian cycle, as its role is taken over by its paralog CRY2, but only shortens it by about 1 h[24]. Thus, the fact that CRY1 depletion shows such a marked effect in resection stimulation strongly suggest that is specifically CRY1, and not the rest of the circadian built-in clock, what modulates this process. In agreement, CRY2 depletion did not cause an RPA foci defect whereas depletion of any PER paralogue showed a reduction in DNA end resection (Suppl. Fig. 2E), reinforcing this idea. But more importantly, CRY1 depletion completely abolished the circadian oscillation of RPA foci formation (Fig. 2B). Interestingly, in the context of prostate cancer, it has been shown that CRY1 acts independently of the rest of the CRY-PER complex regulating DNA repair through controlling the stability of DNA repair factors[25]. To confirm the effect of CRY1 depletion on RPA recruitment upon DSBs formation by different means, we took advantage of the DiVA system, in which AsiSI-mediated restriction creates breaks upon tamoxifen addition[26]. ChIP-seq analysis of RPA recruitment on the best 214 AsiSI cutting sites, as previously described by BLISS[27], showed a stronger accumulation of RPA upon DSB formation in CRY1 depleted cells when compared with control cells (Fig. 2C).

Of note, this effect was underestimated, as CRY1 depletion increased the
number of G1 cells (Suppl. Fig. 2F-G), a position on the cell cycle that is refractory to
resection. To minimize this cell cycle effect, we repeated the immunofluorescence and
analysed RPA foci determining the cell cycle phases by QIBC, using DAPI (DNA
content) and EdU (DNA replication) as surrogates for cell cycle progression. CRY1
depletion increased the average number of RPA foci (above the horizontal dashed line)
in S/G2 cells (right of the vertical dashed line) (Fig. 2D). Overexpression of CRY1 in
cells transfected with a control siRNA reduced the amount of RPA foci upon DNA
damage (Fig. 2A). We repeated the experiment in cells overexpressing human CRY1
but not transfected with any siRNA (Fig. 2E; Suppl. Fig 2H for representative images
and 2I for expression levels) or treated with KL001, a drug that stabilizes CRY1[28] (Fig.
2F; Suppl. Fig. 2J for expression levels and 2K for representative images). In both cases,
resection was dampened. We conclude that we can artificially mimic the "early morning"
or "late afternoon" condition by either depleting CRY1 or overexpressing/stabilizing
CRY1, respectively. We took advantage of this setup to test if DSB repair pathways
efficiency were affected by CRY1 levels. Indeed, low levels of CRY1 stimulate both
RAD51-dependent and -independent homologous recombination (Fig. 2G-H).
Strikingly, this did not cause a reduction in NHEJ (Fig. 2I). Conversely, high levels of
CRY1 hampered homology directed repair (Fig. 2J-K), but again did not affect the bulk
of NHEJ (Fig. 2L). In agreement with this effect in recombination, RAD51 recruitment
to DSB readily react to CRY1 levels (Fig. 2M-N and Suppl. Fig. 2L-M). The foci
formation of the Shieldin complex 1 h after DNA damage induction was also affected
by CRY1 expressions levels (Fig. 2O-P and Suppl. Fig. 2N-O), with an opposite pattern
of RPA foci. In order to understand why the decrease in RIF1 formation might not be
reflected in a decrease in NHEJ we performed kinetic study of RIF1 foci formation. We
discovered that RIF1 recruitment was not really blocked, but delayed, in cells
downregulated for CRY1 in a similar fashion to the depletion of the anti-resection factor
CCAR2 (Fig. 2Q). This suggested that stimulating resection efficiency mostly delay
NHEJ, but such changes do not reflect in a large reduction of the bulk NHEJ. In
agreement, depletion of CRY1 does not abolish the recruitment of 53BP1 (Fig. 2R) or
the bonafide NHEJ factor KU70 to damaged chromatin (Fig. 2S). Hence, it seems that
only the anti-resection factor RIF1 is affected by CRY1, whereas the core NHEJ is
unaffected.
34

CRY1 regulates the antagonist relationship between CtIP and CCAR2

We tested the effect of the depletion of the core resection factor CtIP in cells upon circadian synchronization. As expected, CtIP depletion reduced resection efficiency when compared with siRNA control transfected cells (Fig. 3A; Suppl. Fig 3A for representative images and Suppl. Fig. 3B for statistics). Strikingly, CtIP downregulation not only decreased resection, but mostly abolished the circadian oscillations of the process, suggesting that such regulation might rely on controlling this protein. In fact, at "late afternoon", resection was limited to a degree compatible with an inhibition of CtIP.

CtIP is known to suffer a tight regulation at many different levels, including protein levels, mRNA stability, post-translational modifications, but also by the direct interaction with multiple proteins, including the antagonist factor CCAR2[7,8,23]. We reasoned that an inhibition of resection at "late afternoon" would agree with a peak of activity of this protein. Strikingly, CCAR2 is already linked with the regulation of circadian rhythms[29,30]. CCAR2 depletion also abolished circadian oscillation of DNA end processing, but in this case allowing fully active resection at any given time of the day/night cycle (Fig. 3A; Suppl. Fig. 3A-B). Importantly, despite many circadian-mediated regulation happening by controlling the transcription of target genes, and a role of CRY1 regulating DDR factors expression in prostate cancer[25], neither CCAR2 nor CtIP protein levels change during the circadian cycle (Suppl. Fig. 3C). CCAR2 depletion was completely epistatic over both CRY1 depletion and CRY1 overexpression for RPA and BRCA1 foci formation, suggesting that CRY1 acts fully through stimulation of CCAR2 (Fig. 3B-D; Suppl. Fig. 3D-F). This epistatic effect cannot be attributed to large cell cycle changes (Suppl. Fig. 3G). This fits with the idea that is CRY1 itself, and not the whole intrinsic circadian clock, what controls resection and recombination. Thus, we wondered if CRY1 protein might directly interact with CCAR2. Indeed, both proteins physically interact constitutively, i.e. regardless of the exposure to IR (Fig. 3E). CCAR2 and CtIP also interact constitutively[23]. Indeed, we could co-immunoprecipitate CtIP and CRY1 by reciprocal IPs (Fig. 3F-G) and detect their proximity by Proximity Ligation Assay (PLA; Suppl. Fig. 3H). Furthermore, CtIP depletion reduced the interaction between the other two proteins (Fig. 3H), suggesting that all three proteins might interact together.

Considering this interaction, we wondered if CRY1 was recruited to sites of chromosome breaks as it has been described for both CtIP and CCAR2[23]. Using a GFP-

CRY1 we observed a fast accumulation of this protein at DSBs by laser microirradiation with an even faster kinetic than the early DDR factor MDC1 (Fig. 3I; Suppl. movies 1 and 2). We previously showed that CCAR2 interacts with CtIP, and upon the appearance of DSBs controls CtIP recruitment and spreading from the breaks[23]. In less permissive conditions for resection, CCAR2 is maintained at DSBs, where is phosphorylated by ATM to block the DNA end processing[9]. On the contrary, in resection-proficient conditions it is evicted from the DNA[9]. Considering our data, we wondered if CRY1 could be regulating CCAR2 retention/eviction from sites of damaged chromatin. Strikingly, phosphorylated CCAR2 recruitment/retention was diminished if CRY1 was downregulated, as observed by the fraction bound to chromatin upon DNA damage induction (Fig. 3J) and retained by immunofluorescence upon pre-extraction (Fig. 3K). Then, we checked CCAR2 eviction from the vicinity of the damaged DNA by laser microirradiation, that can be observed as a negative signal upon known as an anti-stripe[23]. Interestingly, CRY1 depletion increases the formation of CCAR2 anti-stripes 7-fold (Fig. 3L, see white arrow for an example of a laser-induced anti-stripe). Furthermore, not only the number of cells showing antistripes was increased upon CRY1 depletion, but quantification of the remaining signal at the laser-induced antistripes showed a stronger reduction of the CCAR2 signal (Fig. 3M), agreeing with a reduced capacity to retain the protein there. On the contrary, when CtIP was downregulated even those fewer cells that showed antistripes still maintain a significant proportion of the GFP-CCAR2 signal, arguing that there is some CCAR2 eviction, but a considerable fraction of the protein is still retained (Fig. 3M). Then, we tested how CtIP recruitment/retention to laser lines responded to CRY1 levels live. Cells transfected with a control siRNA accumulate quickly, within minutes, GFP-CtIP laser stripes (Fig. 3N, Suppl. movie 3). However, when CRY1 was depleted, we observed that GFP-CtIP accumulation was stronger and at a faster rate (Fig. 3N and Suppl. movie 4). The opposite effect was observed upon CRY1 overexpression using a GFP-tagged version of the protein, that causes a reduction on the recruitment of Cherry-CtIP at laser lines (Fig. 3O; Suppl. movies 5 and 6). Furthermore, an increased retention of CtIP at sites of AsiSI-induced breaks could also be observed upon CRY1 depletion by ChIP-Seq (Fig. 3P). Thus, CRY1 levels control the retention or eviction of CCAR2, a requisite to unleash CtIP and allow DNA end resection.

Resection is modulated by DNA-PK-mediated phosphorylation of CRY1

Our data fit with a model in which CRY1 presence reinforces the antagonistic effect of CCAR2 over CtIP, stabilizing CCAR2 presence on damaged chromatin hence blocking CtIP release and, therefore, resection. However, and differently of the cell cycle regulation of resection, this is not an all or nothing mechanism. So, we hypothesized that even when CRY1 levels are high, at "late afternoon", there must be a mechanism that allows some resection. Thus, we decided to analyse how the DDR finetunes the circadian oscillations of DNA end resection. Previously, CRY1 itself has been shown to be a substrate of the kinase DNA-PK[31], a modification conserved in mammals but no other vertebrates. Interestingly, these phosphorylation sites are absent in CRY2. Considering its anti-resection and pro-NHEJ role, it made sense that such phosphorylation stimulates CRY1-mediated inhibition of CtIP-mediated resection. Thus, we analysed the effect of such modification on the modulation of DNA end resection and recombination. Indeed, ectopic expression of human CRY1 or its stabilization with KL001 hampered DNA end processing only if DNA-PK was active (Fig. 4A-B; Suppl. Fig. 4A-B for representative images). Furthermore, DNA-PK inhibition abolished the resection dampening observed 12h after release from DEX (Fig. 4C). CRY1 has been shown to be phosphorylated at three specific sites[31]. So, to test if this effect was direct, we checked the effect of the overexpression of a CRY1 version that cannot be phosphorylated by DNA-PK (CRY1-3A) or that mimicked constitutive phosphorylation (CRY1-3E) in resection in the presence or absence of DNA-PK inhibitor. As shown in figure 4D, overexpression of wildtype CRY1 hampered resection only when DNA-PK was active, in agreement with a requirement of DNA-PK-mediated phosphorylation for its anti-resection role (see Suppl. Fig. 4C for representative images). In stark contrast, a CRY1 that cannot be modified by DNA-PK did not impair resection even when DNA-PK was functional. On the contrary, a constitutive phosphorylation of CRY1 was enough to hamper resection even in the presence of DNA-PK inhibitor. To determine the molecular role of those phosphorylations, we measured CRY1 accumulation at laser-lines in real time in the different mutant backgrounds (Figure 4E; Suppl. movies 7-9). All three versions of the protein were recruited to DSBs with similar kinetics. As a control, we confirmed that all three constructs are expressed to similar levels and maintain a comparable protein stability to the endogenous CRY1 (Suppl. Fig. 4D).

Then, we tested if retention of CRY1 was affected by DNA-PK-mediated phosphorylation by analysing laser lines with longer time courses (Figure 4F; Suppl. movies 10-12). Strikingly, we observed that albeit all three CRY1 variants were readily

recruited, only when CRY1 was phosphorylated it was retained at sites of damaged

DNA. Thus, these data suggest that upon the appearance of DSBs CRY1 is recruited to

sites of broken chromosomes, but only when it is phosphorylated by DNA-PK it is

retained. We wondered if other PIKKs could affect CRY1 accumulation directly or

indirectly. Indeed, as expected, DNA-PK inhibition affects CRY1 retention similarly to

the CRY1-3A mutant (Fig. 4G). Additionally, the inhibition of ATM, but not ATR, also

diminished CRY1 retention (Fig. 4G). If this reflects the effect of ATM on CCAR2

retention or others, more indirect, effects, remains to be studied. Of note, neither DNA-

PKcs nor ATM nor ATR levels change during a circadian cycle nor in response to

CRY1 depletion (Suppl. Fig. 4E).

Circadian regulation of DSB repair affects cell survival to genotoxic agents

Because of CRY1 regulating DNA end resection and, therefore, HDR, cells

become hyper-resistant to several DNA damaging agents when this factor was depleted

(Fig. 5A-B) and hyper-sensitive when it was stabilized using KL001 (Fig. 5C-D). Albeit

this effect might seem modest, it is of a similar level to that observed upon depletion of

core resection factors such as CtIP[32]. So, we conclude that HDR stimulation when

CRY1 was low facilitated repair, whereas high CRY1 induced a reduction in HDR that

hampered DNA repair. Indeed, CRY1 depletion reduced the number of cells with

micronuclei 24 h after irradiation with 10 Gy (Fig. 5E; see Suppl. Fig. 5A for

representative images) and the number of unrepaired breaks in the next mitosis upon

DSB induction (Fig. 5F; see Suppl. Fig. 5B for representative images). On the contrary,

stabilization of CRY1 increases the number of unrepaired breaks in the following

mitosis after exposure to IR (Fig. 5G; see Suppl. Fig. 5C for representative images).

Moreover, when we measured the disappearance of γH2AX 24h after irradiation as a

proxy of DSB repair completion, we observed that cells depleted of CRY1 were more

efficient than control cells (Fig. 5H; see Suppl. Fig. 5D for representative images). To

confirm that CRY1 acts through CCAR2, we tested survival to IR and Etoposide upon

single and double depletions (Fig. 5I-J). As expected, both single depletions affected

survival similarly, and double depletion showed an epistatic effect. The same genetic

relationship was seen on micronuclei formation (Fig. 5K). Furthermore, we have

directly tested for the persistence of DSBs by COMET assay. Indeed, depletion of either

CRY1 or CCAR2 strongly reduced the tail length, indicating a faster elimination of the

breaks (Fig. 5L). As expected for factors involved in the same pathway, double

depletion of both did not render any further effect. Thus, we conclude that the levels of CRY1 are important to maintain the stability of the genome through its regulation of CCAR2.

CRY1 levels affects tumour biology and prognosis

Circadian proteins are commonly deregulated in tumour samples. Thus, we wondered if such relationship with the repair of induced DSBs and genomic instability was maintained in those pathological setups and, furthermore, if it could be exploited in the clinic. First, we reasoned that if CRY1 levels impact genomic instability it might affect the probability to acquire secondary mutations in cancer samples that lead to a new tumour event. Using Pan-Cancer data from The Cancer Genome Atlas (TCGA) we observed that cells with high levels of CRY1, i.e., with a decreased recombination capacity and worse repair according to our model, showed an acceleration in the appearance of a new tumour event (Fig. 6A). According to our data, higher levels of CRY1 were associated with an increased exposure to the mutational signature associated with defective HDR (SBS3; Fig. 6B), i.e. the one defined by BRCA1 or BRCA2 absence. In fact, tumours with low levels of CRY1 present virtually no exposure to this specific mutational signature. Then, we reasoned that despite this increased mutagenicity of cancer samples with high levels of CRY1, this reduced repair ability can be exploited by using therapeutic agents designed to create DSBs. Indeed, using again TCGA data, we observed that breast cancer patients with tumours expressing higher CRY1 levels were more responsive to radiotherapy, in agreement with the idea that high CRY1 levels render the tumour cells more sensitive to radiation (Fig. 6C). On the contrary, breast tumours with lower CRY1 expression were less efficiently treated with radiotherapy (Fig. 6C), with a decrease in the median survival of 1.5 years (571 days). Strikingly, similar results were observed regarding CCAR2 levels (Fig. 6D), albeit with an even bigger increase in median survival of over 2.5 years (957 days), reinforcing that both CCAR2 and CRY1 work in the same pathway. Furthermore, in heterotopic, subcutaneous, xenografts models using HCT116 cells bearing a constitutive shRNA against CRY1 or a control sequence, we observed a faster growth of tumours lacking CRY1 and a better tolerance to etoposide treatment (Fig. 6E, Suppl. Fig 6). Indeed, in samples expressing CRY1, etoposide treatment caused a 25% reduction in tumour growth rate. In agreement with a mild increase in HDR, and a

reduced cytotoxic effect of etoposide, such decrease in tumour growth rate was slightly

reduced to 18% in samples lacking CRY1.

Then, we reasoned that if CRY1 levels affected the response to genotoxic agents

such as radiotherapy, it is possible that the time of the day when patients are irradiated

might impact the efficiency of this treatment. In a retrospective analysis using patient

data from the Radiotherapy Service from the University Hospital Virgen Macarena

(Seville, Spain) we analysed this response stratifying the patients in two groups: those

irradiated in the morning versus those irradiated in the afternoon/evening. Using data

from all cancer types (pancancer) we observed a significant difference between those

two groups (Fig. 6F). In agreement with our CRY1 levels results, irradiation in the

afternoon-evening, when CRY1 levels build up, rendered tumour samples more

sensitive to radiotherapy and improved patient prognosis. We hypothesize that this

effect might be small because many cancers show deregulation of circadian proteins,

and in this case, patients might not always benefit from this chronotherapy. For example,

directly deregulation of CRY1 has been observed in some tumours such as lung cancer,

whereas in other tumour types like prostate cancer deregulation of circadian rhythms

does not rely in changes in CRY1 levels[33]. Strikingly, careful examination of the data

agreed with this idea and, indeed, whereas prostate cancer patients do benefit of

irradiation at specific times of the day (Fig. 6G), lung cancer patients did not (Fig. 6H).

This differential behaviour can be observed also in other cancer types. For example, and

in agreement with our TCGA analysis for CRY1 levels (Fig. 6B), a benefit in an

afternoon irradiation can be seen in breast cancer patients (Fig. 6I). An even stronger

effect is observed in melanoma, maybe reflecting a response to the direct exposure to

sunlight of the tissue, but the low number of patients with this type of cancer that are

treated with radiotherapy precludes the statistical significance of the differences (Fig.

6J). On the contrary, other tumours types such as head and neck or gliomas do not

benefit from a chronotherapeutic approach (Fig 6K-L).

DISCUSSION

Circadian rhythms and the repair of DNA damage are strongly evolutionary

linked. Indeed, CRY1 belongs to the Cryptochrome family, in which bacterial

photolyases are included[34]. Similarly, Timeless acts as a bona fide circadian accessory

factor in Drosophila, whereas in mammals has shifted its role toward a checkpoint

function[35]. Strikingly, in Drosophila there is a second family member, Timeout, that has

been related to the repair of DSBs[35]. So, it seems that during evolution a bleeding

between circadian-repair proteins has occurred multiple times. Although this simple

crossing between circadian-repair pathways might explain why in mammals CRY1

behaves as both a circadian and a repair factor, a more intriguing hypothesis is that this

dual nature reflects an evolutionary benefit of synchronizing DSB repair with the

night/day cycle. So, a likely explanation is that this regulation prepares cells to face a

load of DSBs at specific times of the day. Considering that UV light is a strong inductor

of DNA damage, including DSBs when not repaired by NER, this connection might not

seem surprising. Indeed, NER has been clearly shown to be regulated by the circadian

clock, attuning this repair pathway to the part of the day in which UV-derived DNA

lesions will appear[36]. So, a possibility is that this connection reflects a need to prepare

cells to face unrepaired UV-lesions that can lead to DSBs that are more likely to appear

during the day. However, we do not favour this idea, mostly by two facts. First, most

mammalian cells do not suffer this kind of damage as they are protected from the

mutagenic effect of UV by the skin. And second and more importantly, we have

observed a conservation of the role of CRY1 in mice that disagrees with this hypothesis.

Relevantly, mice are nocturnal animals, so they have a 12-hour shift in their circadian

clock respect to humans and other diurnal mammals, with CRY1 peaking at "early

morning" and sporting a maximal reduction at "late afternoon"[37]. Thus, in mice cells

recombination peaks at "dusk" and it is reduced at "dawn", opposite to what happens in

human cells. So, for mice to maintain the same CRY1-dependent circadian regulation

points out toward a different source of DNA damage that does not appear during the day

but accumulates at night. An alternative explanation will be replication-born damage. It

has been established that the circadian clock and the cell cycles are in phase, and in

human cells that means that replication takes usually place at specific times of the day[38].

Albeit it is not completely clear how this coordination works, and the time of the day in

which replication takes place varies among different cell types, it has been proposed

that replication only takes place at specific times (gates) of the day in specific circadian

windows[38]. So, it is possible that the regulation of resection by CRY1 levels is a way to

reinforce the cell cycle regulation of homologous recombination, ensuring that resection

and HDR only happen in specific circadian windows too. A better understanding on the

synchronization between replication and the circadian clock is required to test this idea.

However, considering all our data together, we favour a different explanation. Indeed,

we propose that this circadian synchronization might prepare mammalian cells to a

source of damage that is more likely to appear during the day in humans and at night in

mice. So, we hypothesize that this circadian regulation might reflect a need to attune

DSB repair with the appearance of damage caused by the cellular metabolism, another

well-stablished source of DNA damage[39]. Interestingly, CCAR2 has been proposed to

coordinate the cell metabolism with the circadian clock through its interaction with the

Rev-erbα receptor[30], suggesting that it might mediate also further coordination of these

two aspects with DSB repair. In this model, during the more active phase of the cells,

i.e., when the metabolism is more active, higher rates of recombination will be required.

So, in humans, cells will benefit of a boost of recombination at "early morning"

whereas mice will favour such increase at "late afternoon" as a preparation of the

initiation of the active phase of their circadian cycle. Then, during this more active

phase recombination can slowly decline until "dusk" (in human) or "dawn" (mice) in

readiness for a more reduced metabolic status, meaning that the chances of encounter

DSBs that require recombination for repair are reduced. Then, the cells will increase

slowly again their ability to activate resection to face the next day/night cycle.

But how does this circadian regulation impinge on the regulation of DNA end

resection? We favour a model in which is the specific presence of CRY1, and no other

circadian factors, what controls resection through CCAR2 and its inhibition of CtIP.

Strikingly, this cryptochrome paralogue specificity might reflect the lack of

conservation of the C-terminal tail between CRY1 and CRY2, including the absence of

the DNA-PK phosphorylation sites in the later. In any case, we do not discard other

alternative mechanisms in other animals that lack this specific phosphorylation or even

CRY1 itself. In our model (Fig. 6M; this reflects the circadian regulation in humans in

an ideal 12 hours day followed by 12 hours night cycle, but in mice it will be shifted 12

hours), we envision that the levels of CRY1 are critical to retain CCAR2 and restrain

CtIP by a physical interaction between these three proteins. Indeed, we think that at

"early morning" (Figure 6M, top left) CCAR2 and CtIP are recruited together to DSBs

as previously proposed[9]. But in the absence of CRY1, CCAR2 cannot stay on damaged

DNA, so it is readily evicted from chromatin unleashing CtIP and creating a pro-

resection environment. On the contrary, at "late afternoon" (Figure 6M, bottom left),

CRY1 will be constitutively interacting with CCAR2. Therefore, when a DSB appear

all three proteins will be recruited on chromatin in a very rapid response. Then, the

activation of DNA-PK will cause the phosphorylation of CRY1, what will lock it on

DNA, fastening CCAR2 also on the broken chromatin and constraining CtIP activity. This is also reinforced by ATM activity, either by affecting CtIP or CCAR2 or by additional mechanisms. During either the day or the night (Figure 6M, right side), cells will react differently depending on the levels of CRY1 that will be accumulating (during the day) or decreasing (during the night), so the breaks will be more or less prone to be resected depending on the CRY1 levels at that specific time of the day. Interestingly, this first regulation represents only a first layer of the modulation of recombination by the circadian proteins. In the accompanying paper, Legube and colleagues, describe an additional mechanism by which the PER proteins affect homology mediated repair. Interestingly, and as for CRY1, this seems to be a non-canonical function of PER proteins independent of others circadian core clock components. Specifically, they see that PER2 reduces the clustering of DNA double strand break by helping their localization at the nuclear envelope, therefore fostering the strand invasion step of recombination and minimizing genetic translocations. Due to an uncoupling on the oscillation of CRY1 and PER proteins, this mechanism reinforces a reduction of recombination during the night. Thus, we collectively support a two-tier model in which resection is dampened at "late afternoon" following CRY1 peak, but even those breaks that have been partially resected will later fail to engage a homologue sequence due to reduce PER levels at night. Additionally, it has been recently proposed that BMAL1-CLOCK stimulate resection and recombination through the acetylation of histone H4[40], hence suggesting a further stimulation of recombination during the day. Importantly, this regulatory mechanism will impact in the quality of repair, with a strong effect in the appearance of genomic translocation when defective. Of note, a previous report discovered that in the context of prostate cancer, the androgen receptor stimulates CRY1, that then affects DNA repair by regulating the expression of DDR[25]. However, they find the opposite role of CRY1 and propose that stimulates recombination. If this is due to differences in cell types or a consequence of hormone induction remains to be resolved.

This built-in mechanism(s) will act at the cellular level. Additionally, there is a secondary control of the circadian rhythms at the level of light sensing that can reset the cellular clock, that in the case of the mammals require the detection of light by the eyes, the processing of this signal at the suprachiasmatic nucleus (SCN), and then the transmission of this information to all the cells in the organism[11]. How this supracellular regulation impinges in this regulatory network we have uncovered should be explored

in the future. Indeed, our data suggest that perturbations of the circadian rhythms might have consequences in the ability of the cells to repair broken chromosomes. Furthermore, it will affect not only their ability to repair DSBs, but also the balance between the more error-free HDR and the error-prone NHEJ, hence putatively impacting both the quantity and quality of the repair. This might, in turn, affect the stability of the genome, altering the mutational load. Indeed, we see an increased exposure to the mutational signature associated with defective HDR in cells with high levels of CRY1. Strikingly, it has been known for decades that people that alternate between different day-night shifts at work or suffering of chronic jetlag are more prone to develop cancer and they generally have a poorer prognosis[41,42], albeit the reasons are far from clear. One tantalizing idea is that these workers have a deregulation between their DSBs repair pathways due to the circadian perturbations, so they have a higher chance toward a mutagenic repair that might contribute, among other factors, to tumour development. Along the same lines, the same explanation can be invoked to understand why circadian regulators are often altered in many cancer cells[43,44]. So, tampering with the built-in circadian clock, either by external effects, such as shift changes, or internal causes, such as mutation of circadian genes, will contribute to an increase in the mutagenic burden that can predispose toward the appearance of cancer. Interestingly, tissue, age and even gender differences in the robustness of the circadian cycle have been documented[15], something that might be of importance in cancer evolution. Our analysis using TCGA data and mouse xenografts support this idea of higher chances of tumour development and a faster evolution when CRY1 protein levels are reduced. Furthermore, this should also be considered for cancer treatment, as cells depleted for CRY1 are more resistant to DSB-inducing agents, even in the xenografts models. The exploitation of the connection between circadian rhythms and cancer treatment efficiency is known as chronotherapy[45], and there is discussion if this might be applied to radiotherapy[46]. We have even observed this effect in a real-life clinical setup, albeit with a very modest impact and, interestingly, only on specific tumour samples. The reasons behind those differences and, specifically, if this reflects the loss of CRY1 regulation in specific tumours must be explored in the future to understand the therapeutic potential of chronoradiotherapy.

REFERENCES

1. Jackson, S. P. & Bartek, J. The DNA-damage response in human biology and

disease. *Nature* **461**, 1071–1078 (2009).

2. Chang, H. H. Y., Pannunzio, N. R., Adachi, N. & Lieber, M. R. Non-homologous DNA end joining and alternative pathways to double-strand break repair. *Nat. Rev. Mol. Cell Biol.* **18**, 495–506 (2017).

3. Ranjha, L., Howard, S. M. & Cejka, P. Main steps in DNA double-strand break repair: an introduction to homologous recombination and related processes. *Chromosoma* **127**, 187–214 (2018).

4. Jasin, M. & Rothstein, R. Repair of strand breaks by homologous recombination. *Cold Spring Harb. Perspect. Biol.* **5**, (2013).

5. Cejka, P. DNA end resection: Nucleases team up with the right partners to initiate homologous recombination. *J. Biol. Chem.* **290**, 22931–22938 (2015).

6. Symington, L. S. Mechanism and regulation of DNA end resection in eukaryotes. *Crit. Rev. Biochem. Mol. Biol.* **51**, 195–212 (2016).

7. Mozaffari, N. L., Pagliarulo, F. & Sartori, A. A. Human CtIP: A 'double agent' in DNA repair and tumorigenesis. *Semin. Cell Dev. Biol.* **113**, 47–56 (2021).

8. Makharashvili, N. & Paull, T. T. CtIP: A DNA damage response protein at the intersection of DNA metabolism. *DNA Repair (Amst).* **32**, 75–81 (2015).

9. López-Saavedra, A. *et al.* A genome-wide screening uncovers the role of CCAR2 as an antagonist of DNA end resection. *Nat. Commun.* **7**, (2016).

10. Iyera, D. R. *et al.* CCAR2 functions downstream of the Shieldin complex to promote double-strand break end-joining. *Proc. Natl. Acad. Sci.* **119**, e2214935119 (2022).

11. Stratmann, M. & Schibler, U. Properties, entrainment, and physiological functions of mammalian peripheral oscillators. *J. Biol. Rhythms* **21**, 494–506 (2006).

12. Bhadra, U., Thakkar, N., Das, P. & Pal Bhadra, M. Evolution of circadian rhythms: from bacteria to human. *Sleep Med.* **35**, 49–61 (2017).

13. Takahashi, J. S. Transcriptional architecture of the mammalian circadian clock. *Nat. Rev. Genet.* **18**, 164–179 (2017).

14. Brown, S. A., Kowalska, E. & Dallmann, R. (Re)inventing the Circadian Feedback Loop. *Dev. Cell* **22**, 477–487 (2012).

15. Talamanca, L., Gobet, C. & Naef, F. Sex-dimorphic and age-dependent organization of 24-hour gene expression rhythms in humans. *Science (80-.).* **379**, 478–483 (2023).

16. Balsalobre, A. *et al.* Resetting of circadian time in peripheral tissues by glucocorticoid signaling. *Science (80-.).* **289**, 2344–2347 (2000).

17. Izumo, M., Sato, T. R., Straume, M. & Johnson, C. H. Quantitative analyses of circadian gene expression in mammalian cell cultures. *PLoS Comput. Biol.* **2**, 1248–1261 (2006).

18. Ferretti, L. P., Lafranchi, L. & Sartori, A. A. Controlling DNA-end resection: A new task for CDKs. *Front. Genet.* **4**, 1–7 (2013).

19. Gaucher, J., Montellier, E. & Sassone-Corsi, P. Molecular Cogs: Interplay between Circadian Clock and Cell Cycle. *Trends Cell Biol.* **28**, 368–379 (2018).

20. Escribano-Díaz, C. *et al.* A Cell Cycle-Dependent Regulatory Circuit Composed of 53BP1-RIF1 and BRCA1-CtIP Controls DNA Repair Pathway Choice. *Mol. Cell* **49**, 872–883 (2013).

21. Chapman, J. R. *et al.* RIF1 Is Essential for 53BP1-Dependent Nonhomologous End Joining and Suppression of DNA Double-Strand Break Resection. *Mol. Cell* **49**, 858–871 (2013).

22. Di Virgilio, M. *et al.* Rif1 prevents resection of DNA breaks and promotes immunoglobulin class switching. *Science (80-.).* **339**, 711–715 (2013).

23. Lopez-Saavedra, A. *et al.* A genome-wide screening uncovers the role of CCAR2 as an antagonist of DNA end resection. *Nat. Commun.* **7**, 12364 (2016).

24. Baggs, J. E. *et al.* Network features of the mammalian circadian clock. *PLoS Biol.* **7**, 0563–0575 (2009).

25. Shafi, A. A. *et al.* The circadian cryptochrome, CRY1, is a pro-tumorigenic factor that rhythmically modulates DNA repair. *Nat. Commun.* **12**, (2021).

26. Aymard, F. *et al.* Transcriptionally active chromatin recruits homologous recombination at DNA double-strand breaks. *Nat. Struct. Mol. Biol.* **21**, 366–374 (2014).

27. Iannelli, F. *et al.* A damaged genome's transcriptional landscape through multilayered expression profiling around in situ-mapped DNA double-strand breaks. *Nat. Commun.* **8**, 15656 (2017).

28. Hirota, T. *et al.* Identification of small molecule activators of cryptochrome. *Science (80-.).* **337**, 1094–1097 (2012).

29. Magni, M., Buscemi, G. & Zannini, L. Cell cycle and apoptosis regulator 2 at the interface between DNA damage response and cell physiology. *Mutat. Res. - Rev. Mutat. Res.* **776**, 1–9 (2018).

30. Chini, E. N., Chini, C. C. S., Nin, V. & Escande, C. Deleted in breast cancer-1 (DBC-1) in the interface between metabolism, aging and cancer. *Biosci. Rep.* **33**, 637–643 (2013).

31. Gao, P. *et al.* Phosphorylation of the cryptochrome 1 C-terminal tail regulates circadian period length. *J. Biol. Chem.* **288**, 35277–35286 (2013).

32. Huertas, P. & Jackson, S. P. Human CtIP Mediates Cell Cycle Control of DNA End Resection and Double Strand Break Repair. *J. Biol. Chem.* **284**, 9558–9565 (2009).

33. Mocellin, S., Tropea, S., Benna, C. & Rossi, C. R. Circadian pathway genetic variation and cancer risk: Evidence from genome-wide association studies. *BMC Med.* **16**, 1–8 (2018).

34. Öztürk, N. *et al.* Structure and function of animal cryptochromes. *Cold Spring Harb. Symp. Quant. Biol.* **72**, 119–131 (2007).

35. Mazzoccoli, G., Laukkanen, M. O., Vinciguerra, M., Colangelo, T. & Colantuoni, V. A Timeless Link Between Circadian Patterns and Disease. *Trends Mol. Med.* **22**, 68–81 (2016).

36. Bee, L. *et al.* Nucleotide excision repair efficiency in quiescent human fibroblasts is modulated by circadian clock. *Nucleic Acids Res.* **43**, 2126–2137 (2015).

37. Wang, J. *et al.* Nuclear Proteomics Uncovers Diurnal Regulatory Landscapes in Mouse Liver. *Cell Metab.* **25**, 102–117 (2017).

38. Masri, S., Cervantes, M. & Sassone-Corsi, P. The circadian clock and cell cycle: Interconnected biological circuits. *Curr. Opin. Cell Biol.* **25**, 730–734 (2013).

39. Moretton, A. & Loizou, J. I. Interplay between cellular metabolism and the dna damage response in cancer. *Cancers (Basel).* **12**, 1–29 (2020).

40. Zhang, C. *et al.* BMAL1 collaborates with CLOCK to directly promote DNA double-strand break repair and tumor chemoresistance. (2023). doi:10.1038/s41388-023-02603-y

41. Shostak, A. Circadian clock, cell division, and cancer: From molecules to organism. *Int. J. Mol. Sci.* **18**, (2017).

42. Lamia, K. A. Ticking time bombs: connections between circadian clocks and cancer. *F1000Research* **6**, 1910 (2017).

43. Yu, H. *et al.* Cryptochrome 1 Overexpression Correlates with Tumor Progression and Poor Prognosis in Patients with Colorectal Cancer. *PLoS One* **8**, (2013).

44. Xiang, R. *et al.* Circadian clock gene Per2 downregulation in non-small cell lung

cancer is associated with tumour progression and metastasis. *Oncol. Rep.* **40**, 3040–3048 (2018).

45. Lévi, F. Circadian chronotherapy for human cancers. *Lancet Oncol.* **2**, 307–315 (2001).

46. Chan, S. *et al.* Does the Time of Radiotherapy Affect Treatment Outcomes? A Review of the Literature. *Clin. Oncol.* **29**, 231–238 (2017).

47. Jimeno-González, S. *et al.* Defective histone supply causes changes in RNA polymerase II elongation rate and cotranscriptional pre-mRNA splicing. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 14840–14845 (2015).

48. Schmidl, C., Rendeiro, A. F., Sheffield, N. C. & Bock, C. ChIPmentation: fast, robust, low-input ChIP-seq for histones and transcription factors. *Nat. Methods* **12**, 963–965 (2015).

ACKNOWLEDGEMENTS.

This work was funded by the R+D+I grants PID2019-104195G and PID2022-136791NB-I00 funded by MICIU/AEI/10.13039/501100011033/ FEDER/UE and the grant P18-RT-1204 from the Consejería de Transformación Económica, Industria, Conocimiento y Universidades, Junta de Andalucía. The mouse work in ALC laboratory was funded by the R+D+I grant PID2020-119329RB-I00 and SJG lab is funded by the grant PID2022-139253N, both funded by MICIU/AEI/10.13039/501100011033/ FEDER/UE. AR-F is funded with a FPU fellowship from the Spanish Ministry of Education and MC-P is the recipient of a MSCA-IF-2020 (coDNAres). PA was supported by a postdoctoral fellowship from the Junta de Andalucía and the European Social Fund (202099908313486), and by a postdoctoral grant from the Scientific Foundation of the Spanish Association Against Cancer (POSTD234422AGUI). CABIMER is supported by the regional government of Andalucía (Junta de Andalucía). We thank José Carlos Reyes-Rosa for critical reading of the manuscript.

DATA AVAILABILITY.

All relevant data are included in the manuscript. Raw data will be provided upon request. ChIP-seq data are deposited at GEO under the accession number GSE261427.

ETHICS&INCLUSION.

All authors are based on Seville (Spain) as a collaboration of the Research Centre CABIMER and the local University Hospital Virgen Macarena. Research using patient clinical data was approved by the Ethical Board of the Hospital. The animal experimentation protocol was approved by the CABIMER Ethics Committee for Animal Experimentation (CEEA) and the Junta de Andalucía (project 26/02/2021/025) and the experiments were performed in compliance with the Spanish and European regulations.

FIGURE LEGENDS

Figure 1. Resection oscillates following a circadian pattern. A, Schematic representation of the experimental setup for circadian synchronization. Cells were exposed to dexamethasone or forskolin for two hours. Then, cells were released by changing the medium, mimicking "sunrise" conditions. Cells were then irradiated at the indicated timepoints that correspond to the depicted times of the day/night cycle. **B**, U2OS were synchronized as mentioned in A with either Dexamethasone (blue line) or EtOH as a control (red line) and irradiated with 10Gy at the indicated time points. 1 h later cells were immunostained for RPA. Then, the percentage of cells with more than 10 RPA foci was scored. At least 250 cells were analyzed for each sample in each replica. The average and standard deviation of four independent experiments is shown. Statistically significance was calculated using a two-way ANOVA. Only statistical different comparisons are depicted. **C**, Same as B, but using Forskolin (blue line) or DMSO as a control (red line). The average and standard deviation of four independent experiments is shown. **D**, Same as B, but cells were followed for 48 h after release from dexamethasone. **E**, Same as in D, but measuring CRY1 levels by immunofluorescence. **F**, Same as B but in RPE1 cells. Dashed line represents the value of an asynchronous control. The average and standard deviation of four independent experiments is shown. **G**, Same as F but in MEFs. **H**, The length of resected DNA was calculated as described in the methods section in cells synchronized and released from dexamethasone at the indicated times or in an asynchronous control. Each dot represents the length of an individual resected DNA fiber, and the median is shown in red. Statistical significance was calculated using a Mann–Whitney test. The experiment was repeated three times with similar results. **I**, Same as F but immunostaining for BRCA1. **H**, Same as I but immunostaining for RIF1. **K**, Same as I but immunostaining for RAD51 3 hours after DSB induction with IR in U2OS cells. Unless stated otherwise, for all main and

supplementary figures, the average and standard deviation of three independent experiments are shown and p values are represented with one ($p < 0.05$), two ($p < 0.01$) or three ($p < 0.001$) asterisks.

Figure 2. Resection oscillates following a circadian pattern. A, CRY1 (siCRY1) or control (siNT) depleted U2OS cells were transfected with the human version of CRY1 (black bars) or an empty vector (white). 1 h after irradiation with 10 Gy cells were immunostained for RPA. Then, the percentage of cells with more than 10 RPA foci was scored (left). Statistically significance was calculated using a two-way ANOVA. Only statistical different comparisons are depicted. Representative images are shown on the right side. **B**, Same as figure 1B, but in cells transfected with an siRNA against CRY1 or a control sequence, as indicated. **C**, Average recruitment of RPA at the 214 best AsiSI cutting sites measured by ChIP-seq in samples transfected with an siRNA against CRY1 or a control sequence, as indicated. One representative experiment, out of 2 with similar results, is shown. **D**, QIBC representation of RPA foci formation. Cells treated as in A were immunostained for RPA. The intensity of RPA or DAPI were plotted in the Y and X axis, respectively. The vertical dashed line represents the threshold of DAPI intensity corresponding to S/G2 cells, whereas the horizontal line represents the threshold for RPA positive cells. The percentage of S/G2 cells showing RPA foci signal is indicated. A representative experiment out of 2 two independent replicates is shown. **E**, Same as A but in cells overexpressing or not CRY1 without siRNA transfection. **F**, Same as A but in cells treated with the CRY1 stabilizer KL001. **G**, Gene conversion was assessed in U2OS cells bearing the DR-GFP reporter in cells transfected with an siRNA against CRY1 (siCRY1) or a control sequence (siNT), as indicated. The amount of GFP positive cells, i.e., cells that have undergone repair by gene conversion, after induction with the nuclease I-SceI, were normalized to the siNT control taken as 1. The statistical significance was calculated using a Student´s t-test. The average and standard deviation of four independent experiments is shown. **H**, Same as G, but using the SSA reporter SA-GFP. **I**, Same as G, but using the NHEJ reporter EJ5-GFP. **J**, Same as G, but in cells bearing an empty vector or a vector harboring an ectopic version of CRY1. The average and standard deviation of four independent experiments is shown. **K**, Same as J, but using the SSA reporter SA-GFP. **L**, Same as J but using the NHEJ reporter EJ5-GFP. **M**, RAD51 foci 3 h after exposure to irradiation were scored by immunofluorescence in cells depleted for CRY1 (siCRY1) or control cells (siNT). The

number of RAD51 foci per cell was counted and represented as a single dot. The median number of Rad51 foci is shown as a red line. Statistical significance was calculated using a Mann–Whitney test. A representative experiment out of three with similar results is shown. **N**, Same as M, but in cells transfected with a plasmid bearing an ectopic copy of CRY1 or an empty vector. **O**, Same as M, but immunostaining for RIF1. **P**, Same as N, but immunostaining for RIF1. **Q**, U2OS cells transfected with an siRNA against CRY1 (siCRY1), CCAR2 (siCCAR2), both (siCRY1siCCAR2) or a control sequence (siNT), were irradiated with 10Gy. Then, the percentage of cells with more than 10 RIF1 foci was scored at the indicated time points after irradiation. **R**, Same as M, but immunostaining for 53BP1. **S**, Protein samples, depleted or not for CRY1 as indicated, were taken 1 h after irradiation. Upon cellular fractionation, both the whole cell extract (WCE) and the chromatin-bound fraction (Chromatin), were resolved in SDS-PAGE and blotted with antibodies against the indicated proteins.

Figure 3. CRY1 levels control DNA end resection through CCAR2 retention at DSBs. A, U2OS bearing shRNA against CCAR2 (shCCAR2), CtIP (shCtIP) or a control sequence (shNT), were synchronized and released as indicated in Figure 1A with dexamethasone. Cells were irradiated at the indicated time points and RPA foci were scored by immunostaining 1 h after irradiation. Statistically significance was determined using a two-way ANOVA. **B**, RPA foci formation in U2OS cells transfected with the indicated siRNAs. A minus sign means transfection with a control sequence. **C**, Same as B, but in cells transfected with a siRNA against CCAR2 or not and bearing a vector for CRY1 overexpression (hCRY1). Minus signs mean transfection with a siRNA control or the empty vector. The average and standard deviation of four independent experiments is shown. **D**, Same as B but immunostaining for BRCA1. **E**, U2OS cells were irradiated with 10G (+) or mock treated (-). Protein samples were isolated in native condition and CRY1 was immunoprecipitated using a specific antibody against CRY1. IgG was used as immunoprecipitation control. Then, proteins were resolved in SDS PAGE and blotted with the indicated antibodies. **F**, Same as E but using an antibody against CCAR2 for the IP. **G**, Same as F but using an antibody against CtIP for the IP. **H**, Same as E but in cells depleted for CtIP using an siRNA. **I**, U2OS cells bearing Cherry-MDC1 and GFP-CRY1 constructs were laser microirradiated and imaged at the indicated times. Representative images of selected timepoints are shown on the right, and quantification of the signal, relative to the peak

intensity taken as 100%, of 6 cells is represented on the right. **J**, U2OS cells transfected with an siRNA against CRY1 or a control sequence were irradiated with 10 Gy. Protein samples were taken at the indicated timepoints, and the chromatin fraction was purified. After resolving the sample in SDS PAGE, the amount of phosphorylated CCAR2 was determined using a specific antibody. Lamin A was used as a control. A representative western blot, out of three with similar results, is shown on top and its quantification of this specific western blot on bottom. **K**, U2OS cells treated as in J were immunostained using an antibody against phosphorylated CCAR2. Representative images are shown on the left side, and the quantification of the phospo-CCAR2 intensity per cell is plotted on the right side in cells irradiates (IR +) or not (IR -). **L**, U2OS cells bearing a constitutively expressed CCAR2-GFP fusion were laser microirradiated. The percentage of cells showing a negative GFP staining 1 h after laser-microirradiated was quantified and plotted (right side) in cells depleted or not for CRY1. Representative image of a cell showing an anti-stripe (white arrow) before and 1 h after laser-microirradiation are also shown (left). **M**, The intensity of CCAR2-GFP at the anti-stripe of cells showing them was calculated, setting the pan-nuclear intensity as 0, in U2OS cells laser-microirradiated upon depletion of CRY1 or CtIP. The average intensity in each case is shown as a red line. At least 60 cells with anti-stripes were analyzed in each sample. **N**, Recruitment of GFP-CtIP was measure upon laser microirradiation in U2OS cells transfected with an siRNA against CRY1 or a control sequence. Representative images are shown on the left side. Images were taken at the indicated times, and the intensity of GFP-CtIP at the laser-induced stripe was quantified and plotted. **O**, Same as N, but in cells stably expressing a Cherry-CtIP and transfected with a GFP-CRY1 plasmid or an empty vector. **P**, Same as figure 2C, but using and antibody against CtIP for ChIP.

Figure 4. DNA-PK-dependent phosphorylation of CRY1 modulates resection. A, U2OS cells were exposed to the CRY1 stabilizer KL001 and/or the DNA-PK inhibitor NU7441 as indicated with the plus and minus signs, and then irradiated and immunostained for RPA. **B**, Same as A but in cells transfected with an ectopic version of CRY1 or the empty vector instead of exposed to KL001. **C**, U2OS cells treated with an inhibitor of DNA-PK or DMSO as a control were treated as in Figure 1B. Other details as Figure 1B. **D**, Cells stable transfected with GFP-fusions of wildtype CRY1 (GFPCRY1-wt), a non-phosphorytable version of CRY1 (GFPCRY1-3A), a phospho-mimicking version (GFPCRY1-3E) or the empty vector (GFP) were treated with

1 NU7441 or not, as indicated, irradiated and immunostained for RPA. **E**, U2OS stably
2 expressing GFP-CRY1 versions were laser microirradiated and imaged at the indicated
3 times. For each timepoint, the average intensity of at least five cell images is plotted. **F**,
4 Same as E, but for longer time points. **G**, Same as in F, but with cells treated with
5 inhibitors for DNA-PK, ATM, ATR or DMSO as a control, as indicated.
6
Figure 5. CRY1 levels modulate the response to DNA damaging agents *in vitro*. A,
U2OS transfected with an siRNA against CRY1 or a control sequence were seeded at
low density and irradiated at the indicated dose. Cells were left to grow for 10 days and
the survival fraction, compared with an unirradiated sample taken at 100%, is plotted. **B**,
Same as A but in cells treated with the indicated doses of etoposide (VP16) for 1 hour.
C, Same as A but in cells pre-exposed to the CRY1 stabilizer KL001 or DMSO. **D**,
Same as B but in cells pre-exposed to the CRY1 stabilizer KL001 or DMSO. **E**, U2OS
cells depleted or not of CRY1, as indicated, were irradiated with 10 Gy. 24 h after
irradiation cells were stained with DAPI and the percentage of cells with micronuclei
was calculated. **F**, Percentage of chromosome breaks present in mitotic cells following
an irradiation with 2 Gy in cells transfected with the indicated siRNAs. **G**, Same as F,
but in cells exposed to KL001 or DMSO. **H**, Number of γH2AX foci in U2OS cells
unchallenged (-) or 24 h after irradiation (+; 10 Gy), in cells transfected with an siRNA
against CRY1 (+) or a control sequence (-), as indicated. **I**, Same as A but in cells
depleted for CRY1, CCAR2 or both proteins simultaneously. **J**, Same as B but in cells
depleted for CRY1, CCAR2 or both proteins simultaneously. **K**, Same as E but in cells
depleted for CRY1, CCAR2 or both proteins simultaneously. **L**, Cells treated as in
panel K were assayed for DNA break presence using neutral comet assay. Left,
quantification of one out of three similar experiment. Each dot represents one nucleus.
Right, a representative image of a nucleus for each condition is shown.
27
28
29
30 **Figure 6. CRY1 levels modulate the behaviour of cancer cells and their response to**
31 **treatment. A**, Elapsed days for a second tumour event in cells stratified for its CRY1
32 levels in two categories: High (red line) or low (blue line). Data from TCGA. The
33 number of tumour samples is indicated. **B**, Pan-cancer data from the TCGA were
34 separated according the levels of CRY1 and the appearance of mutations associated

with the mutational signature 3, caused by defective HDR, was scored and represented. **C**, Survival of breast cancer patients treated with radiotherapy according the CRY1 levels of the tumour cells. Data from TCGA. Other details as in A. **D**, Same as C but regarding CCAR2 levels. **E**, Nude mice were grafted with HCT116 cells depleted for CRY1 on the right flank and control cells on the left flank. Mice were treated with etoposide as indicated in the method section. The average and standard deviation of tumour volume of 6 mice for each condition and timepoint are plotted. Statistical significance of the changes is shown. **F**, Overall survival of cancer patients treated primarily with radiotherapy at the Radiotherapy Service of the University Hospital Virgen Macarena considering if they were irradiated in the morning (blue line) or the afternoon (red line). **G**, Same as F but considering only prostate cancer patients. **H**, Same as F but considering only lung cancer patients. **I**, Same as F but considering only breast cancer patients. **J**, Same as F but considering only melanoma patients. **K**, Same as F but considering only glioma patients. **L**, Same as F but considering only head and neck cancer patients. **M**, Schematic representation of CRY1-mediated regulation of DNA end resection regarding the time of the day in human samples. At "dawn" (top-left), when CRY1 is absent, CtIP and CCAR2 are recruited to DSBs. However, CCAR2 cannot be maintained into chromatin, releasing CtIP and promoting maximum resection. On the contrary, at "dusk" (bottom left), CRY1 interacts with CCAR2. Upon DSB formation, all three proteins are recruited to damaged chromatin, where CRY1 is phosphorylated by DNA-PK. This locks CRY1 and, consequently, CCAR2 on chromatin, hampering CtIP activity, During the day (top right) or the night (bottom night), when CRY1 levels fluctuate, an intermediate situation ensues. I. e. some breaks will be promptly resected and some will not, depending on the presence of CRY1.

Material and Methods

Cell lines and growth conditions

U2OS cells and MEFs were grown in high glucose DMEM (Sigma-Aldrich)

supplemented with 10% foetal bovine serum (Sigma-Aldrich), 2 mM L-glutamine

(Sigma-Aldrich), 100 units/ml penicillin, and 100 μg/ml streptomycin (Sigma-Aldrich).

U2OS cells stably expressing GFP-CtIP, GFP-CCAR2, Cherry-CtIP and Cherry-MDC1

were grown in a standard medium supplemented with 0.5 mg/mL G418 (Gibco,

15140122). U2OS cells harbouring a single copy of the reporter constructs DR-GFP,

SA-GFP and EJ5-GFP were grown in a standard medium supplemented with 1 μg/mL

puromycin (Sigma, P8833).

RPE1 cells were grown in high glucose DMEM/F-12 medium (Sigma-Aldrich)

supplemented with 10% foetal bovine serum (Sigma-Aldrich), 2 mM L-glutamine

(Sigma-Aldrich), 100 units/ml penicillin, and 100 μg/ml streptomycin (Sigma-Aldrich).

RPE1 cells stably expressing shRNAs against CtIP, CCAR2 and a control sequence

(shScr) were grown in this medium supplemented with 5μg/mL puromycin (Sigma,

P8833).

To inhibit DNA-PK we used NU7441(SelleckChem) for 1 HDR prior irradiation.

To stabilize CRY1, cells were pretreated with KL001 (Sigma) for 24 hrs prior the

relevant experiment.

siRNAs, plasmids, and transfections

siRNA duplexes were obtained from Sigma-Aldrich or Dharmacon (Suppl.

Table 2) and were reversely transfected using RNAiMax Lipofectamine Reagent Mix

(Life Technologies), according to the manufacturer's instructions. Plasmid transfection

was carried out using FuGENE 6 Transfection Reagent (Promega) according to the

manufacturer's protocol.

Circadian synchronization experiments

Circadian synchronization was performed by treating cells with either 1 μM of

Dexamethasone (Sigma), or 10 μM of Forskolin (Sigma) or mock treated with EtOH or

DMSO as a control, respectively, for 2 hours. Release from synchronization was

performed by exchanging the medium with fresh one.

Immunoprecipitation

U2OS cells were harvested in lysis buffer (50 mM Tris–HCl, pH 7.5, 50mM

NaCl, 1 mM EDTA, 0.2 % Triton X-100, 1X protease inhibitors (Roche), 1X

phosphatase inhibitor cocktail 1 (Sigma)) and incubated for 30 min on ice with

Benzonase (100 U/mL). Protein extracts (500 μg-1 mg) were then subjected to

preclearing with 20 μL of either magnetic protein G (for Rabbit antibodies) or Protein A

(for Mouse antibodies) Dynabeads (Invitrogen) for 1 hour at 4ºC. Then, precleared

extracts were incubated overnight with 0.5 μg of CRY1 antibody (Rabbit, A302-614A),

1 μg of CCAR2 antibody (Rabbit, A300-433A-1), 1.5 μL of CCAR2 antibody (Mouse,

5857) or the equivalent amount of IgG (Mouse or Rabbit) as the negative control.

Subsequently, extracts were incubated with the corresponding magnetic protein beads

and after that they were washed three times with lysis buffer with 0.02% of Triton X-

100. The precipitate was eluted in 30 μl of Laemmli buffer 2X. The antibodies used for

immunoprecipitation can be found on Suppl. Table 3.

Immunofluorescence and microscopy

For foci visualization, cells were seeded on 12 mm coverslips, and after the

different treatments, cell were washed once with PBS before following the

immunofluorescence procedure.

For RPA and BRCA1 foci analysis, cells were pre-extracted with Pre-extraction

Buffer (25 mM Tris–HCl, pH 7.5, 50 mM NaCl, 1 mM EDTA, 3 mM $MgCl_2$, 300 mM

sucrose, and 0.2% Triton X-100) for 5 min on ice. Then, cells were fixed on ice with

4% paraformaldehyde (w/v) in PBS for 20 min. For RIF1, 53BP1 and γH2AX foci

analysis, cells were treated for 20 min on ice with 4% paraformaldehyde (w/v) in PBS,

and after two washes with PBS 1X, they were permeabilized with 0.25% Triton X-100

in PBS. For RAD51 foci analysis, cells were pre-extracted with 0.1% Triton X-100 in

PBS for 5 min on ice. Then, cells were fixed on ice with 4% paraformaldehyde (w/v) in

PBS for 20 min on ice.

In all cases, following two washes with PBS, cells were blocked for 1 h with 1%

BSA in PBS, co-stained with the appropriate primary antibodies (Suppl. Table 3) in

blocking solution overnight at 4 °C, washed again with PBS and then co-

immunostained with the appropriate secondary antibodies (Suppl. Table 4) in blocking

buffer. After washing with PBS and dehydration with 100% ethanol, coverslips were

mounted into glass slides using Vectashield mounting medium with DAPI (Vector

Laboratories). In the case of QIBC experiments, cells were stained with 0.5 μg/mL

DAPI in PBS for 5 min prior to drying with ethanol and were mounted using Vectashield mounting medium without DAPI (Vector Laboratories). Images were acquired and analysed using a Leica fluorescence microscope. Foci analysis formation was performed using ImageJ.

Single-molecule analysis of resection tracks

To measure the length of the resected DNA fragments SMART (single-molecule analysis of resection tracks) assay was performed or directly lysing and stretching the DNA on coverslip (SMART-tilt) (Altieri et al., 2020). U2OS cells were grown in the presence of 32 µM CldU (Sigma, C6891) for 24 h. They were then irradiated (10 Gy) for 1 h at 37°C and harvested using Trypsin. Cells were centrifuged at 500 g for 3 min at 4 °C, resuspended in PBS and mixed in a 1:8 proportion with unlabelled cells. For DNA stretching, 2.5 µl of this mix were lysed using a spreading buffer (50 mM EDTA, 0.5% SDS, 200 mM Tris-HCl pH 7.4) to later stretch the nucleic acid fibers on silanized slides (Sigma, S4651), tilting the slides to a 15°. After air-drying the DNA fibers for 10 min, they were fixed in 3:1 methanol/acetic acid at -20 °C for 15 min. Slides were then washed twice in PBS, incubated in 70% ethanol overnight at 4°C and washed twice again in PBS. For immunodetection, samples were blocked with 5% BSA (Sigma, A4503) in PBS for 30 min at room temperature (RT) and incubated with anti-BrdU/CldU (Abcam, ab6326) diluted in blocking buffer for 1 h at 37°C. Slides were washed with PBS twice and incubated with the fluorescent secondary antibody (Alexa Fluor 488 anti-rat, Molecular Probes A-11006) diluted in blocking buffer for 1 h at RT. Finally, samples were washed twice in PBS, dried and mounted with ProLong Gold Antifade Reagent (Molecular Probes, P36930). DNA fibers images were acquired with a Leica Thunder Microscope with automatized stage and a 63X oil immersion objective. For each experiment, at least 100 DNA fibers were measured using the Fiji (ImageJ) image software analysis.

SDS-PAGE and western blot analysis

Protein extracts were prepared in 2X Laemmli buffer (4% SDS, 20% glycerol, 125 mM Tris–HCl, pH 6.8) and boiled at 100 °C for 5 min. Then, samples were resolved by SDS–PAGE and transferred to Amersham™ Protran™ Premium NC nitrocellulose membranes (Cytiva Life Sciences). Membranes were blocked with 5% Skim Milk (Difco) in 0.1% Tween-20, 0.01% Sodium Azide supplemented 1X TBS.

After that, membranes were blotted with the appropriate primary antibody (Suppl. Table 3) and infrared dyed secondary antibodies (Suppl. Table 4) diluted in Odyssey Blocking Buffer (LI-COR) supplemented with 0.1% Tween-20 and 0.01% Sodium Azide. Membranes were air-dried in the dark and scanned in an Odyssey Infrared Imaging System (LI-COR), and images were analysed with Image Studio software (LI-COR).

Cell cycle analysis

Cells were fixed with cold 70% ethanol for at least 2 hours, incubated with 250 µg/ml RNase A (Sigma Aldrich) and 10 µg/ml propidium iodide (Sigma Aldrich) at 37 °C for 30 min and analysed with a LSRFortessa™ Cell Analyzer (BD) Flow Cytometer. Cell cycle distribution data were further analysed using ModFit LT 5.0 software (Verity Software House Inc).

HDR and NHEJ analysis

U2OS cells bearing a single copy integration of the reporters DR-GFP (Gene conversion), SA-GFP (SSA) or EJ5-GFP (NHEJ) were used to analyse the different DSB repair pathways. In all cases, 60.000 cells were plated in 6-well plates in duplicate. One day after seeding, cells were transfected with the indicated siRNA or plasmids. The next day, each duplicate culture was infected with lentiviral particles containing I-SceI–BFP expression construct at MOI 10 using 8 µg/ml polybrene in 2 ml of DMEM. Then, cells were left to grow for an additional 24 h before changing the medium for fresh DMEM. 48 h after viral transduction, cells were washed with PBS, trypsinized, neutralized with DMEM, centrifuged for 3 min at 600 g, and collected by centrifugation and resuspended in 300 µl of PBS. Samples were analysed with a LSRFortessa™ Cell Analyzer (BD) Flow Cytometer.

UV laser micro-irradiation

For the analysis of laser-localized fusion proteins, cells were seeded on µ-slide glass bottom plates (Ibidi). Experiments were imaged with an ORCA Flash 4.0 sCMOS camera (Hamamatsu) using a Yokogawa CSU-W1 Confocal Scanner Unit (Gataca Systems) in an Axio Observer 7 microscope (Zeiss) at 37 ℃ and 5% CO_2 conditions. Laser microirradiation was performed using a laser output of 20% power of a 355nm laser line (10mW; 19kHz, iLAS 3 Gataca Systems) through a UV-transmitting 100-X oil immersion objective (Zeiss). Laser stripes were created at the cell nucleus with a

pixel size of 1, and 2 iterations per stripe using MetaXpress software (Molecular Devices). After irradiation, GFP or Cherry tagged protein accumulation at laser tracks was recorded at the indicated times. Signal intensity was measured using Fiji (ImageJ) by subtracting pre-irradiation signal to each time-point and represented as arbitrary units (a.u.).

Clonogenic cell survival assays

Clonogenic survival assays were carried out seeding 500 cells in 6-well plates triplicates 24 hours prior to treatment. DSBs were produced by ionizing radiation (IR) or by acute treatment with topoisomerase inhibitor etoposide (VP16; Sigma). For IR, cells were exposed to 2 or 4 Gy or mock treated. In the case of VP16 treatments, cells were incubated for 1 h with 5 or 10 μM of the drug or vehicle (DMSO) as control. After two washes with PBS, a fresh medium was added, and cells were incubated at 37 °C for 7–14 days to allow colony formation. Afterward, cells were stained and visualized in the solution of 0.5% Crystal Violet (Merck) and 20% ethanol (Merck). Once the colonies were stained, this solution was removed, and plates were washed with distilled water. The portion of surviving colonies at each dose was calculated by dividing the average number of visible colonies in treated versus control dishes.

ICGC data retrieval and analysis

Patient sets were retrieved from the International Cancer Genome Consortium (ICGC) Data Portal TCGA Breast Cancer (BRCA) datasets. CRY1 and CCAR2 expression levels, survival and radiation therapy information from each donor were obtained using the UCSC Xena web tool.

Subcutaneous xenografts in mice

Xenografts were generated by injecting 8×10^6 cells (HCT116) subcutaneously into flanks of female Athymic Nude (Crl:NU(NCr)-*Foxn1[nu]*) mice (Charles River Laboratories). Each animal was injected with cells depleted for CRY1 (right flank) or control cells (left flank). Tumour volume was estimated according to the formula: (Width^2 × Length)/2. The treatment with etoposide (VP16) started when tumours reached 100-200 mm^3. Mice were injected with etoposide (20mg/kg) or DMSO twice per week and tumour volume and mice weight were measured also twice per week for

20 days. At the end of the treatment, mice were sacrificed, and tumours were removed

to weigh them and acquire images.

ChIP-Seq:

Chromatin Immunoprecipitation was performed as previously described[47].

Briefly, DSBs were induced in AsiSI-ER U2OS cells (which stably express AsiSI-ER)

in low-glucose Dulbecco's Modified Eagle Medium (DMEM) + 10% charcoal-stripped

FBS containing 300 nM 4-OHT for 4h. Then, cells were crosslinked with 1%

formaldehyde for 10 minutes at 37°C. Crosslinking reaction was quenched with 125mM

glycine for 5 minutes and cells were washed 3 times with PBS. Cell pellets were

resuspended in 2.5 mL lysis buffer A (5 mM Pipes pH 8.0, 85 mM KCl, 0.5% NP40,

1mM PMSF) supplemented with protease inhibitors and incubated for 10 minutes on ice.

After centrifugation, nuclear fraction was resuspended in 1 mL of lysis buffer B

(50 mM Tris HCl pH 8.1, 1% SDS, 10 mM EDTA, 1 mM PMSF) supplemented with

protease inhibitors. Chromatin was sonicated using a Bioruptor Pico (Diagenode,

B01060010), 5 cycles of 30 seconds "on" and 30 seconds "off" on ice-cold water. For

each immunoprecipitation, 20-100 μg of chromatin and 4 μg of the indicated antibody

was used in IP buffer (0,1% SDS, 1% TX-100, 2mM EDTA, 20 mM TrisHCl pH8,

150 mM NaCl) at 4°C o/n and then with 40 μL of pre-blocked (1 mg/ml BSA)

Dynabeads protein A (ThermoFisher). Beads were sequentially washed with Wash

buffer 1 (20 mM Tris HCl pH 8.1, 0.1% SDS, 1% Triton x-100, 2 mM EDTA, 150 mM

NaC), Wash buffer 2 (20 mM Tris HCl pH 8.1, 0.1% SDS, 1% Triton x-100, 2 mM

EDTA, 500 mM NaCl) Wash buffer 3 (10 mM Tris HCl pH 8.1, 1% NP-40, 1% NaDoc,

1 mM EDTA, 250 mM LiCl), and twice with TE-buffer (10 mM Tris-HCL pH8, 1 mM

EDTA pH8). ChIPmentation was carried out as previously described[48], using Tn5A

Enzyme provided by the Proteomic Service of CABD (Centro Andaluz de Biología del

Desarrollo). Tagmented DNA was then eluted with 1% SDS in TE at 65°C for

10 minutes and crosslinking was reverted at 65ºC overnight. Protein was degraded with

Proteinase K for 2 hours at 37°C. DNA was purified using QIAGEN PCR purification

Kit (QIAGEN, 28106). Libraries were amplified for N-1 cycles (being N the optimum

Cq determined by qPCR reaction) using NEBNext High-Fidelity Polymerase (New

England Biolabs, M0541). Libraries were purified with Sera-Mag Select Beads (GE

Healthcare, 29343052) and sequenced using Illumina NovaSeq 6000 SP300 cycles

2x150 pb long (paired-end), XP protocol. Raw sequencing files (fastq) quality was

assessed   using   FastQC   0.12.0

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and aligned to the reference human genome (version hg19) using bwa 0.7.17 (https://bio-bwa.sourceforge.net/). Mapped sequences were sorted and indexed using samtools 1.19.2 (http://www.htslib.org). Coverage of mapped reads in bigwig format was obtained using deepTools 3.5.3 bamCoverage function, setting normalization method to CPM. Bigwig files were visualized using Integrative Genomics Viewer 2.17.0. Average ChIP-seq plots of 214 experimentally characterized AsiSI sites[27] were generated using deepTools computeMatrix and plotProfile functions.

Neutral comet assays

U2OS cells were treated with ionizing radiation (10 Gy) and incubated for 24 hours or mock treated. After trypsinization and neutralization, cells were resuspended in PBS at a concentration of 500.000 cells per mL and mixed in a 1:1 ratio with low-melting agarose at 37ºC. 20 uL of cell suspension was dropped onto agarose covered microscope slides and, after drying, slides were incubated in lysis buffer (Trevigen) overnight at 4ºC. After lysis, electrophoresis of the DNA trapped in the agarose was performed at 1 V/cm for 45 min in TEB 1X buffer. DNA was stained with SYBR gold for 10 min and slides were washed in water and further dried at 37ºC. At least 200 cells were analyzed using a Leica DMI8 Thunder fluorescence microscope and tail length was analyzed using ImageJ.

Patient data

Survival data was obtained from the Radiotherapy Service at the University Hospital Virgen Macarena, Seville Spain, from all patients treated with radiotherapy as a primary form of treatment between May 2018 to October 2023. In all cases, in addition to the type of cancer, the gender and age (Suppl. Table 5), the date of treatment initiation, the date of the last known interaction (either the death of the patient or the last known follow up), and approximate time of irradiation was recorded. Each patient was always treated at the same time of the day. Patients were split in two cohorts, according to the time of irradiation, those treated in the first half of the day (8:00 to 14:00; morning) or the second half (14:00 to 20:00).

Romero-Franco et.al. Figure 1

Romero-Franco et.al. Figure 2

Romero-Franco et.al. Figure 3

Romero-Franco et.al. Figure 4

Romero-Franco et.al. Figure 5

Romero-Franco et.al. Figure 6