Evolutionarily diverse fungal zoospores show contrasting swimming patterns specific to ultrastructure.

Luis Javier Galindo[1, *], Thomas A. Richards[1], Jasmine A. Nirody[2, *]

[1] Department of Biology, University of Oxford, Oxford, United Kingdom
[2] Department of Organismal Biology and Anatomy, University of Chicago, Chicago, IL, USA

Author for correspondence:
Jasmine A. Nirody (lead contact) or Luis Javier Galindo
*e-mail: jnirody@uchicago.edu; luisjagg92@gmail.com

Keywords: motility, zoospores, ultrastructure, fungi, taxis

Summary (250 words)
Zoosporic fungi, also called chytrids, produce motile spores with flagellar swimming tails (zoospores)[1,2]. These fungi are key components of aquatic food webs, acting as pathogens, saprotrophs and prey[3–8]. Little is known about the swimming behaviour of fungal zoospores, a crucial factor governing dispersal, biogeographical range, ecological function and infection dynamics[6,9]. Here, we track the swimming patterns of zoospores from 12 evolutionary divergent species of zoosporic fungi across seven orders of the Chytridiomycota and the Blastocladiomycota phyla. We report two major swimming patterns which correlate with the cytoskeletal ultrastructure of these zoospores. Specifically, we show that species without major cytoplasmic tubulin components swim in a circular fashion, while species that harbour prominent cytoplasmic tubulin structures swim in a pattern akin to a random walk (move-stop-redirect-move). We confirm cytoskeleton architecture by performing fluorescence confocal microscopy of the zoospores across all 12 species. We then treat representative species with variant swimming behaviours and cytoplasmic-cytoskeletal arrangements with tubulin stabilizing (Taxol) and depolymerizing (Nocodazole) pharmacological-compounds. We observed that when treating the 'random-walk' species with Nocodazole their swimming behaviour changes to a circular swimming pattern. Confocal imaging of the nocodazole-treated zoospores demonstrates these cells maintain flagellum tubulin structures but lack their characteristic cytoplasmatic tubulin arrangement. These data confirm that the capability of zoospores to perform 'complex' movements as a random walk is linked to the presence of prominent cytoplasmatic tubulin structures. We discuss the link between cytology, sensation, and swimming behaviour manifest in zoosporic fungi.

Results

Differential swimming in variant chytrids

Here we analysed swimming patterns of zoospores from 12 different species from the two major chytrid phyla: Chytridiomycota and Blastocladiomycota. These include eight Chytridiomycota species grouped into six orders: *Synchytrium microbalum* (Synchytriales), *Clydaea vesicula* (Lobulomycetales), *Chytriomyces hyalinus* (Chytridiales), *Rhizoclosmatium globosum* (Chytridiales), *Homolaphlyctis polyrhiza* (Rhizophydiales), *Rhizophlyctis rosea* (Rhizophlyctidiales), *Geranomyces variabilis* (Spizellomycetales), and *Spizellomyces punctatus* (Spizellomycetales); and four species of Blastocladiomycota from the order Blastocladiales: *Catenophlyctis* sp. (Catenariaceae), *Blastocladiella emersonii*, *Allomyces macrogynus* and *Allomyces reticulatus* (Blastocladiaceae).

We find that medium-range motility patterns varied markedly among different species (Figure 1), with the zoospores from species of four Chytridiomycota (e.g., *S. microbalum*, *C. hyalinus*, *R. globosum* and *H. polyrhiza*) swimming in a circular mode. In contrast, all zoospores from species of Blastocladiomycota tested and four different species of Chytridiomycota (e.g., *C. vesicula*, *R. rosea*, *G. variabilis* and *S. punctatus*) swim using a pattern akin to a random walk (move-stop-redirect-move; Figure 1A). To investigate this pattern further, we plotted the distribution of reorientation (turn) angles and the mean square displacement (MSD) of these 12 movement patterns to characterize motility dynamics at both shorter and longer timescales, respectively (see the methods section for the mathematical formulae). Such methods can be used to quantitatively differentiate random walk and circular motility patterns[10,11]. Circular swimming cells are expected to reorient themselves more consistently than those swimming in a random walk-like pattern, resulting in a narrower turn angle distribution. For cells exhibiting circular motility patterns, the MSD plot should form a plateau in the later phases of the MSDs plots. Consistent with the conclusions from our primary observations (Figure 1A), we find that reorientation angle distributions are indeed generally tighter for zoosporic fungi without prominent tubulin cytoplasmic structures (Figure 1B), and that the MSD plateaus at shorter timescales for these species in comparison to the other zoosporic fungi (Figure 1C). We note that the random-walk swimmer *A. macrogynus* (*Am*) shows a narrower distribution and the circular swimmer *C. hyalinus* (*Ch*) shows a broader angle range than expected. This is because *Am* tends to swim in more regular, directed trajectories than other random-walkers; this is also apparent from its MSD which scales with a higher exponent than that of other species (the inconsistencies of this species are discussed further below). On the other hand, *Ch* is the circular-swimmer with the most variable radii, resulting in a wider turn angle distribution. We expect that differences in local behaviours will be further heightened in experiments taken at

greater resolution (e.g., using a camera with a higher frame rate). These results provide
support for a primary conclusion that zoosporic fungi show two distinct swimming patterns.
Apart from slightly higher speeds observed in *H. polyrhiza*, no significant differences in
instantaneous swimming speed were observed across the 12 species (Figure S3A),
suggesting the difference in movement behaviour is a product of cellular mechanics and not
propulsion velocity.

The relationship between cell structure and swimming pattern

To gain insights into the cellular mechanisms which putatively underpin these differences in
swimming patterns, we analyse the intracellular organisation of zoospores. We performed
fluorescent confocal microscopy with antibodies and cytological stains against α-tubulin,
actin and lipid components of the zoospores from all 12 species studied (Figure 2). To
further support our observations we reviewed previous electron microscope ultrastructural
studies for each of these species[12–19] (Table S1). These results suggest that observed
variation in motility patterns segregate with cytological characters, specifically the presence
of prominent cytoplasmatic tubulin structures, and not by higher taxonomic classifications
(Figures 2, 3A).

In all four Blastocladiomycota species, we observe random-walk movement patterns
and show that the cytoplasmatic tubulin microtubules create prominent and complex
cytoskeletal arrays or 'ribs' within the cell body which extend from the kinetosome of the
flagellum around the periphery of the main body of the cell. Our microscopy shows that an
actin cytoskeleton is much less apparent (Figure 2, 3A; n ≥ 10 cells observed for each
species). Similarly, for two Chytridiomycota species that show random-walk movement
patterns, *S. punctatus* and *G. variabilis* (Spizellomycetales) we also observe prominent
tubulin microtubules that radiate at random into the zoospore cytoplasm from the kinetosome
(n ≥ 10) i.e., forming rib-like structures similar to those seen in the Blastocladiomycota. For
these Spizellomycetales, actin is present and forms patches in both species (the presence of
these patches indicates possible actin organisation centres[20,21]).

Two additional Chytridiomycota species also showed prominent accumulations of
non-assembled cytoplasmic α-tubulin-stained material; *R. rosea* and *C. vesicula*. Although
these were not organised as clearly defined ribs in our micrograph analyses, such cell forms
also performed random-walk movement behaviours. The zoospores of *R. rosea* (n = 10)
showed both seemingly randomly arrayed α-tubulin accumulations and actin patches
through-out the cytoplasm. Interestingly, microscopy imaging of *C. vesicula* identified an
additional distinct α-tubulin organisation consisting of a peripheral cytoplasmic mesh of
microtubule structures (n = 10). In *C. vesicula* we observe cytoplasmic actin co-localizing
with the canonical vesicles found near the kinetosome of this species[15] (Figure 2).

In contrast to the previously described species, we find no tubulin structures in the
cytoplasm of *S. microbalum* (Synchytriales) and no major structures the other three
Chytridiomycota: *H. polyrhiza* (Rhizophydiales), *C. hyalinus*, and *R. globosum* (Chytridiales)
beyond a single microtubule structure connecting the base of the flagellum to the microbody-
lipid globule complex organelle (MLC) only observed in TEM studies (n ≥ 10 cells observed
for each species) (Figure 2, 3A, Table S1). Furthermore, in all four zoospores forms actin is
the only detected cytoplasmic-cytoskeletal protein using our current confocal microscopy
protocol. These four species swim in a circular pattern.

As for the lipid organelles, in Chytridiomycota we consistently observed one large
lipid globule (n ≥ 10 – for each of these eight species) in their MLCs, with one exception
found in the chytrid *R. rosea* in which we identified numerous small lipid droplets, a
characteristic of this species[18] (Figure 2). In the Blastocladiomycota we consistently
observed clusters of several lipid globules (n ≥ 10 – for each of these four species), usually
located towards the flagellum as part of the side-body complex organelle (SBC). In the case
of *A. reticulatus* the lipid droplet clusters are numerous and run down the entire length of one
side of the zoospore's cell body (Figure 2).

Our video imaging results identified that the random walk patterns are specifically
characterized by straight movements, followed by a stop and then a redirection of the
cellular body. Observations of movement patterns for *B. emersonii* (*Be*) led to the hypothesis
that re-direction occurs by a hinge-like bending of the area between the cytoplasmic
microtubule and the flagellar kinetosome, and in an angle greater than the final angle of
movement (Figure 3B). By performing live-fluorescent imaging of zoospores after staining
with NileRed (lipid stain; n = 50) we observed that in *Be* the bend and subsequent
redirection occurs to one side of the cell, the side where the SBC is located; suggesting this
organelle is a constituent part and/or determinant of the hinge articulation which determines
swimming redirection (Figure 3B).

Pharmacological inhibition demonstrates tubulin structures determines how random
walker cells reorient
To test the involvement of tubulin cytoplasmatic structures in their swimming patterns, we
treated representative species of each swimming behaviour and cytoplasmic cytoskeletal
type with 1 µM of tubulin stabilizing (Taxol) and depolymerizing (Nocodazole) drugs. We
selected *B. emersonii* (*Be*; Blastocladiomycota) and *S. punctatus* (*Sp*; Chytridiomycota)
given their prominent microtubular ribs and marked random-walk movement, and *R.*
globosum (*Rg*; Chytridiomycota) due to its lack of major α-tubulin cytoplasmic structures,
circular swimming and its increasing utility as a model system[22]. We observed that when
treating random-walk species *Be* and *Sp* with Nocodazole their overall swimming behaviour
changes to manifest a circular swimming pattern (Figure 4A). However, there was no

observable change in the swimming behaviour of the circular swimming *Rg* zoospores

treated with nocodazole or taxol. Additionally, after imaging these nocodazole-treated and

taxol-treated zoospores, we confirmed that nocodazole-treated zoospores lack their

characteristic tubulin 'rib' cytoplasmatic arrangements, while there was no change in the

tubulin arrangement of the taxol-treated species (Figure 4B-G). We also note that for the

nocodazole-treated *Be* zoospores the lipid component (an organelle hypothesised to be key

for the light sensing function[23,24]), which is usually located next to the base of the flagellum,

loses its posterior cellular position and is relocated to random positions within the cell. Lastly,

losing their tubulin cytoplasmic structures caused the zoospore cell morphology to change,

from a tear-shape morphology towards a sphere-shape in *Be* and a reversed tear-shape in

Sp (Figure 4C-F). Besides a slightly shorter flagellum in *Be*, we did not observe major

changes in the tubulin structures of the flagellum across a range of drug treatments for the

three fungal zoospores tested.

We performed an additional series of controls to test the involvement of actin in

zoospore swimming behaviour by using actin depolymerizing drugs (Cytochalasin D,

Latrunculin B and Jasplakinolide). As observed previously on *Batrachochytrium*

dendrobatidis[25] these drugs caused mass zoospore encystation in these three species, and

no change in the swimming behaviour of the few remaining swimming zoospores was

observed (Figures S4, S5).

Discussion

An association between zoospore swimming pattern and cytoplasmic tubulin

structures

Swimming patterns have been studied in several unicellular eukaryotes (e.g.,[26–29]). Zoospore

swimming patterns in the protist *Phytophtora*, for example, have been shown to contribute to

their dispersal, host colonisation and activity as plant pathogens[29,30]. These findings have led

to the suggestion that chemical inhibition of motility could be used as a strategy for

controlling the spread of pathogenic zoosporic species[29]. However, little is known about the

swimming behaviour of fungal zoospores despite the crucial role of this life-cycle stage for

dispersal and therefore host infection[5].

Most chytrid species form uniflagellated zoospores, with only a few known

multiflagellated species (e.g., *Neocallimastix* and *Piromyces*[31]) and lineages that form spores

without flagella (due to a secondary loss in *Hyaloraphidium*[32]). Zoospores can move using

two distinct mechanisms: swimming propelled by beating of the tubulin articulated

flagellum[3,33] or amoeboid crawling using actin driven pseudopodia[34,35]. These different

locomotive modes are suited for specific behavioural goals; as such, there are fungal

species which are both swimmers and crawlers (e.g., *Paraphysoderma*[35]), species with

swimming zoospores which under certain conditions will switch to crawlers (e.g.,

Batrachochytrium dendrobatidis[20]) or with zoospores that exclusively crawl (e.g.,

Sanchytriomycota[34]). Broadly speaking, crawling is better adapted for traveling short

distances on surfaces in liquid environments (e.g., exploration of trophic substrate);

however, flagellum-based swimming is ideal for medium range dispersal in larger volumes of

H_2O[9] (e.g., exploration of a water column to colonise a niche habitat). Thus, swimming plays

an important role in facilitating the trophic life cycle of these fungi, a function that is often

governed by chemo- or photo-taxis function[23,36–38] to enable navigation towards, and

colonisation of, niche habitats. These functions therefore drive the mycoloop[5,7,39].

To our knowledge, this is the first systematic study of swimming patterns across a

wide representation of the chytrid fungi with the patterns identified placed into a cellular and

phylogenetic context (Figure 3A). Here we report two distinct swimming patterns among

different fungal zoospore forming species: Chytridiomycota zoospores swim both in circular

tracks (n = 4 spp), and in random-walk movement (n = 4 spp), while the Blastocladiomycota

zoospores studied adopt a random-walk movement pattern (n = 4 spp). These patterns have

previously been noted in a subset of chytrid fungi, providing additional data consistent with

the pattern outlined here[40]. Specifically, circular swimming in the zoospores of the

Rhizophydiales chytrid *Rhizophydium planktonicum*[41] and random-walk swimming patterns

in the zoospores of the Blastocladyomycota *Paraphysoderma sedebokerense*[35] and

Physoderma spp.[42] from the order Physodermatales. These additional results suggest that a

wider diversity of Blastocladiomycota species swim like the Blastocladiales studied here.

To address whether these swimming patterns form a consistent correlation with key

cellular ultrastructure, we first perform fluorescence confocal microscopy of zoospores from

all 12 species (Figure 2). Using this approach, we observed prominent cytoplasmic tubulin

structures in all four Blastocladiomycota species and in four of the Chytridiomycota species

analysed in this study which swim using the random walk patterns (Figure 3A). Similar to

Blastocladiomycota, the Spizellomycetales Chytridiomycota fungi possess a complex

tubulin-based cytoplasmatic structures with radiating microtubules ('ribs'), as shown in

previous ultrastructural TEM-based studies[16,19,43] (Table S1). Despite also being prominent in

our images, neither the cytoplasmatic unassembled tubulin structures observed here in the

zoospores of *R. rosea,* nor the peripheral tubulin based cytoskeleton observed in *C. vesicula*

were previously detected in TEM-based descriptions[15,18]. However, the strains of *R. rosea*

(JEL0764) and *C. vesicula* (JEL0476) used, are different to the strains used for the original

descriptions (Barr186/BR186 and JEL369 respectively[15,18,44]) so some variation in

cytoskeletal structure may be present within these species complexes. *R. rosea* zoospores

also have a large fibrillar rhizoplast extending from kinetosome which may fulfil a related

function by acting as a hinge during the re-direction of movement[18] (Table S1). Additionally,

all these observations were made using TEM[15,18], thus, these analyses would fail to detected

non-electron dense structures, specifically failing to identifying dissociated tubulin material

like the cytoplasmic tubulin structures detected in our analyses of these two species (Figure

2). The demonstration that some tubulin-based material can be observed in TEM, like the

microtubules of Blastocladiomycota/Spizellomycetales (rib-like tubulin arrangements), while

in others cytoplasmic tubulin can only be observed using fluorescent staining (*C. vesicula*

and *R. rosea*); represents evidence of key differences between the cytoplasmic tubulin-

based structures of these different zoosporic species. It is important to note that we refer to

all these cytoplasmic tubulin structures as 'prominent' because we are aware that in the

circular-swimming zoospores of *C. hyalinus*, *R. globosum* and *H. polyrhiza* there is a minor

cytoplasmic tubulin component in the form of small bundles of microtubules connecting the

kinetosome with the lipid-organelle's rumposome (Figure 3A), which have also been

observed using TEM[45] (Table S1).

Taken together, our analyses confirm that the presence of prominent cytoplasmatic

tubulin of various arrangements and densities correlates with the different swimming

patterns observed. Specifically, zoospores from Chytridiomycota species that lack prominent

cytoplasmatic tubulin structures swim using a circular pattern. In contrast, in all four

Blastocladiomycota species and in those Chytridiomycota species in which there are

prominent α-tubulin-stained structures within their cytoplasm, we observe a random-walk

swimming pattern (Figure 3A).

Next, we sought to experimentally test the correlation between swimming behaviours

and cytoplasmic tubulin structures by pharmacologically inhibiting or stabilizing intracellular

tubulin using nocodazole and taxol, respectively. These drugs have successfully been used

on several microbial eukaryotes to study cytoplasmic and flagellar tubulin-based

structures[46–50]. Our results show that zoospores of fungal species with prominent

cytoplasmic tubulin structures treated with nocodazole shifted their swimming patterns from

random-walk to circular pattern (Figure 4). We note neither drug stopped flagellum

swimming and in all three test species tubulin structures of the flagellum showed no major

structural changes besides a putative and minor reduction in flagellar length in nocodazole-

treated *Be* (Figure 4). Flagellar length does not seem to be factor determining the observed

swimming patterns. All described zoospores vary in their flagellar length regardless of their

swimming patterns, and species with the same flagellar length (e.g., ~20 µm) can present

either circular (e.g., *C. hyalinus*) or random-walk patterns (e.g., *S. punctatus*) (Table S1). In

our test species *Rg* retains a long flagella and circular swimming across all treatments and

after nocodazole treatment *Sp* retains its flagellar length, but the movement behaviour is

transformed to a circular swimming mode. However, our results do demonstrate that we

succeeded in inhibiting the polymerization/formation of 'ribs' within the main body of *Be* and

Sp zoospores which became circular swimmers after treatment (Figure 4). Thus, we
conclude that the capability of zoospores to perform 'complex' movements in a random walk
fashion is linked to the presence of prominent cytoplasmatic tubulin structures. These
findings demonstrate that a prominent tubulin-cytoplasmatic component may allow
zoospores to swim using intricate patterns in which they can redirect movement by making
sudden changes in direction; the random-walk pattern. These microtubular structures that
radiate from the base of the flagellum may contribute to this pattern by forming a hinged
articulation which bends the cell to re-orientate direction of travel towards a new direction of
movement (Figure 3B). On the other hand, cells lacking tubulin cytoplasmic structures may
restrict zoospores to swim using a constant circular/spiral-like pattern of movement.

Is movement pattern connected to light-sensing and the lipid organelle in these
zoospores?

Fungal zoospores possess different lipid organellar arrangement globally referred to as the
microbody-lipid globule complex (MLCs) and are structures known to be involved in storage
and production of chemical energy for flagellar beating[45]. There are different 'types' of MLCs
which vary in structural arrangement across chytrid species. These structures are composed
of microbodies, mitochondria, ribosomes and one or more large membrane-enclosed lipid
globules (Figure 2, 3A)[2,45]. Using Powell's (1978) classification[45] all MLCs discussed here
from Chytridiomycota are type 1, which can be classified as A or B; depending on the
respective absence or presence of a rumposome (a cisterna-like disc of membranous
tubules); zoospores can be further classified as A/B_1 or A/B_2 regarding the arrangement of
different cellular components. The lipid-filled MLC organelles from Blastocladiomycota are
type 4, and are alternatively classified as side-body complexes (SBCs) (Figure 2, 3A)[1,13,45,51].

In addition to energetics, these organelles have been hypothesised to be involved in
light perception[23,24,52,53]. Indeed, *Blastocladiella emersonii* and *Allomyces reticulatus*
zoospores show positive phototaxis[23,37,54,55]. In particular, the SBC of *B. emersonii* was found
to be in close spatial association with a layer of RGC (or CyclOp) proteins (a fusion protein
composed of a type I microbial rhodopsin and guanylyl cyclase enzyme domain) which
participates in a cGMP-mediated light-sensing pathway[23,56,57] by controlling flagellar
activation from its localization next to the SBC. Both these systems are in close physical
association at the base of the flagellum. Recent analyses suggest that the molecular
elements necessary for a functional RGC based light sensing pathway are present in the
eight of the twelve fungal species analysed here[52,58,59]. These include six species with
homodimeric RGC variants (all four Blastocladiomycota species, *S. microbalum* and *H.*
polyrhiza) and two species with heterodimeric RGC complexes (Chytridiales: *C. hyalinus*, *R.*
globosum) (Figure 3A), which consist of two subunits with distinct photochemical properties
which evolved from an ancient duplication of the RGC protein[58,59]. We observe no

association between RGC type, swimming pattern and ultrastructure. However, within the
Chytridiomycota analysed here, those species that swim in circular patterns possess RGCs
and MLCs type 1B and those with random-walk movement lack RGCs and have MLCs type
1A (Figure 3A), but wider sampling is needed to further test this pattern.

We see no direct correlation between number of lipid structures (one large lipid body
vs multiple smaller lipid bodies) and either swimming pattern or possession of the
heterodimeric RGC system. This result infers that the lipid body structure/morphology is not
a factor in determining variant swimming behaviours or variance in light-perception
mechanisms. However, some swimming behaviours may be influenced by the presence of
the lipid structure. *Be* produces zoospores with the RGC system and that swim in a random-
walk pattern and the SBC is found at one side of the kinetosome. Our experiments show that
the cell-bending and redirection of motion in *Be* zoospores occurs always towards the side of
the cell in which the SBC is present (Figure 3B). Among the possible explanations could be
that prominent SBC is structurally restraining the movement of the *Be* zoospores towards
that side, or that the zoospore moves towards the direction in which the light sensing
organelle gets its stimulus. The position of MLCs relative to their movement is harder to
assess with the random-walker Chytridiomycota species (e.g., *Sp*), since their MLCs are
found in the centre of the cell and are unattached from the kinetosome. In circular swimming
chytrids, small cellular sizes and large MLCs complicate the task of determining the position
of MLCs relative to their circular motion, thus they remain to be characterized. Lastly,
fluorescent confocal microscopy revealed that nocodazole-treated circular-swimming *Be*
zoospores not only lacked cytoplasmatic tubulin microtubules, but they also loss the typical
arrangement of the SBC organelle at the base of the flagellum, providing further evidence
that both structures interact through the rib-like tubulin structures (Figure 4C). The physical
link between the SBC and the base of the kinetosome may be essential for the transduction
of the signal to flagellar movement which then translates to the observed swimming pattern.

Regarding a light-perception/swimming pattern relationship, we show that *A.*
macrogynus (*Am*) tends to swim in more regular, directed trajectories than other random-
walkers; as shown by its narrower turn angle distribution and its MSD scaling with higher
exponent (Figure 1B-C). One possible explanation is that *Am* is the only Blastocladiomycota
in our sample known to have replaced light sensing with chemo sensing[37]. Thus, if our set up
is triggering light sensing (all observations are under white light) but not chemo sensing (no
amino acid gradients), we could hypothesize that light sensation is constantly triggering
reorientation in random walker Blastocladiomycota except in *Am*, which has no response to
light. Overall, these experiments point towards a link between zoospore ultrastructure,
movement, and the light sensing organelle component. Our findings provide comparative

model systems for understanding the link between cytology, sensing and divergent motility behaviour in single-celled eukaryotic microbes.

Acknowledgements

This work was funded by the Horizon 2020 research and innovation programme under the European Marie Skłodowska-Curie Individual Fellowship H2020-MSCA-IF-2020 (grant agreement no. 101022101—FungEye) and Leverhulme Trust Project Grant (RPG-2022-234; The evolutionary diversification of a sub-cellular fungal eye). T.A.R. is supported by a Royal Society URF (URF/R/191005). We gratefully acknowledge the Micron Advanced Bioimaging Facility (supported by Wellcome Strategic Awards 091911/B/10/Z and 107457/Z/15/Z). We also acknowledge Suely Lopes Gomes (Universidade de São Paulo) for inspiring work on chytrid light perception. J.A.N. acknowledges support from All Souls College, Oxford and the National Science Foundation through the Center for Living Systems at the University of Chicago (grant no. 2317138). We are also grateful for the comments of three anonymous reviewers, which have improved this manuscript from its original form.

Author contributions

L.J.G., T.A.R. and J.N.: conceived the study; L.J.G.: microscopy imaging, drug assays and manuscript writing; T.A.R. manuscript writing; J.N.: imaging analyses and manuscript writing. All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Declaration of Interests

These authors declare no competing interests.

Figure 1. Movement tracks of zoospores of 12 zoosporic fungal species. (A) Movement
tracks show cytology-specific swimming pattern behaviours, with all Blastocladiomycota
species and Chytridiomycota species possessing prominent tubulin-based cytoplasmic
structures swimming in a random walk pattern, while Chytridiomycota zoospores without
prominent cytoplasmic tubulin structures swim in circular movements. Composite tracks are
shown compiled into a single plot with starting points (time = 0) centred at the point of origin.
(B) Distribution of turn (reorientation) angles for each group of zoospores
(Blastocladiomycota species, Chytridiomycota species with prominent tubulin-based
cytoplasmic components, and Chytridiomycota species with no prominent tubulin-based
cytoplasmic structures). Species-averaged distributions are shown as partially transparent
lines, as labelled. (C) Ensemble- and time-averaged mean-squared displacement (MSD) for
each group of zoospores. Species-averaged MSDs are shown as partially transparent lines,
as labelled. For effective comparison between groups, MSD plots are normalised by the
initial value (at time lag 1) for each species. Unnormalized MSDs for each cell are shown in
Figure S3B. Labels: *Catenophlyctis sp.: Cat, Blastocladiella emersonii: Be, Allomyces
macrogynus: Am, Allomyces reticulatus: Ar, Synchytrium microbalum: Sm, Clydaea vesicula:
Cv, Chytriomyces hyalinus: Ch, Rhizoclosmatium globosum: Rg, Homolaphlyctis polyrhiza:
Hp, Rhizophlyctis rosea: Rr, Geranomyces variabilis: Gv,* and *Spizellomyces punctatus: Sp.*

Figure 2. Ultrastructure of Blastocladiomycota and Chytridiomycota zoospores. First
columns show brightfield phase contrast confocal microscopy images; second columns show
fluorescent confocal microscopy micrograph of the zoospore; third columns show the

merged images of brightfield and fluorescent channels. Microscopy images were taken and

stained using the methods described in Galindo et al. (2022)[52] with NileRed to stain lipid

droplets, α-tubulin DM1A + Alexa Fluor 647 to stain tubulin and Alexa Fluor 488 Phalloidin to

stain actin. In micrographs α-tubulin structures are stained magenta, actin stained cyan, and

the lipid bodies (proposed to function as a light-sensing eye-spot) are stained yellow. In the

fluorescent channel panels for those species which possess cytoplasmatic α-tubulin staining

there is a small panel detailing the α-tubulin component. Species/micrographs

bracketed/highlighted with red - belong to the Blastocladiomycota and those species

micrographs with green-to-pink highlight-brackets correspond to Chytridiomycota from the:

Synchytriales (M1-O1), Lobulomycetales (P1-L1), Chytridiales (M1-R1), Rhizophydiales (S1-

U1), Rhizophlyctidiales (V1-X1) and Spizellomycetales (Y1-D2), respectively (the same

colour code is used in Figure 3). Scale bars: A1-O1 = 10 μm; P1-D2 = 5 μm. Scale bars in

small α-tubulin panels: B1, E1, H1 = 10 μm; K1, Q1, W1, Z1 and C2 = 5 μm. Labels:

Catenophlyctis sp.: Cat, Blastocladiella emersonii: Be, Allomyces macrogynus: Am,

Allomyces reticulatus: Ar, Synchytrium microbalum: Sm, Clydaea vesicula: Cv, Chytriomyces

hyalinus: Ch, Rhizoclosmatium globosum: Rg, Homolaphlyctis polyrhiza: Hp, Rhizophlyctis

rosea: Rr, Geranomyces variabilis: Gv, and *Spizellomyces punctatus: Sp.*

Figure 3. Depiction of motility, ultrastructure and phylogenetic relationships of the
zoosporic fungal species from this study. (A) Phylogenetic relationships adapted from
the multigene phylogeny from Amses et al. 2022[60] showing the absence (black branches) or
presence of homodimeric (yellow branches) or heterodimeric (magenta branches) RGCs
opsin complexes adapted from Broser et al. 2023[59]. Drawings depict the overall
ultrastructure of each zoospore including their lipid, actin, and α-tubulin-based components.
A representative trajectory highlighting the overall movement pattern of each species is
shown to the right of each drawing and grouped by a continuous (circular pattern) or
discontinuous line (random-walk pattern). Number in parentheses bellow the species names
and main branches indicate the lipid-organelle MLC classification of each zoospore
according to Powell's (1978) classification[45]. See Table S1 for further references. (B)
Reconstruction based on our finding of the random-walk pattern of *B. emersonii* zoospores.
411

412
Figure 4. Movement tracks and confocal imaging of fungal zoospore from three species after treatment with Nocodazole and Taxol. (A) Representative swimming patterns and ensemble- and time-averaged mean-squared displacement (MSD) for each

treatment in three zoospores fungal species. (B1-G5) Confocal imaging of drug treated zoospores. The first column shows brightfield phase contrast confocal microscopy images (B1-G1); second column shows in yellow the channel for NileRed staining of lipid droplets (B2-G2); third column shows in magenta α-tubulin DM1A + Alexa Fluor 647 staining of α-tubulin (B3-G3); fourth column shows the fluorescent channel merged (B4-G4); fifth column shows the merged images of brightfield and fluorescent channels (B5-G5). Scale bars = 10 µm. *Blastocladiella emersonii: Be, Spizellomyces punctatus: Sp, Rhizoclosmatium globosum: Rg.*

STAR Methods

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Jasmine A. Nirody (jnirody@uchicago.edu).

Materials availability

This study did not generate any new or unique reagents.

Data and code availability

All data are available in the figures, tables, and data files associated with this manuscript. Plots of all tracks for each species are shown in Supplementary Material. Raw videos are available at https://figshare.com/account/home#/projects/207943 and tracking data, detailed protocols, and analysis codes are provided at http://github.com/jnirody/fungalzoospores. Raw confocal microscopic images can be found at: https://figshare.com/projects/FungEye/132734. Any additional information required to reanalyse the data reported in this work paper is available from the Lead Contact upon request.

Experimental model and subject details

Cultures of *Blastocladiella emersonii* ATCC 22665 (ATCC; American Type Culture Collection), *Catenophlyctis* sp. JEL0575 (CZEUM; Collection of Zoosporic Eufungi at University of Michigan; 18S rDNA sequence could indicate it belongs to a species of *Catenaria*[44]), *Clydaea vesicula* JEL0476 (CZEUM), *Chytriomyces hyalinus* CBS 675.73 (previously referred as *C. confervae* given that CBS 675.73 = ATCC 24931 = Barr 97 and Barr 97 is *C. hyalinus* [44,61]; Westerdijk Fungal Biodiversity Institute), *Homolaphlyctis polyrhiza* JEL0142 (CZEUM), *Rhizophlyctis rosea* JEL0764 (CZEUM), *Geranomyces*

variabilis JEL0559 (CZEUM), and *Spizellomyces punctatus* SW1 (CZEUM) were

vegetatively grown in 25 cm2 culture flasks (Sarstedt) filled with 25 mL of PYG liquid media

(0.13% w/v peptone, 0.13% w/v yeast extract, 0.3% w/v glucose). *Rhizoclosmatium*

globosum JEL0800 (CZEUM) vegetative cells were grown on PYG agar plates (0.13% w/v

peptone, 0.13% w/v yeast extract, 0.3% w/v glucose, and 1.5% w/v agar). *Allomyces*

macrogynus Australia_3 (CZEUM) and *Allomyces reticulatus* ATCC 42465 (ATCC) was

grown in 25 cm^2 culture flasks filled with 25 mL of Emerson YpSs/4 liquid media (0.1% w/v

yeast extract, 0.375% w/v soluble starch, 0.025% w/v dipotassium phosphate, 0.01% w/v

magnesium sulfate). *Synchytrium microbalum* JEL0517 (CZEUM) vegetative cells were

grown on PYG agar plates (0.13% w/v peptone, 0.13% w/v yeast extract, 0.3% w/v glucose,

and 1.5% w/v agar) and in PYG liquid media (0.13% w/v peptone, 0.13% w/v yeast extract,

0.3% w/v glucose). All fungal vegetative growth was performed at 21°C and transferred

every two weeks by inoculating 25 µl of previous culture to a new flask/plate containing 25

ml of media.

To induce sporulation of *B. emersonii* ATCC 22665, *C. vesicula* JEL0476, *C. hyalinus*

CBS 675.73, *H. polyrhiza* JEL0142, *R. rosea* JEL0764, *G. variabilis* JEL0559, and *S.*

punctatus SW1, 200 mL of liquid PYG were inoculated with 20 ml of vegetative cells in 175

cm^2 culture flasks (Sarstedt), and incubated, with 150 rpm agitation, for 24 h at 21°C. To

induce sporulation in *A. macrogynus* Australia_3, *A. reticulatus* ATCC 42465 and

Catenophlyctis sp. JEL0575 vegetative growth was performed for ~5 days in 25 ml of liquid

YpSs/4 (*Allomyces*) or PYG (*Catenophlyctis*). Sporangia were then separated from the

culture using sterile tweezers, washed with distilled H_2O, and placed in 20 mL of distilled

H_2O in a 25cm^2 culture flask at 21°C overnight. *R. globosum* JEL0800 and *S. microbalum*

JEL0517 zoospore obtention was performed by growing this strain on PYG agar plates for 4

days. Plates were then flooded with 7 mL of distilled H_2O and after resting for 15 min to allow

sporulation, zoospore-containing H_2O was transferred to a 15 ml falcon tube. Zoospores

were then separated from sporangia using a 20 µm pluriStrainer (pluriSelect) into 50 ml

falcon tubes. All sporulation inductions were performed at 21°C.

Method details

Images acquisition

Out of the 12 obtained liquid zoospore solutions, seven were PYG-based and five were H_2O

-based (*A. macrogynus*, *A. reticulatus*, *Catenophlyctis* sp., *S. microbalum* and *R. globosum*).

After sporulation, both 7 µl and 100 µl of zoospore suspension aliquots were placed

respectively on a slide and coverslip or placed on the bottom of the well of a flat bottom 86-

well plate. *S. microbalum* was recorded only on plates and *A. reticulatus* culture died before

plate videos could be recorded. Thus, we image zoospores both in a physically constrained

and non-constrained environment; we observed no difference between the two conditions
(Figure S6A). Alternatively, Nile-red (Thermo Fisher Scientific, Cat# N1142) was added in a
1:500 v/v concentration for 5 min before live-imaging of zoospores. Cells were imaged on an
Olympus CKX53 inverted microscope with 10X, 20X, and 40X objectives according to the
size of the zoospores and/or their swimming trajectories. Videos of 15 to 30 seconds were
taken at 60 fps with a Std Chromyx HD camera mounted on a U-TV0.5XC-3-8 0.5x C-Mount
Adapter. Exposures, focus, and stage position were kept constant while recording for motility
analyses. The number of trials *n* recorded for each species is shown in Figure 1A.

Both PYG media and H_2O have similar physical conditions and the use of different
liquid media did not seem to affect the swimming behaviours of closely related species with
similar cellular structures (e.g., circular swimming: *R. globosum* in H_2O and *C. hyalinus* in
PYG medium; random walk swimming: *A. reticulatus* in H_2O and *B. emersonii* in PYG
medium). We note that given the small size of the zoospores, minor changes in media
viscosity are unlikely to affect cell mechanics. However, it is possible that environmental
chemistry and the presence of different ion concentrations in the aqueous environment may
modify swimming behaviour[62,63]. To explore this source of variation further, we re-recorded
the swimming behaviour of zoospores of the 7 species which were originally recorded in
PYG media also in distilled H_2O media (by filtering and re-suspension in dH_2O). We
observed no change in the swimming patterns when moved to alternative media (Figure
S6B).
510

Tracking and motility analysis
Movies of swimming zoospores were viewed in FIJI[64], and movement was tracked using a
semi-automated protocol with the TrackMate plugin[65]. Before tracking, videos were pre-
processed as follows: (1) the background was subtracted; (2) artifacts were detected and
removed by subtracting the median image of the video from each frame. Cells were
segmented using TrackMate's Difference of Gaussian detector; manual quality control by
eye was performed on the first frame for each video. The centre-of-mass of segmented
images were joined to create tracks using a Linear Assignment Problem (LAP) tracker;
maximum gap closing was set at 50 pixels and maximum frame gap at 10 frames. Final
quality control of all tracks was performed using TrackScheme to ensure each track
consisted of only one detected spot per frame.
522

Drug assays
Drugs assays were conducted by exposing cultures of *B. emersonii*, *S. punctatus* (grown
overnight) and *R. globosum* (grown for 4-days) for 5 hours at 21°C to 1 µM of Nocodazole
(tubulin polymerization inhibition) and separately Taxol (microtubule stabilizing) solution

resuspended in DMSO. Zoospores were strained-filtered from sporangia as described before

and then fixed in 4% *w/v* paraformaldehyde (for confocal imaging) and/or video recording.

Aliquots of zoospore solution of these three species were also exposed for 10 min at room

temperature to 1 µM Latrunculin B (a drug which sequesters actin monomers), 100 µM

Cytochalasin D (a drug which caps actin filaments), 10 µM Jasplakinolide (a drug which

stabilizes actin filaments) solutions and then resuspended on DMSO for movement

behaviour video recording. Protocols and drug concentrations were based on previous

studies using these same drug treatments on either zoosporic fungi and/or alternative

microbial eukaryotes[25,46–50,66].

Quantification and statistical analysis

Plotting of tracks and quantitative motility metrics were performed from exported COM data

using in-house scripts written in Python. Random walk and circular motility patterns were

quantitatively differentiated using reorientation angle (or turn angle), a measure of local

cellular behaviour, and mean square displacement (MSD), a measure of the deviation of the

position of a cell r over time. Turn angles at time t were calculated as: $\delta\Theta = \tan-$

$1\,[\mathbf{v}(t) \times \mathbf{v}(t+1))\,/\,(\mathbf{v}(t) \bullet \mathbf{v}(t+1)]$. The MSD of a cell at time t is calculated by $MSD(\tau) = <$

$r(t+\tau) - r(t) >,$ as a function of lag time τ. MSD values are calculated across the entire

trajectory for lag times τ ranging from 1 to 800 frames (0.017 to 13.3 s). Cells exhibiting

circular motility patterns are expected to show plateaus at long-time MSDs. Plots of time-

averaged MSD for each cell are shown as grey lines in Figure S16; ensemble-averages are

shown for each species in red. For interspecies comparisons (Figure 1C), ensemble-

averaged MSDs for each species were normalised by dividing by the MSD value at $\tau = 1$.

Fluorescence assays

In all cases, zoospores were collected from the suspension medium by centrifugation at

1000 x g for 5 min followed by removal of the supernatant in order to concentrate the cells.

In all cases the cell pellet was fixed in 0.5 ml of 4% *w/v* paraformaldehyde in 1X PBS and

transferred into a 15 ml Falcon tube for 15 min at room temperature. Cells were

concentrated by centrifugation and resuspended in PBS for the first washing step. The cell

pellet was then resuspended and permeabilized in 0.5 ml of PBS containing 0.1% *v/v* Triton

X-100 in PBS. A second washing step in 0.5 ml of PBS was performed and then the cells

were blocked with 0.5 ml of 1% *w/v* BSA in PBS and incubated for 45 min at room

temperature, followed by the addition of the primary antibody α-tubulin DM1A (Sigma-

Aldrich, Cat# T6199, RRID: AB_477583) at a concentration of 1:500 v/v in 1% *w/v* BSA in

PBS for 180 min at room temperature. After a third washing step of the primary antibody with

1X PBS, the secondary antibody Alexa Fluor 647 Goat anti-mouse IgG1 antibody was added

(Thermo Fisher Scientific, Cat#A-21240, RRID: AB_2535809) to the PBS-resuspended fixed
zoospore solution for 60 min at room temperature. Alexa Fluor 488 Phalloidin (Invitrogen,
Cat# A12379, RRID: AB_2759222) and Nile Red (Thermo Fisher Scientific, Cat# N1142)
were also added at a 1:500 v/v concentration for 60 min at room temperature. After two
washing steps with 1X PBS, the final cell pellets were resuspended in 100 μl of 20% v/v
Citifluor AF2 Antifadent Mountant Solution and 7 μl was placed on a slide and covered with a
coverslip, which was then sealed with transparent nail polish on the edges to avoid
evaporation. Cells were imaged on a Zeiss LSM-780 inverted high-resolution laser scanning
confocal microscope with a Ph3 ×100 oil objective. Exposures were kept constant during
experiments, and images collected using ZEISS ZEN Software (ZEN Digital Imaging for
Light Microscopy), and analysed/formatted with Fiji [64].
575

References

1. James, T.Y., Porter, T.M., and Martin, W.W. (2014). Blastocladiomycota. Syst. Evol.,
177–207.
2. Powell, M.J. (2017). Chytridiomycota. In Handbook of the Protists: Second Edition, pp.
1523–1558.
3. Sparrow, F.K. (1960). Aquatic Phycomycetes. Univ. Michigan Press *132*.
4. Naranjo-Ortiz, M.A., and Gabaldón, T. (2019). Fungal evolution: major ecological
adaptations and evolutionary transitions. Biol. Rev. *94*, 1443–1476.
5. Kagami, M., de Bruin, A., Ibelings, B.W., and Van Donk, E. (2007). Parasitic chytrids:
their effects on phytoplankton communities and food-web dynamics. Hydrobiologia
578, 113–129.
6. Gleason, F.H., Kagami, M., Lefevre, E., and Sime-Ngando, T. (2008). The ecology of
chytrids in aquatic ecosystems: roles in food web dynamics. Fungal Biol. Rev. *22*, 17–
25.
7. Kagami, M., Miki, T., and Takimoto, G. (2014). Mycoloop: Chytrids in aquatic food
webs. Front. Microbiol. *5*, 166.
8. Cheng, T.L., Rovito, S.M., Wake, D.B., and Vredenburg, V.T. (2011). Coincident mass
extirpation of neotropical amphibians with the emergence of the infectious fungal
pathogen *Batrachochytrium dendrobatidis*. Proc. Natl. Acad. Sci. *108*, 9502–9507.
9. Gleason, F.H., and Lilje, O. (2009). Structure and function of fungal zoospores:
ecological implications. Fungal Ecol. *2*, 53–59.
10. Bentley, S.A., Laeverenz-Schlogelhofer, H., Anagnostidis, V., Cammann, J., Mazza,
M.G., Gielen, F., and Wan, K.Y. (2022). Phenotyping single-cell motility in microfluidic
confinement. Elife *11*, e76519.
11. Bondoc-Naumovitz, K.G., Laeverenz-Schlogelhofer, H., Poon, R.N., Boggon, A.K.,

Bentley, S.A., Cortese, D., and Wan, K.Y. (2023). Methods and Measures for
Investigating Microscale Motility. Integr. Comp. Biol. *63*, 1485–1508.

12. Fuller, M.S., and Olson, L.W. (1971). The zoospore of Allomyces. Microbiology *66*,
171–183.

13. Reichle, R.E., and Fuller, M.S. (1967). The fine structure of *Blastocladiella emersonii*
zoospores. Am. J. Bot. *54*, 81–92.

14. Letcher, P.M., Powell, M.J., Churchill, P.F., and Chambers, J.G. (2006).
Ultrastructural and molecular phylogenetic delineation of a new order, the
Rhizophydiales (Chytridiomycota). Mycol. Res. *110*, 898–915.

15. Simmons, D.R., James, T.Y., Meyer, A.F., and Longcore, J.E. (2009).
Lobulomycetales, a new order in the Chytridiomycota. Mycol. Res. *113*, 450–460.

16. Simmons, D.R. (2011). Phylogeny of Powellomycetaceae fam. nov. and description of
Geranomyces variabilis gen. et comb. nov. Mycologia *103*, 1411–1420.

17. Longcore, J.E., Letcher, P.M., and James, T.Y. (2012). *Homolaphlyctis polyrhiza* gen.
et sp. nov., a species in the Rhizophydiales (Chytridiomycetes) with multiple rhizoidal
axes. Mycotaxon *118*, 433–440.

18. Letcher, P.M., Powell, M.J., Barr, D.J.S., Churchill, P.F., Wakefield, W.S., and Picard,
K.T. (2008). Rhizophlyctidales--a new order in Chytridiomycota. Mycol. Res. *112*,
1031–1048.

19. Barr, D.J.S. (1984). Cytological variation in zoospores of Spizellomyces
(Chytridiomycetes). Can. J. Bot. *62*, 1202–1208.

20. Prostak, S.M., Robinson, K.A., Titus, M.A., and Fritz-Laylin, L.K. (2021). The actin
networks of chytrid fungi reveal evolutionary loss of cytoskeletal complexity in the
fungal kingdom. Curr. Biol. *31*, 1192-1205.e6.

21. Venard, C.M., Vasudevan, K.K., and Stearns, T. (2020). Cilium axoneme
internalization and degradation in chytrid fungi. Cytoskeleton *77*, 365–378.

22. Laundon, D., Chrismas, N., Bird, K., Thomas, S., Mock, T., and Cunliffe, M. (2022). A
cellular and molecular atlas reveals the basis of chytrid development. Elife *11*,
e73933.

23. Avelar, G.M., Schumacher, R.I., Zaini, P.A., Leonard, G., Richards, T.A., and Gomes,
S.L. (2014). A Rhodopsin-Guanylyl cyclase gene fusion functions in visual perception
in a fungus. Curr. Biol. *24*, 1234–1240.

24. Richards, T.A., and Gomes, S.L. (2015). How to build a microbial eye. Nature *523*,
166–167.

25. Robinson, K.A., Prostak, S.M., Campbell Grant, E.H., and Fritz-Laylin, L.K. (2022).
Amphibian mucus triggers a developmental transition in the frog-killing chytrid fungus.
Curr. Biol. *32*, 2765-2771.e4.

26.   Miño, G.L., Koehl, M.A.R., King, N., and Stocker, R. (2017). Finding patches in a
heterogeneous aquatic environment: pH-taxis by the dispersal stage of
choanoflagellates. Limnol. Oceanogr. Lett. *2*, 37–46.

27.   Lisicki, M., Velho Rodrigues, M.F., Goldstein, R.E., and Lauga, E. (2019). Swimming
eukaryotic microorganisms exhibit a universal speed distribution. Elife *8*, e44907.

28.   Kinoshita-Terauchi, N., Bellgrove, A., Shiba, K., and Inaba, K. (2021). Flagellar
motility during sperm chemotaxis and phototaxis in fucalean algae. Eur. J. Phycol. *56*,
85–93.

29.   Tran, Q.D., Galiana, E., Thomen, P., Cohen, C., Orange, F., Peruani, F., and Noblin,
X. (2022). Coordination of two opposite flagella allows high-speed swimming and
active turning of individual zoospores. Elife *11*, e71227.

30.   Bassani, I., Larousse, M., Tran, Q.D., Attard, A., and Galiana, E. (2020). Phytophthora
zoospores: From perception of environmental signals to inoculum formation on the
host-root surface. Comput. Struct. Biotechnol. J. *18*, 3766–3773.

31.   Trinci, A.P.J., Davies, D.R., Gull, K., Lawrence, M.I., Bonde Nielsen, B., Rickers, A.,
and Theodorou, M.K. (1994). Anaerobic fungi in herbivorous animals. Mycol. Res. *98*,
129–152.

32.   Ustinova, I., Krienitz, L., and Huss, V.A.R. (2000). *Hyaloraphidium curvatum* is not a
green alga but a lower fungus; *Amoebidium parasiticum* is not a fungus, but a
member of the DRIPs. Protist *151*, 253–262.

33.   Barr, D.J.S. (2001). Chytridiomycota. In The Mycota, D. J. McLaughlin, E. G.
McLaughlin, and P. A. Lemke, eds. (Springer Berlin Heidelberg), pp. 93–112.

34.   Galindo, L.J., López-García, P., Torruella, G., Karpov, S., and Moreira, D. (2021).
Phylogenomics of a new fungal phylum reveals multiple waves of reductive evolution
across Holomycota. Nat. Commun. *12*, 4973.

35.   Strittmatter, M., Guerra, T., Silva, J., and Gachon, C.M.M. (2016). A new flagellated
dispersion stage in *Paraphysoderma sedebokerense*, a pathogen of *Haematococcus*
pluvialis. J. Appl. Phycol. *28*, 1553–1558.

36.   Moss, A.S., Reddy, N.S., Dortaj, I.M., and San Francisco, M.J. (2008). Chemotaxis of
the amphibian pathogen *Batrachochytrium dendrobatidis* and its response to a
variety of attractants. Mycologia *100*, 1–5.

37.   Swafford, A.J.M., and Oakley, T.H. (2018). Multimodal sensorimotor system in
unicellular zoospores of a fungus. J. Exp. Biol. *221*.

38.   Wang, Y., Verbrugghe, E., Meuris, L., Chiers, K., Kelly, M., Strubbe, D., Callewaert,
N., Pasmans, F., and Martel, A. (2021). Epidermal galactose spurs chytrid virulence
and predicts amphibian colonization. Nat. Commun. *12*, 5788.

39.   Johnson, P.T.J., Dobson, A., Lafferty, K.D., Marcogliese, D.J., Memmott, J., Orlofske,

S.A., Poulin, R., and Thieltges, D.W. (2010). When parasites become prey: ecological

and epidemiological significance of eating parasites. Trends Ecol. Evol. *25*, 362–371.

40. Money, N.P. (2016). Chapter 3 - Spore Production, Discharge, and Dispersal. In The

Fungi (Third Edition), S. C. Watkinson, L. Boddy, and N. P. Money, eds. (Academic

Press), pp. 67–97.

41. Canter, H.M., and Jaworski, G.H.M. (1980). Some General Observations on

Zoospores of the Chytrid *Rhizophydium planktonicum* Canter Emend. New Phytol. *84*,

515–531.

42. Sparrow, F.K., and Johns, R.M. (1970). Observations on chytridiaceous parasites of

phanerogams. Arch. Mikrobiol. *70*, 72–81.

43. Barr, D.J.S. (1980). An outline for the reclassification of the Chytridiales, and for a

new order, the Spizellomycetales. Can. J. Bot. *58*, 2380–2394.

44. Simmons, D.R., Bonds, A.E., Castillo, B.T., Clemons, R.A., Glasco, A.D., Myers, J.M.,

Thapa, N., Letcher, P.M., Powell, M.J., Longcore, J.E., et al. (2020). The Collection of

Zoosporic Eufungi at the University of Michigan (CZEUM): introducing a new

repository of barcoded Chytridiomyceta and Blastocladiomycota cultures. IMA Fungus

11, 20.

45. Powell, M.J. (1978). Phylogenetic implications of the microbody-lipid globule complex

in zoosporic fungi. BioSystems *10*, 167–180.

46. Dawson, S.C., Sagolla, M.S., Mancuso, J.J., Woessner, D.J., House, S.A., Fritz-

Laylin, L., and Cande, W.Z. (2007). Kinesin-13 regulates flagellar, interphase, and

mitotic microtubule dynamics in *Giardia intestinalis*. Eukaryot. Cell *6*, 2354–2364.

47. Toba, S., Fox, L.A., Sakakibara, H., Porter, M.E., Oiwa, K., and Sale, W.S. (2011).

Distinct roles of 1alpha and 1beta heavy chains of the inner arm dynein I1 of

Chlamydomonas flagella. Mol. Biol. Cell *22*, 342–353.

48. Kapoor, P., Sachdeva, M., and Madhubala, R. (1999). Effect of the microtubule

stabilising agent taxol on leishmanial protozoan parasites in vitro. FEMS Microbiol.

Lett. *176*, 429–435.

49. Herth, W. (1983). Taxol effects cytoskeletal microtubules, flagella and spindle

structure of the chrysoflagellate alga Poterioochromonas. Protoplasma *115*, 228–239.

50. Wang, L., Piao, T., Cao, M., Qin, T., Huang, L., Deng, H., Mao, T., and Pan, J. (2013).

Flagellar regeneration requires cytoplasmic microtubule depolymerization and

kinesin-13. J. Cell Sci. *126*, 1531–1540.

51. Cantino, E.C., and Truesdell, L.C. (1970). Organization and fine structure of the side

body and its lipid sac in the zoospore of *Blastocladiella emersonii*. Mycologia *62*, 548–

567.

52. Galindo, L.J., Milner, D.S., Gomes, S.L., and Richards, T.A. (2022). A light-sensing

system in the common ancestor of the fungi. Curr. Biol. *32*, 3146-3153.e3.

53. KAZAMA, F.Y. (1972). Ultrastructure and Phototaxis of the Zoospores of

Phlyctochytrium sp., an Estuarine Chytrid. Microbiology *71*, 555–566.

54. Robertson, J.A. (1972). Phototaxis in a new Allomyces. Arch. Mikrobiol.

55. Saranak, J., and Foster, K.W. (1997). Rhodopsin guides fungal phototaxis. Nature

387, 465–466.

56. Avelar, G.M., Glaser, T., Leonard, G., Richards, T.A., Ulrich, H., and Gomes, S.L.

(2015). A cyclic GMP-dependent K+ channel in the blastocladiomycete fungus

Blastocladiella emersonii. Eukaryot. Cell *14*, 958–963.

57. Gao, S., Nagpal, J., Schneider, M.W., Kozjak-Pavlovic, V., Nagel, G., and Gottschalk,

A. (2015). Optogenetic manipulation of cGMP in cells and animals by the tightly light-

regulated guanylyl-cyclase opsin CyclOp. Nat. Commun. *6*, 1–12.

58. Broser, M., Spreen, A., Konold, P.E., Peter, E., Adam, S., Borin, V., Schapiro, I.,

Seifert, R., Kennis, J.T.M., Bernal Sierra, Y.A., et al. (2020). NeoR, a near-infrared

absorbing rhodopsin. Nat. Commun. *11*, 5682.

59. Broser, M., Busse, W., Spreen, A., Reh, M., Sierra, Y.A.B., Hwang, S., Utesch, T.,

Sun, H., and Hegemann, P. (2023). Diversity of rhodopsin cyclases in zoospore-

forming fungi. Proc. Natl. Acad. Sci. *120*, e2310600120.

60. Amses, K.R., Simmons, D.R., Longcore, J.E., Mondo, S.J., Seto, K., Jerônimo, G.H.,

Bonds, A.E., Quandt, C.A., Davis, W.J., Chang, Y., et al. (2022). Diploid-dominant life

cycles characterize the early evolution of Fungi. Proc. Natl. Acad. Sci. U. S. A. *119*,

e2116841119.

61. Powell, M.J. (2014). Hypothesized evolutionary trends in zoospore ultrastructural

characters in Chytridiales (Chytridiomycota). *106*, 379–396.

62. Reid, B., Morris, B.M., and Gow, N.A.R. (1995). Calcium-Dependent, Genus-Specific,

Autoaggregation of Zoospores of Phytopathogenic Fungi. Exp. Mycol. *19*, 202–213.

63. Byrt, P.N., Irving, H.R., and Grant, B.R. (1982). The Effect of Cations on Zoospores of

the Fungus *Phytophthora cinnamomi*. Microbiology *128*, 1189–1198.

64. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,

Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source

platform for biological-image analysis. Nat. Methods *9*, 676–682.

65. Tinevez, J.-Y., Perry, N., Schindelin, J., Hoopes, G.M., Reynolds, G.D., Laplantine, E.,

Bednarek, S.Y., Shorte, S.L., and Eliceiri, K.W. (2017). TrackMate: An open and

extensible platform for single-particle tracking. Methods *115*, 80–90.

66. Anderson, D.P., Whitney, D.S., Hanson-Smith, V., Woznica, A., Campodonico-

Burnett, W., Volkman, B.F., King, N., Thornton, J.W., and Prehoda, K.E. (2016).

Evolution of an ancient protein function involved in organized multicellularity in

animals. Elife *5*, e10147.

751