Main Manuscript for

Antithetic effects of agonists and antagonists on the structural fluctuations of TRPV1 channel

Ayumi Sumino,[1, 2*†] Yimeng Zhao,[3,4†] Daichi Mukai,[5] Takashi Sumikama,[1, 6] Leonardo Puppulin,[1] Motoyuki Hattori,[3*] Mikihiro Shibata.[1, 2]

Nano Life Science Institute (WPI-NanoLSI), Kanazawa University, Kanazawa, 920-1192, Japan

Institute for Frontier Science Initiative, Kanazawa University, Kanazawa, 920-1192, Japan

State Key Laboratory of Genetic Engineering, Collaborative Innovation Center of Genetics and Development, Shanghai Key Laboratory of Bioactive Small Molecules, Department of Physiology and Neurobiology, School of Life Sciences, Fudan University, Shanghai 200438, China

Human Phenome Institute, Fudan University, Shanghai 200438, China

Division of Nano Life Science, Graduate School of Frontier Science Initiative, Kanazawa, 920-1192, Japan

PRESTO/JST

Present address (LP): Department of Molecular Sciences and Nanosystems, Università Ca' Foscari Venezia, Via Torino 155, 30172 Mestre-Venezia, Italy

† Contributed equally.

* Ayumi Sumino, Motoyuki Hattori.

Email: sumino@staff.kanazawa-u.ac.jp, hattorim@fudan.edu.cn

Author Contributions: A.S. and M.H. designed research. A.S. and D.M., performed HS-AFM imaging. A.S., D.M. and T.S. analyzed data. Y.Z. and M.H. prepared purified TRPV1 sample. A.S., T.S., L.P, M.H. and M.S. wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Classification: Brief reports

Keywords: TRPV1 channel, High-speed atomic force microscopy, Fluctuation, Resiniferatoxin, Capsazepine

This file includes:

Main Text
Figures 1 to 2

Abstract

Transient receptor potential vanilloid member 1 (TRPV1) is a heat and capsaisin receptor that
allows cations to permeate and cause pain. As the molecular basis for temperature sensing, the
heat capacity (ΔC_p) model (D. E. Clapham, C. Miller, *Proc. Natl. Acad. Sci. U. S. A.* **108**, 19492–
19497 (2011).) has been proposed and experimentally supported. Theoretically, heat capacity is
proportional to a variance in enthalpy, presumably related to structural fluctuation; however, the
fluctuation of TRPV1 has not been directly visualized. In this study, we directly visualized single-
molecule structural fluctuations of the TRPV1 channels in a lipid bilayer with the ligands
resiniferatoxin (RTX: agonist, 1000 times hotter than capsaicin) and capsazepine (CPZ: antagonist)
by high-speed atomic force microscopy (HS-AFM). We observed the structural fluctuations of
TRPV1 in an apo state, and found that RTX binding enhances structural fluctuations, while CPZ
suppresses fluctuations. These ligand-dependent differences in structural fluctuation would play a
key role in the gating of TRPV1.

14
15
16 **Main Text**
17
18 **Introduction**
19
The TRPV1 cation channel in sensory neuron opens by heat (> 43 degree Celsius) or
binding of capsaicin, generating pain sensation.(1, 2) As the molecular basis of temperature
sensing, the heat capacity (ΔC_p) model has been proposed(3) and experimentally supported.(4)
Recently, cryo-electron microscopy (cryo-EM) has revealed the molecular structure of TRPV1 not
only with and without ligand, but also in a heat-dependent manner.(5–7) Since C_p is proportional to
enthalpy fluctuation that is directly related to conformational fluctuations, the measurement of the
latter, which has not yet been performed in TRPV1, is also instrumental to fully elucidate the
functioning mechanism of this channel. Furthermore, there are several examples showing that
changes in structural fluctuations are closely related to protein activity, including ion channels.(8–
10) Thus, it makes sense to analyze structural fluctuations in each gating state. Built upon this lack
of knowledge, we use here HS-AFM(11) to directly observe structural fluctuations in TRPV1 under
different experimental conditions. Ideally, measurements with temperature-control are the best way
to control the activity of this channel; however, there are technical difficulties to obtain high-
resolution images by heating the sample, such as thermal drift. As such, in this study, we have
directly observed by HS-AFM the structural fluctuations of the TRPV1 channel in a lipid bilayer with
and without two ligands, the RTX(12) and the CPZ(13).(11)
36
37 **Results**
38
39 **HS-AFM of TRPV1 channels in lipid bilayer**
40
Since the TRPV1 channel is homo-tetrameric channel, many clover-like structures were
found in the TRPV1-reconstituted membrane (Figure 1, Movie S1-S3). Considering the height of
the protrusion from the lipid membrane near the channel (1.4 ± 0.5 nm, n = 24), the observed
surface is likely to be the extracellular side of TRPV1 (Figure 1A). The overall structure of the
channel was not significantly changed by the presence of ligands (Figure 1 B to D), which is
consistent with previous cryo-EM studies that have reported relatively small structural change upon

ligand binding.(6) From these HS-AFM movies, we extracted the trajectories of each of the four
domains to analyze fluctuations of the TRPV1, as shown on the bottom right of figures 1 B-C.
49
Agonist/antagonist changed structural fluctuation of the TRPV1
51
52 Using the single-molecule trajectories from a large number of channels, we analyzed
53 structural fluctuations. Distributions of lateral and angular displacement between successive
54 frames in the HS-AFM videos are affected by ligand binding (Figure 2A and B). In both distributions,
55 the RTX binding increases the fluctuations as compared to the Apo state, while, conversely, the
56 CPZ binding suppresses it.
57
Discussion
59
Possible contribution of structural fluctuation for gating of TRPV1
61
In this study, we elucidate that RTX enhances structural fluctuation of TRPV1, while CPZ
suppresses it. This is the first experimental evidence showing the correlation between molecular
fluctuation and gating state (ligand binging) of the TRPV1 channel. One may thought that the
observed fluctuations are too small (angstrom level) to modulate channel function; however, as
well described in literature(10), energetics of ion permeation can be altered significantly by small
structural fluctuations of the narrow pore structure. Given that even such a small change in the
narrow pore alters energetics, small fluctuations at wider locations could alter the energetics more
largely.
Previously, Miller and Clapham have proposed that change in heat capacity (ΔC_p) is
theoretically a key factor for temperature-gating of TRP channels (3), and Chowdhury et al.
supported this concept by mutational experiments.(4) Intriguingly, Kwon et al. have reported that
the predicted ΔC_p from SASA of cryo-EM structures in closed and open states is smaller than that
predicted by Miller and Clapham. Since RTX act as an agonist at TRPV1, the increase in molecular
fluctuation of TRPV1 in the presence of RTX implies an increase in heat capacity during TPRV1
channel activation. It suggests an important role of structural fluctuations in the TRPV1 gating
mechanism, which follows the heat capacity model. This study suggests the significance of
measuring structural fluctuation, which would be another key factor for heat-sensing of TRPV1,
because structural fluctuations are related to a variance in enthalpy, which is directly proportional
to heat capacity.
81
82
Materials and Methods
84
Expression and purification of TRPV1
86 See extended method.
87
HS-AFM observation of TRPV1 reconstituted in a lipid bilayer
We reconstituted the TRPV1 channel into a lipid bilayer by a previously reported
method.(14) To prevent lateral diffusion of the channels, we modified mica surface by 3-
aminopropyltriethoxysilane (APTES) by applying 0.01 % APTES solution onto the mica for 3 min,
then rinsed by pure water. Solubilized TRPV1 channels (10 μg/mL in 50 mM HEPES (pH8), 200
mM NaCl, 0.025 % DDM, 10 µg/ml soybean lipids, 2 mM TCEP, 10 % glycerol) were applied onto
the APTES-modified mica surface for 5 min, then floating channels were washed out by buffer of
solubilized channel. Next, DDM-destabilized DMPC liposomes (100 μg/mL DMPC in 10 mM
HEPES [pH 7.5], 150 mM NaCl, 0.006 % DDM) were applied and incubated for 10 min, then rinsed
by imaging buffer (10 mM HEPES [pH 7.5], 150 mM NaCl) to remove DDM and excess liposomes.
For imaging with ligands, 10 nM RTX (K_i = 43 pM)(12) or 20 μM CPZ (K_i = 0.52 μM) (13) were
added to all buffer from dilution of solubilized TRPV1 to imaging buffer. All reconstitution
procedures were performed at 25°C. For HS-AFM observations, a laboratory-built HS-AFM was

used(11). We used cantilever AC10 (Olympus Co., Tokyo, Japan) with an electron beam deposition
(EBD) tip.(11) The typical resonant frequency of AC10 in water was 400 kHz.
103
Analysis of fluctuation
To analyze structural fluctuation, we tracked position of four subunits by image analysis.
Prior to particle tracking, we removed background, high-frequency noise and lateral drift of HS-
AFM movies using subtract background, FFT filter and template matching plugin on ImageJ,
respectively. Trajectories of particles were analyzed using TrackMate plugin on ImageJ.(15)
Estimated blob diameter was set to 4.0 nm. Obvious particle misrecognitions and trajectory swaps
were corrected manually. We analyzed 91 channels and 95,347 frames in total.
111
112   **Data Availability.**
113

114   The datasets generated and/or analyzed during the current study are available from the
115   corresponding author (sumino@staff.kanazawa-u.ac.jp) on reasonable request.
116
117   **Acknowledgments**
118
A.S. acknowledge to Prof. Toshio Ando, Prof. Noriyuki Kodera and Dr. Ken-ichi Umeda (Kanazawa
University) for sharing scanning electron microscope for fabrication of EBD tip and operating
software for HS-AFM. A.S. is grateful to Prof. Masahiro Sokabe, Prof Hitoshi Tatsumi, Prof. Hiroaki
Hirata (Kanazawa Institute of Technology) and Prof. Yasushi Sako (RIKEN) for fruitful discussion.
A.S. thanks Ms. Yoko Yamamoto (Kanazawa University) for analytical assistance. A.S. thanks
Grant-in-Aids for Scientific Research (B) (22H01919) and Challenging Research (Exploratory)
(22K19290) for funding. This work was supported by World Premier International Research Center
Initiative (WPI), MEXT, Japan. This work was also supported by funding from the National Natural
Science Foundation of China (32071234, 32271244 and 32250610205) to M.H. and by a key
laboratory program of the Education Commission of Shanghai Municipality (ZDSYS14005).

129
130   **References**
131

1.   M. J. Caterina, D. Julius, The vanilloid receptor: A molecular gateway to the pain pathway.
Annu. Rev. Neurosci. **24**, 487–517 (2001).

2.   M. J. Caterina, *et al.*, The capsaicin receptor: A heat-activated ion channel in the pain
pathway. *Nature* **389**, 816–824 (1997).

3.   D. E. Clapham, C. Miller, A thermodynamic framework for understanding temperature
sensing by transient receptor potential (TRP) channels. *Proc. Natl. Acad. Sci.* **108**, 19492–
19497 (2011).

4.   S. Chowdhury, B. W. Jarecki, B. Chanda, A molecular framework for temperature-
dependent gating of ion channels. *Cell* **158**, 1148–1158 (2014).

5.   M. Liao, E. Cao, D. Julius, Y. Cheng, Structure of the TRPV1 ion channel determined by
electron cryo-microscopy. *Nature* **504**, 107–12 (2013).

6.   Y. Gao, E. Cao, D. Julius, Y. Cheng, TRPV1 structures in nanodiscs reveal mechanisms
of ligand and lipid action. *Nature* **534**, 347–351 (2016).

7.   D. H. Kwon, *et al.*, Heat-dependent opening of TRPV1 in the presence of capsaicin. *Nat.*
Struct. Mol. Biol. **28**, 554–563 (2021).

8.   M. Terazima, Enhanced conformational fluctuations during protein reactions. *Chem. Lett.*
48, 802–810 (2019).

9. M. W. Van Der Kamp, *et al.*, Dynamical origins of heat capacity changes in enzyme-
catalysed reactions. *Nat. Commun.* **9**, 1–7 (2018).

10. T. W. Allen, O. S. Andersen, B. Roux, On the Importance of Atomic Fluctuations, Protein
Flexibility, and Solvent in Ion Permeation. *J. Gen. Physiol.* **124**, 679–690 (2004).

11. T. Uchihashi, N. Kodera, T. Ando, Guide to video recording of structure dynamics and
dynamic processes of proteins by high-speed atomic force microscopy. *Nat. Protoc.* **7**,
1193–1206 (2012).

12. J. Szolcsanyi, A. Szallasi, Z. Szallasi, F. Joo, P. M. Blumberg, Resiniferatoxin: An
ultrapotent selective modulator of capsaicin-sensitive primary afferent neurons. *J.*
Pharmacol. Exp. Ther. **255**, 923–928 (1990).

13. J. Lee, *et al.*, Analysis of structure–activity relationships for the 'B-region' of N-(4-t-
butylbenzyl)-N′-[4-(methylsulfonylamino)benzyl]-thiourea analogues as TRPV1
antagonists. *Bioorg. Med. Chem. Lett.* **15**, 4143–4150 (2005).

14. A. Sumino, T. Uchihashi, S. Oiki, Oriented Reconstitution of the Full-Length KcsA
Potassium Channel in a Lipid Bilayer for AFM Imaging. *J. Phys. Chem. Lett.* **8**, 785–793
(2017).

15. J. Y. Tinevez, *et al.*, TrackMate: An open and extensible platform for single-particle
tracking. *Methods* **115**, 80–90 (2017).

Figures

Figure 1. HS-AFM of TRPV1 channels in a lipid bilayer with or without ligands. (A) Schematic illustration of the experiment. (B-D) HS-AFM images and trajectories of TRPV1 without ligand (B), with 10 nM RTX (C) and 20 µM CPZ (D) added to the imaging buffer. Frame rate: 0.3 sec/frame.

176
177 **Figure 2.** Structural fluctuation of TRPV1 depending on binding of ligands. Distribution of lateral
178 (A) and angular (B) displacement of each subunit between successive frames of HS-AFM video.
179 Frame interval (Δt) is 0.3 sec for A and B. Analyzed number of channels for Apo, RTX and CPZ
180 are, (A): 43, 22 and 26, (B): 33, 13 and 14, respectively. Total frames for analysis of Apo, RTX and
181 CPZ are (A): 46702, 23574 and 25071, (B): 37526, 13802 and 13677, respectively.
182