1
2 **The *Medicago truncatula* nodule-specific cysteine-rich peptides, NCR343 and NCR-**
3 **new35 are required for the maintenance of rhizobia in nitrogen-fixing nodules.**
4
5

6 Beatrix Horváth[2*], Berivan Güngör[1,2*], Mónika Tóth[2], Ágota Domonkos[2], Ferhan Ayaydin[3,4],
7 Farheen Saifi[2], Yuhui Chen[5], János Barnabás Biró[1], Mickael Bourge[6], Zoltán Szabó[2], Zoltán
8 Tóth[2], Rujin Chen[5], Péter Kaló[1,2]
9
10
11
1 Institute of Plant Biology, Biological Research Centre, Eötvös Lóránd Research Network,
Szeged, Hungary
2 Institute of Genetics and Biotechnology, Hungarian University of Agriculture and Life
Sciences, Gödöllő, Hungary
3 Hungarian Centre of Excellence for Molecular Medicine (HCEMM) Nonprofit Ltd., Szeged,
Hungary
4 Cellular Imaging Laboratory, Biological Research Centre, Eötvös Lóránd Research
Network, Szeged, Hungary
5 School of Life Sciences, Lanzhou University, Lanzhou, China 730000
6 Cytometry Facility, Imagerie-Gif, Université Paris-Saclay, CEA, CNRS, Institute for
Integrative Biology of the Cell (I2BC), 91198, Gif-sur-Yvette, France
29 * B.H. and B.G. contributed equally to this work.
30
31 Author for correspondence: Péter Kaló, Email: kalo.peter@brc.hu
32
33 ORCID IDs:
34
Peter Kalo: 0000-0002-0404-8904
Beatrix Horvath: 0000-0001-8499-568X
Berivan Güngör: 0000-0002-5612-1130
Mónika Tóth
Agota Domonkos: 0000-0003-4017-0605
Ferhan Ayaydin 0000-0002-9126-0865
Farheen Saifi: 0000-0002-6912-8227
Yuhui Chen
János Barnabás Biró: 0000-0001-8851-0387
Mickael Bourge: 0000-0002-4111-6208
Zoltán Szabó 0000-0002-2001-0375
Zoltán Tóth 0000-0002-3212-2438
Rujin Chen 0000-0001-5444-2144

48
49
50

Summary

The *Medicago truncatula - Sinorhizobium* sp. symbiosis is one of the most effective nitrogen-

fixing interaction. In the Inverted Repeat-Lacking Clade (IRLC) legumes, including *M.*

truncatula, rhizobia undergo terminal differentiation resulting in elongated and

endoreduplicated bacteroids specialised for nitrogen fixation. This irreversible transition of

rhizobia is controlled by the host plant and mediated by nodule-specific cysteine-rich (NCR)

peptides, of which about 700 are encoded by a large gene family in the *M. truncatula* genome.

Recent studies have demonstrated that some NCR peptides, NCR169, NCR211 and NCR247,

are unique and essential for nitrogen-fixing symbiosis. By analysing the symbiotic phenotype

of ineffective *M. truncatula* mutants, *Mtsym19*, *Mtsym20* and NF-FN9363, we found that the

nodulation phenotype, the bacteroid and symbiotic host cell differentiation were defective

likewise in the formerly studied *ncr* mutants, *Mtdnf4-1* (*ncr211*) and *Mtdnf7-2* (*ncr169*). We

demonstrated that the incomplete differentiation of bacteroids triggers premature senescence

of rhizobia in the nitrogen fixation zones of mutant nodules. We proved that *Mtsym19* and

Mtsym20 mutants are allelic and the *MtSYM19/20* gene encodes the peptide NCR-new35. We

identified a deletion of four *NCR* genes in the symbiotic locus of NF-FN9363 and

demonstrated that the lack of *NCR343* is responsible for the ineffective symbiosis of NF-

FN9363. The activity of *NCR-new35* is significantly lower and limited to the transition zone

of the nodule compared with other crucial *NCR* genes, *NCR169*, *NCR211* and *NCR343* which

are highly expressed in the interzone and nitrogen fixation zone. The fluorescent protein-

tagged version of NCR343 and NCR-new35 localize to the symbiotic compartment. We also

demonstrated that the conserved first cysteine residues are required for the symbiotic function

of NCR343 and NCR-new35. Our discovery added two additional members to the group of

NCR genes essential for nitrogen–fixing symbiosis in *M. truncatula*.

Introduction

Legumes can establish nitrogen-fixing endosymbiotic association with soil bacteria,

collectively termed rhizobia. This interaction results in the formation of nodules, generally

formed on the roots, where rhizobia are hosted and convert atmospheric dinitrogen to

ammonia. The interaction is initiated by the mutual recognition of the host legume and

rhizobia mediated by signalling molecules of both symbiotic partners (Oldroyd & Downie,

2008). The perception of the bacterial signalling molecules, the nodulation factors (NFs), and

the attachment of rhizobia on root hair cells induces two synchronous developmental

processes, rhizobial infection and initiation of nodule primordia by activating the mitosis of

certain root cortical cells. Trapped bacteria in root hair curls induce the invagination of the

plasma membrane forming tubular-like structures with rhizobia inside, termed infection

threads (IT). The ITs progress transcellularly towards the nodule primordia and when they

reach the dividing root cortical cells, rhizobia are released and colonize many of the nodule

cells. Within infected nodule cells, bacteria are encompassed by a plant derived membrane

resulting in formation of subcellular compartments called symbiosomes. Bacteria multiply

and undergo morphological and metabolic transition to adapt from free-living life-style to

their advanced endosymbiotic form termed bacteroids (Jones *et al*., 2007).

Nodules formed on legume roots are classified into two types, the so-called indeterminate or

determinate nodules, based on the term of their meristem activity (Hirsch, 1992). The

cylindrical shaped indeterminate nodules, such as those found on *Medicago truncatula* roots,

possess a persistent meristem during the life-span of the nodules, and as a result, the mature

indeterminate nodules consist of a developmental gradient of cells forming distinct zones

(Vasse *et al*., 1990). The mitotic activity of meristem cells (zone I, ZI), located at the nodule

apex, produces new cells for all nodule tissues. Cells leaving the meristem are relatively small

with a large central vacuole (Gavrin *et al*., 2014) and have a four-fold haploid DNA content

(4C) (Nagymihaly *et al*., 2017). In these cells in the distal part of the infection zone (ZIId),

rod-shaped rhizobia are released from ITs to the host cell cytoplasm and from hereafter, both

rhizobia and infected nodule cells undergo differentiation processes through the subsequent

nodule zones. The transition of symbiotic cells and rhizobia involves the replication of their

genomes without cytokinesis resulting in enlargement of invaded host cells and elongation of

bacteria (Vasse *et al*., 1990; Mergaert *et al*., 2006; Nagymihaly *et al*., 2017). The enlarged

symbiotic cells in the proximal part of the infection zone (ZIIp) have a DNA content of

8C/16C and their volume is almost occupied by distended vacuoles and symbiosomes

containing non-dividing bacteroids. In cells of the first few layers of the nitrogen fixation

zone (ZIIId, also termed as transition zone or interzone between ZII and ZIII and referred

hereafter IZ in this study), elongated bacteroids surround the collapsed central vacuoles of

infected host cells (16C/32C) which are filled with starch granules. Rhizobia and host cells

complete their differentiation in IZ and nitrogen fixation begins in the large symbiotic cells

having highly endoreduplicated chromosomes (32C/64C) and large volume vacuoles in the

nitrogen fixation zone (ZIII). As the indeterminate nodule ages, proximal to the root, a

senescence zone (ZIV) evolves wherein the symbiotic interaction terminates and both infected

host cells and rhizobia degrade.

Rhizobia, hosted in the nodules of legumes of the Inverted Repeat-Lacking Clade (IRLC)

(consisting of legume genera such as *Medicago*, *Pisum*, *Trifolium*, *Vicia*, etc.) and

Aeschynomene species undergo terminal differentiation (Mergaert *et al*., 2006; Czernic *et al*.,

2015). These differentiated bacteroids, reaching 5-10 μm in length and DNA content of 24C

in fully differentiated form in *M. truncatula* nodules, are not able to regrow and proliferate

outside the indeterminate nodule. The irreversible transition of rhizobia in *M. truncatula*

nodules is mediated by host-produced defensin-related nodule-specific cysteine-rich (NCR)

peptides (Van De Velde *et al*., 2010). The genome of *M. truncatula* codes for about 700 NCR

peptides and similar genes in different numbers have been identified in other IRLC legume

species (Mergaert *et al*., 2003; Montiel *et al*., 2017; Zorin *et al*., 2022; Wei *et al*., 2022).

Almost all *NCR* genes are specifically expressed in infected cells of *M. truncatula* root

nodules and they are activated in successive waves during nodule organogenesis and

bacteroid differentiation (Nallu *et al*., 2013; Guefrachi *et al*., 2014). NCR peptides possess

highly conserved N-terminal signals which are recognised by the secretion complex of the

nodule cells (Wang *et al*., 2010). The secretory pathway delivers the mature NCR peptides,

usually 35 to 55 residues in length, to the symbiosomes inducing bacteroid differentiation.

The characteristic feature of NCR peptides is the four or six cysteine residues in conserved

positions which are presumed to be involved in the formation of intra- and intermolecular

disulphide bonds (Mergaert *et al*., 2003). The structural analysis of peptides NCR044 and

NCR169 confirmed the presence of different pattern of disulphide bonds that contribute to the

structural stability of the peptides (Velivelli *et al*., 2020; Isozumi *et al*., 2021). *M. truncatula*

Thioredoxin s1 (Trx s1), which is able to reduce disulphide bonds, is also targeted to

symbiosomes and its role has been suggested regulating the activity of NCR peptides by their

redox state (Ribeiro *et al*., 2017). These results indicate the importance of cysteine residues

and the substitution of the conserved cysteine for serine evidenced the requirement of
cysteines and the intracellular disulphide bridges for the symbiotic function of the NCR169
peptide (Horvath *et al*., 2015).

The high number of *NCR* genes in the *M. truncatula* genome implied that they act
redundantly. Forward genetic studies of deletion mutants deficient in single NCR peptides,
NCR169 or NCR211, revealed that these peptides are essential for bacteroid differentiation
and persistence in *M. truncatula* nodules (Horvath *et al*., 2015; Kim *et al*., 2015). Despite
these findings, little knowledge is available about the molecular mechanism of NCR peptides
mediating bacteroid differentiation *in planta*. A recent reverse genetic analysis of *NCR247*
demonstrated that the encoded peptide is required for nodule functioning and provided
experimental data about the operation of one of the NCR peptides (Sankari *et al*., 2022).
Biochemical studies demonstrated that NCR247 is able to form a complex with haem
facilitating iron uptake by rhizobia. In addition to the prominent positive regulatory role of
these NCR peptides in the symbiotic interaction, other peptides, NFS1 and NFS2, control
bacterial survival in a strain- and allele-specific manner and negatively regulate the nitrogen-
fixing symbiosis in *M. truncatula* (Yang *et al*., 2017; Wang *et al*., 2017).

The *M. truncatula* genome contains several hundreds of *NCR* genes but only few of them
have been proven to be required for nitrogen-fixing symbiosis. In this report, we present the
phenotypic analysis of three *M. truncatula* symbiotic nitrogen-fixing mutants *Mtsym19*,
Mtsym20 and NF-FN9363. The failure of four different rhizobial strains to colonise the
nitrogen fixation zone of the symbiotic nodules indicates that the ineffective symbiotic
phenotype is not strain dependent. In the ineffective nodules, the bacteroid differentiation was
incomplete suggesting the function of the impaired plant genes in this developmental step. We
found that *Mtsym19* and *Mtsym20* are both defective in the same gene, *NCR-new35* and the
deletion of *NCR343* in mutant NF-FN9363 resulted in the ineffective symbiotic phenotype.
We show that the substitution of the first cysteine residue either in NCR-new35 or in NCR343
abolishes the symbiotic function of these peptides. Our results indicate that the peptides NCR-
new35 and NCR343 are essential for terminal bacteroid differentiation in *M. truncatula*
nodules.

181
182
183 **Materials and Methods**
184
Plant materials, rhizobia strains and growth conditions

Symbiotic mutant NF-FN9363 was identified during symbiotic screens of fast neutron

bombarded *M. truncatula* A17 plants (The Samuel Roberts Noble Foundation, Ardmore,

Oklahoma, USA). Mutants *Mtsym18* (TR36), *Mtsym19* (TR183) and *Mtsym20* (TRV43) were

obtained following gamma ray irradiation of *M. truncatula* cv. Jemalong (J5) (Sagan *et al.*,

1995; Morandi *et al.*, 2005). *M. truncatula* cv. Jemalong or genotype A17 were used as a

wild-type (wt) controls for characterization of the symbiotic phenotype of *Mtsym19*, *Mtsym20*

and FN-NF9363 mutants. Symbiotic mutants were crossed to *M. truncatula* A20 for genetic

mapping. Seeds were germinated as described in the *M. truncatula* handbook (Garcia *et al.*,

2006) Seedlings were grown under symbiotic conditions in a zeolite substrate (Geoproduct

Kft., Mád, Hungary) in growth chambers and kept under a photoperiod of 16 h of light / 8 h of

darkness at 24 ^{o}C. Four-day-old pre-grown plants were inoculated with the rhizobium strains

Sinorhizobium (*Ensifer*) *medicae* WSM419, ABS7 or *S. meliloti* 1021 carrying the pXLGD4

plasmid expressing the *hemA::lacZ* reporter gene or strain *S. meliloti* FSM-MA carrying the

pMEpTrpGUSGFP plasmid expressing the β-glucuronidase-GFP fusion protein. Rhizobia

were cultured in tryptone-yeast (TY) media (pH 7.2) supplemented with 6 mM $CaCl_2$

(Beringer, 1974). The bacterial liquid culture was pelleted by centrifugation and resuspended

in Gibson liquid media (Gibson & Nutman 1960) at a final dilution 1:50 (~OD_{600} 0.1) and

used for inoculation. Plants were harvested or scored for symbiotic phenotype indicated in

each experiment.

Histological analyses and microscopy

The rhizobium-dependent nodulation phenotype of the ineffective mutant and wild-type

plants was analysed at 3 weeks post inoculation (wpi) with different bacterial strains using a

Leica MZ10 F stereo microscope at 2.5x standard magnification.

***lacZ* and SYTO13 staining**

To assess bacterial colonization, mutant and wt plants were inoculated with rhizobia strains

carrying the pXLGD4 plasmid and 14-day-old nodules post inoculation were harvested in

1xPBS buffer (pH 7.4), vacuum-fixed in 4% (w/v) paraformaldehyde solution for 3 x 30 sec

in an Eppendorf 5301 concentrator and post-fixed on ice for 30 min. Fixed nodules were

embedded in 5% (w/v) agarose and 65 µm thick longitudinal sections were prepared using a

Leica VT1200S vibrotome. In order to visualize the rhizobium-infected cells, nodule sections

were stained for β-galactosidase activity in X-Gal staining solution (Bovin *et al.*, 1990) for 30

minutes at 37 ^{o}C. Stained sections were analysed using an Olympus BX41M light microscope

(Olympus Life Science Europa GmbH, Hamburg, Germany) with 10x objective and images

were taken with an Olympus E-10 digital camera. The symbiotic nodule cell structure and

bacteroid morphology were also analysed on nodule sections stained with 5 µM SYTO13

(Thermo Fisher Scientific, diluted in 1xPBS buffer pH 7.4) for 20 min at room temperature

and imaged using a Leica TCS SP8 confocal laser scanning microscope. The microscope

configuration was as follows: objective lenses HCX PL FLUOTAR 10x dry (NA: 0.3) and

HC PL APO CS2 63x oil (NA: 1.4); zooms: 0.75 and 2.14; excitation: OPSL 488 nm laser

(SYTO13, green) and OPSL 552 nm laser (red autofluorescence); spectral emission detector:

510-540 nm (SYTO13) and 558 nm – 800 nm (red autofluorescence).

GUS reporter assay

The promoter activity of the *NCR* genes was assayed with the promoter-*GUS* reporter gene

constructs generated using the Gateway cloning technology (Invitrogen). The cloned promoter

fragments (819 bp of *pr NCR-new35*, 1620 bp of *prNCR343*, 1178 bp of *prNCR169* and 2055

bp of *prNCR211*) of the four *NCR* genes were recombined from pDONR201 entry vector

constructs into a modified Gateway destination vector pKGWFS7 containing the dsRED

fluorescent transformation marker under the control of a constitutively active *Arabidopsis*

thaliana ubiquitin10 (pUBQ10) promoter (pKGWFS7-pUBQ10::dsRED). The constructs

were introduced into the roots of *M. truncatula* A17 plants using the *Agrobacterium*

rhizogenes-mediated hairy root transformation carried out as described by Boisson-Dernier

and co-workers (2001). To detect GUS activity, 65 µm thick fixed sections of nodules,

harvested at 2 and 3 wpi with *S. medicae* WSM419, were infiltrated with X-Gluc staining

solution (1 mM 5-bromo-4-chloro-3-indolyl-β-D-glucuronic acid substrate, 50 mM sodium

phosphate buffer (pH 7.4), 2.5 mM potassium ferrocyanide, 2.5 mM potassium ferricyanide,

20% methanol and Triton X-100 at 0.1 ml/100 ml) for 30-40 min at 37 °C. GUS stained

nodule sections were analysed and imaged similar to *lacZ*-stained nodule sections.

Scanning electron microscopy

Nodule samples were treated and sections were prepared for scanning electron microscopy as

described earlier (Domonkos *et al*., 2017).

Genetic mapping and allelism test

The F2 segregation populations were generated by self-pollination of the F1 progeny of the

crosses of the NF-FN9363, *Mtsym19* and *Mtsym20* mutant lines, respectively with the *M.*

truncatula A20 genotype. To determine the map position of the mutant loci, linkage analyses

were carried out with the genotypes of the F2 individuals determined for a marker set of *M.*

truncatula (Choi *et al*., 2004; Mun *et al*., 2006). Primers of additional markers developed for

fine mapping and to identify deletion borders are listed in Table S1. To carry out allelism test

between *Mtsym19* and *Mtsym20*, F3 plants homozygous for both symbiotic loci and carrying

the opposite parental homozygous alleles for the PCR-based MtB267 marker were selected

and crossed. The success of the cross was confirmed by genotyping of the offspring and the

allelic relationship was assayed by analysing their nodulation phenotype and growth habit.

The genomic DNA of *M. truncatula* plants was isolated using the ZenoGene40 plant DNA

purification kit (Zenon Bio, Szeged, Hungary).

Identification of deletions in the genome of *M. truncatula* symbiotic mutants

The database of copy number variations identified with whole genome array-based

comparative genomic hybridization (CGH) analysis of *M. truncatula* fast neutron bombarded

mutant lines (https://medicago-mutant.dasnr.okstate.edu/mutant/index.php) were mined to

identify mutagenesis events (In/Del and SNP variations) co-segregating with the symbiotic

mutant locus of FN-NF9363.

To detect genetic alterations in the genome of symbiotic mutants *Mtsym18* (TR36), *Mtsym19*

(TR183) and *Mtsym20* (TRV43), an analysis of RNAseq data was carried out. Nodules of the

symbiotic mutants were harvested at 2 wpi with *S. medicae* WSM419 (pXLGD4) in liquid

nitrogen, total RNA was extracted with TRI Reagent (Sigma) and purified with Direct-zol

RNA MiniPrep Kit (Zymo Research, Irvine, CA, USA). RNA samples were treated with

DNaseI on Zymo-Spin columns (Zymo Research) according to the manufacturer's

instructions to remove the genomic DNA. For RNA sequencing, the quality of total RNA was

tested on an Agilent 2100 bioanalyzer followed by ribosomal RNA depletion using the

Ribominus plant kit (Invitrogen). Libraries were prepared from 1.2 µg rRNA-depleted RNA

using the Illumina TruSeq RNA Sample Preparation Kit v2 and sequenced using Illumina

NextSeq 500/550 with a minimum of 25 million reads in 150 bp paired-end mode. RNA

sequence analysis was carried out using the CLC Genomics workbench (9.5.3) software. Raw

reads (36 million for TRV43 and 35 million for TR183) from the Illumina sequencing were

trimmed (~5% of the raw reads were discarded) to remove the adaptors and low quality

ambiguous nucleotides (Quality limit 0.05, ambiguous limit 2). Trimmed reads were mapped

to mRNA reference sequence subsets obtained from the *M. truncatula* A17r5.0 genome

assembly (MtrunA17r5.0-ANR-EGN-r1.6.cds)

https://medicago.toulouse.inra.fr/MtrunA17r5.0-ANR/). The number of reads of the symbiotic

mutants were analysed between genetic markers Chr4g0020111 and Chr4g0020631.

Protein sequence analysis

Multiple alignments of NCR peptide sequences were created using the CLC Genomics

Workbench 9.5.3 program with default settings. The phylogenetic analysis was conducted on

the Phylogeny.fr online platform (http://www.phylogeny.fr/simple_phylogeny.cgi). The

alignment of the sequences was performed with the MUSCLE program (v3.8.31), the

phylogenetic tree was constructed using the maximum likelihood method using the PhylML

program (v3.1/3.0 aLRT) and visualized by TreeDyn (v198.3).

Gene expression analysis

The transcriptional activity of *NCR-new35, NCR211, NCR343, NCR169* as well as

senescence- and defence-related genes was analysed with reverse transcription quantitative

polymerase chain reaction (RT-qPCR). The expression of *NCR* genes was monitored in wt

nodules harvested at 2 and 3 wpi with *S. medicae* WSM419. For the expression analysis of

senescence- and defence-related genes, nodules of wt and *nad1-3*, NF-FN9363, *Mtsym20*,

Mtdnf4-1 and *Mtdnf7-1* mutants were harvested at 3 wpi with *S. medicae* WSM419. RNA was

extracted in the same way as for RNAseq analysis. Total RNA was quantified on Nanodrop-

1000 spectrophotometer and cDNA was prepared from 1 µg total RNA with SuperScript III

Reverse Transcriptase (Life Technologies, Invitrogen) using oligo-dT primers according to

the manufacturer's protocol. RT-qPCR reactions were performed on a LightCycler 96 (Roche,

Mannheim, Germany) using qPCRBIO SyGreen Mix (PCR Biosystems, London, UK)

according to the manufacturer's instructions. Relative expression of the target genes was

normalized using the housekeeping gene *Polypyrimidine tract-binding protein 2* (*PTB*,

MtrunA17_ Chr3g0126461) and a gene (MtrunA17_Chr3g0126781) with an ubiquitin

domain as reference genes. No RT reaction and the intron sequence of gene

MtrunA17_Chr3g0126781 was used to test cDNA samples for genomic DNA contamination.

Data were analysed with the LightCycler 96 SW1.1 software. Primer sequences used for RT-

qPCR are listed in Table S1.

Analysis of elongation and DNA content of rhizobia

To measure the bacteroid elongation, rhizobia were isolated from nodules at 16 days post

inoculation (dpi). Nodules collected in 1x PBS (pH 7.4) buffer were crushed and

homogenized with a sterile glass pestle, the debris was pelleted with centrifugation for 10 min

at 2000 rpm at 4 °C and the upper phase was filtered through 50 µm and 20 µm filters

(Sysmex/Partec CellTrics filter). Filtered bacteria were pelleted by centrifugation at 7000 rpm

for 15 min at 4 °C and resuspended in 20 µl 1xPBS buffer. Purified bacteroids were stained in

20 µM propidium iodide (PI) solution diluted in 1xPBS buffer (pH 7.4) for 30 min and the

images were captured at 40x magnification using Leica TCS SP8 confocal laser scanning

microscopy. The microscope configuration was as follows: objective lens PL Fluotar 40x oil

(NA: 1); zoom: 0.75; excitation: OPSL 552 nm laser; spectral emission detector: 573 - 696

nm. The length of bacteroids was measured on the microscope images using the ImageJ

software.

The DNA content and the size of the bacteroid populations were analysed by flow cytometry.

Nodules were chopped in BEB buffer (125 mM KCl, 50 mM Na-succinate in 50 mM N-

[Tris(hydroxymethyl)methyl]-2-aminoethanesulfonicacid sodium salt (TES) buffer (pH 7),

1% BSA) for bacteroid extraction, and filtered through a 50-µm nylon mesh. Bacteroids or

free-living bacteria suspensions were heat-shocked at 70°C for 30 min, and labelled for 10

min with 50 µg/ml of PI, a DNA fluorochrome intercalating dye. Data acquisition was made

using a Cytoflex S cytometer (Beckman Coulter). DNA content of 50,000 – 100,000 stained

bacteria/bacteroids was measured by 561 nm excitation, through a 610/20 nm band-pass filter.

Population was first gated on an SSC-Area/PI-Area dot plot; doublets were then discarded on

a PI-Area/PI-Height dot plot. Finally, DNA content or Forward Scatter (FSC) was measured

on their respective histograms.

Complementation experiments using hairy-root transformation

Constructs for genetic complementation experiments were generated using either single-site

or multisite Gateway Recombination Cloning Technology (Thermo Fisher Scientific). The

PCR amplified products containing the native promoter fragments, *NCR* gene and native 3'

UTR   sequences   (*prNCR343-1620bp::NCR343gDNA-3'UTR-543bp,   prNCR345-*

1705bp::NCR345gDNA-3'UTR-732bp,   prNCR341-1659bp::NCR431gDNA-3'UTR-853bp

and *prNCR-new35-819bp::NCR-new35gDNA-3'UTR-902bp*) were cloned into pKGW-RR

binary destination vector from entry clones generated in pDONR201 vector. The *NCR344*

gene construct was generated in pKm43GW-rolD::EGFP destination vector by recombining

the 1476 bp long promoter region, the coding sequence and 472 bp long 3'UTR fragment

cloned into entry vectors pDONRP4-P1R, pDONR221 and pDONRP2R-P3, respectively. To

investigate the effect of the substitution of the first cysteine residues to serine in the mature

peptide of NCR343, NCR-new35 and NCR211, constructs were generated in pKm43GW-

rolD::EGFP destination vector. The 1620, 819 and 2055 bp long promoter regions of

NCR343, NCR-new35 and *NCR211* were inserted into vector pDONRP4P1, respectively. The

fragments coding for NCR343-C34S, NCR-new35-C41S and NCR211-C28S were generated

by overlapping extension PCR. The *NCR211* coding sequence containing the Cys-Ser

substitution was cloned into pDONR221 vector including the first and second exons with the

translational Stop codon and a part of the subsequent second intron (*NCR211-mod.coding 1-*

2.exon-STOP-2.intron - 449bp). The amplified fragment of the rest of the second intron and

the third exon together with the 3'UTR were cloned into pDONRP2R-P3 vector (*NCR211*

2.intron 1081bp-3.exon-3'UTR 618bp). The modified coding sequence with the native 3'UTR

of *NCR343* was generated in pDONR221 vector (*NCR343-mod.coding-3'UTR 543bp*) and

NCR-new35 (*NCR-new35-mod.coding- 3'UTR 289bp*): The third entry clones in pDONRP2R-

P3 vectors carried additional 3'UTR sequences (*NCR343-3'UTR 252bp; NCR-new35-3'UTR*

613bp), respectively. Fragments for the entry clones were amplified with Phusion DNA

polymerase (ThermoFisher) or CloneAmpTM HiFi PCR Premix (Takara Bio) using the

corresponding primers listed in TableS1. Constructs were introduced into the mutant and wt

plants using *A. rhizogenes*-mediated hairy root transformation (Boisson-Dernier *et al*., 2001).

The transgenic roots were selected based on the GFP or dsRED fluorescence using a Leica

MZ10F fluorescence stereomicroscope.

Subcellular localization of NCR peptides

To investigate the localization of NCR343 and NCR-new35 peptides in symbiotic nodule

cells, constructs coding for mCherry tagged versions of the peptides were generated in

pK7m34GW-rolD::EGFP destination vector using Multisite Gateway Technology (Thermo

Fisher Scientific). The entry clones contained the same promoter fragments in pDONRP4-

P1R which were used for complementation experiments. The coding sequences of the

NCR343 and NCR-new35 peptides from the translational start site without stop codon were

inserted into pDONR221 and the mCherry fluorescent tag was cloned into the vector

pDONRP2R-P3. The fragments of the entry vectors were recombined into the destination

vector pK7m34GW-rolD::EGFP. The constructs were introduced into NF-FN9363

(pNCR343::NCR343-mCherry) and *Mtsym20* (pNCR-new35::NCR-new35-mCherry) mutant

plants with *A. rhizogenes*-mediated hairy-root transformation. The signal of fluorescence-

tagged NCR343 and NCR-new35 peptides was analysed at 4 wpi on non-fixed 65 μm thick

longitudinal and 5 μM SYTO13 stained nodule sections using a Visitron spinning disk

confocal microscope equipped with a Yokogawa CSU-W1 spinning disk unit, an Andor Zyla

4.2 Plus camera and an Olympus IX83 inverted microscope (Visitron Ssystem GmBH,

Puchheim, Germany). For imaging SYTO13, 488 nm laser excitation and 500-550 nm

emission range was used. The mCherry signal was imaged with 561 nm laser excitation and

570-640 nm emission detection range. For low magnification nodule imaging 10x UPlanFL N

dry objective (NA: 0.30) and for closeups 100x UPLXAPOXO (NA: 1.45) oil objective was

used. Composite images were created with ImageJ program.

Results

Nodules of *Medicago truncatula* deletion mutants are defective in colonization of the nitrogen fixation zone by rhizobia

398
399

The NF-FN9363, *Mtsym19* (TR183) and *Mtsym20* (TRV43) ineffective symbiotic (Fix-)

mutants were identified formerly in genetic screens for symbiotic nitrogen fixation mutants of

fast neutron bombarded (NF-FN9363) or γ-ray irradiated (*Mtsym19* and *Mtsym20*) *M.*

truncatula A17 (NF-FN9363) or Jemalong (*Mtsym19* and *Mtsym20*) populations (Xi *et al*.,

2013; Sagan *et al*., 1995; Morandi *et al*., 2005). *Mtsym19* and *Mtsym20* were reported

representing distinct symbiotic loci (Morandi *et al*., 2005) but based on the results described

below, we found that these mutants are allelic and therefore only one allele was used in some

experiments. The NF-FN9363, *Mtsym19* and *Mtsym20* ineffective mutants were inoculated

with different rhizobia strains during the mutant screens. To test the strain dependence of the

phenotype of these ineffective symbiotic mutants, the nodulation and growth phenotype of

wild-type and NF-FN9363, *Mtsym20, Mtdnf4-1* (*dnf4-1*) and *Mtdnf7-2* (*dnf7-2*) (Starker *et al*.,

2006; Domonkos *et al*., 2013) mutant plants were analysed 3 wpi with *S. meliloti* strains 1021

and FSM-MA and *S. medicae* strains ABS7 and WSM419. Mutant plants showed the

symptoms of nitrogen starvation (yellow leaves, reduced growth) 3 wpi with all the tested

rhizobia strains, even with the highly compatible *S. medicae* WSM419 strain (Terpolilli *et al*.,

2008) (Figure 1a). Mutant plants developed roundish or slightly cylindrical white nodules,

indicating the absence of leghemoglobin, with each tested rhizobia strain (Figures 1b and S1).

Nitrogen deficiency of the ineffective symbiotic mutants induced the development of

increased number of nodules and also resulted in reduced dry weight of the aerial part of

mutant plants compared with wild-type plants at 3 wpi with each rhizobia strain (Figure S1b).

To analyse the colonization of symbiotic nodules, mutant and wild-type plants were

inoculated with *S. medicae* WSM419 (pXLGD4), which constitutively expresses the *lacZ*

gene (*hemA::lacZ*). Longitudinal sections of nodules were stained for β-galactosidase activity

14 dpi and the presence of rhizobia was analysed by light microscopy. Wild-type nodules

showed the characteristic zonation of indeterminate nodules and were colonized with rhizobia

in zones IId, IIp, II-III (IZ) and III at this time point (Fig. 1b lower panel). Mutant nodules

were reduced in size compared with wild-type ones and in addition, the zones of

indeterminate nodules could be distinguished in mutant nodules. Rhizobia-infected nodule

cells were observed in the infection zone (ZIId and ZIIp) and interzone of mutant nodules

indicating that the function of the impaired symbiotic genes is essential for the differentiation
or the persistence of differentiated rhizobia in mutants NF-FN9363, *Mtsym19* and *Mtsym20*
(Fig. 1b lower panel). The nodule colonization in mutants NF-FN9363, *Mtsym19* and
Mtsym20 was similar to that in *M. truncatula dnf4* and *dnf7* mutants defective in NCR
peptides NCR211 and NCR169, respectively (Kim *et al*., 2015; Horvath *et al*., 2015).
Therefore, and for the reasons described below, the mutants NF-FN9363, *Mtsym19* and
Mtsym20 were analysed along with the symbiotic mutants *Mtdnf4-1* and *Mtdnf7-2*.
To analyse nodule colonization in more detail, mutant and wild-type nodule sections were
stained with nucleic acid-binding dye SYTO13 at 14 dpi with *S. medicae* WSM419 and
examined by laser-scanning confocal microscopy that enabled the visualization of stained
bacteria and plant nuclei fluoresced in green, the autofluorescent cell wall and accumulated
polyphenolic compounds (red channel) simultaneously (Figure 2). Nodules of wild-type
plants showed rhizobia-colonized cells in the infection zone (ZIId, ZIIp), interzone (IZ) and
nitrogen fixation zone (ZIII) (Figure 2a1-a8). The bacterial invasion of ZII and IZ of mutant
and wild-type nodules was similar, albeit the interzone, where bacteria complete their
differentiation into symbiotic form, was more extensive and enlarged in mutant nodules
compared with wild-type nodules (Figure 1b and Figure 2a1-f1). In contrast to wild-type
nodules, the cells in nitrogen fixation zones of mutant nodules were not colonized by rhizobia
but showed sporadic autofluorescence (Figure 2b1-f1 and b8-f8). Autofluorescence in nodule
tissues often arises from phenolic-compounds derived from pathogen-induced lignification
that generally results in robust and extensive defence responses which were characteristics of
some *M. truncatula* symbiotic mutants (Bourcy *et al*., 2013; Wang *et al*., 2016; Domonkos *et*
al., 2017). In contrast, the early induced decomposing process, the premature senescence is
frequently observed in ineffective symbiotic nodules (Van de Velde *et al*., 2006). Early
senescence generates less intensive and sporadic autofluorescence that was found previously
in *Mtdnf4-1* and *Mtdnf7-2* nodules (Kim *et al*., 2015; Horváth *et al*., 2015). In addition to the
difference in the extension and intensity between the defence-reactions and premature
senescence-related autofluorescence, the two responses can be distinguished based on
activation of marker genes specific for each process. The transcriptional activities of
senescence and defence response-specific marker genes were monitored in NF-FN9363,
Mtsym20, *Mtdnf4*, *Mtdnf7-2*, the *nad1-3* (*nodules with activated defense 1-3;* Domonkos *et*
al.,) mutant and Jemalong wild-type nodules at 3 wpi using RT-qPCR. A strong induction of
cysteine proteinase genes *MtCP2* and *MtCP6* associated with nodule senescence in *M.*
truncatula (Guerra *et al*., 2010) was detected in the ineffective symbiotic nodules of all

studied mutants (Figure S2a). In contrast, marker genes of defence responses, a chitinase

(*MtCHI*) and *MtPR10* (Class-10 *PATHOGENESIS-RELATED PROTEIN*) genes were not

induced in the mutants NF-FN9363, *Mtsym20*, *Mtdnf4* and *Mtdnf7-2* but showed high

transcriptional activity in the *M. truncatula nad1-3* mutant (Figure S2b and c), that controls

defence-like responses in symbiotic nodules, indicating that the autofluorescence was the

result of premature senescence in the nitrogen fixation zones of NF-FN9363, *Mtsym19,*

Mtsym20, *Mtdnf4-1* and *Mtdnf7-2* nodules (Figure 2b8-f8).

Higher magnification of SYTO13-stained nodule sections revealed that infected cells in ZII

and IZ exhibited similar morphology in mutant and wild-type nodules (Figure 2a1-f5).

Bacterial release was normal in the distal part of ZII (ZIId) of mutant nodule cells (Figure 2

a3, b3, c3, d3, e3 and f3) and rhizobia commenced to differentiate in the proximal part of ZII

(ZIIp) (Figure 2a4, b4, c4, d4, e4 and f4). These nodule cells in ZIIp, characterized by

moderately elongated rhizobia with intense fluorescence and enlarged vacuoles, were clearly

visible in wild-type (Figure 2a2 and a4), *Mtdnf7-1* (Figure 2b2 and b4) and *Mtdnf4-1* (Figure

2c2 and c4) nodules but vacuole extension was less pronounced in nodules of FN9363 (Figure

2d2 and d4), *Mtsym20* (Figure 2e2 and e4) and *Mtsym19* (Figure 2f2 and f4). Cells in the first

layer of the nitrogen fixation zone showed similar morphology, possessing compressed

vacuoles, in wild-type and all mutant nodules although rhizobia were slightly shorter and

disorganized in mutant symbiotic cells (Figure 2a5, a6, b5, b6, c5, c6, d5, d6, e5, e6, f5 and

f6). Although this region is composed of a single layer of cells in wild-type nodules (Gavrin *et*

al., 2014; Figure 2 a1), we observed the extension of this part of nodules in all mutants

(Figure 2b1, c1, d1, e1, and f1) indicating the arrest of further differentiation of infected

nodule cells. Symbiotic cells with large vacuoles and differentiated bacteroids oriented

towards these vacuoles were observed in the nitrogen fixation zone of wild-type nodules

(Figure 2a6 and a8). In contrast, vacuoles remained reduced in IZ cells of mutant nodules,

resembling to the morphology of cells with compressed vacuoles in interzones (Figure 2b7,

c7, d7, e7 and f7). Mutant nodules did not contain infected cells in the region corresponding

to the mature nitrogen fixation zone of wild-type nodules but sporadic autofluorescence and

non-differentiated rod shape saprophytic bacteria released from the infection threads were

detected occasionally in this zone (Figure 2a8, b8, c8, d8, e8 and f8).

Bacteroid elongation and morphology are impaired in NF-FN9363 and *Mtsym20* nodules

Rhizobia undergo morphological changes including cell enlargement and genome
amplification in the indeterminate nodules of *M. truncatula* (Mergaert *et al*., 2006) resulting
in elongated (E-type) bacteroids (Montiel *et al*., 2017). The morphology of rhizobia in the
interzone cells of mutant nodules indicated the impaired elongation of bacteroids (Figure 2b5-
7, c5-7, d5-7, e5-7 and f5-7). In order to determine the level of differentiation, the length and
ploidy level of bacteria were measured. Bacteria were isolated from mutant and wild-type
nodules 14 dpi, stained with PI and imaged by confocal laser scanning microscopy. The size
of at least 1200 bacterial cells was measured using the ImageJ software and the relative ratio
of bacteria in different size ranges was compared between wild-type and mutant nodules.
Bacterial populations isolated from mutant nodules showed a shift to the smaller size range
compared with wild-type sample indicating an imperfect elongation of bacteroids in *Mtdnf4-*
1, *Mtdnf7-2*, NF-FN9363 and *Mtsym20* nodules (Figure 3a).
The size and DNA content of rhizobia isolated from *Mtdnf4-1*, NF-FN9363 and *Mtsym20*
mutant and wild-type nodules were further analysed by flow cytometry. Isolated bacteroids
and cultured bacteria were purified and stained with PI. The analysis of FSC intensity, that is
proportional to the size of bacteroids, showed that the majority of bacteroids isolated from
wild-type nodules were enlarged compared with cultured rhizobia (Figure 3b). The bacterial
population purified from mutant nodules contained a mixture of smaller sized cells peaked at
the free-living rhizobia and at smaller sized bacteriods compared with wild-type samples. This
finding was consistent with length measurements of bacteriods which showed the higher
proportion of shorter rhizobia in mutant nodule cells compared with bacteria isolated form
wild-type nodules (Figure 3a). The mean DNA content of bacteroids isolated from mutant
nodules was higher than the peak of 1C or 2C ploidy levels of cultured *S. medicae* WSM419
cells indicating that bacteroids isolated from mutant nodules were polyploids (Figure 3b). The
peaks of the DNA content of rhizobia purified from mutant and wild-type nodules were very
similar suggesting an advanced genome amplification of bacteroids in mutant nodules.
However, the population of bacterial cells isolated from mutant nodules was narrower at a
higher fluorescent range compared with rhizobia purified from wild-type nodules indicating
fewer numbers of bacteroids with amplified genomes at the highest degree. The analysis of
bacterial size and DNA content of rhizobia showed the similar positive correlation between
genome amplification and enlargement of bacteroids detected earlier (Mergaert *et al*., 2006).
To detect alterations in mutant nodules in more detail, the ultrastructure of wild-type and NF-
FN9363, *Mtsym20*, *Mtdnf4-1* and *Mtdnf7-2* mutant nodules was analysed by scanning electron
microscopy (SEM) 18 dpi with *S. medicae* WSM419. Consistent with the light and

fluorescent microcopy images, no visible difference was detected in the morphology and
invasion of nodule cells in the distal part of ZII of wild-type and mutant nodules (Figure S3a2,
b2, c2, d2, e2, a3, b3, c3, d3 and e3). However, rhizobia around the vacuoles did not fill up
completely the infected cells in the proximal part of IZ of mutant nodules and this
discontinuity between the rhizobia and plant cell walls probably indicates the induction of
degradation of cytoplasm and rhizobia (Figure S3a4, b4, c4, d4 and e4). The shrinking of cell
content and the aggregation of rhizobia was more apparent in the first cell layer in IZ of
mutant nodules compared with wild-type nodule cells, which were fully packed with rhizobia,
indicating the advanced collapse of infected mutant cells (Figure S3b5-e5). Close-up images
of free-living (Figure S3f1) and symbiotic rhizobia in ZIId and ZIIp of wild-type nodules
showed rough-surfaced rhizobia (Figure S3a2 and a3) that probably corresponds to the
folding of peribacteroid membranes observed in this region on transmission electron
microscopy images of bacteroids of a former study (Vasse *et al.*, 1990). As bacteroid
differentiation progresses, this morphology changes and smooth-surfaced rhizobia are
detected in the IZ and ZIII in wild-type nodules (Figure S3a4 and a6). This alteration of
bacterial surface was apparent in the distal part of IZ of all mutant nodules (IZd; Figure S3b4,
c4, d4 and e4) implying that bacteroids differentiate similarly in IZ cells of mutant and wild-
type nodules. However, disorder and aggregation of bacteroids, often tangled with cell debris,
were observed in the proximal IZ cells of mutant nodules (IZp; Figure S3b5, c5, d5 and e5)
compared with healthy infected cells of wild-type nodules containing elongated bacteroids
orientated towards the vacuoles (Figure S3a4). This disintegration, characterized by
vacuolated cells and degraded bacteroids, was more advanced in the few infected cells of ZIII
of mutant nodules indicating that decayed cell content was resorbed (Figure S3b6, c6, d6 and
e6). These data reveal that rhizobia are able to differentiate to bacteroids in *Mtdnf4-1*, *Mtdnf7-
2*, NF-FN9363 and *Mtsym20* nodules but their complete transition is either disrupted or the
persistence of bacteroids is defective in mutant nodules.
556
557

Ineffective symbiotic *Medicago truncatula* mutants NF-FN9363, *Mtsym19* and *Mtsym20*
are defective in *NCR* genes
Genetic mapping combined with exploiting the high-density genome array-based comparative
genomic hybridization (aCGH) platform of *M. truncatula* (Chen *et al.*, 2017) or facilitated
with transcriptome analysis of symbiotic mutants were applied to identify the impaired genes
in symbiotic mutants NF-FN9363, *Mtsym19* and *Mtsym20*. Mutant plants were crossed to *M.*

truncatula genotype A20 and genetic mapping was carried out using F2 segregating

populations.

The mutant locus of NF-FN9363 was positioned on chromosome 6 (LG 6) below the genetic

marker Crs towards MtB178 (Figure S4a). To facilitate the identification of gene defective in

NF-FN9363, the genomic positions of copy number variations detected in the mutant line

using the array-based CGH (https://medicago-mutant.dasnr.okstate.edu/mutant/index.php)

were analysed. A pile of probesets corresponding to the genomic position of the symbiotic

locus of NF-FN9363 indicated a potential large deletion in the mutant genome. The deletion

was verified by PCR-based markers that defined an almost 500 kb deletion in the genome of

NF-FN9363 (Figure 4A and S5, Table S1).

The mutant loci of *Mtsym19* and *Mtsym20* were positioned in the same genomic region on

chromosome 4 (LG 4) between genetic markers 4g0020111, 4g0020421 and 4g0020631

(Figure S4a). Former genetic analysis suggested that *Mtsym19* and *Mtsym20* belong to

different complementation groups (Morandi *et al*., 2005). However, the similar position of the

mutant loci of *Mtsym19* and *Mtsym20* indicated either the defect of two different

neighbouring genes or the malfunction of the same gene in the two mutants. In order to verify

their allelic relationship, an allelism test was carried out using F3 mutant plants selected from

the populations of *Mtsym19* x A20 and *Mtsym20* x A20 crosses. Two F3 maternal

homozygote plants for the PCR-based genetic marker MtB267 were selected from the

Mtsym19 mapping population and two F3 plants showing paternal homozygous genotype for

the same genetic marker were chosen from the *Mtsym20* mapping population and crossed to

generate hybrid plants. The different genotypes of the parental lines allowed us to test

heterozygosity of the progeny of the F3 mutant plants. The progeny of the three successful

crossings were all Fix– (n=8) indicating that *Mtsym19* and *Mtsym20* belong to the same

complementation group (Figure S4b).

We presumed that the mutations in *Mtsym19* and *Mtsym20* affected the level of transcripts of

the genes impaired in the mutant plants. In order to facilitate gene identification, the transcript

abundance in nodules of *Mtsym19, Mtsym20* and the symbiotic mutant *Mtsym18*, belonging to

an independent symbiotic complementation group (Morandi *et al*., 2005), was analysed at 2

wpi with *S. medicae* WSM419. Filtered sequence reads were mapped against the *M.*

truncatula genome assembly (A17r5.0) and the number of reads was analysed in the genomic

region between genetic markers 4g0020111 and 4g0020631. Absent or reduced number of

reads aligned to a short non-specific sequence, were detected between gene models

4g0020281 and 4g0020321 in *Mtsym19,* and between 4g0020281 and 4g0020301in *Mtsym20*

compared with *Mtsym18*, indicating deletions in these genomic regions (Figure 4B and S6a).

Further analysis by PCR amplifications identified a less than 45-kb and a less than 32 kb

deletion in the *Mtsym19* and *Mtsym20* genomes, respectively. The deletion in *Mtsym20*

spanned three genes encoding for two putative proteins and a Nodule-specific Cysteine-Rich

(NCR) peptide (Figure S6b).

Deletion of gene *NCR343* is responsible for the symbiotic defect of mutant NF-FN9363

The deletion in NF-FN9363 removed more than 30 predicted genes or gene models including

four *NCR* genes, *NCR341* (6g0467551), *NCR343* (6g0467511), *NCR344* (6g0467581) and

NCR345 (6g0467571) which were the primary candidates responsible for the symbiotic

phenotype of NF-FN9363 (Figure S5b). All four *NCR* genes code for peptides with four

cysteines in conserved positions and the sequence analysis of the four mature NCR peptides

revealed that NCR341, NCR344 and NCR345 showed higher similarity to each other

compared with NCR343 (Figure 4c). The most similar mature peptides of NCR341 and

NCR344 are both cationic (pI =8.67 and 7.95), the mature peptide NCR345 is anionic (pI =

4.59), while the cleaved product of NCR343 results in a slightly anionic (pI = 6.33) mature

peptide. In order to define that the lack of what gene or genes caused the symbiotic phenotype

of NF-FN9363, genetic complementation experiments were carried out. The genomic

fragments of the genes *NCR341*, *NCR343*, *NCR344* and *NCR345* containing at least 1.5 kb

native promoters and native 3' UTRs were introduced into NF-FN9363 roots using *A.*

rhizogenes-mediated hairy root transformation. Transformed roots were detected by either

DsRed or green fluorescent protein markers (Figure S7). The roots were inoculated with *S.*

medicae WSM419 (pXLGD4) and nodules were stained for β-galactosidase activity at 4 wpi.

Nodule cells in nitrogen fixation zone of mutant NF-FN9363 were devoid of rhizobia (Figure

1b), therefore we assessed the complementation of nodules based on the presence of rhizobia

in ZIII. White undeveloped nodules with colonized cells by rhizobia in ZII and IZ were

detected on roots transformed with empty vector or with the genes *NCR341*, *NCR344* and

NCR345 indicating that these *NCR* genes were not able to restore the effective symbiotic

interaction in NF-FN9363 (Figure 5a and Figure S7g1-i3 and l1-m). The nodules developed

on the roots of NF-FN9363 transformed with the construct of *NCR343* were elongated and

pink, and the bacterial invasion of ZIII in these nodules was similar to wild-type nodules,

suggesting that these were functional nodules on NF-FN9363 roots (Figure 5a and Figure

S7f1-f3 and k1-k3). The rescue of the symbiotic phenotype of NF-FN9363 with gene *NCR343*

indicated that the loss of this gene caused the ineffective symbiotic phenotype of mutant NF-
FN9363.

The lack of *NCR-new35* is responsible for the ineffective symbiotic phenotype of
Mtsym19* and *Mtsym20

The identified deletions in mutants *Mtsym19* and *Mtsym20* overlapped by about 32 kb
removing three genes including the gene *NCR-new35,* formerly termed *NCR014* (Bang *et al.*,
2017) (Figure 4b and S6B), which made *NCR-new35* the best candidate for *MtSYM19* and
MtSYM20. In order to confirm that the deletion of *NCR-new35* caused the ineffective
symbiotic phenotype of *Mtsym19* and *Mtsym20*, roots of the two mutants were transformed
with the wild-type *NCR-new35* gene controlled by an 819 bp native promoter fragment and
inoculated with *S. medicae* WSM419 (pXLGD4). DsRed-fluorescent transgenic *Mtsym19* and
Mtsym20 roots expressing the gene *NCR-new35* developed elongated and pink nodules
(Figure S7b1-b3, d1-d3 and o1-o3) in contrast to white round-shaped or slightly cylindrical
nodules formed on empty vector-transformed roots of mutant plants (Figure S7c1-c3, e1-e3
and p1-p3). In order to visualize bacterial invasion, longitudinal sections of 4-week-old
nodules were stained for β-galactosidase activity. Rhizobia-colonized cells were only
observed in zones II and IZs of nodules formed on empty vector-transformed *Mtsym19* and
Mtsym20 roots (Figure 5b). Nodules developed on mutant roots transformed with the *NCR-*
new35 construct showed the typical zonation of indeterminate nodules with colonized cells in
the nitrogen fixation zone like nodules on empty vector-transformed wild-type roots (Figure
5b, 5c and Figure S7a1-a3, j1-j3). The restoration of nodule colonization in ZIII of mutant
nodules confirmed that gene *NCR-new35* corresponds to *MtSYM19* and *MtSYM20*.

Replacement of a single cysteine residue in NCR211, NCR343 and NCR-new35 abolishes
their symbiotic function

NCR peptides usually contain four or six cysteine residues in conserved positions (Alunni *et*
al., 2007). We previously demonstrated that the replacement of a single or multiple cysteine
residues of the symbiotic peptide NCR169 ceased its activity *in planta* indicating that each
cysteine residue is essential for the symbiotic function of NCR169 (Horvath *et al.*, 2015). To
confirm the requirement of cysteine residues for the function of NCR-new35, NCR343 and
NCR211, we introduced constructs coding for modified peptides, wherein the first cysteines
were substituted for serines, into the roots of *Mtsym20*, NF-FN9363 and *Mtdnf4* mutant
plants, respectively using *A. rhizogenes*-mediated hairy root transformation. The nodules on

the roots of mutants transformed with the modified *NCRs* driven by their native promoters
were small and white, suggesting that these were malfunctioning nodules (Figure 5). These
nodules did not show the typical zonation of indeterminate nodules and the proximal regions
of the nodules corresponding to the nitrogen fixation zones were devoid of bacteria, indicating
that the first cysteine residues are essential for the function of NCR-new35, NCR343 and
NCR211.

***NCR-new35* is expressed low in symbiotic cells compared with *NCR169*, *NCR211* and**
NCR343
The members of the large family of *NCR* genes in *M. truncatula* are almost exclusively
expressed in symbiotic nodule cells and they are activated in successive waves during nodule
differentiation (Guefrachi *et al*., 2014). In order to investigate the expression of genes *NCR-*
new35 and *NCR343*, their activity was monitored with the *GUS* reporter gene and RT-qPCR
analysis and we also analysed the nodule transcriptome data of different *M. truncatula* nodule
zones obtained by laser-capture microdissection (LCM) (Roux *et al*., 2014).
To analyse and compare the expression pattern of *NCR-new35* and *NCR343* with the
previously identified *NCR169* and *NCR211* genes essential for effective nitrogen fixation
(Horvath *et al*., 2015; Kim *et al*., 2015), the promoters of these *NCR* genes were fused to the
β-glucuronidase (*GUS*) reporter gene and the constructs were introduced into wild-type *M.*
truncatula roots using *A. rhizogenes*-mediated hairy root transformation. Nodules were
subjected to histochemical staining for GUS activity at 2 and 3 wpi with *S. medicae* strain
WSM419. *NCR343*, *NCR211* and *NCR169* promoters showed activity predominantly in the
interzone of two-week-old nodules (Figure 6c-e). In contrast, a low level of *GUS* expression
was restricted mainly to the first few cell layers of the interzone and even weaker activity was
found sporadically in the distal part of nitrogen fixation zone when the reporter gene was
driven by the promoter of *NCR-new35* (Figure 6b). The activity of these promoters is in
agreement with LCM data (Roux *et al*., 2014), confirming that *NCR-new35*, *NCR343*,
NCR211 and *NCR169* are mainly active in the IZs of two-week-old nodules and *NCR-new35*
is expressed at a much lower level compared with the other three *NCR* genes (Figure 6). The
analysis of the promoter activities with the *GUS* reporter gene at 3 wpi revealed that the
expression of *NCR169* and *NCR343* and to a less extent the activity of *NCR211* extended to
the nitrogen fixation zone (Figure 6g-j). In order to further validate their expression in
nodules, we monitored the transcript levels of *NCR-new35*, *NCR343*, *NCR211* and *NCR169*

with RT-qPCR in nodules harvested at 2 and 3 wpi. In agreement with the LCM

transcriptome data, *NCR343*, *NCR211* and *NCR169* genes showed high relative expression

compared with *NCR-new35* (Figure 6k). These transcriptome and *GUS* activity data indicate

that *NCR-new35* shows different spatial activity and lower expression intensity compared

with the other three essential NCR peptide genes.

NCR-new35 and NCR343 peptides localize to symbiosomes

NCR peptides usually have a conserved signal peptide which cleavage is essential for

targeting of NCR peptides to the bacteroids (Wang *et al*., 2010). Previous studies

demonstrated the localization of NCR169 and NCR211 peptides to the bacteroids in the

symbiotic cells in the IZ and nitrogen fixation zone (Horvath *et al*., 2015; Kim *et al*., 2015).

To make certain the subcellular localization of NCR-new35 and NCR343 peptides,

translational fusions to the *mCherry* reporter gene driven by native *NCR-new35* or *NCR343*

promoters were generated and introduced into *Mtsym20* and NF-FN9363 mutant roots,

respectively using hairy-root transformation. The functional complementation of the mutant

nodules indicated that the fusion proteins retained their activity and the fluorescent tag did not

perturb the function of NCR-new35-mCherry and NCR343-mCherry proteins (Figure 7). The

red fluorescence of NCR-new35-mCherry and NCR343-mCherry proteins was detected in the

IZ and nitrogen fixation zone. The signal of the NCR-new35-mCherry fusion protein is

partially contradictory to the expression pattern of *NCR-new35* detected at 3 wpi with rhizobia

(Figure 6g) and points to the sustained stability of the NCR-new35-mCherry fusion protein in

the nitrogen fixation zone. In contrast to NCR-new35-mCherry, the subcellular localization of

NCR343-mCherry corresponded completely to the spatial expression pattern of *NCR343*

(Figure 6i). Higher magnification of symbiotic cells showed co-localization of mCherry-

tagged NCR-new35 and NCR343 with SYTO13-stained rhizobia (Figure 7c and f).

Discussion

The *M. truncatula* genome contains around 700 *NCR* genes which are almost exclusively

expressed in symbiotic nodule cells (Roux et al. 2014). The encoded NCR peptides mediate

terminal differentiation of the symbiotic partner (Van de Velde *et al*., 2010) but the high

number of *NCR* genes raises the question of why *M. truncatula* evolved so many genes and

how many of them are required for the establishment of symbiotic nitrogen fixation. Forward

and reverse genetic analysis identified that peptides NCR169, NCR211 and NCR247 are

crucial for effective nitrogen-fixing symbiosis (Horvath *et al.*, 2015; Kim *et al.*, 2015; Sankari

et al., 2022) which prompted us to continue the analysis of ineffective *M. truncatula*

symbiotic mutants to identify further essential *NCR* genes. Based on the nodulation phenotype

of mutants defective in genes *NCR169*, *NCR211* and *NCR247*, we focused on symbiotic

mutants *Mtsym19*, *Mtsym20* and NF-FN9363 that developed slightly elongated white nodules

with zonation and invaded nodule cells in the distal part of the nodules. The histological

analysis revealed that nodule zonation and the morphology of cells between the meristem and

the transition zone in mutant nodules were very similar to wild-type nodules indicating the

formation of differentiated symbiotic cell. However, the extended transition zone in mutant

nodules compared with wt nodules indicated a defect in the differentiation process. This was

in agreement with a former study detecting a higher ploidy level of nodule cells in *Mtsym19*

and *Mtsym20* mutants compared with TE7 mutant, which is defective in the *IPD3* gene

(Ovchinnikova *et al.*, 2011; Horvath *et al.*, 2011) with non-differentiated symbiotic nodule

cells, but *Mtsym19* and *Mtsym20* nodule cells had lower endoreduplication index than wild

type ones (Maunoury *et al.*, 2010). The analysis of the length and DNA content of bacteroids

isolated from *Mtsym19*, *Mtsym20* and NF-FN9363 nodules showed that mutant nodules

contained lower proportion of longer and endoreduplicated bacteroids compared with wt

nodules. These results indicated that the differentiation of nodule cells and hosted bacteroids

was advanced but not complete in *Mtsym18*, *Mtsym19* and NF-FN9363 mutants. This finding

was in accordance with the histological and transciptome analysis of *Mtsym19* and *Mtsym20*

nodules which detected the activation of a second transcriptome-switch characteristic of late

Fix mutant plants (Maunoury *et al.*, 2010). The expression analysis of senescence and

defence-related marker genes suggests that the lack of colonization of the cells in the mature

part of nitrogen fixation zone and thus the absence of effective nitrogen fixation in mutant

nodules induced premature senescence of symbiotic cells.

The deletion in the NF-FN9363 mutant includes four *NCR* genes in close proximity. *NCR341*,

NCR343, *NCR344* and *NCR345* showed high similarity at amino acid level which implies that

these homologues have recently evolved at the scale of evolution by tandem duplication.

However, only NCR343 is able to restore the symbiotic phenotype of NF-FN9363 indicating

a specialization of this peptide for the symbiotic interaction between *M. truncatula* Jemalong

A17 and the tested rhizobia. Further analysis by exchanging residues or fragments between

NCR341, NCR343, NCR344 and NCR345 peptides could reveal the unique region(s) of

NCR343 responsible for its distinct biological function. The identified deletions in *Mtsym19*

and *Mtsym20* overlapped indicating that they are defective in the same gene which result is in

contradiction to the previous allelism test that defined them as being in distinct

complementation groups (Morandi *et al.*, 2005). Our allelism test and complementation

experiments with gene *NCR-new35* clearly confirmed that *Mtsym19* and *Mtsym20* are allelic.

The spatial expression analysis of *NCR* genes identified by forward genetic approach and

crucial for symbiosis (Horvath *et al.*, 2015; Kim *et al.*, 2015 and this study) was analysed by a

GUS-reporter assay which showed that *NCR169*, *NCR211* and *NCR343* are active in the

infected cells of IZ and ZIII but revealed that *NCR-new35* can be detected only in IZ cells.

This observation is in correlation with RNA sequencing data obtained from laser-capture

microdissected nodule zones (Roux *et al.*, 2014). In addition to the expression of *NCR-new35*

being detected predominantly only in IZ, its activity was much weaker compared with the

other three *NCR* genes suggesting that the transcriptional intensity of an *NCR* gene does not

correlate with the necessity of the *NCR* genes. Although the promoter activity of *NCR-new35*

is barely detectable in nitrogen fixation zone, the encoded peptide is abundant in this zone

suggesting a slow turnover or the enhanced stability of NCR-new35. Formerly, the

identification of several NCR peptides, which are expressed during the early stages of nodule

colonization, in mature bacteroids suggested the sustained stability of these NCR peptides

(Durgo *et al.*, 2015). Correspondingly to the previous finding on NCR169 peptide (Horvath *et*

al., 2015), both NCR343 and NCR-new35 co-localise with bacteroids forming a ring-shaped

fluorescent signal. Apparently, the fluorescent protein fusions of NCR peptides, which are

functional and able to rescue the symbiotic phenotype of the corresponding mutants, are

localised around the bacteroids. However, a previous proteomic study identified NCR343 and

NCR169, but not NCR211 and NCR-new35, in bacteroids (Durgo *et al.*, 2015) indicating that

at least a subset of the fluorescent-tagged NCR fusion molecules, conceivably following

proteolytic cleavage, enter into the bacteroids.

The *NCR343* gene consists of one exon but *NCR-new35* has two exons, which is the most

common gene structure of *M. truncatula NCR* genes. The encoded mature peptides are

composed of 43 and 47 amino acids, respectively, including four cysteine residues at

conserved positions, the characteristic feature of *M. truncatula* NCR peptides. The charge of

the mature peptides is slightly and strongly anionic (isoeletric point (pI)=6.34 for NCR343

and 4.78 for NCR-new35). The charges of the five NCR peptides (NCR169 pI=8.45, NCR211

pI=5.38, NCR247 pI=10.15, NCR343 and NCR-new35), proved to be essential for symbiosis

(Horvath *et al.*, 2015; Kim *et al.*, 2015; Sankari *et al.*, 2022) vary between 4.78 and 10.15

indicating that anionic, neutral and cationic NCR peptides could be essential for nitrogen-

fixing symbiosis in *M. truncatula*. The formation of intramolecular disulfide bonds between
the conserved cysteine residues in NCR044 and NCR169 peptides was experimentally
verified (Velivelli *et al.*, 2020; Isozumi *et al.*, 2021) and the *in planta* functional requirement
of cysteine residues was demonstrated with substitution of any cysteine with serine which
resulted in inactivation of NCR169 (Horvath *et al.*, 2015). The replacement of the first
cysteine residue with serine in NCR343 and NCR-new35 also abolished the symbiotic activity
of the peptides implying that the requirement of cysteine residues for the *in planta* activity of
NCRs is a common feature. Apart from the four cysteine residues in conserved position, the
five crucial NCR peptides show high sequence variation but the presence of a valine, an
isoleucine and one or two aspartic acid residues between the first two cysteines is dominant
(Figure 4d). The relevance of these residues for the structure and symbiotic activity of NCR
peptides requires further investigation. In brief, the only common feature among the five
identified NCR peptides essential for symbiosis thus far (NCR169, NCR211, NCR247,
NCR343 and NCR-new35) is that they belong to the subgroup of peptides with four
conserved cysteine residues.

Based on the observation that terminally differentiated bacteroids are symbiotically more
effective compared with reversibly differentiated ones (Oono *et al.*, 2010), it is widely
accepted that the controlled bacteroid growth in nodules provides increased fitness benefits
for the host. In IRLC legumes and some Dalbergoid legume species, rhizobia undergo
terminal differentiation provoked by NCR peptides (see reviews Pan & Wang, 2017; Downie
& Kondorosi, 2021; Czernic *et al.*, 2015). The large number of *NCR* genes of *M. truncatula*
induced in consecutive steps and showing overlap expression which implies that these
peptides might function together to optimise the interaction between rhizobia and the host,
and potentially some of them compensate the negative effect of other NCR peptides
(Mergaert, 2018). It has been presumed that there is a core set of *M. truncatula* NCRs with
common functions, loss of which results in the termination of bacteroid differentiation and/or
losing the viability of hosted rhizobia (Pan & Dong, 2017; Roy *et al.*, 2020). Till now
NCR169, NCR211 and NCR247 were identified as essential peptides for symbiosis but our
work provides two additional NCR peptides, NCR343 and NCR-new35 required for effective
nitrogen fixation between *M. truncatula* and *Sinorhizobium* species. The identification of
novel crucial NCRs implies that further forward and reverse genetic studies might extend the
cluster of universal and essential NCR peptides of *M. truncatula*.

Acknowledgements

This work was supported by the Hungarian National Research Fund/National Research,

Development and Innovation Office grants OTKA- 67576, PD104334/108923, 106068,

119652 and 120122/120300, PD-121110 and PD-132495 as well as by the Collaborative

Research Programme ICGEB Research Grant HUN17-03. The present work has benefited

from Imagerie☐Gif core facility supported by l'Agence Nationale de la Recherche (ANR-11-

EQPX-0029/Morphoscope, ANR-10-INBS-04/FranceBioImaging ;

ANR☐11☐IDEX☐0003☐02/ Saclay Plant Sciences). We thank H. Cs. Tolnainé and K. Miró

for their skillful technical assistance and A. Farkas for carrying out sample preparation and

SEM imaging,

Author contributions

B.H, Á.D., R.C. and P.K designed the project. B.H, B.G, M.T., Á.D, F.A., F.S., Y.C., M.B.,

J.B.B., and Z.T. performed the experiments, ÁD and Z.S. carried out data analysis. P.K wrote

the manuscript.

Data availability

All data supporting the findings of this study are available within the paper and within its

supplementary materials published online.

Competing interests

None declared.

References

Alunni B, Kevei Z, Redondo-Nieto M, Kondorosi A, Mergaert P, Kondorosi E. 2007.
Genomic organization and evolutionary insights on GRP and NCR genes, two large
nodule-specific gene families in *Medicago truncatula*. *Molecular Plant-Microbe*
Interactions **20**(9): 1138-1148.
Beringer JE 1974. R-factor transfer in *Rhizobium leguminosarum*. J Gen Microbiol. 421-429.
Boisson-Dernier A, Chabaud M, Garcia F, Becard G, Rosenberg C, Barker D. 2001.
Agrobacterium rhizogenes-transformed roots of *Medicago truncatula* for the study of
nitrogen-fixing and endomycorrhizal symbiotic associations. *Molecular Plant-*

Microbe Interactions **14**(6): 695-700.

Boivin C, Camut S, Malpica C, Truchet G, Rosenberg C. 1990. *Rhizobium meliloti* genes
encoding catabolism of trigonelline are induced under symbiotic conditions. *Plant Cell*
2(12): 1157-1170.

Bourcy M, Brocard L, Pislariu CI, Cosson V, Mergaert P, Tadege M, Mysore KS,
Udvardi MK, Gourion B, Ratet P. 2013. *Medicago truncatula* DNF2 is a PI-PLC-
XD-containing protein required for bacteroid persistence and prevention of nodule
early senescence and defense-like reactions. *New Phytologist* **197**(4): 1250-1261.

Chen YH, Wang XF, Lu SF, Wang HC, Li SB, Chen RJ. 2017. An Array-based
Comparative Genomic Hybridization Platform for Efficient Detection of Copy
Number Variations in Fast Neutron-induced *Medicago truncatula* Mutants. *Jove-*
Journal of Visualized Experiments (129).

Czernic P, Gully D, Cartieaux F, Moulin L, Guefrachi I, Patrel D, Pierre O, Fardoux J,
Chaintreuil C, Nguyen P, et al. 2015. Convergent Evolution of Endosymbiont
Differentiation in Dalbergioid and Inverted Repeat-Lacking Clade Legumes Mediated
by Nodule-Specific Cysteine-Rich Peptides. *Plant Physiology* **169**(2): 1254-1265.

de Bang TC, Lundquist PK, Dai XB, Boschiero C, Zhuang ZH, Pant P, Torres-Jerez I,
Roy S, Nogales J, Veerappan V, et al. 2017. Genome-Wide Identification of
Medicago Peptides Involved in Macronutrient Responses and Nodulation. *Plant*
Physiology **175**(4): 1669-1689.

Domonkos A, Horvath B, Marsh JF, Halasz G, Ayaydin F, Oldroyd GED, Kalo P. 2013.
The identification of novel loci required for appropriate nodule development in
Medicago truncatula. *BMC Plant Biology* **13:157**.

Domonkos A, Kovacs S, Gombar A, Kiss E, Horvath B, Kovats GZ, Farkas A, Toth MT,
Ayaydin F, Boka K, et al. 2017. NAD1 Controls Defense-Like Responses in
Medicago truncatula Symbiotic Nitrogen Fixing Nodules Following Rhizobial
Colonization in a BacA-Independent Manner. *Genes* **8**: 387

Downie JA, Kondorosi E 2021. Why Should Nodule Cysteine-Rich (NCR) Peptides Be
Absent From Nodules of Some Groups of Legumes but Essential for Symbiotic N-
Fixation in Others? Front. Agron.

Durgo H, Klement E, Hunyadi-Gulyas E, Szucs A, Kereszt A, Medzihradszky KF,
Kondorosi E. 2015. Identification of nodule-specific cysteine-rich plant peptides in
endosymbiotic bacteria. *Proteomics* **15**(13): 2291-2295.

Garcia J, Barker DG, Journet EP 2007. Seed storage and germination. In: Mathesius U,
Journet EP, Sumner LW eds. *The Medicago truncatula Handbook*. The Samuel
Roberts Noble Foundation, Ardmore, OK: Samuel Roberts Noble Foundation, 1-9.

Gavrin A, Kaiser BN, Geiger D, Tyerman SD, Wen ZY, Bisseling T, Fedorova EE. 2014.
Adjustment of Host Cells for Accommodation of Symbiotic Bacteria: Vacuole
Defunctionalization, HOPS Suppression, and TIP1g Retargeting in *Medicago*. *Plant*
Cell **26**(9): 3809-3822.

Gibson AH, Nutman PS 1960. Studies on the physiology of nodule formation. VII. A
reappraisal of the effect of preplanting. *Annals of Botany* **24:** 420-433.

Guefrachi I, Nagymihaly M, Pislariu CI, de Velde WV, Ratet P, Mars M, Udvardi MK,
Kondorosi E, Mergaert P, Alunni B. 2014. Extreme specificity of NCR gene
expression in *Medicago truncatula*. *BMC Genomics* **15:712**.

Guerra J, Coussens G, De Keyser A, De Rycke R, De Bodt S, Van De Velde W,
Goormachtig S, Holsters M. 2010. Comparison of Developmental and Stress-
Induced Nodule Senescence in *Medicago truncatula*. *Plant Physiology* **152**(3): 1574-
1584.

Horvath B, Domonkos A, Kereszt A, Szucs A, Abraham E, Ayaydin F, Boka K, Chen

921   **YH, Chen RJ, Murray JD, et al. 2015.** Loss of the nodule-specific cysteine rich
922   peptide, NCR169, abolishes symbiotic nitrogen fixation in the *Medicago truncatula*
923   *dnf7* mutant. *Proc. Natl. Acad. Sci. USA* **112**(49): 15232-15237.
924 **Horvath B, Yeun LH, Domonkos A, Halasz G, Gobbato E, Ayaydin F, Miro K, Hirsch S,**
925   **Sun JH, Tadege M, et al. 2011.** *Medicago truncatula* IPD3 Is a Member of the
926   Common Symbiotic Signaling Pathway Required for Rhizobial and Mycorrhizal
927   Symbioses. *Molecular Plant-Microbe Interactions* **24**(11): 1345-1358.
928 **Isozumi N, Masubuchi Y, Imamura T, Mori M, Koga H, Ohki S. 2021.** Structure and
929   antimicrobial activity of NCR169, a nodule-specific cysteine-rich peptide of *Medicago*
930   *truncatula*. *Scientific Reports* **11**(1).
931 **Jones KM, Kobayashi H, Davies BW, Taga ME, Walker GC. 2007.** How rhizobial
932   symbionts invade plants: the *Sinorhizobium-Medicago* model. *Nature Reviews*
933   *Microbiology* **5**(8): 619-633.
934 **Kim M, Chen YH, Xi JJ, Waters C, Chen RJ, Wang D. 2015.** An antimicrobial peptide
935   essential for bacterial survival in the nitrogen-fixing symbiosis. *Proc. Natl. Acad. Sci.*
936   *USA* **112**(49): 15238-15243.
937 **Maunoury N, Redondo-Nieto M, Bourcy M, Van de Velde W, Alunni B, Laporte P,**
938   **Durand P, Agier N, Marisa L, Vaubert D, et al. 2010.** Differentiation of Symbiotic
939   Cells and Endosymbionts in *Medicago truncatula* Nodulation Are Coupled to Two
940   Transcriptome-Switches. *PLOS ONE* **5**(3): e9519.
941 **Mergaert P. 2018.** Role of antimicrobial peptides in controlling symbiotic bacterial
942   populations. *Natural Product Reports* **35**(4): 336-356.
943 **Mergaert P, Nikovics K, Kelemen Z, Maunoury N, Vaubert D, Kondorosi A, Kondorosi**
944   **E. 2003.** A novel family in *Medicago truncatula* consisting of more than 300 nodule-
945   specific genes coding for small, secreted polypeptides with conserved cysteine motifs.
946   *Plant Physiology* **132**(1): 161-173.
947 **Mergaert P, Uchiumi T, Alunni B, Evanno G, Cheron A, Catrice O, Mausset AE,**
948   **Barloy-Hubler F, Galibert F, Kondorosi A, et al. 2006.** Eukaryotic control on
949   bacterial cell cycle and differentiation in the Rhizobium-legume symbiosis. *Proc. Natl.*
950   *Acad. Sci. USA* **103**(13): 5230-5235.
951 **Montiel J, Downie JA, Farkas A, Bihari P, Herczeg R, Balint B, Mergaert P, Kereszt A,**
952   **Kondorosi E. 2017.** Morphotype of bacteroids in different legumes correlates with the
953   number and type of symbiotic NCR peptides. *Proc. Natl. Acad. Sci. USA* **114**(19):
954   5041-5046.
955 **Morandi D, Prado E, Sagan M, Duc G. 2005.** Characterisation of new symbiotic *Medicago*
956   *truncatula* (Gaertn.) mutants, and phenotypic or genotypic complementary
957   information on previously described mutants. *Mycorrhiza*: **15**: 283-289.
958 **Nagymihaly M, Veluchamy A, Gyorgypal Z, Ariel F, Jegu T, Benhamed M, Szucs A,**
959   **Kereszt A, Mergaert P, Kondorosi E. 2017.** Ploidy-dependent changes in the
960   epigenome of symbiotic cells correlate with specific patterns of gene expression. *Proc.*
961   *Natl. Acad. Sci. USA* **114**(17): 4543-4548.
962 **Nallu S, Silverstein KAT, Samac DA, Bucciarelli B, Vance CP, VandenBosch KA. 2013.**
963   Regulatory Patterns of a Large Family of Defensin-Like Genes Expressed in Nodules
964   of *Medicago truncatula*. *Plos One* **8**: e60355.
965 **Oldroyd GED, Downie JA. 2008.** Coordinating nodule morphogenesis with rhizobial
966   infection in legumes. *Annual Review of Plant Biology* **59**: 519-546.
967 **Oono R, Denison RF. 2010.** Comparing Symbiotic Efficiency between Swollen versus
968   Nonswollen Rhizobial Bacteroids. *Plant Physiology* **154**(3): 1541-1548.
969 **Ovchinnikova E, Journet EP, Chabaud M, Cosson V, Ratet P, Duc G, Fedorova E, Liu**
970   **W, den Camp RO, Zhukov V, et al. 2011.** IPD3 Controls the Formation of Nitrogen-

Fixing Symbiosomes in Pea and Medicago Spp. *Mol. Plant-Micr. Interact.* **24**(11):
1333-1344.
Pan HR, Wang D. 2017. Nodule cysteine-rich peptides maintain a working balance during
nitrogen-fixing symbiosis. *Nature Plants* **3**(5).
Ribeiro CW, Baldacci-Cresp F, Pierre O, Larousse M, Benyamina S, Lambert A,
Hopkins J, Castella C, Cazareth J, Alloing G, et al. 2017. Regulation of
Differentiation of Nitrogen-Fixing Bacteria by Microsymbiont Targeting of Plant
Thioredoxin s1. *Current Biology* **27**(2): 250-256.
Roux B, Rodde N, Jardinaud MF, Timmers T, Sauviac L, Cottret L, Carrere S, Sallet E,
Courcelle E, Moreau S, et al. 2014. An integrated analysis of plant and bacterial
gene expression in symbiotic root nodules using laser-capture microdissection coupled
to RNA sequencing. *Plant Journal* **77**(6): 817-837.
Roy P, Achom M, Wilkinson H, Lagunas B, Gifford ML. 2020. Symbiotic Outcome
Modified by the Diversification from 7 to over 700 Nodule-Specific Cysteine-Rich
Peptides. *Genes* **11**(4).
Sagan M, Morandi D, Tarenghi E, Duc G. 1995. Selection of nodulation and mycorrhizal
mutants in the model plant *Medicago truncatula* (Gaertn) after gamma-ray
mutagenesis. *Plant Science* **111**(1): 63-71.
Sankari S, Babu VMP, Bian K, Alhhazmi A, Andorfer MC, Avalos DM, Smith TA,
Yoon K, Drennan CL, Yaffe MB, et al. 2022. A haem-sequestering plant peptide
promotes iron uptake in symbiotic bacteria. *Nature Microbiology* **7**(9): 1453-+.
Starker CG, Parra-Colmenares AL, Smith L, Mitra RM, Long SR. 2006. Nitrogen
fixation mutants of *Medicago truncatula* fail to support plant and bacterial symbiotic
gene expression. *Plant Physiology* **140**(2): 671-680.
Terpolilli J, O'Hara G, Tiwari R, Dilworth M, Howieson J. 2008. The model legume
Medicago truncatula A17 is poorly matched for N-2 fixation with the sequenced
microsymbiont *Sinorhizobium meliloti* 1021. *New Phytologist* **179**(1): 62-66.
Van de Velde W, Zehirov G, Szatmari A, Debreczeny M, Ishihara H, Kevei Z, Farkas A,
Mikulass K, Nagy A, Tiricz H, et al. 2010. Plant Peptides Govern Terminal
Differentiation of Bacteria in Symbiosis. *Science* **327**(5969): 1122-1126.
Vasse J, de Billy F, Camut S, Truchet G. 1990. Correlation between ultrastructural
differentiation of bacteroids and nitrogen fixation in alfalfa nodules. *J Bacteriology*
172(8): 4295-4306.
Velivelli SLS, Czymmek KJ, Li H, Shaw JB, Buchko GW, Shah DM. 2020. Antifungal
symbiotic peptide NCR044 exhibits unique structure and multifaceted mechanisms of
action that confer plant protection. *Proc. Natl. Acad. Sci. USA* **117**(27): 16043-16054.
Wang C, Yu HX, Luo L, Duan LJ, Cai LY, He XX, Wen JQ, Mysore KS, Li GL, Xiao
AF, et al. 2016. NODULES WITH ACTIVATED DEFENSE 1 is required for
maintenance of rhizobial endosymbiosis in Medicago truncatula. *New Phytologist*
212(1): 176-191.
Wang D, Griffitts J, Starker C, Fedorova E, Limpens E, Ivanov S, Bisseling T, Long S.
2010. A Nodule-Specific Protein Secretory Pathway Required for Nitrogen-Fixing
Symbiosis. *Science* **327**(5969): 1126-1129.
Wang Q, Yang SM, Liu JG, Terecskei K, Abraham E, Gombar A, Domonkos A, Szucs
A, Kormoczi P, Wang T, et al. 2017. Host-secreted antimicrobial peptide enforces
symbiotic selectivity in *Medicago truncatula. Proc. Natl. Acad. Sci. USA* **114**(26):
6854-6859.
Wei F, Liu Y, Zhou DL, Zhao WL, Chen ZN, Chen DS, Li YG, Zhang XX. 2022.
Transcriptomic Identification of a Unique Set of Nodule-Specific Cysteine-Rich
Peptides Expressed in the Nitrogen-Fixing Root Nodule of *Astragalus sinicus.*

Molecular Plant-Microbe Interactions **35**(10): 893-905.

Xi JJ, Chen YH, Nakashima J, Wang SM, Chen RJ. 2013. *Medicago truncatula esn1*
Defines a Genetic Locus Involved in Nodule Senescence and Symbiotic Nitrogen
Fixation. *Molecular Plant-Microbe Interactions* **26**(8): 893-902.

Yang SM, Wang Q, Fedorova E, Liu JG, Qin QL, Zheng QL, Price PA, Pan HR, Wang
D, Griffitts JS, et al. 2017. Microsymbiont discrimination mediated by a host-
secreted peptide in *Medicago truncatula*. *Proc. Natl. Acad. Sci. USA* **114**(26): 6848-
6853.

Zorin EA, Kliukova MS, Afonin AM, Gribchenko ES, Gordon ML, Sulima AS,
Zhernakov AI, Kulaeva OA, Romanyuk DA, Kusakin PG, et al. 2022. A variable
gene family encoding nodule-specific cysteine-rich peptides in pea (*Pisum sativum*
L.). *Frontiers in Plant Science* **13**: 884726.

1033

1034

1035

1036 **Figure legends**

1037

Fig. 1 The ineffective nodulation phenotype of *M. truncatula* symbiotic mutants *Mtdnf7-2*,
Mtdnf4-1, NF-FN9363, *Mtsym20* and *Mtsym19*. (a) Ineffective nitrogen-fixing (Fix-)
symbiotic mutants showed the symptoms of the nitrogen starvation (retarded growth, pale
green leaves) compared with wild-type (wt) plant two weeks following inoculation with strain
Sinorhizobium medicae WSM419 carrying the *lacZ* reporter gene. (b) Slightly elongated
white nodules were formed on the roots of the Fix- symbiotic mutant plants and cylindrical
pink nodules were observed on wt roots 14 days post inoculation (dpi) with *S. medicae*
WSM419 (upper images). Longitudinal nodule sections stained for β-galactosidase activity
show the normal colonization of nodule zones characteristic for wt indeterminate nodules but
in ZIII of Fix- nodules are devoid of rhizobia infected cells (lower images).

The zones were labelled with square brackets and the IZ was highlighted in magenta. ZIId,
distal part of infection zone; ZIIp, proximal part of infection zone; IZ, interzone; ZIII,
nitrogen fixation zone; Bars, (a) 2 cm; (b) 200 µm.

1051

Fig. 2 Laser confocal microscopic analyses of symbiotic cell structure and bacteroid
morphology on SYTO13 stained nodule sections of the Fix- mutants NF-FN9363, *Mtsym19*
and *Mtsym20* compared with nodules of wt and *Mtdnf7-2*, *Mtdnf4-1* plants 14 days post
inoculation with *S. medicae* WSM419. (a1-f1) SYTO13-stained longitudinal nodule sections
show the defect of bacterial invasion in ZIII of ineffective symbiotic nodules. The green
channel visualizes SYTO13-stained bacteroids and plant cell nuclei, autofluorescence is
pseudocolored in red. Higher magnification shows the transition between ZII and IZ (a2-f2)

and IZ-ZIII (a5-f5). (a3-f3; a4-f4) No obvious alteration was observed in bacterial occupation
of infected cells or the morphology of bacteroids in distal part of ZII (ZIId) between wt and
mutant nodule cells. (a6-f6; b7-f7) Moderate elongation of bacteroids was found in the
proximal part of IZ (IZp) cells of mutant nodules compared with wt nodule cells. This region
is composed of a single cell layer in wt nodules but it was extended in mutant nodules. Mutant
infected cells in IZp contain disordered bacteroids indicating the inception of their
disintegration. (a8-f8) The cells in ZIII but its first cell layer (IZd) in mutant nodules (b7-f7)
were devoid of bacteroids and contained autofluoresced speckles indicating the presence of
early senescence-related phenolic compounds. Bars, (a1-f1) 200 µm; (a2-f8) 20 µm. it,
infection thread; b, bacteroid; sb, saprophytic bacteria; v, vacuole; n, nucleus; ui, uninfected
cells.

1070

Fig. 3 Bacteroid differentiation is similarly impaired in NF-FN9363 and *Mtsym20* mutant
nodules as in the formerly identified ineffective symbiotic mutants *Mtdnf4-1* and *Mtdnf7-2* at
2 weeks post inoculation with *S. medicae* WSM419. (a) The distribution of size of bacteroids
isolated from wt and mutant nodules at 2 wpi, stained with PI, captured by confocal laser
scanning microscopy and measured on the images using the ImageJ program. The size of
bacteroids from mutant nodules had a higher proportion of shorter bacterial cells compared
with wild-types and rhizobia above 6 µm in length were hardly observed in mutant nodules
although these classes in wt nodules were evident. The size of at least 1200 bacterial cells was
measured and their relative distribution in length classes is presented. Values around the radar
chart indicate bacterial length (µm) and circles show percentage (%). (b) The relative size and
DNA content of *S. medicae* WSM419 bacteroids isolated from wt and mutant nodules at 16
dpi measured by flow-cytometry. Bacteroid relative size was determined by the forward light
scatter (FSC) and endoreduplication was measured by fluorescence intensity. The size and
DNA content of *S. medicae* WSM419 bacteroid populations isolated from mutant nodules
shifted to smaller size and lower ploidy level compared with bacteroids found in wt nodules.

1086

Fig. 4 The identified deleted regions and the gene models in the symbiotic loci of NF-FN9363
and *Mtsym19*, *Mtsym20* mutants. (a) Genetic mapping and high-density genome array-based
comparative genomic hybridization (aCGH) identified a nearly 500 kb deletion including four
NCR genes in the NF-FN9363 genome. (b) The extension of the deleted regions in the mutant
loci of *Mtsym19* and *Mtsym20* was determined by PCR based markers and by analysing reads
obtained from RNAseq. The structure of the *NCR343* (a) and *NCR-new35* (b) genes (coding

sequences, the intron of *NCR-new35* and 5' UTRs and 3' UTRs) are represented by white

boxes, thin excised black line and thick black lines, respectively) and the features of the

encoded NCR343 (a) and NCR-new35 (b) peptides containing a signal peptide and mature

peptides with four cysteine residues in conserved positions. (c) Multiple sequence alignment

and phylogenetic analysis of mature peptides encoded by the *NCR343, NCR341, NCR345* and

NCR344 genes deleted in the symbiotic locus of NF-FN9363 and (d) the sequence analysis of

mature NCR peptides proved to be crucial for differentiation and persistence of rhizobia

during nitrogen-fixing symbiosis. The degree of sequence conservation of the residues

between the four mature peptides is represented below the alignment. Colours are based on

Robert Fletterick's "Shapely models."

1103

Fig. 5 *NCR343* and *NCR-new35* are required for development of functional nodules and the

first cysteine residues of NCR343 and NCR-new35 peptides are essential for their function.

The rescue of the nodulation defect of mutants NF-FN9363, *Mtsym19* and *Mtsym20* was

identified based on the restoration of morphology and colonization of nodules formed on

hairy roots transformed with empty vector or *NCR* peptide genes and inoculated with *S.*

medicae WSM419 carrying the *lacZ* marker gene. Longitudinal sections of 4-week-old

nodules were stained for β-galactosidase activity. The nodules on NF-FN9363 roots

transformed with *NCR343* (a), on *Mtsym19* and *Mtsym20* roots transformed with *NCR-new35*

(b), on wt roots transformed with an empty vector (c) and on *Mtdnf4-1* roots transformed with

NCR211 genes (d) showed characteristic zonation of indeterminate nodules indicating the

restoration of the symbiotic phenotypes. Gene constructs of *NCR345, NCR341*, and *NCR344*

did not restore the symbiotic phenotype of nodules developed on transformed roots of NF-

FN9363 indicating that these peptides are not essential for the symbiotic interaction between

M. truncatula A17 and *S. medicae* WSM419. The substitution of the first cysteine residue for

serine abolished complementation in the mutant NCR343 (a), NCR-new35 (b) and NCR211

(d) genes indicating the essential role of the first cysteines. Bars, (a–d), 200 μm.

1120

Fig. 6 The expression analysis of the *NCR-new35* and *NCR343* genes is compared with the

activity of *NCR211* and *NCR169*. The empty vector (a and f) and the construct of promoter

fragments of *NCR-new35* (b and g), *NCR211* (c and h), *NCR343* (d and i) and *NCR169* (e and

j*)* genes fused to the β-glucoronidase (GUS) gene, respectively were introduced into the roots

of wt Jemalong plants with *A. rhizogenes*-mediated hairy-root transformation. Nodule

sections were stained for GUS activity at two (a-e) and three (f-j) wpi with *S. medicae*

WSM419. The relative spatial expression of *NCR-new35*, *NCR211*, *NCR343* and *NCR169* genes 2 wpi with *S. medicae* WSM419 generated with RNA sequencing of different nodule zones obtained with laser-capture microdissection (LCM) are shown below the images of 2-week-old nodules. (k) Relative expression of *NCR169*, *NCR211* and *NCR343* compared with *NCR-new35* was analysed in wild-type nodules by reverse transcription quantitative polymerase chain reaction (RT-qPCR) 2 and 3 wpi with rhizobia. Magenta square brackets, IZ; black arrowhead, ZIIp. Bars, (a-j) 200 µm.

Fig. 7 NCR-new35-mCherry and NCR343-mCherry fusion proteins restored the symbiotic defects of *Mtsym20* (a) and NF-FN9363 (d) mutant plants, respectively at 4 wpi with *S. medicae* WSM419. (a and d) The red NCR-new35-mCherry and NCR343-mCherry signals overlap with the green fluorescence of SYTO13-stained bacteroids in the infected cells of interzones (magenta brackets) and nitrogen fixation zones indicating the co-localization of bacteroids and peptides NCR-new35 and NCR343. (b, c, and e, f) Higher magnification of cells from the IZ revealed the red mCherry signal of NCR-new35 and NCR343 fusion proteins surrounding the bacteroids in the peribacteroid space. Bars, (a, d) 200 µm; (b, c, e and f) 20 µm.

Fig. S1 Strain dependence of the nodulation and symbiotic phenotype of four ineffective symbiotic mutants NF-FN9363, *Mtsym20, Mtdnf4-1* and *Mtdnf7-2* compared with wt *M. truncatula* Jemalong nodules. (a) Plants were inoculated with *S. meliloti* strains 1021 and FSM-MA as well as with *S. medicae* strains ABS7 and WSM419 (*S. medicae* WSM419) and then nodule images were captured 3 wpi. Pink nodules were formed on wt roots with the four rhizobia strains. Roundish or slightly elongated white nodules developed on *Mtdnf7-2*, *Mtdnf4-1*, NF-FN9363 and *Mtsym20* roots, irrespective of the bacterial strains used for inoculation. Scale bars: 200µm. (b) Box-plot of nodule number and dry weight of wt and four ineffective symbiotic nitrogen-fixing mutants 3 weeks post inoculation with different rhizobia strains.

Fig. S2 Transcriptional activity of senescence and defense-related marker genes in the nodules of four ineffective symbiotic mutants NF-FN9363, *Mtsym20*, *Mtdnf4-1, Mtdnf7-2* and *nad1-3* compared with wt nodules 3 wpi with *S. medicae* WSM419 using RT-qPCR.

1161 Senescence marker genes *MtCP2 and MtCP6* (a) were activated in all mutant nodules but
1162 pathogen-related genes *MtCHI* (b) and *MtPR10* (c) were highly activated only in *nad1-3*
1163 nodules.

1164

Fig. S3 Scanning electron microscopy analysis of symbiotic cells and bacteroid morphology
in NF-FN9363, *Mtsym20*, *Mtdnf7-2*, *Mtdnf4-1* mutant and wt nodules 2 wpi with *S. medicae*
WSM419. SEM images of whole nodule sections of wt plant (a1), mutant lines (b1-e1) free
living *S. medicae* WSM419 bacteria (f1). Higher magnification of symbiotic cells
characteristic for the distal part of infection zone (ZIId, a2-e2), the proximal part of infection
zone (ZIIp, a3-e3), the first few cell layer of interzone (IZd, a4-e4), the proximal part of IZ of
mutant nodules (IZp, b5-e5) and the mature nitrogen fixation zone (ZIII, a6-e6). An enlarged
part of the symbiotic cells showing bacterial morphology are below each images.
Scale bars: 100 µm in (a1-e1), 10 µm in (a2-e6). Scale bars in the images of plant cells and
bacteroids at higher magnification (f1, a2-e6) 1 µm. it: infection thread b: bacteroid, db:
decomposing bacteria, sb: saprophytic bacteria, v: vacuole, n: nucleus, s: starch granules *:
discontinuity between rhizobia and plant cell walls.

1177

Fig. S4 Chromosomal position of the symbiotic loci of NF-FN9363, *Mtsym19* and *Mtsym20*
and the allelism test between *Mtsym19* and *Mtsym20*. (a) The symbiotic locus of mutant NF-
FN9363 was mapped in Linkage Group 6 (LG 6) of *M. truncatula* downwards the genetic
marker Crs towards MtB178. Genetic mapping defined the chromosomal position of the
symbiotic loci of *Mtsym20* and *Mtsym19* on chromosome 4 (LG 4). (b) The allelic
relationship of *Mtsym20* and *Mtsym19* mutants was proved with allelism tests. F3 *Mtsym20*
and *Mtsym19* symbiotic mutant plants, selected from the mapping populations showing
opposite parental homozygous genotypes for the genetic marker MtB267, were crossed and
the offsprings were scored for the symbiotic phenotype. The success of the crossing of the
two mutant lines was detected with the heterozygous genotypes of C1, C3 and C4 hybrid
plants. Plants of C2 and C5 were self-mated. The Fix- symbiotic phenotype of plants of C1,
C3 and C4 crosses revealed the allelic relationship of mutants *Mtsym20* and *Mtsym19*; MM,
Molecular weight Marker

1191

Fig. S5 The deleted genomic region in the symbiotic locus of NF-FN9363. (a) An
approximately 500 kb genomic region deleted in the symbiotic locus in NF-FN9363 was
defined with PCR-based markers designed for genes. The A17 genomic DNA was used as a

wild type control in the PCR reactions. * primer dimers. M, mutant-NF-FN9363; MM,
Molecular weight Marker (b) List of genes or gene models in the deleted genomic region
detected by using microarray-based CGH. Dark grey shading shows genes corresponding to
probesets with log2 CGH ratios lower than -2,5, pale grey shading is for $\log_2$ CGH ratios less
than -1 and greater than -2,5. (+) and (-) indicate the presence or absence of PCR fragments
identified with the genomic DNA of NF-FN9363. Deleted *NCR* genes in the symbiotic locus
of NF-FN9363 are in bold.
1202
Fig. S6 The deleted genomic region of the symbiotic loci of *Mtsym19* and *Mtsym20*. (a) PCR-
based markers identified overlapping deleted genomic regions in *Mtsym19* and *Mtsym20* on
chromosome 4 (Chr 4). PCR-based markers were designed for genes marked with gene IDs
and visualized on agarose gel. The *M. truncatula* Jemalong (J5) genomic DNA was used for a
wild-type control. (b) List of genes and gene models in the deleted genomic region. (+) not
deleted: (-) deleted genes in the symbiotic loci of *Mtsym19* and *Mtsym20* detected with PCR-
based markers. MM, Molecular weight Marker
1210
Fig. S7 Complementation assays of NF-FN9363, *Mtsym20* and *Mtsym19* symbiotic
ineffective mutants with *NCR* gene constructs and the study of the requirement of first
cysteine residues in the function for the peptides NCR343, NCR-new35 and NCR211.
Constructs of *NCR* genes controlled by their native promoter fragments were introduced into
the roots of mutant plants using *A. rhizogenes*-mediated transformation. The transgenic roots
were detected either by dsRed fluorescence, in the case of gene constructs created in single
site gateway (GW) system (a1-i1), or with GFP fluorescence using the constructs generated
by multi-site GW system (j1-t1). Complementation was assessed by the formation of
cylindrical nodules with the characteristic zonation of indeterminate nodules. To compare
zonation of nodules, nodules developed on the wild-type (a1-a3 and j1-j3) and symbiotic
mutant roots (*Mtsym20* - c1-c2 and p1-p3, *Mtsym19* - e1-e2, NF-FN9363 - g1-g3 and l1-l3,
and *Mtdnf4-1* -s1-s3) transformed with empty vector were used. The symbiotic defects of both
mutants *Mtsym20* (b1-b3 and o1-o3) and *Mtsym19* (d1-d3) were rescued with the *NCR-new35*
gene constructs indicating that they are deficient in the same genes. The *NCR343* gene
constructs (f1-f3 and k1-k3) restored but the constructs of *NCR345* (h1-h3), *NCR341* (i1-i3)
and *NCR344* (m1-m3) did not complement the nodulation phenotype of NF-FN-9363. The
modification of the constructs substituting the first cysteines for serines equally abolished the
ability of NCR343-C34S (n1-n3), NCR-new35-C41S (q1-q3), NCR211-C28S (t1-t3) peptides

to restore the symbiotic nodulation phenotype of mutants NF-FN9363, *Mtsym20* and *Mtdnf4-*

1, respectively. Construct containing the wild-type *NCR211* gene restored the symbiotic

nodulation phenotype on *Mtdnf4-1* roots (r1-r3). Bars, (a1-t3) 200 μm.

(a)

Bacteroids length categories (1-12 µm)

Mtsym20 —— *Mtdnf4-1* —— *Mtdnf7-2* —— NF-FN9363 —— wt

(b)

cell size (FSC)

DNA content

Mtsym20
wt
free living WSM

NF-FN9363
wt
free living WSM

Mtdnf4-1
wt
free living WSM

(b)

RNA seq	Total Read Count					
Mtsym18	1032	2282	153	25	39	576
Mtsym20	278		2	0	76	49	854
Mtsym19	179		0	0	0	0	765

	ev	*NCR-new35*	*NCR211*	*NCR343*	*NCR169*
2 wpi	(a)	(b)	(c)	(d)	(e)

(k) Relative expression of *NCRs 343, 211* and *169* compared with *NCR-new35* at 2 wpi

RNA seq at 2 wpi

ZIId%	ZIIp%	IZ%	ZIII%
0,0	7,6	86,1	6,3
total reads LCM		10196,72721	

ZIId%	ZIIp%	IZ%	ZIII%
0,1	22,8	76,0	1,1
total reads LCM		309850,3989	

ZIId%	ZIIp%	IZ%	ZIII%
0,1	1,9	75,3	22,7
total reads LCM		63074,62211	

ZIId%	ZIIp%	IZ%	ZIII%
0,0	2,0	67,9	30,0
total reads LCM		688160,173	

3 wpi	(f)	(g)	(h)	(i)	(j)

Relative expression of *NCRs 343, 211* and *169* compared with *NCR-new35* at 3 wpi

(a) NCR-new35 -mCHERRY+SYTO13

Mtsym20

(b) mCHERRY

(c) mCHERRY+SYTO13

(d) NCR343 -mCHERRY+SYTO13

NF-FN9363

(e) mCHERRY

(f) mCHERRY+SYTO13