The pGinger family of expression plasmids

Allison N. Pearson[1,2,3*], Mitchell G. Thompson[1,4*], Liam D. Kirkpatrick[1,4], Cindy Ho[1,2], Khanh M. Vuu[1,4], Lucas M. Waldburger[1,2,5], Jay D. Keasling[1,2,6,7,8†], Patrick M. Shih[1,3,4,9†]

[1]Joint BioEnergy Institute, 5885 Hollis Street, Emeryville, CA 94608, USA.
[2]Biological Systems & Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA.
[3]Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720, USA
[4]Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory, Berkeley, California, USA
[5]Department of Bioengineering, University of California, Berkeley, California, USA
[6]Department of Chemical and Biomolecular Engineering, University of California, Berkeley, CA 94720, USA
[7]The Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, Denmark
[8]Center for Synthetic Biochemistry, Institute for Synthetic Biology, Shenzhen Institutes for Advanced Technologies, Shenzhen, China
[9]Innovative Genomics Institute, University of California, Berkeley, CA

*Allison N. Pearson and Mitchell G. Thompson contributed equally to this manuscript

†Correspondence should be addressed to either Jay D. Keasling (jdkeasling@lbl.gov) or Patrick M. Shih (pmshih@lbl.gov)

Abstract

The pGinger suite of expression plasmids comprises 43 plasmids that will enable precise constitutive and inducible gene expression in a wide range of gram-negative bacterial species. Constitutive vectors are composed of 16 synthetic constitutive promoters upstream of RFP, with a broad host range BBR1 origin and a kanamycin resistance marker. The family also has seven inducible promoters (Jungle Express, NahR, XylS, RhaS, Lac, LacUV5, and TetR) controlling RFP expression on BBR1/kanamycin plasmid backbones. For four of these inducible promoters (Jungle Express, NahR, Lac, and TetR), we created variants that utilize the RK2 origin and spectinomycin or gentamicin selection. Relevant RFP expression and growth data have been

collected in the model bacterium *Escherichia coli* as well as *Pseudomonas putida*. All pGinger

vectors are available via the Joint BioEnergy Institute (JBEI) Public Registry.

Introduction

Precise and reliable control over gene expression is one of the most fundamental

requirements of synthetic biology (1). Consequently, there has been considerable effort towards

identifying myriad genetic elements that enable researchers to regulate the strength and timing of

transcription across all domains of life (2–4). The end result of these efforts are often

consolidated families of plasmid vectors that facilitate advanced genetic engineering, such as the

BglBrick family of plasmids for *E. coli* (5, 6) and the jStack vectors used in multiple plant

species (7). However, as the field of synthetic biology moves beyond traditional model

organisms, families of expression vectors must be tailored to meet the specific requirements of

particular hosts. Advances in non-model organisms often come in the form of species or genus

specific toolkits (8, 9), though more recently comprehensive plasmid toolkits have been

developed and validated for a wide range of gram-negative organisms (10). Still, given that many

bacteria require very particular combinations of promoters, origins, and selectable markers to

enable controlled gene expression, there remains a need for vectors that will allow rapid

prototyping of genetic circuits in understudied bacteria.

To facilitate the exploration of non-model hosts, we have developed a small suite of

plasmids that permit both constitutive and inducible expression from the broad host-range origin

of replication BBR1 using a kanamycin selection marker. For a subset of the inducible promoters

that are known to work across multiple hosts, we have assembled combinatorial variants that

utilize the compatible broad host-range origin RK2 (11) as well as both spectinomycin and

gentamicin selection markers. This family of plasmids, which we have named the pGinger suite,

requires no assembly of these parts, can be easily cloned into via standard Gibson assembly

techniques, and has both digital sequences and physical samples that can be publicly accessed

through the Joint BioEnergy Institute (JBEI) registry (12).

Results

Design and Architecture of pGinger Plasmids

All pGinger vectors express RFP with a consensus ribosomal binding site (RBS -

TTTAAGAAGGAGATATACAT) derived from the BglBrick plasmid library. The overall

conserved plasmid architecture and naming convention of the pGinger suite are shown in **Figure**

1.

Figure 1: **Plasmids architecture of the pGinger suite**: The pGinger plasmids share a common naming convention

where the first two letters after pGinger correspond to the origin and resistance marker respectively, followed by the

promoter. All plasmids share the same architecture as the above map of pGingerBK-TetR, whereby a conserved

RBS-RFP is downstream of the promoter followed by a strong terminator. All selectable markers are upstream of

the promoter, with the origin between the marker and the RFP cassette.

The BBR1 origin and kanamycin cassette of relevant pGinger vectors were both derived from

plasmid pBADTrfp (13). To develop a family of constitutive expression plasmids, the AraC

coding sequence and promoter of pBADTrfp were replaced with 16 different synthetic promoters

from the Anderson Promoter Library (http://parts.igem.org/Promoters/Catalog/Anderson). For

the inducible vectors, the AraC coding sequence and promoter of pBADTrfp were replaced with

the following seven inducible systems: Jungle Express - derived from pTR_sJExD-rfp (14);

NahR - derived from pPS43 (15); RhaS - derived from pCV203 (15); TetR - derived from

pBbE2a-RFP; XylS - derived from pPS66 (15); Lac - derived from pBbE6a-RFP (6); LacUV5 -

derived from pBbE5a-RFP (6). For four of the inducible promoters (Jungle Express, NahR, Lac,

and TetR), additional vectors were constructed that varied both the origin and antibiotic marker.

All RK2 origins were derived from pBb(RK2)1k-GFPuv (8), while the gentamicin resistance

cassette was derived from pMQ30 (16), and the spectinomycin cassette was derived from

pSR43.6 (17). All BBR1 derivatives of pBADTrfp contain the *mob* element that facilitates

conjugal transfer. A full description of each pGinger vector can be found in **Table 1**.

Name	Origin	Marker	Promoter Class	Promoter	Inducer	JBEI ICE No.
pGingerBK-J23100	BBR1	Kanamycin	Constitutive	J23100	NA	JPUB_020797
pGingerBK-J23101	BBR1	Kanamycin	Constitutive	J23101	NA	JPUB_020799
pGingerBK-J23102	BBR1	Kanamycin	Constitutive	J23102	NA	JPUB_020815
pGingerBK-J23103	BBR1	Kanamycin	Constitutive	J23103	NA	JPUB_020801
pGingerBK-J23104	BBR1	Kanamycin	Constitutive	J23104	NA	JPUB_020803
pGingerBK-J23105	BBR1	Kanamycin	Constitutive	J23105	NA	JPUB_020817
pGingerBK-J23106	BBR1	Kanamycin	Constitutive	J23106	NA	JPUB_020793
pGingerBK-J23107	BBR1	Kanamycin	Constitutive	J23107	NA	JPUB_020819
pGingerBK-J23108	BBR1	Kanamycin	Constitutive	J23108	NA	JPUB_020821

pGingerBK-J23110	BBR1	Kanamycin	Constitutive	J23110	NA	JPUB_020805
pGingerBK-J23111	BBR1	Kanamycin	Constitutive	J23111	NA	JPUB_020807
pGingerBK-J23113	BBR1	Kanamycin	Constitutive	J23113	NA	JPUB_020809
pGingerBK-J23114	BBR1	Kanamycin	Constitutive	J13114	NA	JPUB_020811
pGingerBK-J23117	BBR1	Kanamycin	Constitutive	J13117	NA	JPUB_020795
pGingerBK-J23118	BBR1	Kanamycin	Constitutive	J23118	NA	JPUB_020823
pGingerBK-J23119	BBR1	Kanamycin	Constitutive	J23119	NA	JPUB_020813
pGingerBK-JE	BBR1	Kanamycin	Inducible	Jungle Express	Crystal Violet	JPUB_020825
pGingerBK-NahR	BBR1	Kanamycin	Inducible	NahR	Salicylic acid	JPUB_020831
pGingerBK-RhaS	BBR1	Kanamycin	Inducible	RhaS	Rhamnose	JPUB_020829
pGingerBK-TetR	BBR1	Kanamycin	Inducible	TetR	Oxytetracycline	JPUB_020835
pGingerBK-XylS	BBR1	Kanamycin	Inducible	XylS	Benzoate	JPUB_020827
pGingerBK-Lac	BBR1	Kanamycin	Inducible	Lac	IPTG	JPUB_020833
pGingerBK-LacUV5	BBR1	Kanamycin	Inducible	LacUV5	IPTG	JPUB_020837
pGingerBG-JE	BBR1	Gentamicin	Inducible	Jungle Express	Crystal Violet	JPUB_020847
pGingerBS-JE	BBR1	Spectinomycin	Inducible	Jungle Express	Crystal Violet	JPUB_020855
pGingerRK-JE	RK2	Kanamycin	Inducible	Jungle Express	Crystal Violet	JPUB_020871
pGingerRG-JE	RK2	Gentamicin	Inducible	Jungle Express	Crystal Violet	JPUB_020881
pGingerRS-JE	RK2	Spectinomycin	Inducible	Jungle Express	Crystal Violet	JPUB_020863
pGingerBG-NahR	BBR1	Gentamicin	Inducible	NahR	Salicylic acid	JPUB_020845
pGingerBS-NahR	BBR1	Spectinomycin	Inducible	NahR	Salicylic acid	JPUB_020853
pGingerRK-NahR	RK2	Kanamycin	Inducible	NahR	Salicylic acid	JPUB_020869
pGingerRG-NahR	RK2	Gentamicin	Inducible	NahR	Salicylic acid	JPUB_020879
pGingerRS-NahR	RK2	Spectinomycin	Inducible	NahR	Salicylic acid	JPUB_020859
pGingerBG-TetR	BBR1	Gentamicin	Inducible	TetR	Oxytetracycline	JPUB_020843
pGingerBS-TetR	BBR1	Spectinomycin	Inducible	TetR	Oxytetracycline	JPUB_020851
pGingerRK-TetR	RK2	Kanamycin	Inducible	TetR	Oxytetracycline	JPUB_020865
pGingerRG-TetR	RK2	Gentamicin	Inducible	TetR	Oxytetracycline	JPUB_020877
pGingerRS-TetR	RK2	Spectinomycin	Inducible	TetR	Oxytetracycline	JPUB_020861
pGingerBG-Lac	BBR1	Gentamicin	Inducible	Lac	IPTG	JPUB_020841
pGingerBS-Lac	BBR1	Spectinomycin	Inducible	Lac	IPTG	JPUB_020849

pGingerRK-Lac	RK2	Kanamycin	Inducible	Lac	IPTG	JPUB_020867
pGingerRG-Lac	RK2	Gentamicin	Inducible	Lac	IPTG	JPUB_020875
pGingerRS-Lac	RK2	Spectinomycin	Inducible	Lac	IPTG	JPUB_020857

Table 1: **Plasmids in the pGinger suite**: Relevant characteristics of pGinger plasmids including origin of

replication, antibiotic selection, promoter characteristics, and if applicable inducing molecule. JBEI public registry

numbers are also included for digital accessibility.

Evaluation of Constitutive Expression pGinger Plasmids

To evaluate the relative strength of constitutive Anderson promoters in the context of the

pGinger vectors, plasmids were introduced into both *P. putida* and *E. coli*. Fluorescence was

measured after growth in LB medium after 24 hours. When fluorescence was normalized to cell

density, expression from Anderson promoters showed significant correlation (Spearman's ρ =

0.49, p = 0.045) between *P. putida* and *E. coli* (**Figure 2**). Promoters J23103 and J23113 were

significantly stronger in *E. coli* than in *P. putida*, while promoter J23111 was significantly

stronger in *P. putida*. Promoter sequences and mean expression values in both *E. coli* and *P.*

putida are listed in **Table 2**.

Figure 2: Activity of Constitutive Promoters in *E. coli* and *P. putida*. RFP expression normalized to cell density

from Anderson promoters within either *E. coli* (y-axis) or *P. putida* (x-axis) are shown with standard deviations (

n=3).

Promoter	Promoter Sequence	*E. coli* expression	*P. putida* expression
J23100	ttgacggctagctcagtcctaggtacagtgctagc	13267 (+/- 517)	22343 (+/- 1262)
J23101	tttacagctagctcagtcctaggtattatgctagc	11530 (+/- 2565)	22010 (+/- 3162)
J23102	ttgacagctagctcagtcctaggtactgtgctagc	13300 (+/- 1815)	24067 (+/- 858)
J23103	ctgatagctagctcagtcctagggattatgctagc	18476 (+/- 1857)	565 (+/- 135)
J23104	ttgacagctagctcagtcctaggtattgtgctagc	18522 (+/- 682)	23440 (+/- 4588)
J23105	tttacggctagctcagtcctaggtactatgctagc	2622 (+/- 363)	10220 (+/- 558)

J23106	tttacggctagctcagtcctaggtatagtgctagc	5697 (+/- 369)	14659 (+/- 748)
J23107	tttacggctagctcagccctaggtattatgctagc	2798 (+/- 44)	7429 (+/- 716)
J23108	ctgacagctagctcagtcctaggtataatgctagc	1149 (+/- 84)	1523 (+/- 84)
J23110	tttacggctagctcagtcctaggtacaatgctagc	6402 (+/- 782)	12098 (+/- 2251)
J23111	ttgacggctagctcagtcctaggtatagtgctagc	1547 (+/- 106)	15548 (+/- 3229)
J23113	ctgatggctagctcagtcctagggattatgctagc	16220 (+/- 480)	826 (+/- 92)
J23114	tttatggctagctcagtcctaggtacaatgctagc	625 (+/- 48)	1325 (+/- 289)
J23117	ttgacagctagctcagtcctagggattgtgctagc	229 (+/- 25)	730 (+/- 28)
J23118	ttgacggctagctcagtcctaggtattgtgctagc	9628 (+/- 507)	21237 (+/- 5215)
J23119	ttgacagctagctcagtcctaggtataatgctagc	22157 (+/- 473)	14979 (+/- 3262)
WT	NA	124 (+/- 6)	141 (+/- 17)

Table 2: Expression of pGinger Anderson promoters: For each Anderson promoter the sequence is provided as

well as the mean cell density normalized RFP fluorescence in both *E. coli* and *P. putida*. Standard deviations are

provided in parentheses, n=3.

<u>Evaluation of Inducible pGinger Plasmids</u>

The expression of the seven inducible promoters within the pGinger suite was evaluated

using the BBR1 origin and kanamycin marker (pGingerBK) against a titration of the inducer in

both *E. coli* and *P. putida* (**Figure 3**). All promoters showed inducibility in *E. coli*, and all but

the rhamnose inducible RhaS promoter showed inducibility in *P. putida*. Relevant expression

characteristics of the inducible pGingerBK vectors in both tested bacteria are listed in **Table 3**.

The strongest normalized expression from an inducible promoter in *E. coli* was the TetR system,

while both the strongest promoters in *P. putida* were found to be NahR and Jungle Express

promoters, which showed nearly identical maximal expression. In both bacteria, the Jungle

Express promoter demonstrated the greatest level of induction relative to background expression.

Figure 3: Activity of Inducible Promoters in *E. coli* and *P. putida*. RFP expression normalized to cell density (y-

axis) from inducible promoters within either *E. coli* (top panel in orange) or *P. putida* (bottom panel in blue) as a

function of inducer concentration in mM (x-axis). Fits to the Hill equation are shown as dashed lines and shaded to

show confidence intervals. Raw data points are overlaid (n=3).

Promoter	Organism	Background	Max Exp.	Max Conc.	Induction
LacO	*E. coli*	1510 (+/- 186)	13127 (+/- 1693)	1 mM	9x
	P. putida	137 (+/- 5)	3074 (+/- 30)	5 mM	22x
NahR	*E. coli*	1260 (+/- 57)	16484 (+/- 1693)	500 uM	9x
	P. putida	847 (+/- 244)	25697 (+/- 2976)	5 mM	30x

LacUV5	*E. coli*	2889 (+/- 36)	10137 (+/- 855)	250 uM	4x
	P. putida	203 (+/- 53)	16622 (+/- 3671)	10 mM	82x
Jungle Express	*E. coli*	100 (+/- 2)	28545 (+/- 1890)	125 nM	285x
	P. putida	183 (+/- 1)	16956 (+/- 4002)	313 nM	93x
RhaS	*E. coli*	172 (+/- 42)	9251 (+/- 1389)	5 mM	54x
	P. putida	NA	NA	NA	NA
TetR	*E. coli*	341 (+/- 10)	33631 (+/- 7692)	1 uM	98x
	P. putida	176 (+/- 9)	3214 (+/- 319)	1 uM	18x
XylS	*E. coli*	329 (+/- 57)	15280 (+/- 1590)	313 uM	46x
	P. putida	161 (+/- 7)	8401 (+/- 1877)	5 mM	52x

Table 3: Inducible Promoters in *E. coli* and *P. putida*: For each inducible promoter on a BBR1 origin with a

kanamycin marker, the experimentally observed background (uninduced) fluorescence and maximal fluorescence

are given for both *E. coli* and *P. putida*. Standard deviations are provided in parentheses, n=3. Additionally, the

inducer concentration used to achieve maximal expression and the relative induction levels are listed.

To evaluate the effect of varying origin and selectable markers on expression from

inducible promoters, all six variants of the Jungle Express, LacO, NahR, and TetR were

investigated for their dose-response to their inducer molecules in *E. coli* (**Figure 4**). Relevant

expression parameters are listed in **Table 4**. In general, BBR1 variants showed greater

expression than RK2 origin plasmids, which is expected given the higher copy of BBR1

plasmids in *E. coli* (10). Amongst the pGinger Jungle Express vectors, both pGingerRS-JE and

pGingerRK-JE showed dose-responses distinct from the other vectors (**Figure 4A**). Notably, all

pGingerRS (RK2-Spectinomycin) plasmids showed the lowest expression across each promoter

tested (**Table 4**).

Figure 4: Activity of Inducible pGinger variants in _E. coli_. For origin and selection marker pGinger variants of

Jungle Express (A), LacO (B), NahR (C), and TetR (D), dose-response curves of normalized RFP expression are

shown as a function of mM inducer. Error bars represent standard deviations (n=3). Note that the x-axis is non-

linear.

Promoter	Origin	Marker	Background	Max Exp.	Max Conc.	Induction
LacO	BBR1	Kan	1510 (+/- 186)	13127 (+/- 1693)	1 mM	9x
LacO	BBR1	Gent	1976 (+/- 70)	14143 (+/- 1608)	63 uM	7x
LacO	BBR1	Spec	1830 (+/- 320)	10694 (+/- 943)	250 uM	6x

LacO	RK2	Kan	803 (+/- 216)	8213 (+/- 721)	250 uM	10x
LacO	RK2	Gent	1823 (+/- 78)	7654 (+/- 230)	250 uM	4x
LacO	RK2	Spec	950 (+/- 46)	4967 (+/- 895)	16 uM	5x
NahR	BBR1	Kan	1260 (+/- 57)	16484 (+/- 1693)	500 uM	9x
NahR	BBR1	Gent	1877 (+/- 334)	16317 (+/- 1534)	31 uM	9x
NahR	BBR1	Spec	1372 (+/- 84)	14083 (+/- 984)	1 mM	10x
NahR	RK2	Kan	445 (+/- 39)	8048 (+/- 859)	1 mM	18x
NahR	RK2	Gent	934 (+/- 56)	9299 (+/- 460)	1 mM	10x
NahR	RK2	Spec	482 (+/- 69)	7510 (+/- 817)	1 mM	16x
Jungle Express	BBR1	Kan	100 (+/- 2)	28545 (+/- 1890)	125 nM	285x
Jungle Express	BBR1	Gent	124 (+/- 24)	31409 (+/- 1348)	625 nM	254x
Jungle Express	BBR1	Spec	107 (+/- 9)	31360 (+/- 905)	313 nM	292x
Jungle Express	RK2	Kan	151 (+/- 25)	11702 (+/- 3180)	625 nM	78x
Jungle Express	RK2	Gent	91 (+/- 6)	19413 (+/- 3180)	156 nM	214x
Jungle Express	RK2	Spec	153 (+/- 18)	4978 (+/- 305)	20 nM	32x
TetR	BBR1	Kan	341 (+/- 10)	33631 (+/- 7692)	1 uM	98x
TetR	BBR1	Gent	351 (+/- 32)	21646 (+/- 5579)	500 nM	62x
TetR	BBR1	Spec	232 (+/- 43)	19224 (+/- 3027)	125 nM	83x
TetR	RK2	Kan	184 (+/- 5)	10281 (+/- 967)	31 nM	56x
TetR	RK2	Gent	232 (+/- 39)	7883 (+/- 865)	63 nM	34x
TetR	RK2	Spec	197 (+/- 5)	3399 (+/- 143)	16 nM	17x

151 **Table 4: Inducible pGinger variants in *E. coli*:** For pGinger variants of LacO, NahR, Jungle Express, and TetR

152 promoters, the experimentally observed background (uninduced) fluorescence and maximal fluorescence in *E. coli*

153   are provided. Standard deviations are provided in parentheses, n=3. Additionally, the inducer concentration used to

154   achieve maximal expression and the relative induction levels are listed.

Discussion

The pGinger suite of plasmids offers researchers an array of small, pre-assembled vectors

that will permit rapid identification of useful genetic elements in diverse gram-negative bacteria

due to the use of broad host-range origins (RK2) and selectable markers known to work across

many species (kanamycin, spectinomycin, gentamicin). The compatibility of RK2 and BBR1

origins may also permit researchers to introduce multiple pGinger vectors into a single strain

simultaneously (11). In combination with other recent plasmid suites that have been publicly

released, the pGinger plasmids have the potential to facilitate more advanced synthetic biology

and metabolic engineering efforts in bacterial species that have been traditionally understudied.

Materials & Methods

<u>Strains and Media</u>

Cultures were grown in lysogeny broth (LB) Miller medium (BD Biosciences, USA) at

37 °C for *E. coli* XL1-Blue (QB3 Macrolab, USA) and 30 °C for *P. putida* KT2440 (ATCC

47054). The medium was supplemented with kanamycin (50 mg/L, Sigma Aldrich, USA),

gentamicin (30 mg/L, Fisher Scientific, USA), or spectinomycin (100mg/L, Sigma Aldrich,

USA), when indicated. All other compounds were purchased through Sigma Aldrich (Sigma

Aldrich, USA).

<u>Plasmid Design and Construction</u>

All plasmids were designed using Device Editor and Vector Editor software, while all

primers used for the construction of plasmids were designed using j5 software (12, 18, 19).

Plasmids were assembled via Gibson Assembly using standard protocols (20). Plasmids were

routinely isolated using the Qiaprep Spin Miniprep kit (Qiagen, USA), and all primers were

purchased from Integrated DNA Technologies (IDT, Coralville, IA).

<u>Plasmid and Sequence Availability</u>

All strains and plasmids from Table 1 can be found via the following link to the JBEI

Public Registry: <u>https://public-registry.jbei.org/folders/771</u>.

<u>Characterization assays</u>

To characterize RFP expression from these vectors, we measured optical density and

fluorescence after growth in 96 well plates for 24 hours. First, overnight cultures were inoculated

into 5 mL of LB medium from single colonies and grown at 30 °C or 37 °C. These cultures were

then diluted 1:100 into 500 μL of LB medium with the appropriate antibiotic in 96 square v-

bottom deep well plates (Biotix™ DP22009CVS). For characterization of the inducible systems,

inducer was added to wells in the first column of the plate at the maximum concentration tested

and diluted two-fold across the plate until the last column, which was left as the zero-inducer

control. Plates were sealed with a gas-permeable microplate adhesive film (Axygen™ BF400S)

and grown for 24 hours at either 30 °C or 37 °C with shaking at 200 rpm. Optical density was

measured at 600 nm, and fluorescence was measured at an excitation wavelength of 535 nm and

an emission wavelength of 620 nm. All data was analyzed and visualized using custom Python

scripts using the SciPy (22), NumPy (23), Pandas, Matplotlib, and Seaborn libraries. Fits to the

Hill equation were done as previously described (24).

Acknowledgements

The order of authors was determined by who has the reddest hair. Mitchell Thompson is a Simons Foundation Awardee of the Life Sciences Research Foundation. Lucas Waldburger is funded by the National Science Foundation Graduate Research Fellowship. This work was part of the DOE Joint BioEnergy Institute (https://www.jbei.org) supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, supported by the U.S. Department of Energy, Energy Efficiency and Renewable Energy, Bioenergy Technologies Office, through contract DE-AC02-05CH11231 between Lawrence Berkeley National Laboratory and the U.S. Department of Energy. The views and opinions of the authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. The United States Government retains and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a nonexclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes. The Department of Energy will provide public access to these results of federally sponsored research in accordance with the DOE Public Access Plan (http://energy.gov/downloads/doe-public-access-plan).

Contributions

Conceptualization; A.N.P., M.G.T.; Methodology; A.N.P., M.G.T..; Investigation, A.N.P.,

M.G.T., L.D.K., K.M.V., L.M.W.; Writing – Original Draft, A.N.P., M.G.T..; Writing – Review

and Editing, All authors.; Resources and supervision; P.M.S, J.D.K.

Competing Interests

J.D.K. has financial interests in Amyris, Ansa Biotechnologies, Apertor Pharma, Berkeley Yeast,

Demetrix, Lygos, Napigen, ResVita Bio, and Zero Acre Farms.

References

1.   Khalil AS, Collins JJ. 2010. Synthetic biology: applications come of age. Nat Rev Genet
     11:367–379.

2.   Nielsen AAK, Segall-Shapiro TH, Voigt CA. 2013. Advances in genetic circuit design:
     novel biochemistries, deep part mining, and precision gene expression. Curr Opin Chem
     Biol 17:878–892.

3.   Zhang Y, Ding W, Wang Z, Zhao H, Shi S. 2021. Development of Host-Orthogonal
     Genetic Systems for Synthetic Biology. Advanced Biology 5:e2000252.

4.   Johns NI, Gomes ALC, Yim SS, Yang A, Blazejewski T, Smillie CS, Smith MB, Alm EJ,
     Kosuri S, Wang HH. 2018. Metagenomic mining of regulatory elements enables
     programmable species-selective gene expression. Nat Methods 15:323–329.

5.   Anderson JC, Dueber JE, Leguia M, Wu GC, Goler JA, Arkin AP, Keasling JD. 2010.
     BglBricks: A flexible standard for biological part assembly. J Biol Eng 4:1.

237 6. Lee TS, Krupa RA, Zhang F, Hajimorad M, Holtz WJ, Prasad N, Lee SK, Keasling JD.

238   2011. BglBrick vectors and datasheets: A synthetic biology platform for gene expression.

239   J Biol Eng 5:12.

240 7. Shih PM, Vuu K, Mansoori N, Ayad L, Louie KB, Bowen BP, Northen TR, Loqué D.

241   2016. A robust gene-stacking method utilizing yeast assembly for plant synthetic biology.

242   Nat Commun 7:13215.

243 8. Cook TB, Rand JM, Nurani W, Courtney DK, Liu SA, Pfleger BF. 2018. Genetic tools for

244   reliable gene expression and recombineering in *Pseudomonas putida*. J Ind Microbiol

245   Biotechnol 45:517–527.

246 9. Phelan RM, Sachs D, Petkiewicz SJ, Barajas JF, Blake-Hedges JM, Thompson MG,

247   Reider Apel A, Rasor BJ, Katz L, Keasling JD. 2017. Development of Next Generation

248   Synthetic Biology Tools for Use in *Streptomyces venezuelae*. ACS Synth Biol 6:159–166.

249 10. Schuster LA, Reisch CR. 2021. A plasmid toolbox for controlled gene expression across

250   the Proteobacteria. Nucleic Acids Res 49:7189–7202.

251 11. Pasin F, Bedoya LC, Bernabé-Orts JM, Gallo A, Simón-Mateo C, Orzaez D, García JA.

252   2017. Multiple T-DNA Delivery to Plants Using Novel Mini Binary Vectors with

253   Compatible Replication Origins. ACS Synth Biol 6:1962–1968.

254 12. Ham TS, Dmytriv Z, Plahar H, Chen J, Hillson NJ, Keasling JD. 2012. Design,

255   implementation and practice of JBEI-ICE: an open source biological part registry platform

256   and tools. Nucleic Acids Res 40:e141.

257 13. Bi C, Su P, Müller J, Yeh Y-C, Chhabra SR, Beller HR, Singer SW, Hillson NJ. 2013.

258   Development of a broad-host synthetic biology toolbox for *Ralstonia eutropha* and its

application to engineering hydrocarbon biofuel production. Microb Cell Fact 12:107.

14.   Ruegg TL, Pereira JH, Chen JC, DeGiovanni A, Novichkov P, Mutalik VK, Tomaleri GP,

Singer SW, Hillson NJ, Simmons BA, Adams PD, Thelen MP. 2018. Jungle Express is a

versatile repressor system for tight transcriptional control. Nat Commun 9:3617.

15.   Calero P, Jensen SI, Nielsen AT. 2016. Broad-Host-Range ProUSER Vectors Enable Fast

Characterization of Inducible Promoters and Optimization of p-Coumaric Acid Production

in *Pseudomonas putida KT2440*. ACS Synth Biol 5:741–753.

16.   Shanks RMQ, Kadouri DE, MacEachran DP, O'Toole GA. 2009. New yeast

recombineering tools for bacteria. Plasmid 62:88–97.

17.   Schmidl SR, Sheth RU, Wu A, Tabor JJ. 2014. Refactoring and optimization of light-

switchable *Escherichia coli* two-component systems. ACS Synth Biol 3:820–831.

18.   Chen J, Densmore D, Ham TS, Keasling JD, Hillson NJ. 2012. DeviceEditor visual

biological CAD canvas. J Biol Eng 6:1.

19.   Hillson NJ, Rosengarten RD, Keasling JD. 2012. j5 DNA assembly design automation

software. ACS Synth Biol 1:14–21.

20.   Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison CA, Smith HO. 2009.

Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat Methods

6:343–345.

21.   Thompson MG, Pearson AN, Barajas JF, Cruz-Morales P, Sedaghatian N, Costello Z,

Garber ME, Incha MR, Valencia LE, Baidoo EEK, Martin HG, Mukhopadhyay A,

Keasling JD. 2020. Identification, Characterization, and Application of a Highly Sensitive

Lactam Biosensor from *Pseudomonas putida*. ACS Synth Biol 9:53–62.

22.   Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski

E, Peterson P, Weckesser W, Bright J, van der Walt SJ, Brett M, Wilson J, Millman KJ,

Mayorov N, Nelson ARJ, Jones E, Kern R, Larson E, Carey CJ, SciPy 1.0 Contributors.

2020. SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat Methods

17:261–272.

23.   van der Walt S, Colbert SC, Varoquaux G. 2011. The NumPy Array: A Structure for

Efficient Numerical Computation. Comput Sci Eng 13:22–30.

24.   Thompson MG, Costello Z, Hummel NFC, Cruz-Morales P, Blake-Hedges JM, Krishna

RN, Skyrud W, Pearson AN, Incha MR, Shih PM, Garcia-Martin H, Keasling JD. 2019.

Robust Characterization of Two Distinct Glutarate Sensing Transcription Factors of

Pseudomonas putida l -Lysine Metabolism. ACS Synth Biol 8:2385–2396.