A reliable and efficient adaptive Bayesian method to assess static lower limb proprioception

Jonathan M Wood[1,2], *Susanne M Morton[1,2], *Hyosub E Kim[1,2,3]

*Both authors contributed equally

1. Department of Physical Therapy, University of Delaware, Newark, DE 19711, United States
2. Biomechanics and Movement Sciences Program, University of Delaware, Newark, DE 19711, United States
3. Department of Psychological and Brain Sciences, University of Delaware, Newark, DE 19716, United States

Abstract

Background

Lower limb proprioception is critical for maintaining stability during gait and may impact how individuals modify their movements in response to changes in the environment and body state, a process termed "sensorimotor adaptation". However, the connection between lower limb proprioception and sensorimotor adaptation during human gait has not been established. We suspect this gap is due in part to the lack of reliable, efficient methods to assess global lower limb proprioception in an ecologically valid context.

New Method

We assessed static lower limb proprioception using an alternative forced choice task, administered twice to determine test-retest reliability. Participants stood on a dual-belt treadmill which passively moved one limb to stimulus locations selected by a Bayesian adaptive algorithm. At the stimulus locations, participants judged relative foot positions and the algorithm estimated the point of subjective equality (PSE) and the uncertainty of lower limb proprioception.

Results

Using the Bland-Altman method, combined with Bayesian statistics, we found that both the PSE and uncertainty estimates had good reliability.

Comparison with Existing Method(s)

Current methods assessing static lower limb proprioception do so within a single joint, in non-weight bearing positions, and rely heavily on memory. One exception assessed static lower limb proprioception in standing but did not measure reliability and contained confounds impacting participants' judgments, which we experimentally controlled here.

Conclusions

This efficient and reliable method assessing lower limb proprioception will aid future mechanistic understanding of locomotor adaptation and serve as a useful tool for basic and clinical researchers studying balance and falls.

1. Introduction

Lower limb proprioception is critical for maintaining upright stability and regulating the gait cycle. The proprioceptive sense is formed through a combination of inputs, mostly from muscle spindles and Golgi tendon organs, but also cutaneous and joint capsule receptors (Proske and Gandevia, 2012). Here, we are most interested in proprioception as it relates to static lower limb position sense. Static lower limb position sense contributes to balance and stability as evidenced by studies showing that impaired static lower limb proprioception is associated with higher fall risk (Lord et al., 1991; Lord and Ward, 1994; Ribeiro and Oliveira, 2007). In addition to its contributions to posture and stability, static lower limb proprioception may also impact how individuals implicitly adapt their gait pattern in response to changes in the environment or body state (e.g., fatigue) (Bruijn et al., 2012; Bunday and Bronstein, 2009), a learning process termed "sensorimotor adaptation" (Prokop et al., 1995; Reisman et al., 2005). This link between static proprioception and sensorimotor adaptation is well established in upper-extremity reaching (Cressman and Henriques, 2009; Harris, 1963; Henriques and Cressman, 2012; Mattar et al., 2013; Ostry et al., 2010; Simani et al., 2007; Tsay et al., 2022). However, no causal link between proprioception and locomotor adaptation has been clearly established in humans, which is surprising given how dependent normal walking is on reliable proprioceptive estimates (Dietz, 2002; Hiebert et al., 1996; Kriellaars et al., 1994; Pearson, 2004; Roden-Reynolds et al., 2015; Whelan et al., 1995). We suggest that this discrepancy exists because of the way proprioception is measured in the lower limb. Specifically, most lower limb proprioception assessments are not performed in an ecologically valid context, and those that are lack established reliability and validity (Sombric et al., 2019; Vazquez et al., 2015). In order to address these limitations, we have developed a new psychophysical method for assessing lower limb proprioception.

We first considered the measurement of lower limb proprioception in a context that most closely approximates gait. Current methods assessing static lower limb proprioception often do so within a single joint, in a non-weight bearing position, and rely heavily on remembered positions (Han et al., 2016; Hillier et al., 2015; Horváth et al., 2022). While these methods have proven useful for characterizing specific deficits in joint proprioception after orthopedic injury (e.g., Relph et al., 2014), they cannot be readily translated to functional lower extremity movements. This is because a unified percept of limb location results from the nervous system's integration of proprioceptive signals across multiple limb joints (Bosco et al., 2000; Fuentes and Bastian, 2010; Gandevia, 1985; Proske and Gandevia, 2012; Soechting, 1982). The body position in which proprioception is measured is also important because differences in proprioceptive estimates emerge in weight bearing vs non-weight bearing positions both in the upper limb (Ansems et al., 2006) and in the knee joint (Bullock-Saxton et al., 2001; Stillman and McMeeken, 2001). For these reasons, we measured whole lower limb proprioception while standing as this provides the closest approximation to gait.

A comprehensive way of characterizing a sensory system is using a psychophysical assessment to estimate two of its distinct physiologic characteristics: the point of subjective equality (PSE) and the uncertainty. The PSE is the stimulus that is perceived as equal to another stimulus, reflecting the limits of a sensory system to discriminate between two stimuli. In the context of the current study, the PSE is the location where participants perceive the two lower limbs are symmetrical. The PSE can also be translated to a measure of proprioceptive accuracy by calculating the difference between the PSE and actual equality (i.e., a proprioceptive bias). Uncertainty refers to the variability in responses surrounding the PSE, reflecting the noise within the sensory system. The PSE (in terms of a proprioceptive bias) and uncertainty seem to play distinct roles in sensorimotor adaptation (Ruttle et al., 2021; Tsay et al., 2021). While upper extremity proprioceptive biases shift as a result of sensorimotor adaptation, proprioceptive

102 uncertainty at baseline predicts the magnitude of sensorimotor adaptation. The proprioceptive
103 shift and uncertainty estimates themselves are uncorrelated (Tsay et al., 2021). Thus, it is
104 important to reliably estimate both the PSE and uncertainty.
105
A widely accepted method of estimating both the PSE and uncertainty is to use a two-alternative
forced choice (2AFC) task, where, in the context of the current study, participants judge the
relative positions of their feet when placed at various locations. The response data are then fit
with a psychometric function, where the inflection point (i.e., the probability of a response being
0.5), represents the PSE and the slope of the function is inversely proportional to the uncertainty
(Kingdom and Prins, 2016). We know of two studies that used an AFC task to estimate both PSE
and uncertainty of lower limb proprioception (Vazquez et al., 2015; Waddington and Adams,
1999). However, the Waddington study used active repositioning focused on only ankle inversion
and eversion movements, whereas the Vazquez study did not measure or report the reliability of
their method. Furthermore, in the latter study participants' responses were heavily impacted by
the direction in which the test limb was moved to the stimulus position, which was either always
forward or always backward depending on group assignment. Ideally, in a static assessment of
lower limb proprioception, the movement of the limbs to stimulus positions should not exert any
influence on participants' judgments. Therefore, while a method like the one used in the study by
Vazquez and colleagues offers promise, adjustments are needed to improve the validity of
responses, and the reliability of the estimates must be assessed.
122
Efficiency is also critical for assessing changes in proprioception in single-session motor learning
studies, when considering factors such as clinical feasibility, and to maintain participant safety,
as standing still for long periods can lead to syncope (Jardine et al., 2018). In the most established
psychophysical method, the method of constant stimuli, participants make the same number of
sensory judgements at each pre-determined stimulus location. However, this method is inefficient
because some of the stimulus locations provide little information about the PSE or uncertainty
estimates (Kingdom and Prins, 2016; Leek, 2001; Watson and Fitzhugh, 1990). Indeed, Kingdom
and Prins (Kingdom and Prins, 2016, p. 57) suggested that as many as 400 trials are required to
accurately estimate both the PSE and uncertainty. Adaptive psychophysical methods, where the
stimulus locations are selected based on prior responses, were specifically developed to solve
this efficiency problem (Leek, 2001). The Psi algorithm is one such method that uses Bayesian
estimation to calculate PSE and uncertainty values after each trial. It then chooses the next
stimulus that will maximize the information gained for both estimates on the subsequent trial
(Kontsevich and Tyler, 1999). This method results in more efficient and accurate estimates of
both the PSE and slope compared to the method of constant stimuli (King-smith and Rose, 1997;
Kontsevich and Tyler, 1999; Livesey and Livesey, 2016; Turpin et al., 2010). The Psi algorithm is
most frequently used in visual and auditory psychophysics, and we note only one instance where
is was used to estimate (wrist) proprioception (Elangovan et al., 2018).
141
Here, we implemented an adapted version of the Psi algorithm and measured its reliability in
assessing lower limb proprioception in an ecologically valid context using a test-retest design. We
first wanted to ensure that participant responses were not confounded by the direction of
movement to each stimulus location, a concern we had based on a prior study (Vazquez et al.,
2015). Then, to assess reliability across test sessions, we calculated agreement, defined as the
ability for a test to reproduce the same values when measured at different times, using the Bland-
Altman method (Altman and Bland, 1983; Giavarina, 2015). In addition, we used Bayesian
statistics to fully quantify the probability that the method has good agreement, defined as low
evidence of bias in the PSE and uncertainty estimates across the two tests. Combined, we found
that, with the current method, movement direction did not impact participant responses, and good
agreement for both the PSE and uncertainty could be achieved after only 50 trials.

153
2. Materials and methods

156 *2.1. Participants*

Young healthy participants between the ages of 18 and 35 were recruited from the University of Delaware community. Participants were excluded if they had any chronic or recent musculoskeletal or neurologic diagnoses, pain, or impaired sensation. This work was completed in accordance with the Code of Ethics of the World Medical Association. All participants provided written informed consent prior to being enrolled into the study. This study was approved by the University of Delaware Institutional Review Board.

165 *2.2. Motion capture*

Participants stood on a dual-belt treadmill instrumented with two force plates, one under each belt (Bertec, Columbus OH). We obtained kinematic data, sampled at 100Hz, using an eight-camera Vicon MX40 motion capture system with Nexus software (Vicon Motion Systems Inc., London, UK). Seven retroreflective markers were placed on participants' shoes and ankles in the following locations: bilateral heels, bilateral lateral malleoli, bilateral 5^{th} metatarsal head, and the left 1^{st} metatarsal head. We used custom written MATLAB scripts (version 2022a, MathWorks, Natick, MA) to control the treadmill belts and obtain live kinematic and kinetic data from Nexus software.

175 *2.3. AFC task*

We used a two-AFC task to measure lower limb proprioception. To measure test-retest reliability, each participant performed the test twice on the same day, with a 20-minute break between tests. For each test, participants stood on a treadmill with vision of their legs occluded with a black drape, and auditory feedback (sounds of the treadmill belts/motors) occluded with noise cancelling headphones (Figure 1A). The primary kinematic variable for the test was foot position difference (in millimeters):

$$Foot\ position\ difference = Left\ marker_{Ypos} - Right\ marker_{Ypos} \qquad (1)$$

We used the lateral malleoli markers to measure foot position difference. Thus, positive values indicate the left foot was forward of the right foot, while negative values indicate the left foot was behind. Since the foot is the end-effector for the lower limb, we assume judgements were made by combining proprioceptive information across lower limb joints. The right foot served as the reference foot and did not move throughout the test. The left foot served as the test foot and was passively moved by the left treadmill belt to the stimulus positions, measured in terms of foot position difference. After participants performed two practice trials to orient them to the task, their heels were aligned so that everyone started from the same position relative to the laboratory's y-axis. Since there may still be differences in the ankle markers along the y-axis in this position, we corrected for this baseline difference (rounded to the nearest millimeter) for stimulus positions and PSE estimates. Each test was comprised of 75 trials, and each trial had two parts: 1) movement to a start position, 2) movement to a stimulus position.

The treadmill moved the test foot to a start position at a speed selected from a uniform distribution between 40 and 50 mm/s (Figure 1B, top). We controlled for the potential bias in responses caused by movement direction by providing pseudorandomized start positions so that the test foot started in front of the stimulus position 38 times and behind the stimulus 37 times. Therefore, we assume any bias in responses from always moving to the stimulus from the same direction

204 would be washed out, something we confirmed in our formal analysis (see Section 3.1). The
205 specific start position on a given trial, t, was selected from one of two normal distributions centered
206 around either -100 or +100 mm of foot position difference depending on the movement direction
207 for that trial. Both distributions had a standard deviation of 5 mm:
208

209 $$Start\ position_{[t]} \sim \begin{cases} \mathcal{N}(+100, 5), & if\ movement\ direction_{[t]} == backward \\ \mathcal{N}(-100, 5), & if\ movement\ direction_{[t]} == forward \end{cases} \quad (2)$$

210
After a 0-2 second pause at the start position, the test foot was moved to a stimulus position at a
speed selected from a uniform distribution between 10 and 30 mm/s (Figure 1B, bottom). The
stimulus positions were located at 21 possible foot difference locations: every 10 mm interval
between -100 to +100 mm. Most stimulus positions were selected using the Psi algorithm (see
section 2.5). However, we inserted pre-selected stimulus locations to keep participants engaged
in the task. There were two types of pre-selected stimuli: 1) far stimuli (±100 or ±90) inserted
randomly once every 10 trials, and 2) near stimuli (±10, ±20 or ±30 mm from the current PSE
estimate, rounded to the nearest stimulus location) inserted randomly once every 5 trials. No
preselected stimuli were inserted within the first 5 trials.

220
221 Each time the test foot reached its stimulus position, the treadmill stopped, and a prompt appeared
222 on a monitor in front of the participant: "Do you feel your right or left foot is more forward?" The
223 participant's response was recorded by the experimenter using a custom graphical user interface
224 (GUI) in MATLAB.

225
Before the test, participants were instructed to keep their weight equally distributed between both
feet, measured with the force plates under each treadmill belt and displayed for the experimenter.
If participants consistently kept greater than ~60% of their weight through one foot, a verbal
reminder was provided. We implemented short breaks after 25 and 50 trials to prevent fatigue
and blood pooling in the legs. Before the break, current marker positions were recorded from
Nexus software, then the participant was asked to walk around the lab for ~30 seconds. After the
participant stepped back onto the treadmill, we positioned their feet so the ankle markers were
back to the exact location from which they started and then testing continued.

Figure 1. Task setup, representative data, and movement direction analysis. (A) AFC task setup. Participants stood on the split-belt treadmill with visual feedback of their legs occluded by a black drape and auditory feedback occluded by noise cancelling headphones. The screen prompted participants to verbally respond to the question "Do you feel like your right or left foot is more forward?" when the test foot stopped at the stimulus location. **(B)** Trial sequence. 1) The test foot was moved to a start position either in front of or behind the stimulus position. Start positions were sampled from one of two normal distributions, depending on the movement direction assigned for that specific trial. 2) The test foot was moved to one of 21 possible stimulus positions between -100 and +100 mm. Tick marks (drawn to scale) represent each possible stimulus location. **(C)** Individual participant data. We reconstructed the psychometric functions from the PSE (α) and uncertainty (β) estimates for Test 1 (blue) and Test 2 (red). The dashed black lines, provided as a reference, represent a PSE of 0 mm. The individual dots represent the participant's response data for each test. **(D)** Movement direction analysis. We measured the mean empirical probability of responding left when the movement direction to the stimulus position was forward (gray) vs backward (black). Dots represent individuals. We also used a Bayesian logistic regression model to analyze the unique impact of each movement direction on participant responses by calculating the difference between the posterior estimates of the movement direction coefficients (one for forward movement, one for backwards) of every participant, yielding a posterior contrast distribution (histogram) which we compared to a region of practical equivalence (ROPE). We note that 99.3% of the posterior distribution is within the ROPE, confirming that responses were not biased by movement direction to the stimulus position.

2.4. Psychometric function

Each participant's probability of responding that the test foot was more forward at each stimulus position (x) was represented as a normal cumulative distribution function (cdf) with two parameters:

$$\psi(x) = normcdf(x; \alpha, \beta) \tag{3}$$

The α parameter, the mean or inflection point of the cdf, represents the PSE, corresponding to the position that the participant perceives the feet were in the same location along the laboratory's y-axis (i.e., where the probability of judging "left" is exactly 0.5). The β parameter, the standard deviation of the cdf, represents the uncertainty, which reflects the noise within the sensory system itself. Both are measured in terms of foot position difference. The α and β parameters were estimated adaptively on each trial using the Psi algorithm. Of note, in our pilot testing the probability of responding left at -100 and +100 was consistently 0 or 1, respectively. Thus, we did not include the lapse and guess rate parameters that are sometimes used in psychometric functions for AFC tasks.

2.5. Psi Algorithm

Here we provide a brief description of the Psi algorithm, which is described in detail by Kostovich and Tyler (1999). There are two primary components of the Psi algorithm: 1) stimulus selection and 2) parameter estimation.

2.5.1 Stimulus selection

Before each trial, the Psi algorithm simulates responses at each possible stimulus location for the next trial, calculating a simulated joint posterior distribution using Bayes' rule. The simulated joint posterior distribution (p_{sim}) contains the probability of every possible combination of α and β values given the simulated responses and the stimulus location. Information entropy (H), a measure of the magnitude of uncertainty in a probability distribution, is calculated for each stimulated posterior distribution:

$$H_{[x]} = -\sum_i p_{sim[i]} log_2 p_{sim[i]} \tag{4}$$

289
290 The stimulus position that minimizes the information entropy is selected for the subsequent trial
291 as it provides the most certainty and thus the greatest gain in information for the α and β estimates
292 from the current trial to the next (Shannon, 1948). That is, the stimulus position that will
293 theoretically aid in maximally efficient parameter estimation is selected on each trial.
294
295 *2.5.2 Parameter estimation*
296
297 Once the participant makes their selection at the stimulus location, the Psi algorithm uses Bayes'
298 rule to estimate the most likely α and β values given the prior and the current response (r):
299

$$p_{[t]}(\lambda|x,r) = \frac{p_{[t]}(\lambda)p(r|\lambda,x)}{\sum_\lambda \left(p_{[t]}(\lambda)p(r|\lambda,x)\right)} \tag{5}$$

301
The prior, $p_{[t]}(\lambda)$, is a joint probability distribution representing the initial guess of α and β values.
The prior for the first trial was the same for all participants. We assumed that the most likely α
was 0 with a wide standard deviation: $\alpha \sim N(0,20)$. Similarly, we assumed a reasonably wide prior
distribution for β, with an expected value of 20: $\beta \sim Exponential(20)$. On subsequent trials, the
posterior for trial t became the prior for trial t+1 ($p_{[t+1]}(\lambda) = p_{[t]}(\lambda|x,r)$). The likelihood in equation
5, $p(r|\lambda,x)$, is the probability of the participant's response given each parameter value at the
stimulus position. As suggested by Kostovich and Tyler, we created a pair of lookup tables to
improve computational efficiency, one for each response, which served as indexable likelihoods
for each stimulus:
311
312 $p(r = "left"|\lambda,x) = \psi$ (6a)
313 $p(r = "right"|\lambda,x) = 1 - \psi$ (6b)
314
315 The individual α and β estimates were calculated by marginalizing over the joint posterior
316 distribution (equation 5) and taking the mean of each marginalized distribution (Emerson, 1986).
317 We used the final α and β estimates after the last (75[th]) trial for our agreement analyses.
318
319 *2.7. Statistical analysis*
320
We used a Bayesian approach to make statistical inferences for the validity of responses and the
agreement of PSE and uncertainty estimates. This approach enabled us to calculate the full
posterior probability distribution of model parameters, providing a complete picture that quantifies
our uncertainty. Thus, it naturally emphasizes estimation over binary decision rules in accordance
with recommendations made by the American Statistical Association and others (Kruschke, 2013;
Kruschke and Liddell, 2018; Wasserstein and Lazar, 2016).
327
For each analysis, we started by defining a statistical model that was consistent with our question
and the data structure. Next, we calculated the posterior probability of all parameter values in the
statistical model using Bayes' rule, combining our prior assumptions regarding parameter values
(i.e., the prior) with evidence from our data (i.e., the likelihood). We selected the distribution for
each parameter's prior as reasonably wide and uninformative, using its maximum entropy
distribution (McElreath, 2016). We used the Pymc4 (version 4.3) library (Salvatier et al., 2016) in
Python (version 3.11) to perform Markov Chain Monte Carlo (MCMC) sampling to estimate the
posterior probability distributions. We drew 10,000 samples from the posterior in each of 4 chains
(i.e., 40,000 total samples), using 2,000 tuning samples in each chain. We performed diagnostics
for each model, ensuring parameter values were consistent across chains and checking posterior

estimates for possible errors (Kruschke, 2014; McElreath, 2016). We provide the full models and
code, including detailed information about the priors and diagnostics, online at
(https://osf.io/g8nx4/). We made inferences based on the posterior distribution of the model
parameters, reporting the 95% high density interval (HDI), defined as the narrowest span of
credible values that contains 95% of the posterior distribution (Kruschke, 2014). In cases in which
we wanted to quantify support for the "null" hypothesis, we also calculated the percent of credible
parameter values that fell within a region of practical equivalence (ROPE; Kruschke, 2014).

First, we ensured that responses were not confounded by movement direction, a concern based
on a prior study (Vazquez et al., 2015). We modeled each participant's response data using a
Bayesian logistic regression. Since the response data on each trial (t) were binary, we modeled
them as a Bernoulli distribution where the probability of responding "left" (p_{left}) was impacted by
the stimulus location ($X_{Stimulus}$), the movement direction ($\beta_{Move\ direction}$), and the participant
($\alpha_{Participant}$):

$Response_{[t]} \sim Bernoulli(p_{left[t]})$ (7a)
$logit(p_{left[t]}) = \alpha_{Participant[i]} + \beta_{Movedirection[j]} + \beta_{Stimulus} X_{Stimulus[t]}$ (7b)

We included data from both tests for each participant in this model. Stimulus position was coded
as a continuous variable. Movement direction and participant were coded as indexing variables,
meaning separate $\alpha_{Participant}$ posteriors were computed for each participant (i; 13 total alphas),
representing each participant's bias to judge "left" independent of movement direction or stimulus
position, and two separate group level $\beta_{Move\ direction}$ posteriors were computed for each
movement direction (j; forward and backward; 2 total $\beta_{Move\ direction}$ parameters) to the stimulus
position. We reasoned that if moving forward or backward had no influence on responses, the
two posterior distributions for $\beta_{Move\ direction}$ should be identical. We therefore calculated the
difference in probability between the $\beta_{Move\ direction}$ posterior distributions, setting a ROPE for this
contrast between -0.1 and 0.1 in terms of probability.

To assess test-retest agreement, we used the Bland-Altman method (Altman and Bland, 1983;
Giavarina, 2015). The Bland-Altman method is recommended to assess agreement because it
specifically tests for biases across the range of "true" scores of a given variable, where a proxy
for the true score is the mean of an individual's score on Test 1 and Test 2 (Bland and Altman,
1986). The method involves 3 analyses: 1) calculating the mean bias between the two estimates,
2) calculating the bias across "true" values, and 3) determining the limits of agreement, calculated
as the mean of the difference in estimates ±1.96 times the standard deviation of the difference in
estimate. To test for biases in steps 1 and 2 we applied Bayesian inference instead of the
frequentist methods typically used in the Bland-Altman method. We determined if there was a
mean bias by estimating the distribution of differences between Test 1 and Test 2 for the PSE
and uncertainty estimates separately. We modeled these contrasts as a normal distribution,
estimating the most likely μ and σ values that could have generated each individual's (i) estimated
difference from Test 1 to Test 2:

$Difference_{[i]} \sim Normal(\mu, \sigma)$ (8)

We focused our inference on the posterior distribution for μ, which, if no bias was present, should
be close to zero. We therefore set a ROPE between -5 and 5 mm. We determined if there was a
bias across the range of "true" values of PSE and uncertainty using a Bayesian regression
analysis. The outcome variable, differences between Test 1 and Test 2 estimates for each

participant ($Y_{[i]}$), was modeled as a normal distribution with the mean (μ) being predicted by the
"true" PSE/uncertainty value (X_{true}):
389
390   $Y_{[i]} \sim Normal(\mu_{[i]}, \sigma)$ (9a)
391   $\mu_{[i]} = b + m\, X_{true[i]}$ (9b)
392
Here, a bias will manifest as an intercept (b) and slope (m) with magnitudes greater than 0.
Therefore, we set a ROPE for the intercept between -5 and 5 mm and a ROPE for the slope
between -0.1 and 0.1, which would translate to a 1 mm change in bias for a 10 mm change in
"true" score. We defined good agreement as little to no evidence of a mean bias or a bias across
true scores. Specifically, 90% of the posterior distributions calculated in the Bland-Altman analysis
should not fall outside the ROPEs. We also performed a secondary analysis to determine if perfect
agreement was at least plausible by characterizing the relationship between Test 1 and Test 2 for
both the PSE and uncertainty estimates. Perfect agreement would result in a slope of 1 and an
intercept of 0. We again used a Bayesian regression model, except the estimate on Test 2 was
the outcome variable ($Y_{Test\,2}$) and the estimate on Test 1 was the predictor variable ($X_{Test\,1}$):
403
404   $Y_{Test\,2[i]} \sim Normal(\mu_{[i]}, \sigma)$ (10a)
405   $\mu_{[i]} = b + m\, X_{Test\,1[i]}$ (10b)
406
407   Here the HDIs for m characterize the relationship between Test 1 and 2.
408
409   **3. Results and discussion**
410
411   *3.1. Movement direction did not influence participant responses*
412
Thirteen participants (8 female, 5 male) completed the study. Data for one representative
participant are displayed in Figure 1C. The psychometric functions (solid curves) were produced
from the PSE (α) and uncertainty (β) estimates after the 75th trial and superimposed onto the
empirical data (dots). For both tests, PSE values were positively biased (Test 1 mean [95% HDI]
= 12.0 mm [3.3 20.6]; Test 2 = 15.3 mm [5.3 25.2]) and uncertainty measures (i.e., SDs) were just
under 20mm (Test 1 = 18.2 mm [14.4 21.8]; Test 2 = 17.5 mm [13.5 21.4]). Our first concern,
based on a prior study (Vazquez et al., 2015), was to ensure that we controlled for the potential
confound that movement direction may have had on responses. While the empirical probability of
responding left when moving backward and forward (Figure 1D, left) appear similar, we also
determined the individual impact of movement direction on responses using a Bayesian logistic
regression. If movement direction did play a major role in responses, the difference in probabilities
for $\beta_{Move\ direction\ [Forward]}$ vs $\beta_{Move\ direction\ [Backward]}$ would be large, making the contrast largely
different from 0. However, we found that the movement direction contrast was practically equal to
0 (Figure 1D, right; posterior mean [95% HDI], contrast = 0.04 [-0.01, 0.09], 99.3% in the ROPE).
Thus, randomizing the start positions prevented biased responses.
428
429   *3.2. Bland-Altman analysis revealed good agreement for both PSE and uncertainty estimates*
430
We used the Bland-Altman method to assess agreement (Figure 2A). First, we calculated the
limits of agreement for PSE and uncertainty estimates (-22.3 to 15.5 mm and -10.1 to 11.6 mm,
respectively; dotted lines in Figure 2A). Limits of agreement for both were relatively narrow
considering the scale of the measurement, and the PSE limits were consistent with the range of
values observed in previous work (Vazquez et al., 2015) which used 33 more trials than the
current method. Next, we characterized the mean bias between Test 1 and Test 2 for PSE and

uncertainty estimates (histograms in Figure 2A). Both biases were very close to 0. The average
bias for PSE was -3.3mm [-9.0, 2.8] (72.4% in ROPE). In other words, while the PSE was
nominally lower on Test 2 than Test 1, we can be fairly confident that there is little-to-no bias for
PSE measurements. With regard to uncertainty estimates, there was effectively no bias as the
95% HDI was fully within the ROPE (0.7mm [-2.8, 4.1], 98.8% in ROPE).
442
Next, we tested for evidence of a bias across different "true" PSE and uncertainty estimates,
where the true estimate is represented by the mean of an individual's estimates on Test 1 and
Test 2 (i.e., the x-axis in the Bland-Altman plot), using a Bayesian linear regression model (solid
regression line and shading in Figure 2A). For PSE, there was a small negative slope, although
the most credible estimates surrounded a slope estimate of zero (-0.15 [-0.56, 0.28], 29.6% in
ROPE), and the intercept was unbiased (-1.3 [-9.5 6.9], 76.0% in ROPE). These results indicate
that participants with larger PSE values tend have increased bias between test 1 and 2. However,
even when considering the largest mean PSE of ~40mm, this only produces a bias of -6mm.
Therefore, we interpret this bias as negligible. For uncertainty estimates, the most probable
regression was very similar to a line with 0 slope and 0 intercept (slope = -0.08 [-0.69, 0.58],
24.5% in ROPE; intercept = 2.0 [-9.8, 13.8], 58.4% in ROPE). Combined, both Bland-Altman
results show that estimated PSE and uncertainty values remain largely consistent across test-
retest sessions and across the range values, a sign of good agreement for this method.
456

457
Figure 2: Point of subjective equality (PSE) and uncertainty estimates have good agreement. (A) Bland-Altman
plot for the PSE (α; top) and uncertainty (β; bottom) estimates. The mean of Test 1 and Test 2 for each individual is
plotted on the x-axis and the difference between the two tests is plotted on the y-axis. Thus, each dot represents one
individual. The limits of agreement are plotted as the dotted lines, and the posterior distributions for the mean biases
are plotted on the right sides as histograms. The most probable regression line is the solid line with the shading
representing the 95% high density interval (HDI), with the black dashed line at 0 as a reference **(B)** Regression model
for PSE and uncertainty. Estimates for Test 2 are plotted against estimates for Test 1 for each individual (dots) with the
unity line (black dashed line) provided as a reference for perfect agreement. The most probable regression line is the
solid line with the shading representing the 95% HDI.

As a secondary analysis, we characterized the relationship between estimates on Test 1 and Test
2 to determine if something close to perfect agreement (slope=1, intercept=0) was at least
plausible, again using a Bayesian linear regression model. Whereas the Bland-Altman analysis
uses information from the two tests to estimate a "true" score (the x-axes in Fig 2A), the regression
analysis measures the difference between the two tests without sharing information across tests.
Figure 2B shows the most credible regression lines for PSE (top) and uncertainty (bottom)
estimates, with the shading representing the 95% HDI. While there were credible slope values
that fell both above and below one (slope = 0.92 [0.49, 1.38], intercept = 4.2 [-3.7, 12.4]), the
single best PSE estimate (maximum a posteriori) between Tests 1 and 2 was very nearly one.
The uncertainty estimates indicated that this parameter is less likely to perfectly reproduce the
scores on Test 1 and Test 2 (slope = 0.65 [0.09, 1.19], intercept = 5.6 [-5.1, 15.8]). Interestingly,
the results of this secondary analysis seemingly conflict with the Bland-Altman analysis. However,
we see them as complementary: While the average bias is low and consistent across scores, the
uncertainty values are less likely to be exactly reproduced from Test 1 to Test 2 compared to the
PSE values. This might be interpreted as a reduction in uncertainty from Test 1 to Test 2 indicating
a practice effect, however, individuals with lower uncertainty on Test 1 demonstrated increased
uncertainty on Test 2. Consistent with reports in other psychophysical assessments, measures of
uncertainty are more difficult to recover compared to PSE (King-smith and Rose, 1997;
Kontsevich and Tyler, 1999; Turpin et al., 2010). Together, the results of this secondary analysis
suggest that, while there is evidence of variability across test sessions, it is well within an
acceptable range for psychophysical assessment. Indeed, the 95% HDI for PSE and uncertainty
estimates both included what we would expect if the tests had perfect agreement.
489
As further evidence of good agreement in our method, and to make direct comparisons with other
studies, we also calculated intraclass correlation coefficients ($ICC_{2,1}$), another commonly used
metric of agreement (Berchtold, 2016; Kottner et al., 2011; McGraw and Wong, 1996). The ICC
values for PSE ($ICC_{2,1}$ [95% CI] = 0.80 [0.48, 0.93]) and uncertainty (0.61 [0.05, 0.85]) indicate
good agreement (Portney and Watkins, 2009). These values fall within ranges of other
psychophysics studies using the Psi algorithm and the method of constants, as those studies
report ICCs for PSE estimates that range between 0.48 to 0.96 for the Psi algorithm (Schilling et
al., 2017; Silva et al., 2020), and between 0.77 to 0.88 for the method of constants (Nicholson et
al., 1997). Our method also compares favorably to other proprioception-specific assessments with
ICCs ranging from 0.11 to 0.95 for measures of proprioceptive accuracy (Antcliff et al., 2021; Arvin
et al., 2015; Deshpande et al., 2003; Gorst et al., 2020; Hillier et al., 2015; Rinderknecht et al.,
2018), and from 0.0 to 0.64 for measures of proprioceptive variability (Juul-Kristensen et al., 2008;
Rahlf et al., 2019; Strong et al., 2021). The fact that the current method has good agreement for
both proprioceptive PSE and uncertainty opens the door for future studies to assess the
importance of uncertainty to lower limb function.
505
506 *3.3. Agreement was similar after 50 trials*
507
One of our primary goals for this method was to maximize efficiency. Participants took an average
of 20±1 minutes to complete testing (including the two short walking breaks). The simplest way
to reduce this time would be to reduce the number of trials, however, there is a risk that reducing
the number of trials would also reduce the agreement of the PSE and uncertainty estimates. Since
the Psi algorithm estimates PSE and uncertainty after every trial, we can assess agreement after
any trial during the test. As a post-hoc analysis, we chose to assess agreement after 50 trials as
the test time at this point would have been between 10-15 minutes. We found that all our
measures of agreement were similar after 50 and 75 trials (Table 1). In contrast, examining
agreement after 25 trials revealed a substantial reduction in agreement. Therefore, decreasing
the number of trials to 50 would significantly reduce the time of the test without sacrificing

agreement, providing a reasonable timeframe for a test to be placed in the middle of a motor
learning paradigm or to be used for a balance and falls risk assessment.
520
521 **Table 1: Agreement comparison after 50 and 75 trials**

	Limits of agreement (mm)	Bland Altman Mean Bias (mm; % in ROPE)	Bland Altman Bias Regression (mm; % in ROPE)	Test 1 vs Test 2 Regression	ICC$_{2,1}$ [95% CI]
PSE (α)					
50 trials	-22.3 to 15.7	-3.2 [-9.2 2.6] (73.5%)	Slope = -0.17 [-0.61 0.29] (26.2%) Intercept = -1.1 [-9.3 7.1] (77.0%)	Slope= 0.91 [0.45 1.40] Intercept = 4.1 [-3.9 12.0]	0.78 [0.43, 0.92]
75 trials	-22.3 to 15.5	-3.3 [-9.0 2.8] (72.4%)	Slope = -0.15 [-0.56 0.28] (29.6%) Intercept = -1.3 [-9.5 6.9] (76.0%)	Slope= 0.92 [0.49 1.38] Intercept = 4.2 [-3.7 12.4]	0.80 [0.48, 0.93]
Uncertainty (β)					
50 trials	-9.3 to 10.2	0.5 [-2.7 3.5] (99.5%)	Slope = 0.03 [-0.65 0.69] (24.3%) Intercept = 0.0 [-12.5 12.2] (60.0%)	Slope= 0.57 [0.03 1.09] Intercept = 7.4 [-2.8 17.4]	0.58 [0.05, 0.85]
75 trials	-10.1 to 11.6	0.7 [-2.8 4.1] (98.8%)	Slope = -0.08 [-0.69 0.58] (24.5%) Intercept = 2.0 [-9.8 13.8] (58.4%)	Slope= 0.65 [0.09 1.19] Intercept = 5.6 [-5.6 15.8]	0.61 [0.05, 0.86]

522
523 *3.4. Limitations and future directions*
524
Despite the efficiency and good agreement of this method, we also recognize some of its
limitations. For instance, one way to increase ecological validity when measuring proprioception
would be to assess three-dimensional position sense as opposed to measuring proprioception in
only the sagittal plane as we have done here. However, we believe our results should at least
generalize to locomotor adaptation on the split-belt, given that, in this context, adaptation is often
measured in sagittal plane kinematics like step length (Reisman et al., 2005). Another potential
limitation is that during standing, although participants' limbs were passively moved by the
motorized treadmill, it is impossible to avoid some subtle movements and associated muscle
activity. While no overt movements were observed during testing, in theory, any voluntary muscle
activity might provide additional information about where the limb is located due to an efference
copy of the motor command (Wolpert et al., 1995). However, in our case, such a compromise was
unavoidable, as the primary goal of this study was to develop an assessment of lower limb position
sense in a context that most closely resembles gait.
538
In addition to measuring proprioception in standing, we assessed position sense of the whole
lower limb by asking individuals to focus on the end effector (foot position) when making their
judgements. Interestingly, there are few static lower extremity proprioception studies that ask
individuals to focus on end effector position (but see Sigmundsson et al., 2000; Vazquez et al.,
2015), despite the functional significance of foot position to successful gait (e.g., stepping over
curbs, trail running, etc.). Conversely, upper extremity studies frequently assess whole limb
proprioception by having participants make judgments regarding end effector position (hand or
finger (Jones et al., 2010; Vindras et al., 1998). These methods have characterized the precision
of limb position sense (van Beers et al., 1998), the relationship between static proprioception and
voluntary reaching movements (Jones et al., 2010), and the importance of static proprioception
to sensorimotor adaptation (Clayton et al., 2014; Cressman and Henriques, 2010; Simani et al.,
2007; Tsay et al., 2021; van Beers et al., 2002). Unfortunately, similar lines of research are absent
in locomotion, despite the critical role of proprioception in walking (Hiebert et al., 1996; Kriellaars
et al., 1994; Pearson, 2004; Roden-Reynolds et al., 2015; Whelan et al., 1995). We speculate
this is due at least in part to the previous absence of reliable and efficient methods of assessing
lower limb proprioception in an upright, functional, and multi-joint context. We hope the current
method opens the door to increasing understanding of the relationship between lower limb
proprioception and locomotor adaptation in young, neurotypical adults as well as in older and

neurologic populations, for whom changes in proprioception may have a significant impact on their ability to adapt gait patterns (Bruijn et al., 2012; Bunday and Bronstein, 2009; Lam and Pearson, 2002; Pearson, 2000; Santuz et al., 2022).

Efficient and reliable, whole lower limb proprioceptive measurements during standing should be useful for non-adaptation studies as well. For example, proprioception is an important part of multifactorial falls assessments (Lord et al., 1991), but the proprioception test previously recommended involves active toe position matching in a non-weight bearing position. The current method offers a more ecologically valid lower limb proprioception test since it is much closer in context to when falls most often occur: during weight bearing activities like walking and transfers (Talbot et al., 2005). Furthermore, since most lower limb proprioception tests only measure proprioceptive accuracy or bias (Han et al., 2016; Hillier et al., 2015; Horváth et al., 2022), it is unknown how proprioceptive uncertainty may relate to falls, something that can now be empirically assessed using the current method.

As there are very few studies reporting PSE and uncertainty values for the lower limbs, we were interested in comparing our values to those reported in upper extremity studies. We found that the average uncertainty estimates in the present study were quite similar to values found in upper extremity proprioception studies (Jones et al., 2010), with both falling just under 2 cm. Additionally, we found that PSE estimates here were biased, a finding consistent with intrinsic biases in perceived hand/arm position (Fuentes and Bastian, 2010; Ingram et al., 2019; Jones et al., 2010; van Beers et al., 1996). We note that the bias reported in the present study is not related to footedness, nor due to moving only the left limb, as we randomized the test limb during pilot testing and found no difference in PSE values when the left versus the right limbs were moved. Importantly, biased estimates in upper extremity studies have been linked to biases in reaching direction (Jones et al., 2010; Vindras et al., 1998), suggesting that future work in the lower extremities may benefit from examining whether PSE biases are related to step length or other gait-related movements.

4. Conclusion

We developed and tested the reliability of a lower limb proprioception assessment in a gait-specific context. This method is efficient, requiring only 50 trials to reliably estimate both the PSE and uncertainty of lower limb proprioception. We believe this method will aid future mechanistic understanding of locomotor adaptation and serve as a useful tool for basic and clinical researchers studying balance and falls.

Acknowledgements
This work was supported by the National Institutes of Health [K12- HD055931 (HEK), R01-AG071585 (SMM), S10-RR028114 (SMM)], the National Science Foundation [M3X 1934650 (HEK)], and partial funding from the University of Delaware Graduate College (JMW). We want to extend a special thank you to Saunders Penn for his help creating Figure 1A and B and for his help performing data collections.

References

Altman, D.G., Bland, J.M., 1983. Measurement in Medicine: The Analysis of Method Comparison Studies. J. R. Stat. Soc. Ser. Stat. 32, 307–317. https://doi.org/10.2307/2987937

Ansems, G.E., Allen, T.J., Proske, U., 2006. Position sense at the human forearm in the horizontal plane during loading and vibration of elbow muscles. J. Physiol. 576, 445–455. https://doi.org/10.1113/jphysiol.2006.115097

Antcliff, S., Welvaert, M., Witchalls, J., Wallwork, S.B., Waddington, G., 2021. Assessing Proprioception in an Older Population: Reliability of a Protocol Based on Active Movement Extent Discrimination. Percept. Mot. Skills 128, 2075–2096. https://doi.org/10.1177/00315125211029906

Arvin, M., Hoozemans, M.J.M., Burger, B.J., Verschueren, S.M.P., van Dieën, J.H., Pijnappels, M., 2015. Reproducibility of a knee and hip proprioception test in healthy older adults. Aging Clin. Exp. Res. 27, 171–177. https://doi.org/10.1007/s40520-014-0255-6

Berchtold, A., 2016. Test–retest: Agreement or reliability? Methodol. Innov. 9, 2059799116672875. https://doi.org/10.1177/2059799116672875

Bland, J.M., Altman, D.G., 1986. Statistical methods for assessing agreement between two methods of clinical measurement. Lancet Lond. Engl. 1, 307–310.

Bosco, G., Poppele, R.E., Eian, J., 2000. Reference Frames for Spinal Proprioception: Limb Endpoint Based or Joint-Level Based? J. Neurophysiol. 83, 2931–2945. https://doi.org/10.1152/jn.2000.83.5.2931

Bruijn, S.M., Van Impe, A., Duysens, J., Swinnen, S.P., 2012. Split-belt walking: adaptation differences between young and older adults. J. Neurophysiol. 108, 1149–1157. https://doi.org/10.1152/jn.00018.2012

Bullock-Saxton, J.E., Wong, W.J., Hogan, N., 2001. The influence of age on weight-bearing joint reposition sense of the knee. Exp. Brain Res. 136, 400–406. https://doi.org/10.1007/s002210000595

Bunday, K.L., Bronstein, A.M., 2009. Locomotor adaptation and aftereffects in patients with reduced somatosensory input due to peripheral neuropathy. J. Neurophysiol. 102, 3119–3128. https://doi.org/10.1152/jn.00304.2009

Clayton, H.A., Cressman, E.K., Henriques, D.Y.P., 2014. The effect of visuomotor adaptation on proprioceptive localization: the contributions of perceptual and motor changes. Exp. Brain Res. 232, 2073–2086. https://doi.org/10.1007/s00221-014-3896-y

Cressman, E.K., Henriques, D.Y.P., 2010. Reach Adaptation and Proprioceptive Recalibration Following Exposure to Misaligned Sensory Input. J. Neurophysiol. 103, 1888–1895. https://doi.org/10.1152/jn.01002.2009

Cressman, E.K., Henriques, D.Y.P., 2009. Sensory Recalibration of Hand Position Following Visuomotor Adaptation. J. Neurophysiol. 102, 3505–3518. https://doi.org/10.1152/jn.00514.2009

Deshpande, N., Connelly, D.M., Culham, E.G., Costigan, P.A., 2003. Reliability and validity of ankle proprioceptive measures. Arch. Phys. Med. Rehabil. 84, 883–889. https://doi.org/10.1016/s0003-9993(03)00016-9

Dietz, V., 2002. Proprioception and locomotor disorders. Nat. Rev. Neurosci. 3, 781–790. https://doi.org/10.1038/nrn939

Elangovan, N., Tuite, P.J., Konczak, J., 2018. Somatosensory Training Improves Proprioception and Untrained Motor Function in Parkinson's Disease. Front. Neurol. 9.

Emerson, P.L., 1986. Observations on maximum-likelihood and Bayesian methods of forced-choice sequential threshold estimation. Percept. Psychophys. 39, 151–153. https://doi.org/10.3758/BF03211498

Fuentes, C.T., Bastian, A.J., 2010. Where is your arm? Variations in proprioception across space and tasks. J. Neurophysiol. 103, 164–171. https://doi.org/10.1152/jn.00494.2009

Gandevia, S.C., 1985. Illusory movements produced by electrical stimulation of low-threshold muscle afferents from the hand. Brain 108, 965–981. https://doi.org/10.1093/brain/108.4.965

Giavarina, D., 2015. Understanding Bland Altman analysis. Biochem. Medica 25, 141–151. https://doi.org/10.11613/BM.2015.015

Gorst, T., Freeman, J., Yarrow, K., Marsden, J., 2020. Assessing lower limb position sense in stroke using the gradient discrimination test (GradDT^{TM}) and step-height discrimination test (StepDT^{TM}): a reliability and validity study. Disabil. Rehabil. 42, 2215–2223. https://doi.org/10.1080/09638288.2018.1554008

Han, J., Waddington, G., Adams, R., Anson, J., Liu, Y., 2016. Assessing proprioception: A critical review of methods. J. Sport Health Sci. 5, 80–90. https://doi.org/10.1016/j.jshs.2014.10.004

Harris, C.S., 1963. Adaptation to Displaced Vision: Visual, Motor, or Proprioceptive Change? Science 140, 812–813.

Henriques, D.Y.P., Cressman, E.K., 2012. Visuomotor adaptation and proprioceptive recalibration. J. Mot. Behav. 44, 435–444. https://doi.org/10.1080/00222895.2012.659232

Hiebert, G.W., Whelan, P.J., Prochazka, A., Pearson, K.G., 1996. Contribution of hind limb flexor muscle afferents to the timing of phase transitions in the cat step cycle. J. Neurophysiol. 75, 1126–1137. https://doi.org/10.1152/jn.1996.75.3.1126

Hillier, S., Immink, M., Thewlis, D., 2015. Assessing Proprioception: A Systematic Review of Possibilities. Neurorehabil. Neural Repair 29, 933–949. https://doi.org/10.1177/1545968315573055

Horváth, Á., Ferentzi, E., Schwartz, K., Jacobs, N., Meyns, P., Köteles, F., 2022. The measurement of proprioceptive accuracy: A systematic literature review. J. Sport Health Sci. https://doi.org/10.1016/j.jshs.2022.04.001

Ingram, L.A., Butler, A.A., Gandevia, S.C., Walsh, L.D., 2019. Proprioceptive measurements of perceived hand position using pointing and verbal localisation tasks. PLOS ONE 14, e0210911. https://doi.org/10.1371/journal.pone.0210911

Jardine, D.L., Wieling, W., Brignole, M., Lenders, J.W.M., Sutton, R., Stewart, J., 2018. Pathophysiology of the vasovagal response. Heart Rhythm 15, 921–929. https://doi.org/10.1016/j.hrthm.2017.12.013

Jones, S.A.H., Cressman, E.K., Henriques, D.Y.P., 2010. Proprioceptive localization of the left and right hands. Exp. Brain Res. 204, 373–383. https://doi.org/10.1007/s00221-009-2079-8

Juul-Kristensen, B., Lund, H., Hansen, K., Christensen, H., Danneskiold-Samsøe, B., Bliddal, H., 2008. Test-retest reliability of joint position and kinesthetic sense in the elbow of healthy subjects. Physiother. Theory Pract. 24, 65–72. https://doi.org/10.1080/09593980701378173

Kingdom, F.A.A., Prins, N., 2016. Psychophysics: A Practical Introduction. Academic Press.

King-smith, P.E., Rose, D., 1997. Principles of an Adaptive Method for Measuring the Slope of the Psychometric Function. Vision Res. 37, 1595–1604. https://doi.org/10.1016/S0042-6989(96)00310-0

Kontsevich, L.L., Tyler, C.W., 1999. Bayesian adaptive estimation of psychometric slope and threshold. Vision Res. 39, 2729–2737. https://doi.org/10.1016/S0042-6989(98)00285-5

Kottner, J., Audigé, L., Brorson, S., Donner, A., Gajewski, B.J., Hróbjartsson, A., Roberts, C., Shoukri, M., Streiner, D.L., 2011. Guidelines for Reporting Reliability and Agreement Studies (GRRAS) were proposed. J. Clin. Epidemiol. 64, 96–106. https://doi.org/10.1016/j.jclinepi.2010.03.002

Kriellaars, D.J., Brownstone, R.M., Noga, B.R., Jordan, L.M., 1994. Mechanical entrainment of
fictive locomotion in the decerebrate cat. J. Neurophysiol. 71, 2074–2086.
https://doi.org/10.1152/jn.1994.71.6.2074
Kruschke, J., 2014. Doing Bayesian Data Analysis: A Tutorial with R, JAGS, and Stan.
Academic Press.
Kruschke, J.K., 2013. Bayesian estimation supersedes the t test. J. Exp. Psychol. Gen. 142,
573–603. https://doi.org/10.1037/a0029146
Kruschke, J.K., Liddell, T.M., 2018. The Bayesian New Statistics: Hypothesis testing,
estimation, meta-analysis, and power analysis from a Bayesian perspective. Psychon.
Bull. Rev. 25, 178–206. https://doi.org/10.3758/s13423-016-1221-4
Lam, T., Pearson, K.G., 2002. The Role of Proprioceptive Feedback in the Regulation and
Adaptation of Locomotor Activity, in: Gandevia, S.C., Proske, U., Stuart, D.G. (Eds.),
Sensorimotor Control of Movement and Posture, Advances in Experimental Medicine
and Biology. Springer US, Boston, MA, pp. 343–355. https://doi.org/10.1007/978-1-
4615-0713-0_40
Leek, M.R., 2001. Adaptive procedures in psychophysical research. Percept. Psychophys. 63,
1279–1292. https://doi.org/10.3758/BF03194543
Livesey, E.J., Livesey, D.J., 2016. Validation of a Bayesian Adaptive Estimation Technique in
the Stop-Signal Task. PLOS ONE 11, e0165525.
https://doi.org/10.1371/journal.pone.0165525
Lord, S.R., Clark, R.D., Webster, I.W., 1991. Physiological Factors Associated with Falls in an
Elderly Population. J. Am. Geriatr. Soc. 39, 1194–1200. https://doi.org/10.1111/j.1532-
5415.1991.tb03574.x
Lord, S.R., Ward, J.A., 1994. Age-associated differences in sensori-motor function and balance
in community dwelling women. Age Ageing 23, 452–460.
Mattar, A.A.G., Darainy, M., Ostry, D.J., 2013. Motor learning and its sensory effects: time
course of perceptual change and its presence with gradual introduction of load. J.
Neurophysiol. 109, 782–791. https://doi.org/10.1152/jn.00734.2011
McElreath, R., 2016. Statistical Rethinking: A Bayesian Course with Examples in R and Stan.
CRC Press.
McGraw, K.O., Wong, S.P., 1996. Forming inferences about some intraclass correlation
coefficients. Psychol. Methods 1, 30–46. https://doi.org/10.1037/1082-989X.1.1.30
Nicholson, L., Adams, R., Maher, C., 1997. Reliability of a discrimination measure for
judgements of non-biological stiffness. Man. Ther. 2, 150–156.
https://doi.org/10.1054/math.1997.0295
Ostry, D.J., Darainy, M., Mattar, A.A.G., Wong, J., Gribble, P.L., 2010. Somatosensory Plasticity
and Motor Learning. J. Neurosci. 30, 5384–5393.
https://doi.org/10.1523/JNEUROSCI.4571-09.2010
Pearson, K.G., 2004. Generating the walking gait: role of sensory feedback. Prog Brain Res
143, 123–9. https://doi.org/10.1016/S0079-6123(03)43012-4
Pearson, K.G., 2000. Neural Adaptation in the Generation of Rhythmic Behavior. Annu. Rev.
Physiol. 62, 723–753. https://doi.org/10.1146/annurev.physiol.62.1.723
Portney, L.G., Watkins, M.P., 2009. Foundations of clinical research: applications to practice.
Pearson/Prentice Hall Upper Saddle River, NJ.
Prokop, T., Berger, W., Zijlstra, W., Dietz, V., 1995. Adaptational and learning processes during
human split-belt locomotion: interaction between central mechanisms and afferent input.
Exp. Brain Res. 106, 449–456. https://doi.org/10.1007/BF00231067
Proske, U., Gandevia, S.C., 2012. The Proprioceptive Senses: Their Roles in Signaling Body
Shape, Body Position and Movement, and Muscle Force. Physiol. Rev. 92, 1651–1697.
https://doi.org/10.1152/physrev.00048.2011

Rahlf, A.L., Petersen, E., Rehwinkel, D., Zech, A., Hamacher, D., 2019. Validity and Reliability
of an Inertial Sensor-Based Knee Proprioception Test in Younger vs. Older Adults.
Front. Sports Act. Living 1, 27. https://doi.org/10.3389/fspor.2019.00027
Reisman, D.S., Block, H.J., Bastian, A.J., 2005. Interlimb coordination during locomotion: what
can be adapted and stored? J. Neurophysiol. 94, 2403–2415.
https://doi.org/10.1152/jn.00089.2005
Relph, N., Herrington, L., Tyson, S., 2014. The effects of ACL injury on knee proprioception: a
meta-analysis. Physiotherapy 100, 187–195.
https://doi.org/10.1016/j.physio.2013.11.002
Ribeiro, F., Oliveira, J., 2007. Aging effects on joint proprioception: the role of physical activity in
proprioception preservation. Eur. Rev. Aging Phys. Act. 4, 71–76.
https://doi.org/10.1007/s11556-007-0026-x
Rinderknecht, M.D., Lambercy, O., Raible, V., Büsching, I., Sehle, A., Liepert, J., Gassert, R.,
2018. Reliability, validity, and clinical feasibility of a rapid and objective assessment of
post-stroke deficits in hand proprioception. J. NeuroEngineering Rehabil. 15, 47.
https://doi.org/10.1186/s12984-018-0387-6
Roden-Reynolds, D.C., Walker, M.H., Wasserman, C.R., Dean, J.C., 2015. Hip proprioceptive
feedback influences the control of mediolateral stability during human walking. J.
Neurophysiol. 114, 2220–2229. https://doi.org/10.1152/jn.00551.2015
Ruttle, J.E., 't Hart, B.M., Henriques, D.Y.P., 2021. Implicit motor learning within three trials. Sci.
Rep. 11, 1627. https://doi.org/10.1038/s41598-021-81031-y
Salvatier, J., Wiecki, T.V., Fonnesbeck, C., 2016. Probabilistic programming in Python using
PyMC3. PeerJ Comput. Sci. 2, e55. https://doi.org/10.7717/peerj-cs.55
Santuz, A., Laflamme, O.D., Akay, T., 2022. The brain integrates proprioceptive information to
ensure robust locomotion. J. Physiol. https://doi.org/10.1113/JP283181
Schilling, T., Ohlendorf, A., Leube, A., Wahl, S., 2017. TuebingenCSTest – a useful method to
assess the contrast sensitivity function. Biomed. Opt. Express 8, 1477–1487.
https://doi.org/10.1364/BOE.8.001477
Shannon, C.E., 1948. A mathematical theory of communication. Bell Syst. Tech. J. 27, 379–
423. https://doi.org/10.1002/j.1538-7305.1948.tb01338.x
Sigmundsson, H., Whiting, H.T.A., Loftesnes, J.M., 2000. Development of proprioceptive
sensitivity. Exp. Brain Res. 135, 348–352. https://doi.org/10.1007/s002210000531
Silva, G.M., Souto, J.J., Fernandes, T.P., Bonifacio, T.A., Almeida, N.L., Gomes, G.H.,
Felisberti, F.M., Santos, N.A., 2020. Impairments of facial detection in tobacco use
disorder: baseline data and impact of smoking duration. Braz. J. Psychiatry 43, 376–384.
https://doi.org/10.1590/1516-4446-2020-1107
Simani, M.C., McGuire, L.M.M., Sabes, P.N., 2007. Visual-Shift Adaptation Is Composed of
Separable Sensory and Task-Dependent Effects. J. Neurophysiol. 98, 2827–2841.
https://doi.org/10.1152/jn.00290.2007
Soechting, J.F., 1982. Does position sense at the elbow reflect a sense of elbow joint angle or
one of limb orientation? Brain Res. 248, 392–395. https://doi.org/10.1016/0006-
8993(82)90601-1
Sombric, C., Gonzalez-Rubio, M., Torres-Oviedo, G., 2019. Split-Belt walking induces changes
in active, but not passive, perception of step length. Sci. Rep. 9, 16442.
https://doi.org/10.1038/s41598-019-52860-9
Stillman, B.C., McMeeken, J.M., 2001. The role of weightbearing in the clinical assessment of
knee joint position sense. Aust. J. Physiother. 47, 247–253.
https://doi.org/10.1016/S0004-9514(14)60272-5
Strong, A., Srinivasan, D., Häger, C.K., 2021. Development of supine and standing knee joint
position sense tests. Phys. Ther. Sport Off. J. Assoc. Chart. Physiother. Sports Med. 49,
112–121. https://doi.org/10.1016/j.ptsp.2021.02.010

Talbot, L.A., Musiol, R.J., Witham, E.K., Metter, E.J., 2005. Falls in young, middle-aged and
older community dwelling adults: perceived cause, environmental factors and injury.
BMC Public Health 5, 86. https://doi.org/10.1186/1471-2458-5-86
Tsay, J.S., Kim, H., Haith, A.M., Ivry, R.B., 2022. Understanding implicit sensorimotor
adaptation as a process of proprioceptive re-alignment. eLife 11, e76639.
https://doi.org/10.7554/eLife.76639
Tsay, J.S., Kim, H.E., Parvin, D.E., Stover, A.R., Ivry, R.B., 2021. Individual differences in
proprioception predict the extent of implicit sensorimotor adaptation. J. Neurophysiol.
125, 1307–1321. https://doi.org/10.1152/jn.00585.2020
Turpin, A., Jankovic, D., McKendrick, A., 2010. Identifying steep psychometric function slope
quickly in clinical applications. Vision Res., Vision Research Reviews 50, 2476–2485.
https://doi.org/10.1016/j.visres.2010.08.032
van Beers, R.J., Sittig, A.C., Denier van der Gon, J.J., 1998. The precision of proprioceptive
position sense. Exp. Brain Res. 122, 367–377. https://doi.org/10.1007/s002210050525
van Beers, R.J., Sittig, A.C., van der Gon Denier, J.J., 1996. How humans combine
simultaneous proprioceptive and visual position information. Exp. Brain Res. 111, 253–
261. https://doi.org/10.1007/BF00227302
van Beers, R.J., Wolpert, D.M., Haggard, P., 2002. When feeling Is more important than seeing
in sensorimotor adaptation. Curr. Biol. 12, 834–837. https://doi.org/10.1016/S0960-
9822(02)00836-9
Vazquez, A., Statton, M.A., Busgang, S.A., Bastian, A.J., 2015. Split-belt walking adaptation
recalibrates sensorimotor estimates of leg speed but not position or force. J.
Neurophysiol. https://doi.org/10.1152/jn.00302.2015
Vindras, P., Desmurget, M., Prablanc, C., Viviani, P., 1998. Pointing Errors Reflect Biases in the
Perception of the InitialHand Position. J. Neurophysiol. 79, 3290–3294.
https://doi.org/10.1152/jn.1998.79.6.3290
Waddington, G., Adams, R., 1999. Discrimination of active plantarflexion and inversion
movements after ankle injury. Aust. J. Physiother. 45, 7–13.
https://doi.org/10.1016/s0004-9514(14)60335-4
Wasserstein, R.L., Lazar, N.A., 2016. The ASA Statement on p-Values: Context, Process, and
Purpose. Am. Stat. 70, 129–133. https://doi.org/10.1080/00031305.2016.1154108
Watson, A.B., Fitzhugh, A., 1990. The method of constant stimuli is inefficient. Percept.
Psychophys. 47, 87–91. https://doi.org/10.3758/BF03208169
Whelan, P.J., Hiebert, G.W., Pearson, K.G., 1995. Stimulation of the group I extensor afferents
prolongs the stance phase in walking cats. Exp. Brain Res. 103, 20–30.
https://doi.org/10.1007/BF00241961
Wolpert, D.M., Ghahramani, Z., Jordan, M.I., 1995. An internal model for sensorimotor
integration. Science 269, 1880–1882. https://doi.org/10.1126/science.7569931
840
841

Figures and Tables

Figure 1:

846
847
848

Figure 2:

850
851
852

Table 1: Agreement comparison after 50 and 75 trials

	Limits of agreement (mm)	Bland Altman Mean Bias (mm; % in ROPE)	Bland Altman Bias Regression (mm; % in ROPE)	Test 1 vs Test 2 Regression	ICC$_{2,1}$ [95% CI]
PSE (α)					
50 trials	-22.3 to 15.7	-3.2 [-9.2 2.6] (73.5%)	Slope = -0.17 [-0.61 0.29] (26.2%) Intercept = -1.1 [-9.3 7.1] (77.0%)	Slope= 0.91 [0.45 1.40] Intercept = 4.1 [-3.9 12.0]	0.78 [0.43, 0.92]
75 trials	-22.3 to 15.5	-3.3 [-9.0 2.8] (72.4%)	Slope = -0.15 [-0.56 0.28] (29.6%) Intercept = -1.3 [-9.5 6.9] (76.0%)	Slope= 0.92 [0.49 1.38] Intercept = 4.2 [-3.7 12.4]	0.80 [0.48, 0.93]
Uncertainty (β)					
50 trials	-9.3 to 10.2	0.5 [-2.7 3.5] (99.5%)	Slope = 0.03 [-0.65 0.69] (24.3%) Intercept = 0.0 [-12.5 12.2] (60.0%)	Slope= 0.57 [0.03 1.09] Intercept = 7.4 [-2.8 17.4]	0.58 [0.05, 0.85]
75 trials	-10.1 to 11.6	0.7 [-2.8 4.1] (98.8%)	Slope = -0.08 [-0.69 0.58] (24.5%) Intercept = 2.0 [-9.8 13.8] (58.4%)	Slope= 0.65 [0.09 1.19] Intercept = 5.6 [-5.6 15.8]	0.61 [0.05, 0.86]

Figure Captions

Figure 3. Task setup, representative data, and movement direction analysis. (A) AFC task setup. Participants stood on the split-belt treadmill with visual feedback of their legs occluded by a black drape and auditory feedback occluded by noise cancelling headphones. The screen prompted participants to verbally respond to the question "Do you feel like your right or left foot is more forward?" when the test foot stopped at the stimulus location. **(B)** Trial sequence. 1) The test foot was moved to a start position either in front of or behind the stimulus position. Start positions were sampled from one of two normal distributions, depending on the movement direction assigned for that specific trial. 2) The test foot was moved to one of 21 possible stimulus positions between -100 and +100 mm. Tick marks (drawn to scale) represent each possible stimulus location. **(C)** Individual participant data. We reconstructed the psychometric functions from the PSE (α) and uncertainty (β) estimates for Test 1 (blue) and Test 2 (red). The dashed black lines, provided as a reference, represent a PSE of 0 mm. The individual dots represent the participant's response data for each test. **(D)** Movement direction analysis. We measured the mean empirical probability of responding left when the movement direction to the stimulus position was forward (gray) vs backward (black). Dots represent individuals. We also used a Bayesian logistic regression model to analyze the unique impact of each movement direction on participant responses by calculating the difference between the posterior estimates of the movement direction coefficients (one for forward movement, one for backwards) of every participant, yielding a posterior contrast distribution (histogram) which we compared to a region of practical equivalence (ROPE). We note that 99.3% of the posterior distribution is within the ROPE, confirming that responses were not biased by movement direction to the stimulus position.

Figure 4: Point of subjective equality (PSE) and uncertainty estimates have good agreement. (A) Bland-Altman plot for the PSE (α; top) and uncertainty (β; bottom) estimates. The mean of Test 1 and Test 2 for each individual is plotted on the x-axis and the difference between the two tests is plotted on the y-axis. Thus, each dot represents one individual. The limits of agreement are plotted as the dotted lines, and the posterior distributions for the mean biases are plotted on the right sides as histograms. The most probable regression line is the solid line with the shading representing the 95% high density interval (HDI), with the black dashed line at 0 as a reference **(B)** Regression model for PSE and uncertainty. Estimates for Test 2 are plotted against estimates for Test 1 for each individual (dots) with the unity line (black dashed line) provided as a reference for perfect agreement. The most probable regression line is the solid line with the shading representing the 95% HDI.