Title

Greater hippocampal atrophy in developmental amnesia is associated with better recall and working memory performance

Authors/Affiliations

Loïc J. Chareyron[a,b], W.K. Kling Chong[c], Tina Banks[c], Neil Burgess[d], Richard C. Saunders[e], Faraneh Vargha-Khadem[a]*

[a] Cognitive Neuroscience and Neuropsychiatry, Developmental Neurosciences, University College London Great Ormond Street Institute of Child Health, London, UK

[b] Laboratory of Brain and Cognitive Development, Institute of Psychology, University of Lausanne, Switzerland

[c] Developmental Imaging & Biophysics, Developmental Neurosciences, University College London Great Ormond Street Institute of Child Health, London, UK

[d] Institute of Cognitive Neuroscience, University College London, London, UK.

[e] Laboratory of Neuropsychology, National Institute of Mental Health, National Institutes of Health, Bethesda, MD 20892.

Corresponding Author

* Faraneh Vargha-Khadem.
Email: f.vargha-khadem@ucl.ac.uk

Abstract

Despite early-life bilateral hippocampal damage, developmental amnesic (DA) patients exhibit well-developed semantic memory, but severely impaired episodic retrieval. It is unclear, however, whether the residual hippocampus plays a role in encoding and/or retrieval of new information. Here, we report that in DA patients the extent of atrophy in CA-DG subregions and subicular complex is above 40% while the atrophy of the uncus is only moderate (-23%). Importantly, the perihippocampal cortices are unaffected. Paradoxically, patients' recall, Verbal IQ, Working Memory, and Processing Speed scores correlate *negatively* with the volume of the uncus. We propose that in DA patients, the uncus serves as "gatekeeper" providing signals that may inhibit compensatory mnemonic functions by the preserved surrounding cortical areas. Our findings highlight the role of uncus in recall, and document not only the extent, but also the limits of functional plasticity and circuit reorganization in the young brain after early bilateral hippocampal damage.

Introduction

Develomental amnesia

We previously identified a group of patients with hippocampal atrophy who had suffered hypoxic-ischaemic episodes in infancy or childhood and later developed a memory disorder differing from that commonly described in patients with adult-onset amnesia of temporal-lobe origin [1,2]. In the latter cases, the disorder most often takes the form of a global anterograde memory loss (i.e. one affecting both episodic and new post-injury semantic memory, thus severely restricting recall and recognition processes [3–6]). In patients with hippocampal damage of early-

onset, by contrast, the disorder is more limited, being characterized by markedly
impaired episodic memory, recall/recollection, spatial processing and navigation
deficits, but relatively preserved semantic memory, working memory, and
recognition performance. We have labeled this dissociated form of memory
function 'developmental amnesia' (DA) [1,7–10]. Given the early onset of
hippocampal damage, the compensatory reorganization that has occurred in
patients with DA, is likely to be distinct from what is observed in patients with
hippocampal damage acquired in adulthood. The differential maturation of
distinct hippocampal circuits during normal development might underlie the
differential emergence of specific hippocampus-dependent memory processes [11].
In patients with adult-acquired hippocampal lesions, the damage has occurred to
already-established, and normally-functioning memory circuits, whereas in
patients with DA, the early bilateral lesions have probably led to the development
of a differently-organized memory system. For these reasons, it is difficult, and
somewhat inappropriate, to compare memory function and cognitive profiles in
patients with DA to those of patients with adult-onset amnesia [12].
To date, a cohort of 18 patients with DA has been reported, each patient
having suffered an episode of hypoxia-ischaemia early in life [2]. Importantly,
irrespective of the underlying aetiology [13], or age at onset [14], the hypoxic-
ischaemic event was the cause of hippocampal damage and cognitive memory
impairment in each case[15] .

Memory circuit reorganization
Among the many questions that have arisen from the study of patients with DA,
the foremost concerns the regions of the medial temporal lobe (MTL) and/or
preserved hippocampal subregions that could be involved in memory circuit
reorganization following early hippocampal damage.
In the model of hierarchical organization of cognitive memory proposed
by Mishkin and colleagues [15], recollection and familiarity depend on different
MTL structures [16,17]. According to this model, the perihippocampal gyrus, and in
particular the perirhinal cortex, may support familiarity, while the hippocampus
may be critical for recollection [18]. The selectivity of impairment observed in
patients with DA has provided strong support for this dual-process model [12,16].
Because patients with DA have both neuroradiological and quantitatively-
confirmed bilateral hippocampal atrophy, but no visible abnormality within the
parahippocampal region, it has been proposed that their episodic memory
impairment, and recall/recollection deficits are due to their hippocampal
pathology, whereas their relatively intact semantic memory, and
recognition/familiarity processes are due in part to the preservation of the
perihippocampal gyrus.
Hippocampus
Hypoxia-ischemia is known to differentially affect the hippocampal fields and
subdivisions. Studies of animal models of cerebral ischemia have reported that a
brief episode results in selective neuronal death in the CA1 field of the
hippocampus [19,20] while the adjacent CA3 remains less vulnerable [21]. In humans,
while damage to CA1 is consistently reported following cerebral ischemia,
damage to additional hippocampal regions is much more variable [22]. Postmortem
examinations of patients who suffered ischemic episodes in adulthood have
reported neural cell loss either restricted to the CA1 region, or distributed in the

dentate gyrus, subiculum, uncus and entorhinal cortex [23,24]. The preservation of
circumscribed hippocampal subregions following injury of early-onset in DA
could account for the increased activation of the residual hippocampus during
different memory tasks [25,26], although activation *per se* does not necessarily imply
correspondingly intact function [27].
In the study reported by Dzieciol and colleagues [2], the degree of
hippocampal atrophy in patients with DA ranged from 28% to 62% compared to
healthy controls. This large range of hippocampal volume reduction in this cohort
suggests that there may indeed be variability in the hippocampal response to
hypoxia-ischemia at the level of hippocampal subregions. It would thus be
possible to assess whether the compensatory response in patients with DA is
dependent on the extent of atrophy in the hippocampal subregions. One way to
address this is to evaluate the association between task-dependent
neuropsychological performance and volumes of hippocampal subregions.
Perihippocampal gyrus
While postmortem examinations have reported some neural cell loss in layers of
entorhinal cortex in patients who suffered ischemic episodes in adulthood [24], it
was reported that the volume of the parahippocampal gyrus was not significantly
reduced in patients with adult-onset amnesia presenting with bilateral
hippocampal damage [28]. A cortical thickness analysis performed in a single
patient with DA didn't find abnormality in the subhippocampal structures within
the medial temporal lobe, including the perirhinal and ventral entorhinal cortices
[29]. Also, a recent study demonstrated that parahippocampal activity during scene
reinstatement in patients with DA was similar to controls [27]. However, the
structural integrity of the perihippocampal gyrus, and by implication, the integrity
of the hypothesized substrate of semantic memory and other preserved mnemonic
processes remain to be quantitatively assessed in patients with DA.
119
Here, we studied the largest cohort of patients with the rare condition of
developmental amnesia ever assembled to test the impact of plastic changes and
circuit reorganization in the compensation of hippocampal function. We have used
manual segmentation of MRI scans to obtain estimates of the volume of three
hippocampal subregions: the uncus (uncal portion of the hippocampus) [30,31], the
CA-DG (including CA fields and dentate gyrus) and the subicular complex
(including subiculum, presubiculum and parasubiculum), and three surrounding
cortical areas (entorhinal, perirhinal and parahippocampal cortices) in patients
with DA and healthy controls. Controls and patients were assessed with
neuropsychological tests of intelligence (WISC or WAIS), and recall/recognition
for verbal and visual material (Doors and People Test). Finally, we investigated
the relationships between degree of atrophy of hippocampal subregions and
cognitive deficits.
We predicted that: (i) residual hippocampal subregions and/or surrounding
cortical areas were sufficiently preserved to rescue some aspects of hippocampal
function, and (ii) plastic changes in these regions occurred as a function of extent
of hippocampal damage in all patients with DA.
We reported various level of atrophy of the hippocampal subregions and
an absence of structural damage in the surrounding cortical areas in patients with
DA. Also, we documented several paradoxical inverse relationships between the

volume of the residual uncus and patients' recall memory performance, Verbal
Comprehension Index, Working memory Index and Processing Speed Index.

142   Results

143   *Volumetric estimates*
144   Three hippocampal subregions (uncus, CA-DG and subicular complex) and three
145   surrounding cortical areas (entorhinal, perirhinal and parahippocampal cortices)
146   were manually segmented in healthy controls (N = 32) and DA (N = 23) groups
147   (Fig. S1-S2).
148

Fig. 1. Volume of the hippocampus and its subregions. Volume of the hippocampus (**A.**)
and its subregions (**B.**, uncus, CA-DG, and subicular complex) in control (grey; N = 32) and
DA (orange; N = 23) groups. Data are represented as mean ± SD. Percentage values refer to
the mean difference to control group. Volumes are calculated as the average of left and right
hemisphere volumes. All volumes are corrected for intracranial (ICV) volume. *: P < .05;
***: P < .001.

The volume of the hippocampus as a whole was 40% smaller in the DA than in
the control group (range: −26% to −53%; $t_{(53)} = -15.32$; P < .001) (Fig. 1; Table
S1). The volume of the hippocampal subregions was also significantly smaller in
the DA than in the control group (2-way ANOVA, $F_{(1,159)} = 392.5$, P < .001). The
uncus was 23% smaller (range: +4% to −49%; Tukey's HSD post-hoc test: P =
.037), the CA-DG subregions were 45% smaller (range: −28% to −59%; P <
.001), and the subicular complex was 41% smaller (range: −22% to −56%; P <
.001). The level of atrophy of the patients' hippocampal subregions differed
significantly from each other ($F_{(2,66)} = 23.0$, P < .001). Tukey's HSD post-hoc test
indicated that the CA-DG and subicular complex showed greater atrophy than the
uncus (both P < .001).
Atrophy of the hippocampus in the DA group was significantly lower in its
anterior (−36%) than its posterior (−44%) segment (paired t test, $t_{(22)} = 2.72$, P =
.012; Fig. S3). Because no differences were observed between the anterior and the
posterior segment of the CA-DG (paired t test, $t_{(22)} = 0.94$, P = .36) or subicular
complex ($t_{(22)} = 1$, P = .33) subregions, the lower level of atrophy observed in the

anterior part of the whole hippocampus can be attributed to the relatively more
preserved uncus.
In contrast, there was no significant difference in the volume of the entorhinal,
perirhinal and parahippocampal cortices between control and DA groups (Table
S1; 2-way ANOVA, $F_{(1,159)}$ = 6.7, P = .010; Tukey's HSD post-hoc tests, all P >
.35). Also, we didn't observe any correlation between the volume of the
hippocampus or hippocampal subregions and the volume of surrounding cortical
areas in patients with DA (Pearson's correlation analysis, all P > .19).

Neuroradiological assessment
Neuroradiological ratings based on visual inspection of the MR images provided
independently by one of the authors (WKC) revealed abnormally small fornix and
mammillary bodies in all but two patients with DA (Table S2). No abnormalities
were detected in the parahippocampal gyrus (perirhinal, entorhinal, and
parahippocampal cortices), or the thalamus, and the basal ganglia. Additional
visible abnormality was observed in some cases in the white matter and the
cerebellum. The pattern of visible abnormality detected within the hippocampal
circuit, and the notable absence thereof in other cortical and subcortical structures,
was consistent across individual patients irrespective of their varying aetiology,
and age at hypoxic-ischaemic-induced hippocampal damage.

Doors and People Test
All patients with DA (N = 23) and 21 of the control participants were assessed on
the Doors and People Test, which provides equated measures of recognition and
recall in the visual and verbal domains. All patients' subtest scores were
significantly below zero (Wilcoxon signed rank test: all Z < −2.4, all P < .016),
indicating that patients were impaired relative to controls (who represent zero, as
their scores were used to convert raw scores into z-scores). The patients' scores
for the four Doors and People subtests differed significantly from each other
(Friedman's test: $\chi^2_{(3,n=22)}$ = 43.7, P < .001). The visual recall ('Shapes') subtest
yielded a significantly greater deficit than all of the other subtests (all P < .001).
The degree of deficit on the verbal recall ('People') subtest was greater than on
the two recognition subtests (both P < .02). The degree of deficit was greater on
the visual recognition ('Doors') subtest than on the verbal recognition ('Names')
subtest (P = .001). Collapsing across subtests, we found a greater deficit in recall
compared with recognition (Z = −4.04; P < .001), and a greater deficit in memory
for visual compared with memory for verbal material (Z = −4.11; P < .001).
We observed significant negative correlations between the two recall
subtest scores and the volume of the uncus in the DA group [Verbal recall
('People'): Pearson's $r_{(22)}$ = –0.47; P = .02; Visual recall ('Shapes'): $r_{(22)}$ = –0.55;
P = .007; Fig.2AB]. No other correlations for other subtests or other regions of
interest in the DA group were significant (all P > .09; Table S3). Collapsing
across test material confirmed that scores on recall were inversely correlated with
the volume of the uncus in the DA group ($r_{(22)}$ = –0.58; P = .004; Fig. 2C). As a
corollary, we observed that patients with more preserved uncus had significantly
lower scores on recall than patients with more damaged uncus (Z = −2.74; P =
.006). In the control group, no correlations for any subtests or regions of interest
were significant (all P > .11; Table S3).

Fig. 2. *Doors and People* **recall scores correlations with the volumes of the uncus in patients with DA.** The patients' scores (N = 23) on recall subtests of the *Doors and People Test* correlated negatively with the volume of the uncus. **A.** Verbal recall ('People') subtest. **B.** Visual recall ('Shapes') subtest. **C.** Collapsed scores. Pearson's correlation coefficient. All volumes are corrected for intracranial (ICV) volume. *: P < .05; **: P < .01.

During *Doors and People* testing, the procedures for recall subtests are repeated until all items (names or shapes) are correctly recalled, or for a maximum of three presentations. Delayed (cued) verbal and visual recall were tested after completion of the verbal and visual recognition subtests, respectively. Collapsed recall scores were inversely correlated with the volume of the uncus in the DA group for all of the three trials as well as for the delayed recall (all $r_{(22)} < -0.43$; all P < .05; Fig. S4)

The patients' scores differed significantly from each other between the three trials of the visual recall ('Shapes') subtest (Friedman's test: $\chi^2_{(2,n=20)} = 8.00$, P = .018) and between the three trials of the verbal recall ('People') subtest ($\chi^2_{(2,n=20)} = 14.15$, P < .001). Post hoc Wilcoxon signed-ranks tests indicated that scores obtained on trial-1 were significantly lower than scores obtained on trial-3 for the visual recall ('Shapes') subtest (Z = −2.72; P = .006) and for the verbal recall ('People') subtest (Z = −2.88; P = .004). Interested to determine potential differences in learning between patients depending on whether they presented with severe or moderate uncus atrophy, we compared increases in patients' scores between trial-1 and trial-3. There was no difference in learning scores between patients with a larger uncus (above group mean volume) and patients with a smaller uncus (Mann-Whitney U test; visual recall ('Shapes'): Z = −1.26; P = .21; verbal recall ('People'): Z = 0.61; P = .54).

We found a correlation between patients' scores for the delayed verbal recall ('People') subtest and the volume of the uncus (Pearson's $r_{(20)} = -0.5$; P = .02) while the correlation with delayed visual recall ('Shapes') scores was not significant ($r_{(20)} = -0.3$; P = .18).

WISC/WAIS
All patients with DA (N = 23) and 23 of the control participants were assessed on the WISC or WAIS. Patients were impaired relative to our group of controls only for the Perceptual Reasoning Index scores (2-way ANOVA, $F_{(1,176)} = 10.61$, P = .001; Tukey's HSD post-hoc test, P = 0.04; all other P > .25). There were significant inverse correlations between the volume of the uncus and patients'

scores for the Verbal Comprehension Index (VCI), the Working Memory Index (WMI) and the Processing Speed Index (PSI) (all $r_{(22)} < -0.42$; all $P < .05$; Fig 3; Table S4). No other correlations for other ROIs in the DA group were significant. In controls, the WMI scores were positively associated with the volume of the subicular complex ($r_{(22)} = 0.46$; $P = .029$).

Fig. 3. Inverse correlations between WISC/WAIS indexes scores and the volume of the uncus in patients with DA. Patients' scores on Wechsler Intelligence Scales WISC or WAIS (N = 23). **A.** Verbal comprehension Index; **B.** Perceptual Reasoning Index; **C.** Working Memory Index; **D.** Processing Speed Index. Pearson's correlation coefficient. All volumes are corrected for intracranial (ICV) volume. *: $P < .05$.

Discussion

In this study, we addressed the question of whether the volumes of residual hippocampal subregions and parahippocampal areas are associated with cognitive memory performance in patients with hypoxic-ischemic-induced hippocampal damage of early-onset. Using a combination of neuropsychological test results and volumetric estimations in a large cohort of patients with developmental amnesia (DA), we documented several inverse relationships between the volume of the residual uncus and patients' performance on recall memory, Verbal comprehension Index, Working memory Index and Processing speed Index. The effects reported here suggest that in patients with DA, the uncus, either via its connections with the contralateral uncus or with other anterior temporal lobe areas, and depending on its extent of damage, provides signals that may inhibit compensatory mnemonic development by the preserved surrounding cortical areas. This process could occur either because normally silent connections promote compensation, or because signals (either typical or pathological) inhibit compensation. However, whatever the extent of hippocampal damage in DA, and despite these compensatory reorganizations, hippocampal-dependent episodic memory recall remains irreversibly compromised.

Limitations
Although manual tracing is still the gold standard for measuring the volumes of hippocampal subregions and cortical areas on MRI acquisitions, especially in the case of patients, manual segmentation is subjective and susceptible to measurement errors. Also, the precision of the measure is limited by the resolution

of the MRI scans. However, analyses of MR images for estimating the extent of
hippocampal damage in amnesic patients have been validated by postmortem
examinations [24]. In particular, a clear concordance has been reported in the
appearance of the damaged hippocampus on the MRI and in the histological
sections [24]. The border between cytoarchitectonic hippocampal subregions were
estimated based on delineation of these regions on histological sections [32]. The
delineation on MR images of the perihippocampal and parahippocampal cortical
areas were based on that described in [33]. It is not clear to what extent these borders
would be affected in the brains of DA patients with significant hippocampal
volume reduction. Our observations of an absence of significant volume reduction
of the entorhinal, perirhinal and parahippocampal cortices in patients with DA is
consistent with previous report showing an absence of significant volume
reduction in patients with adult-onset amnesia [28], and in a patient with DA [29].
Also, intact parahippocampal activity during scene reinstatement has been
described in patients with DA [27].

Differential effect of early hypoxia-ischemia on hippocampal subregions
Our observation that the CA-DG were the most affected hippocampal subregions
in patients with DA is congruent with documented reports of greater sensitivity of
this subregion to hypoxia-ischemia. It has long been known that hypoxia and
ischemia result in selective neuronal death in the CA1 field [19], although the reason
for this selectivity is still debated [21]. Neurohistological studies of patients who
suffered ischemic episodes in adulthood have reported cases with neural cell loss
restricted to the CA1 region bilaterally whilst the CA1' region of the uncus, the
subiculum, CA3 and surrounding cortices were all preserved [23,24]. High-resolution
magnetic resonance imaging showed that CA1 was always affected in patients
with hippocampal damage [22].
In contrast, the uncus showed a low sensitivity to hypoxia in our patient cohort
which could be a reflection of the distinct vascularization of this hippocampal
subregion. The branches of the posterior cerebral artery (PCA) coming from the
vertebral artery irrigate the posterior part of the hippocampus, while the branches
of the anterior choroidal artery (AchA) originating from the internal carotid artery
irrigate the uncal portion of the hippocampus [34–36]. High-resolution angiography
and anatomical studies have highlighted that the anterior hippocampal region
could receive a mixed blood supply (from both the PCA and the AchA) in about
half of the individuals [34,37]. A mixed blood supply could provide more vascular
reserve and lower vulnerability to neuronal injury and atrophy [38]. The uncus,
which receives a distinct blood supply compared to the rest of the hippocampus,
might thus exhibit greater resistance to the adverse effects of hypoxic-ischemic
events than the other hippocampal regions.

Uncus and recall memory
The residual uncus could be partly functional in patients with DA and could be
recruited for recall memory. We reported here that the volume of the uncus
correlated negatively with recall scores as well as working memory performance
in patients with DA. In the WAIS/WISC test, Working Memory Index is made up
of two subtests (Digit Span and Letter Number Sequencing) both requiring verbal
recall within the span of working memory. The function of the uncus is not well
understood mainly because its structure-function mapping remains to be

determined in rodents [39], but in humans, the uncus has been proposed to be
involved in the long-term recall of scenes [40]. Connectivity studies in nonhuman
primates have shown that the uncus is less connected to other regions of the
hippocampus, and that it has its own associational system of connections [41].
Notably, the uncus is also the only hippocampal region reciprocally connected to
its contralateral counterpart [42]. The uncus' unique connectivity could allow this
region to support some information processing despite severe damage to the other
hippocampal areas.
Notwithstanding the importance of the uncus in relation to severity of
atrophy during delayed recall, this subregion maintains its role during encoding
and learning of memoranda in both low and high severity subgroups equally.
Thus, we noted that despite their poor performance overall, both patient subgroups
incrementally increased their scores across the three learning trials of the *People*
and *Shapes* subtests of the *Doors and People Test*. The absence of a difference in
performance between the subgroups during encoding and learning, but not during
delayed recall, highlights the role of the uncus in retrieval of auditory verbal
memory after a delay in patients with DA.
Interestingly, we also reported an inverse relationship between the volume of
the uncus and patients' performance in delayed verbal, but not visual recall.
Impairment in delayed verbal recall in relation to extent of atrophy of the uncus
highlights the vulnerability of auditory verbal information to recall after a delay [43].
The above, together with ontogenetic lateralization of verbal output via the left
hemisphere language network, and the unique reciprocal connection of the uncus
to its contralateral counterpart, could account for the 'gatekeeping' role that the
uncus might play in verbal recall.

Working memory
We reported here, in addition to an absence of working memory impairment, a
negative association between uncus volume and working memory performance in
patients with DA. The more established view proposes that the hippocampus plays
an important role in episodic memory, spatial cognition, and navigation, while it
is not critically contributing to working memory [44,45]. But an alternative view
proposes that the hippocampus may contribute to any memory task, including
working memory [46,47], in particular in supporting high-precision binding [48]. This
latter possibility is challenged however by the fact that patients with DA
presenting with severe hippocampal damage have repeatedly shown unimpaired
working memory performance [49–51]. Recent work extended these findings by
showing that high resolution working memory performance was unimpaired, and
even often superior, relative to control group in a patient with DA [52]. Our
observation of a negative correlation between uncus volume and working memory
performance suggests that the perirhinal, entorhinal, and parahippocampal cortices
normally involved in the binding of information in the short term [53] may be
subject to plastic changes in patients with DA and that these plastic changes may
depend on the degree of hippocampal damage. The possibility that the
hippocampal damage in patients with DA may not influence working memory
performance directly, but indirectly via reorganization of the MTL circuit is
explored below.

Functional hypotheses

Experimental studies in nonhuman primates have reported similar paradoxical findings to those reported in the present study. Monkeys with small hippocampal lesions exhibit greater memory loss on a Delayed Nonmatching-to-Sample (DNMS) task than that found in animals with greater damage [54]. Furthermore, a positive correlation between the extent of damage to the hippocampus and scores on a DNMS task has been reported in experimentally lesioned monkeys [55]. Several hypotheses have been proposed to explain why discrete lesions result in greater memory impairment [54,56,57]. However, it is difficult to relate these findings obtained following adult experimental lesions in the monkey to observations in patients with early hippocampal damage whose circuits might have undergone massive compensation. One hypothesis to explain our observations is that the level of hippocampal damage would likely have a strong direct influence on the organization of the medial temporal lobe memory circuits in the adult. Following an early event of hypoxia-ischemia, however, greater hippocampal damage might induce greater compensatory reconfigurations in the neural circuits and enable other structures, in particular the surrounding cortical areas normally recruited by the hippocampus, to assume important aspects of memory function. In contrast, when the hippocampus is only partially damaged, the information flow would remain present in the hippocampus and could result in incomplete, and possibly disruptive information processing. Our observation could thus reveal the existence of multiple, redundant routes within the residual hippocampal subregions and surrounding cortical areas in patients with developmental amnesia [58]. These parallel circuits could compete for the control of behavior and disrupt memory performance [57].

Our findings are consistent with the hypothesis that a close relationship can be evidenced between the volume of remaining hippocampal subregions and the severity of cognitive deficits in patients with hippocampal damage, in the case of patients with DA this relationship was inverse. Our observations also provide further support for the model of hierarchical organization of cognitive memory [15] by proposing that surrounding cortical areas normally involved in the encoding of associative relations between decontextualized information [53] could reorganize to contribute to hippocampal function following early hippocampal damage. The limits of this reorganization, however, may be determined by the extent of damage to the hippocampal subregions.

Evidence for such reorganization has already been reported in a nonhuman primate model of developmental amnesia. Monkeys with early bilateral hippocampal lesion can learn new spatial relational information in striking contrast to monkeys with adult lesion [59,60]. Interestingly, in these monkeys with early hippocampal lesions, the surrounding cortical areas were preserved and were shown to contribute to spatial learning in the absence of functional hippocampal circuits [61]. A recent report showed that the number of immature neurons in the entorhinal and perirhinal cortices was higher in monkeys with early hippocampal lesions than in controls [62]. This increase in the number of immature neurons was paralleled by an increase in the number of mature neurons in the entorhinal cortex of neonate-lesioned monkeys compared to controls. These structural changes may contribute to some functional recovery after early hippocampal injury [62].

One could speculate that the functional reorganization of the perihippocampal structures (normally devoted to the processing of

decontextualized information) to compensate hippocampal function could, in turn,
benefit the processing of semantic memory and lead to better context-free memory
performance in patients (see: [63]). Indications of increased memory performance
have indeed been reported in two different patients with DA. Jonin et al. [29] studied
patient 'KA' presenting with 55% hippocampal volume reduction and preserved
perihippocampal structures. While patient KA displayed few, if any, residual
episodic abilities, this patient was able to accurately retrieve semantic memories,
and show evidence of superior or even very superior access to these memories
compared to controls. Another patient with DA, patient 'Jon', has often produced
response accuracy levels that are at least higher than those of control participants
in previous experimental explorations of visuospatial working memory [49,51,52]. Jon
was at least numerically superior to the control mean on recognition-based
measures of shape-color binding [51]. In measures of color-location memory, Jon's
recognition accuracy matched the highest achieving control participant, while his
reconstruction performance was superior to 6 of the 7 controls [49]. Across four
tasks measuring binding between color and orientation, or color and location
using simultaneous or sequential presentation of stimuli, Jon's response accuracy
was high, and always numerically superior to the control mean [52].
These observations from single patients studies together with the present
findings, obtained in a large cohort of patients with DA, support the hypothesis of
a reorganization of the preserved perihippocampal areas following early
hippocampal damage.

Conclusions

Our results have important implications for understanding the organization of the
memory circuit following early hippocampal injury. These observations highlight
the paradoxical finding that in patients with DA greater hippocampal damage can
be associated with better performance in recall memory, Verbal comprehension
Index, Working memory Index, and Processing speed Index. In contrast, less
severe damage, leaving behind relatively preserved hippocampal subregions, can
lead to more pronounced cognitive deficits. These unexpected findings can
disentangle the functional organization of the medial temporal lobe memory
system, and attribute a central role in the compensation of cognitive memory to
extra-hippocampal structures in patients with early hippocampal damage.

Materials and methods

Participants
The patients with DA were characterized by (i) a history of episodes of hypoxia-
ischemia in early life, (ii) quantified hippocampal volume reduction above 25%
relative to a group of controls, and (iii) severely impaired episodic memory [2]. The
most common aetiology was perinatal hypoxic-ischemic encephalopathy (Table
S2). While long-term neurodevelopmental outcome depends on the severity of the
hypoxic-ischemic episode, subtle cognitive deficits and behavioral alterations
have been revealed through long-term evaluations even in mild forms of hypoxia-
ischemia [64].
When all the volumetric estimates were completed, one of the patients,

who presented with moderate memory deficits and only 11.0% hippocampal
volume reduction associated with two unconfirmed episodes of prolonged seizures
at age 4, was excluded from the DA group.
Also, we excluded from the DA group a patient with temporal lobe
epilepsy presenting with a 25.8% hippocampal volume reduction, but without a
documented episode of early hypoxia-ischemia and/or status epilepticus.
Additionally, blinded neuroradiological examination carried out independently by
one of the authors (WKC) revealed that this patient's MRI scan was entirely
normal. The hippocampus, fornix and mammillary bodies were noted to have a
normal appearance in contrast to all other patients with DA.
The DA group included three patients (DA05, DA06, DA10) who had
sustained hypoxic–ischemic episodes between the ages of 9 and 12 years in
contrast to other patients who had suffered such episodes perinatally or within the
first 3 months of life. However, it has been previously demonstrated that the early
(below age one), and late (between 6-14 years) groups exhibit similar profiles, and
thus concluded that the effective age at injury for developmental amnesia extends
from birth to puberty [14].
We report here hippocampal volume estimates in 32 control participants
(16 females; mean age: 17.7 years, SD 7.7, range 8−38) and 23 patients with DA
(11 females; mean age: 19.5 years, SD 9.0, range 8−40).
498
The study was approved by the London-Bentham Research Ethics Committee
(REC Reference No. 05/Q0502/88) and all participants and their carers read an
age-appropriate information sheet and gave written informed consent according to
the Declaration of Helsinki, before commencing the study.

Structural MRI
Participants were either scanned with a 1.5T-MRI scanner (24 control
participants, 16 patients with DA) or with a 3T-MRI scanner (8 control
participants, 7 patients with DA).
Whole brain structural 1.5T-MRI scans were obtained using a Siemens
Avanto Scanner, with a T1-weighted 3D FLASH sequence with the following
parameters: repetition time 11ms; echo time 4.94ms; in-plane resolution
1mm×1mm; slice thickness of 1mm [65].
Whole brain structural 3T-MRI scans were obtained using a 3T Siemens
MRI system with a 20 channel head coil. A T1-weighted magnetization-prepared
rapid gradient-echo (MPRAGE) scan was acquired with the following parameters:
in-plane resolution of 1mm×1mm; slice thickness of 1mm; repetition time of
2,300ms; echo time of 2.74ms [66].
For the measurement of regions of interest (ROI) volumes, the data were
reformatted into 1 mm-thick contiguous slices in the coronal plane. ROI cross-
sectional areas were measured in all slices along the entire length of the
hippocampus, entorhinal, perirhinal and parahippocampal cortices using ITK-
SNAP (version 3.8.0; www.itksnap.org) [67]. Manual tracing is still the gold
standard for measuring hippocampal volume and is particularly appropriate for
samples of relatively small size [68], presenting with various levels of hippocampal
atrophy [69], or including children and adolescents [70]. All manual measurements of
the ROIs were carried out by one of the authors (LJC) who remained blind to
participant identity (Fig. S1-S2).

The hippocampus comprises the CA fields (CA3, CA2, CA1), the dentate
gyrus (DG), subiculum, presubiculum, and parasubiculum [71]. The human uncus
[also termed the "uncal portion of the hippocampal formation" [42]] is situated
antero-medially and consists of slightly modified DG, CA3, CA2, CA1, and
subiculum [72]. For the segmentation of the hippocampus and its subregions, we
followed the guide provided by Dalton et al. [73]. Given the difficulty to distinguish
hippocampal CA fields on 1.5T and 3T-MRI acquisitions, we restricted our
volumetric analyses to three ROIs: uncus [defined according to [72]], CA-DG (CA
fields and dentate gyrus), and subicular complex (subiculum, presubiculum, and
parasubiculum). The segmentation of the entorhinal, perirhinal and
parahippocampal cortices was based on structural descriptions of human medial
temporal lobe and segmentations protocols [33,74–77].
The volumes in cubic millimeters were calculated by summing the number
of voxels ($1\times1\times1$ mm) for each ROI. The volume of the hippocampus as a whole
was calculated as the sum of hippocampal subregions' volumes (uncus, CA-DG,
subicular complex). We then used parametric tests as volumetric estimates of the
hippocampal subregions and cortical areas were normally distributed (Shapiro-
Wilk normality test; control: all $P > .06$; DA: all $P > .09$). There was no
significant difference between volumetric estimates of the whole hippocampus or
cortical areas obtained with the 1.5T-MRI scanner and the 3T-MRI scanner in
control and DA groups (two-way ANOVA, both $F < 1.3$; both $P > .25$). Also there
was no significant gender difference for the volumes of the whole hippocampus or
cortical areas in control and DA groups (both $F < 0.08$; both $P > .79$). Volume
estimates on 1.5T and 3T-MRI scans and volume estimates in males and females
were thus presented together. There was a significant right larger than left volume
asymmetry for the volume of the whole hippocampus in both control and DA
groups (repeated measures two-way ANOVA: $F_{(1,53)} = 19.2$; $P < .001$; Tukey's
HSD post-hoc test: both $P < .004$). The right CA-DG subregions were
significantly larger than the left in both control and DA groups ($F_{(1,53)} = 26.1$; $P <$
$.001$; both $P < .004$). There was no significant asymmetry for the volume of the
other hippocampal subregions (all $P > .05$). Also, there was no significant
asymmetry for the volume of the entorhinal or perirhinal cortices (all $P > .12$) but
there was a significant right larger than left volume asymmetry for the volume of
the parahippocampal cortex in the control group ($F_{(1,53)} = 7.0$; $P = .011$; Tukey's
HSD post-hoc test: $P = .009$). The volume of all the ROIs was calculated as the
mean of the left and right hemisphere measurements.
Intracranial volume (ICV) was manually measured on the sagittal dataset with
a 1-in-10 random and systematic sampling strategy. The ICV was normally
distributed (Shapiro-Wilk normality test; control: $P = .05$; DA: $P = .23$). The ICV
was 6.8% smaller in the DA group than in the control group ($t_{(53)} = -2.70$; $P =$
$.009$). Corrections, derived from the regression line of control ROI volume (V)
versus ICV, were then made for ICV based on the formula [$\Delta V = a\times(\Delta ICV)$] where
'a' is the regression coefficient.

WISC/WAIS
The Wechsler Adult Intelligence Scale, 3rd Ed. (WAIS-III) and the Wechsler
Intelligence Scale for Children, 4th Ed. (WISC-IV) provide four index scores:
Verbal Comprehension Index, measuring understanding and expression of verbal
concepts; Perceptual Reasoning Index, reflecting visuospatial perception and

perceptuomotor manipulation; Working Memory Index, tracking on-line
immediate memory and executive control; and Processing Speed Index,
measuring speed of visuoperceptual discrimination, and copying of number-
symbol associations. The four index scores are expressed as standard scores with a
mean of 100 and a standard deviation of 15.
Scoring. The WISC-IV or WAIS-III were administered to all patients with DA (N
= 23, 11 females, mean age: 19.5 years, SD: 9.0, range: 8−40) and to 23 controls
(13 females, mean age: 18.0 years, SD: 8.4, range: 8.2–38). For the analysis of the
data, we then used parametric tests as the data were normally distributed in all
four indexes (Shapiro–Wilk test of normality; Control: all P > .36; DA: all P >
.10).

The Doors and People Test
The *Doors and People Test* was administered according to the instructions in the
published manual [78], as described in detail in [8,79]. Briefly, for the verbal recall
subtest ('People'), participants were presented with four pictures of people and
after three learning trials asked to recall their names cued by their profession. For
the visual recall subtest ('Shapes'), participants first copied, and drew the shapes
from memory after three learning trials. For verbal recognition ('Names'),
participants were presented with a list of names, and then asked to recognize each
one from a list of four alternatives. Finally, in the visual recognition subtest
('Doors'), participants viewed a list of doors and then identified each one from a
list of four alternatives. See Supplementary Information for more details.
Scoring. The *Doors and People Test* was administered to all patients with DA (N
= 23, 11 females, mean age: 19.5 years, SD: 9.0, range: 8−40) and to 21 controls
(11 females, mean age: 17.6 years, SD: 8.6, range: 8−38). One of the patients
performed all but the 'Names' subtest ('People', 'Doors', 'Shapes': N = 23;
'Names': n = 22). Given that the DA group included individuals under the age of
16, which is the first age band at which standard scores are available on the *Doors*
and People Test, we used standard scores based on the scores of our control
group; raw scores (SD): People, 28.2 (6.9); Doors, 18.9 (3.8); Shapes, 34.6 (2.7);
Names, 18.6 (3.7). We converted patients' and controls' raw scores to z-scores
relative to the control group's scores. For the analysis of the data, we then used
nonparametric tests as the data were not normally distributed in all four subtests
(Shapiro–Wilk test of normality; Control: P < .001; DA: P = .044).

Statistical analysis
We followed the recommendations of Rothman [80], arguing that "not making
adjustments for multiple comparisons is preferable because it will lead to fewer
errors of interpretation when the data under evaluation are not random numbers
but actual observations on nature," and Saville [81], who argued that a procedure
without correction is preferable because it provides greater consistency to
compare results between studies.

Acknowledgments

This work was supported by the UK Medical Research Council (program grant
numbers G03000117 and G1002276).

We thank the late Mortimer Mishkin, Laboratory of Neuropsychology NIMH,
Bethesda, MD, for his unfailing inspiration, encouragement and support enabling
the realization of this research. We also thank the participants and their families
for their support of our research.
623

Data availability

625 Anonymized participant data will be available upon reasonable request.
626

Competing interests

628 The authors declare no conflict of interest.

References

1.   Vargha-Khadem, F. *et al.* Differential effects of early hippocampal
pathology on episodic and semantic memory. *Science* **277**, 376–380 (1997).
2.   Dzieciol, A. M. *et al.* Hippocampal and diencephalic pathology in
developmental amnesia. *Cortex* **86**, 33–44 (2017).
3.   Bayley, P. J. & Squire, L. R. Failure to acquire new semantic knowledge
in patients with large medial temporal lobe lesions. *Hippocampus* **15**, 273–280
(2005).
4.   Manns, J. R., Hopkins, R. O. & Squire, L. R. Semantic Memory and the
Human Hippocampus. *Neuron* **38**, 127–133 (2003).
5.   Scoville, W. B. & Milner, B. Loss of recent memory after bilateral
hippocampal lesions. *J. Neurol. Neurosurg. Psychiatr.* **20**, 11–21 (1957).
6.   Verfaellie, M., Koseff, P. & Alexander, M. P. Acquisition of novel
semantic information in amnesia: effects of lesion location. *Neuropsychologia* **38**,
484–492 (2000).
7.   Guderian, S. *et al.* Hippocampal Volume Reduction in Humans Predicts
Impaired Allocentric Spatial Memory in Virtual-Reality Navigation. *J. Neurosci.*
35, 14123–14131 (2015).
8.   Patai, E. Z. *et al.* Extent of hippocampal atrophy predicts degree of deficit
in recall. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 12830–12833 (2015).
9.   Picard, L. *et al.* Functional independence within the self-memory system:
new insights from two cases of developmental amnesia. *Cortex* **49**, 1463–1481
(2013).
10.   Rosenbaum, R. S., Cassidy, B. N. & Herdman, K. A. Patterns of preserved
and impaired spatial memory in a case of developmental amnesia. *Front Hum*
Neurosci **9**, 196 (2015).
11.   Lavenex, P. & Banta Lavenex, P. Building hippocampal circuits to learn
and remember: insights into the development of human memory. *Behav Brain Res*
254, 8–21 (2013).
12.   Elward, R. L. & Vargha-Khadem, F. Semantic memory in developmental
amnesia. *Neurosci. Lett.* **680**, 23–30 (2018).
13.   Geva, S. *et al.* Volume reduction of caudate nucleus is associated with

movement coordination deficits in patients with hippocampal atrophy due to
perinatal hypoxia-ischaemia. *Neuroimage Clin* **28**, 102429 (2020).
14.   Vargha-Khadem, F. *et al.* Developmental amnesia: effect of age at injury.
Proc Natl Acad Sci U S A **100**, 10055–10060 (2003).
15.   Mishkin, M., Suzuki, W. A., Gadian, D. G. & Vargha-Khadem, F.
Hierarchical Organization of Cognitive Memory. *Philosophical Transactions:*
Biological Sciences **352**, 1461–1467 (1997).
16.   Baddeley, A. D. On trying to prove Endel Tulving wrong: A revised modal
model of amnesia. *Neuropsychologia* **148**, 107590 (2020).
17.   Vargha-Khadem, F. & Cacucci, F. A brief history of developmental
amnesia. *Neuropsychologia* **150**, 107689 (2021).
18.   Argyropoulos, G. P. D. *et al.* Functional Specialization of the Medial
Temporal Lobes in Human Recognition Memory: Dissociating Effects of
Hippocampal versus Parahippocampal Damage. *Cereb Cortex* **32**, 1637–1652
(2022).
19.   Schmidt-Kastner, R. & Freund, T. F. Selective vulnerability of the
hippocampus in brain ischemia. *Neuroscience* **40**, 599–636 (1991).
20.   Lavenex, P., Sugden, S. G., Davis, R. R., Gregg, J. P. & Lavenex, P. B.
Developmental regulation of gene expression and astrocytic processes may
explain selective hippocampal vulnerability. *Hippocampus* **21**, 142–149 (2011).
21.   Wang, X. & Michaelis, E. K. Selective neuronal vulnerability to oxidative
stress in the brain. *Front Aging Neurosci* **2**, 12 (2010).
22.   Bartsch, T. *et al.* Selective neuronal vulnerability of human hippocampal
CA1 neurons: lesion evolution, temporal course, and pattern of hippocampal
damage in diffusion-weighted MR imaging. *J. Cereb. Blood Flow Metab.* **35**,
1836–1845 (2015).
23.   Zola-Morgan, S., Squire, L. R. & Amaral, D. G. Human amnesia and the
medial temporal region: enduring memory impairment following a bilateral lesion
limited to field CA1 of the hippocampus. *J. Neurosci.* **6**, 2950–2967 (1986).
24.   Rempel-Clower, N. L., Zola, S. M., Squire, L. R. & Amaral, D. G. Three
cases of enduring memory impairment after bilateral damage limited to the
hippocampal formation. *J. Neurosci.* **16**, 5233–5255 (1996).
25.   Maguire, E. A., Vargha-Khadem, F. & Mishkin, M. The effects of bilateral
hippocampal damage on fMRI regional activations and interactions during
memory retrieval. *Brain* **124**, 1156–1170 (2001).
26.   Rabin, J. S., Olsen, R. K., Gilboa, A., Buchsbaum, B. R. & Rosenbaum, R.
S. Using fMRI to understand event construction in developmental amnesia.
Neuropsychologia **90**, 261–273 (2016).
27.   Elward, R. L., Rugg, M. D. & Vargha-Khadem, F. When the brain, but not
the person, remembers: Cortical reinstatement is modulated by retrieval goal in
developmental amnesia. *Neuropsychologia* **154**, 107788 (2021).
28.   Shrager, Y., Levy, D. A., Hopkins, R. O. & Squire, L. R. Working
memory and the organization of brain systems. *J Neurosci* **28**, 4818–4822 (2008).
29.   Jonin, P.-Y. *et al.* Superior explicit memory despite severe developmental
amnesia: In-depth case study and neural correlates. *Hippocampus* **28**, 867–885
(2018).
30.   Zeidman, P. & Maguire, E. A. Anterior hippocampus: the anatomy of
perception, imagination and episodic memory. *Nat. Rev. Neurosci.* **17**, 173–182
(2016).

31. Clark, I. A., Dalton, M. A. & Maguire, E. A. Posterior hippocampal CA2/3
volume is associated with autobiographical memory recall ability in lower
performing individuals. *Sci Rep* **13**, 7924 (2023).
32. Rosene, D. L. & Van Hoesen, G. W. Hippocampal efferents reach
widespread areas of cerebral cortex and amygdala in the rhesus monkey. *Science*
198, 315–317 (1977).
33. Insausti, R. *et al.* MR volumetric analysis of the human entorhinal,
perirhinal, and temporopolar cortices. *AJNR Am J Neuroradiol* **19**, 659–671
(1998).
34. Erdem, A., Yaşargil, G. & Roth, P. Microsurgical anatomy of the
hippocampal arteries. *J. Neurosurg.* **79**, 256–265 (1993).
35. Huther, G., Dörfl, J., Van der Loos, H. & Jeanmonod, D. Microanatomic
and vascular aspects of the temporomesial region. *Neurosurgery* **43**, 1118–1136
(1998).
36. Marinković, S., Milisavljević, M. & Puskas, L. Microvascular anatomy of
the hippocampal formation. *Surg Neurol* **37**, 339–349 (1992).
37. Perosa, V. *et al.* Hippocampal vascular reserve associated with cognitive
performance and hippocampal volume. *Brain* **143**, 622–634 (2020).
38. Liebeskind, D. S. Collateral circulation. *Stroke* **34**, 2279–2284 (2003).
39. Ding, S.-L. Comparative anatomy of the prosubiculum, subiculum,
presubiculum, postsubiculum, and parasubiculum in human, monkey, and rodent.
J. Comp. Neurol. **521**, 4145–4162 (2013).
40. Zeidman, P., Lutti, A. & Maguire, E. A. Investigating the functions of
subregions within anterior hippocampus. *Cortex* **73**, 240–256 (2015).
41. Kondo, H., Lavenex, P. & Amaral, D. G. Intrinsic connections of the
macaque monkey hippocampal formation: II. CA3 connections. *J. Comp. Neurol.*
515, 349–377 (2009).
42. Amaral, D. G., Insausti, R. & Cowan, W. M. The commissural connections
of the monkey hippocampal formation. *J. Comp. Neurol.* **224**, 307–336 (1984).
43. Schulze, K., Vargha-Khadem, F. & Mishkin, M. Test of a motor theory of
long-term auditory memory. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 7121–7125 (2012).
44. Eichenbaum, H. Remembering: functional organization of the declarative
memory system. *Curr Biol* **16**, R643-645 (2006).
45. Eichenbaum, H., Otto, T. & Cohen, N. J. Two functional components of
the hippocampal memory system. *Behavioral and Brain Sciences* **17**, 449–517
(1994).
46. Hannula, D. E. & Ranganath, C. Medial temporal lobe activity predicts
successful relational memory binding. *J Neurosci* **28**, 116–124 (2008).
47. Libby, L. A., Hannula, D. E. & Ranganath, C. Medial temporal lobe
coding of item and spatial information during relational binding in working
memory. *J Neurosci* **34**, 14233–14242 (2014).
48. Borders, A. A., Ranganath, C. & Yonelinas, A. P. The hippocampus
supports high-precision binding in visual working memory. *Hippocampus* **32**,
217–230 (2022).
49. Allen, R. J., Vargha-Khadem, F. & Baddeley, A. D. Item-location binding
in working memory: is it hippocampus-dependent? *Neuropsychologia* **59**, 74–84
(2014).
50. Baddeley, A., Jarrold, C. & Vargha-Khadem, F. Working memory and the
hippocampus. *J Cogn Neurosci* **23**, 3855–3861 (2011).

51.   Baddeley, A., Allen, R. & Vargha-Khadem, F. Is the hippocampus
necessary for visual and verbal binding in working memory? *Neuropsychologia*
48, 1089–1095 (2010).
52.   Allen, R. J., Atkinson, A. L., Vargha-Khadem, F. & Baddeley, A. D. Intact
high-resolution working memory binding in a patient with developmental amnesia
and selective hippocampal damage. *Hippocampus* **32**, 597–609 (2022).
53.   Miyashita, Y. Perirhinal circuits for memory processing. *Nature Reviews*
Neuroscience **20**, 577–592 (2019).
54.   Bachevalier, J. & Mishkin, M. Mnemonic and neuropathological effects of
occluding the posterior cerebral artery in Macaca mulatta. *Neuropsychologia* **27**,
83–105 (1989).
55.   Murray, E. A. & Mishkin, M. Object recognition and location memory in
monkeys with excitotoxic lesions of the amygdala and hippocampus. *J. Neurosci.*
18, 6568–6582 (1998).
56.   Beason-Held, L. L., Rosene, D. L., Killiany, R. J. & Moss, M. B.
Hippocampal formation lesions produce memory impairment in the rhesus
monkey. *Hippocampus* **9**, 562–574 (1999).
57.   Baxter, M. G. & Murray, E. A. Opposite relationship of hippocampal and
rhinal cortex damage to delayed nonmatching-to-sample deficits in monkeys.
Hippocampus **11**, 61–71 (2001).
58.   Aggleton, J. P., Vann, S. D., Oswald, C. J. & Good, M. Identifying cortical
inputs to the rat hippocampus that subserve allocentric spatial processes: a simple
problem with a complex answer. *Hippocampus* **10**, 466–474 (2000).
59.   Banta Lavenex, P., Amaral, D. G. & Lavenex, P. Hippocampal lesion
prevents spatial relational learning in adult macaque monkeys. *J. Neurosci.* **26**,
4546–4558 (2006).
60.   Lavenex, P., Lavenex, P. B. & Amaral, D. G. Spatial relational learning
persists following neonatal hippocampal lesions in macaque monkeys. *Nat.*
Neurosci. **10**, 234–239 (2007).
61.   Chareyron, L. J., Banta Lavenex, P., Amaral, D. G. & Lavenex, P.
Functional organization of the medial temporal lobe memory system following
neonatal hippocampal lesion in rhesus monkeys. *Brain Struct Funct* **222**, 3899–
3914 (2017).
62.   Villard, J. *et al.* Structural plasticity in the entorhinal and perirhinal
cortices following hippocampal lesions in rhesus monkeys. *Hippocampus* (2023)
doi:10.1002/hipo.23567.
63.   Kapur, N. *et al.* Positive clinical neuroscience: explorations in positive
neurology. *Neuroscientist* **19**, 354–369 (2013).
64.   de Vries, L. S. & Jongmans, M. J. Long-term outcome after neonatal
hypoxic-ischaemic encephalopathy. *Arch. Dis. Child. Fetal Neonatal Ed.* **95**,
F220-224 (2010).
65.   Cooper, J. M., Vargha-Khadem, F., Gadian, D. G. & Maguire, E. A. The
effect of hippocampal damage in children on recalling the past and imagining new
experiences. *Neuropsychologia* **49**, 1843–1850 (2011).
66.   Buck, S., Bastos, F., Baldeweg, T. & Vargha-Khadem, F. A Functional
MRI Paradigm Suitable for Language and Memory Mapping in Pediatric
Temporal Lobe Epilepsy. *Front Neurol* **10**, 1384 (2019).
67.   Yushkevich, P. A. *et al.* User-guided 3D active contour segmentation of
anatomical structures: significantly improved efficiency and reliability.

Neuroimage **31**, 1116–1128 (2006).

68.   Akudjedu, T. N. *et al.* A comparative study of segmentation techniques for

the quantification of brain subcortical volume. *Brain Imaging Behav* **12**, 1678–

1695 (2018).

69.   Schmidt, M. F. *et al.* A comparison of manual tracing and FreeSurfer for

estimating hippocampal volume over the adult lifespan. *Hum Brain Mapp* **39**,

2500–2513 (2018).

70.   Herten, A., Konrad, K., Krinzinger, H., Seitz, J. & von Polier, G. G.

Accuracy and bias of automatic hippocampal segmentation in children and

adolescents. *Brain Struct Funct* **224**, 795–810 (2019).

71.   Amaral, D. G. & Lavenex, P. Hippocampal neuroanatomy. in *The*

hippocampus book 37–114 (Oxford University Press, 2007).

72.   Ding, S.-L. & Van Hoesen, G. W. Organization and Detailed Parcellation

of Human Hippocampal Head and Body Regions Based on a Combined Analysis

of Cyto- and Chemoarchitecture. *J. Comp. Neurol.* **523**, 2233–2253 (2015).

73.   Dalton, M. A., Zeidman, P., Barry, D. N., Williams, E. & Maguire, E. A.

Segmenting subregions of the human hippocampus on structural magnetic

resonance image scans: An illustrated tutorial. *Brain Neurosci Adv* **1**,

2398212817701448 (2017).

74.   Blaizot, X. *et al.* The human parahippocampal region: I. Temporal pole

cytoarchitectonic and MRI correlation. *Cereb Cortex* **20**, 2198–2212 (2010).

75.   Ding, S.-L. & Van Hoesen, G. W. Borders, extent, and topography of

human perirhinal cortex as revealed using multiple modern neuroanatomical and

pathological markers. *Hum Brain Mapp* **31**, 1359–1379 (2010).

76.   Frankó, E., Insausti, A. M., Artacho-Pérula, E., Insausti, R. & Chavoix, C.

Identification of the human medial temporal lobe regions on magnetic resonance

images. *Hum Brain Mapp* **35**, 248–256 (2014).

77.   Kivisaari, S. L., Probst, A. & Taylor, K. I. The Perirhinal, Entorhinal, and

Parahippocampal Cortices and Hippocampus: An Overview of Functional

Anatomy and Protocol for Their Segmentation in MR Images. in *fMRI: Basics*

and Clinical Applications (eds. Ulmer, S. & Jansen, O.) 355–383 (Springer

International Publishing, 2020). doi:10.1007/978-3-030-41874-8_24.

78.   Baddeley, A. D., Emslie, H. & Nimmo-Smith, I. *Doors and people : a test*

of visual and verbal recall and recognition. (Bury St Edmunds : Thames Valley

Test Company, 1994).

79.   Adlam, A.-L. R., Malloy, M., Mishkin, M. & Vargha-Khadem, F.

Dissociation between recognition and recall in developmental amnesia.

Neuropsychologia **47**, 2207–2210 (2009).

80.   Rothman, K. J. No adjustments are needed for multiple comparisons.

Epidemiology **1**, 43–46 (1990).

81.   Saville, D. J. Multiple Comparison Procedures: The Practical Solution.

The American Statistician **44**, 174–180 (1990).