Title
Paradoxical consequences of early hippocampal damage: greater
atrophy is associated with better recall, working memory and
visuospatial perception in developmental amnesia

Authors/Affiliations
Loïc J. Chareyron[a], W.K. Kling Chong[b], Tina Banks[c], Mortimer Mishkin[d], Neil Burgess[e],
Richard C. Saunders[d], Faraneh Vargha-Khadem[a]*
[a] Cognitive Neuroscience and Neuropsychiatry, Developmental Neurosciences, University
College London Great Ormond Street Institute of Child Health, London, UK
[b] Developmental Imaging & Biophysics, University College London Great Ormond Street
Institute of Child Health, London, UK
[c] Developmental Imaging & Biophysics, University College London Great Ormond Street
Institute of Child Health, London, UK
[d] Laboratory of Neuropsychology, National Institute of Mental Health, National Institutes of
Health, Bethesda, MD 20892.
[e] Institute of Cognitive Neuroscience, University College London, London, UK.

Corresponding Author
* Faraneh Vargha-Khadem.
Email: f.vargha-khadem@ucl.ac.uk

Abstract
Despite bilateral hippocampal damage dating to perinatal or early-childhood
period, patients with developmental amnesia (DA) exhibit well-developed
semantic memory, but severely impaired episodic memory. It is not clear,
however, whether the residual hippocampus plays a role in encoding and/or
retrieval of new information, or the surrounding cortical areas rescue aspects of
these critical cognitive memory processes after early injury. We used manual MRI
segmentation to estimate the volume of three hippocampal subregions and three
surrounding cortical areas in 23 patients with DA and 32 controls. In patients, the
level of atrophy in CA-DG subregions and subicular complex was more than 40%
while the atrophy of the uncus was moderate (−23%). In contrast, volumes of
entorhinal, perirhinal and parahippocampal cortices were smaller than those of
controls, but not statistically different. Patients' recall, Verbal IQ, Working
Memory Index and Processing Speed Index scores correlated negatively with the
volume of the uncus while spatial perception/immediate memory performance
correlated negatively with the volume of the subicular complex. We propose that
in patients with DA, no other structure can perform the function of the
hippocampus in episodic memory. However, cognitive memory processing is
compromised as a function of extent of atrophy in hippocampal subregions, such
that the greater the damage, the more likely that preserved surrounding cortical
areas will be recruited to rescue the putative functions of the damaged subregions.
Our findings document for the first time not only the extent, but also the limits of
circuit reorganization occurring in the young brain following bilateral
hippocampal damage.

Introduction

Developmental amnesia

We previously identified a group of patients with hippocampal atrophy who had suffered hypoxic-ischaemic episodes in infancy or childhood and later developed a memory disorder differing from that commonly described in patients with adult-onset amnesia of temporal-lobe origin. In the latter cases, the disorder most often takes the form of a global anterograde memory loss [i.e. one affecting both episodic and new post-injury semantic memory, and severely restricting recall and recognition processes (Bayley and Squire, 2005; Manns et al., 2003; Scoville and Milner, 1957; Verfaellie et al., 2000)]. In patients with hippocampal damage of early-onset, by contrast, the disorder is more limited, being characterized by markedly impaired episodic memory, recall/recollection, and spatial processing deficits, but relatively preserved semantic memory, working memory, and recognition performance. We have labeled this dissociated form of memory function 'developmental amnesia' (DA, Guderian et al., 2015; Patai et al., 2015; Picard et al., 2013; Rosenbaum et al., 2015; Vargha-Khadem et al., 1997). Given the early onset of hippocampal damage, the compensatory organization that has occurred in patients with DA is likely to be distinct from what is observed in patients with hippocampal damage acquired in adulthood. Also, in patients with acquired hippocampal lesions, the damage has occurred to already-established, and normally-functioning memory circuits, whereas in patients with DA, the early bilateral lesions have probably led to the development of a different memory system. For these reasons, it is difficult, and somewhat inappropriate, to compare memory function and cognitive profiles in patients with DA to those of patients with adult-onset amnesia (Elward and Vargha-Khadem, 2018, P: 24-26).

To date, a cohort of 18 patients with DA has been reported, each patient having suffered an episode of hypoxia-ischaemia in early life (Dzieciol et al., 2017). Importantly, irrespective of the underlying aetiology (Geva et al., 2020), or age at onset (Vargha-Khadem et al., 2003), the hypoxic-ischaemic event was the cause of hippocampal damage and cognitive impairment (Cooper et al., 2015).

Memory circuit reorganization

Among the many questions that have arisen from the study of patients with DA, the foremost concerns the regions of the medial temporal lobe (MTL) and/or preserved hippocampal subregions that could be involved in memory circuit reorganization following early hippocampal damage.

In the model of hierarchical organization of cognitive memory proposed by Mishkin and colleagues (1997; see also: Baddeley, 2020; Vargha-Khadem and Cacucci, 2021), recollection and familiarity depend on different MTL structures. According to this model, the perihippocampal gyrus, and in particular the perirhinal cortex, may support familiarity, while the hippocampus may support recollection (Argyropoulos et al., 2021). The selectivity of impairment observed in patients with DA has provided strong support for this dual-process model.

Because patients with DA have both neuroradiological and quantitatively-confirmed bilateral hippocampal atrophy, but no visible abnormality within the parahippocampal region, it has been proposed that their episodic memory impairment, and recall/recollection deficits are due to their hippocampal pathology, whereas their relatively intact semantic memory, and

recognition/familiarity processes are due in part to the preservation of the
perihippocampal gyrus.

Hippocampus
Hypoxia-ischemia is known to differentially affect the hippocampal fields and
subdivisions. Studies of animal models of cerebral ischemia have reported that a
brief episode results in selective neuronal death in the CA1 field of the
hippocampus (Schmidt-Kastner and Freund, 1991) while the adjacent CA3
remains less vulnerable (Wang and Michaelis, 2010). In humans, while damage to
CA1 is consistently reported following cerebral ischemia, damage to additional
hippocampal regions is much more variable (Bartsch et al., 2015). Postmortem
examinations of patients who suffered ischemic episodes in adulthood have
reported neural cell loss either restricted to the CA1 region, or distributed in the
dentate gyrus, subiculum, uncus and entorhinal cortex (Rempel-Clower et al.,
1996; Zola-Morgan et al., 1986). The preservation of circumscribed hippocampal
subregions following injury of early-onset in DA could account for the increased
activation of the residual hippocampus during different memory tasks (Maguire et
al., 2001; Rabin et al., 2016), although activation *per se* does not necessarily
imply correspondingly intact function (Elward et al., 2021).
In the study reported by Dzieciol and colleagues (Dzieciol et al., 2017), the
degree of hippocampal atrophy in patients with DA ranged from 28% to 62%
compared to healthy controls. This large range of hippocampal volume reduction
in this cohort suggests that there may indeed be variability in the hippocampal
response to hypoxia-ischemia at the level of hippocampal subregions. It would
thus be possible to assess whether the compensatory response in patients with DA
is dependent on the extent of atrophy in the hippocampal subregions. One way to
address this is to evaluate the association between task-dependent
neuropsychological performance and volumes of hippocampal subregions

Perihippocampal gyrus
While postmortem examinations have reported some neural cell loss in layers of
entorhinal cortex in patients who suffered ischemic episodes in adulthood
(Rempel-Clower et al., 1996), it was reported that the volume of the
parahippocampal gyrus was not significantly reduced in patients with adult-onset
amnesia presenting with bilateral hippocampal damage (Shrager et al., 2008). A
cortical thickness analysis performed in a single patient with DA didn't find
abnormality in the subhippocampal structures within the medial temporal lobe,
including the perirhinal and ventral entorhinal cortices (Jonin et al., 2018). Also, a
recent study demonstrated that parahippocampal activity during scene
reinstatement in patients with DA was similar to controls (Elward et al., 2021).
However, the structural integrity of the perihippocampal gyrus, and by
implication, the integrity of the hypothesized substrate of semantic memory and
other preserved mnemonic processes remain to be quantitatively assessed in
patients with DA.
133
Here, we have used manual segmentation of MRI scans to obtain estimates of the
volume of three hippocampal subregions: the uncus [uncal portion of the
hippocampus (Zeidman and Maguire, 2016)], the CA-DG (including CA fields
and dentate gyrus) and the subicular complex (including subiculum, presubiculum
and parasubiculum), and three surrounding cortical areas (entorhinal, perirhinal

and parahippocampal cortices) in patients with DA and healthy controls. Controls
and patients were assessed with neuropsychological tests of intelligence (WISC or
WAIS), recall/recognition for verbal and visual material (Doors and People Test)
and spatial perception/memory (Four Mountains Test). Finally, we investigated
the relationships between degree of atrophy of hippocampal subregions and
cognitive deficits.
We predicted that: (i) residual hippocampal subregions and/or surrounding
cortical areas were sufficiently preserved to rescue some aspects of hippocampal
function, and (ii) plastic changes in these regions occurred as a function of
hippocampal damage in all patients with DA.
We documented several paradoxical inverse relationships: i) between the
volume of the residual uncus and patients' recall memory performance, Verbal
Comprehension Index, Working memory Index and Processing Speed Index, and
ii) between the volume of the residual subicular complex and patients'
performance on spatial perception/memory. We also reported an absence of
structural damage in the surrounding cortical areas in patients with DA.

Results

Volumetric estimates

157 Three hippocampal subregions (uncus, CA-DG and subicular complex) and three
158 surrounding cortical areas (entorhinal, perirhinal and parahippocampal cortices)
159 were manually segmented in healthy controls (N = 32) and DA (N = 23) groups
160 (Fig. S1).
161

Fig. 1. Volume of the hippocampus and its subregions. Volume of the hippocampus (**A.**)
and its subregions (**B.**, uncus, CA-DG, and subicular complex) in control (grey; N = 32) and
DA (orange; N = 23) groups. Data are represented as mean ± SD. Percentage values refer to
the mean difference to control group. Volumes are calculated as the average of left and right
hemisphere volumes. All volumes are corrected for intracranial (ICV) volume. *: P < .05;
***: P < .001.

The volume of the hippocampus as a whole was 40% smaller in the DA than in
the control group (range: −26% to −53%; $t_{(53)} = -15.32$; P < .001) (Fig. 1; Table
S1). The volume of the hippocampal subregions was also significantly smaller in
the DA than in the control group (2-way ANOVA, $F_{(1,159)} = 392.5$, P < .001). The

uncus was 23% smaller (range: +4% to −49%; Tukey's HSD post-hoc test: P =
.037), the CA-DG subregions were 45% smaller (range: −28% to −59%; P <
.001), and the subicular complex was 41% smaller (range: −22% to −56%; P <
.001). The level of atrophy of the patients' hippocampal subregions differed
significantly from each other ($F_{(2,66)} = 23.0$, P < .001). Tukey's HSD post-hoc test
indicated that the CA-DG and subicular complex showed greater atrophy than the
uncus (both P < .001).
Atrophy of the hippocampus in the DA group was significantly lower in its
anterior (−36%) than its posterior (−44%) segment (paired t test, $t_{(22)} = 2.72$, P =
.012; Fig. S2). Because no differences were observed between the anterior and the
posterior segment of the CA-DG (paired t test, $t_{(22)} = 0.94$, P = .36) or subicular
complex ($t_{(22)} = 1$, P = .33) subregions, the lower level of atrophy observed in the
anterior part of the whole hippocampus can be attributed to the relatively more
preserved uncus.
In contrast, there was no significant difference in the volume of the entorhinal,
perirhinal and parahippocampal cortices between control and DA groups (Table
S1; 2-way ANOVA, $F_{(1,159)} = 6.7$, P = .010; Tukey's HSD post-hoc tests, all P >
.35). Also, we didn't observe any correlation between the volume of the
hippocampus or hippocampal subregions and the volume of surrounding cortical
areas in patients with DA (Pearson's correlation analysis, all P > .19).

Neuroradiological assessment
Neuroradiological ratings based on visual inspection of the MR images provided
independently by one of the authors (WKC) revealed abnormally small fornix and
mammillary bodies in all but two patients with DA (Table S2). No abnormalities
were detected in the parahippocampal gyrus (perirhinal, entorhinal, and
parahippocampal cortices), or the thalamus, and the basal ganglia. Additional
visible abnormality was observed in some cases in the white matter and the
cerebellum. The pattern of visible abnormality detected within the hippocampal
circuit, and the notable absence thereof in other cortical and subcortical structures,
was consistent across individual patients irrespective of their varying aetiology,
and age at hypoxic-ischaemic-induced hippocampal damage.

WISC/WAIS
All patients with DA (N = 23) and 23 of the control participants were assessed on
the WISC or WAIS. Patients were impaired relative to our group of controls only
for the Perceptual Reasoning Index scores (2-way ANOVA, $F_{(1,176)} = 10.61$, P =
.001; Tukey's HSD post-hoc test, P = 0.04; all other P > .25). There were
significant inverse correlations between the volume of the uncus and patients'
scores for the Verbal Comprehension Index (VCI), the Working Memory Index
(WMI) and the Processing Speed Index (PSI) (all $r_{(22)} < -0.42$; all P < .05; Fig 2).
Looking at PSI scores, we observed an inverse correlations between "Digit
Symbol-Coding" score and the volume of the uncus ($r_{(22)} = -0.44$; P = .037) and
between "Symbol Search" and the volume of the uncus and subicular complex
($r_{(22)} = -0.43$; P = .04 and $r_{(22)} = -0.44$; P = .037, respectively; Table S3). No
other correlations for other ROIs in the DA group were significant. In controls, the
WMI scores were positively associated with the volume of the subicular complex
($r_{(22)} = 0.46$; P = .029).
220

Fig. 2. Inverse correlations between WISC/WAIS indexes scores and the volume of the uncus in patients with DA. Participants' scores on Wechsler Intelligence Scales WISC or WAIS (patients: orange circles, N = 23; controls: grey triangles, n = 23). **A.** Verbal comprehension Index; **B.** Perceptual Reasoning Index; **C.** Working Memory Index; **D.** Processing Speed Index. Pearson's correlation coefficient. All volumes are corrected for intracranial (ICV) volume. *: P < .05.

Doors and People Test

All patients with DA (N = 23) and 21 of the control participants were assessed on the Doors and People Test, which provides equated measures of recognition and recall in the visual and verbal domains. All patients' subtest scores were significantly below zero (Wilcoxon signed rank test: all Z < −2.4, all P < .016), indicating that patients were impaired relative to controls (who represent zero, as their scores were used to convert raw scores into z-scores). The patients' scores for the four Doors and People subtests differed significantly from each other (Friedman's test: $\chi^2_{(3,n=22)}$ = 43.7, P < .001). The visual recall ('Shapes') subtest yielded a significantly greater deficit than all of the other subtests (all P < .001). The degree of deficit on the verbal recall ('People') subtest was greater than on the two recognition subtests (both P < .02). The degree of deficit was greater on the visual recognition ('Doors') subtest than on the verbal recognition ('Names') subtest (P = .001). Collapsing across subtests, we found a greater deficit in recall compared with recognition (Z = −4.04; P < .001), and a greater deficit in memory for visual compared with memory for verbal material (Z = −4.11; P < .001).

We observed significant negative correlations between the two recall subtest scores and the volume of the uncus in the DA group [Verbal recall ('People'): Pearson's $r_{(22)}$ = –0.47; P = .02; Visual recall ('Shapes'): $r_{(22)}$ = –0.55; P = .007]. No other correlations for other subtests or other regions of interest in the DA group were significant (all P > .09; Table S4). Collapsing across test material confirmed that scores on recall were inversely correlated with the volume of the uncus in the DA group ($r_{(22)}$ = –0.58; P = .004; Fig. 3A). As a corollary, we observed that patients with more preserved uncus had significantly lower scores on recall than patients with more damaged uncus (Z = −2.74; P = .006). In the control group, no correlations for any subtests or regions of interest were significant (all P > .11; Table S4).

Fig. 3. *Doors and People* recall scores correlations with the volumes of hippocampal subregions. The patients' scores (orange circles; N = 23) on recall subtests of the *Doors and People Test* correlated negatively with the volume of the uncus (**A.**) but not with the volumes of the subicular complex (**B.**) or CA-DG (**C.**). The controls' scores (grey triangles; n = 21) did not correlate with the volume of any subregion. Pearson's correlation coefficient. All volumes are corrected for intracranial (ICV) volume. **: P < .01.

During *Doors and People* testing, the procedures for recall subtests are repeated until all items (names or shapes) are correctly recalled, or for a maximum of three presentations. The patients' scores differed significantly from each other between the three trials of the visual recall ('Shapes') subtest (Friedman's test: $\chi^2_{(2,n=20)}$ = 8.00, P = .018) and between the three trials of the verbal recall ('People') subtest ($\chi^2_{(2,n=20)}$ = 14.15, P < .001). Post hoc Wilcoxon signed-ranks tests indicated that scores obtained on trial-1 were significantly lower than scores obtained on trial-3 for the visual recall ('Shapes') subtest (Z = −2.72; P = .006) and for the verbal recall ('People') subtest (Z = −2.88; P = .004). Interested to determine potential differences in learning between patients depending on whether they presented with severe or moderate uncus atrophy, we compared increases in patients' scores between trial-1 and trial-3. There was no difference in learning scores between patients with a larger uncus (above group mean volume) and patients with a smaller uncus (Mann-Whitney U test; visual recall ('Shapes'): Z = −1.26; P = .21; verbal recall ('People'): Z = 0.61; P = .54).

Delayed (cued) verbal and visual recall were tested after completion of the verbal and visual recognition subtests, respectively. We found a correlation between patients' scores for the delayed verbal recall ('People') subtest and the volume of the uncus (Pearson's $r_{(20)}$ = −0.5; P = .02) while the correlation with delayed visual recall ('Shapes') scores was not significant ($r_{(20)}$ = −0.3; P = .18).

Four Mountains Test

Fourteen patients with DA and 19 of the control participants were assessed on the *Four Mountains Test* to examine their perception and immediate memory for topographical information in spatial scenes. Patients' subtest scores converted to z-scores were not significantly below zero (Wilcoxon signed rank test: 'Spatial Perception': P = .75; 'Spatial Memory': P = .13), indicating that many of the patients were not impaired relative to our group of controls. Patients' scores were significantly lower for the 'Spatial Memory' than for the 'Spatial Perception' subtest (Z = −2.67; P = .008). There were significant inverse correlations between patients' scores for the 'Spatial Perception' and 'Spatial Memory' subtests and the volume of the subicular complex ($r_{(13)}$ = –0.59 ; P = .025 and $r_{(13)}$ = –0.58 ; P =

.03, respectively). Collapsing across test material confirmed that scores were
strongly inversely correlated with the volume of the subicular complex in the DA
group ($r_{(13)}$ = –0.63; P = .015) (Fig. 4B). Patients' scores for the 'Spatial
Perception' subtest were also negatively associated with the volume of the uncus
($r_{(13)}$ = –0.54 ; P = .047). In the control group, no correlations for any ROIs were
significant (all P > .49; Table S5).

Fig. 4. *Four Mountains* **scores correlations with the volumes of hippocampal subregions.**
The patients' scores (orange circles; n = 14) correlated negatively with the volume of the
subicular complex (**B.**) but not with the volumes of the uncus (**A.**) or CA-DG (**C.**). The
controls' scores (grey triangles; n = 19) did not correlate with the volume of any subregion.
Pearson's correlation coefficient. All volumes are corrected for intracranial (ICV) volume.

Discussion

In this study, we addressed the question of whether the volumes of residual
hippocampal subregions and parahippocampal areas are associated with memory
and spatial processing performance in patients with hypoxic-ischaemic-induced
hippocampal damage of early-onset. Using a combination of neuropsychological
test results and volumetric estimations in a large cohort of patients with
developmental amnesia (DA), we documented several inverse relationships: i)
between the volume of the residual uncus and patients' performance on recall
memory, Verbal comprehension Index, Working memory Index and Processing
speed Index, and ii) between the volume of the residual subicular complex and
patients' performance on spatial perception/memory. The effects reported here
suggest that following an early-life exposure to hypoxia-ischemia, the function of
hippocampal subregions showing greater damage will more likely be compensated
by preserved extra-hippocampal structures. By contrast, hippocampal subregions
with milder injury could increase system disruption by maintaining their putative
functions. Despite these compensatory reorganizations, hippocampal-dependent
episodic memory recall remains irreversibly compromised.

Limitations
Although manual tracing is still the gold standard for measuring the volumes of
hippocampal subregions and cortical areas on MRI acquisitions, manual
segmentation is subjective and susceptible to measurement errors. Also, the
precision of the measure is limited by the resolution of the MRI scans. However,
analyses of MR images for estimating the extent of hippocampal damage in

amnesic patients have been validated by postmortem examinations (Rempel-
Clower et al., 1996). In particular, a clear concordance has been reported in the
appearance of the damaged hippocampus on the MRI and in the histological
sections (Rempel-Clower et al., 1996). The border between cytoarchitectonic
hippocampal subregions were estimated based on delineation of these regions on
histological sections (Rosene and Van Hoesen, 1977). The delineation on MR
images of the perihippocampal and parahippocampal cortical areas were based on
that described in (Insausti et al., 1998). It is not clear to what extent these borders
would be affected in the brains of DA patients with significant hippocampal
volume reduction. Our observations of an absence of significant volume reduction
of the entorhinal, perirhinal and parahippocampal cortices in patients with DA is
consistent with previous report showing an absence of significant volume
reduction in patients with adult-onset amnesia (Shrager et al., 2008), and in a
patient with DA (Jonin et al., 2018). Also, intact parahippocampal activity during
scene reinstatement has been described in patients with DA (Elward et al., 2021).

Differential effect of early hypoxia-ischemia on hippocampal subregions
Our observation that the CA-DG were the most affected hippocampal subregions
in patients with DA is congruent with documented reports of greater sensitivity of
this subregion to hypoxia-ischemia. It has long been known that hypoxia and
ischemia result in selective neuronal death in the CA1 field (Schmidt-Kastner and
Freund, 1991), although the reason for this selectivity is still debated (Wang and
Michaelis, 2010). Neurohistological studies of patients who suffered ischemic
episodes in adulthood have reported cases with neural cell loss restricted to the
CA1 region bilaterally whilst the CA1' region of the uncus, the subiculum, CA3
and surrounding cortices were all preserved (Rempel-Clower et al., 1996; Zola-
Morgan et al., 1986). High-resolution magnetic resonance imaging showed that
CA1 was always affected in patients with hippocampal damage (Bartsch et al.,
2015).
The subicular complex was less affected than the CA-DG subregion in
patients with DA and this difference could be related to the different levels of
maturation of the hippocampal subregions at birth. Quantitative volumetric
measurements in non-human primates revealed that the subiculum, presubiculum
and parasubiculum develop earlier than the dentate gyrus, CA3, and CA1 (Jabès et
al., 2011) and this progressive maturation of the hippocampal circuits may
participate in the differential protracted maturation of relational memory processes
(Jabès and Nelson, 2015). The level of maturation of the hippocampal subregions
at birth could also influence their vulnerability to hypoxia-ischemia (Vannucci
and Hagberg, 2004).
In contrast, the uncus showed a low sensitivity to hypoxia in our patient cohort
which could be a reflection of the distinct vascularization of this hippocampal
subregion. The branches of the posterior cerebral artery (PCA) coming from the
vertebral artery irrigate the posterior part of the hippocampus, while the branches
of the anterior choroidal artery (AChA) originating from the internal carotid artery
irrigate the uncal portion of the hippocampus (Erdem et al., 1993; Huther et al.,
1998; Marinković et al., 1992). High-resolution angiography and anatomical
studies have highlighted that the anterior hippocampal region could receive a
mixed blood supply (from both the PCA and the AChA) in about half of the
individuals (Erdem et al., 1993; Perosa et al., 2020). A mixed blood supply could

provide more vascular reserve and lower vulnerability to neuronal injury and
atrophy (Liebeskind, 2003). The uncus, which receives a distinct blood supply
compared to the rest of the hippocampus, might thus exhibit greater resistance to
the adverse effects of hypoxic-ischemic events than the other hippocampal
regions.

Uncus and recall memory
The residual uncus could be partly functional in patients with DA and could be
recruited for recall memory. We reported here that the volume of the uncus
correlated negatively with recall scores as well as working memory performance
in patients with DA. In the WAIS/WISC test, Working Memory Index is made up
of two subtests (Digit Span and Letter Number Sequencing) both requiring verbal
recall in the span of working memory. The function of the uncus is not well
understood mainly because its structure-function mapping remains to be
determined in rodents (Ding, 2013) but in humans, the uncus has been proposed to
be involved in the long-term recall of scenes (Zeidman et al., 2015). Connectivity
studies in nonhuman primates have shown that the uncus is less connected to
other regions of the hippocampus, and that it has its own associational system of
connections (Kondo et al., 2009). Notably, the uncus is also the only hippocampal
region reciprocally connected to its contralateral counterpart (Amaral et al., 1984).
The uncus' unique connectivity could allow this region to support some
information processing despite severe damage to the other hippocampal areas.
Notwithstanding the importance of the uncus in relation to severity of
atrophy during delayed recall, this subregion maintains its role during encoding
and learning of memoranda in both low and high severity subgroups equally.
Thus, we noted that despite their poor performance overall, both patient subgroups
incrementally increased their scores across the three learning trials of the *People*
and *Shapes* subtests of the *Doors and People Test*. The absence of a difference in
performance between the subgroups during encoding and learning, but not during
delayed recall, highlights the role of the uncus in retrieval of auditory verbal
memory after a delay in patients with DA.
Interestingly, we also reported an inverse relationship between the volume of
the uncus and patients' performance in delayed verbal, but not visual recall.
Impairment in delayed verbal recall in relation to extent of atrophy of the uncus
highlights the vulnerability of auditory verbal information to recall after a delay
(Schulze et al., 2012).

Subiculum
Residual circuits of the subicular complex might participate in spatial perception
and spatial memory in patients with DA. Here, we reported inverse correlations
between spatial perception/memory performance and the volume of the residual
subicular complex in patients with DA. The subiculum receives its major
excitatory input from CA1, but it also receives a strong projection from the
entorhinal cortex (Amaral and Lavenex, 2007; Matsumoto et al., 2019; Witter and
Amaral, 2020). The subiculum is also the main source of subcortical projections
directed to the hippocampus, and the presubiculum receives the only direct input
from the anterior thalamic nucleus (Amaral and Lavenex, 2007; Matsumoto et al.,
2019). The subiculum, presubiculum and parasubiculum send dense projections to
the entorhinal cortex (Saunders and Rosene, 1988; Witter and Amaral, 2020).

Thus, in the absence of functional CA-DG, the residual subicular complex could
marginally participate in input processing through its other interconnections, and
in particular, through interaction with the structurally preserved entorhinal cortex
which constitutes the main entryway for much of the neocortical input reaching
the hippocampus (Amaral et al., 1987).
Animal and human studies suggest that the subiculum plays an important role
in spatial processing and spatial representation (Matsumoto et al., 2019). Grid,
border, and head direction cells have been found in the subiculum of rodents and
primates including humans (Boccara et al., 2010; Lee et al., 2018; Lever et al.,
2009; Robertson et al., 1999). In the rat, cross-structural interaction between the
subiculum and the ventral striatum has been reported to be critical for spatial
memory consolidation (Torromino et al., 2019). In humans, the subiculum is
suggested to be specifically involved in scene-based cognitive processing,
including scene perception and scene construction during memory recall (Dalton
and Maguire, 2017; Hodgetts et al., 2017; Zeidman et al., 2015).
Interestingly, we did not observe correlations between the volume of the
subicular complex and patients' performance on the non-spatial tasks of the *Doors*
and People Test, suggesting that the subicular complex might be playing a more
prominent role in the processing of spatial inputs.
The Processing Speed Index of the Wechsler Scales is made up of two
subtests ('Coding' which requires learning and memory and 'Symbol Search'
which requires visual scanning/cancellation, and visual stimulus matching, but
without memory demands). Interestingly, the Coding scores are negatively
correlated with the uncus volume while the Symbol Search scores are negatively
correlated with both the uncus and the subicular complex. Our results tend to
demonstrate that the hippocampus is involved in two cognitive processes of visual
perception, and cognitive memory.

Working memory
We reported here, in addition to an absence of working memory impairment, a
negative association between uncus volume and working memory performance in
patients with DA. The more established view proposes that the hippocampus plays
an important role in episodic memory, spatial cognition, and navigation, while it
is not critically contributing to working memory (Eichenbaum, 2006; Eichenbaum
et al., 1994). But an alternative view proposes that the hippocampus may
contribute to any memory task, including working memory (Hannula and
Ranganath, 2008; Libby et al., 2014), in particular in supporting high-precision
binding (Borders et al., 2021). This latter possibility is challenged however by the
fact that patients with DA presenting with severe hippocampal damage have
repeatedly shown unimpaired working memory performance (Allen et al., 2014;
Baddeley et al., 2011, 2010). Recent work extended these findings by showing
that high resolution working memory performance was unimpaired, and even
often superior, relative to control group in a patient with DA (Allen et al., 2022).
Our observation of a negative correlation between uncus volume and working
memory performance suggests that the perirhinal, entorhinal, and
parahippocampal cortices normally involved in the binding of information in the
short term (Miyashita, 2019) may be subject to plastic changes in patients with
DA and that these plastic changes may depend on the degree of hippocampal
damage. The possibility that the hippocampal damage in patients with DA may

not influence working memory performance directly, but indirectly via
reorganization of the MTL circuit is explored below.

Functional hypotheses
Experimental studies in nonhuman primates have reported similar paradoxical
findings to those reported in the present study. Monkeys with small hippocampal
lesions exhibit greater memory loss on Delayed Nonmatching-to-Sample (DNMS)
task than that found in animals with greater damage (Bachevalier and Mishkin,
1989). Furthermore, a positive correlation between the extent of damage to the
hippocampus and scores on a DNMS task has been reported in experimentally
lesioned monkeys (Murray and Mishkin, 1998). Several hypotheses have been
proposed to explain why discrete lesions result in greater memory impairment
(Bachevalier and Mishkin, 1989; Baxter and Murray, 2001; Beason-Held et al.,
1999). However, it is difficult to relate these findings obtained following adult
experimental lesions in the monkey to observations in patients with early
hippocampal damage whose circuits might have undergone massive
compensation. One hypothesis to explain our observations is that the level of
hippocampal damage would likely have a strong direct influence on the
organization of the medial temporal lobe memory circuits in the adult. Following
an early event of hypoxia-ischemia, however, greater hippocampal damage might
induce greater compensatory reconfigurations in the neural circuits and enable
other structures, in particular the surrounding cortical areas normally recruited by
the hippocampus, to assume important aspects of memory function. In contrast,
when the hippocampus is only partially damaged, the information flow would
remain present in the hippocampus and could result in incomplete, and possibly
disruptive information processing. Our observation could thus reveal the existence
of multiple, redundant routes within the residual hippocampal subregions and
surrounding cortical areas in patients with developmental amnesia (Aggleton et
al., 2000). These parallel circuits could compete for the control of behavior and
disrupt perception and memory performance (Baxter and Murray, 2001).
Our findings are consistent with the hypothesis that a close relationship
can be evidenced between the volume of remaining hippocampal subregions and
the severity of cognitive deficits in patients with hippocampal damage, in the case
of patients with DA this relationship was inverse. Our observations also provide
further support for the model of hierarchical organization of cognitive memory
(Mishkin et al., 1997) by proposing that surrounding cortical areas normally
involved in the encoding of associative relations between decontextualized
information (Miyashita, 2019) could reorganize to contribute to hippocampal
function following early hippocampal damage. The limits of this reorganization,
however, may be determined by the extent of damage to the hippocampal
subregions.
Evidence for such reorganization has already been reported in a nonhuman
primate model of developmental amnesia. Monkeys with early bilateral
hippocampal lesion can learn new spatial relational information in striking
contrast to monkeys with adult lesion (Banta Lavenex et al., 2006; Lavenex et al.,
2007). Interestingly, in these monkeys with early hippocampal lesions, the
surrounding cortical areas were preserved and were shown to contribute to spatial
learning in the absence of functional hippocampal circuits (Chareyron et al.,
2017).

One could speculate that the functional reorganization of the perihippocampal structures (normally devoted to the processing of decontextualized information) to compensate hippocampal function could, in turn, benefit the processing of semantic memory and lead to better context-free memory performance in patients (see: Kapur et al., 2013). Indications of increased memory performance have indeed been reported in two different patients with DA. Jonin et al. (2018) studied patient 'KA' presenting with 55% hippocampal volume reduction and preserved perihippocampal structures. While patient KA displayed few, if any, residual episodic abilities, this patient was able to accurately retrieve semantic memories, and show evidence of superior or even very superior access to these memories than controls. Another patient with DA, patient 'Jon', has often produced response accuracy levels that are at least higher than those of control participants in previous experimental explorations of visuospatial working memory (Allen et al., 2022, 2014; Baddeley et al., 2010). Jon was at least numerically superior to the control mean on recognition-based measures of shape-color binding (Baddeley et al., 2010). In measures of color-location memory, Jon's recognition accuracy matched the highest achieving control participant, while his reconstruction performance was superior to 6 of the 7 controls (Allen et al., 2014). Across four tasks measuring binding between color and orientation or color and location using simultaneous or sequential presentation of stimuli, Jon's response accuracy was high, and always numerically superior to the control mean (Allen et al., 2022).

These observations from single patients studies together with the present findings, obtained in a large cohort of patients with DA, support the hypothesis of a reorganization of the preserved perihippocampal areas following early hippocampal damage.

Conclusions

Our results have important implications for understanding the organization of memory following early hippocampal injury. These observations highlight the paradoxical finding that in patients with DA greater hippocampal damage can be associated with better performance in recall memory, Verbal comprehension Index, Working memory Index, Processing speed Index and spatial perception. In contrast, less severe damage leaving behind relatively preserved hippocampal subregions can lead to more pronounced cognitive deficits. These unexpected findings can disentangle the functional organization of the medial temporal lobe memory system, and attribute a central role in the compensation of memory and spatial cognition to extra-hippocampal structures in patients with early hippocampal damage.

Materials and methods

Participants
The patients with DA were characterized by (i) a history of episodes of hypoxia-ischemia in early life, (ii) quantified hippocampal volume reduction above 25% relative to a group of controls, and (iii) severely impaired episodic memory (Dzieciol et al., 2017). The most common aetiology was perinatal hypoxic-

565   ischemic encephalopathy (Table S2). While long-term neurodevelopmental
566   outcome depends on the severity of the hypoxic-ischemic episode, subtle
567   cognitive deficits and behavioral alterations have been revealed through long-term
568   evaluations even in mild forms of hypoxia-ischemia (de Vries and Jongmans,
569   2010).
570   When all the volumetric estimates were completed, one of the patients,
571   who presented with moderate memory deficits and only 11.0% hippocampal
572   volume reduction associated with two unconfirmed episodes of prolonged seizures
573   at age 4, was excluded from the DA group.
574   Also, we excluded from the DA group a patient with temporal lobe
epilepsy presenting with a 25.8% hippocampal volume reduction, but without a
documented episode of early hypoxia-ischemia and/or status epilepticus.
Additionally, blinded neuroradiological examination carried out independently by
one of the authors (WKC) revealed that this patient's MRI scan was entirely
normal. The hippocampus, fornix and mammillary bodies were noted to have a
normal appearance in contrast to all other patients with DA.
581   We report here hippocampal volume estimates in 32 control participants
(16 females; mean age: 17.7 years, SD 7.7, range 8−38) and 23 patients with DA
(11 females; mean age: 19.5 years, SD 9.0, range 8−40).

Structural MRI
Participants were either scanned with a 1.5T-MRI scanner (24 control
participants, 16 patients with DA) or with a 3T-MRI scanner (8 control
participants, 7 patients with DA).
588   Whole brain structural 1.5T-MRI scans were obtained using a Siemens
589   Avanto Scanner, with a T1-weighted 3D FLASH sequence with the following
590   parameters: repetition time 11ms; echo time 4.94ms; in-plane resolution
591   1mm×1mm; slice thickness of 1mm (Cooper et al., 2011).
592   Whole brain structural 3T-MRI scans were obtained using a 3T Siemens
593   MRI system with a 20 channel head coil. A T1-weighted magnetization-prepared
594   rapid gradient-echo (MPRAGE) scan was acquired with the following parameters:
595   in-plane resolution of 1mm×1mm; slice thickness of 1mm; repetition time of
596   2,300ms; echo time of 2.74ms (Buck et al., 2019).
597   For the measurement of regions of interest (ROI) volumes, the data were
reformatted into 1 mm-thick contiguous slices in the coronal plane. ROI cross-
sectional areas were measured in all slices along the entire length of the
hippocampus, entorhinal, perirhinal and parahippocampal cortices using ITK-
SNAP (version 3.8.0; www.itksnap.org; Yushkevich et al., 2006). Manual tracing
is still the gold standard for measuring hippocampal volume and is particularly
appropriate for samples of relatively small size (Akudjedu et al., 2018), presenting
with various levels of hippocampal atrophy (Schmidt et al., 2018), or including
children and adolescents (Herten et al., 2019). All manual measurements of the
ROIs were carried out by one of the authors (LJC) who remained blind to
participant identity.
608   The hippocampus comprises the CA fields (CA3, CA2, CA1), the dentate
gyrus (DG), subiculum, presubiculum, and parasubiculum (Amaral and Lavenex,
2007). The human uncus [also termed the "uncal portion of the hippocampal
formation" (Amaral et al., 1984)] is situated antero-medially and consists of
slightly modified DG, CA3, CA2, CA1, and subiculum (Ding and Van Hoesen,

2015). For the segmentation of the hippocampus and its subregions, we followed
the guide provided by Dalton et al. (Dalton et al., 2017). Given the difficulty to
distinguish hippocampal CA fields on 1.5T and 3T-MRI acquisitions, we
restricted our volumetric analyses to three ROIs: uncus [defined according to
(Ding and Van Hoesen, 2015)], CA-DG (CA fields and dentate gyrus), and
subicular complex (subiculum, presubiculum, and parasubiculum). The
segmentation of the entorhinal, perirhinal and parahippocampal cortices was
based on structural descriptions of human medial temporal lobe and
segmentations protocols (Blaizot et al., 2010; Ding and Van Hoesen, 2010;
Frankó et al., 2014; Insausti et al., 1998; Kivisaari et al., 2020).
The volumes in cubic millimeters were calculated by summing the number
of voxels ($1 \times 1 \times 1$ mm) for each ROI. The volume of the hippocampus as a whole
was calculated as the sum of hippocampal subregions' volumes (uncus, CA-DG,
subicular complex). We then used parametric tests as volumetric estimates of the
hippocampal subregions and cortical areas were normally distributed (Shapiro-
Wilk normality test; control: all $P > .06$; DA: all $P > .09$). There was no
significant difference between volumetric estimates of the whole hippocampus or
cortical areas obtained with the 1.5T-MRI scanner and the 3T-MRI scanner in
control and DA groups (two-way ANOVA, both $F < 1.3$; both $P > .25$). Also there
was no significant gender difference for the volumes of the whole hippocampus or
cortical areas in control and DA groups (both $F < 0.08$; both $P > .79$). Volume
estimates on 1.5T and 3T-MRI scans and volume estimates in males and females
were thus presented together. There was a significant right larger than left volume
asymmetry for the volume of the whole hippocampus in both control and DA
groups (repeated measures two-way ANOVA: $F_{(1,53)} = 19.2$; $P < .001$; Tukey's
HSD post-hoc test: both $P < .004$). The right CA-DG subregions were
significantly larger than the left in both control and DA groups ($F_{(1,53)} = 26.1$; $P <$
.001; both $P < .004$). There was no significant asymmetry for the volume of the
other hippocampal subregions (all $P > .05$). Also, there was no significant
asymmetry for the volume of the entorhinal or perirhinal cortices (all $P > .12$) but
there was a significant right larger than left volume asymmetry for the volume of
the parahippocampal cortex in the control group ($F_{(1,53)} = 7.0$; $P = .011$; Tukey's
HSD post-hoc test: $P = .009$). The volume of all the ROIs was calculated as the
mean of the left and right hemisphere measurements.
Intracranial volume (ICV) was manually measured on the sagittal dataset with
a 1-in-10 random and systematic sampling strategy. The ICV was normally
distributed (Shapiro-Wilk normality test; control: $P = .05$; DA: $P = .23$). The ICV
was 6.8% smaller in the DA group than in the control group ($t_{(53)} = -2.70$; $P =$
.009). Corrections, derived from the regression line of control ROI volume (V)
versus ICV, were then made for ICV based on the formula $[\Delta V = a \times (\Delta ICV)]$ where
'a' is the regression coefficient.

WISC/WAIS
The Wechsler Adult Intelligence Scale, 3rd Ed. (WAIS-III) and the Wechsler
Intelligence Scale for Children, 4th Ed. (WISC-IV) provide four index scores:
Verbal Comprehension Index, measuring understanding and expression of verbal
concepts; Perceptual Reasoning Index, reflecting visuospatial perception and
perceptuomotor manipulation; Working Memory Index, tracking on-line
immediate memory and executive control; and Processing Speed Index,

measuring speed of visuoperceptual discrimination, and copying of number-symbol associations. The four index scores are expressed as standard scores with a mean of 100 and a standard deviation of 15.

Scoring. The WISC-IV or WAIS-III were administered to all patients with DA (N = 23, 11 females, mean age: 19.5 years, SD: 9.0, range: 8−40) and to 23 controls (13 females, mean age: 18.0 years, SD: 8.4, range: 8.2–38). For the analysis of the data, we then used parametric tests as the data were normally distributed in all four indexes (Shapiro–Wilk test of normality; Control: all $P > .36$; DA: all $P > .10$).

The Doors and People Test

The *Doors and People Test* was administered according to the instructions in the published manual (Baddeley et al., 1994), as described in detail in (Adlam et al., 2009; Patai et al., 2015). Briefly, for the verbal recall subtest ('People'), participants were presented with four pictures of people and after three learning trials asked to recall their names cued by their profession. For the visual recall subtest ('Shapes'), participants first copied, and drew the shapes from memory after three learning trials. For verbal recognition ('Names'), participants were presented with a list of names, and then asked to recognize each one from a list of four alternatives. Finally, in the visual recognition subtest ('Doors'), participants viewed a list of doors and then identified each one from a list of four alternatives. See Supplementary Information for more details.

Scoring. The *Doors and People Test* was administered to all patients with DA (N = 23, 11 females, mean age: 19.5 years, SD: 9.0, range: 8−40) and to 21 controls (11 females, mean age: 17.6 years, SD: 8.6, range: 8−38). One of the patients performed all but the 'Names' subtest ('People', 'Doors', 'Shapes': N = 23; 'Names': n = 22). Given that the DA group included individuals under the age of 16, which is the first age band at which standard scores are available on the *Doors and People Test*, we used standard scores based on the scores of our control group; raw scores (SD): People, 28.2 (6.9); Doors, 18.9 (3.8); Shapes, 34.6 (2.7); Names, 18.6 (3.7). We converted patients' and controls' raw scores to z-scores relative to the control group's scores. For the analysis of the data, we then used nonparametric tests as the data were not normally distributed in all four subtests (Shapiro–Wilk test of normality; Control: $P < .001$; DA: $P = .044$).

The Four Mountains Test

The *Four Mountains Test* was developed to examine hippocampal contribution to perception and short-term memory for topographical and nonspatial information in spatial scenes (Hartley et al., 2007). Briefly, this test involves computer-generated landscapes containing four mountains. In the 'Spatial Perception' task, participants selected one picture that matched the sample image (the same place from a different viewpoint) from four alternatives. In the 'Spatial Memory' task, the sample image was presented for approximately 8 sec. After a 2 sec. delay, participants were presented with four choices and selected the one previously viewed. See Supplementary Information for more details.

Scoring. The *Four Mountains Test* was administered to 14 patients with DA (7 females, mean age: 21.6 years, SD: 10.3, range: 8−40) and to 19 controls (11 females, mean age: 16.9 years, SD: 7.8, range: 8−35). We converted patients' raw scores to z-scores relative to the scores of our control group; raw scores (SD):

709 Perception, 11.6 (2.3); Memory, 9.95 (2.8). We then used nonparametric tests as
710 the data in the Control group were not normally distributed [Shapiro–Wilk test of
711 normality; $P < .05$].

Acknowledgments

713 This work was supported by the UK Medical Research Council (program grant
714 numbers G03000117 and G1002276).
715
716 We thank the participants and their families for their support of our research.
717

Competing interests

719 The authors declare no conflict of interest.

References

Adlam, A.-L.R., Malloy, M., Mishkin, M., Vargha-Khadem, F., 2009.
Dissociation between recognition and recall in developmental amnesia.
Neuropsychologia 47, 2207–2210.
https://doi.org/10.1016/j.neuropsychologia.2009.01.038

Aggleton, J.P., Vann, S.D., Oswald, C.J., Good, M., 2000. Identifying cortical
inputs to the rat hippocampus that subserve allocentric spatial processes: a simple
problem with a complex answer. Hippocampus 10, 466–474.
https://doi.org/10.1002/1098-1063(2000)10:4<466::AID-HIPO13>3.0.CO;2-Y

Akudjedu, T.N., Nabulsi, L., Makelyte, M., Scanlon, C., Hehir, S., Casey, H.,
Ambati, S., Kenney, J., O'Donoghue, S., McDermott, E., Kilmartin, L., Dockery,
P., McDonald, C., Hallahan, B., Cannon, D.M., 2018. A comparative study of
segmentation techniques for the quantification of brain subcortical volume. Brain
Imaging Behav 12, 1678–1695. https://doi.org/10.1007/s11682-018-9835-y

Allen, R.J., Atkinson, A.L., Vargha-Khadem, F., Baddeley, A.D., 2022. Intact
high-resolution working memory binding in a patient with developmental amnesia
and selective hippocampal damage. Hippocampus 32, 597–609.
https://doi.org/10.1002/hipo.23452

Allen, R.J., Vargha-Khadem, F., Baddeley, A.D., 2014. Item-location binding in
working memory: is it hippocampus-dependent? Neuropsychologia 59, 74–84.
https://doi.org/10.1016/j.neuropsychologia.2014.04.013

Amaral, D.G., Insausti, R., Cowan, W.M., 1987. The entorhinal cortex of the
monkey: I. Cytoarchitectonic organization. J Comp Neurol 264, 326–55.

Amaral, D.G., Insausti, R., Cowan, W.M., 1984. The commissural connections of
the monkey hippocampal formation. J. Comp. Neurol. 224, 307–336.
https://doi.org/10.1002/cne.902240302

Amaral, D.G., Lavenex, P., 2007. Hippocampal neuroanatomy, in: The
Hippocampus Book. Oxford University Press, Oxford, pp. 37–114.

Argyropoulos, G.P.D., Dell'Acqua, C., Butler, E., Loane, C., Roca-Fernandez, A.,
Almozel, A., Drummond, N., Lage-Martinez, C., Cooper, E., Henson, R.N.,
Butler, C.R., 2021. Functional Specialization of the Medial Temporal Lobes in
Human Recognition Memory: Dissociating Effects of Hippocampal versus
Parahippocampal Damage. Cereb Cortex bhab290.
https://doi.org/10.1093/cercor/bhab290

Bachevalier, J., Mishkin, M., 1989. Mnemonic and neuropathological effects of
occluding the posterior cerebral artery in Macaca mulatta. Neuropsychologia 27,
83–105. https://doi.org/10.1016/0028-3932(89)90092-4

Baddeley, A., Allen, R., Vargha-Khadem, F., 2010. Is the hippocampus necessary
for visual and verbal binding in working memory? Neuropsychologia 48, 1089–
1095. https://doi.org/10.1016/j.neuropsychologia.2009.12.009

Baddeley, A., Jarrold, C., Vargha-Khadem, F., 2011. Working memory and the
hippocampus. J Cogn Neurosci 23, 3855–3861.
https://doi.org/10.1162/jocn_a_00066

Baddeley, A.D., 2020. On trying to prove Endel Tulving wrong: A revised modal
model of amnesia. Neuropsychologia 148, 107590.
https://doi.org/10.1016/j.neuropsychologia.2020.107590

Baddeley, A.D., Emslie, H., Nimmo-Smith, I., 1994. Doors and people : a test of
visual and verbal recall and recognition. Bury St Edmunds : Thames Valley Test
Company, UK.

Banta Lavenex, P., Amaral, D.G., Lavenex, P., 2006. Hippocampal lesion
prevents spatial relational learning in adult macaque monkeys. J. Neurosci. 26,
4546–4558. https://doi.org/10.1523/JNEUROSCI.5412-05.2006

Bartsch, T., Döhring, J., Reuter, S., Finke, C., Rohr, A., Brauer, H., Deuschl, G.,
Jansen, O., 2015. Selective neuronal vulnerability of human hippocampal CA1
neurons: lesion evolution, temporal course, and pattern of hippocampal damage in
diffusion-weighted MR imaging. J. Cereb. Blood Flow Metab. 35, 1836–1845.
https://doi.org/10.1038/jcbfm.2015.137

Baxter, M.G., Murray, E.A., 2001. Opposite relationship of hippocampal and
rhinal cortex damage to delayed nonmatching-to-sample deficits in monkeys.
Hippocampus 11, 61–71. https://doi.org/10.1002/1098-1063(2001)11:1<61::AID-
HIPO1021>3.0.CO;2-Z

Bayley, P.J., Squire, L.R., 2005. Failure to acquire new semantic knowledge in
patients with large medial temporal lobe lesions. Hippocampus 15, 273–280.
https://doi.org/10.1002/hipo.20057

Beason-Held, L.L., Rosene, D.L., Killiany, R.J., Moss, M.B., 1999. Hippocampal
formation lesions produce memory impairment in the rhesus monkey.
Hippocampus 9, 562–574. https://doi.org/10.1002/(SICI)1098-
1063(1999)9:5<562::AID-HIPO10>3.0.CO;2-X

Blaizot, X., Mansilla, F., Insausti, A.M., Constans, J.M., Salinas-Alamán, A., Pró-
Sistiaga, P., Mohedano-Moriano, A., Insausti, R., 2010. The human
parahippocampal region: I. Temporal pole cytoarchitectonic and MRI correlation.
Cereb Cortex 20, 2198–2212. https://doi.org/10.1093/cercor/bhp289

Boccara, C.N., Sargolini, F., Thoresen, V.H., Solstad, T., Witter, M.P., Moser,
E.I., Moser, M.-B., 2010. Grid cells in pre- and parasubiculum. Nat. Neurosci. 13,
987–994. https://doi.org/10.1038/nn.2602

Borders, A.A., Ranganath, C., Yonelinas, A.P., 2021. The hippocampus supports
high-precision binding in visual working memory. Hippocampus.
https://doi.org/10.1002/hipo.23401

Buck, S., Bastos, F., Baldeweg, T., Vargha-Khadem, F., 2019. A Functional MRI
Paradigm Suitable for Language and Memory Mapping in Pediatric Temporal
Lobe Epilepsy. Front Neurol 10, 1384. https://doi.org/10.3389/fneur.2019.01384

Chareyron, L.J., Banta Lavenex, P., Amaral, D.G., Lavenex, P., 2017. Functional
organization of the medial temporal lobe memory system following neonatal
hippocampal lesion in rhesus monkeys. Brain Struct Funct 222, 3899–3914.
https://doi.org/10.1007/s00429-017-1441-z

Cooper, J.M., Gadian, D.G., Jentschke, S., Goldman, A., Munoz, M., Pitts, G.,
Banks, T., Chong, W.K., Hoskote, A., Deanfield, J., Baldeweg, T., de Haan, M.,
Mishkin, M., Vargha-Khadem, F., 2015. Neonatal hypoxia, hippocampal atrophy,
and memory impairment: evidence of a causal sequence. Cereb Cortex 25, 1469–
1476. https://doi.org/10.1093/cercor/bht332

Cooper, J.M., Vargha-Khadem, F., Gadian, D.G., Maguire, E.A., 2011. The effect
of hippocampal damage in children on recalling the past and imagining new
experiences. Neuropsychologia 49, 1843–1850.
https://doi.org/10.1016/j.neuropsychologia.2011.03.008

Dalton, M.A., Maguire, E.A., 2017. The pre/parasubiculum: a hippocampal hub
for scene-based cognition? Curr Opin Behav Sci 17, 34–40.
https://doi.org/10.1016/j.cobeha.2017.06.001

Dalton, M.A., Zeidman, P., Barry, D.N., Williams, E., Maguire, E.A., 2017.
Segmenting subregions of the human hippocampus on structural magnetic
resonance image scans: An illustrated tutorial. Brain Neurosci Adv 1,
2398212817701448. https://doi.org/10.1177/2398212817701448

de Vries, L.S., Jongmans, M.J., 2010. Long-term outcome after neonatal hypoxic-
ischaemic encephalopathy. Arch. Dis. Child. Fetal Neonatal Ed. 95, F220-224.
https://doi.org/10.1136/adc.2008.148205

Ding, S.-L., 2013. Comparative anatomy of the prosubiculum, subiculum,
presubiculum, postsubiculum, and parasubiculum in human, monkey, and rodent.
J. Comp. Neurol. 521, 4145–4162. https://doi.org/10.1002/cne.23416

Ding, S.-L., Van Hoesen, G.W., 2015. Organization and Detailed Parcellation of
Human Hippocampal Head and Body Regions Based on a Combined Analysis of
Cyto- and Chemoarchitecture. J. Comp. Neurol. 523, 2233–2253.
https://doi.org/10.1002/cne.23786

Ding, S.-L., Van Hoesen, G.W., 2010. Borders, extent, and topography of human
perirhinal cortex as revealed using multiple modern neuroanatomical and
pathological markers. Hum Brain Mapp 31, 1359–1379.
https://doi.org/10.1002/hbm.20940

Dzieciol, A.M., Bachevalier, J., Saleem, K.S., Gadian, D.G., Saunders, R., Chong,

W.K.K., Banks, T., Mishkin, M., Vargha-Khadem, F., 2017. Hippocampal and diencephalic pathology in developmental amnesia. Cortex 86, 33–44. https://doi.org/10.1016/j.cortex.2016.09.016

Eichenbaum, H., 2006. Remembering: functional organization of the declarative memory system. Curr Biol 16, R643-645. https://doi.org/10.1016/j.cub.2006.07.026

Eichenbaum, H., Otto, T., Cohen, N.J., 1994. Two functional components of the hippocampal memory system. Behavioral and Brain Sciences 17, 449–517. https://doi.org/10.1017/S0140525X00035391

Elward, R.L., Rugg, M.D., Vargha-Khadem, F., 2021. When the brain, but not the person, remembers: Cortical reinstatement is modulated by retrieval goal in developmental amnesia. Neuropsychologia 154, 107788. https://doi.org/10.1016/j.neuropsychologia.2021.107788

Elward, R.L., Vargha-Khadem, F., 2018. Semantic memory in developmental amnesia. Neurosci. Lett. 680, 23–30. https://doi.org/10.1016/j.neulet.2018.04.040

Erdem, A., Yaşargil, G., Roth, P., 1993. Microsurgical anatomy of the hippocampal arteries. J. Neurosurg. 79, 256–265. https://doi.org/10.3171/jns.1993.79.2.0256

Frankó, E., Insausti, A.M., Artacho-Pérula, E., Insausti, R., Chavoix, C., 2014. Identification of the human medial temporal lobe regions on magnetic resonance images. Hum Brain Mapp 35, 248–256. https://doi.org/10.1002/hbm.22170

Geva, S., Jentschke, S., Argyropoulos, G.P.D., Chong, W.K., Gadian, D.G., Vargha-Khadem, F., 2020. Volume reduction of caudate nucleus is associated with movement coordination deficits in patients with hippocampal atrophy due to perinatal hypoxia-ischaemia. Neuroimage Clin 28, 102429. https://doi.org/10.1016/j.nicl.2020.102429

Guderian, S., Dzieciol, A.M., Gadian, D.G., Jentschke, S., Doeller, C.F., Burgess, N., Mishkin, M., Vargha-Khadem, F., 2015. Hippocampal Volume Reduction in Humans Predicts Impaired Allocentric Spatial Memory in Virtual-Reality Navigation. J. Neurosci. 35, 14123–14131. https://doi.org/10.1523/JNEUROSCI.0801-15.2015

Hannula, D.E., Ranganath, C., 2008. Medial temporal lobe activity predicts successful relational memory binding. J Neurosci 28, 116–124. https://doi.org/10.1523/JNEUROSCI.3086-07.2008

Hartley, T., Bird, C.M., Chan, D., Cipolotti, L., Husain, M., Vargha-Khadem, F., Burgess, N., 2007. The hippocampus is required for short-term topographical memory in humans. Hippocampus 17, 34–48. https://doi.org/10.1002/hipo.20240

Herten, A., Konrad, K., Krinzinger, H., Seitz, J., von Polier, G.G., 2019. Accuracy and bias of automatic hippocampal segmentation in children and adolescents. Brain Struct Funct 224, 795–810. https://doi.org/10.1007/s00429-018-1802-2

Hodgetts, C.J., Voets, N.L., Thomas, A.G., Clare, S., Lawrence, A.D., Graham, K.S., 2017. Ultra-High-Field fMRI Reveals a Role for the Subiculum in Scene Perceptual Discrimination. J. Neurosci. 37, 3150–3159. https://doi.org/10.1523/JNEUROSCI.3225-16.2017

Huther, G., Dörfl, J., Van der Loos, H., Jeanmonod, D., 1998. Microanatomic and
vascular aspects of the temporomesial region. Neurosurgery 43, 1118–1136.
https://doi.org/10.1097/00006123-199811000-00065

Insausti, R., Juottonen, K., Soininen, H., Insausti, A.M., Partanen, K., Vainio, P.,
Laakso, M.P., Pitkänen, A., 1998. MR volumetric analysis of the human
entorhinal, perirhinal, and temporopolar cortices. AJNR Am J Neuroradiol 19,
659–671.

Jabès, A., Lavenex, P.B., Amaral, D.G., Lavenex, P., 2011. Postnatal
development of the hippocampal formation: a stereological study in macaque
monkeys. J. Comp. Neurol. 519, 1051–1070. https://doi.org/10.1002/cne.22549

Jabès, A., Nelson, C.A., 2015. 20 Years after "The ontogeny of human memory:
A cognitive neuroscience perspective," where are we? International Journal of
Behavioral Development 39, 293–303.
https://doi.org/10.1177/0165025415575766

Jonin, P.-Y., Besson, G., La Joie, R., Pariente, J., Belliard, S., Barillot, C.,
Barbeau, E.J., 2018. Superior explicit memory despite severe developmental
amnesia: In-depth case study and neural correlates. Hippocampus 28, 867–885.
https://doi.org/10.1002/hipo.23010

Kapur, N., Cole, J., Manly, T., Viskontas, I., Ninteman, A., Hasher, L., Pascual-
Leone, A., 2013. Positive clinical neuroscience: explorations in positive
neurology. Neuroscientist 19, 354–369.
https://doi.org/10.1177/1073858412470976

Kivisaari, S.L., Probst, A., Taylor, K.I., 2020. The Perirhinal, Entorhinal, and
Parahippocampal Cortices and Hippocampus: An Overview of Functional
Anatomy and Protocol for Their Segmentation in MR Images, in: Ulmer, S.,
Jansen, O. (Eds.), FMRI: Basics and Clinical Applications. Springer International
Publishing, Cham, pp. 355–383. https://doi.org/10.1007/978-3-030-41874-8_24

Kondo, H., Lavenex, P., Amaral, D.G., 2009. Intrinsic connections of the
macaque monkey hippocampal formation: II. CA3 connections. J. Comp. Neurol.
515, 349–377. https://doi.org/10.1002/cne.22056

Lavenex, P., Lavenex, P.B., Amaral, D.G., 2007. Spatial relational learning
persists following neonatal hippocampal lesions in macaque monkeys. Nat.
Neurosci. 10, 234–239. https://doi.org/10.1038/nn1820

Lee, S.A., Miller, J.F., Watrous, A.J., Sperling, M.R., Sharan, A., Worrell, G.A.,
Berry, B.M., Aronson, J.P., Davis, K.A., Gross, R.E., Lega, B., Sheth, S., Das,
S.R., Stein, J.M., Gorniak, R., Rizzuto, D.S., Jacobs, J., 2018.
Electrophysiological Signatures of Spatial Boundaries in the Human Subiculum.
J. Neurosci. 38, 3265–3272. https://doi.org/10.1523/JNEUROSCI.3216-17.2018

Lever, C., Burton, S., Jeewajee, A., O'Keefe, J., Burgess, N., 2009. Boundary
vector cells in the subiculum of the hippocampal formation. J. Neurosci. 29,
9771–9777. https://doi.org/10.1523/JNEUROSCI.1319-09.2009

Libby, L.A., Hannula, D.E., Ranganath, C., 2014. Medial temporal lobe coding of
item and spatial information during relational binding in working memory. J
Neurosci 34, 14233–14242. https://doi.org/10.1523/JNEUROSCI.0655-14.2014

Liebeskind, D.S., 2003. Collateral circulation. Stroke 34, 2279–2284.
https://doi.org/10.1161/01.STR.0000086465.41263.06

Maguire, E.A., Vargha-Khadem, F., Mishkin, M., 2001. The effects of bilateral
hippocampal damage on fMRI regional activations and interactions during
memory retrieval. Brain 124, 1156–1170.
https://doi.org/10.1093/brain/124.6.1156

Manns, J.R., Hopkins, R.O., Squire, L.R., 2003. Semantic Memory and the
Human Hippocampus. Neuron 38, 127–133. https://doi.org/10.1016/S0896-
6273(03)00146-6

Marinković, S., Milisavljević, M., Puskas, L., 1992. Microvascular anatomy of the
hippocampal formation. Surg Neurol 37, 339–349. https://doi.org/10.1016/0090-
3019(92)90001-4

Matsumoto, N., Kitanishi, T., Mizuseki, K., 2019. The subiculum: Unique
hippocampal hub and more. Neurosci. Res. 143, 1–12.
https://doi.org/10.1016/j.neures.2018.08.002

Mishkin, M., Suzuki, W.A., Gadian, D.G., Vargha-Khadem, F., 1997.
Hierarchical Organization of Cognitive Memory. Philosophical Transactions:
Biological Sciences 352, 1461–1467.

Miyashita, Y., 2019. Perirhinal circuits for memory processing. Nature Reviews
Neuroscience 20, 577–592. https://doi.org/10.1038/s41583-019-0213-6

Murray, E.A., Mishkin, M., 1998. Object recognition and location memory in
monkeys with excitotoxic lesions of the amygdala and hippocampus. J. Neurosci.
18, 6568–6582.

Patai, E.Z., Gadian, D.G., Cooper, J.M., Dzieciol, A.M., Mishkin, M., Vargha-
Khadem, F., 2015. Extent of hippocampal atrophy predicts degree of deficit in
recall. Proc. Natl. Acad. Sci. U.S.A. 112, 12830–12833.
https://doi.org/10.1073/pnas.1511904112

Perosa, V., Priester, A., Ziegler, G., Cardenas-Blanco, A., Dobisch, L., Spallazzi,
M., Assmann, A., Maass, A., Speck, O., Oltmer, J., Heinze, H.-J., Schreiber, S.,
Düzel, E., 2020. Hippocampal vascular reserve associated with cognitive
performance and hippocampal volume. Brain 143, 622–634.
https://doi.org/10.1093/brain/awz383

Picard, L., Mayor-Dubois, C., Maeder, P., Kalenzaga, S., Abram, M., Duval, C.,
Eustache, F., Roulet-Perez, E., Piolino, P., 2013. Functional independence within
the self-memory system: new insights from two cases of developmental amnesia.
Cortex 49, 1463–1481. https://doi.org/10.1016/j.cortex.2012.10.003

Rabin, J.S., Olsen, R.K., Gilboa, A., Buchsbaum, B.R., Rosenbaum, R.S., 2016.
Using fMRI to understand event construction in developmental amnesia.
Neuropsychologia 90, 261–273.
https://doi.org/10.1016/j.neuropsychologia.2016.07.036

Rempel-Clower, N.L., Zola, S.M., Squire, L.R., Amaral, D.G., 1996. Three cases
of enduring memory impairment after bilateral damage limited to the hippocampal
formation. J. Neurosci. 16, 5233–5255.

Robertson, R.G., Rolls, E.T., Georges-François, P., Panzeri, S., 1999. Head
direction cells in the primate pre-subiculum. Hippocampus 9, 206–219.
https://doi.org/10.1002/(SICI)1098-1063(1999)9:3<206::AID-HIPO2>3.0.CO;2-
H

Rosenbaum, R.S., Cassidy, B.N., Herdman, K.A., 2015. Patterns of preserved and
impaired spatial memory in a case of developmental amnesia. Front Hum
Neurosci 9, 196. https://doi.org/10.3389/fnhum.2015.00196

Rosene, D.L., Van Hoesen, G.W., 1977. Hippocampal efferents reach widespread
areas of cerebral cortex and amygdala in the rhesus monkey. Science 198, 315–
317.

Saunders, R.C., Rosene, D.L., 1988. A comparison of the efferents of the
amygdala and the hippocampal formation in the rhesus monkey: I. Convergence
in the entorhinal, prorhinal, and perirhinal cortices. Journal of Comparative
Neurology 271, 153–184. https://doi.org/10.1002/cne.902710202

Schmidt, M.F., Storrs, J.M., Freeman, K.B., Jack, C.R., Turner, S.T., Griswold,
M.E., Mosley, T.H., 2018. A comparison of manual tracing and FreeSurfer for
estimating hippocampal volume over the adult lifespan. Hum Brain Mapp 39,
2500–2513. https://doi.org/10.1002/hbm.24017

Schmidt-Kastner, R., Freund, T.F., 1991. Selective vulnerability of the
hippocampus in brain ischemia. Neuroscience 40, 599–636.
https://doi.org/10.1016/0306-4522(91)90001-5

Schulze, K., Vargha-Khadem, F., Mishkin, M., 2012. Test of a motor theory of
long-term auditory memory. Proc. Natl. Acad. Sci. U.S.A. 109, 7121–7125.
https://doi.org/10.1073/pnas.1204717109

Scoville, W.B., Milner, B., 1957. Loss of recent memory after bilateral
hippocampal lesions. J. Neurol. Neurosurg. Psychiatr. 20, 11–21.

Shrager, Y., Levy, D.A., Hopkins, R.O., Squire, L.R., 2008. Working memory
and the organization of brain systems. J Neurosci 28, 4818–4822.
https://doi.org/10.1523/JNEUROSCI.0710-08.2008

Torromino, G., Autore, L., Khalil, V., Mastrorilli, V., Griguoli, M., Pignataro, A.,
Centofante, E., Biasini, G.M., De Turris, V., Ammassari-Teule, M., Rinaldi, A.,
Mele, A., 2019. Offline ventral subiculum-ventral striatum serial communication
is required for spatial memory consolidation. Nat Commun 10, 5721.
https://doi.org/10.1038/s41467-019-13703-3

Vargha-Khadem, F., Cacucci, F., 2021. A brief history of developmental amnesia.
Neuropsychologia 150, 107689.
https://doi.org/10.1016/j.neuropsychologia.2020.107689

Vargha-Khadem, F., Gadian, D.G., Watkins, K.E., Connelly, A., Van Paesschen,
W., Mishkin, M., 1997. Differential effects of early hippocampal pathology on
episodic and semantic memory. Science 277, 376–380.
https://doi.org/10.1126/science.277.5324.376

Vargha-Khadem, F., Salmond, C.H., Watkins, K.E., Friston, K.J., Gadian, D.G.,
Mishkin, M., 2003. Developmental amnesia: effect of age at injury. Proc Natl
Acad Sci U S A 100, 10055–10060. https://doi.org/10.1073/pnas.1233756100

Verfaellie, M., Koseff, P., Alexander, M.P., 2000. Acquisition of novel semantic
information in amnesia: effects of lesion location. Neuropsychologia 38, 484–492.
https://doi.org/10.1016/S0028-3932(99)00089-5

Wang, X., Michaelis, E.K., 2010. Selective neuronal vulnerability to oxidative
stress in the brain. Front Aging Neurosci 2, 12.
https://doi.org/10.3389/fnagi.2010.00012

Witter, M.P., Amaral, D.G., 2020. The entorhinal cortex of the monkey: VI.
Organization of projections from the hippocampus, subiculum, presubiculum, and
parasubiculum. Journal of Comparative Neurology doi: 10.1002/cne.24983.
https://doi.org/10.1002/cne.24983

Yushkevich, P.A., Piven, J., Hazlett, H.C., Smith, R.G., Ho, S., Gee, J.C., Gerig,
G., 2006. User-guided 3D active contour segmentation of anatomical structures:
significantly improved efficiency and reliability. Neuroimage 31, 1116–1128.
https://doi.org/10.1016/j.neuroimage.2006.01.015

Zeidman, P., Lutti, A., Maguire, E.A., 2015. Investigating the functions of
subregions within anterior hippocampus. Cortex 73, 240–256.
https://doi.org/10.1016/j.cortex.2015.09.002

Zeidman, P., Maguire, E.A., 2016. Anterior hippocampus: the anatomy of
perception, imagination and episodic memory. Nat. Rev. Neurosci. 17, 173–182.
https://doi.org/10.1038/nrn.2015.24

Zola-Morgan, S., Squire, L.R., Amaral, D.G., 1986. Human amnesia and the
medial temporal region: enduring memory impairment following a bilateral lesion
limited to field CA1 of the hippocampus. J. Neurosci. 6, 2950–2967.

1034

1035