Title

Paradoxical consequences of early hippocampal damage: greater atrophy is associated with better recall, working memory and visuospatial perception in developmental amnesia

Authors/Affiliations

Loïc J. Chareyron[a], W.K. Kling Chong[b], Tina Banks[c], Mortimer Mishkin[d], Neil Burgess[e], Richard C. Saunders[d], Faraneh Vargha-Khadem[a]*

[a] Cognitive Neuroscience and Neuropsychiatry, Developmental Neurosciences, University College London Great Ormond Street Institute of Child Health, London, UK

[b] Developmental Imaging & Biophysics, University College London Great Ormond Street Institute of Child Health, London, UK

[c] Developmental Imaging & Biophysics, University College London Great Ormond Street Institute of Child Health, London, UK

[d] Laboratory of Neuropsychology, National Institute of Mental Health, National Institutes of Health, Bethesda, MD 20892.

[e] Institute of Cognitive Neuroscience, University College London, London, UK.

Corresponding Author

* Faraneh Vargha-Khadem.

Email: f.vargha-khadem@ucl.ac.uk

Abstract

Despite bilateral hippocampal damage dating to perinatal or early-childhood period, patients with developmental amnesia (DA) exhibit well-developed semantic memory, but severely impaired episodic memory. It is not clear, however, whether the residual hippocampus plays a role in encoding and/or retrieval of new information, or the surrounding cortical areas rescue aspects of these critical cognitive memory processes after early injury. We used manual MRI segmentation to estimate the volume of three hippocampal subregions and three surrounding cortical areas in 23 patients with DA and 32 controls. In patients, the level of atrophy in CA-DG subregions and subicular complex was more than 40% while the atrophy of the uncus was moderate (-23%). In contrast, volumes of entorhinal, perirhinal and parahippocampal cortices were smaller than those of controls, but not statistically different. Patients' recall, Verbal IQ, Working Memory Index and Processing Speed Index scores correlated negatively with the volume of the uncus while spatial perception/immediate memory performance correlated negatively with the volume of the subicular complex. We propose that in patients with DA, no other structure can perform the function of the hippocampus in episodic memory. However, cognitive memory processing is compromised as a function of extent of atrophy in hippocampal subregions, such that the greater the damage, the more likely that preserved surrounding cortical areas will be recruited to rescue the putative functions of the damaged subregions. Our findings document for the first time not only the extent, but also the limits of circuit reorganization occurring in the young brain following bilateral hippocampal damage.

Introduction

Developmental amnesia

We previously identified a group of patients with hippocampal atrophy who had suffered hypoxic-ischaemic episodes in infancy or childhood and later developed a memory disorder differing from that commonly described in patients with adult-onset amnesia of temporal-lobe origin. In the latter cases, the disorder most often takes the form of a global anterograde memory loss [i.e. one affecting both episodic and new post-injury semantic memory, and severely restricting recall and recognition processes (Bayley and Squire, 2005; Manns et al., 2003; Scoville and Milner, 1957; Verfaellie et al., 2000)]. In patients with hippocampal damage of early-onset, by contrast, the disorder is more limited, being characterized by markedly impaired episodic memory, recall/recollection, and spatial processing deficits, but relatively preserved semantic memory, working memory, and recognition performance. We have labeled this dissociated form of memory function 'developmental amnesia' (DA, Guderian et al., 2015; Patai et al., 2015; Picard et al., 2013; Rosenbaum et al., 2015; Vargha-Khadem et al., 1997). Given the early onset of hippocampal damage, the compensatory organization that has occurred in patients with DA is likely to be distinct from what is observed in patients with hippocampal damage acquired in adulthood. Also, in patients with acquired hippocampal lesions, the damage has occurred to already-established, and normally-functioning memory circuits, whereas in patients with DA, the early bilateral lesions have probably led to the development of a different memory system. For these reasons, it is difficult, and somewhat inappropriate, to compare memory function and cognitive profiles in patients with DA to those of patients with adult-onset amnesia (Elward and Vargha-Khadem, 2018, P: 24-26).

To date, a cohort of 18 patients with DA has been reported, each patient having suffered an episode of hypoxia-ischaemia in early life (Dzieciol et al., 2017). Importantly, irrespective of the underlying aetiology (Geva et al., 2020), or age at onset (Vargha-Khadem et al., 2003), the hypoxic-ischaemic event was the cause of hippocampal damage and cognitive impairment (Cooper et al., 2015).

Memory circuit reorganization

Among the many questions that have arisen from the study of patients with DA, the foremost concerns the regions of the medial temporal lobe (MTL) and/or preserved hippocampal subregions that could be involved in memory circuit reorganization following early hippocampal damage.

In the model of hierarchical organization of cognitive memory proposed by Mishkin and colleagues (1997; see also: Baddeley, 2020; Vargha-Khadem and Cacucci, 2021), recollection and familiarity depend on different MTL structures. According to this model, the perihippocampal gyrus, and in particular the perirhinal cortex, may support familiarity, while the hippocampus may support recollection (Argyropoulos et al., 2021). The selectivity of impairment observed in patients with DA has provided strong support for this dual-process model.

Because patients with DA have both neuroradiological and quantitatively-confirmed bilateral hippocampal atrophy, but no visible abnormality within the parahippocampal region, it has been proposed that their episodic memory impairment, and recall/recollection deficits are due to their hippocampal pathology, whereas their relatively intact semantic memory, and recognition/familiarity processes are due in part to the preservation of the perihippocampal gyrus.

Hippocampus
Hypoxia-ischemia is known to differentially affect the hippocampal fields and
subdivisions. Studies of animal models of cerebral ischemia have reported that a brief
episode results in selective neuronal death in the CA1 field of the hippocampus (Schmidt-
Kastner and Freund, 1991) while the adjacent CA3 remains less vulnerable (Wang and
Michaelis, 2010). In humans, while damage to CA1 is consistently reported following
cerebral ischemia, damage to additional hippocampal regions is much more variable
(Bartsch et al., 2015). Postmortem examinations of patients who suffered ischemic
episodes in adulthood have reported neural cell loss either restricted to the CA1 region, or
distributed in the dentate gyrus, subiculum, uncus and entorhinal cortex (Rempel-Clower
et al., 1996; Zola-Morgan et al., 1986). The preservation of circumscribed hippocampal
subregions following injury of early-onset in DA could account for the increased
activation of the residual hippocampus during different memory tasks (Maguire et al.,
2001; Rabin et al., 2016), although activation *per se* does not necessarily imply
correspondingly intact function (Elward et al., 2021).
In the study reported by Dzieciol and colleagues (Dzieciol et al., 2017), the degree
of hippocampal atrophy in patients with DA ranged from 28% to 62% compared to
healthy controls. This large range of hippocampal volume reduction in this cohort
suggests that there may indeed be variability in the hippocampal response to hypoxia-
ischemia at the level of hippocampal subregions. It would thus be possible to assess
whether the compensatory response in patients with DA is dependent on the extent of
atrophy in the hippocampal subregions. One way to address this is to evaluate the
association between task-dependent neuropsychological performance and volumes of
hippocampal subregions
Perihippocampal gyrus
While postmortem examinations have reported some neural cell loss in layers of
entorhinal cortex in patients who suffered ischemic episodes in adulthood (Rempel-
Clower et al., 1996), it was reported that the volume of the parahippocampal gyrus was
not significantly reduced in patients with adult-onset amnesia presenting with bilateral
hippocampal damage (Shrager et al., 2008). A cortical thickness analysis performed in a
single patient with DA didn't find abnormality in the subhippocampal structures within
the medial temporal lobe, including the perirhinal and ventral entorhinal cortices (Jonin et
al., 2018). Also, a recent study demonstrated that parahippocampal activity during scene
reinstatement in patients with DA was similar to controls (Elward et al., 2021). However,
the structural integrity of the perihippocampal gyrus, and by implication, the integrity of
the hypothesized substrate of semantic memory and other preserved mnemonic processes
remain to be quantitatively assessed in patients with DA.
127
Here, we have used manual segmentation of MRI scans to obtain estimates of the volume
of three hippocampal subregions: the uncus [uncal portion of the hippocampus (Zeidman
and Maguire, 2016)], the CA-DG (including CA fields and dentate gyrus) and the
subicular complex (including subiculum, presubiculum and parasubiculum), and three
surrounding cortical areas (entorhinal, perirhinal and parahippocampal cortices) in
patients with DA and healthy controls. Controls and patients were assessed with
neuropsychological tests of intelligence (WISC or WAIS), recall/recognition for verbal
and visual material (Doors and People Test) and spatial perception/memory (Four
Mountains Test). Finally, we investigated the relationships between degree of atrophy of
hippocampal subregions and cognitive deficits.

We predicted that: (i) residual hippocampal subregions and/or surrounding
cortical areas were sufficiently preserved to rescue some aspects of hippocampal
function, and (ii) plastic changes in these regions occurred as a function of hippocampal
damage in all patients with DA.
We documented several paradoxical inverse relationships: i) between the volume
of the residual uncus and patients' recall memory performance, Verbal Comprehension
Index, Working memory Index and Processing Speed Index, and ii) between the volume
of the residual subicular complex and patients' performance on spatial
perception/memory. We also reported an absence of structural damage in the surrounding
cortical areas in patients with DA.

148   Results

149   *Volumetric estimates*
150   Three hippocampal subregions (uncus, CA-DG and subicular complex) and three
151   surrounding cortical areas (entorhinal, perirhinal and parahippocampal cortices) were
152   manually segmented in healthy controls (N = 32) and DA (N = 23) groups (Fig. S1).
153

154
155   **Fig. 1. Volume of the hippocampus and its subregions.** Volume of the hippocampus (**A.**) and its
156   subregions (**B.**, uncus, CA-DG, and subicular complex) in control (grey; N = 32) and DA (orange; N =
157   23) groups. Data are represented as mean ± SD. Percentage values refer to the mean difference to
158   control group. Volumes are calculated as the average of left and right hemisphere volumes. All
159   volumes are corrected for intracranial (ICV) volume. *: P < .05; ***: P < .001.

160
The volume of the hippocampus as a whole was 40% smaller in the DA than in the
control group (range: −26% to −53%; $t_{(53)}$ = −15.32; P < .001) (Fig. 1; Table S1). The
volume of the hippocampal subregions was also significantly smaller in the DA than in
the control group (2-way ANOVA, $F_{(1,159)}$ = 392.5, P < .001). The uncus was 23%
smaller (range: +4% to −49%; Tukey's HSD post-hoc test: P = .037), the CA-DG
subregions were 45% smaller (range: −28% to −59%; P < .001), and the subicular
complex was 41% smaller (range: −22% to −56%; P < .001). The level of atrophy of the
patients' hippocampal subregions differed significantly from each other ($F_{(2,66)}$ = 23.0, P
< .001). Tukey's HSD post-hoc test indicated that the CA-DG and subicular complex
showed greater atrophy than the uncus (both P < .001).
Atrophy of the hippocampus in the DA group was significantly lower in its
anterior (−36%) than its posterior (−44%) segment (paired t test, $t_{(22)}$ = 2.72, P = .012;
Fig. S2). Because no differences were observed between the anterior and the posterior

segment of the CA-DG (paired t test, $t_{(22)} = 0.94$, $P = .36$) or subicular complex ($t_{(22)} = 1$,
$P = .33$) subregions, the lower level of atrophy observed in the anterior part of the whole
hippocampus can be attributed to the relatively more preserved uncus.
In contrast, there was no significant difference in the volume of the entorhinal,
perirhinal and parahippocampal cortices between control and DA groups (Table S1; 2-
way ANOVA, $F_{(1,159)} = 6.7$, $P = .010$; Tukey's HSD post-hoc tests, all $P > .35$). Also, we
didn't observe any correlation between the volume of the hippocampus or hippocampal
subregions and the volume of surrounding cortical areas in patients with DA (Pearson's
correlation analysis, all $P > .19$).

Neuroradiological assessment
Neuroradiological ratings based on visual inspection of the MR images provided
independently by one of the authors (WKC) revealed abnormally small fornix and
mammillary bodies in all but two patients with DA (Table S2). No abnormalities were
detected in the parahippocampal gyrus (perirhinal, entorhinal, and parahippocampal
cortices), or the thalamus, and the basal ganglia. Additional visible abnormality was
observed in some cases in the white matter and the cerebellum. The pattern of visible
abnormality detected within the hippocampal circuit, and the notable absence thereof in
other cortical and subcortical structures, was consistent across individual patients
irrespective of their varying aetiology, and age at hypoxic-ischaemic-induced
hippocampal damage.

WISC/WAIS
All patients with DA ($N = 23$) and 23 of the control participants were assessed on the
WISC or WAIS. Patients were impaired relative to our group of controls only for the
Perceptual Reasoning Index scores (2-way ANOVA, $F_{(1,176)} = 10.61$, $P = .001$; Tukey's
HSD post-hoc test, $P = 0.04$; all other $P > .25$). There were significant inverse
correlations between the volume of the uncus and patients' scores for the Verbal
Comprehension Index (VCI), the Working Memory Index (WMI) and the Processing
Speed Index (PSI) (all $r_{(22)} < –0.42$; all $P < .05$; Fig 2). Looking at PSI scores, we
observed an inverse correlations between "Digit Symbol-Coding" score and the volume
of the uncus ($r_{(22)} = –0.44$; $P = .037$) and between "Symbol Search" and the volume of
the uncus and subicular complex ($r_{(22)} = –0.43$; $P = .04$ and $r_{(22)} = –0.44$; $P = .037$,
respectively; Table S3). No other correlations for other ROIs in the DA group were
significant. In controls, the WMI scores were positively associated with the volume of the
subicular complex ($r_{(22)} = 0.46$; $P = .029$).
208

Fig. 2. Inverse correlations between WISC/WAIS indexes scores and the volume of the uncus in patients with DA. Participants' scores on Wechsler Intelligence Scales WISC or WAIS (patients: orange circles, N = 23; controls: grey triangles, n = 23). **A.** Verbal comprehension Index; **B.** Perceptual Reasoning Index; **C.** Working Memory Index; **D.** Processing Speed Index. Pearson's correlation coefficient. All volumes are corrected for intracranial (ICV) volume. *: P < .05.

Doors and People Test

All patients with DA (N = 23) and 21 of the control participants were assessed on the Doors and People Test, which provides equated measures of recognition and recall in the visual and verbal domains. All patients' subtest scores were significantly below zero (Wilcoxon signed rank test: all Z < −2.4, all P < .016), indicating that patients were impaired relative to controls (who represent zero, as their scores were used to convert raw scores into z-scores). The patients' scores for the four Doors and People subtests differed significantly from each other (Friedman's test: $\chi^2_{(3,n=22)}$ = 43.7, P < .001). The visual recall ('Shapes') subtest yielded a significantly greater deficit than all of the other subtests (all P < .001). The degree of deficit on the verbal recall ('People') subtest was greater than on the two recognition subtests (both P < .02). The degree of deficit was greater on the visual recognition ('Doors') subtest than on the verbal recognition ('Names') subtest (P = .001). Collapsing across subtests, we found a greater deficit in recall compared with recognition (Z = −4.04; P < .001), and a greater deficit in memory for visual compared with memory for verbal material (Z = −4.11; P < .001).

　　We observed significant negative correlations between the two recall subtest scores and the volume of the uncus in the DA group [Verbal recall ('People'): Pearson's $r_{(22)}$ = –0.47; P = .02; Visual recall ('Shapes'): $r_{(22)}$ = –0.55; P = .007]. No other correlations for other subtests or other regions of interest in the DA group were significant (all P > .09; Table S4). Collapsing across test material confirmed that scores on recall were inversely correlated with the volume of the uncus in the DA group ($r_{(22)}$ = –0.58; P = .004; Fig. 3A). As a corollary, we observed that patients with more preserved uncus had significantly lower scores on recall than patients with more damaged uncus (Z = −2.74; P = .006). In the control group, no correlations for any subtests or regions of interest were significant (all P > .11; Table S4).

Fig. 3. *Doors and People* recall scores correlations with the volumes of hippocampal subregions. The patients' scores (orange circles; N = 23) on recall subtests of the *Doors and People Test* correlated negatively with the volume of the uncus (**A.**) but not with the volumes of the subicular complex (**B.**) or CA-DG (**C.**). The controls' scores (grey triangles; n = 21) did not correlate with the volume of any subregion. Pearson's correlation coefficient. All volumes are corrected for intracranial (ICV) volume. **: P < .01.

During *Doors and People* testing, the procedures for recall subtests are repeated until
all items (names or shapes) are correctly recalled, or for a maximum of three
presentations. The patients' scores differed significantly from each other between the
three trials of the visual recall ('Shapes') subtest (Friedman's test: $\chi^2_{(2,n=20)} = 8.00$, P =
.018) and between the three trials of the verbal recall ('People') subtest ($\chi^2_{(2,n=20)} = 14.15$,
P < .001). Post hoc Wilcoxon signed-ranks tests indicated that scores obtained on trial-1
were significantly lower than scores obtained on trial-3 for the visual recall ('Shapes')
subtest (Z = −2.72; P = .006) and for the verbal recall ('People') subtest (Z = −2.88; P =
.004). Interested to determine potential differences in learning between patients
depending on whether they presented with severe or moderate uncus atrophy, we
compared increases in patients' scores between trial-1 and trial-3. There was no
difference in learning scores between patients with a larger uncus (above group mean
volume) and patients with a smaller uncus (Mann-Whitney U test; visual recall
('Shapes'): Z = −1.26; P = .21; verbal recall ('People'): Z = 0.61; P = .54).
Delayed (cued) verbal and visual recall were tested after completion of the verbal and
visual recognition subtests, respectively. We found a correlation between patients' scores
for the delayed verbal recall ('People') subtest and the volume of the uncus (Pearson's
$r_{(20)} = -0.5$; P = .02) while the correlation with delayed visual recall ('Shapes') scores
was not significant ($r_{(20)} = -0.3$; P = .18).

Four Mountains Test
Fourteen patients with DA and 19 of the control participants were assessed on the *Four*
Mountains Test to examine their perception and immediate memory for topographical
information in spatial scenes. Patients' subtest scores converted to z-scores were not
significantly below zero (Wilcoxon signed rank test: 'Spatial Perception': P = .75;
'Spatial Memory': P = .13), indicating that many of the patients were not impaired
relative to our group of controls. Patients' scores were significantly lower for the 'Spatial
Memory' than for the 'Spatial Perception' subtest (Z = −2.67; P = .008). There were
significant inverse correlations between patients' scores for the 'Spatial Perception' and
'Spatial Memory' subtests and the volume of the subicular complex ($r_{(13)} = -0.59$; P =
.025 and $r_{(13)} = -0.58$; P = .03, respectively). Collapsing across test material confirmed
that scores were strongly inversely correlated with the volume of the subicular complex
in the DA group ($r_{(13)} = -0.63$; P = .015) (Fig. 4B). Patients' scores for the 'Spatial
Perception' subtest were also negatively associated with the volume of the uncus ($r_{(13)} = -$
0.54 ; P = .047). In the control group, no correlations for any ROIs were significant (all P
> .49; Table S5).

Fig. 4. *Four Mountains* **scores correlations with the volumes of hippocampal subregions.** The
patients' scores (orange circles; n = 14) correlated negatively with the volume of the subicular
complex (**B.**) but not with the volumes of the uncus (**A.**) or CA-DG (**C.**). The controls' scores (grey
triangles; n = 19) did not correlate with the volume of any subregion. Pearson's correlation coefficient.
All volumes are corrected for intracranial (ICV) volume.

Discussion

In this study, we addressed the question of whether the volumes of residual hippocampal
subregions and parahippocampal areas are associated with memory and spatial processing
performance in patients with hypoxic-ischaemic-induced hippocampal damage of early-
onset. Using a combination of neuropsychological test results and volumetric estimations
in a large cohort of patients with developmental amnesia (DA), we documented several
inverse relationships: i) between the volume of the residual uncus and patients'
performance on recall memory, Verbal comprehension Index, Working memory Index
and Processing speed Index, and ii) between the volume of the residual subicular
complex and patients' performance on spatial perception/memory. The effects reported
here suggest that following an early-life exposure to hypoxia-ischemia, the function of
hippocampal subregions showing greater damage will more likely be compensated by
preserved extra-hippocampal structures. By contrast, hippocampal subregions with milder
injury could increase system disruption by maintaining their putative functions. Despite
these compensatory reorganizations, hippocampal-dependent episodic memory recall
remains irreversibly compromised.

Limitations
Although manual tracing is still the gold standard for measuring the volumes of
hippocampal subregions and cortical areas on MRI acquisitions, manual segmentation is
subjective and susceptible to measurement errors. Also, the precision of the measure is
limited by the resolution of the MRI scans. However, analyses of MR images for
estimating the extent of hippocampal damage in amnesic patients have been validated by
postmortem examinations (Rempel-Clower et al., 1996). In particular, a clear
concordance has been reported in the appearance of the damaged hippocampus on the
MRI and in the histological sections (Rempel-Clower et al., 1996). The border between
cytoarchitectonic hippocampal subregions were estimated based on delineation of these
regions on histological sections (Rosene and Van Hoesen, 1977). The delineation on MR
images of the perihippocampal and parahippocampal cortical areas were based on that
described in (Insausti et al., 1998). It is not clear to what extent these borders would be
affected in the brains of DA patients with significant hippocampal volume reduction. Our
observations of an absence of significant volume reduction of the entorhinal, perirhinal
and parahippocampal cortices in patients with DA is consistent with previous report
showing an absence of significant volume reduction in patients with adult-onset amnesia
(Shrager et al., 2008), and in a patient with DA (Jonin et al., 2018). Also, intact
parahippocampal activity during scene reinstatement has been described in patients with
DA (Elward et al., 2021).

Differential effect of early hypoxia-ischemia on hippocampal subregions
Our observation that the CA-DG were the most affected hippocampal subregions in
patients with DA is congruent with documented reports of greater sensitivity of this
subregion to hypoxia-ischemia. It has long been known that hypoxia and ischemia result
in selective neuronal death in the CA1 field (Schmidt-Kastner and Freund, 1991),

although the reason for this selectivity is still debated (Wang and Michaelis, 2010).
Neurohistological studies of patients who suffered ischemic episodes in adulthood have
reported cases with neural cell loss restricted to the CA1 region bilaterally whilst the
CA1' region of the uncus, the subiculum, CA3 and surrounding cortices were all
preserved (Rempel-Clower et al., 1996; Zola-Morgan et al., 1986). High-resolution
magnetic resonance imaging showed that CA1 was always affected in patients with
hippocampal damage (Bartsch et al., 2015).
The subicular complex was less affected than the CA-DG subregion in patients with
DA and this difference could be related to the different levels of maturation of the
hippocampal subregions at birth. Quantitative volumetric measurements in non-human
primates revealed that the subiculum, presubiculum and parasubiculum develop earlier
than the dentate gyrus, CA3, and CA1 (Jabès et al., 2011) and this progressive maturation
of the hippocampal circuits may participate in the differential protracted maturation of
relational memory processes (Jabès and Nelson, 2015). The level of maturation of the
hippocampal subregions at birth could also influence their vulnerability to hypoxia-
ischemia (Vannucci and Hagberg, 2004).
In contrast, the uncus showed a low sensitivity to hypoxia in our patient cohort which
could be a reflection of the distinct vascularization of this hippocampal subregion. The
branches of the posterior cerebral artery (PCA) coming from the vertebral artery irrigate
the posterior part of the hippocampus, while the branches of the anterior choroidal artery
(AChA) originating from the internal carotid artery irrigate the uncal portion of the
hippocampus (Erdem et al., 1993; Huther et al., 1998; Marinković et al., 1992). High-
resolution angiography and anatomical studies have highlighted that the anterior
hippocampal region could receive a mixed blood supply (from both the PCA and the
AChA) in about half of the individuals (Erdem et al., 1993; Perosa et al., 2020). A mixed
blood supply could provide more vascular reserve and lower vulnerability to neuronal
injury and atrophy (Liebeskind, 2003). The uncus, which receives a distinct blood supply
compared to the rest of the hippocampus, might thus exhibit greater resistance to the
adverse effects of hypoxic-ischemic events than the other hippocampal regions.

Uncus and recall memory
The residual uncus could be partly functional in patients with DA and could be recruited
for recall memory. We reported here that the volume of the uncus correlated negatively
with recall scores as well as working memory performance in patients with DA. In the
WAIS/WISC test, Working Memory Index is made up of two subtests (Digit Span and
Letter Number Sequencing) both requiring verbal recall in the span of working
memory. The function of the uncus is not well understood mainly because its structure-
function mapping remains to be determined in rodents (Ding, 2013) but in humans, the
uncus has been proposed to be involved in the long-term recall of scenes (Zeidman et al.,
2015). Connectivity studies in nonhuman primates have shown that the uncus is less
connected to other regions of the hippocampus, and that it has its own associational
system of connections (Kondo et al., 2009). Notably, the uncus is also the only
hippocampal region reciprocally connected to its contralateral counterpart (Amaral et al.,
1984). The uncus' unique connectivity could allow this region to support some
information processing despite severe damage to the other hippocampal areas.
Notwithstanding the importance of the uncus in relation to severity of atrophy
during delayed recall, this subregion maintains its role during encoding and learning of
memoranda in both low and high severity subgroups equally. Thus, we noted that despite
their poor performance overall, both patient subgroups incrementally increased their

scores across the three learning trials of the *People* and *Shapes* subtests of the *Doors and*
People Test. The absence of a difference in performance between the subgroups during
encoding and learning, but not during delayed recall, highlights the role of the uncus in
retrieval of auditory verbal memory after a delay in patients with DA.
Interestingly, we also reported an inverse relationship between the volume of the
uncus and patients' performance in delayed verbal, but not visual recall. Impairment in
delayed verbal recall in relation to extent of atrophy of the uncus highlights the
vulnerability of auditory verbal information to recall after a delay (Schulze et al., 2012).

Subiculum
Residual circuits of the subicular complex might participate in spatial perception and
spatial memory in patients with DA. Here, we reported inverse correlations between
spatial perception/memory performance and the volume of the residual subicular complex
in patients with DA. The subiculum receives its major excitatory input from CA1, but it
also receives a strong projection from the entorhinal cortex (Amaral and Lavenex, 2007;
Matsumoto et al., 2019; Witter and Amaral, 2020). The subiculum is also the main source
of subcortical projections directed to the hippocampus, and the presubiculum receives the
only direct input from the anterior thalamic nucleus (Amaral and Lavenex, 2007;
Matsumoto et al., 2019). The subiculum, presubiculum and parasubiculum send dense
projections to the entorhinal cortex (Saunders and Rosene, 1988; Witter and Amaral,
2020). Thus, in the absence of functional CA-DG, the residual subicular complex could
marginally participate in input processing through its other interconnections, and in
particular, through interaction with the structurally preserved entorhinal cortex which
constitutes the main entryway for much of the neocortical input reaching the
hippocampus (Amaral et al., 1987).
Animal and human studies suggest that the subiculum plays an important role in
spatial processing and spatial representation (Matsumoto et al., 2019). Grid, border, and
head direction cells have been found in the subiculum of rodents and primates including
humans (Boccara et al., 2010; Lee et al., 2018; Lever et al., 2009; Robertson et al., 1999).
In the rat, cross-structural interaction between the subiculum and the ventral striatum has
been reported to be critical for spatial memory consolidation (Torromino et al., 2019). In
humans, the subiculum is suggested to be specifically involved in scene-based cognitive
processing, including scene perception and scene construction during memory recall
(Dalton and Maguire, 2017; Hodgetts et al., 2017; Zeidman et al., 2015).
Interestingly, we did not observe correlations between the volume of the subicular
complex and patients' performance on the non-spatial tasks of the *Doors and People*
Test, suggesting that the subicular complex might be playing a more prominent role in the
processing of spatial inputs.
The Processing Speed Index of the Wechsler Scales is made up of two subtests
('Coding' which requires learning and memory and 'Symbol Search' which requires
visual scanning/cancellation, and visual stimulus matching, but without memory
demands). Interestingly, the Coding scores are negatively correlated with the uncus
volume while the Symbol Search scores are negatively correlated with both the uncus and
the subicular complex. Our results tend to demonstrate that the hippocampus is involved
in two cognitive processes of visual perception, and cognitive memory.

Working memory
We reported here, in addition to an absence of working memory impairment, a negative
association between uncus volume and working memory performance in patients with

DA. The more established view proposes that the hippocampus plays an important role in
episodic memory, spatial cognition, and navigation, while it is not critically contributing
to working memory (Eichenbaum, 2006; Eichenbaum et al., 1994). But an alternative
view proposes that the hippocampus may contribute to any memory task, including
working memory (Hannula and Ranganath, 2008; Libby et al., 2014), in particular in
supporting high-precision binding (Borders et al., 2021). This latter possibility is
challenged however by the fact that patients with DA presenting with severe hippocampal
damage have repeatedly shown unimpaired working memory performance (Allen et al.,
2014; Baddeley et al., 2011, 2010). Recent work extended these findings by showing that
high resolution working memory performance was unimpaired, and even often superior,
relative to control group in a patient with DA (Allen et al., 2022). Our observation of a
negative correlation between uncus volume and working memory performance suggests
that the perirhinal, entorhinal, and parahippocampal cortices normally involved in the
binding of information in the short term (Miyashita, 2019) may be subject to plastic
changes in patients with DA and that these plastic changes may depend on the degree of
hippocampal damage. The possibility that the hippocampal damage in patients with DA
may not influence working memory performance directly, but indirectly via
reorganization of the MTL circuit is explored below.

Functional hypotheses
Experimental studies in nonhuman primates have reported similar paradoxical findings to
those reported in the present study. Monkeys with small hippocampal lesions exhibit
greater memory loss on Delayed Nonmatching-to-Sample (DNMS) task than that found
in animals with greater damage (Bachevalier and Mishkin, 1989). Furthermore, a positive
correlation between the extent of damage to the hippocampus and scores on a DNMS task
has been reported in experimentally lesioned monkeys (Murray and Mishkin, 1998).
Several hypotheses have been proposed to explain why discrete lesions result in greater
memory impairment (Bachevalier and Mishkin, 1989; Baxter and Murray, 2001; Beason-
Held et al., 1999). However, it is difficult to relate these findings obtained following adult
experimental lesions in the monkey to observations in patients with early hippocampal
damage whose circuits might have undergone massive compensation. One hypothesis to
explain our observations is that the level of hippocampal damage would likely have a
strong direct influence on the organization of the medial temporal lobe memory circuits
in the adult. Following an early event of hypoxia-ischemia, however, greater
hippocampal damage might induce greater compensatory reconfigurations in the neural
circuits and enable other structures, in particular the surrounding cortical areas normally
recruited by the hippocampus, to assume important aspects of memory function. In
contrast, when the hippocampus is only partially damaged, the information flow would
remain present in the hippocampus and could result in incomplete, and possibly
disruptive information processing. Our observation could thus reveal the existence of
multiple, redundant routes within the residual hippocampal subregions and surrounding
cortical areas in patients with developmental amnesia (Aggleton et al., 2000). These
parallel circuits could compete for the control of behavior and disrupt perception and
memory performance (Baxter and Murray, 2001).
Our findings are consistent with the hypothesis that a close relationship can be
evidenced between the volume of remaining hippocampal subregions and the severity of
cognitive deficits in patients with hippocampal damage, in the case of patients with DA
this relationship was inverse. Our observations also provide further support for the model
of hierarchical organization of cognitive memory (Mishkin et al., 1997) by proposing that

surrounding cortical areas normally involved in the encoding of associative relations
between decontextualized information (Miyashita, 2019) could reorganize to contribute
to hippocampal function following early hippocampal damage. The limits of this
reorganization, however, may be determined by the extent of damage to the hippocampal
subregions.
Evidence for such reorganization has already been reported in a nonhuman
primate model of developmental amnesia. Monkeys with early bilateral hippocampal
lesion can learn new spatial relational information in striking contrast to monkeys with
adult lesion (Banta Lavenex et al., 2006; Lavenex et al., 2007). Interestingly, in these
monkeys with early hippocampal lesions, the surrounding cortical areas were preserved
and were shown to contribute to spatial learning in the absence of functional hippocampal
circuits (Chareyron et al., 2017).
One could speculate that the functional reorganization of the perihippocampal
structures (normally devoted to the processing of decontextualized information) to
compensate hippocampal function could, in turn, benefit the processing of semantic
memory and lead to better context-free memory performance in patients (see: Kapur et
al., 2013). Indications of increased memory performance have indeed been reported in
two different patients with DA. Jonin et al. (2018) studied patient 'KA' presenting with
55% hippocampal volume reduction and preserved perihippocampal structures. While
patient KA displayed few, if any, residual episodic abilities, this patient was able to
accurately retrieve semantic memories, and show evidence of superior or even very
superior access to these memories than controls. Another patient with DA, patient 'Jon',
has often produced response accuracy levels that are at least higher than those of control
participants in previous experimental explorations of visuospatial working memory
(Allen et al., 2022, 2014; Baddeley et al., 2010). Jon was at least numerically superior to
the control mean on recognition-based measures of shape-color binding (Baddeley et al.,
2010). In measures of color-location memory, Jon's recognition accuracy matched the
highest achieving control participant, while his reconstruction performance was superior
to 6 of the 7 controls (Allen et al., 2014). Across four tasks measuring binding between
color and orientation or color and location using simultaneous or sequential presentation
of stimuli, Jon's response accuracy was high, and always numerically superior to the
control mean (Allen et al., 2022).
These observations from single patients studies together with the present findings,
obtained in a large cohort of patients with DA, support the hypothesis of a reorganization
of the preserved perihippocampal areas following early hippocampal damage.

Conclusions

Our results have important implications for understanding the organization of memory
following early hippocampal injury. These observations highlight the paradoxical finding
that in patients with DA greater hippocampal damage can be associated with better
performance in recall memory, Verbal comprehension Index, Working memory Index,
Processing speed Index and spatial perception. In contrast, less severe damage leaving
behind relatively preserved hippocampal subregions can lead to more pronounced
cognitive deficits. These unexpected findings can disentangle the functional organization
of the medial temporal lobe memory system, and attribute a central role in the
compensation of memory and spatial cognition to extra-hippocampal structures in
patients with early hippocampal damage.

Materials and methods

Participants

The patients with DA were characterized by (i) a history of episodes of hypoxia-ischemia in early life, (ii) quantified hippocampal volume reduction above 25% relative to a group of controls, and (iii) severely impaired episodic memory (Dzieciol et al., 2017). The most common aetiology was perinatal hypoxic-ischemic encephalopathy (Table S2). While long-term neurodevelopmental outcome depends on the severity of the hypoxic-ischemic episode, subtle cognitive deficits and behavioral alterations have been revealed through long-term evaluations even in mild forms of hypoxia-ischemia (de Vries and Jongmans, 2010).

When all the volumetric estimates were completed, one of the patients, who presented with moderate memory deficits and only 11.0% hippocampal volume reduction associated with two unconfirmed episodes of prolonged seizures at age 4, was excluded from the DA group.

Also, we excluded from the DA group a patient with temporal lobe epilepsy presenting with a 25.8% hippocampal volume reduction, but without a documented episode of early hypoxia-ischemia and/or status epilepticus. Additionally, blinded neuroradiological examination carried out independently by one of the authors (WKC) revealed that this patient's MRI scan was entirely normal. The hippocampus, fornix and mammillary bodies were noted to have a normal appearance in contrast to all other patients with DA.

We report here hippocampal volume estimates in 32 control participants (16 females; mean age: 17.7 years, SD 7.7, range 8−38) and 23 patients with DA (11 females; mean age: 19.5 years, SD 9.0, range 8−40).

The study was approved by the London-Bentham Research Ethics Committee (REC Reference No. 05/Q0502/88) and all participants and their carers read an age-appropriate information sheet and gave written informed consent according to the Declaration of Helsinki, before commencing the study.

Structural MRI

Participants were either scanned with a 1.5T-MRI scanner (24 control participants, 16 patients with DA) or with a 3T-MRI scanner (8 control participants, 7 patients with DA).

Whole brain structural 1.5T-MRI scans were obtained using a Siemens Avanto Scanner, with a T1-weighted 3D FLASH sequence with the following parameters: repetition time 11ms; echo time 4.94ms; in-plane resolution 1mm×1mm; slice thickness of 1mm (Cooper et al., 2011).

Whole brain structural 3T-MRI scans were obtained using a 3T Siemens MRI system with a 20 channel head coil. A T1-weighted magnetization-prepared rapid gradient-echo (MPRAGE) scan was acquired with the following parameters: in-plane resolution of 1mm×1mm; slice thickness of 1mm; repetition time of 2,300ms; echo time of 2.74ms (Buck et al., 2019).

For the measurement of regions of interest (ROI) volumes, the data were reformatted into 1 mm-thick contiguous slices in the coronal plane. ROI cross-sectional areas were measured in all slices along the entire length of the hippocampus, entorhinal, perirhinal and parahippocampal cortices using ITK-SNAP (version 3.8.0; www.itksnap.org; Yushkevich et al., 2006). Manual tracing is still the gold standard for measuring hippocampal volume and is particularly appropriate for samples of relatively small size (Akudjedu et al., 2018), presenting with various levels of hippocampal atrophy

568   (Schmidt et al., 2018), or including children and adolescents (Herten et al., 2019). All
569   manual measurements of the ROIs were carried out by one of the authors (LJC) who
570   remained blind to participant identity.

571   The hippocampus comprises the CA fields (CA3, CA2, CA1), the dentate gyrus
(DG), subiculum, presubiculum, and parasubiculum (Amaral and Lavenex, 2007). The
human uncus [also termed the "uncal portion of the hippocampal formation" (Amaral et
al., 1984)] is situated antero-medially and consists of slightly modified DG, CA3, CA2,
CA1, and subiculum (Ding and Van Hoesen, 2015). For the segmentation of the
hippocampus and its subregions, we followed the guide provided by Dalton et al. (Dalton
et al., 2017). Given the difficulty to distinguish hippocampal CA fields on 1.5T and 3T-
MRI acquisitions, we restricted our volumetric analyses to three ROIs: uncus [defined
according to (Ding and Van Hoesen, 2015)], CA-DG (CA fields and dentate gyrus), and
subicular complex (subiculum, presubiculum, and parasubiculum). The segmentation of
the entorhinal, perirhinal and parahippocampal cortices was based on structural
descriptions of human medial temporal lobe and segmentations protocols (Blaizot et al.,
2010; Ding and Van Hoesen, 2010; Frankó et al., 2014; Insausti et al., 1998; Kivisaari et
al., 2020).

585   The volumes in cubic millimeters were calculated by summing the number of
voxels (1×1×1 mm) for each ROI. The volume of the hippocampus as a whole was
calculated as the sum of hippocampal subregions' volumes (uncus, CA-DG, subicular
complex). We then used parametric tests as volumetric estimates of the hippocampal
subregions and cortical areas were normally distributed (Shapiro-Wilk normality test;
control: all $P > .06$; DA: all $P > .09$). There was no significant difference between
volumetric estimates of the whole hippocampus or cortical areas obtained with the 1.5T-
MRI scanner and the 3T-MRI scanner in control and DA groups (two-way ANOVA, both
$F < 1.3$; both $P > .25$). Also there was no significant gender difference for the volumes of
the whole hippocampus or cortical areas in control and DA groups (both $F < 0.08$; both P
$> .79$). Volume estimates on 1.5T and 3T-MRI scans and volume estimates in males and
females were thus presented together. There was a significant right larger than left
volume asymmetry for the volume of the whole hippocampus in both control and DA
groups (repeated measures two-way ANOVA: $F_{(1,53)} = 19.2$; $P < .001$; Tukey's HSD post-
hoc test: both $P < .004$). The right CA-DG subregions were significantly larger than the
left in both control and DA groups ($F_{(1,53)} = 26.1$; $P < .001$; both $P < .004$). There was no
significant asymmetry for the volume of the other hippocampal subregions (all $P > .05$).
Also, there was no significant asymmetry for the volume of the entorhinal or perirhinal
cortices (all $P > .12$) but there was a significant right larger than left volume asymmetry
for the volume of the parahippocampal cortex in the control group ($F_{(1,53)} = 7.0$; $P = .011$;
Tukey's HSD post-hoc test: $P = .009$). The volume of all the ROIs was calculated as the
mean of the left and right hemisphere measurements.

Intracranial volume (ICV) was manually measured on the sagittal dataset with a 1-in-
10 random and systematic sampling strategy. The ICV was normally distributed
(Shapiro-Wilk normality test; control: $P = .05$; DA: $P = .23$). The ICV was 6.8% smaller
in the DA group than in the control group ($t_{(53)} = −2.70$; $P = .009$). Corrections, derived
from the regression line of control ROI volume (V) versus ICV, were then made for ICV
based on the formula [$\Delta V = a \times (\Delta ICV)$] where 'a' is the regression coefficient.

WISC/WAIS
The Wechsler Adult Intelligence Scale, 3rd Ed. (WAIS-III) and the Wechsler Intelligence
Scale for Children, 4th Ed. (WISC-IV) provide four index scores: Verbal Comprehension
Index, measuring understanding and expression of verbal concepts; Perceptual Reasoning

Index, reflecting visuospatial perception and perceptuomotor manipulation; Working
Memory Index, tracking on-line immediate memory and executive control; and
Processing Speed Index, measuring speed of visuoperceptual discrimination, and copying
of number-symbol associations. The four index scores are expressed as standard scores
with a mean of 100 and a standard deviation of 15.
Scoring. The WISC-IV or WAIS-III were administered to all patients with DA (N = 23,
11 females, mean age: 19.5 years, SD: 9.0, range: 8−40) and to 23 controls (13 females,
mean age: 18.0 years, SD: 8.4, range: 8.2–38). For the analysis of the data, we then used
parametric tests as the data were normally distributed in all four indexes (Shapiro–Wilk
test of normality; Control: all P > .36; DA: all P > .10).
627

The Doors and People Test
The *Doors and People Test* was administered according to the instructions in the
published manual (Baddeley et al., 1994), as described in detail in (Adlam et al., 2009;
Patai et al., 2015). Briefly, for the verbal recall subtest ('People'), participants were
presented with four pictures of people and after three learning trials asked to recall their
names cued by their profession. For the visual recall subtest ('Shapes'), participants first
copied, and drew the shapes from memory after three learning trials. For verbal
recognition ('Names'), participants were presented with a list of names, and then asked to
recognize each one from a list of four alternatives. Finally, in the visual recognition
subtest ('Doors'), participants viewed a list of doors and then identified each one from a
list of four alternatives. See Supplementary Information for more details.
Scoring. The *Doors and People Test* was administered to all patients with DA (N = 23,
11 females, mean age: 19.5 years, SD: 9.0, range: 8−40) and to 21 controls (11 females,
mean age: 17.6 years, SD: 8.6, range: 8−38). One of the patients performed all but the
'Names' subtest ('People', 'Doors', 'Shapes': N = 23; 'Names': n = 22). Given that the
DA group included individuals under the age of 16, which is the first age band at which
standard scores are available on the *Doors and People Test*, we used standard scores
based on the scores of our control group; raw scores (SD): People, 28.2 (6.9); Doors, 18.9
(3.8); Shapes, 34.6 (2.7); Names, 18.6 (3.7). We converted patients' and controls' raw
scores to z-scores relative to the control group's scores. For the analysis of the data, we
then used nonparametric tests as the data were not normally distributed in all four
subtests (Shapiro–Wilk test of normality; Control: P < .001; DA: P = .044).

The Four Mountains Test
The *Four Mountains Test* was developed to examine hippocampal contribution to
perception and short-term memory for topographical and nonspatial information in spatial
scenes (Hartley et al., 2007). Briefly, this test involves computer-generated landscapes
containing four mountains. In the 'Spatial Perception' task, participants selected one
picture that matched the sample image (the same place from a different viewpoint) from
four alternatives. In the 'Spatial Memory' task, the sample image was presented for
approximately 8 sec. After a 2 sec. delay, participants were presented with four choices
and selected the one previously viewed. See Supplementary Information for more details.
Scoring. The *Four Mountains Test* was administered to 14 patients with DA (7 females,
mean age: 21.6 years, SD: 10.3, range: 8−40) and to 19 controls (11 females, mean age:
16.9 years, SD: 7.8, range: 8−35). We converted patients' raw scores to z-scores relative
to the scores of our control group; raw scores (SD): Perception, 11.6 (2.3); Memory, 9.95
(2.8). We then used nonparametric tests as the data in the Control group were not
normally distributed [Shapiro–Wilk test of normality; P < .05].

Acknowledgments

This work was supported by the UK Medical Research Council (program grant numbers G03000117 and G1002276).

We thank the participants and their families for their support of our research.

Data availability

Anonymized participant data will be available upon reasonable request.

Competing interests

The authors declare no conflict of interest.

References

Adlam, A.-L.R., Malloy, M., Mishkin, M., Vargha-Khadem, F., 2009. Dissociation between recognition and recall in developmental amnesia. Neuropsychologia 47, 2207–2210. https://doi.org/10.1016/j.neuropsychologia.2009.01.038

Aggleton, J.P., Vann, S.D., Oswald, C.J., Good, M., 2000. Identifying cortical inputs to the rat hippocampus that subserve allocentric spatial processes: a simple problem with a complex answer. Hippocampus 10, 466–474. https://doi.org/10.1002/1098-1063(2000)10:4<466::AID-HIPO13>3.0.CO;2-Y

Akudjedu, T.N., Nabulsi, L., Makelyte, M., Scanlon, C., Hehir, S., Casey, H., Ambati, S., Kenney, J., O'Donoghue, S., McDermott, E., Kilmartin, L., Dockery, P., McDonald, C., Hallahan, B., Cannon, D.M., 2018. A comparative study of segmentation techniques for the quantification of brain subcortical volume. Brain Imaging Behav 12, 1678–1695. https://doi.org/10.1007/s11682-018-9835-y

Allen, R.J., Atkinson, A.L., Vargha-Khadem, F., Baddeley, A.D., 2022. Intact high-resolution working memory binding in a patient with developmental amnesia and selective hippocampal damage. Hippocampus 32, 597–609. https://doi.org/10.1002/hipo.23452

Allen, R.J., Vargha-Khadem, F., Baddeley, A.D., 2014. Item-location binding in working memory: is it hippocampus-dependent? Neuropsychologia 59, 74–84. https://doi.org/10.1016/j.neuropsychologia.2014.04.013

Amaral, D.G., Insausti, R., Cowan, W.M., 1987. The entorhinal cortex of the monkey: I. Cytoarchitectonic organization. J Comp Neurol 264, 326–55.

Amaral, D.G., Insausti, R., Cowan, W.M., 1984. The commissural connections of the monkey hippocampal formation. J. Comp. Neurol. 224, 307–336. https://doi.org/10.1002/cne.902240302

Amaral, D.G., Lavenex, P., 2007. Hippocampal neuroanatomy, in: The Hippocampus Book. Oxford University Press, Oxford, pp. 37–114.

Argyropoulos, G.P.D., Dell'Acqua, C., Butler, E., Loane, C., Roca-Fernandez, A.,
Almozel, A., Drummond, N., Lage-Martinez, C., Cooper, E., Henson, R.N., Butler, C.R.,
2021. Functional Specialization of the Medial Temporal Lobes in Human Recognition
Memory: Dissociating Effects of Hippocampal versus Parahippocampal Damage. Cereb
Cortex bhab290. https://doi.org/10.1093/cercor/bhab290

Bachevalier, J., Mishkin, M., 1989. Mnemonic and neuropathological effects of
occluding the posterior cerebral artery in Macaca mulatta. Neuropsychologia 27, 83–105.
https://doi.org/10.1016/0028-3932(89)90092-4

Baddeley, A., Allen, R., Vargha-Khadem, F., 2010. Is the hippocampus necessary for
visual and verbal binding in working memory? Neuropsychologia 48, 1089–1095.
https://doi.org/10.1016/j.neuropsychologia.2009.12.009

Baddeley, A., Jarrold, C., Vargha-Khadem, F., 2011. Working memory and the
hippocampus. J Cogn Neurosci 23, 3855–3861. https://doi.org/10.1162/jocn_a_00066

Baddeley, A.D., 2020. On trying to prove Endel Tulving wrong: A revised modal model
of amnesia. Neuropsychologia 148, 107590.
https://doi.org/10.1016/j.neuropsychologia.2020.107590

Baddeley, A.D., Emslie, H., Nimmo-Smith, I., 1994. Doors and people : a test of visual
and verbal recall and recognition. Bury St Edmunds : Thames Valley Test Company, UK.

Banta Lavenex, P., Amaral, D.G., Lavenex, P., 2006. Hippocampal lesion prevents
spatial relational learning in adult macaque monkeys. J. Neurosci. 26, 4546–4558.
https://doi.org/10.1523/JNEUROSCI.5412-05.2006

Bartsch, T., Döhring, J., Reuter, S., Finke, C., Rohr, A., Brauer, H., Deuschl, G., Jansen,
O., 2015. Selective neuronal vulnerability of human hippocampal CA1 neurons: lesion
evolution, temporal course, and pattern of hippocampal damage in diffusion-weighted
MR imaging. J. Cereb. Blood Flow Metab. 35, 1836–1845.
https://doi.org/10.1038/jcbfm.2015.137

Baxter, M.G., Murray, E.A., 2001. Opposite relationship of hippocampal and rhinal
cortex damage to delayed nonmatching-to-sample deficits in monkeys. Hippocampus 11,
61–71. https://doi.org/10.1002/1098-1063(2001)11:1<61::AID-HIPO1021>3.0.CO;2-Z

Bayley, P.J., Squire, L.R., 2005. Failure to acquire new semantic knowledge in patients
with large medial temporal lobe lesions. Hippocampus 15, 273–280.
https://doi.org/10.1002/hipo.20057

Beason-Held, L.L., Rosene, D.L., Killiany, R.J., Moss, M.B., 1999. Hippocampal
formation lesions produce memory impairment in the rhesus monkey. Hippocampus 9,
562–574. https://doi.org/10.1002/(SICI)1098-1063(1999)9:5<562::AID-
HIPO10>3.0.CO;2-X

Blaizot, X., Mansilla, F., Insausti, A.M., Constans, J.M., Salinas-Alamán, A., Pró-
Sistiaga, P., Mohedano-Moriano, A., Insausti, R., 2010. The human parahippocampal
region: I. Temporal pole cytoarchitectonic and MRI correlation. Cereb Cortex 20, 2198–
2212. https://doi.org/10.1093/cercor/bhp289

Boccara, C.N., Sargolini, F., Thoresen, V.H., Solstad, T., Witter, M.P., Moser, E.I.,
Moser, M.-B., 2010. Grid cells in pre- and parasubiculum. Nat. Neurosci. 13, 987–994.
https://doi.org/10.1038/nn.2602

Borders, A.A., Ranganath, C., Yonelinas, A.P., 2021. The hippocampus supports high-

precision binding in visual working memory. Hippocampus.
https://doi.org/10.1002/hipo.23401

Buck, S., Bastos, F., Baldeweg, T., Vargha-Khadem, F., 2019. A Functional MRI
Paradigm Suitable for Language and Memory Mapping in Pediatric Temporal Lobe
Epilepsy. Front Neurol 10, 1384. https://doi.org/10.3389/fneur.2019.01384

Chareyron, L.J., Banta Lavenex, P., Amaral, D.G., Lavenex, P., 2017. Functional
organization of the medial temporal lobe memory system following neonatal
hippocampal lesion in rhesus monkeys. Brain Struct Funct 222, 3899–3914.
https://doi.org/10.1007/s00429-017-1441-z

Cooper, J.M., Gadian, D.G., Jentschke, S., Goldman, A., Munoz, M., Pitts, G., Banks, T.,
Chong, W.K., Hoskote, A., Deanfield, J., Baldeweg, T., de Haan, M., Mishkin, M.,
Vargha-Khadem, F., 2015. Neonatal hypoxia, hippocampal atrophy, and memory
impairment: evidence of a causal sequence. Cereb Cortex 25, 1469–1476.
https://doi.org/10.1093/cercor/bht332

Cooper, J.M., Vargha-Khadem, F., Gadian, D.G., Maguire, E.A., 2011. The effect of
hippocampal damage in children on recalling the past and imagining new experiences.
Neuropsychologia 49, 1843–1850.
https://doi.org/10.1016/j.neuropsychologia.2011.03.008

Dalton, M.A., Maguire, E.A., 2017. The pre/parasubiculum: a hippocampal hub for
scene-based cognition? Curr Opin Behav Sci 17, 34–40.
https://doi.org/10.1016/j.cobeha.2017.06.001

Dalton, M.A., Zeidman, P., Barry, D.N., Williams, E., Maguire, E.A., 2017. Segmenting
subregions of the human hippocampus on structural magnetic resonance image scans: An
illustrated tutorial. Brain Neurosci Adv 1, 2398212817701448.
https://doi.org/10.1177/2398212817701448

de Vries, L.S., Jongmans, M.J., 2010. Long-term outcome after neonatal hypoxic-
ischaemic encephalopathy. Arch. Dis. Child. Fetal Neonatal Ed. 95, F220-224.
https://doi.org/10.1136/adc.2008.148205

Ding, S.-L., 2013. Comparative anatomy of the prosubiculum, subiculum, presubiculum,
postsubiculum, and parasubiculum in human, monkey, and rodent. J. Comp. Neurol. 521,
4145–4162. https://doi.org/10.1002/cne.23416

Ding, S.-L., Van Hoesen, G.W., 2015. Organization and Detailed Parcellation of Human
Hippocampal Head and Body Regions Based on a Combined Analysis of Cyto- and
Chemoarchitecture. J. Comp. Neurol. 523, 2233–2253. https://doi.org/10.1002/cne.23786

Ding, S.-L., Van Hoesen, G.W., 2010. Borders, extent, and topography of human
perirhinal cortex as revealed using multiple modern neuroanatomical and pathological
markers. Hum Brain Mapp 31, 1359–1379. https://doi.org/10.1002/hbm.20940

Dzieciol, A.M., Bachevalier, J., Saleem, K.S., Gadian, D.G., Saunders, R., Chong,
W.K.K., Banks, T., Mishkin, M., Vargha-Khadem, F., 2017. Hippocampal and
diencephalic pathology in developmental amnesia. Cortex 86, 33–44.
https://doi.org/10.1016/j.cortex.2016.09.016

Eichenbaum, H., 2006. Remembering: functional organization of the declarative memory
system. Curr Biol 16, R643-645. https://doi.org/10.1016/j.cub.2006.07.026

Eichenbaum, H., Otto, T., Cohen, N.J., 1994. Two functional components of the

hippocampal memory system. Behavioral and Brain Sciences 17, 449–517.
https://doi.org/10.1017/S0140525X00035391

Elward, R.L., Rugg, M.D., Vargha-Khadem, F., 2021. When the brain, but not the
person, remembers: Cortical reinstatement is modulated by retrieval goal in
developmental amnesia. Neuropsychologia 154, 107788.
https://doi.org/10.1016/j.neuropsychologia.2021.107788

Elward, R.L., Vargha-Khadem, F., 2018. Semantic memory in developmental amnesia.
Neurosci. Lett. 680, 23–30. https://doi.org/10.1016/j.neulet.2018.04.040

Erdem, A., Yaşargil, G., Roth, P., 1993. Microsurgical anatomy of the hippocampal
arteries. J. Neurosurg. 79, 256–265. https://doi.org/10.3171/jns.1993.79.2.0256

Frankó, E., Insausti, A.M., Artacho-Pérula, E., Insausti, R., Chavoix, C., 2014.
Identification of the human medial temporal lobe regions on magnetic resonance images.
Hum Brain Mapp 35, 248–256. https://doi.org/10.1002/hbm.22170

Geva, S., Jentschke, S., Argyropoulos, G.P.D., Chong, W.K., Gadian, D.G., Vargha-
Khadem, F., 2020. Volume reduction of caudate nucleus is associated with movement
coordination deficits in patients with hippocampal atrophy due to perinatal hypoxia-
ischaemia. Neuroimage Clin 28, 102429. https://doi.org/10.1016/j.nicl.2020.102429

Guderian, S., Dzieciol, A.M., Gadian, D.G., Jentschke, S., Doeller, C.F., Burgess, N.,
Mishkin, M., Vargha-Khadem, F., 2015. Hippocampal Volume Reduction in Humans
Predicts Impaired Allocentric Spatial Memory in Virtual-Reality Navigation. J. Neurosci.
35, 14123–14131. https://doi.org/10.1523/JNEUROSCI.0801-15.2015

Hannula, D.E., Ranganath, C., 2008. Medial temporal lobe activity predicts successful
relational memory binding. J Neurosci 28, 116–124.
https://doi.org/10.1523/JNEUROSCI.3086-07.2008

Hartley, T., Bird, C.M., Chan, D., Cipolotti, L., Husain, M., Vargha-Khadem, F.,
Burgess, N., 2007. The hippocampus is required for short-term topographical memory in
humans. Hippocampus 17, 34–48. https://doi.org/10.1002/hipo.20240

Herten, A., Konrad, K., Krinzinger, H., Seitz, J., von Polier, G.G., 2019. Accuracy and
bias of automatic hippocampal segmentation in children and adolescents. Brain Struct
Funct 224, 795–810. https://doi.org/10.1007/s00429-018-1802-2

Hodgetts, C.J., Voets, N.L., Thomas, A.G., Clare, S., Lawrence, A.D., Graham, K.S.,
2017. Ultra-High-Field fMRI Reveals a Role for the Subiculum in Scene Perceptual
Discrimination. J. Neurosci. 37, 3150–3159. https://doi.org/10.1523/JNEUROSCI.3225-
16.2017

Huther, G., Dörfl, J., Van der Loos, H., Jeanmonod, D., 1998. Microanatomic and
vascular aspects of the temporomesial region. Neurosurgery 43, 1118–1136.
https://doi.org/10.1097/00006123-199811000-00065

Insausti, R., Juottonen, K., Soininen, H., Insausti, A.M., Partanen, K., Vainio, P., Laakso,
M.P., Pitkänen, A., 1998. MR volumetric analysis of the human entorhinal, perirhinal,
and temporopolar cortices. AJNR Am J Neuroradiol 19, 659–671.

Jabès, A., Lavenex, P.B., Amaral, D.G., Lavenex, P., 2011. Postnatal development of the
hippocampal formation: a stereological study in macaque monkeys. J. Comp. Neurol.
519, 1051–1070. https://doi.org/10.1002/cne.22549

Jabès, A., Nelson, C.A., 2015. 20 Years after "The ontogeny of human memory: A
cognitive neuroscience perspective," where are we? International Journal of Behavioral
Development 39, 293–303. https://doi.org/10.1177/0165025415575766

Jonin, P.-Y., Besson, G., La Joie, R., Pariente, J., Belliard, S., Barillot, C., Barbeau, E.J.,
2018. Superior explicit memory despite severe developmental amnesia: In-depth case
study and neural correlates. Hippocampus 28, 867–885.
https://doi.org/10.1002/hipo.23010

Kapur, N., Cole, J., Manly, T., Viskontas, I., Ninteman, A., Hasher, L., Pascual-Leone,
A., 2013. Positive clinical neuroscience: explorations in positive neurology.
Neuroscientist 19, 354–369. https://doi.org/10.1177/1073858412470976

Kivisaari, S.L., Probst, A., Taylor, K.I., 2020. The Perirhinal, Entorhinal, and
Parahippocampal Cortices and Hippocampus: An Overview of Functional Anatomy and
Protocol for Their Segmentation in MR Images, in: Ulmer, S., Jansen, O. (Eds.), FMRI:
Basics and Clinical Applications. Springer International Publishing, Cham, pp. 355–383.
https://doi.org/10.1007/978-3-030-41874-8_24

Kondo, H., Lavenex, P., Amaral, D.G., 2009. Intrinsic connections of the macaque
monkey hippocampal formation: II. CA3 connections. J. Comp. Neurol. 515, 349–377.
https://doi.org/10.1002/cne.22056

Lavenex, P., Lavenex, P.B., Amaral, D.G., 2007. Spatial relational learning persists
following neonatal hippocampal lesions in macaque monkeys. Nat. Neurosci. 10, 234–
239. https://doi.org/10.1038/nn1820

Lee, S.A., Miller, J.F., Watrous, A.J., Sperling, M.R., Sharan, A., Worrell, G.A., Berry,
B.M., Aronson, J.P., Davis, K.A., Gross, R.E., Lega, B., Sheth, S., Das, S.R., Stein, J.M.,
Gorniak, R., Rizzuto, D.S., Jacobs, J., 2018. Electrophysiological Signatures of Spatial
Boundaries in the Human Subiculum. J. Neurosci. 38, 3265–3272.
https://doi.org/10.1523/JNEUROSCI.3216-17.2018

Lever, C., Burton, S., Jeewajee, A., O'Keefe, J., Burgess, N., 2009. Boundary vector cells
in the subiculum of the hippocampal formation. J. Neurosci. 29, 9771–9777.
https://doi.org/10.1523/JNEUROSCI.1319-09.2009

Libby, L.A., Hannula, D.E., Ranganath, C., 2014. Medial temporal lobe coding of item
and spatial information during relational binding in working memory. J Neurosci 34,
14233–14242. https://doi.org/10.1523/JNEUROSCI.0655-14.2014

Liebeskind, D.S., 2003. Collateral circulation. Stroke 34, 2279–2284.
https://doi.org/10.1161/01.STR.0000086465.41263.06

Maguire, E.A., Vargha-Khadem, F., Mishkin, M., 2001. The effects of bilateral
hippocampal damage on fMRI regional activations and interactions during memory
retrieval. Brain 124, 1156–1170. https://doi.org/10.1093/brain/124.6.1156

Manns, J.R., Hopkins, R.O., Squire, L.R., 2003. Semantic Memory and the Human
Hippocampus. Neuron 38, 127–133. https://doi.org/10.1016/S0896-6273(03)00146-6

Marinković, S., Milisavljević, M., Puskas, L., 1992. Microvascular anatomy of the
hippocampal formation. Surg Neurol 37, 339–349. https://doi.org/10.1016/0090-
3019(92)90001-4

Matsumoto, N., Kitanishi, T., Mizuseki, K., 2019. The subiculum: Unique hippocampal
hub and more. Neurosci. Res. 143, 1–12. https://doi.org/10.1016/j.neures.2018.08.002

Mishkin, M., Suzuki, W.A., Gadian, D.G., Vargha-Khadem, F., 1997. Hierarchical
Organization of Cognitive Memory. Philosophical Transactions: Biological Sciences 352,
1461–1467.

Miyashita, Y., 2019. Perirhinal circuits for memory processing. Nature Reviews
Neuroscience 20, 577–592. https://doi.org/10.1038/s41583-019-0213-6

Murray, E.A., Mishkin, M., 1998. Object recognition and location memory in monkeys
with excitotoxic lesions of the amygdala and hippocampus. J. Neurosci. 18, 6568–6582.

Patai, E.Z., Gadian, D.G., Cooper, J.M., Dzieciol, A.M., Mishkin, M., Vargha-Khadem,
F., 2015. Extent of hippocampal atrophy predicts degree of deficit in recall. Proc. Natl.
Acad. Sci. U.S.A. 112, 12830–12833. https://doi.org/10.1073/pnas.1511904112

Perosa, V., Priester, A., Ziegler, G., Cardenas-Blanco, A., Dobisch, L., Spallazzi, M.,
Assmann, A., Maass, A., Speck, O., Oltmer, J., Heinze, H.-J., Schreiber, S., Düzel, E.,
2020. Hippocampal vascular reserve associated with cognitive performance and
hippocampal volume. Brain 143, 622–634. https://doi.org/10.1093/brain/awz383

Picard, L., Mayor-Dubois, C., Maeder, P., Kalenzaga, S., Abram, M., Duval, C.,
Eustache, F., Roulet-Perez, E., Piolino, P., 2013. Functional independence within the
self-memory system: new insights from two cases of developmental amnesia. Cortex 49,
1463–1481. https://doi.org/10.1016/j.cortex.2012.10.003

Rabin, J.S., Olsen, R.K., Gilboa, A., Buchsbaum, B.R., Rosenbaum, R.S., 2016. Using
fMRI to understand event construction in developmental amnesia. Neuropsychologia 90,
261–273. https://doi.org/10.1016/j.neuropsychologia.2016.07.036

Rempel-Clower, N.L., Zola, S.M., Squire, L.R., Amaral, D.G., 1996. Three cases of
enduring memory impairment after bilateral damage limited to the hippocampal
formation. J. Neurosci. 16, 5233–5255.

Robertson, R.G., Rolls, E.T., Georges-François, P., Panzeri, S., 1999. Head direction
cells in the primate pre-subiculum. Hippocampus 9, 206–219.
https://doi.org/10.1002/(SICI)1098-1063(1999)9:3<206::AID-HIPO2>3.0.CO;2-H

Rosenbaum, R.S., Cassidy, B.N., Herdman, K.A., 2015. Patterns of preserved and
impaired spatial memory in a case of developmental amnesia. Front Hum Neurosci 9,
196. https://doi.org/10.3389/fnhum.2015.00196

Rosene, D.L., Van Hoesen, G.W., 1977. Hippocampal efferents reach widespread areas
of cerebral cortex and amygdala in the rhesus monkey. Science 198, 315–317.

Saunders, R.C., Rosene, D.L., 1988. A comparison of the efferents of the amygdala and
the hippocampal formation in the rhesus monkey: I. Convergence in the entorhinal,
prorhinal, and perirhinal cortices. Journal of Comparative Neurology 271, 153–184.
https://doi.org/10.1002/cne.902710202

Schmidt, M.F., Storrs, J.M., Freeman, K.B., Jack, C.R., Turner, S.T., Griswold, M.E.,
Mosley, T.H., 2018. A comparison of manual tracing and FreeSurfer for estimating
hippocampal volume over the adult lifespan. Hum Brain Mapp 39, 2500–2513.
https://doi.org/10.1002/hbm.24017

Schmidt-Kastner, R., Freund, T.F., 1991. Selective vulnerability of the hippocampus in
brain ischemia. Neuroscience 40, 599–636. https://doi.org/10.1016/0306-4522(91)90001-
5

Schulze, K., Vargha-Khadem, F., Mishkin, M., 2012. Test of a motor theory of long-term
auditory memory. Proc. Natl. Acad. Sci. U.S.A. 109, 7121–7125.
https://doi.org/10.1073/pnas.1204717109

Scoville, W.B., Milner, B., 1957. Loss of recent memory after bilateral hippocampal
lesions. J. Neurol. Neurosurg. Psychiatr. 20, 11–21.

Shrager, Y., Levy, D.A., Hopkins, R.O., Squire, L.R., 2008. Working memory and the
organization of brain systems. J Neurosci 28, 4818–4822.
https://doi.org/10.1523/JNEUROSCI.0710-08.2008

Torromino, G., Autore, L., Khalil, V., Mastrorilli, V., Griguoli, M., Pignataro, A.,
Centofante, E., Biasini, G.M., De Turris, V., Ammassari-Teule, M., Rinaldi, A., Mele,
A., 2019. Offline ventral subiculum-ventral striatum serial communication is required for
spatial memory consolidation. Nat Commun 10, 5721. https://doi.org/10.1038/s41467-
019-13703-3

Vargha-Khadem, F., Cacucci, F., 2021. A brief history of developmental amnesia.
Neuropsychologia 150, 107689. https://doi.org/10.1016/j.neuropsychologia.2020.107689

Vargha-Khadem, F., Gadian, D.G., Watkins, K.E., Connelly, A., Van Paesschen, W.,
Mishkin, M., 1997. Differential effects of early hippocampal pathology on episodic and
semantic memory. Science 277, 376–380. https://doi.org/10.1126/science.277.5324.376

Vargha-Khadem, F., Salmond, C.H., Watkins, K.E., Friston, K.J., Gadian, D.G., Mishkin,
M., 2003. Developmental amnesia: effect of age at injury. Proc Natl Acad Sci U S A 100,
10055–10060. https://doi.org/10.1073/pnas.1233756100

Verfaellie, M., Koseff, P., Alexander, M.P., 2000. Acquisition of novel semantic
information in amnesia: effects of lesion location. Neuropsychologia 38, 484–492.
https://doi.org/10.1016/S0028-3932(99)00089-5

Wang, X., Michaelis, E.K., 2010. Selective neuronal vulnerability to oxidative stress in
the brain. Front Aging Neurosci 2, 12. https://doi.org/10.3389/fnagi.2010.00012

Witter, M.P., Amaral, D.G., 2020. The entorhinal cortex of the monkey: VI. Organization
of projections from the hippocampus, subiculum, presubiculum, and parasubiculum.
Journal of Comparative Neurology doi: 10.1002/cne.24983.
https://doi.org/10.1002/cne.24983

Yushkevich, P.A., Piven, J., Hazlett, H.C., Smith, R.G., Ho, S., Gee, J.C., Gerig, G.,
2006. User-guided 3D active contour segmentation of anatomical structures: significantly
improved efficiency and reliability. Neuroimage 31, 1116–1128.
https://doi.org/10.1016/j.neuroimage.2006.01.015

Zeidman, P., Lutti, A., Maguire, E.A., 2015. Investigating the functions of subregions
within anterior hippocampus. Cortex 73, 240–256.
https://doi.org/10.1016/j.cortex.2015.09.002

Zeidman, P., Maguire, E.A., 2016. Anterior hippocampus: the anatomy of perception,
imagination and episodic memory. Nat. Rev. Neurosci. 17, 173–182.
https://doi.org/10.1038/nrn.2015.24

Zola-Morgan, S., Squire, L.R., Amaral, D.G., 1986. Human amnesia and the medial
temporal region: enduring memory impairment following a bilateral lesion limited to
field CA1 of the hippocampus. J. Neurosci. 6, 2950–2967.