Bacteriophage therapy for the treatment of *Mycobacterium tuberculosis* infections

in humanized mice

Fan Yang[1]*, Alireza Labani-Motlagh[1,5]*, Josimar Dornelas Moreira[1], Danish Ansari[1], Jose

Alejandro Bohorquez[1], Sahil Patel[1], Fabrizio Spagnolo[2], Jon Florence[1], Abhinav

Vankayalapati[1], Ramakrishna Vankayalapati[1], John J. Dennehy[3,4,#], Buka Samten[1#], and

Guohua Yi[1#]

[1]Center for Biomedical Research, The University of Texas Health Science Center at Tyler

[2]Life Sciences Department, Long Island University Post, Brookville, NY

[3]Biology Department, Queens College of The City University of New York, Flushing, NY

[4]The Graduate Center of The City University of New York, New York, NY

[5]Current address: Center for Discovery and Innovation, Hackensack Meridian Health,

New Jersey

*Contribute equally

#Corresponding authors. Guohua Yi (Guohua.Yi@uthct.edu), Buka Samten

(Buka.Samten@uthct.edu), and John J. Dennehy (John.Dennehy@qc.cuny.edu)

Abstract

The continuing emergence of new strains of antibiotic-resistant bacteria has renewed interest in phage therapy; however, there has been limited progress in applying phage therapy to multi-drug resistant *Mycobacterium tuberculosis* (*Mtb*) infections. In this study, we tested seven bacteriophage strains able to lyse *Mycobacterium smegmatis* for their *Mtb*-killing activities and found that four efficiently lysed *Mtb* H37Rv in 7H10 agar plates. However, only phage DS6A efficiently killed H37Rv in liquid culture and in *Mtb*-infected human primary macrophages. In subsequent experiments, we infected humanized mice with aerosolized H37Rv, then treated these mice with DS6A intravenously to test its *in vivo* efficacy. We found that DS6A treated mice showed increased body weight and improved pulmonary function relative to control mice. Furthermore, DS6A reduced *Mtb* load in mouse organs with greater efficacy in the spleen. These results demonstrated the feasibility of developing phage therapy as an effective therapeutic against *Mtb* infection.

In 2020, while all focus was on COVID-19, *Mycobacterium tuberculosis* (*Mtb*), the causative pathogen of Tuberculosis (TB), infected 10 million people and caused 1.5 million deaths globally [1], making it one of the leading causes of death by a single infectious agent. More importantly, one-third (>2 billion) of the world's population has been latently infected with *Mtb* (LTBI) [2]. When complicated with other diseases/infections, such as HIV and diabetes, these co-morbidities can cause LTBI patients to develop active TB and increase mortality [3]. Although these devastating facts highlight the threat posed by this deadly bacterium, the emergence of drug resistant *Mtb* strains in recent years has worsened the situation in terms of *Mtb* prevention and control.

Upon establishment of infection, *Mtb* exhibits remarkable abilities to adapt to the local environment to evade the host's immune responses due to its unique and dynamic four-layer cell envelope structure, which can help it adapt to hostile lung microenvironments and facilitate its entry into a non-replicating drug-tolerant persister state. In this state, the bacilli are well protected against antibiotic therapy [4, 5], and are potentially a source of new strains of drug-resistant *Mtb* (DR-TB)[6]. In 2019, around half a million people developed DR-TB, and 78% of these cases developed multidrug-resistant TB (MDR-TB) or extensively drug-resistant TB (XDR-TB), which are resistant to the first-line drugs, rifampicin, and isoniazid, or to second-line drugs [1]. MDR-TB and XDR-TB are more challenging to treat and are associated with increased morbidity and mortality [7]. Therefore, the development of alternative therapies in addition to antibiotics is critical for the advancement of TB therapy in the era of *Mtb* drug resistance.

Phage therapy has emerged as a renewed approach to eliminating bacterial infections [8, 9]. Phages, viruses that infect bacteria, are bacteria's natural enemies and

have been used to control bacterial infections since even before the discovery of antibiotics [10]. In the age of antibiotic resistance, phage therapy has drawn tremendous attention. Recent clinical cases and trials demonstrated that phages can be used to treat antibiotic-resistant bacterial infections with positive clinical outcomes [11-16], and patients with drug-resistant *M. abscessus* and *M. chelonae* have also been successfully treated [17-19]. These prominent bodies of work on using phage therapy to treat other mycobacterial infections inspired us to pursue phage therapy as a potentially effective treatment for *Mtb* infection. However, as to the phage therapy for *Mtb* treatment, the significant advancements have focused on screening lytic phage strains for effective killing of *Mtb* in bacterial culture plates, demonstrated by plaque formation [20-22], but no systematic study has characterized the effectiveness of phages in killing *Mtb* in primary human macrophages or in reliable animal models. Therefore, extensive preclinical studies, especially using primary human macrophages or animal models that resemble human clinical settings, are of exceptional clinical importance in the development of phage therapy for *Mtb* treatment.

One of the major challenges to eliminating *Mtb* infection is *Mtb*'sinduction of granuloma formation once host defenses fail to kill the bacteria. While granuloma formation limits *Mtb* growth, it provides a survival niche for *Mtb* replication when the immune system is weakened. Granulomas, the hallmark of TB pathology, consist of macrophages, neutrophils, and lymphoid cells, including T and B cells, and are formed upon *Mtb* infection in patients [23]. The commonly employed mouse model can be infected with *Mtb*, but *Mtb* can only form a granuloma-like structure in mice. Mouse *Mtb* infections lack the caseous necrotic granulomas that are often observed in TB patients

[24], suggesting human immune cells may be needed to recapitulate real clinical settings.

Recently humanized mice based on NOD-scid IL2Rgamma^{KO} (NSG) mice were

developed. However, the mouse innate immune system in these mice was deficient and

transplanted human hematopoietic stem cells (hHSCs) were generally not well-

developed; thus, the human B, T, and myeloid cells were immature, and NK cells lost

functions [25, 26]. These mouse models provide valuable tools to study *Mtb* infection [24,

27]; however, the myeloid barrier in these NSG mice causes a relative lack of leukocyte

differentiation [26]. The difficulty of forming sufficient granulomas (which need mature

macrophages) makes these mice less able to recapitulate *Mtb* infection in humans.

Nevertheless, the newly developed NSG-SGM3 mice transgenically express three

human cytokine/chemokine genes IL-3, GM-CSF, and KITLG, which can enhance the

differentiation and maturation of the myeloid-lineage cells [26, 28-31]. Moreover, these

three transgenic genes can improve the homeostasis of human CD34+ HSCs, increase

neutrophil and macrophage numbers and function, and stimulate hCD34+ cells to

differentiate into myeloid progenitor cells [30, 32]. Thus, this humanized mouse model

can generate sufficient and fully functional myeloid cells, including macrophages, to study

Mtb infection in the context of an entire human immune system.

In this study, we show that phage DS6A can efficiently kill wild-type *Mtb* H37Rv *in*

vitro (in both 7H10 agar plates and 7H9 liquid culture), in primary human macrophages,

and in humanized NSG-SGM3 mice, demonstrating the feasibility of developing phage

DS6A as an effective therapeutic against *Mtb* infection.

Results

Phage infectivity in *M. smegmatis* and in *M. tuberculosis* H37Rv.

We tested seven bacteriophages for their lysis capacities of *M. smegmatis* and *Mtb*

H37Rv by plaque assay. Serially diluted phages (10^2-10^4 pfu) were incubated with *M.*

smegmatis or H37Rv in 12-well plates with 7H10 agar. The results showed that five phage

strains (Cali, D29, BPs, Adephagia and TM4) lysed *M. smegmatis* (Fig. 1a), and four

phage strains (BPs, Cali, D29, and DS6A) lysed H37Rv and formed plaques (Fig. 1b).

We then tested the *Mtb* killing ability of these seven phage strains in liquid culture

by measuring the optical density at 600 nm (OD_{600}) of the H37Rv suspensions at different

time points after incubation with phages at a multiplicity of infection (MOI, ratio of phage

to bacteria) of 1 by inoculation of 7H9 media with 5×10^4 CFUs of H37Rv and 5×10^4 pfu of

phages. We found that only DS6A significantly reduced *Mtb* OD_{600} on day 9 (Fig. 2a and

2b), while other phages did not reduce *Mtb* OD_{600} on day 9, although several of them

showed the ability to kill *Mtb* in agar plates. We then plated the cultures from different

time points to determine the CFU of the bacilli (Fig. 2c). The CFU results were consistent

with the OD_{600} measurements.

We further confirmed this outcome using a GFP-expressing H37Rvstrain, H37Rv-

GFP, in a liquid culture condition as described above. Due to the highly efficient

expression of GFP gene, the live bacteria are visible under fluorescent microscopy. The

microscopy images of the day 9 cultures clearly showed that there was no GFP

expression in the wells with H37Rv-GFP co-cultured with DS6A, whereas H37Rv in the

wells co-cultured with other phages remain fluorescent, further demonstrating the highly

effective *Mtb*-killing ability of DS6A (Fig. 2d).

Phage DS6A can efficiently eliminate *Mtb* H37Rv in primary human macrophages.

To test whether the phages can kill *Mtb* in infected primary human macrophages,

which is critical for therapeutic effectiveness in clinical settings, we isolated peripheral

blood mononuclear cells (PBMCs) from the blood of a healthy donor and isolated the

hCD14+ monocytes using microbeads to above 95% purity as determined by flow

cytometry analysis (Supplementary Fig. 1) and differentiated the purified CD14+ cells into

macrophages by incubation of the cells with hM-CSF, hGM-CSF, and hIL-4 for 5 days.

The macrophages were then infected with H37Rv and treated with different phages. On

days 5 and 10, we plated the macrophage lysates on 7H10 agar plates to determine the

CFUs (Fig. 3a). The results showed no reduction in *Mtb* CFUs on day 5 when compared

to the *Mtb*-infected macrophages without phage treatment as a control. However, on day

10, phage DS6A completely eradicated the bacilli from the infected macrophages, while

other phages still failed to show any reduction in *Mtb* growth in macrophages (Fig. 3b and

3c).

We further tested whether the phages can also kill *Mtb* in infected macrophages

from different donors. For this purpose, we isolated monocytes from four healthy donors'

PBMCs, and repeated the above experiment in these four samples using phage DS6A,

and phage BPs was used as a negative control. Meanwhile, we wanted to see whether

Mtb could be killed in a shorter time rather than ten days, so we set the sampling time

points to days 4 and 7. The results showed that all four donors displayed the same pattern

in which *Mtb* in the infected macrophages were killed within seven days by phage DS6A,

but not in the BPs treatment (Fig. 3d and 3e). Interestingly, some of the phage-treated

cultures have a significantly higher bacillary load than the control bacteria cultures (Fig.

3b and 3d), which we address further in the discussion section.

149   ***Mtb* infection of humanized NSG-SGM3 mice.**

150   Compared to the commonly used Hu-PBL and BLT mouse models, the humanized

151   NSG-SGM3 mouse model has several advantages: 1) Humanized NSG-SGM3 mice

152   exhibit improved reconstitution with an increased general population of human immune

153   cells (hCD45+); 2) Cytokine expression supports human myeloid cell differentiation and

154   maturation; thus the functional macrophages and dendritic cells are significantly elevated

155   in the bone marrow compared with NSG recipients [26, 29]; 3) There is a significant

156   increase of regulatory T cells (Treg) [29], which are important in regulating *Mtb*

157   pathogenesis [33, 34]; 4) B cells can be developed to mature phenotypes with the ability

158   of class-switching [25], which is important to evaluate the IgM and IgG responses against

159   phages; 5) They are relatively easy to generate by intravenous injection of human HSC

160   into irradiated adult mice for humanization.

161   We obtained NSG-SGM3 mice from the Jackson Laboratory and are breeding

162   them in-house. For humanization, 4-6-week-old NSG-SGM3 mice were irradiated at

163   150cGy and then infused with human CD34+ HSC. Ten to fifteen weeks after HSC

164   injection, the animals developed a full complement of human immune cell types, including

165   CD4+ and CD8+ T cells, B cells, myeloid cells, and natural killer (NK) cells (Fig. 4a).

166   To establish humanized NSG-SGM3 mouse model of *Mtb* infection, we infected

167   the humanized NSG-SGM3 mice with H37Rv via a Madison chamber to deposit about

168   50-100 CFUs per mouse lung. At 25 days post-infection, we sacrificed the mice and used

169   the homogenates from the lungs and the spleens to analyze the immune cell profile after

170   infection. Additionally, we determined *Mtb* growth in lungs and spleens. The results

171   showed that, when compared with the PBMCs analysis before infection, the spleen

immune cells have undergone substantial changes after infection, and the CD8+ T cells

and myeloid cells have significantly expanded (Fig. 4b). Meanwhile, from both the lung

and spleen homogenates, we can detect a high number of bacilli (Fig. 4c), indicating that

the mice established the *Mtb* reservoirs not only in the lungs but also systematically.

Phage DS6A effectively eradicates *Mtb* in humanized mice.

To investigate if phage DS6A can eradicate *Mtb in vivo*, we performed the animal

experiment in the humanized NSG-SGM3 mouse model of tuberculosis (Fig. 5a). We

infected the humanized NSG-SGM3 mice with low dose aerosolized H37Rv and as

confirmed by an average ~100 bacilli per lung deposited at day one post infection (Fig.

5c). From day 3 of *Mtb* infection, we treated the mice every other day with ten doses of

DS6A (10^{11} phage particles/dose) via intravenous administration.

After the treatments, we determined the growth of *Mtb* in mouse lungs and spleens,

performed pulmonary function tests (PFTs) and overall health evaluation by monitoring

mouse body weight changes in association with changes in bacillary burden of the mice.

The results showed that DS6A-treated *Mtb*-infected mice gained significantly more weight

than control *Mtb*-infected mice (Fig. 5b). The PFTs also showed that two of the four

parameters tested in the phage-treated group had significantly improved, with less

respiratory resistance and better elasticity, demonstrating improved pulmonary function

after treatments (Fig. 5d). The CT scan results showed that there were fewer high-density

areas with much cleaner lungs in the phage-treated group of mice, which indicates less

inflammation and other pathological changes (Fig. 5e). Furthermore, the lung volume of

the phage-treated mice was larger than untreated *Mtb* control mice, even though the

difference didn't reach significance (p=0.1782) (Fig. 5d). The CFU counts showed that

the bacilli were completely eradicated in the spleens in six of nine mice of the phage-

treated group, and the average bacillary load in the spleen was significantly lower than in

the *Mtb*-infected control group (Fig. 5c). Moreover, the lung bacillary load showed three-

times higher in *Mtb*-infected control mice than in phage-treated mice, while the difference

didn't reach significance (Fig. 5c). These results suggest that, after intravenous

administration, phage DS6A has the capacity to kill the replicating *Mtb* within/or outside

the lung that are disseminated to the blood circulation.

To explain why the phage DS6A performed better in killing *Mtb* bacilli in spleens

than in lungs, we determined the phage genome copies in the lung and spleen

homogenates. The quantitative PCR result showed that the spleen had three times more

phage copies in spleens than in lungs (Supplementary Fig. 2), suggesting that the

effectiveness in tissues may be positively correlated to the phage copies, and that phages

intravenously administered preferentially distributed and replicated more in the spleen

than in the lungs, which may explain the organ-dependent differences in *Mtb* eradication

in humanized mice.

Given that phage DS6A could not wholly eradicate the spleen bacteria in several

mice, we wondered whether phage resistance developed in the bacteria within these

mice. For this purpose, we picked colonies from the plates that spread the lung and spleen

homogenates of phage-treated mice, and then mixed them with phage DS6A to see if the

phage could kill the bacteria *in vitro*. The results showed that the negative control phage

could not kill both bacteria colonies picked from DS6A-treated or untreated mice, while

the DS6A could kill both (Fig. 5f), indicating that there was no phage resistance developed

during the 20-day phage treatment period.

Phage therapy elicits antibody responses in humanized NSG-SGM3 mice, and it doesn't affect the function of immune cells.

There is always a concern that the immune responses, especially antibody responses, will stymie the effect of phage therapy. Therefore, we determined the phage DS6A-specific antibody responses in sera from our experimental mice by ELISA. We detected a significantly higher level of human IgM in phage DS6A-treated mouse sera (Fig. 6b). However, we could only detect weak IgA and IgG responses in these mice (Fig. 6a and c).

We further tested whether phage therapy could affect the cytokine profiles that will be in association with reduced *Mtb* growth and improved lung function due to the phage treatment. We determined the levels of 27 cytokines including cytokines of human lymphocytes and myeloid cells in the lung and spleen homogenates of the mice by a multiplex cytokine assay (Supplementary Fig. 3). Overall, the cytokine results showed that the cytokine levels in the spleen were higher than that of the lungs, and that the phage treatment induced a few myeloid cells-derived cytokines significantly lower than that of the control *Mtb*-infected mice, consistent with reduced bacilli burden in the mice treated with phage. We did not see significantly elevated levels of T cell cytokines in the mice after four weeks of *Mtb* infection, consistent with significantly increased CD14+ monocytes, but not lymphocytes, in the humanized mice after *Mtb* infection.

Discussion

The development of alternative treatments to antibiotics for *Mtb* infection is an urgent task in the era of antibiotic resistance. In this study, we have shown that several bacteriophages can efficiently kill *Mtb* bacilli on 7H10 agar plates, while only phage DS6A significantly reduced *Mtb* OD_{600} in liquid cultures. Similarly, only phage DS6A was able to kill *Mtb* bacilli in primary human macrophages. In addition, we showed that phage DS6A could effectively eradicate *Mtb* bacilli in spleens and reduced the bacilli load in the lungs in a humanized mouse model of *Mtb* infection. Thus, phage DS6A appears to have the potential to be developed as an effective therapeutic for *Mtb* treatment.

Phage therapy has shown great potential in controlling some intransigent drug-resistant bacterial infections. Compared to typical antibiotic treatment, phage therapy has advantages such as: host specificity, low toxicity, the ability to lyse the bacteria, increased effectiveness (since they can self-replicate *in situ*), and a narrower potential for inducing resistance, all of which render it an attractive treatment strategy in the era of antibiotic resistance. Several phage strains have shown potential in eliminating *Mtb* infection [20]. Unfortunately, there has been little progress in *Mtb* phage therapy, with only two studies published to date that reported on the use of phage therapy for the treatment of *Mtb* in guinea pigs, with limited success [35, 36]. Therefore, our study showing the effectiveness of phage therapy in treating *Mtb* infections of clinically relevant humanized mice represents a significant advancement in this field.

Humanized mouse models provide versatile tools to study various infectious diseases. For *Mtb* infection, the myeloid cell lineage is especially important because the macrophages are the main targets of the bacilli. In most NSG-based humanized mice,

myeloid cell differentiation is not efficient, and the macrophages are not mature, thus the

physiological processes such as phagocytosis and endocytosis, which may be involved

in *Mtb* infection and bacteriophage uptake, may not be completed as properly as they are

in humans. However, due to the transgenic expression of the three human cytokine

genes, the myeloid cells were well-differentiated in NSG-SGM3 humanized mice that we

have used in this study. After *Mtb* infection, the myeloid cells were successfully

propagated (Fig. 4b). Moreover, after phage treatments, the phage was shown capable

of killing *Mtb* in human macrophages engrafted in the humanized mice, suggesting that

NSG-SGM3 mice can recapitulate human *Mtb* infections, and thus NSG-SGM3 mice are

an appropriate and clinically relevant animal model for *Mtb* phage therapy.

Although D29, BPs, Cali and DS6A each showed the ability to kill *Mtb* on bacterial

lawns, only DS6A efficiently killed *Mtb* in liquid cultures, in human primary macrophages,

and in humanized mice, suggesting that DS6A is the most appropriate phage therapeutic

for *Mtb* infections. The reasons why other the phages tested can kill H37Rv efficiently on

bacterial lawns, but not in liquid culture, are not fully clear (Fig. 2). The reasons may be

different for each phage. The most likely explanation for why D29, BPs, and Cali kill *Mtb*

on 7H10 plates, but not in liquid culture is that *Mtb* readily acquires resistance to D29,

BPs, and Cali, but not to DS6A (Fig. 5f). When inoculated in liquid culture at an MOI = 1,

it may be that at least some *Mtb* cells are genetically incapable of being infected by D29,

BPs, and Cali, thus these resistant *Mtb* cells eventually dominate the culture. Additionally,

it appears that D29, BPs, and Cali cannot easily break this resistance. Nevertheless,

these results indicate that liquid cultures, rather than bacteria lawns, appears to be a

better prediction model for the *in vivo* efficacy of *Mtb* phage therapy.

Of note, we also observed that once the phage cannot kill *Mtb* in liquid culturing

condition, it sometimes will stimulate *Mtb* growth, as shown in Fig. 3. The observation

that, in some cultures, bacteria load is higher in the presence of phages Adephagia,

BPs, D29, Pixie, and TM4 (but not Cali and DS6A), than in phage-free controls (Fig. 3b

and 3d) is puzzling.

Nevertheless, our experiments provide clear evidence that phage DS6A was

effective in killing *Mtb* in the spleen via the intravenous route. However, the bacterial

clearance in the lungs was not as effective as in spleens. This discrepancy is possibly

due to the administration route, as this is one of the most important factors that affects *in*

vivo efficacy of phage therapy [37]. Through the intravenous route, phages can easily

reach the spleen and liver through circulation as blood filtering organs, but the lungs are

much more difficult to reach. A previous study showed that the intravenous delivery of a

pneumonia bacteria-specific phage led to a 2-log lower prevalence (<100 times) in the

lungs when compared to an intratracheal delivery route [37]. Additionally, phages need

to penetrate the lung epithelial barrier [38] in order to access the alveolar macrophages

that are parasitized by *Mtb*, thus further increasing the difficulty of efficient intravenous

phage administration.

To test this hypothesis, we compared phage copy number in spleens and in lungs,

and the results indicated there are more phage copies in spleen than that in lungs. It is

worth noting that our original dose contained 10^{11} phages per dose, while we only

detected thousands of copies at the termination time point. This discrepancy may be due

to the pharmacodynamical changes of the phages, as the phage's half-life might be as

low as 2.2 h and was undetectable after 36 h of injection [39, 40]. Even though we need

307 to further investigate the pharmacokinetics of the phage DS6A, our qPCR result offered

308 a reasonable explanation of why the intravenous administration of phage is more efficient

309 in spleens than in lungs. Nevertheless, our results suggest that optimization of

310 administration will be helpful to improve lung bacterial clearance (e.g., via intratracheal

311 route or nebulizer delivery, etc.) as tuberculosis, in majority of cases, are infections of the

312 lungs.

313 Phage resistance has always been a concern as it may dampen the therapeutic

314 efficacy of phage therapy. In our study, it seems not to be a case for phage DS6A within

315 the treatment period (Fig. 5f). However, in the clinical settings, the treatment time may be

316 as long as several months, and *Mtb*-resistant strains may still appear. Therefore,

317 screening more effective bacteriophage strains to form a phage cocktail would be

318 essential to overcome phage resistance by *Mtb*.

319 Moreover, our study showed that significant IgM but weaker IgA and IgG

320 responses were induced after 20 days of phage treatment (Fig. 6). This is because the

321 short-term administration of phage may lead to IgM as dominant antibodies, and with the

322 prolonged administration, the IgG response will become dominant. Given that phage

323 treatment still improved disease progression in a human clinical case of *M. chelonae*

324 despite a significant anti-phage antibody response [18], it may be that are not a great

325 concern. Indeed, in our study, the antibody responses did not adversely affect the efficacy

326 of phage therapy to eradicate *Mtb* in the spleen. This may be because our treatment

327 regimen was not deployed long enough to enable to the development of sufficient quantity

328 and quality of antibody responses to counteract the high dose of phages we administered.

However, in clinical settings where a long treatment regimen might be needed, a high

dose might be warranted to reduce the effects of antibody responses to the phages.

Taken together, our study showed that bacteriophage DS6A could effectively

eliminate *Mtb* in agar plates, in liquid culture, in infected macrophages, as well as in

humanized mouse model. Therefore, our study demonstrated the feasibility of developing

phage therapy as an effective therapeutic against *Mtb* infection.

Materials and Methods

<u>Regulatory and Ethical statement</u>

Healthy donors were recruited for the collection of blood samples from the

employees and students at the University of Texas Health Science Center at Tyler

(UTHSCT) following the protocol approved by the Institutional Review Board (IRB) ethics

committee of the UTHSCT. All animal procedures were approved by the UTHSCT

Institutional Animal Care and Use Committee (IACUC).

<u>Bacterial strains, phage strains, media, and culture conditions.</u>

Bacteriophage DS6A was purchased from the ATCC (Cat# 25618-B2), and other

phages Adephagia, BPs, Cali, D29, Pixie, and TM4 were provided by Dr. Dennehy at

Queens College, The City University of New York. *M. smegmatis* mc2 155 was purchased

from the ATCC (Cat# 700084). *Mtb* H37Rv and *Mtb* H37Rv-GFP were provided by Dr.

Samten's at UTHSCT. Except for DS6A, all mycobacteriophages were amplified in *M.*

smegmatis; DS6A was amplified in *Mtb* H37Rv in the BSL-3 Laboratory at UTHSCT.

<u>Amplification and quantification of bacteriophages</u>

Bacteriophages, except DS6A, were amplified in *M. smegmatis. M. smegmatis*

was streaked on 7H10 agar plate supplemented with 10% Middlebrook Oleic Albumin

Dextrose Catalase (OADC), 1mM $CaCl_2$, 10 µg/mL of carbenicillin, and 100 µg/mL of

cycloheximide and cultured at 37°C for about 3 - 4 days for single colony growth. A single

colony of *M. smegmatis* was cultured in 7H9 media supplemented with 10% Middlebrook

Albumin Dextrose Catalase enrichment supplement (ADC), 1mM $CaCl_2$, 10 µg/mL of

carbenicillin, 100 µg/mL of cycloheximide, and 0.25% Tween-80 at 37°C for 3-4 days until

the OD at 600 nm was over 2.0. A hundred mLs of *M. smegmatis* from the above culture

was inoculated into 5 mL of 7H9 media supplemented with 10% ADC, 1mM $CaCl_2$, 10

µg/mL of carbenicillin, 100 µg/mL of cycloheximide without Tween-80, and incubated

overnight with shaking at 200 rpm at 37°C. Next day, 10^8 pfu of bacteriophages with 100

µL of *M. smegmatis* were mixed and incubated at 37°C for 10 min, then the phage-bacilli

mixture was added to 7mL of top agar at 60°C, and immediately applied onto 150-mm

7H10 agar plates. The phage-bacilli agar plates were incubated at 37°C overnight, then

the bacilli were collected in 10 mL of phage buffer (10 mM Tris, pH 7.5, 10 mM MgSO4,

0.4% NaCl) and stored at 4°C overnight. After centrifugation, the supernatants were

collected and filtered through a 0.22 µM PVDF filter, and the phages were stored at -80°C

until use.

Mtb was cultured in 7H9 with 10% ADC following the standard *Mtb* culture

procedures. *Mtb* single cell suspensions at 10^8 cells per ml were infected with phage

DS6A using an MOI of 1 for 10 min at 37 °C. The infection mixture was mixed with 7 mL

of top agar (0.8% agarose in 7H9 with 10% ADC) and applied onto 150 mm 7H10 agar

plate (supplemented with 10% OADC, 1 mL of $CaCl_2$). The plates were incubated at 37°C

for ~8 days. After plaques formed, the phage DS6A was collected in phage buffer and

filtered through a 0.22 μM PVDF filter.

Bacteriophages' titers were determined by plaque assays by infection of H37Rv

with phage DS6A or *M. smegmatis* with other phages, respectively, after serially diluted

the phages. Alternatively, quantitative PCR was also used to quantify the viral genomes.

<u>Testing *Mtb*-killing ability on 7H10 agar plates</u>

To test whether the bacteriophages can kill *Mtb* on the solid agar plate culture, we

performed the plaque assays. Briefly, 10^4, 10^3, 10^2 pfu of Adephagia, BPs, Cali, D29,

Pixie, TM4 (titer based on *M. smegmatis*), and DS6A (titer based on *Mtb*) were mixed

with 1 x 10^5 *Mtb* bacilli and incubated at 37°C for 1 h with shaking. The *Mtb*-phage mixture

was then mixed with 500 μL of 0.8% top agar in 7H9 with 10% ADC and applied to 12-

well 7H10 agar plates. The plates were incubated at 37°C for ~7-8 days until the plaques

developed.

<u>Testing *Mtb*-killing ability in 7H9 liquid culture</u>

The *Mtb* killing activities of the bacteriophages were determined by infecting 1 x

10^5 H37Rv in 1 mL 7H9 media with various bacteriophages at an MOI of 1 for 10 min,

then inoculated into 20 mL 7H9 media supplemented with 10% ADC and incubated at

37°C for 9 days. At days 3, 6, and 9, 10 μL of 10-fold serial dilutions of the culture were

plated on 7H10 agar plates, and CFU were counted after cultured at 37°C for 2-3 weeks.

<u>Flow cytometry</u>

Cells from one well were collected and stained for macrophage markers. The

antibodies used were APC-conjugated CD11b (cat# 301350), FITC-conjugated CD14

(cat# 325604), BV711-conjugated CD16 (cat# 302044). The isotype controls included

APC-mouse IgG1 (cat# 400142), FITC-mouse IgG1 (cat# 400108), BV711-mouse IgG1

(cat# 400168). All antibodies were purchased from BioLegend. Briefly, the cells were

washed with PBS and incubated with the antibodies at room temperature for 15 min,

avoiding light. The cells were then washed and resuspended with a buffer of 0.5%

BSA/PBS + 3mM EDTA prior to analysis with Attune NXT (ThermoFisher). Compensation

was performed using UltraComp eBeads Plus (01-3333-42, Invitrogen). The data were

then analyzed with FlowJo version 10.6.1, and the graphs were created with GraphPad

Prism 8.

<u>Testing the *Mtb*-killing ability of the bacteriophages in macrophages</u>

Blood was drawn from five healthy donors and PBMCs were isolated using Ficoll-

Paque Plus (17-1440-02, GE Healthcare, Danderyd, Sweden) gradient centrifugation.

Then, CD14+ cells were isolated from the PBMCs using human CD14+ micro beads

conjugated with anti-human CD14 mAbs (130-118-906, Miltenyi Biotec, Bergisch

Gladbach, Germany) LC columns (130-042-401, Miltenyi Biotec, Bergisch Gladbach,

Germany) following the manufacturing instructions. The purity of the cells was checked

with flow cytometry.

Half a million cells were seeded in each well of 24-well plates containing 1.5 mL of

RPMI-1640 supplemented with 10% FBS and 1% penicillin + streptomycin. The cells were

then incubated in the presence of 50 ng/mL of GM-CSF (02532, Stem Cells, Vancouver,

Canada) and 50 ng/mL of M-CSF (216-MC, R&D Systems, Minneapolis, USA) at 37°C

and 5% CO_2 for differentiation. Fresh cytokines were replenished on day 3 once. On day

6, some cells were collected and stained for macrophage markers followed by flow

cytometry analysis. The rest of the cells were washed with PBS and rested overnight in

fresh medium without serum and antibiotics. The cells were then infected with H37Rv at

an MOI of 1. Four hours post infection, the cells were washed with warm PBS thrice to

remove extracellular bacilli. The cells were then incubated in fresh RPMI-1640 with 10%

FBS with no antibiotics as controls and with either DS6A or BPs phages at 37°C. The

growth of H37Rv in macrophages were followed for different time points after infection by

plating serially diluted cellular lysates on 7H10 agar plates as follows: After removing the

culture medium, the cells were treated with ddH$_2$O for 10 minutes followed by 0.25% SDS

in HBSS for 10 min. At the end, 20% BSA in PBS was added to stop the lysis. The content

of each well was collected into a tube and sonicated three times with 33% amplitude for

10 seconds. After making dilutions, 10 µL of cellular lysate was plated on 7H10 agar

plates in duplicates. The plates were incubated at 37°C for 2-3 weeks and the bacterial

colonies were counted for CFU determination.

<u>Generation of humanized mice</u>

Humanized NSG-SGM3 mice (NSG mice transgenically expressing three human

cytokine/chemokine genes IL-3, GM-CSF, and KITLG, RRID: IMSR_JAX:013062) were

obtained from Jackson Laboratories and bred in UTHSCT vivarium. To increase scientific

rigor, the experiments have been done in a blinded manner. To exclude sex bias, each

experiment was considered to contain an equal number of male and female mice. For

humanization of NSG-SGM3 mice, the mice were irradiated with a dose of 100 cGy per

mouse and injected intravenously with 0.2 million of HSCs (purchased from Stemcell

Technologies Inc.). After 8–10 weeks, the human cell reconstitution was confirmed by

flow cytometry analysis of human immune cells in the mice by staining the PBMCs with

fluorescent human antibodies specific for CD45, CD3, CD4, CD8, CD14, CD19, and

CD56.

Mtb infection and phage treatment of humanized mice and other animal experiments

Humanized mice were infected with a low dose of aerosolized H37Rv in a Madison

chamber as previously described [41] by properly diluted H37Rv stocks in 10 mL normal

saline to deposit about 100 CFUs per mouse lung as confirmed by the determination of

Lung CFUs 1 day post infection. All animal experiments were done in a blinded manner

to increase scientific rigor. Wherever possible, each experiment contains equal numbers

of male and female mice to exclude sex bias.

Mtb-infected mice were treated with 1 x 10^{11} pfu phage DS6A per mouse or equal

volumes of PBS as a control by intravenous administration every other day for a total of

8 treatments. The mice's body weight was measured every other day during the

treatment. Three days after the last treatment, the mice were subjected to pulmonary

function testing and CT scan as described previously [42]. Briefly, mice were anesthetized

with a ketamine/xylazine mixture, and the anesthetized mice were cannulated with a

sterile, 20-gauge intravenous cannula through the vocal cords into the trachea. The

snapshot perturbation method was used to evaluate lung compliance by collecting the

measurements of elastance, compliance, and total lung resistance for each mouse using

the FlexiVent system (SCIREQ, Tempe, AZ). The flexiVent was set to a tidal volume of

30 mL/kg at a frequency of 150 breaths/min against 2–3 cm H_2O positive end-expiratory

pressure, according to manufacturer's instructions.

After lung function testing, the mice were immediately subjected to CT scans for

the measurements of lung volume. The Explore Locus Micro-CT Scanner (General

Electric, GE Healthcare, Wauwatosa, WI) was used for CT imaging. CT scans were
performed during full inspiration and at a resolution of 93 μm. Lung volumes were
calculated from lung renditions collected at full inspiration. Microview software 2.2
(http://microview.sourceforge.net) was used to analyze lung volumes and render three-
dimensional images. The mice were maintained under anesthesia using isofluorane
throughout the pulmonary function testing.

At the end of testing, the mice were euthanized, and the lungs and spleens were
collected and homogenized. The serially diluted lung and spleen homogenates were
plated on 7H10 agar plates with OADC and incubated at 37°C for 2-3 weeks for the
determination of the organ CFUs.

<u>ELISA for detection of phage-specific antibodies</u>

Phage DS6A were coated in 96-well Nunc-Immuno plates (Thermofisher) with a
concentration of 5×10^9 pfu/mL in coating buffer (Na2CO3, pH8.0) at 4°C overnight. After
washing with PBST (0.1% Tween-20 in PBS) four times, the plate was blocked with PBSM
(5% skimmed milk in PBS) and followed by applying 10-fold serially diluted humanized
mouse sera and incubated for 2 h at room temperature, then washed with PBST for 5
times. 100 mL of HRP-labeled secondary antibodies were applied to the plates and
incubated at room temperature for 1 h. The secondary antibodies included: (1) goat anti-
human IgG Fc (HRP) pre-adsorbed (catalog no. ab98624; Abcam); (2) goat anti-human
IgA alpha chain (HRP) pre-adsorbed (catalog no. ab98558; Abcam); and (3) goat anti-
human IgM mu chain (HRP) pre-adsorbed (catalog no. ab98549; Abcam). After washing
away the unbound secondary antibodies, TMB substrate was added and the OD values
at 450 nm were recorded using a BioTek Synergy 2 plate reader.

Multiplex assay for cytokine profiling

Lung and spleen tissue samples from both experimental groups were homogenized in PBS to a final volume of 1.5mL, using a 70 µm cell strainer (MTC Bio, Sayreville, NJ). The homogenate was passed through a 0.2 µm filter (Sigma-Aldrich, St. Louis, MO), and the flowthrough was used to evaluate the cytokine profile in these tissues. The samples were analyzed in duplicates using the Bio-Plex Pro™ Human Cytokine panel (Bio-Rad, Hercules, CA), according to the manufacturer's instructions. Briefly, 50 µL of filtered tissue homogenate were dispensed in a 96-well plate containing magnetic beads conjugated with antibodies for the detection of 27 different cytokines. After further incubation with detection antibodies and streptavidin-PE, the samples were analyzed in the Bio-Plex MAGPIX multiplex reader (Bio-Rad Laboratories Inc., CA). Fluorescence values were converted into cytokine concentrations (expressed as pg/mL), using a regression curve based on the values obtained from a set of standard dilutions.

The cytokines reported by the Bio-Plex Pro™ Human Cytokine panel were: Basic FGF, Eotaxin, G-CSF, GM-CSF, IFN-γ, IL-1β, IL-1Ra, IL-2, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-12, IL-13, IL-15, IL-17, IP-10, MCP-1, MIP-1α, MIP-1β, PDGF-BB, RANTES, TNF-α and VEGF.

Statistical analysis

Each treatment was triplicated (PCR) or duplicated (all other experiments), and the experiments were repeated at least once to ensure reproducibility. Power analysis was done to determine the sample size to ensure biological significance. The data were analyzed using GraphPad Prism software. Unpaired student T-tests were used to analyze the differences between treated and control groups. All statistical data are represented

511   as mean ± SEM. Statistical significance was defined as *P≤0.05, **P≤0.01, and

512   ***P≤0.001.

Data availability

All data supporting the findings of this study are available in the manuscript. If there are any special requests or questions for the data, please contact the corresponding author (G.Y.).

Acknowledgments

We thank Dr. Amy Tvinnereim for helping perform the following experiments: *Mtb* infection of the humanized mice and the pulmonary function test.

Funding:

This work was partially supported by the NIH Common funds and the National Institute of Allergy and Infectious Diseases grant UG3AI150550, and the National Heart, Lung, and Blood Institute grant R01HL125016 to G.Y., and NIH grant no. R21AI156798 to J.J.D..

Contributions

Guohua Yi: Conceived and guided the study, designed and performed experiments, analyzed data, edited figures, and wrote and finalized the manuscript.

Fan Yang: Performed the experiments and analyzed data.

Alireza Labani-Motlagh: Performed the experiments and analyzed data.

Josimar Dornelas Moreira: Performed the experiments.

Danish Ansari: Performed the experiments.

Jose Alejandro Bohorquez Garzon: Performed the experiments.

Sahil Patel: Performed the experiments.

Fabrizio Spagnolo: Performed the experiments and edited the manuscript.

Jon Florence: Performed the experiments.

Abhinav Vankayalapati: Performed the experiments.

Ramakrishna Vankayalapati: Participated in experiment discussions, provided suggestions and edited the manuscript.

John J. Dennehy: Designed experiments, participated in experiment discussions, provided phage strains, and edited the manuscript.

544 Buka Samten: Designed experiments, guided experiments, performed experiments,
545 participated in experiment discussions, and edited the manuscript.

547 **Competing interests**

548 All the authors declare no competing interests.

Figure legends

Figure 1. Phage infectivity of *M. smegmatis* and *Mtb* H37Rv in solid agar plates.

Serial dilutions (10^2-10^4 pfu) of different strains of bacteriophages were mixed with 1 x

10^5 CFUs *M. smegmatis* **(a)** or H37Rv **(b)** in 7H9 medium and incubated at 37°C with

shaking for 1 h. The infection cultures were then mixed with 0.8% top agar and spread

on 7H10 agar plates supplemented with 10% OADC enrichment.

Figure 2. Phage infectivity of *Mtb* in liquid culture. H37Rv (1 x 10^5 CFUs) were

infected with various bacteriophages at an MOI of 1 for 1 h, then inoculated into 20 mL

7H9 media supplemented with 10% ADC and incubated at 37°C for 9 days. The cultures

were sampled at days 3, 6, and 9 post infection by plating 10 µL of 10-fold serial dilutions

of on the 7H10 agar plates and cultured at 37°C for CFU determination. **(a)** OD_{600} of

sampled liquid cultures was measured at different time points as indicated. **(b)** Statistics

of the CFUs of each sampled liquid culture sampled at various time points. **(c)** Titering

assay of phage-*Mtb* liquid culture sampled at different time points. The data are

representative results of two independent experiments. **(d)** Phage infection of H37Rv that

expresses a GFP reporter gene. An MOI of 1 was used to infect H37Rv-GFP. GFP

expression was pictured under fluorescence microscopy after nine days of phage

infection.

Figure 3. Phage DS6A eliminates *Mtb* in infected primary human macrophages. (a)

Experimental procedures of the assay. Panels **(b)** and **(c):** Half million microphages were

infected with H37Rv at an MOI of 1. After 4 h of *Mtb* infection, seven different phages

($5x10^6$, MOI of 10) were applied to the *Mtb*-infected primary human macrophages, and

cultured at 37°C. The cells were sampled at days 0, 5 and 10 to determine the bacillary

load for the evaluation the *Mtb*-killing ability of the phages by plating the ultrasound-

broken cells on 7H10 plates (supplemented with 10% OADC) and the CFUs were counted

on Day 8. Panel **(b)** shows the statistics of the CFUs of all phages, and panel **(c)** shows

the representative pictures (duplicated) of the 7H10 plates taken at day 5 and day 10.

Panels **(d)** and **(e):** Testing the *Mtb*-killing ability in *Mtb*-infected macrophages derived

from five different healthy donors. The infected macrophages were sampled at different

time points, sonicated, and plated on 7H10 agar plates using different dilutions to titer the

bacillary load. Panel **(d)** shows the statistics of five donors, and panel **(e)** shows the

representative pictures taken from 10x diluted plates for days 4 and 7 cultures.

Figure 4. *Mtb*-infection of humanized NSG-SGM3 mice. (a) Reconstitution of human

immune cells in humanized NSG-SGM3 mice. NSG-SGM3 mice were irradiated and

intravenously injected with 2×10^5 hCD34+ HSC. After 10-12 weeks, reconstitution of

human T cells, B cells, myeloid cells, and natural killer (NK) cells within human CD45-

gated population was analyzed by flow cytometry. **(b)** Statistics of different immune cells

before and after infection. **(c)** Five mice were sacrificed to check *Mtb* infection in lungs

and spleens by determining the organ CFUs by plating serially diluted organ

homogenates on 7H10 agar plates as described above.

Figure 5. Phage DS6A is effective at eradicating *Mtb* in humanized NSG-SGM3 mice.

(a) Schematic presentation of the animal experiment procedures. H37Rv aerosol infected

humanized mice were treated three days after infection with 1×10^{11} pfu of phage DS6A

per mouse or equal volume of sterile PBS every other day for eight times. The mouse

body weights were monitored over time, and the body weight change over the initial body

weight of the mice are shown as percentages **(b)** The growth of H37Rv in the mice lung

and spleen homogenates together with initial infection dose (left panel in purple) are

shown (**c**) Before termination of the experiment, the mice were subjected to pulmonary

function testing. The measurements of elastance, compliance, total lung resistance, and

total lung volume were collected, and the statistics are shown **(d)**. CT scans were

performed for each mouse, and two representative figures show the *Mtb*-infected control

mouse and phage-treated mouse lungs, respectively **(e)**. The left panel of each mouse

CT scan figure shows the 3D image, the white areas represent the high-density scan

(e.g., tissues), while the black areas represent low density scan (e.g., air). The three small

figures in each mouse scan show different angles of scan results. The colonies from the

plates that spread with lung or spleen homogenates of the H37Rv infected control or

phage-treated mice were picked and mixed with 1 x 10^6 pfu of phage DS6A or a negative

control phage (TM4) in 10 µl 7H9 media and incubated at 37°C for 1h. Then the infection

cultures were pipetted onto the 7H10 agar plates OADC and incubated at 37°C for

determination of CFUs. (**f**) The animal experiments were performed four times, and the

3^{rd} and 4^{th} experiments were shown. Except the pulmonary functions were tested once

(3^{rd} animal experiment), the other data are shown as the combined results from the 3^{rd}

and 4^{th} experiments. All the statistics are shown as mean ± SEM.

Figure 6. Humanized NSG-SGM3 mice developed antibody responses against

phage DS6A. ELISA was performed to determine the phage DS6A-specific human IgA,

IgM, and IgG titers in sera of the humanized mice infected with H37Rv and treated with

phage DS6A or PBS buffer control. The experiment was repeated once, and the results

from one experiment with five mice each group are shown.

Figures

623   **Figure 1**

629 **Figure 2**

Figure 3

Figure 4

Figure 5

658   **Figure 6**

(A)

(B)

(C)

References

1 WHO. Global Tuberculosis Report2021.

2 Venketaraman V. Recent Advances in Mycobacterial Research. *J Clin Med* 2020; 9.

3 Bates M, Marais BJ, Zumla A. Tuberculosis Comorbidity with Communicable and
Noncommunicable Diseases. *Cold Spring Harb Perspect Med* 2015; 5.

4 Allue-Guardia A, Garcia JI, Torrelles JB. Evolution of Drug-Resistant Mycobacterium tuberculosis
Strains and Their Adaptation to the Human Lung Environment. *Front Microbiol* 2021; 12: 612675.

5 Lee JJ, Lee SK, Song N, Nathan TO, Swarts BM, Eum SY *et al*. Transient drug-tolerance and
permanent drug-resistance rely on the trehalose-catalytic shift in Mycobacterium tuberculosis.
Nat Commun 2019; 10: 2928.

6 Trubenova B, Roizman D, Moter A, Rolff J, Regoes RR. Population genetics, biofilm recalcitrance,
and antibiotic resistance evolution. *Trends Microbiol* 2022; 30: 841-852.

7 Ali MH, Alrasheedy AA, Kibuule D, Godman B, Hassali MA, Ali HMH. Assessment of multidrug-
resistant tuberculosis (MDR-TB) treatment outcomes in Sudan; findings and implications. *Expert
Rev Anti Infect Ther* 2019; 17: 927-937.

8 Liu C, Hong Q, Chang RYK, Kwok PCL, Chan HK. Phage-Antibiotic Therapy as a Promising Strategy
to Combat Multidrug-Resistant Infections and to Enhance Antimicrobial Efficiency. *Antibiotics
(Basel)* 2022; 11.

9 Allue-Guardia A, Saranathan R, Chan J, Torrelles JB. Mycobacteriophages as Potential Therapeutic
Agents against Drug-Resistant Tuberculosis. *Int J Mol Sci* 2021; 22.

10 Lin DM, Koskella B, Lin HC. Phage therapy: An alternative to antibiotics in the age of multi-drug
resistance. *World J Gastrointest Pharmacol Ther* 2017; 8: 162-173.

11 Duplessis C, Biswas B, Hanisch B, Perkins M, Henry M, Quinones J *et al*. Refractory Pseudomonas
Bacteremia in a 2-Year-Old Sterilized by Bacteriophage Therapy. *J Pediatric Infect Dis Soc* 2018; 7:
253-256.

12 Khawaldeh A, Morales S, Dillon B, Alavidze Z, Ginn AN, Thomas L *et al*. Bacteriophage therapy for
refractory Pseudomonas aeruginosa urinary tract infection. *J Med Microbiol* 2011; 60: 1697-1700.

13 LaVergne S, Hamilton T, Biswas B, Kumaraswamy M, Schooley RT, Wooten D. Phage Therapy for
a Multidrug-Resistant Acinetobacter baumannii Craniectomy Site Infection. *Open Forum Infect Dis*
2018; 5: ofy064.

707
708 14 Schooley RT, Biswas B, Gill JJ, Hernandez-Morales A, Lancaster J, Lessor L *et al*. Development and
709 Use of Personalized Bacteriophage-Based Therapeutic Cocktails To Treat a Patient with a
710 Disseminated Resistant Acinetobacter baumannii Infection. *Antimicrob Agents Chemother* 2017;
711 61.

712
713 15 Wright A, Hawkins CH, Anggard EE, Harper DR. A controlled clinical trial of a therapeutic
714 bacteriophage preparation in chronic otitis due to antibiotic-resistant Pseudomonas aeruginosa;
715 a preliminary report of efficacy. *Clin Otolaryngol* 2009; 34: 349-357.

716
717 16 Chan BK, Turner PE, Kim S, Mojibian HR, Elefteriades JA, Narayan D. Phage treatment of an aortic
718 graft infected with Pseudomonas aeruginosa. *Evol Med Public Health* 2018; 2018: 60-66.

719
720 17 Dedrick RM, Guerrero-Bustamante CA, Garlena RA, Russell DA, Ford K, Harris K *et al*. Engineered
721 bacteriophages for treatment of a patient with a disseminated drug-resistant Mycobacterium
722 abscessus. *Nat Med* 2019; 25: 730-733.

723
724 18 Little JS, Dedrick RM, Freeman KG, Cristinziano M, Smith BE, Benson CA *et al*. Bacteriophage
725 treatment of disseminated cutaneous Mycobacterium chelonae infection. *Nat Commun* 2022; 13:
726 2313.

727
728 19 Nick JA, Dedrick RM, Gray AL, Vladar EK, Smith BE, Freeman KG *et al*. Host and pathogen response
729 to bacteriophage engineered against Mycobacterium abscessus lung infection. *Cell* 2022; 185:
730 1860-1874 e1812.

731
732 20 Guerrero-Bustamante CA, Dedrick RM, Garlena RA, Russell DA, Hatfull GF. Toward a Phage
733 Cocktail for Tuberculosis: Susceptibility and Tuberculocidal Action of Mycobacteriophages against
734 Diverse Mycobacterium tuberculosis Strains. *mBio* 2021; 12.

735
736 21 Gan Y, Wu T, Liu P, Guo S. Characterization and classification of Bo4 as a cluster G
737 mycobacteriophage that can infect and lyse M. tuberculosis. *Arch Microbiol* 2014; 196: 209-218.

738
739 22 Azimi T, Mosadegh M, Nasiri MJ, Sabour S, Karimaei S, Nasser A. Phage therapy as a renewed
740 therapeutic approach to mycobacterial infections: a comprehensive review. *Infect Drug Resist*
741 2019; 12: 2943-2959.

742
23 Pai M, Behr MA, Dowdy D, Dheda K, Divangahi M, Boehme CC *et al*. Tuberculosis. *Nat Rev Dis*
Primers 2016; 2: 16076.

24   Arrey F, Lowe D, Kuhlmann S, Kaiser P, Moura-Alves P, Krishnamoorthy G *et al*. Humanized Mouse
Model Mimicking Pathology of Human Tuberculosis for in vivo Evaluation of Drug Regimens. *Front
Immunol* 2019; 10: 89.

25   Jangalwe S, Shultz LD, Mathew A, Brehm MA. Improved B cell development in humanized NOD-
scid IL2Rgamma(null) mice transgenically expressing human stem cell factor, granulocyte-
macrophage colony-stimulating factor and interleukin-3. *Immun Inflamm Dis* 2016; 4: 427-440.

26   Coughlan AM, Harmon C, Whelan S, O'Brien EC, O'Reilly VP, Crotty P *et al*. Myeloid Engraftment
in Humanized Mice: Impact of Granulocyte-Colony Stimulating Factor Treatment and Transgenic
Mouse Strain. *Stem Cells Dev* 2016; 25: 530-541.

27   Calderon VE, Valbuena G, Goez Y, Judy BM, Huante MB, Sutjita P *et al*. A humanized mouse model
of tuberculosis. *PLoS One* 2013; 8: e63331.

28   Yu CI, Martinek J, Wu TC, Kim KI, George J, Ahmadzadeh E *et al*. Human KIT+ myeloid cells facilitate
visceral metastasis by melanoma. *J Exp Med* 2021; 218.

29   Billerbeck E, Barry WT, Mu K, Dorner M, Rice CM, Ploss A. Development of human CD4+FoxP3+
regulatory T cells in human stem cell factor-, granulocyte-macrophage colony-stimulating factor-
, and interleukin-3-expressing NOD-SCID IL2Rgamma(null) humanized mice. *Blood* 2011; 117:
3076-3086.

30   Janke LJ, Imai DM, Tillman H, Doty R, Hoenerhoff MJ, Xu JJ *et al*. Development of Mast Cell and
Eosinophil Hyperplasia and HLH/MAS-Like Disease in NSG-SGM3 Mice Receiving Human CD34+
Hematopoietic Stem Cells or Patient-Derived Leukemia Xenografts. *Vet Pathol* 2021; 58: 181-204.

31   Terahara K, Iwabuchi R, Tsunetsugu-Yokota Y. Perspectives on Non-BLT Humanized Mouse
Models for Studying HIV Pathogenesis and Therapy. *Viruses* 2021; 13.

32   Willinger T, Rongvaux A, Takizawa H, Yancopoulos GD, Valenzuela DM, Murphy AJ *et al*. Human
IL-3/GM-CSF knock-in mice support human alveolar macrophage development and human
immune responses in the lung. *Proc Natl Acad Sci U S A* 2011; 108: 2390-2395.

33   Cardona P, Cardona PJ. Regulatory T Cells in Mycobacterium tuberculosis Infection. *Front Immunol*
2019; 10: 2139.

34   Larson RP, Shafiani S, Urdahl KB. Foxp3(+) regulatory T cells in tuberculosis. *Adv Exp Med Biol*
2013; 783: 165-180.

35   Peng LL, Yongai; Chen, Baowen; Li, Youlun; Shen, Xiaobing; Su, Cheng and Wang, Guozhi.
Therapeutic effect of bacteriophage D29 in the treatment for guinea pigs infected with sensitive
strain of Mycobacterium tuberculosis. *Chinese Journal of Zoonoses* 2009; 25: 733-736.

36   Sula L, Sulova J, Stolcpartova M. Therapy of experimental tuberculosis in guinea pigs with
mycobacterial phages DS-6A, GR-21 T, My-327. *Czech Med* 1981; 4: 209-214.

37   Delattre R, Seurat J, Haddad F, Nguyen TT, Gaborieau B, Kane R *et al*. Combination of in vivo phage
therapy data with in silico model highlights key parameters for pneumonia treatment efficacy.
Cell Rep 2022; 39: 110825.

38   Brune K, Frank J, Schwingshackl A, Finigan J, Sidhaye VK. Pulmonary epithelial barrier function:
some new players and mechanisms. *Am J Physiol Lung Cell Mol Physiol* 2015; 308: L731-745.

39   Cerveny KE, DePaola A, Duckworth DH, Gulig PA. Phage therapy of local and systemic disease
caused by Vibrio vulnificus in iron-dextran-treated mice. *Infect Immun* 2002; 70: 6251-6262.

40   Chhibber S, Kaur S, Kumari S. Therapeutic potential of bacteriophage in treating Klebsiella
pneumoniae B5055-mediated lobar pneumonia in mice. *J Med Microbiol* 2008; 57: 1508-1513.

41   Feng Y, Kong Y, Barnes PF, Huang FF, Klucar P, Wang X *et al*. Exposure to cigarette smoke inhibits
the pulmonary T-cell response to influenza virus and Mycobacterium tuberculosis. *Infect Immun*
2011; 79: 229-237.

42   Tucker TA, Jeffers A, Alvarez A, Owens S, Koenig K, Quaid B *et al*. Plasminogen activator inhibitor-
1 deficiency augments visceral mesothelial organization, intrapleural coagulation, and lung
restriction in mice with carbon black/bleomycin-induced pleural injury. *Am J Respir Cell Mol Biol*
2014; 50: 316-327.