Title: Endothelial vacuole membrane enriched aquaporins regulate microvascular lumenization in development and hyperglycemia

Changsheng Chen[1, *], Yinyin Qin[1, 2, *], Yidan Xu[3, 4, *], Lixun Chen[3, 4], Linnong Wang[3, 4], Dong Liu[1, #]

[1] School of Life Sciences, Nantong Laboratory of Development and Diseases; Second Affiliated Hospital; Key Laboratory of Neuroregeneration of Jiangsu and Ministry of Education, Co-innovation Center of Neuroregeneration, Nantong University, Nantong, China

[2] Suzhou Science & Technology Town Hospital, Suzhou, China

[3] Department of Ophthalmology, Nanjing First Hospital, Nanjing Medical University, Nanjing, China.

[4] Nanjing Red Cross Eye Bank, Nanjing, China.

*, these authors contributed equally to this work

#, author for correspondence

Contact details for Correspondence:

Dong Liu , Ph.D

Nantong Laboratory of Development and Diseases, Nantong University

Qixiu Road 19, Nantong, China, 226001

Phone: + (86) - 18605133927

Fax: + (86)-513-85012809

Email: liudongtom@gmail.com; tom@ntu.edu.cn

Abstract

In patients with diabetic microvascular complications, reduced vessel perfusion or vascular occlusion is a common characteristic which will cause the insufficient blood supply. However, identification of novel regulators involved in microvascular lumenization defects is hindered by the lacking of a model for imaging the blood vessels at high resolution *in vivo*. Taking advantage of the transparency of zebrafish, we observed the reduction of vascular diameter and compromised perfusion in high glucose treated embryos. RNA sequencing and whole-mount *in situ* hybridization analysis indicated that two aquaporins (*aqp1a.1* and *aqp8a.1*) were significant down-regulated, which was further confirmed by endothelial specific Q-PCR. It was also shown that the two aqps were spatio-temporally enriched in the endothelial cells (ECs) of vascular system. Zebrafish with loss of *aqp1a.1* or *aqp8a.1* displayed lumenization defects in intersegmental vessels, recapitulating the phenotype in hyperglycemic zebrafish model. While overexpressing the aquaporins in zebrafish promoted the enlargement of the vascular diameter. Moreover, the defective vasculature induced by high-glucose treatment could be rescued by *aqp1a.1* upregulation. In addition, both *aqp1a.1* and *apq8a.1* were localized in the intracellular vacuoles in cultured ECs as well as in the ECs of sprouting ISVs, and loss of Aqps caused the reduction of those vacuoles, which was required for lumenization. Finally, we found that the expression of human *AQP1* was downregulated in diabetic human retina samples and high-glucose treated human retinal microvascular endothelial cells. All these results suggest that EC-enriched aquaporins have a role in developmental and pathological blood vessel lumenization, and they might be potential targets for gene therapy to cure diabetes-related vascular lumenization defects.

Key words: diabetic retinopathy; Aquaporin; lumenization; rescue

Introduction

Hyperglycemia caused microvascular complications, including hemorrhages, microaneurysms, capillary non-perfusion, and neovascularization (Agemy et al., 2015; Beckman and Creager, 2016; Duh et al., 2017; Toh et al., 2019; Wang and Lo, 2018), are the major source of morbidity and mortality in diabetes. However, it remains unclear how does the reduced perfusion of blood vessels form under hyperglycemic conditions. To address this, the zebrafish (*Danio Rerio*) model was utilized in our study, which offers superior properties in cardiovascular research (Gore et al., 2012; Liu and Stainier, 2012). The transparency of zebrafish embryos allows to perform high-resolution imaging analysis of vascular development *in vivo*. Importantly, the zebrafish can live up to the first week of development without a functional vasculature, allowing a detailed analysis even in animals with severe cardiovascular defects; by contrast, avian and mammalian embryos die rapidly in the absence of a functional cardiovascular system. Moreover, a variety of established zebrafish diabetes mellitus (DM) models are available and easy to manipulate (Chen and Liu, 2022).

Aquaporin (AQPs) are a superfamily of tetrameric membrane proteins that involve in transportation of water across cell plasma membranes (Papadopoulos et al., 2008; Smith and Agre, 1991). Several of these small, integral membrane proteins are also known for transfer of other small, uncharged solutes (e.g. glycerol), to maintain fluid homeostasis (Verkman, 2005). Accumulating evidences suggest that specific classes of AQPs are involved in angiogenesis and endothelial cell (EC) function under physiological and pathological conditions (Ribatti et al., 2014; Verkman, 2002). In current work, the zebrafish embryos were immersed into a high glucose solution to mimic a hyperglycemic state. Under the treatment, the narrowed and reduced perfusion of blood vessels in both eyes and trunks of zebrafish embryos were observed. It has been revealed that two endothelial enriched aquaporins, *aqp1a.1* and *aqp8a.1*, were significantly down-regulated upon high glucose treatment. These results suggest a role of aquaporins in both developmental and hyperglycemic blood vessels. Here, we found that *apq1a.1* or *aqp8a.1* loss-of-function led to the reduction of diameter and perfusion

of intersegmental vessels, which recapitulated the vascular phenotype of hyperglycemic zebrafish embryos. However, gain-of-function of *apq1a.1* or *aqp8a.1* in zebrafish embryos caused the increased diameter of blood vessels. Moreover, overexpression of *apq1a.1* in hyperglycemic zebrafish embryos rescued the defective vascular phenotype. In addition, gene expression analysis of retina samples from 49 non-diabetic and 54 diabetic donors showed that the human AQP1 (orthologous to zebrafish *aqp1a.1*) was also down-regulated under diabetic condition and high-glucose treated human retinal microvascular endothelial cells. All taken together, AQPs are important regulators in blood vessel lumenization during vascular system development and hyperglycemia.

Results

Hyperglycemia leads to vascular lumenization defect in zebrafish during embryogenesis

To examine the effects of high glucose on vascular development, a hyperglycemia zebrafish model was established using the transgenic line *Tg(fli1ep:EGFP-CAAX)^{ntu666}*, in which an endothelial enhancer/promoter fragment (fli1ep) was employed to drive the specific expression of the enhanced green fluorescent protein (EGFP) in ECs, and the CAAX membrane targeting motif was able to guarantee the localization of EGFP at the surface of ECs (Fig.S1). In this line, a direct non-invasive visualization of the zebrafish vasculature can be achieved under a conventional fluorescence microscopy or a confocal microscopy. A hyperglycemic state in zebrafish embryos was achieved by immersing them into a high glucose solution from 2 days post fertilization (dpf), and fluorescence imaging of the vasculature was performed at 3 dpf (Fig.1A and S2). The high glucose treatment did not yield phenotypically differences and developmental defects on zebrafish embryos, but the vasculature was substantially influenced. Under this condition, the vessel diameters were significantly decreased compared with the untreated control counterparts, including the inner optic circle (IOC) in the eyes as well as the intersegmental vessels (ISVs), dorsal aorta (DA) and posterior cardinal vein (PCV) in the trunk (Fig.1B-C'' and Fig.1E-F''). The IOC in zebrafish eyes and the ISVs in zebrafish trunk were declined by 27.9% and 36.8%, respectively (Fig.1D and H). Also,

microangiography analysis revealed that 59% of the narrowed ISVs were not perfused based on a (Fig.1G), suggesting that the dysfunctional vascular lumens were formed after high glucose induction.

Figure 1. High glucose treatment affects vascular lumenization in zebrafish embryos. A, Schematic representation of establishing zebrafish hyperglycemia/DM model. Zebrafish embryos are immersed into high glucose solution from 2 to 3 dpf. Microangiography and confocal imaging are performed afterwards. **B-C'',** Representative confocal images of IOC phenotypes in 3-dpf *Tg(fli1ep:EGFP-CAAX)^ntu666* transgenic line with or without glucose treatment. White arrowheads indicate the IOC. **D,** Quantification of IOC diameter. **E-F'',** Representative confocal images of ISV phenotypes in 3-dpf *Tg(fli1ep:EGFP-CAAX)^ntu666* transgenic line with or without glucose treatment. White arrowheads indicate the ISV. **G,** The incidence of normal and non-perfused ISVs in 3-dpf *Tg(fli1ep:EGFP-CAAX)^ntu666* transgenic line with or without glucose treatment. **H,** Quantification of ISV diameter. Data are shown as mean ± s.e.m. A two-tailed, unpaired Student's t-test is applied. ***, $p < 0.001$. ****, $p < 0.0001$.

High glucose treatment caused the downregulation of expression of *aqp1a.1* and *aqp8a.1*

To further explore which genes are altered upon glucose treatment, transcriptome sequencing analysis was performed. Compared to those in control zebrafish, the expression of a serials of genes were significantly altered in 72 and 96 hours post fertilization (hpf) embryos treated with high glucose, with 642 genes upregulated and 1,149 genes downregulated at 72 hpf, as well as 1,259 genes upregulated and 1,851 genes downregulated at 96 hpf (Fig.2A and B). Approximately 30% of the upregulated genes and 40% of the downregulated genes were overlapped between 72 and 96 hpf (Fig.2C). KEGG analysis revealed that a variety of biological processes or signaling pathways related to glucose and lipid metabolism, immune responses and vascular regulation, were enriched in embryos under hyperglycemic condition (Fig.2D). Several EC-enriched genes, like *cxcr4a* and *esm1*, were found to be upregulated after high glucose treatment, while the aquaporins (*aqp1a.1* and *aqp8a.1*) exhibited reduced expressions (Fig.2E and F). Further whole-mount ISH assay demonstrated that the zebrafish *aqp1a.1* and *aqp8a.1* were highly enriched in ECs of vascular system (Fig.S3). Additionally, the expressions of *aqp1a.1* and *aqp8a.1* in hyperglycemic zebrafishes were validated by real-time qPCR and ISH assay, and the results were consistent with the transcriptomic data (Fig.3A-H). To further examine whether *aqp1a.1* and *aqp8a.1*

were down-regulated in ECs of high glucose treated embryos, we performed qPCR
assay with isolated ECs. The ECs in control and high glucose-treated group at 96 hpf
were respectively sorted by FACs and subsequently used for RNA isolation (Fig.3I).
Upon high glucose induction, both *aqp1a.1* and *aqp8a.1* in ECs were significantly
downregulated (Fig.3J and K).

**Figure 2. Transcriptomic profiling of *Tg(fli1ep:EGFP-CAAX)^(ntu666)* transgenic line with or
without glucose treatment. A and B,** Volcano plots of downregulated genes (blue dots),
upregulated genes (red dots), and unchanged genes (grey dots) in high glucose-treated group
versus the control group at 72 and 96 hpf, respectively. **C,** Venn diagram shows the overlapping
upregulated and downregulated DEGs upon high glucose treatment at 72 and 96 hpf. **D,** KEGG

enrichment analysis reveals the signaling pathways and the cellular and biological processes that the DEGs are involved. **E and F,** Heatmaps based on RNA-seq data demonstrate that *aqp1a.1* and *aqp8a.1* are downregulated upon high glucose treatment.

Figure 3. Zebrafish *aqp1a.1* and *aqp8a.1* are differentially expressed upon high glucose treatment. A-H, Expressions of *aqp1a.1* and *aqp8a.1* in zebrafish after high glucose treatment are examined by Q-PCR and ISH. **I,** Schematic representation of isolation of ECs from control and high glucose-treated *Tg(kdrl:EGFP)* zebrafish embryos for RNA extraction and Q-PCR. **J and K,** Quantification of *aqp1a.1* and *aqp8a.1* expressions in isolated ECs. Data are shown as mean ± s.e.m. A two-tailed, unpaired Student's t-test is applied. **, p < 0.01.

aqp1a.1 or *aqp8a.1* are essential for blood vessel lumenization

To gain an insight into the functions of EC-expressed AQPs in vascular system, we performed gene functional analysis using zebrafish model. We first used CRISPR/Cas9 genome editing method to knockout either *aqp1a.1* or *aqp8a.1* in *Tg(fli1ep:EGFP-*

CAAX)^{ntu666} transgenic zebrafish (Fig.S4 and S5). Homozygous *aqp1a.1*-knockout

mutants displayed reduced lumen size of the small intersegmental vessels (ISVs),

which were generated from angiogenesis. However, the diameters of large vessels,

including dorsal aortae (DA) and posterior cardinal vein (PCV), which are generated

through vasculogenesis, were not significantly affected (Fig.4A). Upon *apq1a.1* gene

deletion, the diameters of DV and PCV only decreased by 4.2% and 4.7% at 48 hpf,

respectively, whereas the average diameter of ISVs showed an approximately 20%

decrease (Fig.4B). To examine whether the vascular defects were caused by off-target

effects of the CRISPR system, we knocked down *aqp1a.1* expression by injecting the

aqp1a.1 splicing morpholino into *Tg(fli1ep:EGFP-CAAX)^{ntu666}* embryos. The vascular

defects were recapitulated and even enhanced within *aqp1a.1*-MO-injected zebrafish

embryos (Fig.S6). In addition, the injection of *aqp1a.1* mRNA rescued the vascular

defects in *aqp1a.1* morphants. These results suggested that Aqp1a.1 exerted a role in

regulation of vascular lumenization. However, the homozygous or even heterozygous

aqp8a.1-knockout mutants have not been obtained since all harvested embryos were

genotyped as multiples of 3 base pairs deletion (Fig.S5). We were of the opinion that

targeted deletion of the *aqp8a.1* gene resulted in embryonic lethality in zebrafish.

Although it was hardly to get *aqp8a.1*-knokout mutant, the G0 generation chimeras,

which contained the *aqp8a.1*-knockout cells, also displayed narrowed vessel lumens

(Fig.4A). Compared with those of control, the diameters of DA, PCV and ISVs in

aqp8a.1-KO G0 mutants decreased 11.2%, 9.8% and 13.3%, respectively (Fig. 4B). In

order to test whether the vascular phenotypic change was result from *aqp8a.1* loss of

function, we injected *aqp8a.1*-specific morpholino into *Tg(fli1ep:EGFP-CAAX)^{ntu666}*

embryos to block *aqp8a.1* pre-RNA splicing. The morpholino-mediated *aqp8a.1*

knockdown led to reduced lumen diameters of blood vessels, which mimicked the

phenotype caused by *aqp1a.1* loss-of-function (Fig.S6). Also, non-perfused or

perfusion-defective ISVs were more frequently observed in *aqp1a.1* or *aqp8a.1*

knockout embryos based on microangiography analysis (Fig.4C and D). All the results

suggested that loss-of-function of either *aqp1a.1* or *aqp8a.1* impaired the vascular

lumen formation during blood vessel development.

Figure 4. Knockout of EC-enriched aquaporins (*aqp1a.1* or *aqp8a.1*) leads to defective vascular lumenization in zebrafish. A, Confocal images of ISV phenotypes in 48-hpf *Tg(fli1ep:EGFP-CAAX)*[ntu666] control embryos and *aqp1a.1*-knockout and *aqp8a.1*-knockout embryos. Red and blue square brackets indicate artery and vein, respectively. Magenta, red and blue arrowheads in the representative sketch indicate the positions for vessel diameter measurement for ISV, DA and PCV, respectively. **B,** Quantification of the diameters of ISV, DA and PCV. Data are shown as mean ± s.e.m. One-way ANOVA analysis is applied. **C,** Microangiography of the trunk region in 72-hpf *Tg(fli1ep:EGFP-CAAX)*[ntu666] control embryos and *aqp1a.1*-knockout and *aqp8a.1*-knockout embryos. Blue arrowheads indicate the perfusion-deficient ISVs. **D,** The incidence of normal and defective perfused ISVs in 72-hpf

Tg(fli1ep:EGFP-CAAX)^{ntu666} control embryos and *aqp1a.1*-knockout and *aqp8a.1*-knockout

embryos. ns, not significant. *, p<0.05. **, p < 0.01. ****, p < 0.0001.

Next, we performed Aqp1a.1/Aqp8a.1 gain of function analysis to further confirm the

contribution of EC-expressed aquaporins on vascular lumenization. *Tg(fli1ep:aqp1a.1-*

mCherry)^{ntu667} and *Tg(fli1ep:aqp8a.1-EGFP)^{ntu668}* lines were constructed to overexpress

fluorescently-labeled Aqp1a.1 and Aqp8a.1 in zebrafish, respectively. Overexpressing

either of them induced the enlargement of the blood vessel lumens (Fig.5A).

Overexpression of *aqp1a.1* led to increases in DA and PCV diameter by 15.2% and

17.3%, respectively (Fig.5B and C), while the diameter of ISVs increased almost half of

the control (approx. 47%) upon *aqp1a.1* overexpression (Fig.5D). The morphology of

blood vessels in *aqp8a.1* gain of function embryos resembled the vascular phenotype of

aqp1a.1 overexpression (Fig.5A). Diameters of DA, PCV, and ISVs increased 22.4%,

23.2% and 32.2% in *aqp8a.1* overexpressed zebrafish embryos, respectively (Fig.5B-D).

Moreover, co-overexpression of *aqp1a.1* and *aqp8a.1* in zebrafish endothelial cells

resulted in further dilation of DA, but the diameters of PCV and ISVs did not present

significant increase compare to single aquaporin (*aqp1a.1* or *aqp8a.1*) overexpression

(Fig.5A-D). The diameters of DA, PCV and ISVs in *aqp1a.1* and *aqp8a.1* co-

overexpressed line exhibited 30.9%, 19.8% and 39.0% elevation compared to control

(Fig.5A-D). These results suggested a role of Aqp1a.1 and Aqp8a.1 in promoting

vascular lumen formation.

Figure 5. Overexpression of EC-enriched aquaporins (*aqp1a.1* or *aqp8a.1*) in ECs results in vessel lumen enlargement in zebrafish. A, Confocal images of ISV phenotypes in 48-hpf *Tg(fli1ep:EGFP-CAAX)^{ntu666}* transgenic line (control), *Tg(fli1ep:aqp1a.1-mCherry)^{ntu667}* transgenic line (*aqp1a.1* overexpression), *Tg(fli1ep:aqp8a.1-EGFP)^{ntu668}* transgenic line (*aqp8a.1* overexpression), and *Tg(fli1ep:aqp8a.1-EGFP::fli1ep:aqp1a.1-mCherry)* double transgenic line (*aqp1a.1* and *aqp8a.1* co-overexpression). Red and blue square brackets indicate artery and vein, respectively. **B-D,** Quantification of the diameters of ISV, DA and PCV. Data are shown as mean ± s.e.m. One-way ANOVA analysis is applied. **, $p < 0.01$. ****, $p < 0.0001$.

In addition, we applied $HgCl_2$ treatment on *Tg(fli1ep:EGFP-CAAX)^{ntu666}* zebrafish embryo to inhibit aquaporin function and check its effect on the vascular lumenization. Mercury has been commonly used to specifically block the water-channel activity of Aqp1a.1 and Aqp8a.1(Khan et al., 2013; Liu et al., 2006, p. 8; Preston et al., 1992; Yang et al., 2006). To screen the proper dosage for the treatment, we tried varied doses of $HgCl_2$. It was shown that the embryos exposed to 1.6 μM $HgCl_2$ from 22 to 48 hpf had no obvious morphological defects. Here, lower concentration (0.4 μM) of $HgCl_2$ was

sufficient to narrow the diameter of the blood vessels (Fig.S7). Under this concentration,
there were 15.4%, 15.7%, and 16.9% decreases in the diameters of DV, PCV, and ISVs,
respectively. With further increase in $HgCl_2$ concentration, the diameters of all the
aforementioned blood vessels reduced more remarkably (Fig.S7). As a result, blocking
the channel activity of aquaporins led to defective vascular tube formation.

Overexpression of *aqp1a.1* rescues hyperglycemia-induced vascular
lumenization defect

To elucidate whether the hyperglycemia-induced vascular lumenization defect is AQP
dependent, we establish a transgenic construct, in which *apq1a.1* was specifically
overexpressed in ECs. As shown in Fig.6A, the *pTol2-fli1ep-aqp1a.1-mCherry* plasmid
and Tol2 transposase mRNA were co-injected into 4-cell stage embryos of
Tg(fli1ep:EGFP-CAAX)^{ntu666} progeny. By using this strategy, the mCherry-labeled
Aqp1a.1 was mosaically expressed in the zebrafish vascular system. After injection, the
embryos were first selected for mosaic mCherry fluorescence and subsequently
immersed into glucose solution from 48 hpf for 24 h prior to confocal imaging analysis
(Fig.6B). As expected, a few ISVs exhibited both mCherry and GFP expression, and
these ISVs displayed wider lumens than the ones without mCherry expression (Fig.6C
and D). These results demonstrated that overexpression of *aqp1a.1* recovered the
defective lumen formation caused by hyperglycemia.

Figure 6. Overexpression of zebrafish *aqp1a.1* rescues hyperglycemia-caused defective lumenization. A, Schematic representation of establishment of *aqp1a.1*-overexpressed transgenic line under *Tg(fli1ep:EGFP-CAAX)*[ntu666] background. DNA construct and Tol2 transposase mRNA are co-injected at 4-cell stage to establish a chimera line, in which the mCherry-labeled Aqp1a.1 is mosaically expressed in vascular system. **B,** Embryos are selected and immersed into glucose solution from 2 to 3 dpf prior to imaging. **C,** Representative confocal images of ISVs in 3-dpf *Tg(fli1ep:EGFP-CAAX::fli1ep:apq1.1-mCherry)* double transgenic line. **D,** Quantification of the lumen diameters in ISVs with or without Aqp1a.1 expression. Data are shown as mean ± s.e.m. A two-tailed, unpaired Student's t-test is applied. ****, p < 0.0001.

AQPs contribute to vascular lumenization through intracellular vacuoles

Given that intracellular vacuoles fusion is an essential way of vascular lumenization in zebrafish ISVs (Kamei et al., 2006), whether AQPs involves in this process is undetermined. By expressing the fluorescently-labeled *aqp1a.1* and *aqp8a.1* in human umbilical vein endothelial cells (HUVECs), we found that both Aqps were localized in the endothelial cell membrane and highly enriched in vacuoles membrane subcellularly (Fig.7A and B). Moreover, Aqp1a.1 was co-localized with Cdc42, a marker for intracytoplasmic vesicles (Kamei et al., 2006; Osmani et al., 2010), in sprouting ISVs (Fig.7C). Meanwhile, loss-of-function of *aqp1a.1* and *aqp8a.1* led to the reduction of intracellular vacuoles and defective endothelial tube assembly in ISVs (Fig.7D). In addition, the area of endothelial vacuoles was significantly declined upon loss of Aqps in endothelial cells of ISVs (Fig.7E and F). These results indicated that *aqp1a.1* and *aqp8a.1* displayed cytoplasmic vacuolar localization and required for the intracellular vacuole-mediated endothelial tube formation.

Figure 7. Aqp1a.1 and Aqp8a.1 localize in intracellular vacuoles of ECs *in vitro* and *in vivo*. A and B, Fluorescently-labeled Aqps are expressed on the surface of cytoplasmic vacuoles (blue arrowheads) in HUVECs. **C,** Confocal imaging of zebrafish ISVs in *Tg(fli1:EGFP-cdc42wt::fli1a:aqp1a.1-mCherry)* double transgenic line, in which the EGFP-labeled Cdc42 protein marks the vacuolar structures (white arrowheads) and Aqp1a.1 is labeled by mCherry. Aqp1a.1 co-localizes with intracellular vacuoles (dashed circles). Scale bars, 10 µm. **D,** Knockout aqp1a.1 and knockdown aqp8a.1 in zebrafish leads to defective vascular lumen formation in ISVs. **E,** Schematic representation of intracellular vacuole change upon loss of Aqp. **F,** Quantification of the area of intracellular vacuoles in zebrafish ISVs. Data are shown as mean ± s.e.m. A two-tailed, unpaired Student's t-test is applied. **, $p < 0.01$.

Expression of *AQP1* in human retina specimens and cultured HRMECs

To examine whether the AQP1 expression is correlated with diabetes in patients, we measured its expression in diabetic and non-diabetic retina samples. In this study, a total of 103 donors (n=54 in diabetic group and n=49 in non-diabetic group) were enrolled. Fig.8A showed the representative fundus images and fluorescein angiograms of non-diabetic retinas and diabetic subjects with different grades of DR, starting from mild non-proliferative diabetic retinopathy (NPDR) to severe proliferative diabetic retinopathy (PDR). The mean diameters of central retinal vessels (central retinal artery and vein) and the retinal vessel branches (retinal arterioles and venules) were measured (Fig.8B). There was no difference in the central retinal vessel between diabetic and non-diabetic retina samples, whereas the diameter of the retinal vessel branches was significantly decreased in diabetic retinas (Fig.8C and D). Next, the mRNAs of both groups were isolated and applied to quantitative-PCR. The statistical results demonstrated that the relative expression of *AQP1* was significantly lower in diabetic samples than in non-diabetic samples (Fig.8E). The percentages of donors with downregulated *AQP1* were 77.8% (42/54) and 42.9% (21/49) in diabetic and non-diabetic groups, respectively (Table 1). The *AQP1* expression was negatively correlated with diabetes (P=0.0003). Moreover, there was no difference between the groups in terms of age and sex (P=0.3983 and P=0.3439, respectively) (Table 2). In the diabetic group, 19 patients were diagnosed with NPDR and 35 patients were diagnosed with

PDR. However, there was no *AQP1* expression difference between NPDR patients (16

samples showed down-regulated *AQP1* expression) and PDR patients (26 samples

showed down-regulated *AQP1* expression).

Figure 8. *AQP1* expression is downregulated in retina samples from diabetic patients. A,

Representative fundus images (upper) and fluorescein angiography (lower) of normal retinas

and retina with diabetic retinopathy (DR) at different stages (from mild to severe). **B,**

Representative sketch of human retinal vessels. **C,** Diameters of the central retinal vessels

(central retinal artery and vein) and the branching retinal vessels (retinal arterioles and venules).

Data are shown as mean ± s.e.m. A two-tailed, unpaired Student's t-test is applied. ns, not

significant. ****, p < 0.0001. **D,** Quantification of *AQP1* expression in healthy (n=49) and diabetic (n=54) retina samples. Data are shown as mean ± s.e.m. A two-tailed, unpaired Student's t-test is applied. ***, p < 0.001. **E,** Quantification of *AQP1* expression in HRMECs cultured in normal and high glucose medium. One-way ANOVA analysis is applied. ****, p < 0.0001.

The expression of *AQP1* in endothelial cells under high glucose condition was further assessed in cultured human retinal microvascular endothelial cells (HRMECs) (Fig.8F). Compared with cells cultured in complete classic medium, high glucose-treated cells displayed a lower *AQP1* expression (Fig.8G), consisting with our previous results.

Table 1 Differential expression of *AQP1* mRNA in non-diabetic and diabetic retinas

Type	number	Relative expression		Negative	P
		<0.5[a]	>0.5		
Retinas from non-diabetic donors (Healthy subjects)	49	21	28	42.9%	0.0003
Retinas from diabetic donors (Diabetic subjects)	54	42	12	77.8%	χ^2=13.190

[a]Two-folds downregulated.

Table 2 Correlation of *AQP1* expression with age and sex

Group		number	Relative expression		Negative	P
			<0.5[a]	>0.5		
Age	≤64[b]	50	28	22	56.0%	0.3983
	>64	53	34	19	64.1%	χ^2=0.7143
Sex	Male	70	45	25	64.3%	0.3439
	Female	33	18	15	54.5%	χ^2=0.8958

[a]Two-folds downregulated; [b]Mean age.

Discussion

Diabetic retinopathy (DR) is a common vascular complication of diabetes and also a leading cause of vision loss among working-age populations (Forbes and Cooper, 2013).

It is often accompanied by a series of retinal vascular lesions, including reduced vessel
perfusion, damaged vasculature, increased vascular permeability, and
neovascularization (Duh et al., 2017). Hyperglycemia is an inextricable reason for DR,
and the diameter changes in retinal vessels has been recognized as a clinical sign of
DR (Klein et al., 2012; Wong, 2011). In our study, we observed a significant
downregulation of *AQP1* in retinas from patients with diabetes and high glucose treated
HRMECs. These observations indicate a novel pathophysiological link between AQP1
and DR. Aquaporins (AQPs) belongs to a superfamily of major intrinsic proteins (MIPs)
and serves as channel proteins to transport water and/or small solutes across plasma
membrane (Papadopoulos et al., 2008; Reizer et al., 1993; Verkman, 2005). Based on
their abilities of transporting water or other solutes, such as glycerol, the aquaporin
superfamily can be divided into two subfamilies: aquaporins and aquaglyceroprins.
However, whether AQPs play roles in vascular luminization and pathological vascular
complications progression are poorly understood. Discoveries in vascular research have
been limited by a shortage of a model for imaging the blood vessels at high resolution *in*
vivo. To address this, the zebrafish model was utilized in our study, which offers
superior properties in cardiovascular research. In zebrafish, 14 of mammalian AQP
homologues have been cloned and their expression have been examined with WISH
(Qin et al., 2019). The unique spatio-temporal distribution of individual aquaporins
during zebrafish embryogenesis implied their distinct roles in the embryonic
development. Although aquaporins, particularly AQP1, have been reported to be
strongly expressed in ECs and proliferating tumor microvessels in humans (Nico and
Ribatti, 2010; Ribatti et al., 2014; Verkman, 2002; Verkman et al., 2008), their functions
in vascular system remain to be elucidated. Among all these aquaporin homologues,
zebrafish *aqp1a.1* and *aqp8a.1* were endothelium-enriched. To further investigate the
effects of EC-expressed Aqp1a.1 and Aqp8a.1 on angiogenesis during zebrafish
development, we examined the vascular phenotypic changes upon *aqp1a.1*/*aqp8a.1*
loss-of-function and gain-of-function. Either *aqp1a.1* or *aqp8a.1* disruption in zebrafish
embryos narrowed dorsal aorta, cardinal vein and ISVs, which were more sensitive to
AQPs loss of function. In contrast, overexpressing *aqp1a.1* or *aqp8a.1* leaded to
increases in vascular diameters. Moreover, blockage the function of AQPs by mercury

resulted in the luminization defect of blood vessel as well. Taken together, these
observations suggested that AQPs modulated the blood vessel lumenization during
both vasculogenesis and angiogenesis. In addition, both aqp1a.1 and apq8a.1 were
localized in the intracellular vacuoles in cultured ECs as well as the ECs of sprouting
ISVs, and loss of Aqps caused the reduction of those vacuoles. Since zebrafish ISVs
adopted the intracellular vacuoles fusion as the main strategy for vascular lumenization,
the -enriched Aqps might play a critical role in this biological process, supporting two
previous notions that the vascular lumen formation was driven by intercellular fusion of
endothelial vacuoles (Kamei et al., 2006) and AQP was expressed on cytoplasmic
vesicles and required for lumen extension (Khan et al., 2013).

Zebrafish has been widely used in diabetic studies (Jung et al., 2016; Lee and Yang,
2021; Wiggenhauser et al., 2020), and a hyperglycemic state and a series of diabetic
vascular complications can be easily simulated through immersing the zebrafish
embryos or adults in high concentrated glucose solution (Gleeson et al., 2007; Liang et
al., 2010). In our data, high glucose treatment led to the vascular defects which were
well matched with the luminization phenotype in *aqp1a.1/aqp8a.1* loss-of-function.
Under hyperglycemic condition, the IOC in eyes and ISVs were narrowed significantly.
Furthermore, we discovered that zebrafish *aqp1a.1* and *aqp8a.1* were downregulated
upon high glucose exposure, which was consistent with our observation that *AQP1* was
downregulated in human diabetic retinas and high glucose treated HRMECs. Most
importantly, overexpressing *aqp1a.1*, orthologous to human *AQP1*, in ECs of zebrafish
DM model could rescue the vascular defects. All these results were in support of the
viewpoint that EC-expressed aquaporins are correlated with hyperglycemia-related
vascular complications. Sada et al. have ever mentioned that overexpression of
AQP1 attenuates the hyperglycemia-induced vascular complications (Sada et al., 2016,
p. 1). In their proposed working model, hyperglycemia leads to high oxygen
consumption which promotes mitochondrial ROS production and induces cellular
hypoxia. Afterwards, the mitochondrial ROS and cellular hypoxia promote
the overproduction of endothelin-1 and fibronectin, and induce the apoptosis, which

further leads to diabetic vascular complications. Based on our data, the attenuated
hypoxia might due to improvement of the vascular perfusion induced by the
overexpression of AQP1. However, overexpression of AQP1 is able to suppress the
hypoxic state through functioning as the oxygen channel in ECs (Echevarría et al.,
2007), and then prevent the aforementioned hyperglycemia-induced vascular damages.
In addition to AQP1, other non-EC-expressed aquaporins are also associated with
diabetes. Zhang et al. have found that AQP11 was down-regulated in DR and resulted
in intracellular edema in DR subsequently (Zhang et al., 2022). AQP2 was also reported
to be involved in diabetic kidney diseases (Hadidi et al., 2022; Hinrichs et al., 2022).
However, to our knowledge, it was the first time that AQP1 was reported to be involved
in physiological as well as pathological vascular lumen formation.

Our investigations on retina samples from diabetic patients and zebrafish DM model
emphasized that the EC-enriched aquaporins are associated with vascular lumenization
defect in hyperglycemia. Functional inhibition of aquaporins recapitulated vascular
alterations of high-glucose treated zebrafish model, and overexpression of aquaporin in
ECs rescued the defect. In summary, current study suggests that EC-enriched
aquaporins have a role in developmental and pathological blood vessel lumenization,
and they might be potential targets for gene therapy to cure diabetes-related vascular
lumenization defects.

Materials and Methods

Study approval

Experiments involving human samples were approved by Affiliated Hospital of Nantong
University (2020-K013). Retina samples from 103 donors were obtained from Nanjing
Red Cross Eye Bank (Nanjing, China) between 2019 and 2021, and informed written
consent was obtained from each individual. The 49 non-diabetic retina donors included
31 males and 18 females (average age, 62; age range, 23-85 years), and the 54
diabetic donors included 39 males and 15 females (average age, 66; age range, 37-88
years). All animal experimentation was carried out in accordance with NIH Guidelines
for care and use of laboratory animals (http://oacu.od.nih.gov/regs/index.htm) and

ethically approved by the Administration Committee of Experimental Animals, Jiangsu
Province, China (Approval ID: SYXK(SU) 2017–111). Best efforts were made to
minimize the number of animals used and prevent their suffering.

Zebrafish husbandry and strains

The wild-type AB line and transgenic lines *Tg(kdrl:EGFP)* (Wang et al., 2016),
Tg(fli1ep:EGFP-CAAX)^{ntu666}, *Tg(fli1ep:aqp1a.1-mCherry)^{ntu667}*, *Tg(fli1ep:aqp8a.1-*
EGFP)^{ntu668}, and *Tg(fli1:EGFP–cdc42wt)* (Kamei et al., 2006) were used in this study.
All zebrafish embryos and adult fishes were raised and maintained at 28.5 ºC.

Construction of transgenic lines

Tg(fli1ep:EGFP-CAAX)^{ntu666} transgenic line was generated by injecting *pTol2-fli1ep-*
EGFP-CAAX into the progeny of wild-type AB line together with Tol2 transposase
mRNA into 1-cell stage wildtype fertilized eggs (1 ng / embryo) according to the Tol2kit
protocol (Kwan et al., 2007). *Tg(fli1ep:aqp1a.1-mCherry)^{ntu667}* and *Tg(fli1ep:aqp8a.1-*
mCherry)^{ntu668} transgenic lines were generated by using the same strategy (Fig.S6).

High glucose treatment

The zebrafish embryos were incubated at high glucose solution for 24 hours from 48 hpf
and imaged at 72 hpf. The embryos incubated in E3 medium were used as control. To
measure the glucose level in zebrafish embryos after glucose treatment, the embryos
were first sacrificed with rapid chilling (submerge fish embryos into 2-4 ºC chilled water).
Following confirmation of the absence of heart beat under a using a stereo microscope,
the embryos were rinsed with E3 medium twice and then dried with paper towel and
placed into 2-ml microcentrifuge tube. One-hundred embryos were collected for each
time point (48, 60, 72, 84, and 96 hpf, respectively) and divided into 5 groups (5
biological replicates). The embryos were homogenized and centrifuged to collect the
supernatant. Two microliter of the supernatant was placed on a glucometer strip (One-

Touch Ultra 2) and the glucose level was measured with a glucometer (One-Touch Ultra

2).

ECs sorting, RNA extraction, RNA-seq, and qPCR

To isolate ECs, the control and high-glucose treated *Tg(kdrl:EGFP)* embryos (300-400

embryos for each) were collected at 96 hpf, and followed by washing with PBC for three

times and digested with 0.25% trypsin at 37˚C. The digested cells were collected by

centrifugation and allowed to pass through a 40 mm FACS tube (BD Falcon, 352340).

The EGFP positive cells were sorted by fluorescence-activated cell sorting on FACS

Aria3 (BD Biosciences). Total mRNAs of the retina samples, zebrafish embryos (30

embryos a pool), or ECs were extracted with TRIzol Reagent (Invitrogen, USA) and

quantified and qualified by NanoDrop (ThermoFisher Scientific, USA) and 1% agarose

gel. To study gene expression changes after high glucose treatment, RNA-seq was

performed with the prepared mRNAs from 72 and 96 hpf-control and high glucose-

treated zebrafish embryos. Three independent replicates of the samples were analyzed

for each treatment. All RNA samples were submitted to GENEWIZ Science (Suzhou,

China), and deep sequencings were performed on an Illumina Hiseq2500. Quantitative

RT-PCR was performed using the corresponding primers listed in Supplementary Table

S1.

Whole-mount in situ hybridization (WISH)

WISH with antisense RNA probes was performed as described previously (Krueger et

al., 2011). Templates for making probe to detect the expression of aqp1a.1

(NM_207059.1) and aqp8a.1 (NM_001004661.1), was cloned from cDNA fragments.

Primers for WISH were listed in Supplementary Table S1. After hybridization, images of

the embryos were acquired with an Olympus stereomicroscope MVX10 equipped with

an Olympus DP71 camera.

Generation of *aqp1a.1* or *aqp8a.1* knockout mutants

To generate *aqp1a.1* or *aqp8a.1* gene mutant zebrafish, we used a CRISPR/Cas9-mediated approach. CRISPR/Cas9 target sites designed to identify the sequences in the second exon of *aqp1a.1* or *aqp8a.1*. Primers for single-RNA (sgRNA) synthesis were listed in Supplementary Table S1. Each sgDNA were individually subcloned into pT7 vector and transcribed into sgRNA in vitro using the T7 mMessage mMachine kit (Ambion). Microinjection was performed with 1-cell stage zebrafish embryos and each embryo was co-injected with 100 pg sgRNA and 200 pg Cas9 mRNA. G0 generations were examined by PCR and subsequently Sanger sequencing. A mutant with 7 bp deletion at the second exon of *aqp1a.1* which leads to the occurrence of a stop codon at downstream of the Cas9 cutting site was identified. The heterozygous *aqp1a.1*$^{+/-}$ mutants were incrossed to obtain the homozygous F2 progenies.

Morpholino-mediated gene knockdown of *aqp1a.1* or *aqp8a.1* in zebrafish

The gene-specific morpholinos (Gene Tools, LLC) were used to block the splicing of the pre-mRNA. Morpholino antisense oligomers (refer to as MO) were prepared according to the manufacturer's instruction. The MO sequences used were listed Supplementary Table S1. In this experiment, 2 nL of 0.3 mM morpholino oligo was microinjected into the embryos at the 1-cell stage.

HgCl$_2$ treatment

Functional inhibition of Aqp1a.1 and Aqp8a.1 using HgCl2 was performed with dechorionated embryos at 22 hpf in E3 medium. Three different HgCl2 concentrations (0.2, 0.4, and 1.6 μM) were used. The embryos were incubated in the HgCl2-containing E3 medium from 22 hpf and imaged at 48 hpf.

Microangiography and confocal imaging

For confocal imaging of blood vessel development, zebrafish embryos treated with 1-
phenyl-2-thiourea (Sigma Aldrich) to inhibit the pigmentation. After manually
dechorionation, embryos were anesthetized with egg water/0.16 mg/ml tricaine/1% 1-
phenyl-2-thiourea (Sigma) and embedded in 0.6% low melting agarose. For
microangiography, 5 μg/μl Dextran TexasRed (Invitrogen) was injected into the common
cardinal vein (CCV) at 72 hpf and immediately processed for optical imaging. Living
imaging was performed with Nikon A1R confocal microscopy.

Quantitative analysis of blood vessel diameter

The vessel diameters were quantified in Fiji using the Vessel Analysis plugin (Schindelin
et al., 2012). For zebrafish, 5 locations on IOC, DA and PCV were selected and
measured. For zebrafish ISV diameter measurement, 5 ISVs beyond the yolk yac
extension were selected and an individual ISV was measured at 5 positions. A total of 5
fishes were applied in each group. The average of these data points was statistically
used as the diameter of the analyzed vessel. For human retinal vessels, measurements
were carried out on fluorescein angiograms. The mean diameters of the central retinal
vessels in the optic disk and the branching retinal vessels were separately calculated.
The vascular perfusion analysis in zebrafish was based on the microangiographic
images. Completely or partially lost the red fluorescence in ISV was defined as
defective-perfused ISV.

Cell culture and transfection

Human umbilical vein endothelial cells (HUVECs, Cell System) were cultured in in
endothelial cell growth medium according to the protocol provided by the manufacturer
(VascuLife, Cell System). Lipofectamine™ 3000 Transfection Reagent (ThermoFisher
Scientific) was used for plasmid transfection. The *aqp1a.1* or *aqp8a.1* cDNA was
subcloned into pmCherry-N1 or pEGFP-N1 vector (Addgene). Human retinal
microvascular endothelial cells (HRMECs, Cell System) were cultured in complete
classic medium supplemented with/without high content glucose (50 or 100 mM). Cells

were incubated in normal medium and high glucose medium for 48 h and subjected to
RNA extraction.

Statistical analysis

Statistical analysis employed a two-tailed, unpaired Student's t-test, One-way ANOVA,
or Fisher's exact test are described in figure legends. Chi square test (χ2-value) was
used for intergroup comparison of categorical data. All data are presented as mean ±
s.e.m. P<0.05 was considered to be statistically significant.

Acknowledgements

This study was supported by grants from the National Natural Science Foundation of
China (https://www.nsfc.gov.cn/; 81870359 and 2018YFA0801004 received by DL;
82000458 received by CC). The funders had no role in study design, data collection and
analysis, decision to publish, or preparation of the manuscript.

Author Contributions

DL supervised and designed this project. CC, DL , YQ wrote the manuscript. DL, CC,
and YX analyzed the data. CC, YQ and YX performed the experiments; YX, LC and LW
collected the clinical data.

Conflicts of Interest

The Authors declare that there is no conflict of interest.

References

Agemy SA, Scripsema NK, Shah CM, Chui T, Garcia PM, Lee JG, Gentile RC, Hsiao Y-
S, Zhou Q, Ko T, Rosen RB. 2015. Retinal vascular perfusion density mapping

using optical coherence tomography angiography in normals and diabetic retinopathy patients. *Retina* **35**:2353–2363. doi:10.1097/IAE.0000000000000862

Beckman JA, Creager MA. 2016. Vascular Complications of Diabetes. *Circ Res* **118**:1771–1785. doi:10.1161/CIRCRESAHA.115.306884

Chen C, Liu D. 2022. Establishment of Zebrafish Models for Diabetes Mellitus and Its Microvascular Complications. *J Vasc Res* **59**:1–9. doi:10.1159/000522471

Duh EJ, Sun JK, Stitt AW. 2017. Diabetic retinopathy: current understanding, mechanisms, and treatment strategies. *JCI Insight* **2**:e93751. doi:10.1172/jci.insight.93751

Echevarría M, Muñoz-Cabello AM, Sánchez-Silva R, Toledo-Aral JJ, López-Barneo J. 2007. Development of cytosolic hypoxia and hypoxia-inducible factor stabilization are facilitated by aquaporin-1 expression. *J Biol Chem* **282**:30207–30215. doi:10.1074/jbc.M702639200

Forbes JM, Cooper ME. 2013. Mechanisms of Diabetic Complications. *Physiological Reviews* **93**:137–188. doi:10.1152/physrev.00045.2011

Gleeson M, Connaughton V, Arneson LS. 2007. Induction of hyperglycaemia in zebrafish (Danio rerio) leads to morphological changes in the retina. *Acta Diabetol* **44**:157–163. doi:10.1007/s00592-007-0257-3

Gore AV, Monzo K, Cha YR, Pan W, Weinstein BM. 2012. Vascular development in the zebrafish. *Cold Spring Harbor Perspectives in Medicine* **2**:1–21. doi:10.1101/cshperspect.a006684

Hadidi H, Kamali R, Binesh A. 2022. Dynamics and energetics of water transport through aquaporin mutants causing nephrogenic diabetes insipidus (NDI): A molecular dynamics study. *J Biomol Struct Dyn* **40**:1273–1284. doi:10.1080/07391102.2020.1824813

Hinrichs GR, Baltzer S, Pallien T, Svenningsen P, Dalgaard EB, Hertz JM, Bistrup C, Jensen BL, Klussmann E. 2022. A Novel AQP2 Sequence Variant Causing Aquaporin-2 Retention in the Cytoplasm and Autosomal Dominant Nephrogenic Diabetes Insipidus. *Kidney Int Rep* **7**:2289–2294. doi:10.1016/j.ekir.2022.07.001

Jung S-H, Kim YS, Lee Y-R, Kim JS. 2016. High glucose-induced changes in hyaloid-retinal vessels during early ocular development of zebrafish: a short-term animal

model of diabetic retinopathy: HG-induced changes in hyaloid-retinal vessels in

ZF. *British Journal of Pharmacology* **173**:15–26. doi:10.1111/bph.13279

Kamei M, Brian Saunders W, Bayless KJ, Dye L, Davis GE, Weinstein BM. 2006.

Endothelial tubes assemble from intracellular vacuoles in vivo. *Nature* **442**:453–

456. doi:10.1038/nature04923

Khan LA, Zhang H, Abraham N, Sun L, Fleming JT, Hall DH, Gobel V. 2013.

Intracellular lumen extension requires ERM-1-dependent apical membrane

expansion and AQP-8-mediated flux **15**:143–156.

Klein R, Myers CE, Lee KE, Gangnon R, Klein BEK. 2012. Changes in Retinal Vessel

Diameter and Incidence and Progression of Diabetic Retinopathy. *Arch*

Ophthalmol **130**. doi:10.1001/archophthalmol.2011.2560

Krueger J, Liu D, Scholz K, Zimmer A, Shi Y, Klein C, Siekmann A, Schulte-merker S,

Cudmore M, Ahmed A, Noble F. 2011. Flt1 acts as a negative regulator of tip cell

formation and branching morphogenesis in the zebrafish embryo **2120**:2111–

2120. doi:10.1242/dev.063933

Kwan KM, Fujimoto E, Grabher C, Mangum BD, Hardy ME, Campbell DS, Parant JM,

Yost HJ, Kanki JP, Chien C-B. 2007. The Tol2kit: a multisite gateway-based

construction kit for Tol2 transposon transgenesis constructs. *Dev Dyn* **236**:3088–

3099. doi:10.1002/dvdy.21343

Lee Y, Yang J. 2021. Development of a zebrafish screening model for diabetic

retinopathy induced by hyperglycemia: Reproducibility verification in animal

model. *Biomedicine & Pharmacotherapy* **135**:111201.

doi:10.1016/j.biopha.2020.111201

Liang J, Gui Y, Wang W, Gao S, Li J, Song H. 2010. Elevated glucose induces

congenital heart defects by altering the expression of tbx5, tbx20, and has2 in

developing zebrafish embryos. *Birth Defects Res A Clin Mol Teratol* **88**:480–486.

doi:10.1002/bdra.20654

Liu J, Stainier DYR. 2012. Zebrafish in the study of early cardiac development.

Circulation Research **110**:870–874. doi:10.1161/CIRCRESAHA.111.246504

Liu K, Nagase H, Huang CG, Calamita G, Agre P. 2006. Purification and functional characterization of aquaporin-8. *Biology of the Cell* **98**:153–161. doi:https://doi.org/10.1042/BC20050026

Nico B, Ribatti D. 2010. Aquaporins in tumor growth and angiogenesis. *Cancer Letters* **294**:135–138. doi:10.1016/j.canlet.2010.02.005

Osmani N, Péglion F, Chavrier P, Etienne-Manneville S. 2010. Cdc42 localization and cell polarity depend on membrane traffic. *The Journal of cell biology* **191**:1261–9. doi:10.1083/jcb.201003091

Papadopoulos MC, Saadoun S, Verkman AS. 2008. Aquaporins and cell migration. *Pflugers Arch - Eur J Physiol* **456**:693–700. doi:10.1007/s00424-007-0357-5

Preston GM, Carroll TP, Guggino WB, Agre P. 1992. Appearance of water channels in Xenopus oocytes expressing red cell CHIP28 protein. *Science* **256**:385–387. doi:10.1126/science.256.5055.385

Qin Y, Wang S, Duan X, Liu D. 2019. Expression analysis of the aquaporins during zebrafish embryonic development. *Gene Expression Patterns* **32**:38–43. doi:10.1016/j.gep.2019.04.001

Reizer J, Reizer A, Saier MH. 1993. The MIP family of integral membrane channel proteins: sequence comparisons, evolutionary relationships, reconstructed pathway of evolution, and proposed functional differentiation of the two repeated halves of the proteins. *Crit Rev Biochem Mol Biol* **28**:235–257. doi:10.3109/10409239309086796

Ribatti D, Ranieri G, Annese T, Nico B. 2014. Aquaporins in cancer. *Biochimica et Biophysica Acta (BBA) - General Subjects* **1840**:1550–1553. doi:10.1016/j.bbagen.2013.09.025

Sada K, Nishikawa T, Kukidome D, Yoshinaga T, Kajihara N, Sonoda K, Senokuchi T, Motoshima H, Matsumura T, Araki E. 2016. Hyperglycemia Induces Cellular Hypoxia through Production of Mitochondrial ROS Followed by Suppression of Aquaporin-1. *PLoS One* **11**:e0158619. doi:10.1371/journal.pone.0158619

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,

Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**:676–682. doi:10.1038/nmeth.2019

Smith BL, Agre P. 1991. Erythrocyte Mr 28,000 transmembrane protein exists as a multisubunit oligomer similar to channel proteins. *Journal of Biological Chemistry* **266**:6407–6415. doi:10.1016/S0021-9258(18)38133-X

Toh H, Smolentsev A, Bozadjian RV, Keeley PW, Lockwood MD, Sadjadi R, Clegg DO, Blodi BA, Coffey PJ, Reese BE, Thomson JA. 2019. Vascular changes in diabetic retinopathy—a longitudinal study in the Nile rat. *Lab Invest* **99**:1547–1560. doi:10.1038/s41374-019-0264-3

Verkman AS. 2005. More than just water channels: unexpected cellular roles of aquaporins. *Journal of Cell Science* **118**:3225–3232. doi:10.1242/jcs.02519

Verkman AS. 2002. Aquaporin water channels and endothelial cell function*. *J Anatomy* **200**:617–627. doi:10.1046/j.1469-7580.2002.00058.x

Verkman AS, Hara-Chikuma M, Papadopoulos MC. 2008. Aquaporins—new players in cancer biology. *J Mol Med* **86**:523–529. doi:10.1007/s00109-008-0303-9

Wang W, Lo ACY. 2018. Diabetic Retinopathy: Pathophysiology and Treatments. *Int J Mol Sci* **19**:1816. doi:10.3390/ijms19061816

Wang X, Ling CC, Li L, Qin Y, Qi J, Liu X, You B, Shi Y, Zhang J, Jiang Q, Xu H, Sun C, You Y, Chai R, Liu D. 2016. MicroRNA-10a / 10b represses a novel target gene mib1 to regulate angiogenesis **1**:140–150. doi:10.1093/cvr/cvw023

Wiggenhauser LM, Qi H, Stoll SJ, Metzger L, Bennewitz K, Poschet G, Krenning G, Hillebrands J-L, Hammes H-P, Kroll J. 2020. Activation of Retinal Angiogenesis in Hyperglycemic $pdx1^{-/-}$ Zebrafish Mutants. *Diabetes* **69**:1020–1031. doi:10.2337/db19-0873

Wong TY. 2011. Retinal Vessel Diameter as a Clinical Predictor of Diabetic Retinopathy Progression: Time to Take Out the Measuring Tape. *Arch Ophthalmol* **129**:95. doi:10.1001/archophthalmol.2010.347

Yang B, Kim JK, Verkman AS. 2006. Comparative efficacy of $HgCl_2$ with candidate aquaporin-1 inhibitors DMSO, gold, TEA^+ and acetazolamide. *FEBS Letters* **580**:6679–6684. doi:10.1016/j.febslet.2006.11.025

708   Zhang C, Luo D, Xie H, Yang Q, Liu D, Tang L, Zhang Jingting, Li W, Tian H, Lu L, Sun

709   X, Xu G-T, Zhang Jingfa. 2022. Aquaporin 11 alleviates retinal Müller intracellular

710   edema through water efflux in diabetic retinopathy. *Pharmacol Res* **187**:106559.

711   doi:10.1016/j.phrs.2022.106559

A High glucose solution

Tg(fli1ep:EGFP-CAAX)^ntu666

vessel diameter quantification

microangiography

Glucose → microangiography + imaging

0 1 2 3 4 5 (dpf)

B ctrl **C** high glucose

fli1ep:EGFP-CAAX

B' microangiography **C'**

B'' merge **C''**

D raletive vessel diameter (IOC)

ctrl   high glu

***

E *fli1ep:EGFP-CAAX* **F**

E' microangiography **F'**

E'' merge **F''**

G phenotype (%)

no perfusion
perfusion

ctrl   high glu

****

H raletive vessel diameter (ISV)

ctrl   high glu

****

A

tol2 — fli1ep — aqp1a.1 — mCherry — tol2

+ tol2 transposase mRNA

*Tg(fli1ep:EGFP-CAAX)*ntu666 embryos

EGFP

EGFP + mCherry

Genetic Mosaicism of *aqp1a.1* in *Tg(fli1ep:EGFP-CAAX)*ntu666 background

B

mosac selection   glucose   imaging

0   1   2   3 (dpf)

C

fli1ep:EGFP-CAAX

fli1ep:aqp1a.1-mCherry

merge

D

raletive vessel diameter

****

EGFP+ mCherry-   EGFP+ mCherry+

A aqp8a.1-EGFP

Endothelail cell

10 μm

B aqp8a.1-EGFP   aqp1a.1-mCherry   merge

Endothelail cell

10 μm

C *Tg(fli1:EGFP–cdc42wt)*   *Tg(fli1a:aqp1a.1-mCherry)*   merge

10 μm

D   ctrl   aqp1a.1 KO
8a.1 MO

Tg(fli1:EGFP–cdc42wt)

10 μm

E   ctrl   loss of Aqp

Endothelail cell

F

area of vacuoles/ISV (μm²)

**

ctrl   loss of Aqp

Supplementary Figure 1. The diagram for showing the constructs of three transgenic lines.

Supplementary Figure 2. The glucose treatment of zebrafish embryo. A. The time line of glucose treatment of zebrafish embryos. B. The glucose level of body fluid of zebrafish embryos.

Supplementary Figure 3. WISH analysis of *aqp1a.1* and *aqp8a.1*. A-L. The *aqp1a.1* expression in 20 hpf, 22 hpf, 24 hpf, 36 hpf, 48 hpf and 72 hpf embryos. M-S. The *aqp8a.1* expression in 20 hpf, 24 hpf, 36 hpf, 48 hpf and 72 hpf embryos.

A

CRISPR/Cas9-mediated *aqp1a.1* knockout

ATG

exon1 — exon2 — exon3 — exon4

sgRNA   PAM

ACAGACAAAAGGCGGCGGGACGTGTCGGGCTCCGCTCCTCTGGCCATCGGCCTCAGTGTTT
TGTCTGTTTTCCGCCGCCCTGCACAGCCCGAGGCGAGGAGACCGGTAGCCGGAGTCACAAA

B

WT | ACAGACAAAAGGCGGCGGGACGTGTCGGGCTCCGCTCCTCTGGCCATCGGCCTCAGTGTTT

Genotypes of G0 generations

ACAGACAAAAGGCGGCGGGACGTGTCGGGCTCCGCTCCTC —————ATCGGCCTCAGTGTTT (-5)
ACAGACAAAAGGCGGCGGGACGTGTCGGGCTCCGCTC ———————CATCGGCCTCAGTGTTT (-7)
ACAGACAAAAGGCGGCGGGACGTGTCGGGCTCCGCTCCTCTGGCC ———————TCAGTGTTT (-7)
ACAGACAAAAGGCGGCGGGACGTGTCGGGCTCCGCTCCTCTGGC —————————————GTTT (-13)
A—————————————————————————— GGCTCCGCTCCTCAGTTTAGCGGCCTCAGTGTTT (-26)
ACAGACAAAAGGCGGCGGGACGTGTCGGGCTCCGCTCCTCTGGCCTCATCGGCCTCAGTGTTT (+2)

C   Predicted protein sequence of homozygous F3 line

(260 aa)

(169 aa)

Supplementary Figure 4. Generation of zebrafish *aqp1a.1* knockout using CRISPR/Cas9. A. Schematic diagram showing gRNA targeting site on the second exon of *aqp1a.1* gene. Starting codon (ATG) site is indicated by arrow. B. Genotypes of G0 generations. Numbers in the brackets show the number of nucleotides were deleted (−) or inserted (+). Inserted nucleotide is in red. WT, wild-type. C. Schematic diagram showing the predicted proteins encoded by the two homozygous F3 line. The mutants are reading frameshift mutations that result in truncated proteins. The gray rectangles indicate the wrong coded amino acid sequences.

A

CRISPR/Cas9-mediated *aqp8a.1* knockout

B

C

Supplementary Figure 5. Generation of zebrafish *aqp8a.1* knockout using CRISPR/Cas9. A. Schematic diagram showing gRNA targeting site on the second exon of *aqp8a.1* gene. Starting codon (ATG) site is indicated by arrow. B. Genotypes of G0 generations. Numbers in the brackets show the number of nucleotides were deleted (−) or inserted (+). Inserted nucleotide is in red. C. Genotypes of F1 generations.

Supplementary Figure 6. Knockdown of EC-enriched aquaporins (*aqp1a.1* or *aqp8a.1*) leads to defective vascular lumenization in zebrafish. A, Confocal images of ISV phenotypes in 48-hpf *Tg(kdrl:EGFP)* control embryos and *aqp1a.1*-knockdown and *aqp8a.1*- knockdown embryos. Red and blue square brackets indicate artery and vein, respectively. Blue arrowheads indicate the narrowed ISVs. **B,** Quantification of the diameters of ISV, DA and PCV. Data are shown as mean ± s.e.m. One-way ANOVA analysis is applied. *, p<0.05. **, p < 0.01. ****, p < 0.0001.

Supplementary Figure 7. HgCl₂ treatment impairs blood vessel lumenization. A, Representative confocal images of ISV phenotypes in 48-hpf *Tg(kdrl:ras-mCherry)* upon HgCl2 treatment. **B-D,** Quantification of the diameters of ISV, DA and PCV. Data are shown as mean ± s.e.m. One-way ANOVA analysis is applied. ns, not significant. ***, p < 0.001. ****, p < 0.0001.

Supplementary Table S1. Primer list

Primers for ISH probe synthesis		
	Left primer	Right primer
aqp1a.1	5'-CCCAGACCAGGAGATCAAGG-3'	5'-CGACCCAGTAGACCCAGTA-3'
aqp8a.1	5'-CTGCCCTTCTTCGAGCACTA-3'	5'-ATTTGCCGTCACAGTGAGTC-3'
Primers for RT-PCR		
	Left primer	Right primer
AQP1(Human)	5'-CCTCTCTGTAGCCCTTGGAC-3'	5'-GGCCAGGATGAAGTCGTAGA-3'
aqp1a.1(Zebrafish)	5'-CCCAGACCAGGAGATCAAGG-3'	5'-CGACCCAGTAGACCCAGTA-3'
aqp8a.1(Zebrafish)	5'-CTGCCCTTCTTCGAGCACTA-3'	5'-ATTTGCCGTCACAGTGAGTC-3'
sgRNA sequence		
aqp1a.1 sgRNA	5'-GGCTCCGCTCCTCTGGCCAT-3'	
aqp8a.1 sgRNA	5'-AGGAATGGAGGTGATCCTCC-3'	
Primers for sgRNA synthesis		
	Left primer	Right primer
aqp1a.1 sgRNA	5'-TGATACAAGTAGCATTCTGTGTCG-3'	5'-TGAGCATTCTCTAGCTGCA-3'
aqp8a.1 sgRNA	5'-CTATTCCTATTAGACCAAAGT-3'	5'-TGATGTGACCTGGAATC-3'
Morpholino (MO) sequence		
aqp1a.1 splicing MO	5'-GTGTGAATGCTGATTACTCACGGCT-3'	
aqp8a.1 splicing MO	5'-ACCACCACTGGAAAACAACAAGTGT-3'	