Evaluating the mouse neural precursor line, SN4741, as a suitable proxy for midbrain dopaminergic neurons

Rachel J. Boyd[1,3], Sarah A. McClymont[1,3], Nelson B. Barrientos[1], Paul W. Hook[1], William D. Law[1], Rebecca J. Rose[1], Eric L. Waite,[1] Dimitrios Avramopoulos[1], and Andrew S. McCallion*[1,2]

1. McKusick-Nathans Department of Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA.

2. Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD 21287, USA.

3. These authors contributed equally.

*To whom correspondence should be addressed

Electronic addresses:

Rachel J. Boyd - rboyd25@jhmi.edu, Sarah A. McClymont - sarahmcclymont@gmail.com, Nelson B. Barrientos - nbarrie1@jhu.edu, Paul W. Hook - phook2@jhmi.edu, William D. Law - williamdlaw@gmail.com, Rebecca J. Rose – rrose10@jhmi.edu, Eric L. Waite – eric.waite@pennmedicine.upenn.edu, Dimitrios Avramopoulos - adimitr1@jhmi.edu, Andrew S. McCallion - andy@jhmi.edu

KEYWORDS:

Parkinson disease, mouse-derived cell lines, immortalized cell lines, chromatin accessibility,

RNA-seq, ATAC-seq, scRNA-seq, genomic characterization, disease-relevant model systems

ABSTRACT:

To overcome the ethical and technical limitations of *in vivo* human disease models, the broader

scientific community frequently employs model organism-derived cell lines to investigate of

disease mechanisms, pathways, and therapeutic strategies. Despite the widespread use of certain

in vitro models, many still lack contemporary genomic analysis supporting their use as a proxy

for the affected human cells and tissues. Consequently, it is imperative to determine how

accurately and effectively any proposed biological surrogate may reflect the biological processes

it is assumed to model. One such cellular surrogate of human disease is the established mouse

neural precursor cell line, SN4741, which has been used to elucidate mechanisms of

neurotoxicity in Parkinson disease for over 25 years. Here, we are using a combination of classic

and contemporary genomic techniques – karyotyping, RT-qPCR, single cell RNA-seq, bulk

RNA-seq, and ATAC-seq – to characterize the transcriptional landscape, chromatin landscape,

and genomic architecture of this cell line, and evaluate its suitability as a proxy for midbrain

dopaminergic neurons in the study of Parkinson disease. We find that SN4741 cells possess an

unstable triploidy and consistently exhibits low expression of dopaminergic neuron markers

across assays, even when the cell line is shifted to the non-permissive temperature that drives

differentiation. The transcriptional signatures of SN4741 cells suggest that they are maintained in

an undifferentiated state at the permissive temperature and differentiate into immature neurons at

the non-permissive temperature; however, they may not be dopaminergic neuron precursors, as

previously suggested. Additionally, the chromatin landscapes of SN4741 cells, in both the

differentiated and undifferentiated states, are not concordant with the open chromatin profiles of

ex vivo, mouse E15.5 forebrain- or midbrain-derived dopaminergic neurons. Overall, our data

suggest that SN4741 cells may reflect early aspects of neuronal differentiation but are likely not

a suitable a proxy for dopaminergic neurons as previously thought. The implications of this study

extend broadly, illuminating the need for robust biological and genomic rationale underpinning

the use of *in vitro* models of molecular processes.

BACKGROUND:

in vitro cellular surrogates present an excellent opportunity for elucidating the molecular

mechanisms behind human disease without the ethical and technical limitations of *in vivo*

systems. As such, most studies of human disease that employ genomic or cellular manipulations

or assays that require high cell quantity and quality, are often conducted *in vitro* to ensure

biological and statistical robustness[1–3]. For example, *in vitro* models are frequently employed

in studies of the role of genomic regulation in human disease, identification of candidate genes

and regulatory elements and evaluation of their functional characteristics through genetic

manipulations and high-throughput assays[4–7]. As genome-wide association studies (GWASs)

continue to indict human disease-associated variants, it is becoming evident that most of them lie

within non-coding regions of the genome[8]. Such regions frequently represent cis-regulatory

elements (CREs), required for the transcriptional modulation of cognate genes. The assays

required to evaluate their function[4,9,10], or connect CREs with the promoters they

modulate[11], often require large cell numbers, making, *in vitro* cellular systems the preferred

strategy.

Prioritizing non-coding GWAS variants and disease-relevant sequences for extensive

investigation requires knowledge of their chromatin accessibility status. Open chromatin is prone

to harbor functional sequences; and since chromatin accessibility profiles vary across cell types

and developmental time, it is important to prioritize disease-associated variants that lie within

open chromatin regions in the disease-relevant cell type(s) [8,12,13]. It is also critical to

functionally evaluate the biological consequences of disease-associated variation, test the

efficacy of potential therapeutics, and observe the effects of disease-relevant insults in the

appropriate cellular context[8,14,15]. Therefore, when studying disease associated variation, the

most effective *in vitro* cellular surrogates should ideally mimic the chromatin architecture and

transcriptional profiles of the *in vivo* cell types affected by disease.

In Parkinson disease (PD), midbrain (MB) dopaminergic (DA) neurons in the substantia

nigra (SN) are the primary affected cell type[16]. Preferential degeneration of these neurons

elicits a progressive neurodegenerative disorder characterized by motor deficits[16]. As the

second most common neurodegenerative disorder, affecting approximately 1% of adults over 70

years old[17,18], PD is the focus of extensive research efforts. As such, various cell lines have

been used as *in vitro* proxies of MB DA neurons to study the cellular impacts of PD-relevant

insults, as well as candidate PD-associated sequences, their functions, and their potential as

therapeutic targets[19].

One such cell line, SN4741, is reported to be a clonal DA neuronal progenitor line that

was established in 1999 from mouse embryonic day 13.5 (E13.5) SN tissue[20]. The SN was

dissected from transgenic mice containing 9.0 kb of the 5' promoter region of tyrosine

hydroxylase (*TH*), fused to the temperature-sensitive mutant Simian Virus 40 T antigen

(SV40Tag-tsA58) oncogene[20]. The goal of this *TH* promoter transgene was to enable selective

acquisition of DA neurons, while the purpose of the SV40Tag oncogene was to facilitate

conditional immortalization of the cell line. The temperature sensitive mutant form of this

immortalizing gene (tsA58) should permit uncontrolled differentiation and proliferation at the

permissive temperature (33°C), maintain cells in an undifferentiated state at 37°C, and since

tsA58 displays diminished activity at 39°C, it should direct differentiation that more closely

resembles primary cells when the culture is shifted to this non-permissive temperature[20].

As an established mouse neural precursor line, SN4741 cells have since been used to

elucidate mechanisms of neurotoxicity in PD[21–25], test the efficacy of therapeutic targets

against PD relevant insults[26,27], and assay the impacts of PD-associated genetic

mutation[28,29] and transcriptional regulation[30–32]. Important technological advances have

also arisen since the genesis and implementation of the SN4741 cell line, including chromatin

conformation capture technologies[11,33,34], RNA-sequencing (RNA-seq)[35], and assay for

transposase-accessible chromatin using sequencing (ATAC-seq)[36]. In this study, we exploit

these modern approaches to assess the suitability of SN4741 as an *in vitro* proxy for DA neurons

and determine the extent to which this cell line is appropriate for prioritizing and investigating

the mechanisms by which PD-associated variation confers disease risk.

Through a combination of karyotyping, single-cell (sc)RNA-seq, and RT-qPCR, we

evaluate the genomic integrity of this immortalized cell line, determine how the transcriptional

profile and expression of DA neuron marker genes in this line changes between undifferentiated

(37°C) and differentiated (39°C) states, and evaluate whether these transcriptional changes are

consistent throughout the differentiation process. The data we collect suggests that while these

cells show evidence that they are exiting a proliferative state and entering a more differentiated

state, they are an unsuitable model of SN DA neurons, as they possess aneuploidy and structural

abnormalities, as well as consistently low expression of DA neuron markers upon differentiation. We employ bulk RNA-seq to quantify transcriptional differences between differentiated and undifferentiated SN4741 cells and determine that, while transcriptional profiles change to reflect differentiation, they do not show strong evidence that these cells are entering a DA state. We then compare chromatin accessibility profiles of undifferentiated and differentiated SN4741 cells with those of *ex vivo* mouse E15.5 midbrain (MB) and forebrain (FB) neurons and determine that the chromatin accessibility profiles of SN4741 cells do not reflect the cellular population from which they were derived. Collectively, cytogenetic, chromatin, and transcriptional data suggest that the SN4741 cell line is not as strong a cellular surrogate for DA neurons as previously thought. Ultimately, this work underscores the importance of leveraging technological advances in genomic and cellular analyses to evaluate, and re-evaluate, the suitability of established model systems in disease biology.

RESULTS:

SN4741 is an Unstable Polyploid Cell Line:

G-band karyotyping was performed on 20 SN4741 metaphase spreads and a representative karyogram (**Figure 1A**) was generated. The karyotype was interpreted as an abnormal, polyploid, karyotype with complex numerical abnormalities and unbalanced, structural abnormalities. While most, but not all, abnormalities were consistently present in these cells; none of the 20 cells assessed had the same chromosome complement, and no normal cells were observed. All cells possessed at least one copy of each mouse autosome (1 through 19) and female sex chromosomes; however, most chromosomes were triploid in each cell (**Figure 1B**). These karyotypic abnormalities already call into question the viability of these cells as a

surrogate for human neurodegenerative disease. Since these cells are genetically unstable, there

may be large experimental batch effects as the cell populations shift across divisions.

Furthermore, gene dosage effects that severely deviate from normal copy number in DA neurons

may lead to confounding and unreliable results.

Undifferentiated and Differentiated SN4741 Cells Express Similar Levels of Dopaminergic

Neuron Marker Genes by RT-qPCR:

Preliminary expression analysis by RT-qPCR confirmed expression of a variety of DA

neuron markers: forkhead box A2 *(Foxa2),* nuclear receptor subfamily 4 group A, member 2

(Nr4a2), solute carrier family 6 member 3 *(Slc6a3),* and tyrosine hydroxylase *(Th).* Compared to

the expression of these markers in the undifferentiated SN4741 cell culture (37°C), relative

expression of all markers remained at similar levels when the cells were shifted to the higher

temperature condition (39°C), except for an increase in *Th* expression (**Figure 1C**). While

elevated *Th* has been used as a marker of differentiation into DA neurons in previous work with

SN4741 cells[20,37,38], an increase in *Th* expression is not exclusively associated with DA

neurons. *Th* is a marker for all catecholaminergic neurons (dopaminergic and adrenergic)[39],

and evidence suggests that *Th* expression is transient in other neurons throughout embryonic

development[40–42]. These results indicate that at the non-permissive temperature, SN4741 cells

may not be fully differentiating into DA neuron progenitors.

scRNA-seq Reveals that SN4741 Cells Differentiate at the Non-Permissive Temperature,

but Lack Expression of DA Neuron Marker Genes:

To assess the consistency of the differentiation protocol, transcriptomes were generated

from ≥17,000 cells across four replicates cultured at the permissive temperature (37°C) and four

replicates cultured at the non-permissive temperature (39°C). Analysis of the single-cell

transcriptomes reveal that the cells cluster by growth temperature (**Figure 1D**). This separation

of cells by temperature is accompanied by changes to the cell cycle, with cells at the permissive

(37°C) temperature mostly in either G2M or S phase, while cells at the non-permissive

temperature (39°C) are mostly in G1 phase (**Figure 1E**), indicating that they may be

differentiated. In expression analysis, markers of proliferation that are expressed in G2M phase,

like Marker of Proliferation Ki-67 (*Mki67*), are predominantly expressed in cells at the

permissive temperature (**Figure 1F**), corroborating the cell cycle analysis. When shifted to the

non-permissive temperature, SN4741 cells appear to robustly differentiate, exemplified by a

decrease in the expression of Nestin (*Nes*), a neural stem cell marker (**Figure 1F**). Additional

transcriptional changes at this non-permissive temperature include an increase in the expression

of a neural marker CUGBP Elav-Like Family Member 5 (*Celf5*)[43], as well as genes that have

been found to regulate neural stem cell self-renewal (Inhibitor of DNA Binding 2, *Id2;* High

Mobility Group AT-Hook 2, *Hmga2*)[44,45], neurogenesis (Iroquois Homeobox 3, *Irx3*)[46],

and arborization of neurons (Sodium Voltage-Gated Channel Beta Subunit 1, *Scn1b*)[47],

indicating that these cells may be differentiating into neural precursor cells (**Figure 1G**).

Furthermore, Cadherin 13 (*Cdh13*), a modulator of GABAergic neurons, is significantly

upregulated at this non-permissive temperature (**Figure 1G**), while the expression of a variety of

DA neuron markers fail to be detected in either the permissive or non-permissive temperatures.

Markers, including Aldehyde Dehydrogenase 1 Family Member A1 (*Aldh1a1), Foxa2,* LIM

Homeobox Transcription Factor 1 Beta *(Lmx1b), Nr4a2*, Paired-like homeodomain 3 (*Pitx3),*

Slc6a3, and *Th* have few to no reads assigned to them **(Figure 1H)**. Collectively, these results

suggest that while SN4741 cells are differentiating towards a neuronal fate when shifted to the

nonpermissive temperature, they may not be entering a clear DA trajectory under these

conditions.

ATAC-seq Identifies Differential Open Chromatin Profiles in SN4741 Cells at the

Permissive and Non-Permissive Temperatures:

To consider how chromatin accessibility changes between the two temperatures, we

performed ATAC-seq on SN4741 cells in both the undifferentiated and differentiated states.

Libraries were confirmed to be technically and biologically relevant (**Supplemental Figure 1**),

and well correlated between replicates (**Supplemental Figure 2; Figure 2A-B**).

A total of 83,778 consensus open chromatin regions were identified, with 70% of peaks

shared between the two temperatures (**Figure 2C**). Principal component analysis of these

consensus regions suggests a clear separation in the chromatin state between the two

temperatures (**Figure 2D**). To explore these differences, we performed differential accessibility

analysis with DiffBind[48], to find a total of 5,055 differentially accessible regions: 2,654

enriched in the permissive temperature and 2,401 enriched at the non-permissive temperature

($\log_2$FC > 1, FDR < 0.05; **Figure 2E**).

Gene ontology of genes adjacent to differentially accessible regions largely recapitulate

the scRNA-seq analysis; functions associated with regions preferentially open at the permissive

temperature suggest the maintenance of the undifferentiated, cell-cycling state (**Figure 2F**). The

gene ontology of genes adjacent to those regions preferentially accessible at the non-permissive

temperature is less coherent and suggest cell differentiation towards several fates (blood vessels,

cartilage, tooth), none of which are neuronal and, perhaps unsurprisingly, demonstrate evidence

of response to temperature stress[49] (**Figure 2G**).

Overall, there is a shift in the chromatin accessibility between the two temperatures that

indicate the cells transition from an undifferentiated to differentiated state as the cells move from

the permissive to non-permissive temperature. The differences in chromatin accessibility further

confirm that SN4741 cells are not differentiating towards a neuronal lineage.

Comparison of chromatin accessibility in SN4741 cells fails to recapitulate the chromatin

landscape of *ex vivo* mouse DA neurons:

To evaluate the potential relationship between SN4741 cells and DA neurons they are

presumably modelling, we compared the chromatin accessibility between the SN4741 cells at

both temperatures to previously generated *ex vivo* mouse embryonic DA neuron chromatin

accessibility profiles (NCBI GEO: GSE122450;[50]).

Considering the consensus peak set of 165,334 regions generated from all *in vivo* and *ex*

vivo samples and their normalized read counts, we observe a clear separation between the

SN4741 cell culture model and the *ex vivo* DA neurons by correlation and principal component

analysis (**Figure 3A, B**). Examining the raw overlap of peaks between the SN4741 cells and *ex*

vivo neurons, just 12.5% (20,667) are present in all four cell types/conditions (**Figure 3C**). The

chromatin profiles are largely exclusive between the SN4741 cell culture model and the *ex vivo*

DA neurons: 41.3% (68,304) of regions are accessible solely in the *ex vivo* neuron populations

and 40% (65,857) are exclusively accessible in the SN4741 cell culture models. There is little

overlap between the *ex vivo* and cultured samples. In comparison of the *ex vivo* midbrain DA

neurons to the non-permissive, differentiated temperature, only 183 peaks are restricted to these

populations.

The chromatin profiles between *ex vivo* embryonic DA neurons and their prospective *in*

vitro cell culture surrogate are virtually independent. They exhibit scant overlap in their global

chromatin profiles and bear little resemblance to each other at regulatory regions of key DA

neuron genes (**Figure 3D**). Neither the analysis of the SN4741 chromatin accessibility profiles in

isolation or in comparison with *ex vivo* neurons would suggest these cells to be appropriate

models of embryonic DA neurons.

Transcriptional Changes in SN4741 Cells Indicate Differentiation from Pluripotent Stem

Cells into Brain Cells that do not Fully Resemble MB DA Neurons:

Bulk RNA-seq data were also generated for SN4741 cells, at both the permissive and

non-permissive temperatures, to determine whether transcriptome changes reflect differentiation

into DA neurons, or other neural cell types. To evaluate the RNA-seq libraries, quality-control

measures were performed *in silico* (**Supplemental Figure 3**). PCA (**Supplemental Figure 3B**),

and sample-sample distances (**Supplemental Figure 3C**) reaffirmed that samples cultured at the

same temperature are more like one-another than samples cultured at the alternate temperature.

We found that 735 genes were upregulated at the non-permissive temperature (adjusted

p-value < 0.01 and $\log_2$ FC > 1.5), and 954 genes were downregulated (adjusted p-value < 0.01

and $\log_2$ FC< -1.5) at the non-permissive temperature. The list of genes significantly

downregulated   at   the   non-permissive   temperature   was   submitted   to   Enrichr

(https://maayanlab.cloud/Enrichr/) [51–53] for gene ontology (GO) and analysis of cell type

markers. Consistent with the observation that cells at the non-permissive temperature are

differentiated and in G1 phase of the cell cycle, downregulated genes resulted in GO terms

strongly enriched for mitotic and DNA replication processes (**Figure 4A)**. Additionally,

significantly downregulated genes at the non-permissive temperature overlap with subsets of

PanglaoDB[54] cell type marker genes, suggesting that these cells are shifting away from a state

that resembles neural stem cells (**Figure 4B**).

Similarly, the list of significantly upregulated genes was submitted to Enrichr for GO and

analysis of cell type markers. As expected, upregulated genes resulted in GO terms for biological

processes that indicate a more terminally differentiated cell type **(Figure 4C**): "synaptic vesicle

docking," "negative regulation of osteoblast proliferation," "lens fiber cell differentiation,"

"regulation of osteoblast proliferation," and "forebrain regionalization". While not included in

the top 10 terms by combined score ranking, "neuron remodeling," "synaptic transmission,

glutamatergic," "neuron maturation," and "synaptic transmission, cholinergic" were also

identified as significantly associated terms. Notably, "synaptic transmission, dopaminergic" and

"dopaminergic neuron differentiation" were also listed as insignificant terms (**Figure 4C**), as *Th*

was the lone overlapping marker gene for these terms.

In line with GO terms enriched for biological processes involving differentiation,

possibly in neuronal cells, overlapping PanglaoDB[54] cell type marker genes suggest that

SN4741 cells at the non-permissive temperature most significantly resemble immature neurons

(**Figure 4D**). "Oligodendrocytes," "retinal progenitor cells," "satellite glial cells," "dopaminergic

neurons," "adrenergic neurons," "GABAergic neurons," and "glutamatergic neurons" were also

listed as cell types with significant marker gene overlap.

The distribution of various cell type marker genes on a volcano plot, indicating the

$\log_2$FC in expression and -$\log_{10}$ adjusted p-values of DE genes, reveals that the specific genes

overlapping "pluripotent stem cell" markers (26/112), cluster as the most highly significantly

downregulated genes **(Figure 4E)**. In contrast, only two of the upregulated marker genes

overlapping "immature neurons" (16/136, **Figure 4F**) and "oligodendrocytes" (17/178, **Figure**

4G) cluster in a similarly strong way. Plotting the 11/119 overlapping upregulated genes for

"dopaminergic neurons" (**Figure 4H**) reveals that 7/11 overlapping genes (*Celf5, Dpys15,*

Cacna1b, Tmem179, Nova2, Nrx1, and *Cntn2*) are also marker genes for immature neurons.

Plotting the DA neuron markers also assayed by RT-qPCR validates that the relative expression

of these markers is consistent between these highly sensitive assays. At 39°C, *Th* expression

increases ($\log_2$FC = 1.552651); *Nr4a2* expression decreases ($\log_2$FC = -1.042794), and

expression of *Slc6a3* and *Foxa2* is not significantly different between conditions: (*Slc6a3* was

filtered out due to low read counts across both temperature conditions and *Foxa2* $\log_2$FC = -

0.4142541).

To confirm the GO-indicted cell types, normalized read counts for select marker genes

were plotted for each temperature replicate: *Celf5*[43]*, Nrxn1*[55]*, Ntrk1*[56]*,* and *Unc13a*[57]*,*

for "immature neurons" were upregulated at 39°C relative to cells at 37°C (**Figure 5A**);

Olig3[58]*, Il33*[59]*, Hdac11*[60]*,* and *Ptgds*[61] for "oligodendrocytes" were upregulated at

39°C relative to cells at 37°C (**Figure 5B**); and *Ccna2*[62]*, Cdc6*[63]*, Cenpf*[64]*,* and *Gins1*[65]

for "pluripotent stem cells" were downregulated at 39°C relative to cells at 37°C (**Figure 5C**).

The differentially expressed gene sets were then analyzed using STRING (https://string-

db.org/)[66]. The set of upregulated genes was enriched for protein-protein interactions (number

of edges = 705; expected number of edges = 438; PPI enrichment p-value = < 1.0e-16) and GO

terms such as "Neuron differentiation" (GO:0030182; FDR = 0.0016), "Neuron development"

(GO:0048666; FDR = 0.0070), and "Neurogenesis" (GO:0022008; FDR = 0.0085) supporting

that this upregulated gene set is a meaningful group likely belonging to a network involved in

neuronal maturation. The set of downregulated genes was also enriched for protein-protein

interactions (number of edges = 9251; expected number of edges = 1910; PPI enrichment p-

value = < 1.0e-16) and GO terms such as "Cell cycle" (GO:0007049; FDR = 5.90e-36), "Mitotic

cell cycle" (GO:0000278; FDR = 6.97e-34), and "Cell division" (GO:0051301; 6.35e-25) further

confirming that these cells are no longer undergoing cell division.

Finally, previously generated reads per kilobase of exon per million reads mapped

(RPKM) from *ex vivo* E15.5 mouse embryonic DA neuron bulk RNA-seq[50] were used to

compare how closely the SN4741 transcriptome resembles the neuronal populations they are

expected to model. Similar to our results comparing chromatin accessibility between these two

datasets, correlation of RPKM shows a clear separation between the SN4741 cell culture model

and the *ex vivo* DA neurons (**Figure 5D**). Collectively, these results confirm that at the non-

permissive temperature, SN4741 cells are no longer rapidly dividing, neural stem cells.

However, while the transcriptional profile of these cells indicates that they are differentiating

towards cell types present in the brain, these cells do not fully possess characteristics of the MB

DA neurons they are meant to model.

DISCUSSION:

It is critically important that studies of human disease generate biologically accurate data,

whether aimed at elucidating molecular mechanisms, onset and progression, or management and

therapeutics. In the context of discovery biology or the illumination of human health and disease

mechanisms, misattribution of cellular identity, or other deviations from biological accuracy,

may result the misinterpretation of biological findings or misdirected research efforts. When

studying human disease, cellular surrogates are often used to overcome the ethical and technical

limitations of employing animal models. Therefore, it is imperative that disease-relevant insights

are predicated on robust data generated from model systems representing human biology as

accurately as possible.

Here, we demonstrate the importance of assessing *in vitro* models of disease to determine

the extent to which they can yield biologically accurate data that can be used to inform aspects of

human disease. The SN4741 cell line has been used to study neurotoxicity and therapeutic

interventions [21–27], PD-associated genetic mutation[28,29] and cell signaling and

transcriptional regulation[30–32,37], since it was initially characterized as an immortal, mouse

MB-derived cell line that differentiates into DA neurons at a non-permissive temperature[20].

However, contemporary genomic analyses have not been leveraged to characterize and evaluate

the SN4741 cell line as a suitable proxy for DA neurons in PD, until now.

We employed karyotyping, RT-qPCR, and scRNA-seq to assess the genomic stability of

these cells and determine how consistently they differentiate into DA neurons at the non-

permissive temperature. We generated bulk RNA-seq and ATAC-seq data from this cell line at

both the permissive and non-permissive temperatures, to extensively characterize this cell line

and document how transcriptional landscapes and chromatin accessibility profiles shift in

response to temperature-induced differentiation and compare to known profiles of *ex vivo* DA

neurons. Our results suggest that SN4741 is an unstable, polyploid cell line that is unlikely to be

a viable differentiation model of DA neurons; and thus, is likely not a robust proxy by which to

study MB DA neurons in the context of human phenotypes, including PD, schizophrenia,

addiction, memory, or movement disorders.

The results of karyotyping alone indicate that any data generated using SN4741 cells may

be biologically inaccurate due to extreme variability in chromosome complement and therefore,

copy number variation, between individual cells. Consequently, the results of previous studies

evaluating   neurotoxicity[21,23,24,26,27,30,32,38,67],   cellular   signaling

pathways[22,25,28,68,69], and transcriptional profiling[37] in these cells may have been unduly

influenced by the extreme imbalance in gene dosage that we found to vary from cell to cell. For

example, alpha-synuclein (*SNCA)* has been consistently implicated in PD risk[70–73],

particularly due to variants that promote α-synuclein misfolding[74] and overexpression[50] or

events that result in gene amplification [75,76]. *Snca* is present on mouse chromosome 6 and the

karyotypes generated for SN4741 cells show that chromosome 6 is triploid in most assayed cells

(**Figure 1B**). Therefore, using the SN4741 cell line to model neurodegeneration in PD may result

in inaccurate data due to an exaggerated vulnerability towards degeneration imposed by elevated

Snca copy number, by gene dosage effects of other interacting gene products in relevant

pathways, or by the structural instability of this line.

Even if this cell line could be adopted to study *Snca* overexpression/amplification,

ATAC-seq profiling of open chromatin regions in this cell line at the permissive and non-

permissive temperatures indicates that these cells do not possess chromatin accessibility profiles

similar to those of *ex vivo,* mouse E15.5 MB neurons. In PD, disease is characterized by the

degeneration of MB DA neurons, while DA neurons of the FB are spared. Therefore, the

chromatin profiles of MB DA neurons, as well as the differentially accessible regions of the

genome between MB and FB neurons, may influence the preferential vulnerability of MB

neurons in PD[50]. In the context of exploiting these chromatin profiles to study PD-associated

variability and neurotoxicity, SN4741 cells are likely a poor model, as the open chromatin

regions of these cells are not a reliable proxy for mouse E15.5 MB or FB DA neurons.

The chromatin accessibility profiles of SN4741 cells not only fail to cluster with *ex vivo*

populations of mouse MB neurons, but the transcriptional landscapes of these cells suggest that

these cells have shifted towards a more differentiated state that may be less DA than previously

thought. Examination of cell cycle markers by scRNA-seq demonstrates that SN4741 cells at the

non-permissive temperature are more differentiated than cells at the permissive temperature, as

expected[20]. GO terms for genes that are significantly downregulated at the non-permissive

temperature reinforces that these cells are no longer rapidly dividing, pluripotent stem cells.

However, RT-qPCR, scRNA-seq, and bulk RNA-seq in these cells fail to detect significant

upregulation of most key DA neuron markers in the differentiated cells, except for *Th*. *Th* is not

exclusively expressed by DA neurons at embryonic timepoints[40–42]. In fact, significantly

upregulated genes in SN4741 cells at the non-permissive temperature that overlap with GO terms

and cell cycle marker genes suggests that *Th* is the only significantly upregulated gene

overlapping with biological processes involving DA neurons. Rather, additional overlapping cell

type marker genes suggest that these cells more closely resemble immature neurons.

In parallel, we generated promoter capture (pc)Hi-C data at the non-permissive

temperature with the intention of exploring how non-coding disease-relevant variants interact

with promoters and potentially regulate gene expression in MD DA neurons. As our group is

focused on PD-associated variation, which is unlikely to act broadly in immature neurons, our

group has not analyzed the resulting data, beyond basic quality control (**Supplemental Figure**

4). While the SN4741 cells at the non-permissive temperature fail to recapitulate the

transcriptomic or chromatin state of DA neurons, it is of potential interest for follow-up studies

that they do resemble some immature neuron types. Although not analyzed by our group, we

generated output files for interaction detection, and this data has been made publicly available

for others to explore (accessible through: https://github.com/rachelboyd/SN4741_pcHiC), as it

may be useful to study genomic interactions at promoters driving an immature neuronal state.

However, the cell type best represented by SN4741 cells at the non-permissive temperature still

requires deeper characterization.

Any data generated using SN4741 cells in the context of DA neuron modeling and/or PD

must be interpreted with caution and in light of the appropriate caveats. Due to the instability and

polyploidy of this cell line, we recommend that the use of SN4741 cells for PD- related research

be re-evaluated. Future studies designed to fine-tune the classification of these cells may support

the use of SN4741 cells as a model of other neuronal or non-neuronal cells. Additionally, the

differentiation trajectory of these cells may be amenable to intervention(s) that could drive their

molecular state towards one that resembles DA neurons more closely.

CONCLUSIONS:

This study establishes a valuable precedent with broad implications across biological and

disease-related research. Prior to using SN4741 cells to study non-coding regulatory variation in

PD, we characterized this cell line to determine its suitability as a model of DA neurons in PD,

and found that these cells are unstable, polyploid cells that do not demonstrate strong molecular

characteristics of MB DA neurons. These cells express low levels of DA neuron markers, and

chromatin landscapes in differentiated SN4741 cells scarcely overlap open chromatin regions in

ex vivo mouse E15.5 midbrain neurons. We demonstrate the importance of genomic

characterization of *in vitro* model systems prior to generating data and valuable resources that

may be used to inform aspects of human disease. In future studies that utilize *in vitro* models of

any human disease, due diligence to confirm their suitability as surrogates could save time,

resources, and possibly lives, by avoiding misdirection and advancing successful therapeutic

development.

METHODS:

Cell Culture

SN4741 cells were obtained from the Ernest Arenas group at the Karolinska Institutet.

SN4741 cells were confirmed to be mycoplasma free using a MycoAlert® Mycoplasma

Detection Assay (Lonza) and were maintained in high glucose Dulbecco's Modified Eagle

Medium (DMEM; Gibco 1196502), supplemented with 1% penicillin–streptomycin and 10%

fetal bovine serum (FBS) in a humidified 5% CO_2 incubator at 37°C. Cells at 80% confluence

were passaged by trypsinization approximately every 2-3 days. To induce differentiation, 24

hours after the cells were passaged, media was replaced by DMEM supplemented with 1%

penicillin–streptomycin and 0.5% FBS at 39°C. Cells were allowed to grow and differentiate in

these conditions for 48 hours before harvesting for experimentation.

G-Band Karyotyping

At passage 21, undifferentiated SN4741 cells were sent to the WiCell Research Institute

(Madison, Wisconsin), at 40-60% confluency, for chromosomal G-band analyses. Karyotyping

was conducted on 20 metaphase spreads, at a band resolution of >230, according to the

International System for Human Cytogenetic Nomenclature.

cDNA Synthesis and RT-qPCR for DA neuron markers

RNA was extracted from both differentiated and undifferentiated cells by following the RNeasy Mini Kit (QIAGEN) protocol, as written. 1 μg of each RNA sample underwent first-strand cDNA synthesis using the SuperScript III First-Strand Synthesis System for RT-PCR (Invitrogen) according to the oligo(dT) method. qPCR was performed with Power SYBR Green Master Mix (Applied Biosystems), using primers for β-actin (*Actb*), *Foxa2, Nr4a2, Slc6a3,* and *Th* (**Table S1**). Reactions were run in triplicate under default SYBR Green Standard cycle specifications on the Viia7 Real-Time PCR System (Applied Biosystems). Normalized relative quantification and error propagation followed the data analysis and associated calculations proposed by Taylor *et al*. (2019)[77], with results normalized to *Actb*.

Single cell RNA-seq Library Preparation, Sequencing, and Alignment

Both differentiated (39°C) and undifferentiated (37°C) cells were trypsinized, and scRNA-seq libraries were generated following the Chromium 10X pipeline[78]. Four replicates at each temperature across >17,000 cells were assayed. Cell capture, cDNA generation, and library preparation were performed with the standard protocol for the Chromium Single Cell 3' V3 reagent kit. Libraries were quantified with the Qubit dsDNA High Sensitivity Assay (Invitrogen) in combination with the High Sensitivity DNA Assay (Agilent) on the Agilent 2100 Bioanalyzer. Single-cell RNA-sequencing libraries were pooled and sequenced on an Illumina NovaSeq 6000 (SP flow cells), using 2x50 bp reads per library, to a combined depth of 1.6 billion reads. The quality of sequencing was evaluated via FastQC. Paired-end reads were aligned to the mouse reference genome (mm10) using the CellRanger v3.0.1 pipeline. Unique

molecular identifier (UMI) counts were quantified per gene per cell ("cellranger count") and

aggregated ("cellranger aggr") across samples with no normalization.

Single Cell RNA-seq Analysis

Using Seurat[79](v4.2.0), cells were filtered to remove stressed/dying cells (% of reads

mapping to the mitochondria > 15%) and empty droplets or doublets (number of unique genes

detected <200 or >6,000). Cells were scored for their stage in the cell cycle using

"CellCycleScoring()" on cell cycle genes provided by Seurat ("cc.genes"). Cells were then

normalized using "SCTransform" (vst.flavor = "v2") and corrected for percent mitochondrial

reads and sequence depth. Principal component (PC) analysis was performed and a PC cut-off

was identified using "ElbowPlot()." Using this PC cutoff and a minimum distance of 0.001,

UMAP clustering was used for dimensionality reduction. Expression was plotted on a log scale

with "VlnPlot()" for a variety of proliferation and DA neuron markers.

ATAC-seq Library Preparation and Quantification

ATAC-seq libraries were generated for four replicates of undifferentiated (37°C) and

differentiated (39°C) SN4741 cells, according to the Omni-ATAC protocol[80], with minor

modifications. Aliquots of 50,000 cells were centrifuged at 2000 x g for 20 mins at 4°C, and the

resulting pellets were resuspended in 50μL of resuspension buffer. Cells were left to lyse for 3

minutes on ice before being centrifuged again at 2000 x g for 20 minutes at 4°C. The resulting

nuclei pellets were then tagmented, as written, using 50μL of transposition mixture and then

incubated at 37°C for 30 mins in a 1000 RPM thermomixer. After transposition, DNA was

purified with the Zymo DNA Clean and Concentrator -5 Kit and eluted in 21μL of elution buffer.

Pre-amplification of the transposed fragments was performed according to the conditions

outlined in the Omni-ATAC protocol[80]; however, 12 pre-amplification cycles were run in lieu

of qPCR amplification to determine additional cycles. The amplified libraries were prepared

according to the Nextera DNA Library Prep Protocol Guide, except that libraries were purified

with 40.5μL AMPure XP beads (Beckman Coulter), and 27.5μL of resuspension buffer was

added to each sample. All libraries were quantified with the Qubit dsDNA High Sensitivity

Assay (Invitrogen) in combination with the High Sensitivity DNA Assay (Agilent) on the

Agilent 2100 Bioanalyzer.

ATAC-seq Sequencing, Alignment, and Peak Calling

Libraries were sequenced on Illumina NovaSeq 6000 (SP flow cells), using 2x50 bp reads

per library, to a total combined depth of 1.6 billion reads. The quality of sequencing was

evaluated with FastQC (v0.11.9)[81] and summarized with MultiQC (v1.13)[82]. Reads were

aligned to the mouse reference genome (mm10) in local mode with Bowtie2[83](v2.4.1), using –

X 1000 to specify an increased pair distance to 1000bp. Samtools (v1.15.1)[84] and Picard

(v2.26.11; http://broadinstitute.github.io/picard/) were used to sort, deduplicate and index reads.

Peaks were called with MACS3 (v3.0.0a7; https://github.com/macs3-project/MACS)[85] and

specifying --nomodel and --nolambda for the `callpeaks()` command. Peaks overlapping mm10

blacklisted/block listed regions called by ENCODE[86,87] and in the original ATAC-seq

paper[36] were also removed with BEDTools (v2.30.0)[88].

For visualization with IGV, IGVTools (v2.15.2) was used to convert read pileups to

TDFs. The fraction of reads in peaks was calculated with DeepTools (v3.5.1)[89] using the

plotEnrichment command. The average mapping distance flag was extracted from the SAM files

with   a   custom   script   available   at   our   GitHub   repo

(https://github.com/sarahmcclymont/SN4741_ATAC/) to generate the fragment length plot.

Mouse (mm10) transcriptional start site (TSS) coordinates were downloaded from the UCSC

Genome Browser[90] (Mouse genome; mm10 assembly; Genes and Gene Predictions; RefSeq

Genes track using the table refGene), and DeepTools (v3.5.1)[89] was used to plot the pileup of

reads overtop of these TSSs. Conservation under peaks (phastCons)[91] and the genomic

distribution of peaks were calculated using the Cis-regulatory Element Annotation System

(CEAS)[92] and conservation tool of the Cistrome[93] pipeline. Analysis can be found at

http://cistrome.org/ap/u/smcclymont/h/sn4741-atac-seq-ceas-and-conservation.

ATAC-seq Normalization and Differential Peak Analysis

Each sample's peak file and BAMs were read into and analysed with DiffBind

(v3.8.1)[48]. Peaks present in two or more libraries were considered in the consensus peakset.

Reads overlapping these consensus peaks were counted with `dba.count()` specifying summits =

100, bRemoveDuplicates = TRUE. These read counts were normalized with `dba.normalize()` on

the full library size using the RLE normalization method as it is native to the DESeq2 analysis

we employed in the following `dba.analyze()` step. The volcano plot was generated using a

custom script using the output of the `dba.report()` command, where th = 1 and fold = 0 and

bCounts = T, to output all peaks regardless of their foldchange or significance. Significantly

differentially accessible regions (filtered for abs(Fold)>1 & FDR < 0.05) were submitted to

GREAT (v4.0.4)[94] and the gene ontology of the nearest gene, as identified with the

basal+extension method (where proximal was considered to be ±5kb) was assessed and plotted.

ATAC-seq comparison to *ex vivo* MB and FB DA neurons

Previously generated ATAC-seq libraries from *ex vivo* E15.5 mouse embryonic DA

neurons[50] were re-analyzed in parallel, following all the above alignment and filtering steps.

DiffBind (v3.8.1) was used to compare the samples, as above, and the R package UpSetR

(v1.4.0)[95] was used to plot the overlap of peaks between conditions.

Bulk RNA-seq Library Preparation, Sequencing, and Alignment

Cells were run through QIAshredder (Qiagen) and total RNA was extracted using the

RNeasy Mini Kit (Qiagen) according to the manufacturer's recommendations, except that RNA

was eluted twice in 50μL of water. Total RNA integrity was determined with the RNA Pico Kit

(Agilent) on the Agilent 2100 Bioanalyzer. RNA samples were sent to the Johns Hopkins

University Genetic Resources Core Facility (GRCF) for library prep (NEBNext Ultra II

directional library prep kit with poly-A selection) and sequencing. The libraries were pooled and

sequenced on an Illumina NovaSeq 6000 (SP flow cells), using 2x50 bp reads per library, to a

combined depth of 1.6 billion reads. The quality of sequencing was evaluated via FastQC.

FASTQ files were aligned to the mouse reference genome (mm10) with HISAT2[96] (v2.0.5)

and sample reads from different lanes were merged using samtools[84](v.1.10) function

"merge.". Aligned reads from individual samples were quantified against the mm10 reference

transcriptome with the subread[97–99](v1.6.1) function "featureCounts" [100], using -t exon and

-g gene_id, (**Supplemental Figure 3A**).

Bulk RNA-seq Analysis

The DESeq2 (v3.15) package was used for data quality assessment and analyses. A

DESeqDataSet of count data was generated using "DESeqDataSetFromMatrix" (design = ~

temp). The data underwent variance stabilizing transformation (vst) prior to using "plotPCA" to

visualize experimental covariates/batch effects (**Supplemental Figure 3B**) and R package

"pheatmap" (v1.0.12; https://CRAN.R-project.org/package=pheatmap) to visualize the sample-

to-sample distances (**Supplemental Figure 3C**).

Genes with an average of at least 1 read for each sample were analyzed to identify

differentially expressed (DE) genes between temperature conditions, using the function

"DESeq." P-value distribution after differential expression (DE) analysis **(Supplemental Figure**

3D) verified that the majority of called DE genes are significant. Results (alpha = 0.01) were

generated and subjected to log fold change shrinkage using the function "lfcShrink" (type =

"apeglm")[101] for subsequent visualization and ranking. The function "plotMA" was used to

generate MA plots, both before and after LFC shrinkage, to visualize the $\log_2$ fold changes

attributable to the non-permissive temperature shift over the mean of normalized counts for all

the samples in the DESeqDataSet (**Supplemental Figure 3E-F**). MA plots demonstrated that log

fold change shrinkage of the data successfully diminished the effect size of lowly expressed

transcripts with relatively high levels of variability.

Volcano plots were generated using a custom function to visualize $\log_2$ fold changes of

specific genes in the dataset. A gene was considered significantly differentially expressed if it

demonstrated an adjusted p-value < 0.01 and $|\log_2 \text{FC}| > 1.5$. These significantly differentially

expressed genes were submitted to Enrichr[51–53] for analyses within the "ontologies" and "cell

types" categories. The upregulated and downregulated gene sets were passed to STRING [66] for

analysis of protein-protein interactions and network relationships.

Bulk RNA-seq comparison to *ex vivo* MB and FB DA neurons

Read counts from the SN4741 bulk RNA-seq dataset were converted to RPKM and

compared to bulk RNA-seq data from previously generated *ex vivo* mouse embryonic DA

neurons (NCBI GEO: GSE122450;[50]). A Pearson correlation heatmap was generated using

ggplot2[102].

Promoter capture HiC library generation

PcHiC was performed as previously described[103], with minor modifications. Briefly,

SN4741 cells were cultured at the non-permissive temperature and plated at five million cells per

10cm dish. The cells were crosslinked using 1% formaldehyde, snap frozen using liquid

nitrogen, and stored at -80°C. The cells were dounce homogenized and restriction enzyme

digestion, using 400 units *HindIII-HF* overnight at 37°C. The total volume was maintained at

500μL, through addition of 1X NEBuffer 2.1. Heat inactivation was performed at 80°C for 20

minutes, and biotinylated-dCTP was used for biotin fill-in reaction. Blunt-end ligation was

performed using Thermo T4 DNA ligase, with cohesive end units maintained at 15,000 and

buffer and water volumes adjusted to ensure a total volume of 665μL was added to each Hi-C

tube. Cross-linking was performed overnight, with additional (50μL) proteinase K added for two

hours the following day. DNA purification was split across two reactions using 2mL PhaseLock

tubes, and volumes were adjusted accordingly. Each PhaseLock reaction was split again into two

vials for ethanol purification, and centrifugation at step 6.3.8 was performed at room

temperature. The pellets were dissolved in 450μL 1X TLE and transferred to a 0.5mL 30kD

Amicon Column. After washing, the column was inverted into a new container, and no

additional liquid was added to raise the volume to 100μL. All four reactions were combined, the

total volume determined, and RNAseA (1mg/mL) equal to 1% of the total volume was added for

30 minutes at 37°C.

The libraries were assessed for successful blunt-end ligation by a ClaI restriction enzyme

digest of PCR products, as previously described[103]. Biotin was removed from un-ligated ends

and DNA was sheared to a size of 200-300bp using the Covaris M220 (High setting, 35 cycles of

30s "on" and 90s "off"; vortexing/spinning down samples and changing sonicator water every 5

cycles). Size selection was performed using AMpure XP magnetic beads, as previously

described[103] except that all resuspension steps were increased by 5μL, so that 5μL could be

used for QC with the High Sensitivity DNA Assay (Agilent) on the Agilent 2100 Bioanalyzer (at

three stages: post-sonication, post-0.8x size-selection, and post-1.1x size-selection). The

remaining protocol was performed as described. Capture probes (Arbor Biosciences;

https://github.com/nbbarrientos/SN4741_pcHiC) were designed against mouse (mm10) RefSeq

transcription start sites, filtering out "XM" and "XR" annotated genes. The remaining promoters

were intersected with the *in silico* digested *HindIII* mouse genome, to retain all *HindIII*

fragments containing a promoter. Potential probes sites were assessed ±330bp of the *HindIII* cut

site on either end of the fragment and finalized probe sets were filtered using no repeats and

"strict" criteria, as defined by Arbor Biosciences. After generating a uniquely indexed HiC

library with complete Illumina adapters, probes targeting promoter containing fragments were

hybridized following Arbor Biosciences capture protocol (v4) at 65°C, 1μg DNA, and one round

of capture. The library was PCR amplified before sequencing on an Illumina NovaSeq 6000 (SP

flow cells), using 2x50 bp reads per library, to a combined depth of 1.6 billion reads.

Promoter capture HiC data analysis

Raw pcHiC reads for each replicate (n=4) were evaluated for quality via FastQC. FASTQ

files were mapped to mm10 using Bowtie2[83] (v.2.4.1) and filtered using HiCUP[104](v. 0.8).

The HiCUP pipeline was configured with the following parameters: FASTQ format (Format:

Sanger), maximum di-tag length (Longest: 700), minimum di-tag length (shortest: 50), and

filtering and alignment statistics were reported (**Supplemental Figure 4A-C**). BAM files were

generated for each replicate using samtools[84](v.1.10). DeepTools[89](v.3.5.1) before read

coverage similarities and replicate correlation was assessed using the function

"multiBamSummary" (in *bins* mode) to analyze the entire genome. A Pearson correlation

heatmap was generated using the function "plotCorrelation" (**Supplemental Figure 4D**). As a

result of high Pearson correlation coefficient among replicates ($r > 0.93$), library replicates were

combined. The CHiCAGO[105](v. 1.18.0) pipeline was used to convert the merged BAM file

into CHiCAGO format. The digested mm10 reference genome was used to generate a restriction

map file, a baited restriction map file, and the rest of required input files (.npb, .nbpb, and .poe)

required to run the CHiCAGO pipeline.

<u>AVAILABILITY OF DATA AND MATERIALS</u>:

All data and analysis pipelines are available at <u>https://github.com/rachelboyd</u>. ATAC-

sequencing, RNA-sequencing, and promoter-capture Hi-C data will be available at the Gene

Expression Omnibus (GEO) under the accession number GSEXXXXXX upon manuscript

acceptance.

<u>ABBREVIATIONS</u>:

ActB: β-actin
Aldh1a1:   Aldehyde Dehydrogenase 1 Family Member A1
ATAC-seq: Assay for Transposase-Accessible Chromatin using Sequencing
BAM: Binary Alignment and Map
Cacna1b: Calcium channel, voltage-dependent, N type, alpha 1B subunit
Ccna2:   Cyclin A2
Cdc6:   Cell division cycle 6

Cdh13:	Cadherin 13
CEAS:	Cis-Regulatory Element Annotation System
Celf5:	CUGBP Elav-Like Family Member 5
Cenpf:	Centromere protein F
Cntn2:	Contactin 2
CO2:	Carbon Dioxide
CRE:	Cis Regulatory Element
DA:	Dopaminergic
DMEM:	Dulbecco's Modified Eagle Medium
Dpysl5:	Dihydropyrimidinase-like 5
E13.5/15.5:	Embryonic Day 13.5/15.5
ENCODE:	Encyclopedia of DNA Elements
FB:	Forebrain
FBS:	Fetal Bovine Serum
FDR:	False Discovery Rate
Foxa2:	Forkhead Box A2
Gins1:	GINS complex subunit 1 (Psf1 homolog)
GO:	Gene Ontology
GRCF:	Genetics Core Research Facility
GWAS:	Genome-Wide Association study
Hdac11:	Histone deacetylase 11
Hmga2:	High Mobility Group AT-Hook 2
Id2:	Inhibitor Of DNA Binding 2
IGV:	Integrative Genomics Viewer
Il33:	Interleukin 33
Irx3:	Iroquois Homeobox 3
KEGG:	Kyoto Encyclopedia of Genes and Genomes
LFC:	Log Fold-Change
Lmx1b:	LIM Homeobox Transcription Factor 1 Beta
MB:	Midbrain
Mki67:	Marker of Proliferation Ki-67
Nes:	Nestin
Nova2:	NOVA alternative splicing regulator 2
Nr4a2:	Nuclear Receptor Subfamily 4 Group A, Member 2
Nrx1:	Neurexin 1
Ntrk1:	Neurotrophic receptor tyrosine kinase 1
OCR:	Open Chromatin Region
Olig3:	Oligodendrocyte transcription factor 3
PC(A):	Principal Component (Analysis)
pcHi-C:	Promoter-Capture Hi-C
PD:	Parkinson Disease
Pitx3:	Paired-like homeodomain 3
Ptgds :	Prostaglandin D2 synthase
(q)PCR:	(Quantitative) Polymerase Chain Reaction

QC:	Quality Control
RNA:	Ribonucleic Acid
RPKM:	Reads per kilobase of exon per million reads mapped
RT:	Reverse Transcriptase
Scn1b:	Sodium Voltage-Gated Channel Beta Subunit 1
scRNA-seq:	Single Cell RNA sequencing
Slc6a3:	Solute Carrier Family 6 Member 3
SN:	Substantia Nigra
SNCA/Snca:	Alpha-synuclein
SV40Tag:	Simian Virus 40 T antigen
TH/Th:	Tyrosine Hydroxylase
Tmem179:	Transmembrane protein 179
ts:	Temperature-Sensitive
TSS:	Transcriptional Start Site
Unc13a:	Unc-13 homolog A
vst:	Variance Stabilizing Transformation

ACKNOWLEDGEMENTS:

The authors would like to acknowledge Ernest Arenas (Karolinska Institutet), for providing

SN4741 cells, as well as the Johns Hopkins Genomics Core Research Facility (GCRF) and the

WiCell Research Institute, for providing technical services.

FUNDING:

This research, undertaken at Johns Hopkins University School of Medicine, was supported in

part by awards from the National Institutes of Health (NS62972 and MH106522) to A.S.M., by

T32 GM007814-40 to R.J.B. and N.B.B., and by the Canadian Institutes of Health Research

(DFD-181599) to R.J.B.

AUTHOR INFORMATION:

Authors and Affiliations:

McKusick-Nathans Department of Genetic Medicine, Johns Hopkins University School of

Medicine, Baltimore, Maryland, USA.

Rachel J. Boyd, Sarah A. McClymont, Nelson B. Barrientos, Paul W. Hook, William D. Law,

Rebecca J. Rose, Eric L. Waite, Dimitrios Avramopoulos, & Andrew S. McCallion

Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland,

USA.

Andrew S. McCallion

Contributions:

New data was generated by S.A.M., P.W.H., W.D.L., and E.W.L.; analyzed by R.J.B., S.A.M.,

P.W.H., N.B.B., W.D.L., and A.S.M. The manuscript was written by R.J.B., S.A.M., N.B.B.,

W.D.L., and A.S.M. Figures were created by R.J.B., S.A.M., and N.B.B. All authors reviewed

and approved the manuscript.

ETHICS DECLARATIONS:

Ethics approval and consent to participate:

Not applicable.

Consent for publication:

Not applicable.

Competing interests:

The authors declare no competing interests.

REFERENCES:

1.   Ormond KE, Mortlock DP, Scholes DT, Bombard Y, Brody LC, Faucett WA, et al.
Human germline genome editing. Am J Hum Genet. 2017;101(2):167–76.

2.   Barbosa DJ, Capela JP, de Lourdes Bastos M, Carvalho F. In vitro models for
neurotoxicology research. Toxicol Res. 2015;4(4):801–42.

3.   Hirsch C, Schildknecht S. In vitro research reproducibility: Keeping up high standards.
Front Pharmacol. 2019;10.

4.   Fisher S, Grice EA, Vinton RM, Bessling SL, Urasaki A, Kawakami K, et al. Evaluating
the biological relevance of putative enhancers using Tol2 transposon-mediated
transgenesis in zebrafish. Nat Protoc. 2006;1(3):1297–305.

5.   Gorkin DU, Lee D, Reed X, Fletez-Brant C, Bessling SL, Loftus SK, et al. Integration of
ChIP-seq and machine learning reveals enhancers and a predictive regulatory sequence
vocabulary in melanocytes. Genome Res. 2012;22(11):2290–301.

6.   Shlyueva D, Stampfel G, Stark A. Transcriptional enhancers: From properties to genome-
wide predictions. Nat Rev Genet. 2014;15(4):272–86.

7.   Gasperini M, Findlay GM, McKenna A, Milbank JH, Lee C, Zhang MD, et al.
CRISPR/Cas9-mediated scanning for regulatory elements required for HPRT1 expression
via thousands of large, programmed genomic deletions. Am J Hum Genet.
2017;101(2):192–205.

8.   Maurano MT, Humbert R, Rynes E, Thurman RE, Haugen E, Wang H, et al. Systematic
localization of common disease-associated variation in regulatory DNA. Science (1979).
2012;337(6099):1190–5.

9.   Kheradpour P, Ernst J, Melnikov A, Rogov P, Wang L, Zhang X, et al. Systematic
dissection of regulatory motifs in 2000 predicted human enhancers using a massively
parallel reporter assay. Genome Res. 2013;23(5):800–11.

10. Shim S, Kwan KY, Li M, Lefebvre V, Nenad &, Estan Sˇ. Cis-regulatory control of
corticospinal system development and evolution. Nature. 2012;486:74–9.

11. Schoenfelder S, Javierre BM, Furlan-Magaril M, Wingett SW, Fraser P. Promoter Capture
Hi-C: High-resolution, genome-wide profiling of promoter interactions. JoVE (Journal of
Visualized Experiments). 2018;2018(136):e57320.

12. Schaub MA, Boyle AP, Kundaje A, Batzoglou S, Snyder M. Linking disease associations
with regulatory information in the human genome. Genome Res. 2012;22(9):1748–59.

13. Ernst J, Kheradpour P, Mikkelsen TS, Shoresh N, Ward LD, Epstein CB, et al. Mapping
and analysis of chromatin state dynamics in nine human cell types. Nature.
2011;473(7345):43–9.

14. Boyle EA, Li YI, Pritchard JK. An expanded view of complex traits: From polygenic to
omnigenic. Cell. 2017;169(7):1177–86.

15. Lee D, Gorkin DU, Baker M, Strober BJ, Asoni AL, McCallion AS, et al. A method to
predict the impact of regulatory variants from DNA sequence. Nat Genet.
2015;47(8):955–61.

16. Fearnley JM, Lees AJ. Ageing and Parkinson's disease: Substantia nigra regional
selectivity. Brain. 1991;114:2283–301.

17. Marras C, Beck JC, Bower JH, Roberts E, Ritz B, Ross GW, et al. Prevalence of
Parkinson's disease across North America. NPJ Parkinsons Dis. 2018;4(1):21.

18. Dorsey ER, Bloem BR. The Parkinson pandemic - A call to action. JAMA Neurol.
2018;75(1):9–10.

19. Ferrari E, Cardinale A, Picconi B, Gardoni F. From cell lines to pluripotent stem cells for
modelling Parkinson's Disease. J Neurosci Methods. 2020;340:108741.

20. Son JH, Chun HS, Joh TH, Cho S, Conti B, Lee JW. Neuroprotection and neuronal
differentiation studies using substantia nigra dopaminergic cells derived from transgenic
mouse embryos. J Neurosci. 1999;19(1):10.

21. Chang J, Zhang X Le, Yu H, Chen J. Downregulation of RTN1-C attenuates MPP+-
induced neuronal injury through inhibition of mGluR5 pathway in SN4741 cells. Brain
Res Bull. 2019;146:1–6.

22. Chen J, Li M, Zhou X, Xie A, Cai Z, Fu C, et al. Rotenone-induced neurodegeneration is
enabled by a p38-Parkin-ROS signaling feedback loop. J Agric Food Chem.
2021;69(46):13942–52.

23. Guiney SJ, Adlard PA, Lei P, Mawal CH, Bush AI, Finkelstein DI, et al. Fibrillar α-
synuclein toxicity depends on functional lysosomes. J Biol Chem. 2020;295(51):17497–
513.

24. Chun HS, Gibson GE, Degiorgio LA, Zhang H, Kidd VJ, Son JH. Dopaminergic cell
death induced by MPP+, oxidant and specific neurotoxicants shares the common
molecular mechanism. J Neurochem. 2001;76(4):1010–21.

25. Chun HS, Lee H, Son JH. Manganese induces endoplasmic reticulum (ER) stress and
activates multiple caspases in nigral dopaminergic neuronal cells, SN4741. Neurosci Lett.
2001;316(1):5–8.

26. Zeng W, Zhang W, Lu F, Gao L, Gao G. Resveratrol attenuates MPP+-induced
mitochondrial dysfunction and cell apoptosis via AKT/GSK-3β pathway in SN4741 cells.
Neurosci Lett. 2017;637:50–6.

27. Cai Z, Zeng W, Tao K, Lu F, Gao G, Yang Q. Myricitrin alleviates MPP+-induced
mitochondrial dysfunction in a DJ-1-dependent manner in SN4741 cells. Biochem
Biophys Res Commun. 2015;458(2):227–33.

28. Mao K, Chen J, Yu H, Li H, Ren Y, Wu X, et al. Poly (ADP-ribose) polymerase 1
inhibition prevents neurodegeneration and promotes α-synuclein degradation via
transcription factor EB-dependent autophagy in mutant α-synucleinA53T model of
Parkinson's disease. Aging Cell. 2020;19(6).

29. Gui C, Ren Y, Chen J, Wu X, Mao K, Li H, et al. p38 MAPK-DRP1 signaling is involved
in mitochondrial dysfunction and cell death in mutant A53T α-synuclein model of
Parkinson's disease. Toxicol Appl Pharmacol. 2020;388:114874.

30. Dong Y, Xiong J, Ji L, Xue X. MiR-421 aggravates neurotoxicity and promotes cell death
in Parkinson's disease models by directly targeting MEF2D. Neurochem Res.
2021;46(2):299–308.

31. Yoo MS, Chun HS, Son JJ, DeGiorgio LA, Kim DJ, Peng C, et al. Oxidative stress
regulated genes in nigral dopaminergic neuronal cells: correlation with the known
pathology in Parkinson's disease. Mol Brain Res. 2003;110(1):76–84.

32. Wang B, Cai Z, Lu F, Li C, Zhu X, Su L, et al. Destabilization of survival factor MEF2D
mRNA by neurotoxin in models of Parkinson's disease. J Neurochem. 2014;130(5):720–
8.

33. Dekker J, Rippe K, Dekker M, Kleckner N. Capturing chromosome conformation.
Science. 2002;295(5558):1306–11.

34. Lieberman-Aiden E, Van Berkum NL, Williams L, Imakaev M, Ragoczy T, Telling A, et
al. Comprehensive mapping of long-range interactions reveals folding principles of the
human genome. Science. 2009;326(5950):289–93.

35. Mortazavi A, Williams BA, McCue K, Schaeffer L, Wold B. Mapping and quantifying
mammalian transcriptomes by RNA-Seq. Nat Methods. 2008;5(7):621–8.

36. Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. Transposition of native
chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding
proteins and nucleosome position. Nat Methods. 2013;10(12):1213–8.

37. Nissim-Eliraz E, Zisman S, Schatz O, Ben-Arie N. Nato3 integrates with the Shh-Foxa2
transcriptional network regulating the differentiation of midbrain dopaminergic neurons. J
Mol Neurosci. 2013;51:13–27.

38. Fishman-Jacob T, Reznichenko L, Youdim MBH, Mandel SA. A sporadic Parkinson
disease model via silencing of the ubiquitin-proteasome/E3 ligase component SKP1A. J
Biol Chem. 2009;284(47):32835–46.

39. Weihe E, Depboylu C, Schu□z B, Schäfer MKH, Eiden LE. Three types of tyrosine
hydroxylase-positive CNS neurons distinguished by dopa decarboxylase and VMAT2 co-
expression. Cell Mol Neurobiol. 2006;26(4–6):659–78.

40. Jonakait GM, Markey KA, Goldstein M, Dreyfus CF, Black IB. Selective expression of
high-affinity uptake of catecholamines by transiently catecholaminergic cells of the rat
embryo: studies in vivo and in vitro. Dev Biol. 1985;108(1):6–17.

41. Cochard P, Goldstein M, Black IB. Ontogenetic appearance and disappearance of tyrosine
hydroxylase and catecholamines in the rat embryo. Proc Natl Acad Sci U S A.
1978;75(6):2986–90.

42. Asmus SE, Parsons S, Landis SC. Developmental changes in the transmitter properties of
sympathetic neurons that innervate the periosteum. J Neurosci. 2000;20(4):1495–504.

43. Ladd AN, Charlet-B. N, Cooper TA. The CELF family of RNA binding proteins is
implicated in cell-specific and developmentally regulated alternative splicing. Mol Cell
Biol. 2001;21(4):1285–96.

44. Nishino J, Kim I, Chada K, Morrison SJ. Hmga2 promotes neural stem cell self-renewal in
young, but not old, mice by reducing p16Ink4a and p19Arf expression. Cell.
2008;135(2):227.

45. Park HJ, Hong M, Bronson RT, Israel MA, Frankel WN, Yun K. Elevated Id2 expression
results in precocious neural stem cell depletion and abnormal brain development. Stem
Cells. 2013;31(5):1010.

46. Dou Z, Son JE, Hui CC. Irx3 and Irx5 - Novel regulatory factors of postnatal
hypothalamic neurogenesis. Front Neurosci. 2021;15:1447.

47. Reid CA, Leaw B, Richards KL, Richardson R, Wimmer V, Yu C, et al. Reduced
dendritic arborization and hyperexcitability of pyramidal neurons in a Scn1b-based model
of Dravet syndrome. Brain. 2014;137:1701–15.

48. Stark R, Brown G. DiffBind: Differential binding analysis of ChIP-Seq peak data.
http://bioconductor.org/packages/devel/bioc/vignettes/DiffBind/inst/doc/DiffBind.pdf

49. Li H, Liu Y, Gu Z, Li L, Liu Y, Wang L, et al. p38 MAPK-MK2 pathway regulates the
heat-stress-induced accumulation of reactive oxygen species that mediates apoptotic cell
death in glial cells. Oncol Lett. 2018;15(1):775.

50. McClymont SA, Hook PW, Soto AI, Reed X, Law WD, Kerans SJ, et al. Parkinson-
associated SNCA enhancer variants revealed by open chromatin in mouse dopamine
neurons. Am J Hum Genet. 2018;103(6):874–92.

51. Xie Z, Bailey A, Kuleshov M v., Clarke DJB, Evangelista JE, Jenkins SL, et al. Gene set
knowledge discovery with Enrichr. Curr Protoc. 2021;1(3):e90.

52. Chen EY, Tan CM, Kou Y, Duan Q, Wang Z, Meirelles G v., et al. Enrichr: Interactive
and collaborative HTML5 gene list enrichment analysis tool. BMC Bioinformatics.
2013;14.

53. Kuleshov M v., Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, et al. Enrichr:
A comprehensive gene set enrichment analysis web server 2016 update. Nucleic Acids
Res. 2016;44(W1):W90–7.

54. Franzén O, Gan LM, Björkegren JLM. PanglaoDB: a web server for exploration of mouse
and human single-cell RNA sequencing data. Database. 2019;2019(1):46.

55. Zeng L, Zhang P, Shi L, Yamamoto V, Lu W, Wang K. Functional impacts of NRXN1
knockdown on neurodevelopment in stem cell models. PLoS One. 2013;8(3).

56. Kaplan DR, Miller FD. Neurotrophin signal transduction in the nervous system. Curr Opin
Neurobiol. 2000;10(3):381–91.

57. Reddy-Alla S, Böhme MA, Reynolds E, Beis C, Grasskamp AT, Mampell MM, et al.
Stable positioning of Unc13 restricts synaptic vesicle fusion to defined release sites to
promote synchronous neurotransmission. Neuron. 2017;95(6):1350-1364.e12.

58. Storm R, Cholewa-Waclaw J, Reuter K, Bröhl D, Sieber M, Treier M, et al. The bHLH
transcription factor Olig3 marks the dorsal neuroepithelium of the hindbrain and is
essential for the development of brainstem nuclei. Development. 2009;136(2):295–305.

59. Sung HY, Chen WY, Huang HT, Wang CY, Chang S bin, Tzeng SF. Down-regulation of
interleukin-33 expression in oligodendrocyte precursor cells impairs oligodendrocyte
lineage progression. J Neurochem. 2019;150(6):691–708.

60. Liu H, Hu Q, D'Ercole AJ, Ye P. Histone Deacetylase 11 regulates oligodendrocyte-
specific gene expression and cell development in OL-1 oligodendroglia cells. Glia.
2009;57(1):1.

61. Sakry D, Yigit H, Dimou L, Trotter J. Oligodendrocyte precursor cells synthesize
neuromodulatory factors. PLoS One. 2015;10(5).

62. Yam CH, Fung TK, Poon RYC. Cyclin A in cell cycle control and cancer. Cellular and
Mol Life Sci. 2002;59(8):1317–26.

63. Borlado LR, Méndez J. CDC6: from DNA replication to cell cycle checkpoints and
oncogenesis. Carcinogenesis. 2008;29(2):237–43.

64. Ma L, Zhao X, Zhu X. Mitosin/CENP-F in mitosis, transcriptional control, and
differentiation. J Biomed Sci. 2006;13(2):205–13.

65. Nagahama Y, Ueno M, Miyamoto S, Morii E, Minami T, Mochizuki N, et al. PSF1, a
DNA replication factor expressed widely in stem and progenitor cells, drives tumorigenic
and metastatic properties. Cancer Res. 2010;70(3):1215–24.

66. Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, et al. The STRING
database in 2021: customizable protein-protein networks, and functional characterization
of user-uploaded gene/measurement sets. Nucleic Acids Res. 2021;49(D1):D605–12.

67. Chun HS, Lee H, Son JH. Manganese induces endoplasmic reticulum (ER) stress and
activates multiple caspases in nigral dopaminergic neuronal cells, SN4741. Neurosci Lett.
2001;316(1):5–8.

68. Choi KC, Kim SH, Ha JY, Kim ST, Son JH. A novel mTOR activating protein protects
dopamine neurons against oxidative stress by repressing autophagy related cell death. J
Neurochem. 2010;112(2):366–76.

69. Bryja V, Čajánek L, Grahn A, Schulte G. Inhibition of endocytosis blocks Wnt signalling
to β-catenin by promoting dishevelled degradation. Acta Physiologica. 2007;190(1):55–
61.

70. Nalls MA, Blauwendraat C, Vallerga CL, Heilbron K, Bandres-Ciga S, Chang D, et al.
Identification of novel risk loci, causal insights, and heritable risk for Parkinson's disease:
a meta-analysis of genome-wide association studies. Lancet Neurol. 2019;18(12):1091–
102.

71. Blauwendraat C, Nalls MA, Singleton AB, Blauwendraat C, Singleton AB, Nalls MA.
The genetic architecture of Parkinson's disease. Lancet Neurol. 2020;19:170–8.

72. Chang D, Nalls MA, Hallgrímsdóttir IB, Hunkapiller J, Brug M van der, Cai F, et al. A
meta-analysis of genome-wide association studies identifies 17 new Parkinson's disease
risk loci. Nat Genet. 2017;49(10):1511–6.

73. Blauwendraat C, Heilbron K, Vallerga CL, Bandres Ciga S, von Coelln R, Pihlstrøm L,
et al. Parkinson's disease age at onset genome wide association study: Defining
heritability, genetic loci, and α synuclein mechanisms. Movement Disorders.
2019;34(6):866–75.

74. Polymeropoulos MH, Lavedan C, Leroy E, Ide SE, Dehejia A, Dutra A, et al. Mutation in
the α-synuclein gene identified in families with Parkinson's disease. Science.
1997;276(5321):2045–7.

75. Singleton AB, Farrer M, Johnson J, Singleton A, Hague S, Kachergus J, et al. Alpha-
synuclein locus triplication causes Parkinson's disease. Science. 2003;302(5646):841.

76. Ibáñez P, Bonnet AM, Débarges B, Lohmann E, Tison F, Pollak P, et al. Causal relation
between alpha-synuclein gene duplication and familial Parkinson's disease. The Lancet.
2004;364(9440):1169–71.

77. Taylor SC, Nadeau K, Abbasi M, Lachance C, Nguyen M, Fenrich J. The ultimate qPCR
experiment: Producing publication quality, reproducible data the first time. Trends
Biotechnol. 2019;37(7):761–74.

78. Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, et al. Massively
parallel digital transcriptional profiling of single cells. Nat Commun. 2017;8(1):1–12.

79.   Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell
878   transcriptomic data across different conditions, technologies, and species. Nat Biotechnol.
879   2018;36(5):411–20.

880   80.   Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA, Vesuna S,
881   et al. An improved ATAC-seq protocol reduces background and enables interrogation of
882   frozen tissues. Nat Methods. 2017;14(10):959–62.

883   81.   Andrews S. FASTQC: A quality control tool for high throughput sequence data. 2010.
884   http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

885   82.   Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for
886   multiple tools and samples in a single report. Bioinformatics. 2016;32(19):3047–8.

887   83.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
888   2012;9(4):357.

889   84.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
890   Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078.

891   85.   Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-based
892   analysis of ChIP-Seq (MACS). Genome Biol. 2008;9(9):1–9.

893   86.   Amemiya HM, Kundaje A, Boyle AP. The ENCODE blacklist: Identification of
894   problematic regions of the genome. Sci Rep. 2019;9(1).

895   87.   Dunham I, Kundaje A, Aldred SF, Collins PJ, Davis CA, Doyle F, et al. An integrated
896   encyclopedia of DNA elements in the human genome. Nature. 2012;489(7414):57–74.

897   88.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
898   features. Bioinformatics. 2010;26(6):841–2.

899   89.   Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, et al. deepTools2: A
900   next generation web server for deep-sequencing data analysis. Nucleic Acids Res.
901   2016;44(W1):W160–5.

902   90.   Karolchik D, Hinricks AS, Furey TS, Roskin KM, Sugnet CW, Haussler D, et al. The
903   UCSC Table Browser data retrieval tool. Nucleic Acids Res. 2004;32(Database
904   issue):D493.

905   91.   Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, Rosenbloom K, et al.
906   Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes.
907   Genome Res. 2005;15(8):1034–50.

908   92.   Shin H, Liu T, Manrai AK, Liu SX. CEAS: cis-regulatory element annotation system.
909   Bioinformatics. 2009;25(19):2605–6.

910   93.   Liu T, Ortiz JA, Taing L, Meyer CA, Lee B, Zhang Y, et al. Cistrome: An integrative
911   platform for transcriptional regulation studies. Genome Biol. 2011;12(8):1–10.

912   94.   McLean CY, Bristor D, Hiller M, Clarke SL, Schaar BT, Lowe CB, et al. GREAT
913   improves functional interpretation of cis-regulatory regions. Nat Biotechnol.
914   2010;28(5):495–501.

915   95.   Conway JR, Lex A, Gehlenborg N. UpSetR: an R package for the visualization of
916   intersecting sets and their properties. Bioinformatics. 2017;33(18):2938–40.

917   96.   Kim D, Langmead B, Salzberg SL. HISAT: A fast spliced aligner with low memory
918   requirements. Nat Methods. 2015;12(4):357–60.

919   97.   Gentleman RC, Carey VJ, Bates DM, Bolstad B, Dettling M, Dudoit S, et al.
920   Bioconductor: Open software development for computational biology and bioinformatics.
921   Genome Biol. 2004;5(10).

922   98.   Huber W, Carey VJ, Gentleman R, Anders S, Carlson M, Carvalho BS, et al.
923   Orchestrating high-throughput genomic analysis with Bioconductor. Nat Methods.
924   2015;12(2):115–21.

925   99.   Liao Y, Smyth GK, Shi W. The Subread aligner: fast, accurate and scalable read mapping
926   by seed-and-vote. Nucleic Acids Res. 2013;41(10):e108–e108.

927   100.   Liao Y, Smyth GK, Shi W. Sequence analysis featureCounts: an efficient general purpose
928   program for assigning sequence reads to genomic features. Bioinformatics.
929   2014;30(7):923–30.

930   101.   Zhu A, Ibrahim JG, Love MI. Heavy-tailed prior distributions for sequence count data:
931   removing the noise and preserving large differences. Bioinformatics. 2019;35(12):2084–
932   92.

933   102.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York;
934   2016. https://ggplot2.tidyverse.org.

935   103.   Belaghzal H, Dekker J, Gibcus JH. Hi-C 2.0: An optimized Hi-C procedure for high-
936   resolution genome-wide mapping of chromosome conformation. Methods. 2017;123:56–
937   65.

938   104.   Wingett S, Ewels P, Furlan-Magaril M, Nagano T, Schoenfelder S, Fraser P, et al. HiCUP:
939   Pipeline for mapping and processing Hi-C data. F1000Res. 2015;4.

940   105.   Cairns J, Freire-Pritchett P, Wingett SW, Várnai C, Dimond A, Plagnol V, et al.
941   CHiCAGO: Robust detection of DNA looping interactions in Capture Hi-C data. Genome
942   Biol. 2016;17(1):1–17.

Figure 1. Characterizing the genomic stability and differentiation consistency of the temperature sensitive SN4741 cell line. A) A representative karyogram of SN4741 cells, indicating structural instability (M; marker chromosomes) and unstable triploidy. **B)** A stacked bar plot summarizing the aneuploidy frequency of each chromosome over 20 SN4741 karyotypes. **C)** Assaying expression of dopaminergic neuron markers by RT-qPCR indicates that *Foxa2, Nr4a2*, and *Slc6a3* remain at similar expression levels, but *Th* expression increases, when SN4741 cells are shifted from the permissive temperature (37°C) to the non-permissive temperature (39°C). **D)** UMAP plot of scRNA-seq at the permissive and non-permissive temperatures indicates that cells at each temperature are transcriptionally distinct. **E)** Analysis of scRNA-seq data demonstrates that shifting the cells to the non-permissive temperature is accompanied by a shift in cell cycle stage from G2M and S phases to primarily G1 phase. **F)** Violin plots generated with scRNA-seq data show that *Mki67*, a marker of cellular proliferation, and *Nes*, a neural stem cell marker, are both expressed at the permissive temperature (37°C), with little to no expression at the non-permissive temperature (39°C). **G)** Violin plots generated with scRNA-seq data show that transcripts associated with immature neurons are upregulated when SN4741 cells are shifted to the non-permissive temperature. **H)** Violin plots generated with scRNA-seq data show that expression of DA neural markers, *Aldh1a1, Foxa2, Lmx1b, Nr4a2, Pitx3, Slc6a3*, and *Th*, remain at similar levels when SN4741 cells are shifted to the non-permissive temperature.

Figure 2: Changes in chromatin accessibility suggest a reduction in potency at the non-permissive temperature. A, B) Replicates are highly similar within temperature conditions, with the majority of peaks present in all four replicates. **C)** The two temperatures share 58,251 regions of open chromatin but do not overlap completely. **D)** Principal component analysis resolves the two temperatures on the first principal component. **E)** Differential accessibility analysis identifies 5,055 differentially accessible regions, with 2,654 preferentially open in the permissive temperature (37°C) and 2,401 preferentially accessible at the non-permissive temperature (39°C). **F)** Gene ontology (GO) of genes adjacent regions that are preferentially open at the permissive temperature are associated with regulation of the cell cycle and negative regulation of differentiation, as is appropriate for this temperature. **G)** Gene ontology of genes adjacent regions that are preferentially open at the non-permissive temperature are associated with a variety of differentiation fates (blood vessels, muscle cells, cartilage/chondrocytes). Additionally, two of the top gene ontology terms relate to the p38 MAPK cascade, which has been found to be activated as a cellular response to heat stress.

Figure 3: Chromatin accessibility of SN4741 cells do not resemble *ex vivo* dopaminergic neurons. A) SN4741 samples are highly correlated with each other but very poorly correlate with the open chromatin landscape of either midbrain (MB) or forebrain (FB) embryonic mouse dopaminergic neurons. **B**) Principal component analysis shows a clear separation between the *ex vivo* and *in vitro* samples along PC1, representing 86% of the variance. **C**) An upset plot and associated Venn diagram quantify the overlap of peaks between the four conditions and show the poor relationship between the SN4741 cells and the *ex vivo* mouse dopaminergic neurons. Most peaks are specific to a single cell type/temperature or are restricted to either the *ex vivo* or *in vitro* samples. Few peaks are specifically shared between the non-permissive temperature and the *ex vivo* samples; for example, there are just 183 peaks that are shared exclusively by the MB dopaminergic neurons and the SN4741 cells at the non-permissive temperature. **D**) A genome track showing the normalized read pile up and called consensus peaks in each of the cell types/temperatures at the key dopaminergic neuron specification gene, *Th*. The chromatin accessibility is largely similar within *ex vivo* or *in vitro* cells but bear little resemblance to each other.

Figure 4. Gene Ontology and Differential Expression Analysis of Bulk RNA-seq Data: A) Top 10 GO terms for downregulated DE genes in SN4741 cells at the non-permissive temperature, followed by GO terms of interest (below dotted line). Terms were evaluated using Combined Score Ranking = (p-value computed using the Fisher exact test)*(z-score computed by assessing the deviation from the expected rank), based on enrichment of DE genes that overlap with Enrichr input genes for each term (the end of each bar). **B)** Top 10 predicted cell types based on downregulated DE genes in SN4741 cells at the non-permissive temperature, followed by predicted cell types of interest (below dotted line). Terms were evaluated using Combined Score Ranking = (p-value computed using the Fisher exact test)*(z-score computed by assessing the deviation from the expected rank), based on enrichment of DE genes that overlap with PanglaoDB input genes for each term (the end of each bar). **C)** Top 10 GO terms for downregulated DE genes in SN4741 cells at the non-permissive temperature. **D)** Top 10 predicted cell types based on upregulated DE genes in SN4741 cells at the non-permissive temperature. **E)** Volcano plot of −log10 adjusted p-value versus $\log_2$ fold change with DESeq2 after lfc shrinkage, contrasting the fold change in expression of SN4741 cells at 39°C, using SN4741 cells at 37°C as reference. Red points = genes that are statistically differentially expressed (adjusted p-value < 0.01, $|\log_2 FC| > 1.5$). Blue points = Overlapping immature neuron marker genes. **F)** Blue points = Overlapping immature neuron marker genes **G)** Blue points = Overlapping oligodendrocyte marker genes. **H)** Blue points = Overlapping DA neuron marker genes.

Figure 5. Validation of gene ontology and comparison to bulk-RNA-seq from ex vivo DA neurons. A) A bar chart showing normalized bulk RNA-seq read counts from genes upregulated at 39°C that overlap the "immature neurons" predicted cell type. **B)** A bar chart showing normalized bulk RNA-seq read counts from genes upregulated at 39°C that overlap the "oligodendrocyte" predicted cell type. **C)** A bar chart showing normalized bulk RNA-seq read counts from genes downregulated at 39°C that overlap the "pluripotent stem cell" predicted cell type. **D)** A Pearson correlation heatmap comparing the transcriptomes of SN4741 cells at 37°C and 39°C to midbrain (MB) or forebrain (FB) embryonic mouse dopaminergic neurons.