1 RIPK3 promotes brain region-specific interferon signaling and restriction of tick-borne flavivirus
2 infection
3

4 Marissa Lindman[a], Juan P Angel[a], Irving Estevez[a], Nydia P Chang[a], Tsui-Wen Chou[a], Micheal
5 McCourt[a], Colm Atkins[a], and Brian P. Daniels[a#]
6
7 [a]Department of Cell Biology and Neuroscience, Rutgers University, Piscataway, NJ, USA
8
#Correspondence:
Brian Daniels
604 Allison Road
Room B314
Piscataway, NJ 08854
(848) 445-2709
b.daniels@rutgers.edu

16
17
18

Abstract

Innate immune signaling in the central nervous system (CNS) exhibits many remarkable specializations that vary across cell types and CNS regions. In the setting of neuroinvasive flavivirus infection, neurons employ the immunologic kinase receptor-interacting kinase 3 (RIPK3) to promote an antiviral transcriptional program, independently of the traditional function of this enzyme in promoting necroptotic cell death. However, while recent work has established roles for neuronal RIPK3 signaling in controlling mosquito-borne flavivirus infections, including West Nile virus and Zika virus, functions for RIPK3 signaling in the CNS during tick-borne flavivirus infection have not yet been explored. Here, we use a model of Langat virus (LGTV) encephalitis to show that RIPK3 signaling is specifically required in neurons of the cerebellum to control LGTV replication and restrict disease pathogenesis. This effect did not require the necroptotic executioner molecule mixed lineage kinase domain like protein (MLKL), a finding similar to previous observations in models of mosquito-borne flavivirus infection. However, control of LGTV infection required a unique, region-specific dependence on RIPK3 to promote expression of key antiviral interferon-stimulated genes (ISG) in the cerebellum. This RIPK3-mediated potentiation of ISG expression was associated with robust cell-intrinsic restriction of LGTV replication in cerebellar granule cell neurons. These findings further illuminate the complex roles of RIPK3 signaling in the coordination of neuroimmune responses to viral infection, as well as provide new insight into the mechanisms of region-specific innate immune signaling in the CNS.

Importance

Interactions between the nervous and immune systems are very carefully orchestrated in order to protect the brain and spinal cord from immune-mediated damage, while still maintaining protective defenses against infection. These specialized neuro-immune interactions have been shown to vary significantly across regions of the brain, with innate antiviral signaling being particularly strong in the cerebellum, although the reasons for this are poorly understood. Here, we show a specialized adaptation of programmed cell death signaling that uniquely protects the cerebellum from tick-borne flavivirus infection. These findings provide important new insight into the molecular mechanisms that promote the uniquely robust antiviral immunity of the cerebellum. They also provide new clues into the pathogenesis of tick-borne encephalitis, a zoonosis of significant global concern.

Introduction

Flaviviruses are a family of positive sense RNA viruses which include several notable pathogens associated with neuroinvasive infection in humans, including West Nile virus (WNV), Zika virus (ZIKV), and Japanese Encephalitis virus (1). While nearly all major flaviviruses are transmitted by mosquito vectors, a small but significant number of flaviviruses are transmitted by ticks, including Tick-borne encephalitis virus (TBEV) and its close relatives that together make up a single TBEV serocomplex. Tick borne encephalitis is a significant and growing threat to public health, particularly in Europe and northern Asia, where TBEV constitutes the most prevalent tick-borne zoonotic disease (2-4). Notably, some TBEV strains elicit mortality rates up to 40% in humans (5), underscoring the urgent need to better understand the mechanisms underlying the pathogenesis of tick-borne flavivirus infections.

Effective control of flavivirus infection in the central nervous system (CNS) requires robust innate immune signaling in neural cells, particularly neurons, which are the predominantly infected cell type in most cases of flavivirus encephalitis (6-9). Effective type I interferon (IFN) signaling is of particular importance for innate control of viral replication in neurons (10-12). Notably, differences in type I IFN signaling across neural cell types and brain regions are associated with differential susceptibility to flavivirus infection. For example, previous reports suggest that the enhanced type I IFN signaling observed in hindbrain regions compared to the forebrain is an underlying determinant of the enhanced susceptibility of forebrain regions to WNV infection (12, 13). However, the unique signaling mechanisms that promote differential interferon-mediated control of viral infection in the hindbrain have not been extensively characterized.

A potential regulator of neuronal interferon signaling during flavivirus infection is receptor interacting protein kinase-3 (RIPK3). RIPK3 is an enzyme traditionally associated with necroptosis, a form of lytic programmed cell death (14). Necroptosis occurs via the RIPK3-dependent activation of mixed lineage kinase domain like protein (MLKL), which forms oligomeric pore complexes that induce cellular lysis (15). However, many recent studies have identified complex roles for RIPK3 signaling in the coordination of inflammation, including the regulation of inflammatory transcriptional responses that occur independently of necroptosis (16-24). We and others have demonstrated that RIPK3 signaling in neurons is of particular importance for the control of neurotropic viral infections, as neuronal RIPK3 promotes a robust antimicrobial transcriptional program, including many interferon stimulated genes (ISGs), that restricts viral infection without inducing neuronal necroptosis (16, 17). Other recent studies have identified unexpected roles for RIPK3 in the regulation of type I IFN signaling, via mechanisms which include the regulation of pattern recognition receptor signaling and protein kinase-R (PKR)-mediated stabilization of *Ifnb* mRNA (18, 19).

In this study, we interrogated roles for RIPK3 in controlling tick-borne flavivirus infection. To do so, we used Langat virus (LGTV), a naturally attenuated member of the TBEV serocomplex that can be studied under BSL2 containment. *Ripk3*[-/-] mice exhibited enhanced clinical disease following subcutaneous LGTV infection, while *Mlkl*[-/-] mice were indistinguishable from littermate controls, suggesting a necroptosis-independent function for RIPK3 in restricting LGTV pathogenesis. Notably, *Ripk3*[-/-] mice exhibited increased viral burden in the cerebellum, along with diminished expression of inflammatory chemokines and ISGs in the cerebellum, but not the cerebral cortex. *In vitro* analysis of cultured primary cortical and cerebellar cell types showed that pharmacologic inhibition of RIPK3 resulted in enhanced LGTV replication in cerebellar granule cell neurons but not in cortical neurons or in astrocytes derived from either

brain region. Transcriptional profiling showed that RIPK3 signaling was uniquely required for the full induction of ISG expression in cerebellar granule cell neurons, demonstrating a previously unknown, region-specific function for RIPK3 in coordinating innate antiviral immunity within the CNS.

Results

RIPK3 controls LGTV pathogenesis independently of MLKL and peripheral immunity

To assess the role of RIPK3 in controlling LGTV pathogenesis, we subcutaneously infected *Ripk3*[-/-] mice, along with heterozygous littermate controls, with 3×10^4 plaque forming units (pfu) of the Malaysian LGTV strain TP21. We note that *Ripk3*[+/-] animals do not exhibit haploinsufficiency and are routinely used as littermate controls in studies of this pathway (25-27). Control animals exhibited limited mortality following LGTV infection (Figure 1A), consistent with previous reports (28, 29). Mice lacking *Ripk3* expression exhibited a modest increase in mortality, although this difference did not reach statistical significance (Figure 1A). However, a higher proportion of *Ripk3*[-/-] mice did develop clinical signs of neurologic disease, including paresis or full hindlimb paralysis, by 14 days post infection (dpi) (Figure 1B), and this difference persisted to at least 21 dpi. These data suggested that although most *Ripk3*[-/-] animals were able to survive subcutaneous LGTV infection, RIPK3 was nevertheless important for the control of neuropathogenesis in this model.

To better understand this phenotype, we first assessed whether RIPK3 was required for early control of systemic infection. Spleens of infected mice exhibited low levels of LGTV RNA that were not impacted by *Ripk3* expression (Figure 1C). To test whether *Ripk3*[-/-] mice exhibited any deficiencies in peripheral immune responses, we performed flow cytometric analysis of major immune cell subsets in the spleens of infected animals at 8 dpi. *Ripk3*[-/-] animals exhibited similar frequencies (Figure 1D) and total numbers (Figure 1E-F) of CD4 and CD8 T cells among all splenocytes compared to littermate controls, as well as similar rates of CD44 expression (a key T cell activation marker) across both subsets (Figure 1G-H). Numbers of B cells (Figure 1I) and natural killer (NK) cells (Figure 1J) were also similar between genotypes. In the myeloid compartment, we observed similar numbers of CD11c[+] MHCII[+] dendritic cells (Figure 1K) between genotypes, as well as similar numbers of myeloid subsets expressing F4/80 (Figure 1L), Ly6G (Figure 1M), and Ly6C (Figure 1N). Both CD11c[+] MHCII[+] and F4/80[+] antigen presenting cell subsets also exhibited similar rates of expression of the costimulation signal CD80 between genotypes (Figure 1O-P). These data suggest that *Ripk3*[-/-] mice mounted normal peripheral immune responses to subcutaneous LGTV challenge, similar to our previous observations with WNV and ZIKV (16, 17). Thus, the increased pathogenesis observed in mice lacking *Ripk3* was unlikely to arise from a failure in peripheral virologic control.

A potential mechanism by which RIPK3 signaling might restrict LGTV pathogenesis is through the induction of necroptosis in infected cells. We thus tested whether loss of the necroptotic executioner molecule MLKL would impact disease course following subcutaneous LGTV infection. Notably, *Mlkl*[-/-] mice exhibited no difference in either survival or development of clinical disease signs compared to littermate controls (Figure 2A-B). We saw similarly that *Mlkl*[-/-] did not exhibit altered splenic viral burden at 8dpi (Figure 2C). Flow cytometric analysis also revealed essentially identical numbers and frequencies of all major immune cell subsets in the spleen at this time point (Figure 2C-P). Multistep growth curve analysis also demonstrated that neither RIPK3 nor MLKL impacted the low levels of LGTV replication observed in primary

156 leukocyte cultures, including bone marrow derived macrophages and dendritic cells
157 (Supplemental Figure 1A-B). These data suggest that MLKL, and therefore necroptosis, is not a
158 major contributor to peripheral virologic control or overall disease pathogenesis in the setting of
159 LGTV infection, and thus that RIPK3 exerts its protective effect in this model through an
160 alternative mechanism.
161
162 **RIPK3 is required for CNS-intrinsic restriction of LGTV infection**
163
Because we did not observe differences in peripheral virologic control in *Ripk3-/-* mice, we
next questioned whether RIPK3 acted in a CNS-intrinsic manner to limit LGTV infection. To
assess this, we next used an intracranial infection route in order to assess local effects of RIPK3
signaling on LGTV pathogenesis. *Ripk3-/-* mice exhibited accelerated and enhanced mortality
compared to littermate controls following intracranial infection (Figure 3A). *Ripk3*-deficient mice
also exhibited worsened clinical disease prior to death, as evidenced by earlier and more
dramatic weight loss following infection (Figure 3B). In contrast, *Mlkl-/-* mice were
indistinguishable from littermate controls in terms of overall mortality (Figure 3C) and weight loss
(Figure 3D) following intracranial infection. These data further supported the idea that RIPK3
restricts LGTV neuropathogenesis via CNS-intrinsic mechanisms, independently of necroptosis.
174
175 **RIPK3 promotes neuronal chemokine expression in a region-specific manner following**
176 **LGTV infection**
177
We and other previously showed that neuronal RIPK3 signaling was required for the
expression of key inflammatory chemokines that served to restrict WNV pathogenesis by
coordinating the recruitment of leukocytes into the infected CNS. We thus questioned whether
RIPK3 also promotes chemokine expression in the CNS during LGTV infection. Surprisingly,
transcriptional profiling in the cerebral cortex of *Ripk3-/-* mice following subcutaneous LGTV
infection revealed no differences in expression of major chemokines compared to littermate
controls (Figure 4A). However, we did observe significantly diminished chemokine responses in
cerebellar tissues derived from *Ripk3-/-* animals (Figure 4B). To understand which cell types
were driving this region-specific deficit in chemokine expression, we next cultured primary
neurons and astrocytes derived specifically from either cerebral cortex or cerebellum and
infected with LGTV, with or without a small molecule inhibitor of RIPK3 (GSK 872). Consistent
with our *in vivo* findings, blockade of RIPK3 in cerebral cortical neurons did not impact
chemokine expression following LGTV infection (Figure 4C). In contrast, infected cerebellar
granule cell neuron cultures exhibited significantly diminished chemokine expression when
RIPK3 was inhibited by GSK 872 (Figure 4D). Notably, we did not observe a dependence on
RIPK3 for the expression of chemokines in astrocytes derived from either region (Figure 4E-F).
These data suggest that RIPK3 serves an unexpected, region-specific transcriptional function in
neurons of the cerebellum during neuroinvasive LGTV infection.
196
197 **RIPK3 is not required for immune cell recruitment to the LGTV-infected CNS**
198
We next questioned whether diminished chemokine expression in the cerebellum of
Ripk3-/- mice would result in a failure to recruit antiviral leukocytes into this brain region. We thus
performed flow cytometric analysis of leukocytes derived from either cerebral cortex or
cerebellum following subcutaneous LGTV infection. Remarkably, we saw no evidence of
changes in lymphocyte recruitment in either brain region of *Ripk3-/-* mice compared to littermate
controls on either 6 or 8 dpi (Figure 5A). This lack of difference extended across all major
CD45hi infiltrating leukocyte subsets, including CD4+ and CD8+ T cells (Figure 5B-C), NK cells

(Figure 5D), CD11c+ MHCII+ dendritic cells (Figure 5E) and myeloid subsets expressing F4/80
(Figure 5F), Ly6G (Figure 5G), and Ly6C (Figure 5H). We similarly did not observe differences
in numbers of CD45^{int} microglia (Figure 5I) in either region. These data suggested that, despite
significant differences in the expression of major leukocyte chemoattractants in the cerebellum,
differences in immune cell recruitment did not account for the increased pathogenesis observed
in *Ripk3*$^{-/-}$ mice during LGTV infection.
212
213 **RIPK3 promotes cell-intrinsic restriction of LGTV replication in cerebellar neurons**
214
Given these observations, we next questioned whether *Ripk3*$^{-/-}$ mice fail to control LGTV
infection due to impaired innate immune restriction of LGTV replication. Assessment of viral
burdens in brains of *Ripk3*$^{-/-}$ mice following subcutaneous LGTV infection revealed that *Ripk3*$^{-/-}$
mice exhibited significantly elevated CNS viral titers, particularly in the cerebellum, at both 8 and
12 dpi (Figure 6A). In contrast, *Mlkl*$^{-/-}$ exhibited no such difference in viral burden in either brain
region (Figure 6B). Differences in viral burden did not appear to be linked to deficits in blood-
brain barrier integrity, as both *Ripk3*$^{-/-}$ mice and littermate controls exhibited similar levels of
sodium fluorescein extravasation into the CNS following infection (Figure 6C). We thus
questioned whether RIPK3 was required for cell-intrinsic restriction of viral replication in
susceptible CNS cell types. Multistep growth curve analysis in primary CNS cells revealed that
pharmacologic inhibition of RIPK3 had no effect on LGTV replication in neurons derived from
cerebral cortex (Figure 6D). In contrast, inhibition of RIPK3 significantly enhanced LGTV
replication in primary cerebellar granule cell neurons cultures (Figure 6E). This effect was
unique to neurons, as GSK 872 treatment had no impact on LGTV replication in primary
astrocytes derived from either brain region (Figure 6F-G). Together, these data suggested that
the enhanced pathogenesis observed in *Ripk3*$^{-/-}$ mice was due to a specific failure to control
infection in neurons of the cerebellum, resulting in enhanced overall CNS viral burden.
232
Our previous observation of diminished chemokine expression in cerebellar neurons
derived from *Ripk3*$^{-/-}$ mice suggested that these cells may exhibit broader deficits in innate
immune signaling, resulting in poor control of LGTV replication. We, therefore, next questioned
whether interferon signaling was perturbed in the cerebellum of mice lacking RIPK3 expression.
Transcriptional profiling in brain tissues following subcutaneous LGTV infection revealed that,
indeed, the cerebella of *Ripk3*$^{-/-}$ mice exhibited diminished expression of many interferon
stimulated genes (ISGs) known to be critical for control of flavivirus replication (30-35), including
Ifit1, Isg15, Mx1, Mx2, Oas1b, and *Rsad2,* while this phenotype was not observed in the
cerebral cortex (Figure 7A-B). Similar analyses in primary cell cultures confirmed that cerebellar
granule cell neurons, but not neurons derived from cerebral cortex, exhibited diminished
expression of ISGs when RIPK3 signaling was blocked via GSK 872 treatment (Figure 7C-D). In
contrast, we observed little to no impact of RIPK3 blockade on ISG expression in astrocytes
derived from either brain region (Figure 7E-F). Together, these data demonstrate that RIPK3
signaling is required for the robust induction of type I IFN responses in neurons of the
cerebellum, which is required for cell-intrinsic restriction of LGTV replication.
248
249 **Discussion**
250
251　　　　Our findings identify a previously unknown function for RIPK3 in the coordination of brain
252 region-specific innate immunity. The study of regional differences in neuroimmune signaling is a
253 growing field, and there is accumulating evidence to suggest that resident neural cells exhibit
254 differential responses to viral infection and cytokine stimulation across distinct anatomical
255 regions of the CNS (36-39). Neurons and astrocytes in the cerebellum, in particular, have been

shown to exhibit higher responsiveness to stimulation by type-I IFN, as well as to express higher basal levels of pathogen sensor molecules compared to other brain regions, suggesting a key evolutionary importance of innate antiviral defense in this tissue (12, 13). This regional difference in type I IFN signaling appears to underlie, at least in part, the relatively lower susceptibility of the cerebellum to flavivirus infection compared to susceptible regions of the forebrain, such as the cerebral cortex and hippocampus. However, the molecular mechanisms that determine the enhanced innate immune signaling observed in the cerebellum remain poorly understood. Our study suggests that RIPK3 signaling is required for the robust induction of ISG expression in cerebellar neurons during LGTV infection, although ongoing work is needed to understand the specific signaling interactions that mediate this effect.

Previous studies have described a highly complex interplay between RIPK3 and type I IFN signaling that varies significantly by cell type and disease model (17-19, 40). It is relatively clear that type I IFN signaling is capable of activating RIPK3 through various mechanisms, resulting in necroptosis and/or necroptosis-independent transcriptional activation (40-43). However, how RIPK3 operates *upstream* of (or synergistically with) type I IFN signaling to influence expression of ISGs is less clear. We and others have shown that ISG expression is significantly diminished in a variety of settings when RIPK3 signaling is ablated (17, 18), including in cerebellar granule cell neurons during LGTV infection in this study. One possible explanation for this effect is RIPK3-mediated activation of NF-κB, a transcription factor strongly associated with RIPK signaling with known roles in potentiating type I IFN signaling and ISG expression (22, 44-46). We and others also previously showed that RIPK3 activation in cortical neurons following ZIKV infection leads to interferon regulatory factor 1 (IRF1) activation, which was required for expression of at least a subset of RIPK3-induced genes in that setting, although this effect is likely indirect, as IRF1 is not a known RIPK3 substrate (17). Additional work will be needed to fully characterize the regulatory mechanisms that are invoked in the interplay between RIPK3 and type I IFN signaling in the CNS.

Our study also further expands our understanding of the necroptosis-independent functions for RIPK3 signaling in the CNS. Many studies have now firmly established the importance of RIPK3 in promoting host defense through mechanisms independent of its canonical role in necroptosis (16-21). However, these necroptosis-independent functions appear to vary significantly by disease state, including CNS infection with distinct neuroinvasive flaviviruses (47, 48). We and others previously showed that the primary role for RIPK3 in restricting WNV encephalitis was the induction of chemokine expression and the recruitment of antiviral leukocytes into the infected CNS (16). Notably, while we did observe RIPK3-mediated chemokine expression in the cerebellum during LGTV infection, this chemokine expression was apparently dispensable for CNS immune cell recruitment. Instead, the transcriptional activation of antiviral effector genes, including ISGs, was required for cell-intrinsic restriction of LGTV replication in neurons, a phenotype more similar to our findings with ZIKV (17), although we did not observe evidence for a regional specification of this response during ZIKV infection. In contrast to these observations, Bian and colleagues have observed quite distinct phenotypes in a model of JEV encephalitis, wherein both RIPK3 and MLKL appeared to exacerbate rather than restrict disease pathogenesis (49, 50). RIPK3 also appeared to *suppress* rather than promote ISG expression in JEV infected neurons. The factors that determine such distinct outcomes of RIPK3 signaling across this family of closely related viruses are mysterious and are the subject of ongoing investigation by our laboratory and others.

Materials and Methods

Mouse lines

$Ripk3^{-/-}$ (51) $Mlkl^{-/-}$ (52) mouse lines were bred and housed under specific-pathogen free conditions in Nelson Biological Laboratories at Rutgers University. $Ripk3^{-/-}$ mice were generously provided by Genentech, Inc. Wild-type C57BL/6J mice were either obtained commercially (Jackson Laboratories) or bred in-house. Mice used for subcutaneous infections were 5 weeks old; mice used for intracranial infections were 8-15 weeks old.

Virus and titer determination

Langat virus strain TP21 was used throughout the study. Founder stocks were obtained from the World Reference Center for Emerging Viruses and Arboviruses (WRCEVA). Laboratory stocks were generated using Vero E6 cells (ATCC, #CRL-1586) and frozen at -80°C until needed. Virus titers were determined by plaque assay on Vero E6 cells. Cells were maintained in DMEM (Corning #10-013-CV) supplemented with 10% Heat Inactivated FBS (Gemini Biosciences #100-106), 1% Penicillin–Streptomycin-Glutamine (Gemini Biosciences #400-110), 1% Amphotericin B (Gemini Biosciences #400–104), 1% Non-Essential Amino Acids (Cytiva, #SH30238.01), and 1% HEPES (Cytiva, #SH30237.01). Plaque assay media was composed of 1X EMEM (Lonza # 12-684F) supplemented with 2% Heat Inactivated FBS (Gemini Biosciences #100-106), 1% Penicillin–Streptomycin-Glutamine (Gemini Biosciences, #400-110), 1% Amphotericin B (Gemini Biosciences #400-104), 1% Non-Essential Amino Acids (Cytiva, #SH30238.01), and 1% HEPES (Cytiva, SH30237.01), 0.75% Sodium Bicarbonate (VWR, #BDH9280) and 0.5% Methyl Cellulose (VWR, #K390). Plaque assays were developed at 5dpi by removal of overlay media and staining/fixation using 10% neutral buffered formalin (VWR, #89370) and 0.25% crystal violet (VWR, #0528). Plaque assays were performed by adding 100uL of serially diluted sample for 1 hour at 37°C to 12-well plates containing 200,000 Vero E6 cells per well. Plates were further incubated with plaque assay media at 37°C and 5% CO_2 for 5 days. Medium was removed from the wells and replaced with fixative containing crystal violet for approximately 20-30 minutes. Plates were washed repeatedly in H_2O and allowed to dry before counting visible plaques.

Mouse infections and tissue harvesting

Isoflurane anesthesia was used for all procedures. Mice were inoculated subcutaneously (50uL) with 3×10^4 PFU or injected intracranially (10uL) with 50 PFU of LGTV-TP21 using insulin syringes (BD Medical, #BD-329461). At appropriate times post infection, mice underwent cardiac perfusions with 30 mL cold sterile 1X phosphate-buffered saline (PBS). Extracted tissues were weighed and homogenized using 1.0 mm diameter zirconia/silica beads (Biospec Products, #11079110z) in sterile PBS for plaque assay or TRI Reagent (Zymo, #R2050-1) for gene expression analysis. Homogenization was performed in an Omni Beadrupter Elite for 2 sequential cycles of 20 s at a speed of 4 m/s.

Primary cell infections

Cortical and cerebellar astrocytes were harvested from P1-P2 pups and cortical neurons were harvested at E13.5-E15.5. Tissues were dissociated using the Neural Dissociation Kit (T) following manufacturer's instructions (Miltenyi, #130-093-231). Astrocytes were expanded in AM-a medium (ScienCell, #1831) supplemented with 10% FBS in fibronectin-coated cell culture flasks and seeded into plates coated with 20 µg/mL Poly-L-Lysine (Sigma-Aldrich, #9155) before experiments. Neurons were seeded into PLL-coated cell culture treated plates and grown in Neurobasal Plus + B-27 supplement medium (Thermo-Fisher Scientific, #A3582901) prior to use in experiments 7-9 days in vitro (DIV). Mouse cerebellar granule cells from C57BL/6 mice

355 (ScienCell, # M1530-57) were seeded into cell culture treated plates coated with 10 ug/mL Poly-
356 D-Lysine (ThermoFisher, #A3890401) containing prewarmed Neuronal Medium (ScienCell,
357 #1521) following manufacturer recommendations and used for experiments 6 DIV.
358
Macrophages and dendritic cells were isolated from bone marrow of euthanized mice.
Femurs were isolated and bone marrow pushed out using a sterile needle and syringe loaded
with RPMI supplemented with 10% FBS, 1% Penicillin–Streptomycin-Glutamine, 1% HEPES,
1% Glutamax (ThermoFisher, #35050061). Bone marrow was plated into non-cell-culture
treated 10cm petri dishes in 8mL supplemented RPMI medium containing either 20ng/mL
recombinant M-CSF (Peprotech, #315-02) or 20ng/mL recombinant GM-CSF (Peprotech, #315-
03) and 20ng/mL IL-4 (Peprotech, #214-14) for differentiation into macrophages or dendritic
cells, respectively. Cells were fed with additional medium containing the appropriate cytokines
four days later and used for experiments at 6-7 DIV. Cells were seeded into cell-culture treated
dishes prior to experimentation. For viral replication determination, all cultures were infected
with LGTV TP21 at an MOI of 0.01. For the purpose of qRT-PCR, cortical and cerebellar neuron
cultures were infected at a MOI of 0.5, while astrocyte cultures were harvested from infections
using an MOI of 0.01.
372
Quantitative real-time PCR
Total RNA from harvested tissues was extracted using Zymo Direct-zol RNA Miniprep kit,
as per manufacturer instructions (Zymo, #R2051). Total RNA extraction from cultured cells,
cDNA synthesis, and subsequent qRT-PCR were performed as previously described (22, 53).
Cycle threshold (CT) values for analyzed genes were normalized to CT values of the
housekeeping gene 18 S ($CT_{Target} − CT_{18S} = \Delta CT$). Data from primary cell culture experiments
were further normalized to baseline control values ($\Delta CT_{experimental} − \Delta CT_{control} = \Delta\Delta CT$ (DDCT)). A
list of primers used in this study can be found in **Supplemental Table 1**.
381
Flow Cytometry
The cerebella and cerebral cortices of mouse brains were dissected from freshly
perfused mice and placed into tubes containing 1X PBS. Brain tissues were incubated with
10mL buffer containing 0.05% Collagenase Type I (Sigma-Aldrich, #C0130), 10ug/mL DNase I
(Sigma-Aldrich, #D4527) and 10mM HEPES (Cytiva, #SH30237.01) in 1X Hanks' Balanced Salt
Solution (VWR, #02-1231-0500) for one hour at room temperature under constant rotation.
Brain tissues were transferred to a 70um strainer on 50mL conical tubes and mashed through
the strainer using the plunger of 3-5mL syringes. Tissue was separated in 8 mL 37% Isotonic
Percoll (Percoll: Cytiva, #17-0891-02; RPMI 1640: Corning, #10-040-CV, supplemented with 5%
FBS) by centrifugation at 1200xg for 30 minutes with a slow break. The myelin layer and
supernatant were discarded. Leukocytes were incubated in 1X RBC Lysis Buffer (Tonbo
Biosciences, #TNB-4300-L100) for 10 minutes at room temperature. Cells were centrifuged and
resuspended in FACS buffer composed of 1X PBS, 2% sodium azide and 5% FBS. Samples
were transferred into a U-bottomed 96-well plate. Leukocytes were blocked with 2% normal
mouse serum and 1% FcX Block (BioLegend, #101320) in FACS buffer for 30 minutes at 4°C
prior to being stained with fluorescently conjugated antibodies to CD3e (Biolegend, clone 17A2),
CD44 (Biolegend, clone IM7), CD19 (Biolegend, clone 6D5), CD8a (Biolegend, clone 53-6.7),
CD4 (Biolegend, clone RM4-5), CD45.2 (Biolegend, clone 104), MHC-II (Biolegend, clone
M5/114.15.2), NK1.1 (Biolegend, clone PK136), CD11c (Biolegend, clone N418), F4/80
(Biolegend, clone BM8), CD11b (Biolegend, clone M1/70), Ly6G (Biolegend, clone 1A8), Ly6C
(Biolegend, clone HK1.4), CD80 (Biolegend, clone 16-10A1), and Zombie NIR (Biolegend,
#423105). Leukocytes were stained for 30 minutes at 4C prior to washing in FACS buffer and
fixation with 1% PFA in PBS (ThermoFisher, #J19943-K2). Data collection and analysis were

performed using a Cytek Northern Lights Cytometer (Cytek, Fremont, California) and FlowJo software (Treestar). Data were normalized using a standard bead concentration counted by the cytometer with each sample (ThermoFisher, #C36950). Spleens were crushed between two slides, filtered through a 70um cell strainer, and washed with FACS buffer. Isolated splenocytes were incubated with 1X RBC Lysis Buffer as done for leukocytes isolated from the brain prior to blocking and staining.

In vivo assessment of blood brain barrier permeability

In vivo assessment of blood brain barrier permeability was carried out as described (54). Mice were injected intraperitoneally with 100uL of 100mg/mL fluorescein sodium salt (Sigma, #F6377) dissolved in sterile 1X PBS. After 45 minutes, blood was collected followed by cardiac perfusion. Tissues were dissected and homogenized in 1X PBS as described above. Serum and supernatant from homogenized tissues were incubated overnight at 4°C with 2% Trichloroacetic acid solution (Sigma, #T0699) at a 1:1 dilution. Precipitated protein was pelleted by 10 minutes of centrifugation at 2,823xg at 4°C. Supernatants were diluted with borate buffer, pH 11 (Sigma, #1094621000) to achieve a neutral pH. Fluorescein emission at 538nm was measured for samples in an optically clear black-walled 96-well plate (Corning, #3904) using a SpectraMax iD3 plate reader (Molecular Devices, San Jose, CA). Tissue fluorescence values were standardized against plasma values for individual mice.

Statistical analysis

Normally distributed data were analyzed using appropriate parametric tests: two-way analysis of variance (ANOVA) with Sidak's correction for multiple comparisons and Log-rank (Mantel-Cox) test for survival comparison, both using GraphPad Prism Software v8 (GraphPad Software, San Diego, CA). Chi square tests for comparison of clinical disease signs was performed using Excel v2211 (Microsoft). $P < 0.05$ was considered statistically significant.

Acknowledgements

This work was supported by R01 NS120895 (to BPD). JPA and IE were supported by NIH Supplement to Promote Diversity (R01 NS120895-S1 and NS120895-S2). NPC was supported by F31 NS124242.

440
441

Figure 1. RIPK3 limits LGTV pathogenesis independently of peripheral immunity

(A-B) Survival analysis (A) and presentation of clinical signs of disease (B) in *Ripk3*[-/-] mice and littermate controls following subcutaneous inoculation with 3×10^4 PFU LGTV TP21. Data are pooled from two experiments. (C) *Ripk3*[-/-] and littermate control mice were infected subcutaneously with LGTV TP21. On indicated days following infection, splenic viral burden was measured via qRT-PCR. Data was normalized against a standard curve of known viral titers to generate plaque-forming unit (PFU) equivalents. Data for each day post infection are pooled from 2-3 experiments. LOD, limit of detection. (D-P) *Ripk3*[-/-] and littermate control mice were infected subcutaneously with LGTV TP21 for 8 days prior to harvesting splenocytes and profiling leukocytes by flow cytometry. (D) Representative flow cytometry plots showing CD8+ and CD4+ T cells among CD3+ leukocytes in the spleen. Numbers represent percentage of cells in each gate relative to total plotted cells. (E-F) Numbers of CD8+ T cells (E) and CD4+ T cells (F) among CD3+ leukocytes. (G-H) Percentage of CD44+ cells among CD8+ T cells (G) and CD4+ T cells (H). (I-N) Numbers of CD19+ B cells (I), CD11b+ NK1.1+ Natural Killer cells (J), CD11c+ MHC-II+ dendritic cells (K), CD45high CD11b+ F4/80+ macrophages (L), CD11b+ Ly6G+ neutrophils (M), and CD45high CD11b+ Ly6C+ monocytes (N) among total leukocytes in the spleen. (O-P) Percentage of CD80+ cells among CD11c+ MHC-II+ dendritic cells (O) and CD11b+ F4/80+ macrophages (P). ns, not significant. *p<0.05, **p < 0.01, ***p < 0.001. Error bars represent SEM.

Figure 2. MLKL signaling does not influence Langat virus pathogenesis.

(A-B) Survival analysis (A) and presentation of clinical signs of disease (B) in *Mlkl⁻/⁻* mice and littermate controls following subcutaneous inoculation with 3X10⁴ PFU LGTV TP21. Data are pooled from two experiments. (C) *Mlkl⁻/⁻* and littermate control mice were infected subcutaneously with LGTV TP21. On indicated days following infection, splenic viral burden was measured via qRT-PCR. Data was normalized against a standard curve of known viral titers to generate plaque-forming unit (PFU) equivalents. Data for each day post infection are pooled from 2-3 experiments. LOD, limit of detection. (D-P) *Mlkl⁻/⁻* and littermate control mice were infected subcutaneously with LGTV TP21 for 8 days prior to harvesting splenocytes and profiling leukocytes by flow cytometry. (D) Representative flow cytometry plots showing CD8+ and CD4+ T cells among CD3+ leukocytes in the spleen. Numbers represent percentage of cells in each gate relative to total plotted cells. (E-F) Numbers of CD8+ T cells (E) and CD4+ T cells (F) among CD3+ leukocytes. (G-H) Percentage of CD44+ cells among CD8+ T cells (G) and CD4+ T cells (H). (I-N) Numbers of CD19+ B cells (I), CD11b+ NK1.1+ Natural Killer cells (J), CD11c+ MHC-II+ dendritic cells (K), CD45high CD11b+ F4/80+ macrophages (L), CD11b+ Ly6G+ neutrophils (M), and CD45high CD11b+ Ly6C+ monocytes (N) among total leukocytes in the spleen. (O-P) Percentage of CD80+ cells among CD11c+ MHC-II+ dendritic cells (O) and CD11b+ F4/80+ macrophages (P). ns, not significant. *p<0.05, **p < 0.01, ***p < 0.001. Error bars represent SEM.

479
480 **Figure 3. RIPK3, but not MLKL, restricts Langat virus pathogenesis following intracranial infection.**
481 Survival and body weight analysis from $Ripk3^{-/-}$ (A-B) and $Mlkl^{-/-}$ (C-D) mice and their respective
482 littermate controls following intracranial inoculation with 50 PFU LGTV TP21. Data are pooled from two
483 (A-B) or three (C-D) experiments. ns, not significant. *p<0.05, **p < 0.01, ***p < 0.001.
484

485
Figure 4. RIPK3 promotes chemokine expression in the cerebellum during LGTV encephalitis.
486
(A-B) *Ripk3* [-/-] and littermate control mice were infected subcutaneously with LGTV TP21. At 8dpi
487
488 cerebral cortical (A) and cerebellar tissues (B) were harvested and assayed for chemokine transcripts via
489 qRT-PCR. (C-F) *Ccl2* and *Cxcl10* expression in wildtype (C57BL/6J) cultures of primary cortical neurons (C),
490 cerebellar granule cell neurons (D), cortical astrocytes (E), and cerebellar astrocytes (F) following 2-hour
491 pretreatment with GSK872 or vehicle and 24h infection with 0.5 (C-D) or 0.01 (E-F) MOI LGTV TP21,
492 measured via qRT-PCR. ns, not significant. *p<0.05, **p < 0.01, ***p < 0.001. Error bars represent SEM.
493
494

Figure 5. Leukocyte recruitment to the CNS occurs independently of RIPK3 signaling during LGTV encephalitis.

(A-I) *Ripk3* [-/-] and littermate control mice were infected subcutaneously with LGTV TP21. Cerebral cortical and cerebellar tissues were harvested and leukocytes isolated for flow cytometric profiling at indicated days post infection (dpi). (A) Representative flow cytometry plots showing CD8+ and CD4+ T cells among CD3+ leukocytes in the brain. Numbers represent percentage of cells in each gate relative to total plotted cells. (B-I) Numbers of CD8+ T cells (B), CD4+ T cells (C), CD11b+ NK1.1+ natural killer cells (D), CD11c+ MHC-II+ dendritic cells (E), CD45[high] CD11b+ F4/80+ macrophages (F), CD11b+ Ly6G+ neutrophils (G), CD45[high] CD11b+ Ly6C+ monocytes (H), and CD45.2[lo] CD11b+ microglia (I) among total brain leukocytes. No comparisons are statistically significant.

507
508
Figure 6. RIPK3 limits LGTV replication in cerebellar granule cell neurons.
(A-B) *Ripk3* [-/-] (A) or *Mlkl* [-/-] (B) mice and littermate controls were infected subcutaneously with LGTV
TP21. At 8 or 12 days post infection (dpi), viral loads in cerebral cortical and cerebellar tissues were
determined by plaque assay. Data are pooled from 2-3 independent experiments. (C) *Ripk3* [-/-] and
littermate control mice were subcutaneously infected with LGTV TP21. BBB permeability was measured
at 8 dpi by detection of sodium fluorescein accumulation in tissue homogenates derived from cerebral
cortex or cerebellum. Data represent individual brain fluorescence values normalized to serum sodium
fluorescein concentration. Individual mouse values were then normalized to the mean values for
uninfected controls. (D-G) Multistep growth curve analysis following infection with 0.01 MOI LGTV TP21
in cortical neurons (D), cerebellar granule cell neurons (E), cortical astrocytes (F), and cerebellar
astrocytes (G). n=3 (cerebellar granule cell neurons) or 4 (astrocytes and cortical neurons) for growth
curve experiments. ns, not significant. *p<0.05, **p < 0.01, ***p < 0.001. Error bars represent SEM.
521
522
523

524
Figure 7. **RIPK3 promotes ISG expression in cerebellar granule cell neurons.**
(A-B) *Ripk3* [-/-] and littermate control mice were infected subcutaneously with LGTV TP21. Transcriptional
expression of indicated genes was assessed via qRT-PCR in cerebral cortical (A) and cerebellar (B) tissues
at 8dpi. (C-D) Transcriptional expression of ISGs in wildtype (C57BL/6J) cultures of primary cortical
neurons (C), cerebellar granule cell neurons (D), cortical astrocytes (E), and cerebellar astrocytes (F)
following 2-hour pretreatment with GSK872 or vehicle and 24-hour infection with 0.5 (C-D) or 0.01 (E-F)
MOI LGTV TP21, measured via qRT-PCR. ns, not significant. *p<0.05, **p < 0.01, ***p < 0.001. Error bars
represent SEM.
533

534
535
536 Supplemental Figure 1: **Neither RIPK3 nor MLKL is required for restriction of LGTV replication in bone**
537 **marrow-derived macrophages and dendritic cells**
538 (A-B) Multistep growth curve analysis following infection with 0.01 MOI LGTV TP21 in primary
539 macrophages (BMDMs) (A) and dendritic cells (BMDCs) (B) cultured from bone marrow of C57BL/6J
540 (WT), $Ripk3^{-/-}$, or $Mlkl^{-/-}$ mice. (n=4) No comparisons are statistically significant.
541
542

543   Supplemental Table 1:
544

Primer sequences for qRT-PCR

Gene	Direction	Sequence (5'-3')
18S	Forward	CTTAGAGGGACAAGTGGCG
18S	Reverse	ACGCTGAGCCAGTCAGTGTA
Ccl2	Forward	TGGCTCAGCCAGATGCAGT
Ccl2	Reverse	TTGGGATCATCTTGCTGGTG
Ccl3	Forward	CCAAGTCTTCTCAGCGCCAT
Ccl3	Reverse	TCCGGCTGTAGGAGAAGCAG
Ccl4	Forward	TCTTGCTCGTGGCTGCCT
Ccl4	Reverse	GGGAGGGTCAGAGCCCA
Ccl5	Forward	CCTGCTGCTTTGCCTACCTCTC
Ccl5	Reverse	ACACACTTGGCGGTTCCTTCGA
Cxcl1	Forward	ATCCAGAGCTTGAAGGTGTTG
Cxcl1	Reverse	GTCTGTCTTCTTTCTCCGTTACTT
Cxcl10	Forward	CCCACGTGTTGAGATCATTG
Cxcl10	Reverse	CACTGGGTAAAGGGGAGTGA
Ifit1	Forward	CCCAGAGAACAGCTACCACC
Ifit1	Reverse	TGTGAAGTGACATCTCAGCTGA
Isg15	Forward	TGCCTGCAGTTCTGTACCAC
Isg15	Reverse	AGTGCTCCAGGACGGTCTTA
LGTV	Forward	GGAACTAGGCCTTGCAGAAT
LGTV	Reverse	TGTTCTCCATTGTCGGGTTAG
Mx1	Forward	ACTATGAGGAGAAGGTGCGG
Mx1	Reverse	ACTTTGCCTCTCCACTCCTC
Mx2	Forward	GCCACGTTCCCTTGATCATC
Mx2	Reverse	AGCCAGCTTAACCAGGGAAT
Oas1b	Forward	TTCTACGCCAATCTCATCAGTG
Oas1b	Reverse	GGTCCCCCAGCTTCTCCTTAC
Rsad2	Forward	TCAAAAGCTGAGGAGGTGGTG
Rsad2	Reverse	TAGGAGGCACTGGAAAACCTTC

545
546
547
548

References

1. Schultz JS, Sparks H, Beckham JD. 2021. Arboviral central nervous system infections. Curr Opin Infect Dis 34:264-271.
2. Ruzek D, Avsic Zupanc T, Borde J, Chrdle A, Eyer L, Karganova G, Kholodilov I, Knap N, Kozlovskaya L, Matveev A, Miller AD, Osolodkin DI, Overby AK, Tikunova N, Tkachev S, Zajkowska J. 2019. Tick-borne encephalitis in Europe and Russia: Review of pathogenesis, clinical features, therapy, and vaccines. Antiviral Res 164:23-51.
3. Riccardi N, Antonello RM, Luzzati R, Zajkowska J, Di Bella S, Giacobbe DR. 2019. Tick-borne encephalitis in Europe: a brief update on epidemiology, diagnosis, prevention, and treatment. Eur J Intern Med 62:1-6.
4. Abdiyeva K, Turebekov N, Yegemberdiyeva R, Dmitrovskiy A, Yeraliyeva L, Shapiyeva Z, Nurmakhanov T, Sansyzbayev Y, Froeschl G, Hoelscher M, Zinner J, Essbauer S, Frey S. 2020. Vectors, molecular epidemiology and phylogeny of TBEV in Kazakhstan and central Asia. Parasit Vectors 13:504.
5. Beaute J, Spiteri G, Warns-Petit E, Zeller H. 2018. Tick-borne encephalitis in Europe, 2012 to 2016. Euro Surveill 23.
6. Fares M, Cochet-Bernoin M, Gonzalez G, Montero-Menei CN, Blanchet O, Benchoua A, Boissart C, Lecollinet S, Richardson J, Haddad N, Coulpier M. 2020. Pathological modeling of TBEV infection reveals differential innate immune responses in human neurons and astrocytes that correlate with their susceptibility to infection. J Neuroinflammation 17:76.
7. Szretter KJ, Daffis S, Patel J, Suthar MS, Klein RS, Gale M, Jr., Diamond MS. 2010. The innate immune adaptor molecule MyD88 restricts West Nile virus replication and spread in neurons of the central nervous system. J Virol 84:12125-38.
8. Iwasaki Y, Zhao JX, Yamamoto T, Konno H. 1986. Immunohistochemical demonstration of viral antigens in Japanese encephalitis. Acta Neuropathol 70:79-81.
9. Klein RS, Lin E, Zhang B, Luster AD, Tollett J, Samuel MA, Engle M, Diamond MS. 2005. Neuronal CXCL10 directs CD8+ T-cell recruitment and control of West Nile virus encephalitis. J Virol 79:11457-66.
10. Lindqvist R, Upadhyay A, Overby AK. 2018. Tick-Borne Flaviviruses and the Type I Interferon Response. Viruses 10.
11. Samuel MA, Diamond MS. 2005. Alpha/beta interferon protects against lethal West Nile virus infection by restricting cellular tropism and enhancing neuronal survival. J Virol 79:13350-61.
12. Cho H, Proll SC, Szretter KJ, Katze MG, Gale M, Jr., Diamond MS. 2013. Differential innate immune response programs in neuronal subtypes determine susceptibility to infection in the brain by positive-stranded RNA viruses. Nat Med 19:458-64.
13. Daniels BP, Jujjavarapu H, Durrant DM, Williams JL, Green RR, White JP, Lazear HM, Gale M, Jr., Diamond MS, Klein RS. 2017. Regional astrocyte IFN signaling restricts pathogenesis during neurotropic viral infection. J Clin Invest 127:843-856.
14. Morgan MJ, Kim YS. 2022. Roles of RIPK3 in necroptosis, cell signaling, and disease. Exp Mol Med 54:1695-1704.
15. Samson AL, Zhang Y, Geoghegan ND, Gavin XJ, Davies KA, Mlodzianoski MJ, Whitehead LW, Frank D, Garnish SE, Fitzgibbon C, Hempel A, Young SN, Jacobsen AV, Cawthorne W, Petrie EJ, Faux MC, Shield-Artin K, Lalaoui N, Hildebrand JM, Silke J, Rogers KL, Lessene G, Hawkins ED, Murphy JM. 2020. MLKL trafficking and accumulation at the plasma membrane control the kinetics and threshold for necroptosis. Nat Commun 11:3151.

598   16.   Daniels BP, Snyder AG, Olsen TM, Orozco S, Oguin TH, 3rd, Tait SWG, Martinez J, Gale
599   M, Jr., Loo YM, Oberst A. 2017. RIPK3 Restricts Viral Pathogenesis via Cell Death-
600   Independent Neuroinflammation. Cell 169:301-313 e11.
601   17.   Daniels BP, Kofman SB, Smith JR, Norris GT, Snyder AG, Kolb JP, Gao X, Locasale JW,
602   Martinez J, Gale M, Jr., Loo YM, Oberst A. 2019. The Nucleotide Sensor ZBP1 and
603   Kinase RIPK3 Induce the Enzyme IRG1 to Promote an Antiviral Metabolic State in
604   Neurons. Immunity 50:64-76 e4.
605   18.   Downey J, Pernet E, Coulombe F, Allard B, Meunier I, Jaworska J, Qureshi S, Vinh DC,
606   Martin JG, Joubert P, Divangahi M. 2017. RIPK3 interacts with MAVS to regulate type I
607   IFN-mediated immunity to Influenza A virus infection. PLoS Pathog 13:e1006326.
608   19.   Saleh D, Najjar M, Zelic M, Shah S, Nogusa S, Polykratis A, Paczosa MK, Gough PJ,
609   Bertin J, Whalen M, Fitzgerald KA, Slavov N, Pasparakis M, Balachandran S, Kelliher M,
610   Mecsas J, Degterev A. 2017. Kinase Activities of RIPK1 and RIPK3 Can Direct IFN-beta
611   Synthesis Induced by Lipopolysaccharide. J Immunol 198:4435-4447.
612   20.   Guo H, Koehler HS, Mocarski ES, Dix RD. 2022. RIPK3 and caspase 8 collaborate to
613   limit herpes simplex encephalitis. PLoS Pathog 18:e1010857.
614   21.   Peng R, Wang CK, Wang-Kan X, Idorn M, Kjaer M, Zhou FY, Fiil BK, Timmermann F,
615   Orozco SL, McCarthy J, Leung CS, Lu X, Bagola K, Rehwinkel J, Oberst A, Maelfait J,
616   Paludan SR, Gyrd-Hansen M. 2022. Human ZBP1 induces cell death-independent
617   inflammatory signaling via RIPK3 and RIPK1. EMBO Rep 23:e55839.
618   22.   Chou TW, Chang NP, Krishnagiri M, Patel AP, Lindman M, Angel JP, Kung PL, Atkins C,
619   Daniels BP. 2021. Fibrillar alpha-synuclein induces neurotoxic astrocyte activation via
620   RIP kinase signaling and NF-kappaB. Cell Death Dis 12:756.
621   23.   Li S, Zhang Y, Guan Z, Ye M, Li H, You M, Zhou Z, Zhang C, Zhang F, Lu B, Zhou P,
622   Peng K. 2023. SARS-CoV-2 Z-RNA activates the ZBP1-RIPK3 pathway to promote virus-
623   induced inflammatory responses. Cell Res doi:10.1038/s41422-022-00775-y.
624   24.   Najjar M, Saleh D, Zelic M, Nogusa S, Shah S, Tai A, Finger JN, Polykratis A, Gough PJ,
625   Bertin J, Whalen M, Pasparakis M, Balachandran S, Kelliher M, Poltorak A, Degterev A.
626   2016. RIPK1 and RIPK3 Kinases Promote Cell-Death-Independent Inflammation by Toll-
627   like Receptor 4. Immunity 45:46-59.
628   25.   Hubbard NW, Ames JM, Maurano M, Chu LH, Somfleth KY, Gokhale NS, Werner M,
629   Snyder JM, Lichauco K, Savan R, Stetson DB, Oberst A. 2022. ADAR1 mutation causes
630   ZBP1-dependent immunopathology. Nature 607:769-775.
631   26.   Xie Y, Zhao Y, Shi L, Li W, Chen K, Li M, Chen X, Zhang H, Li T, Matsuzawa-Ishimoto Y,
632   Yao X, Shao D, Ke Z, Li J, Chen Y, Zhang X, Cui J, Cui S, Leng Q, Cadwell K, Li X, Wei
633   H, Zhang H, Li H, Xiao H. 2020. Gut epithelial TSC1/mTOR controls RIPK3-dependent
634   necroptosis in intestinal inflammation and cancer. J Clin Invest 130:2111-2128.
635   27.   Sai K, Parsons C, House JS, Kathariou S, Ninomiya-Tsuji J. 2019. Necroptosis mediators
636   RIPK3 and MLKL suppress intracellular Listeria replication independently of host cell
637   killing. J Cell Biol 218:1994-2005.
638   28.   Baker DG, Woods TA, Butchi NB, Morgan TM, Taylor RT, Sunyakumthorn P, Mukherjee
639   P, Lubick KJ, Best SM, Peterson KE. 2013. Toll-like receptor 7 suppresses virus
640   replication in neurons but does not affect viral pathogenesis in a mouse model of Langat
641   virus infection. J Gen Virol 94:336-347.
642   29.   Michlmayr D, Bardina SV, Rodriguez CA, Pletnev AG, Lim JK. 2016. Dual Function of
643   Ccr5 during Langat Virus Encephalitis: Reduction in Neutrophil-Mediated Central
644   Nervous System Inflammation and Increase in T Cell-Mediated Viral Clearance. J
645   Immunol 196:4622-31.

30. Kimura T, Katoh H, Kayama H, Saiga H, Okuyama M, Okamoto T, Umemoto E, Matsuura
Y, Yamamoto M, Takeda K. 2013. Ifit1 inhibits Japanese encephalitis virus replication
through binding to 5' capped 2'-O unmethylated RNA. J Virol 87:9997-10003.
31. Szretter KJ, Daniels BP, Cho H, Gainey MD, Yokoyama WM, Gale M, Jr., Virgin HW,
Klein RS, Sen GC, Diamond MS. 2012. 2'-O methylation of the viral mRNA cap by West
Nile virus evades ifit1-dependent and -independent mechanisms of host restriction in
vivo. PLoS Pathog 8:e1002698.
32. Dai J, Pan W, Wang P. 2011. ISG15 facilitates cellular antiviral response to dengue and
west nile virus infection in vitro. Virol J 8:468.
33. Singh PK, Singh S, Farr D, Kumar A. 2019. Interferon-stimulated gene 15 (ISG15)
restricts Zika virus replication in primary human corneal epithelial cells. Ocul Surf 17:551-
559.
34. Bigham AW, Buckingham KJ, Husain S, Emond MJ, Bofferding KM, Gildersleeve H,
Rutherford A, Astakhova NM, Perelygin AA, Busch MP, Murray KO, Sejvar JJ, Green S,
Kriesel J, Brinton MA, Bamshad M. 2011. Host genetic risk factors for West Nile virus
infection and disease progression. PLoS One 6:e24745.
35. Vonderstein K, Nilsson E, Hubel P, Nygard Skalman L, Upadhyay A, Pasto J, Pichlmair
A, Lundmark R, Overby AK. 2018. Viperin Targets Flavivirus Virulence by Inducing
Assembly of Noninfectious Capsid Particles. J Virol 92.
36. Williams JL, Manivasagam S, Smith BC, Sim J, Vollmer LL, Daniels BP, Russell JH, Klein
RS. 2020. Astrocyte-T cell crosstalk regulates region-specific neuroinflammation. Glia
68:1361-1374.
37. Simmons SB, Liggitt D, Goverman JM. 2014. Cytokine-regulated neutrophil recruitment is
required for brain but not spinal cord inflammation during experimental autoimmune
encephalomyelitis. J Immunol 193:555-63.
38. Pierson ER, Goverman JM. 2017. GM-CSF is not essential for experimental autoimmune
encephalomyelitis but promotes brain-targeted disease. JCI Insight 2:e92362.
39. Durrant DM, Daniels BP, Pasieka T, Dorsey D, Klein RS. 2015. CCR5 limits cortical viral
loads during West Nile virus infection of the central nervous system. J Neuroinflammation
12:233.
40. McComb S, Cessford E, Alturki NA, Joseph J, Shutinoski B, Startek JB, Gamero AM,
Mossman KL, Sad S. 2014. Type-I interferon signaling through ISGF3 complex is
required for sustained Rip3 activation and necroptosis in macrophages. Proc Natl Acad
Sci U S A 111:E3206-13.
41. Brault M, Olsen TM, Martinez J, Stetson DB, Oberst A. 2018. Intracellular Nucleic Acid
Sensing Triggers Necroptosis through Synergistic Type I IFN and TNF Signaling. J
Immunol 200:2748-2756.
42. Ingram JP, Thapa RJ, Fisher A, Tummers B, Zhang T, Yin C, Rodriguez DA, Guo H,
Lane R, Williams R, Slifker MJ, Basagoudanavar SH, Rall GF, Dillon CP, Green DR,
Kaiser WJ, Balachandran S. 2019. ZBP1/DAI Drives RIPK3-Mediated Cell Death Induced
by IFNs in the Absence of RIPK1. J Immunol 203:1348-1355.
43. Lee SA, Chang LC, Jung W, Bowman JW, Kim D, Chen W, Foo SS, Choi YJ, Choi UY,
Bowling A, Yoo JS, Jung JU. 2023. OASL phase condensation induces amyloid-like
fibrillation of RIPK3 to promote virus-induced necroptosis. Nat Cell Biol
doi:10.1038/s41556-022-01039-y.
44. Yatim N, Jusforgues-Saklani H, Orozco S, Schulz O, Barreira da Silva R, Reis e Sousa
C, Green DR, Oberst A, Albert ML. 2015. RIPK1 and NF-kappaB signaling in dying cells
determines cross-priming of CD8(+) T cells. Science 350:328-34.

45.   Snyder AG, Hubbard NW, Messmer MN, Kofman SB, Hagan CE, Orozco SL, Chiang K,
Daniels BP, Baker D, Oberst A. 2019. Intratumoral activation of the necroptotic pathway
components RIPK1 and RIPK3 potentiates antitumor immunity. Sci Immunol 4.
46.   Wegner KW, Saleh D, Degterev A. 2017. Complex Pathologic Roles of RIPK1 and
RIPK3: Moving Beyond Necroptosis. Trends Pharmacol Sci 38:202-225.
47.   Angel JP, Daniels BP. 2022. Paradoxical roles for programmed cell death signaling
during viral infection of the central nervous system. Curr Opin Neurobiol 77:102629.
48.   Daniels BP, Oberst A. 2020. Outcomes of RIP Kinase Signaling During Neuroinvasive
Viral Infection. Curr Top Microbiol Immunol doi:10.1007/82_2020_204.
49.   Bian P, Ye C, Zheng X, Luo C, Yang J, Li M, Wang Y, Yang J, Zhou Y, Zhang F, Lian J,
Zhang Y, Jia Z, Lei Y. 2020. RIPK3 Promotes JEV Replication in Neurons via
Downregulation of IFI44L. Front Microbiol 11:368.
50.   Bian P, Zheng X, Wei L, Ye C, Fan H, Cai Y, Zhang Y, Zhang F, Jia Z, Lei Y. 2017. MLKL
Mediated Necroptosis Accelerates JEV-Induced Neuroinflammation in Mice. Front
Microbiol 8:303.
51.   Newton K, Sun X, Dixit VM. 2004. Kinase RIP3 is dispensable for normal NF-kappa Bs,
signaling by the B-cell and T-cell receptors, tumor necrosis factor receptor 1, and Toll-like
receptors 2 and 4. Mol Cell Biol 24:1464-9.
52.   Murphy JM, Czabotar PE, Hildebrand JM, Lucet IS, Zhang JG, Alvarez-Diaz S, Lewis R,
Lalaoui N, Metcalf D, Webb AI, Young SN, Varghese LN, Tannahill GM, Hatchell EC,
Majewski IJ, Okamoto T, Dobson RC, Hilton DJ, Babon JJ, Nicola NA, Strasser A, Silke
J, Alexander WS. 2013. The pseudokinase MLKL mediates necroptosis via a molecular
switch mechanism. Immunity 39:443-53.
53.   Kung PL, Chou TW, Lindman M, Chang NP, Estevez I, Buckley BD, Atkins C, Daniels
BP. 2022. Zika virus-induced TNF-alpha signaling dysregulates expression of neurologic
genes associated with psychiatric disorders. J Neuroinflammation 19:100.
54.   Daniels BP, Holman DW, Cruz-Orengo L, Jujjavarapu H, Durrant DM, Klein RS. 2014.
Viral pathogen-associated molecular patterns regulate blood-brain barrier integrity via
competing innate cytokine signals. mBio 5:e01476-14.
723