1   RIPK3 promotes brain region-specific interferon signaling and restriction of tick-borne flavivirus
2   infection
3

4   Marissa Lindman[a], Juan P Angel[a], Irving Estevez[a], Nydia P Chang[a], Tsui-Wen Chou[a], Micheal
5   McCourt[a], Colm Atkins[a], and Brian P. Daniels[a#]
6
7   [a]Department of Cell Biology and Neuroscience, Rutgers University, Piscataway, NJ, USA
8
#Correspondence:
Brian Daniels
604 Allison Road
Room B314
Piscataway, NJ 08854
(848) 445-2709
b.daniels@rutgers.edu

16
17
18

Abstract

Innate immune signaling in the central nervous system (CNS) exhibits many remarkable specializations that vary across cell types and CNS regions. In the setting of neuroinvasive flavivirus infection, neurons employ the immunologic kinase receptor-interacting kinase 3 (RIPK3) to promote an antiviral transcriptional program, independently of the traditional function of this enzyme in promoting necroptotic cell death. However, while recent work has established roles for neuronal RIPK3 signaling in controlling mosquito-borne flavivirus infections, including West Nile virus and Zika virus, functions for RIPK3 signaling in the CNS during tick-borne flavivirus infection have not yet been explored. Here, we use a model of Langat virus (LGTV) encephalitis to show that RIPK3 signaling is specifically required in neurons of the cerebellum to control LGTV replication and restrict disease pathogenesis. This effect did not require the necroptotic executioner molecule mixed lineage kinase domain like protein (MLKL), a finding similar to previous observations in models of mosquito-borne flavivirus infection. However, control of LGTV infection required a unique, region-specific dependence on RIPK3 to promote expression of key antiviral interferon-stimulated genes (ISG) in the cerebellum. This RIPK3-mediated potentiation of ISG expression was associated with robust cell-intrinsic restriction of LGTV replication in cerebellar granule cell neurons. These findings further illuminate the complex roles of RIPK3 signaling in the coordination of neuroimmune responses to viral infection, as well as provide new insight into the mechanisms of region-specific innate immune signaling in the CNS.

Importance

Interactions between the nervous and immune systems are very carefully orchestrated in order to protect the brain and spinal cord from immune-mediated damage, while still maintaining protective defenses against infection. These specialized neuro-immune interactions have been shown to vary significantly across regions of the brain, with innate antiviral signaling being particularly strong in the cerebellum, although the reasons for this are poorly understood. Here, we show a specialized adaptation of programmed cell death signaling that uniquely protects the cerebellum from tick-borne flavivirus infection. These findings provide important new insight into the molecular mechanisms that promote the uniquely robust antiviral immunity of the cerebellum. They also provide new clues into the pathogenesis of tick-borne encephalitis, a zoonosis of significant global concern.

Introduction

        Flaviviruses are a family of positive sense RNA viruses which include several notable pathogens associated with neuroinvasive infection in humans, including West Nile virus (WNV), Zika virus (ZIKV), and Japanese Encephalitis virus (1). While nearly all major flaviviruses are transmitted by mosquito vectors, a small but significant number of flaviviruses are transmitted by ticks, including Tick-borne encephalitis virus (TBEV) and its close relatives that together make up a single TBEV serocomplex. Tick borne encephalitis is a significant and growing threat to public health, particularly in Europe and northern Asia, where TBEV constitutes the most prevalent tick-borne zoonotic disease (2-4). Notably, some TBEV strains elicit mortality rates up to 40% in humans (5), underscoring the urgent need to better understand the mechanisms underlying the pathogenesis of tick-borne flavivirus infections.

        Effective control of flavivirus infection in the central nervous system (CNS) requires robust innate immune signaling in neural cells, particularly neurons, which are the predominantly infected cell type in most cases of flavivirus encephalitis (6-9). Effective type I interferon (IFN) signaling is of particular importance for innate control of viral replication in neurons (10-12). Notably, differences in type I IFN signaling across neural cell types and brain regions are associated with differential susceptibility to flavivirus infection. For example, previous reports suggest that the enhanced type I IFN signaling observed in hindbrain regions compared to the forebrain is an underlying determinant of the enhanced susceptibility of forebrain regions to WNV infection (12, 13). However, the unique signaling mechanisms that promote differential IFN-mediated control of viral infection in the hindbrain have not been extensively characterized.

        A potential regulator of neuronal IFN signaling during flavivirus infection is receptor interacting protein kinase-3 (RIPK3). RIPK3 is an enzyme traditionally associated with necroptosis, a form of lytic programmed cell death (14). Necroptosis occurs via the RIPK3-dependent activation of mixed lineage kinase domain like protein (MLKL), which forms oligomeric pore complexes that induce cellular lysis (15). However, many recent studies have identified complex roles for RIPK3 signaling in the coordination of inflammation, including the regulation of inflammatory transcriptional responses that occur independently of necroptosis (16-24). We and others have demonstrated that RIPK3 signaling in neurons is of particular importance for the control of neurotropic viral infections, as neuronal RIPK3 promotes a robust antimicrobial transcriptional program, including many IFN stimulated genes (ISGs), that restricts viral infection without inducing neuronal necroptosis (16, 17). Other recent studies have identified unexpected roles for RIPK3 in the regulation of type I IFN signaling, via mechanisms which include the regulation of pattern recognition receptor signaling and protein kinase-R (PKR)-mediated stabilization of *Ifnb* mRNA (18, 19).

        In this study, we interrogated roles for RIPK3 in controlling tick-borne flavivirus infection. To do so, we used Langat virus (LGTV), a naturally attenuated member of the TBEV serocomplex that can be studied under BSL2 containment. *Ripk3*[-/-] mice exhibited enhanced clinical disease following subcutaneous LGTV infection, while *Mlkl*[-/-] mice were indistinguishable from littermate controls, suggesting a necroptosis-independent function for RIPK3 in restricting LGTV pathogenesis. Notably, *Ripk3*[-/-] mice exhibited increased viral burden in the cerebellum, along with diminished expression of inflammatory chemokines and ISGs in the cerebellum, but not the cerebral cortex. *In vitro* analysis of cultured primary cortical and cerebellar cell types showed that pharmacologic inhibition of RIPK3 resulted in enhanced LGTV replication in cerebellar granule cell neurons but not in cortical neurons or in astrocytes derived from either brain region. Transcriptional profiling showed that RIPK3 signaling was uniquely required for the

106 full induction of ISG expression in cerebellar granule cell neurons, demonstrating a previously
107 unknown, region-specific function for RIPK3 in coordinating innate antiviral immunity within the
108 CNS.
109
110
111 **Results**
112
113 **RIPK3 controls LGTV pathogenesis independently of MLKL and peripheral immunity**
114
To assess the role of RIPK3 in controlling LGTV pathogenesis, we subcutaneously
infected *Ripk3⁻ᐟ⁻* mice, along with heterozygous littermate controls, with 3x10⁴ plaque forming
units (pfu) of the Malaysian LGTV strain TP21. We note that *Ripk3⁺ᐟ⁻* animals do not exhibit
haploinsufficiency and are routinely used as littermate controls in studies of this pathway (25-
27). Control animals exhibited limited mortality following LGTV infection (Figure 1A), consistent
with previous reports (28, 29). However, mice lacking *Ripk3* expression exhibited a significantly
accelerated and enhanced rate of mortality (Figure 1A). In addition, a higher proportion of *Ripk3⁻*
ᐟ⁻ mice exhibited clinical signs of neurologic disease, including paresis or full hindlimb paralysis,
by 14 days post infection (dpi) (Figure 1B), and this difference persisted to at least 21 dpi. These
data suggest that *Ripk3* is essential for restricting neuropathogenesis during LGTV infection.
125
To better understand this phenotype, we first assessed whether RIPK3 was required for
early control of systemic infection. Spleens of infected mice exhibited low levels of LGTV RNA
that were not impacted by *Ripk3* expression (Figure 1C). To test whether *Ripk3⁻ᐟ⁻* mice exhibited
any deficiencies in peripheral immune responses, we performed flow cytometric analysis of
major immune cell subsets in the spleens of infected animals at 8 dpi. *Ripk3⁻ᐟ⁻* animals exhibited
similar frequencies (Figure 1D) and total numbers (Figure 1E-F) of CD4 and CD8 T cells among
all splenocytes compared to littermate controls, as well as similar rates of CD44 expression (a
key T cell activation marker) across both subsets (Figure 1G-H). Numbers of B cells (Figure 1I)
and natural killer (NK) cells (Figure 1J) were also similar between genotypes. In the myeloid
compartment, we observed similar numbers of CD11c⁺ MHCII⁺ dendritic cells (Figure 1K)
between genotypes, as well as similar numbers of myeloid subsets expressing F4/80 (Figure
1L), Ly6G (Figure 1M), and Ly6C (Figure 1N). Both CD11c⁺ MHCII⁺ and F4/80⁺ antigen
presenting cell subsets also exhibited similar rates of expression of the costimulation signal
CD80 between genotypes (Figure 1O-P). These data suggest that *Ripk3⁻ᐟ⁻* mice mounted normal
peripheral immune responses to subcutaneous LGTV challenge, similar to our previous
observations with WNV and ZIKV (16, 17). Thus, the increased pathogenesis observed in mice
lacking *Ripk3* was unlikely to arise from a failure in peripheral virologic control.
143
A potential mechanism by which RIPK3 signaling might restrict LGTV pathogenesis is
through the induction of necroptosis in infected cells. We thus tested whether loss of the
necroptotic executioner molecule MLKL would impact disease course following subcutaneous
LGTV infection. Notably, *Mlkl⁻ᐟ⁻* mice exhibited no difference in either survival or development of
clinical disease signs compared to littermate controls (Figure 2A-B). We saw similarly that *Mlkl⁻ᐟ⁻*
did not exhibit altered splenic viral burden at 8dpi (Figure 2C). Flow cytometric analysis also
revealed essentially identical numbers and frequencies of all major immune cell subsets in the
spleen at this time point (Figure 2C-P). Multistep growth curve analysis also demonstrated that
neither RIPK3 nor MLKL impacted the low levels of LGTV replication observed in primary
leukocyte cultures, including bone marrow derived macrophages and dendritic cells
(Supplemental Figure 1A-B). These data suggest that MLKL, and therefore necroptosis, is not a
major contributor to peripheral virologic control or overall disease pathogenesis in the setting of

LGTV infection, and thus that RIPK3 exerts its protective effect in this model through an
alternative mechanism.
158
159 **RIPK3 is required for CNS-intrinsic restriction of LGTV infection**
160
Because we did not observe differences in peripheral virologic control in *Ripk3*[-/-] mice, we
next questioned whether RIPK3 acted in a CNS-intrinsic manner to limit LGTV infection. To
assess this, we next used an intracranial infection route in order to assess local effects of RIPK3
signaling on LGTV pathogenesis. *Ripk3*[-/-] mice exhibited accelerated and enhanced mortality
compared to littermate controls following intracranial infection (Figure 3A). *Ripk3*-deficient mice
also exhibited worsened clinical disease prior to death, as evidenced by earlier and more
dramatic weight loss following infection (Figure 3B). In contrast, *Mlkl*[-/-] mice were
indistinguishable from littermate controls in terms of overall mortality (Figure 3C) and weight loss
(Figure 3D) following intracranial infection. These data further supported the idea that RIPK3
restricts LGTV neuropathogenesis via CNS-intrinsic mechanisms, independently of necroptosis.
171
172 **RIPK3 promotes neuronal chemokine expression in a region-specific manner following**
173 **LGTV infection**
174
We and other previously showed that neuronal RIPK3 signaling was required for the
expression of key inflammatory chemokines that served to restrict WNV pathogenesis by
coordinating the recruitment of leukocytes into the infected CNS. We thus questioned whether
RIPK3 also promotes chemokine expression in the CNS during LGTV infection. Surprisingly,
transcriptional profiling in the cerebral cortex of *Ripk3*[-/-] mice following subcutaneous LGTV
infection revealed no differences in expression of major chemokines compared to littermate
controls (Figure 4A). However, we did observe significantly diminished chemokine responses in
cerebellar tissues derived from *Ripk3*[-/-] animals (Figure 4B). To understand which cell types
were driving this region-specific deficit in chemokine expression, we next cultured primary
neurons and astrocytes derived specifically from either cerebral cortex or cerebellum and
infected with LGTV, with or without a small molecule inhibitor of RIPK3 (GSK 872). Consistent
with our *in vivo* findings, blockade of RIPK3 in cerebral cortical neurons did not impact
chemokine expression following LGTV infection (Figure 4C). In contrast, infected cerebellar
granule cell neuron cultures exhibited significantly diminished chemokine expression when
RIPK3 was inhibited by GSK 872 (Figure 4D). Notably, we did not observe a dependence on
RIPK3 for the expression of chemokines in astrocytes derived from either region (Figure 4E-F).
These data suggest that RIPK3 serves an unexpected, region-specific transcriptional function in
neurons of the cerebellum during neuroinvasive LGTV infection.
193
194 **RIPK3 is not required for immune cell recruitment to the LGTV-infected CNS**
195
We next questioned whether diminished chemokine expression in the cerebellum of
Ripk3[-/-] mice would result in a failure to recruit antiviral leukocytes into this brain region. We thus
performed flow cytometric analysis of leukocytes derived from either cerebral cortex or
cerebellum following subcutaneous LGTV infection. Remarkably, we saw no evidence of
changes in lymphocyte recruitment in either brain region of *Ripk3*[-/-] mice compared to littermate
controls on either 6 or 8 dpi (Figure 5A). This lack of difference extended across all major
CD45[hi] infiltrating leukocyte subsets, including CD4[+] and CD8[+] T cells (Figure 5B-C), NK cells
(Figure 5D), CD11c[+] MHCII[+] dendritic cells (Figure 5E) and myeloid subsets expressing F4/80
(Figure 5F), Ly6G (Figure 5G), and Ly6C (Figure 5H). We similarly did not observe differences
in numbers of CD45[lo] microglia (Figure 5I), suggesting no major differences in microglial

proliferation between genotypes in either region. These data suggested that, despite significant
differences in the expression of major leukocyte chemoattractants in the cerebellum, differences
in immune cell recruitment did not account for the increased pathogenesis observed in *Ripk3⁻/⁻*
mice during LGTV infection.
210
211 **RIPK3 promotes cell-intrinsic restriction of LGTV replication in cerebellar neurons**
212
Given these observations, we next questioned whether *Ripk3⁻/⁻* mice fail to control LGTV
infection due to impaired innate immune restriction of LGTV replication. Assessment of viral
burdens in brains of *Ripk3⁻/⁻* mice following subcutaneous LGTV infection revealed that *Ripk3⁻/⁻*
mice exhibited significantly elevated CNS viral titers, particularly in the cerebellum, at both 8 and
12 dpi (Figure 6A). In contrast, *Mlkl⁻/⁻* exhibited no such difference in viral burden in either brain
region (Figure 6B). Differences in viral burden did not appear to be linked to deficits in blood-
brain barrier integrity, as both *Ripk3⁻/⁻* mice and littermate controls exhibited similar levels of
sodium fluorescein extravasation into the CNS following infection (Figure 6C). We thus
questioned whether RIPK3 was required for cell-intrinsic restriction of viral replication in
susceptible CNS cell types. Multistep growth curve analysis in primary CNS cells revealed that
pharmacologic inhibition of RIPK3 had no effect on LGTV replication in neurons derived from
cerebral cortex (Figure 6D). In contrast, inhibition of RIPK3 significantly enhanced LGTV
replication in primary cerebellar granule cell neurons cultures (Figure 6E). This effect was
unique to neurons, as GSK 872 treatment had no impact on LGTV replication in primary
astrocytes derived from either brain region (Figure 6F-G). Together, these data suggested that
the enhanced pathogenesis observed in *Ripk3⁻/⁻* mice was due to a specific failure to control
infection in neurons of the cerebellum, resulting in enhanced overall CNS viral burden.
230
231 **RIPK3 potentiates Type I IFN signaling in cerebellar neurons during LGTV infection**
232
Our previous observation of diminished chemokine expression in cerebellar neurons
derived from *Ripk3⁻/⁻* mice suggested that these cells may exhibit broader deficits in innate
immune signaling, resulting in poor control of LGTV replication. We, therefore, next questioned
whether IFN signaling was perturbed in the cerebellum of mice lacking RIPK3 expression.
Transcriptional profiling in brain tissues following subcutaneous LGTV infection revealed that,
indeed, the cerebella of *Ripk3⁻/⁻* mice exhibited diminished expression of many ISGs known to
be critical for control of flavivirus replication (30-35), including *Ifit1, Isg15, Mx1, Mx2, Oas1b,* and
Rsad2, while this phenotype was not observed in the cerebral cortex (Figure 7A-B). Similar
analyses in primary cell cultures confirmed that cerebellar granule cell neurons, but not neurons
derived from cerebral cortex, exhibited diminished expression of ISGs when RIPK3 signaling
was blocked via GSK 872 treatment (Figure 7C-D). In contrast, we observed little to no impact
of RIPK3 blockade on ISG expression in astrocytes derived from either brain region (Figure 7E-
F). Together, these data demonstrate that RIPK3 signaling is required for the robust induction of
type I IFN responses in neurons of the cerebellum, which is required for cell-intrinsic restriction
of LGTV replication.
248
To better understand the role of RIPK3 signaling in potentiating ISG expression, we next
questioned whether RIPK3 acts downstream of IFN receptor signaling. To assess this, we
treated neuron cultures with exogenous IFNβ for 1 hour following pretreatment with GSK 872 or
vehicle control. As expected, IFNβ treatment resulted in robust induction of multiple ISGs
(Figure 8A, Supplemental Figure 2A). However, pharmacologic blockade of RIPK3 did not
impact ISG expression induced by IFNβ treatment in either cerebellar granule cell neurons
(Figure 8A) or in cerebral cortical neurons (Supplemental Figure 2A), suggesting that RIPK3

likely does not act directly downstream of the type I IFN receptor (IFNAR) to modulate gene expression and/or that type I IFN alone is not sufficient to induce RIPK3 activation. We next tested the alternative hypothesis that RIPK3 regulates IFN signaling during LGTV infection by directly influencing the expression of IFN ligands and receptors. Surprisingly, transcriptional analysis revealed that pharmacologic blockade of RIPK3 did not influence the expression of the type I IFN ligands *Ifna* and *Ifnb* in cerebellar granule cell neurons following infection (Figure 8B). In contrast, GSK 843 treatment significantly blunted infection-induced upregulation of IFN receptor subunits, including the type I IFN receptor subunits *Ifnar1* and *Ifnar2*, and the type II IFN receptor subunits *Ifngr1*, and *Infgr2*. Other IFN ligands and receptors, including *Ifng* and the type III IFN ligands *Ifnl2* and *Ifnl3* were undetectable in all conditions (data not shown). Importantly, we did not observe this RIPK3-dependency in IFN receptor expression in cerebral cortical neuron cultures (Supplemental Figure 2B), suggesting that RIPK3 functions uniquely in cerebellar granule cell neurons to enhance type I IFN signaling during LGTV infection.

To further investigate a role for RIPK3 in IFN-mediated gene expression in cerebellar neurons, we next performed experiments in which we blocked RIPK3 activity with or without simultaneous blockade of type I IFN signaling using a neutralizing antibody against IFNAR1. We reasoned that this paradigm would allow us to assess the differential influence of RIPK3 on IFNAR-dependent and IFNAR-independent gene expression following LGTV infection. Perhaps unsurprisingly, we observed that expression of most ISGs was completely dependent on IFNAR1 signaling, making it difficult to distinguish a specific role for RIPK3 in the absence of intact type I IFN signaling (Supplemental Figure 2C). We thus identified several alternative inflammatory genes whose expression was either completely (*Cxcl1*) or partially (*Ccl2 and Il6*) independent of IFNAR1 signaling following infection. Notably, pharmacologic blockade of RIPK3 only impacted the IFNAR1-depenent portion of the induced expression of these genes, while having no effect on the IFNAR1-independent portion, as indicated by a lack of effect in αIFNAR1-treated cultures (Figure 8C). Together, these data further support our observation of synergistic signaling between type I IFN and RIPK3 signaling in cerebellar granule cell neurons during LGTV infection.

Discussion

Our findings identify a previously unknown function for RIPK3 in the coordination of brain region-specific innate immunity. The study of regional differences in neuroimmune signaling is a growing field, and there is accumulating evidence to suggest that resident neural cells exhibit differential responses to viral infection and cytokine stimulation across distinct anatomical regions of the CNS (36-39). Neurons and astrocytes in the cerebellum, in particular, have been shown to exhibit higher responsiveness to stimulation by type-I IFN, as well as to express higher basal levels of pathogen sensor molecules compared to other brain regions, suggesting a key evolutionary importance of innate antiviral defense in this tissue (12, 13). This regional difference in type I IFN signaling appears to underlie, at least in part, the relatively lower susceptibility of the cerebellum to flavivirus infection compared to susceptible regions of the forebrain, such as the cerebral cortex and hippocampus. However, the molecular mechanisms that determine the enhanced innate immune signaling observed in the cerebellum remain poorly understood. Our study suggests that RIPK3 signaling is required for the robust induction of ISG expression in cerebellar neurons during LGTV infection, although ongoing work is needed to understand the specific signaling interactions that mediate this effect.

Previous studies have described a highly complex interplay between RIPK3 and type I IFN signaling that varies significantly by cell type and disease model (17-19, 40). It is relatively

clear that type I IFN signaling is capable of activating RIPK3 through various mechanisms,
resulting in necroptosis and/or necroptosis-independent transcriptional activation (40-43).
However, how RIPK3 operates *upstream* of (or synergistically with) type I IFN signaling to
influence expression of ISGs is less clear. We and others have shown that ISG expression is
significantly diminished in a variety of settings when RIPK3 signaling is ablated (17, 18),
including in cerebellar granule cell neurons during LGTV infection in this study. One possible
explanation for this effect is RIPK3-mediated activation of NF-κB, a transcription factor strongly
associated with RIPK signaling with known roles in potentiating type I IFN signaling and ISG
expression (22, 44-46). We and others also previously showed that RIPK3 activation in cortical
neurons following ZIKV infection leads to interferon regulatory factor 1 (IRF1) activation, which
was required for expression of at least a subset of RIPK3-induced genes in that setting,
although this effect is likely indirect, as IRF1 is not a known RIPK3 substrate (17). Additional
work will be needed to fully characterize the regulatory mechanisms that are invoked in the
interplay between RIPK3 and type I IFN signaling in the CNS.
320
Our study also further expands our understanding of the necroptosis-independent
functions for RIPK3 signaling in the CNS. Many studies have now firmly established the
importance of RIPK3 in promoting host defense through mechanisms independent of its
canonical role in necroptosis (16-21). However, these necroptosis-independent functions appear
to vary significantly by disease state, including CNS infection with distinct neuroinvasive
flaviviruses (47, 48). We and others previously showed that the primary role for RIPK3 in
restricting WNV encephalitis was the induction of chemokine expression and the recruitment of
antiviral leukocytes into the infected CNS (16). Notably, while we did observe RIPK3-mediated
chemokine expression in the cerebellum during LGTV infection, this chemokine expression was
apparently dispensable for CNS immune cell recruitment. Instead, the transcriptional activation
of antiviral effector genes, including ISGs, was required for cell-intrinsic restriction of LGTV
replication in neurons, a phenotype more similar to our findings with ZIKV (17), although we did
not observe evidence for a regional specification of this response during ZIKV infection. In
contrast to these observations, Bian and colleagues have observed quite distinct phenotypes in
a model of JEV encephalitis, wherein both RIPK3 and MLKL appeared to exacerbate rather than
restrict disease pathogenesis (49, 50). RIPK3 also appeared to *suppress* rather than promote
ISG expression in JEV infected neurons. The factors that determine such distinct outcomes of
RIPK3 signaling across this family of closely related viruses are mysterious and are the subject
of ongoing investigation by our laboratory and others.
340
341

Materials and Methods

Mouse lines

Ripk3[-/-] (51) *Mlkl*[-/-] (52) mouse lines were bred and housed under specific-pathogen free conditions in Nelson Biological Laboratories at Rutgers University. *Ripk3*[-/-] mice were generously provided by Genentech, Inc. Wild-type C57BL/6J mice were either obtained commercially (Jackson Laboratories) or bred in-house. Mice used for subcutaneous infections were 5 weeks old; mice used for intracranial infections were 8-15 weeks old.

Virus and titer determination

Langat virus strain TP21 was used throughout the study. Founder stocks were obtained from the World Reference Center for Emerging Viruses and Arboviruses (WRCEVA). Laboratory stocks were generated using Vero E6 cells (ATCC, #CRL-1586) and frozen at -80°C until needed. Virus titers were determined by plaque assay on Vero E6 cells. Cells were maintained in DMEM (Corning #10-013-CV) supplemented with 10% Heat Inactivated FBS (Gemini Biosciences #100-106), 1% Penicillin–Streptomycin-Glutamine (Gemini Biosciences #400-110), 1% Amphotericin B (Gemini Biosciences #400–104), 1% Non-Essential Amino Acids (Cytiva, #SH30238.01), and 1% HEPES (Cytiva, #SH30237.01). Plaque assay media was composed of 1X EMEM (Lonza # 12-684F) supplemented with 2% Heat Inactivated FBS (Gemini Biosciences #100-106), 1% Penicillin–Streptomycin-Glutamine (Gemini Biosciences, #400-110), 1% Amphotericin B (Gemini Biosciences #400-104), 1% Non-Essential Amino Acids (Cytiva, #SH30238.01), and 1% HEPES (Cytiva, SH30237.01), 0.75% Sodium Bicarbonate (VWR, #BDH9280) and 0.5% Methyl Cellulose (VWR, #K390). Plaque assays were developed at 5dpi by removal of overlay media and staining/fixation using 10% neutral buffered formalin (VWR, #89370) and 0.25% crystal violet (VWR, #0528). Plaque assays were performed by adding 100uL of serially diluted sample for 1 hour at 37°C to 12-well plates containing 200,000 Vero E6 cells per well. Plates were further incubated with plaque assay media at 37°C and 5% CO2 for 5 days. Medium was removed from the wells and replaced with fixative containing crystal violet for approximately 20-30 minutes. Plates were washed repeatedly in H_2O and allowed to dry before counting visible plaques.

Mouse infections and tissue harvesting

Isoflurane anesthesia was used for all procedures. Mice were inoculated subcutaneously (50uL) with $3x10^4$ PFU or injected intracranially (10uL) with 50 PFU of LGTV-TP21 using insulin syringes (BD Medical, #BD-329461). At appropriate times post infection, mice underwent cardiac perfusions with 30 mL cold sterile 1X phosphate-buffered saline (PBS). Extracted tissues were weighed and homogenized using 1.0 mm diameter zirconia/silica beads (Biospec Products, #11079110z) in sterile PBS for plaque assay or TRI Reagent (Zymo, #R2050-1) for gene expression analysis. Homogenization was performed in an Omni Beadrupter Elite for 2 sequential cycles of 20 s at a speed of 4 m/s.

Primary cell infections

Cortical and cerebellar astrocytes were harvested from P1-P2 pups and cortical neurons were harvested at E13.5-E15.5. Tissues were dissociated using the Neural Dissociation Kit (T) following manufacturer's instructions (Miltenyi, #130-093-231). Astrocytes were expanded in AM-a medium (ScienCell, #1831) supplemented with 10% FBS in fibronectin-coated cell culture flasks and seeded into plates coated with 20 μg/mL Poly-L-Lysine (Sigma-Aldrich, #9155) before experiments. Neurons were seeded into PLL-coated cell culture treated plates and grown in Neurobasal Plus + B-27 supplement medium (Thermo-Fisher Scientific, #A3582901) prior to use in experiments 7-9 days in vitro (DIV). Mouse cerebellar granule cells from C57BL/6 mice

392 (ScienCell, # M1530-57) were seeded into cell culture treated plates coated with 10 ug/mL Poly-
393 D-Lysine (ThermoFisher, #A3890401) containing prewarmed Neuronal Medium (ScienCell,
394 #1521) following manufacturer recommendations and used for experiments 6 DIV.
395
Macrophages and dendritic cells were isolated from bone marrow of euthanized mice.
Femurs were isolated and bone marrow pushed out using a sterile needle and syringe loaded
with RPMI supplemented with 10% FBS, 1% Penicillin–Streptomycin-Glutamine, 1% HEPES,
1% Glutamax (ThermoFisher, #35050061). Bone marrow was plated into non-cell-culture
treated 10cm petri dishes in 8mL supplemented RPMI medium containing either 20ng/mL
recombinant M-CSF (Peprotech, #315-02) or 20ng/mL recombinant GM-CSF (Peprotech, #315-
03) and 20ng/mL IL-4 (Peprotech, #214-14) for differentiation into macrophages or dendritic
cells, respectively. Cells were fed with additional medium containing the appropriate cytokines
four days later and used for experiments at 6-7 DIV. Cells were seeded into cell-culture treated
dishes prior to experimentation. For viral replication determination, all cultures were infected
with LGTV TP21 at an MOI of 0.01. For qRT-PCR experiments, cortical and cerebellar neuron
cultures were infected at an MOI of 0.5, while astrocyte cultures were infected using an MOI of
0.01. The pharmacologic inhibitor of RIPK3, GSK872, was added to cultures at 100nM for 2
hours prior to infection or subsequent treatments. Interferon-β was added to neuron cultures at
10ng/mL for 1 hour prior to harvesting cell lysates. IFNAR-1 monoclonal antibody (MAR1-5A3,
Leinco Technologies) or isotype control (GIR-208, Leinco Technologies) were added to cultures
at 5µg/mL 45 minutes prior to Langat virus infection.
413
Quantitative real-time PCR
Total RNA from harvested tissues was extracted using Zymo Direct-zol RNA Miniprep kit,
as per manufacturer instructions (Zymo, #R2051). Total RNA extraction from cultured cells,
cDNA synthesis, and subsequent qRT-PCR were performed as previously described (22, 53).
Cycle threshold (CT) values for analyzed genes were normalized to CT values of the
housekeeping gene 18 S ($CT_{Target} - CT_{18S} = \Delta CT$). Data from primary cell culture experiments
were further normalized to baseline control values ($\Delta CT_{experimental} - \Delta CT_{control} = \Delta\Delta CT$ (DDCT)). A
list of primers used in this study can be found in **Supplemental Table 1**.
422
Flow Cytometry
The cerebella and cerebral cortices of mouse brains were dissected from freshly
perfused mice and placed into tubes containing 1X PBS. Brain tissues were incubated with
10mL buffer containing 0.05% Collagenase Type I (Sigma-Aldrich, #C0130), 10ug/mL DNase I
(Sigma-Aldrich, #D4527) and 10mM HEPES (Cytiva, #SH30237.01) in 1X Hanks' Balanced Salt
Solution (VWR, #02-1231-0500) for one hour at room temperature under constant rotation.
Brain tissues were transferred to a 70um strainer on 50mL conical tubes and mashed through
the strainer using the plunger of 3-5mL syringes. Tissue was separated in 8 mL 37% Isotonic
Percoll (Percoll: Cytiva, #17-0891-02; RPMI 1640: Corning, #10-040-CV, supplemented with 5%
FBS) by centrifugation at 1200xg for 30 minutes with a slow break. The myelin layer and
supernatant were discarded. Leukocytes were incubated in 1X RBC Lysis Buffer (Tonbo
Biosciences, #TNB-4300-L100) for 10 minutes at room temperature. Cells were centrifuged and
resuspended in FACS buffer composed of 1X PBS, 2% sodium azide and 5% FBS. Samples
were transferred into a U-bottomed 96-well plate. Leukocytes were blocked with 2% normal
mouse serum and 1% FcX Block (BioLegend, #101320) in FACS buffer for 30 minutes at 4°C
prior to being stained with fluorescently conjugated antibodies to CD3e (Biolegend, clone 17A2),
CD44 (Biolegend, clone IM7), CD19 (Biolegend, clone 6D5), CD8a (Biolegend, clone 53-6.7),
CD4 (Biolegend, clone RM4-5), CD45.2 (Biolegend, clone 104), MHC-II (Biolegend, clone
M5/114.15.2), NK1.1 (Biolegend, clone PK136), CD11c (Biolegend, clone N418), F4/80

(Biolegend, clone BM8), CD11b (Biolegend, clone M1/70), Ly6G (Biolegend, clone 1A8), Ly6C
(Biolegend, clone HK1.4), CD80 (Biolegend, clone 16-10A1), and Zombie NIR (Biolegend,
#423105). Leukocytes were stained for 30 minutes at 4C prior to washing in FACS buffer and
fixation with 1% PFA in PBS (ThermoFisher, #J19943-K2). Data collection and analysis were
performed using a Cytek Northern Lights Cytometer (Cytek, Fremont, California) and FlowJo
software (Treestar). Data were normalized using a standard bead concentration counted by the
cytometer with each sample (ThermoFisher, #C36950). Spleens were crushed between two
slides, filtered through a 70um cell strainer, and washed with FACS buffer. Isolated splenocytes
were incubated with 1X RBC Lysis Buffer as done for leukocytes isolated from the brain prior to
blocking and staining.
452
In vivo assessment of blood brain barrier permeability
In vivo assessment of blood brain barrier permeability was carried out as described (54).
Mice were injected intraperitoneally with 100uL of 100mg/mL fluorescein sodium salt (Sigma,
#F6377) dissolved in sterile 1X PBS. After 45 minutes, blood was collected followed by cardiac
perfusion. Tissues were dissected and homogenized in 1X PBS as described above. Serum and
supernatant from homogenized tissues were incubated overnight at 4°C with 2% Trichloroacetic
acid solution (Sigma, #T0699) at a 1:1 dilution. Precipitated protein was pelleted by 10 minutes
of centrifugation at 2,823xg at 4°C. Supernatants were diluted with borate buffer, pH 11 (Sigma,
#1094621000) to achieve a neutral pH. Fluorescein emission at 538nm was measured for
samples in an optically clear black-walled 96-well plate (Corning, #3904) using a SpectraMax
iD3 plate reader (Molecular Devices, San Jose, CA). Tissue fluorescence values were
standardized against plasma values for individual mice.
465
Statistical analysis
Normally distributed data were analyzed using appropriate parametric tests: two-way
analysis of variance (ANOVA) with Sidak's correction for multiple comparisons and Log-rank
(Mantel-Cox) test for survival comparison, both using GraphPad Prism Software v8 (GraphPad
Software, San Diego, CA). Chi square tests for comparison of clinical disease signs was
performed using Excel v2211 (Microsoft). $P < 0.05$ was considered statistically significant.
472

Acknowledgements

This work was supported by R01 NS120895 (to BPD). JPA and IE were supported by NIH Supplement to Promote Diversity (R01 NS120895-S1 and NS120895-S2). NPC was supported by F31 NS124242.

479 Figures
480

481
482

Figure 1. RIPK3 limits LGTV pathogenesis independently of peripheral immunity

(A-B) Survival analysis (A) and presentation of clinical signs of disease (B) in *Ripk3*-/- mice and littermate

controls following subcutaneous inoculation with 3x10⁴ PFU LGTV TP21. Data are pooled from two

experiments. (C) *Ripk3*-/- and littermate control mice were infected subcutaneously with LGTV TP21. On

indicated days following infection, splenic viral burden was measured via qRT-PCR. Data was normalized

against a standard curve of known viral titers to generate plaque-forming unit (PFU) equivalents. Data

for each day post infection are pooled from 2-3 experiments. LOD, limit of detection. (D-P) *Ripk3*-/- and

littermate control mice were infected subcutaneously with LGTV TP21 for 8 days prior to harvesting

splenocytes and profiling leukocytes by flow cytometry. (D) Representative flow cytometry plots

showing CD8+ and CD4+ T cells among CD3+ leukocytes in the spleen. Numbers represent percentage of

cells in each gate relative to total plotted cells. (E-F) Numbers of CD8+ T cells (E) and CD4+ T cells (F)

among CD3+ leukocytes. (G-H) Percentage of CD44+ cells among CD8+ T cells (G) and CD4+ T cells (H).

(I-N) Numbers of CD19+ B cells (I), CD11b+ NK1.1+ Natural Killer cells (J), CD11c+ MHC-II+ dendritic cells

(K), CD45high CD11b+ F4/80+ macrophages (L), CD11b+ Ly6G+ neutrophils (M), and CD45high CD11b+

Ly6C+ monocytes (N) among total leukocytes in the spleen. (O-P) Percentage of CD80+ cells among

CD11c+ MHC-II+ dendritic cells (O) and CD11b+ F4/80+ macrophages (P). ns, not significant. **p < 0.01,

***p < 0.001. Error bars represent SEM.

501
502

Figure 2. MLKL signaling does not influence Langat virus pathogenesis.
(A-B) Survival analysis (A) and presentation of clinical signs of disease (B) in $Mlkl^{-/-}$ mice and littermate
controls following subcutaneous inoculation with $3X10^4$ PFU LGTV TP21. Data are pooled from two
experiments. (C) $Mlkl^{-/-}$ and littermate control mice were infected subcutaneously with LGTV TP21. On
indicated days following infection, splenic viral burden was measured via qRT-PCR. Data was normalized
against a standard curve of known viral titers to generate plaque-forming unit (PFU) equivalents. Data
for each day post infection are pooled from 2-3 experiments. LOD, limit of detection. (D-P) $Mlkl^{-/-}$ and
littermate control mice were infected subcutaneously with LGTV TP21 for 8 days prior to harvesting
splenocytes and profiling leukocytes by flow cytometry. (D) Representative flow cytometry plots
showing CD8+ and CD4+ T cells among CD3+ leukocytes in the spleen. Numbers represent percentage of
cells in each gate relative to total plotted cells. (E-F) Numbers of CD8+ T cells (E) and CD4+ T cells (F)
among CD3+ leukocytes. (G-H) Percentage of CD44+ cells among CD8+ T cells (G) and CD4+ T cells (H).
(I-N) Numbers of CD19+ B cells (I), CD11b+ NK1.1+ Natural Killer cells (J), CD11c+ MHC-II+ dendritic cells
(K), CD45high CD11b+ F4/80+ macrophages (L), CD11b+ Ly6G+ neutrophils (M), and CD45high CD11b+
Ly6C+ monocytes (N) among total leukocytes in the spleen. (O-P) Percentage of CD80+ cells among
CD11c+ MHC-II+ dendritic cells (O) and CD11b+ F4/80+ macrophages (P). ns, not significant. Error bars
represent SEM.

520
521 **Figure 3. RIPK3, but not MLKL, restricts Langat virus pathogenesis following intracranial infection.**
522 Survival and body weight analysis from *Ripk3⁻/⁻* (A-B) and *Mlkl⁻/⁻* (C-D) mice and their respective
523 littermate controls following intracranial inoculation with 50 PFU LGTV TP21. Data are pooled from two
524 (A-B) or three (C-D) experiments. ns, not significant. **$p < 0.01$, ***$p < 0.001$.
525

Figure 4. RIPK3 promotes chemokine expression in the cerebellum during LGTV encephalitis.
(A-B) *Ripk3*[-/-] and littermate control mice were infected subcutaneously with LGTV TP21. At 8dpi
cerebral cortical (A) and cerebellar tissues (B) were harvested and assayed for chemokine transcripts via
qRT-PCR. (C-F) *Ccl2* and *Cxcl10* expression in wildtype (C57BL/6J) cultures of primary cortical neurons (C),
cerebellar granule cell neurons (D), cortical astrocytes (E), and cerebellar astrocytes (F) following 2-hour
pretreatment with GSK872 or vehicle and 24h infection with 0.5 (C-D) or 0.01 (E-F) MOI LGTV TP21,
measured via qRT-PCR. ns, not significant. *p<0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001. Error bars
represent SEM.

537
Figure 5. Leukocyte recruitment to the CNS occurs independently of RIPK3 signaling during LGTV
encephalitis.
(A-I) *Ripk3* $^{-/-}$ and littermate control mice were infected subcutaneously with LGTV TP21. Cerebral
cortical and cerebellar tissues were harvested and leukocytes isolated for flow cytometric profiling at
indicated days post infection (dpi). (A) Representative flow cytometry plots showing CD8+ and CD4+ T
cells among CD3+ leukocytes in the brain. Numbers represent percentage of cells in each gate relative to
total plotted cells. (B-I) Numbers of CD8+ T cells (B), CD4+ T cells (C), CD11b+ NK1.1+ natural killer cells
(D), CD11c+ MHC-II+ dendritic cells (E), CD45^{high} CD11b+ F4/80+ macrophages (F), CD11b+ Ly6G+
neutrophils (G), CD45^{high} CD11b+ Ly6C+ monocytes (H), and CD45.2^{lo} CD11b+ microglia (I) among total
brain leukocytes. No comparisons are statistically significant.
548

549
550
Figure 6. RIPK3 limits LGTV replication in cerebellar granule cell neurons.
(A-B) *Ripk3* [-/-] (A) or *Mlkl* [-/-] (B) mice and littermate controls were infected subcutaneously with LGTV
TP21. At 8 or 12 days post infection (dpi), viral loads in cerebral cortical and cerebellar tissues were
determined by plaque assay. Data are pooled from 2-3 independent experiments. (C) *Ripk3* [-/-] and
littermate control mice were subcutaneously infected with LGTV TP21. BBB permeability was measured
at 8 dpi by detection of sodium fluorescein accumulation in tissue homogenates derived from cerebral
cortex or cerebellum. Data represent individual brain fluorescence values normalized to serum sodium
fluorescein concentration. Individual mouse values were then normalized to the mean values for
uninfected controls. (D-G) Multistep growth curve analysis following infection with 0.01 MOI LGTV TP21
in cortical neurons (D), cerebellar granule cell neurons (E), cortical astrocytes (F), and cerebellar
astrocytes (G). n=3 (cerebellar granule cell neurons) or 4 (astrocytes and cortical neurons) for growth
curve experiments. ns, not significant. *p<0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001. Error bars
represent SEM.
564
565
566

567
Figure 7. **RIPK3 promotes ISG expression in cerebellar granule cell neurons.**
(A-B) *Ripk3 -/-* and littermate control mice were infected subcutaneously with LGTV TP21. Transcriptional
expression of indicated genes was assessed via qRT-PCR in cerebral cortical (A) and cerebellar (B) tissues
at 8dpi. (C-D) Transcriptional expression of ISGs in wildtype (C57BL/6J) cultures of primary cortical
neurons (C), cerebellar granule cell neurons (D), cortical astrocytes (E), and cerebellar astrocytes (F)
following 2-hour pretreatment with GSK872 or vehicle and 24-hour infection with 0.5 (C-D) or 0.01 (E-F)
MOI LGTV TP21, measured via qRT-PCR. ns, not significant. *p<0.05, **p < 0.01, ***p < 0.001, ****p <
0.0001. Error bars represent SEM.
576

577
Figure 8. **RIPK3 promotes expression of IFN receptors and IFN-dependent inflammatory genes in**
cerebellar granule cell neurons.
A-B) Transcriptional expression of indicated genes in wildtype (C57BL/6J) cultures of cerebellar granule
cell neurons in the setting of 2-hour pretreatment with GSK872 or vehicle followed by 1 hour treatment
with 10ng/ml IFNβ (A) or 24-hour infection with 0.5 MOI LGTV TP21 (B). C) Expression of indicated genes
in wildtype cerebellar granule cell neurons pretreated for 45 minutes with an anti-IFNAR1 neutralizing
antibody or isotype control +/- cotreatment with GSK872 or vehicle, followed by 24-hour infection with
0.5 MOI LGTV TP21. ns, not significant. *p<0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001. Error bars
represent SEM.
587
588
589

598
Supplemental Figure 2. **RIPK3 does not impact IFN-mediated responses to LGTV in cortical neurons.**
A-B) Transcriptional expression of indicated genes in wildtype (C57BL/6J) cultures of cerebral cortical
neurons in the setting of 2-hour pretreatment with GSK872 or vehicle followed by 1 hour treatment with
10ng/ml IFNβ (A) or 24-hour infection with 0.5 MOI LGTV TP21 (B). C) Expression of indicated genes in
wildtype cerebral cortical neurons pretreated for 45 minutes with an anti-IFNAR1 neutralizing antibody
or isotype control +/- cotreatment with GSK872 or vehicle, followed by 24-hour infection with 0.5 MOI
LGTV TP21. ns, not significant. *p<0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001. Error bars represent
SEM.
607
608

Supplemental Table 1: Primer sequences for qRT-PCR

Gene	Direction	Sequence (5'-3')
18S	Forward	CTTAGAGGGACAAGTGGCG
18S	Reverse	ACGCTGAGCCAGTCAGTGTA
Ccl2	Forward	TGG CTC AGC CAG ATG CAG T
Ccl2	Reverse	TTG GGA TCA TCT TGC TGG TG
Ccl3	Forward	CCA AGT CTT CTC AGC GCC AT
Ccl3	Reverse	TCC GGC TGT AGG AGA AGC AG
Ccl4	Forward	TCT TGC TCG TGG CTG CCT
Ccl4	Reverse	GGG AGG GTC AGA GCC CA
Ccl5	Forward	CCTGCTGCTTTGCCTACCTCTC
Ccl5	Reverse	ACACACTTGGCGGTTCCTTCGA
Cxcl1	Forward	ATCCAGAGCTTGAAGGTGTTG
Cxcl1	Reverse	GTCTGTCTTCTTTCTCCGTTACTT
Cxcl10	Forward	CCCACGTGTTGAGATCATTG
Cxcl10	Reverse	CACTGGGTAAAGGGGAGTGA
Ifit1	Forward	CCCAGAGAACAGCTACCACC
Ifit1	Reverse	TGTGAAGTGACATCTCAGCTGA
Ifna	Forward	CTTCCACAGGATCACTGTGTACCT
Ifna	Reverse	TTCTGCTCTGACCACCTCCC
Ifnar1	Forward	TGTGCTTCCCACCACTCAAG
Ifnar1	Reverse	AGGCGCGTGCTTTACTTCTA
Ifnar2	Forward	AGACTCTTCGGGTCGCGG
Ifnar2	Reverse	GTTTTTCTGCTCTCACACCTGA
Ifnb	Forward	CTGGAGCAGCTGAATGGAAAG
Ifnb	Reverse	CTTCTCCGTCATCTCCATAGGG
Ifng	Forward	CCTCATGGCTGTTTCTGGCT
Ifng	Reverse	TCATGTCACCATCCTTTTGCC
Ifngr1	Forward	GTGGAGCTTTGACGAGCACT
Ifngr1	Reverse	TCAGTCCAGGAACCCGAATA
Ifngr2	Forward	CGTCCTCGCCAGACTCGTT
Ifngr2	Reverse	AGCAACCTATGCCAAGAGCC
Il6	Forward	ACACATGTTCTCTGGGAAATCGT
Il6	Reverse	AAGTGCATCATCGTTGTTCATACA
Isg15	Forward	TGCCTGCAGTTCTGTACCAC
Isg15	Reverse	AGTGCTCCAGGACGGTCTTA
Mx1	Forward	ACTATGAGGAGAAGGTGCGG
Mx1	Reverse	ACTTTGCCTCTCCACTCCTC
Mx2	Forward	GCCACGTTCCCTTGATCATC
Mx2	Reverse	AGCCAGCTTAACCAGGGAAT
Oas1b	Forward	TTCTACGCCAATCTCATCAGTG
Oas1b	Reverse	GGTCCCCCAGCTTCTCCTTAC
Rsad2	Forward	TCAAAAGCTGAGGAGGTGGTG
Rsad2	Reverse	TAGGAGGCACTGGAAAACCTTC

References

1. Schultz JS, Sparks H, Beckham JD. 2021. Arboviral central nervous system infections. Curr Opin Infect Dis 34:264-271.
2. Ruzek D, Avsic Zupanc T, Borde J, Chrdle A, Eyer L, Karganova G, Kholodilov I, Knap N, Kozlovskaya L, Matveev A, Miller AD, Osolodkin DI, Overby AK, Tikunova N, Tkachev S, Zajkowska J. 2019. Tick-borne encephalitis in Europe and Russia: Review of pathogenesis, clinical features, therapy, and vaccines. Antiviral Res 164:23-51.
3. Riccardi N, Antonello RM, Luzzati R, Zajkowska J, Di Bella S, Giacobbe DR. 2019. Tick-borne encephalitis in Europe: a brief update on epidemiology, diagnosis, prevention, and treatment. Eur J Intern Med 62:1-6.
4. Abdiyeva K, Turebekov N, Yegemberdiyeva R, Dmitrovskiy A, Yeraliyeva L, Shapiyeva Z, Nurmakhanov T, Sansyzbayev Y, Froeschl G, Hoelscher M, Zinner J, Essbauer S, Frey S. 2020. Vectors, molecular epidemiology and phylogeny of TBEV in Kazakhstan and central Asia. Parasit Vectors 13:504.
5. Beaute J, Spiteri G, Warns-Petit E, Zeller H. 2018. Tick-borne encephalitis in Europe, 2012 to 2016. Euro Surveill 23.
6. Fares M, Cochet-Bernoin M, Gonzalez G, Montero-Menei CN, Blanchet O, Benchoua A, Boissart C, Lecollinet S, Richardson J, Haddad N, Coulpier M. 2020. Pathological modeling of TBEV infection reveals differential innate immune responses in human neurons and astrocytes that correlate with their susceptibility to infection. J Neuroinflammation 17:76.
7. Szretter KJ, Daffis S, Patel J, Suthar MS, Klein RS, Gale M, Jr., Diamond MS. 2010. The innate immune adaptor molecule MyD88 restricts West Nile virus replication and spread in neurons of the central nervous system. J Virol 84:12125-38.
8. Iwasaki Y, Zhao JX, Yamamoto T, Konno H. 1986. Immunohistochemical demonstration of viral antigens in Japanese encephalitis. Acta Neuropathol 70:79-81.
9. Klein RS, Lin E, Zhang B, Luster AD, Tollett J, Samuel MA, Engle M, Diamond MS. 2005. Neuronal CXCL10 directs CD8+ T-cell recruitment and control of West Nile virus encephalitis. J Virol 79:11457-66.
10. Lindqvist R, Upadhyay A, Overby AK. 2018. Tick-Borne Flaviviruses and the Type I Interferon Response. Viruses 10.
11. Samuel MA, Diamond MS. 2005. Alpha/beta interferon protects against lethal West Nile virus infection by restricting cellular tropism and enhancing neuronal survival. J Virol 79:13350-61.
12. Cho H, Proll SC, Szretter KJ, Katze MG, Gale M, Jr., Diamond MS. 2013. Differential innate immune response programs in neuronal subtypes determine susceptibility to infection in the brain by positive-stranded RNA viruses. Nat Med 19:458-64.
13. Daniels BP, Jujjavarapu H, Durrant DM, Williams JL, Green RR, White JP, Lazear HM, Gale M, Jr., Diamond MS, Klein RS. 2017. Regional astrocyte IFN signaling restricts pathogenesis during neurotropic viral infection. J Clin Invest 127:843-856.
14. Morgan MJ, Kim YS. 2022. Roles of RIPK3 in necroptosis, cell signaling, and disease. Exp Mol Med 54:1695-1704.
15. Samson AL, Zhang Y, Geoghegan ND, Gavin XJ, Davies KA, Mlodzianoski MJ, Whitehead LW, Frank D, Garnish SE, Fitzgibbon C, Hempel A, Young SN, Jacobsen AV, Cawthorne W, Petrie EJ, Faux MC, Shield-Artin K, Lalaoui N, Hildebrand JM, Silke J, Rogers KL, Lessene G, Hawkins ED, Murphy JM. 2020. MLKL trafficking and accumulation at the plasma membrane control the kinetics and threshold for necroptosis. Nat Commun 11:3151.

16. Daniels BP, Snyder AG, Olsen TM, Orozco S, Oguin TH, 3rd, Tait SWG, Martinez J, Gale M, Jr., Loo YM, Oberst A. 2017. RIPK3 Restricts Viral Pathogenesis via Cell Death-Independent Neuroinflammation. Cell 169:301-313 e11.

17. Daniels BP, Kofman SB, Smith JR, Norris GT, Snyder AG, Kolb JP, Gao X, Locasale JW, Martinez J, Gale M, Jr., Loo YM, Oberst A. 2019. The Nucleotide Sensor ZBP1 and Kinase RIPK3 Induce the Enzyme IRG1 to Promote an Antiviral Metabolic State in Neurons. Immunity 50:64-76 e4.

18. Downey J, Pernet E, Coulombe F, Allard B, Meunier I, Jaworska J, Qureshi S, Vinh DC, Martin JG, Joubert P, Divangahi M. 2017. RIPK3 interacts with MAVS to regulate type I IFN-mediated immunity to Influenza A virus infection. PLoS Pathog 13:e1006326.

19. Saleh D, Najjar M, Zelic M, Shah S, Nogusa S, Polykratis A, Paczosa MK, Gough PJ, Bertin J, Whalen M, Fitzgerald KA, Slavov N, Pasparakis M, Balachandran S, Kelliher M, Mecsas J, Degterev A. 2017. Kinase Activities of RIPK1 and RIPK3 Can Direct IFN-beta Synthesis Induced by Lipopolysaccharide. J Immunol 198:4435-4447.

20. Guo H, Koehler HS, Mocarski ES, Dix RD. 2022. RIPK3 and caspase 8 collaborate to limit herpes simplex encephalitis. PLoS Pathog 18:e1010857.

21. Peng R, Wang CK, Wang-Kan X, Idorn M, Kjaer M, Zhou FY, Fiil BK, Timmermann F, Orozco SL, McCarthy J, Leung CS, Lu X, Bagola K, Rehwinkel J, Oberst A, Maelfait J, Paludan SR, Gyrd-Hansen M. 2022. Human ZBP1 induces cell death-independent inflammatory signaling via RIPK3 and RIPK1. EMBO Rep 23:e55839.

22. Chou TW, Chang NP, Krishnagiri M, Patel AP, Lindman M, Angel JP, Kung PL, Atkins C, Daniels BP. 2021. Fibrillar alpha-synuclein induces neurotoxic astrocyte activation via RIP kinase signaling and NF-kappaB. Cell Death Dis 12:756.

23. Li S, Zhang Y, Guan Z, Ye M, Li H, You M, Zhou Z, Zhang C, Zhang F, Lu B, Zhou P, Peng K. 2023. SARS-CoV-2 Z-RNA activates the ZBP1-RIPK3 pathway to promote virus-induced inflammatory responses. Cell Res doi:10.1038/s41422-022-00775-y.

24. Najjar M, Saleh D, Zelic M, Nogusa S, Shah S, Tai A, Finger JN, Polykratis A, Gough PJ, Bertin J, Whalen M, Pasparakis M, Balachandran S, Kelliher M, Poltorak A, Degterev A. 2016. RIPK1 and RIPK3 Kinases Promote Cell-Death-Independent Inflammation by Toll-like Receptor 4. Immunity 45:46-59.

25. Hubbard NW, Ames JM, Maurano M, Chu LH, Somfleth KY, Gokhale NS, Werner M, Snyder JM, Lichauco K, Savan R, Stetson DB, Oberst A. 2022. ADAR1 mutation causes ZBP1-dependent immunopathology. Nature 607:769-775.

26. Xie Y, Zhao Y, Shi L, Li W, Chen K, Li M, Chen X, Zhang H, Li T, Matsuzawa-Ishimoto Y, Yao X, Shao D, Ke Z, Li J, Chen Y, Zhang X, Cui J, Cui S, Leng Q, Cadwell K, Li X, Wei H, Zhang H, Li H, Xiao H. 2020. Gut epithelial TSC1/mTOR controls RIPK3-dependent necroptosis in intestinal inflammation and cancer. J Clin Invest 130:2111-2128.

27. Sai K, Parsons C, House JS, Kathariou S, Ninomiya-Tsuji J. 2019. Necroptosis mediators RIPK3 and MLKL suppress intracellular Listeria replication independently of host cell killing. J Cell Biol 218:1994-2005.

28. Baker DG, Woods TA, Butchi NB, Morgan TM, Taylor RT, Sunyakumthorn P, Mukherjee P, Lubick KJ, Best SM, Peterson KE. 2013. Toll-like receptor 7 suppresses virus replication in neurons but does not affect viral pathogenesis in a mouse model of Langat virus infection. J Gen Virol 94:336-347.

29. Michlmayr D, Bardina SV, Rodriguez CA, Pletnev AG, Lim JK. 2016. Dual Function of Ccr5 during Langat Virus Encephalitis: Reduction in Neutrophil-Mediated Central Nervous System Inflammation and Increase in T Cell-Mediated Viral Clearance. J Immunol 196:4622-31.

30. Kimura T, Katoh H, Kayama H, Saiga H, Okuyama M, Okamoto T, Umemoto E, Matsuura Y, Yamamoto M, Takeda K. 2013. Ifit1 inhibits Japanese encephalitis virus replication through binding to 5' capped 2'-O unmethylated RNA. J Virol 87:9997-10003.

31. Szretter KJ, Daniels BP, Cho H, Gainey MD, Yokoyama WM, Gale M, Jr., Virgin HW, Klein RS, Sen GC, Diamond MS. 2012. 2'-O methylation of the viral mRNA cap by West Nile virus evades ifit1-dependent and -independent mechanisms of host restriction in vivo. PLoS Pathog 8:e1002698.

32. Dai J, Pan W, Wang P. 2011. ISG15 facilitates cellular antiviral response to dengue and west nile virus infection in vitro. Virol J 8:468.

33. Singh PK, Singh S, Farr D, Kumar A. 2019. Interferon-stimulated gene 15 (ISG15) restricts Zika virus replication in primary human corneal epithelial cells. Ocul Surf 17:551-559.

34. Bigham AW, Buckingham KJ, Husain S, Emond MJ, Bofferding KM, Gildersleeve H, Rutherford A, Astakhova NM, Perelygin AA, Busch MP, Murray KO, Sejvar JJ, Green S, Kriesel J, Brinton MA, Bamshad M. 2011. Host genetic risk factors for West Nile virus infection and disease progression. PLoS One 6:e24745.

35. Vonderstein K, Nilsson E, Hubel P, Nygard Skalman L, Upadhyay A, Pasto J, Pichlmair A, Lundmark R, Overby AK. 2018. Viperin Targets Flavivirus Virulence by Inducing Assembly of Noninfectious Capsid Particles. J Virol 92.

36. Williams JL, Manivasagam S, Smith BC, Sim J, Vollmer LL, Daniels BP, Russell JH, Klein RS. 2020. Astrocyte-T cell crosstalk regulates region-specific neuroinflammation. Glia 68:1361-1374.

37. Simmons SB, Liggitt D, Goverman JM. 2014. Cytokine-regulated neutrophil recruitment is required for brain but not spinal cord inflammation during experimental autoimmune encephalomyelitis. J Immunol 193:555-63.

38. Pierson ER, Goverman JM. 2017. GM-CSF is not essential for experimental autoimmune encephalomyelitis but promotes brain-targeted disease. JCI Insight 2:e92362.

39. Durrant DM, Daniels BP, Pasieka T, Dorsey D, Klein RS. 2015. CCR5 limits cortical viral loads during West Nile virus infection of the central nervous system. J Neuroinflammation 12:233.

40. McComb S, Cessford E, Alturki NA, Joseph J, Shutinoski B, Startek JB, Gamero AM, Mossman KL, Sad S. 2014. Type-I interferon signaling through ISGF3 complex is required for sustained Rip3 activation and necroptosis in macrophages. Proc Natl Acad Sci U S A 111:E3206-13.

41. Brault M, Olsen TM, Martinez J, Stetson DB, Oberst A. 2018. Intracellular Nucleic Acid Sensing Triggers Necroptosis through Synergistic Type I IFN and TNF Signaling. J Immunol 200:2748-2756.

42. Ingram JP, Thapa RJ, Fisher A, Tummers B, Zhang T, Yin C, Rodriguez DA, Guo H, Lane R, Williams R, Slifker MJ, Basagoudanavar SH, Rall GF, Dillon CP, Green DR, Kaiser WJ, Balachandran S. 2019. ZBP1/DAI Drives RIPK3-Mediated Cell Death Induced by IFNs in the Absence of RIPK1. J Immunol 203:1348-1355.

43. Lee SA, Chang LC, Jung W, Bowman JW, Kim D, Chen W, Foo SS, Choi YJ, Choi UY, Bowling A, Yoo JS, Jung JU. 2023. OASL phase condensation induces amyloid-like fibrillation of RIPK3 to promote virus-induced necroptosis. Nat Cell Biol doi:10.1038/s41556-022-01039-y.

44. Yatim N, Jusforgues-Saklani H, Orozco S, Schulz O, Barreira da Silva R, Reis e Sousa C, Green DR, Oberst A, Albert ML. 2015. RIPK1 and NF-kappaB signaling in dying cells determines cross-priming of CD8(+) T cells. Science 350:328-34.

45. Snyder AG, Hubbard NW, Messmer MN, Kofman SB, Hagan CE, Orozco SL, Chiang K,
Daniels BP, Baker D, Oberst A. 2019. Intratumoral activation of the necroptotic pathway
components RIPK1 and RIPK3 potentiates antitumor immunity. Sci Immunol 4.
46. Wegner KW, Saleh D, Degterev A. 2017. Complex Pathologic Roles of RIPK1 and
RIPK3: Moving Beyond Necroptosis. Trends Pharmacol Sci 38:202-225.
47. Angel JP, Daniels BP. 2022. Paradoxical roles for programmed cell death signaling
during viral infection of the central nervous system. Curr Opin Neurobiol 77:102629.
48. Daniels BP, Oberst A. 2020. Outcomes of RIP Kinase Signaling During Neuroinvasive
Viral Infection. Curr Top Microbiol Immunol doi:10.1007/82_2020_204.
49. Bian P, Ye C, Zheng X, Luo C, Yang J, Li M, Wang Y, Yang J, Zhou Y, Zhang F, Lian J,
Zhang Y, Jia Z, Lei Y. 2020. RIPK3 Promotes JEV Replication in Neurons via
Downregulation of IFI44L. Front Microbiol 11:368.
50. Bian P, Zheng X, Wei L, Ye C, Fan H, Cai Y, Zhang Y, Zhang F, Jia Z, Lei Y. 2017. MLKL
Mediated Necroptosis Accelerates JEV-Induced Neuroinflammation in Mice. Front
Microbiol 8:303.
51. Newton K, Sun X, Dixit VM. 2004. Kinase RIP3 is dispensable for normal NF-kappa Bs,
signaling by the B-cell and T-cell receptors, tumor necrosis factor receptor 1, and Toll-like
receptors 2 and 4. Mol Cell Biol 24:1464-9.
52. Murphy JM, Czabotar PE, Hildebrand JM, Lucet IS, Zhang JG, Alvarez-Diaz S, Lewis R,
Lalaoui N, Metcalf D, Webb AI, Young SN, Varghese LN, Tannahill GM, Hatchell EC,
Majewski IJ, Okamoto T, Dobson RC, Hilton DJ, Babon JJ, Nicola NA, Strasser A, Silke
J, Alexander WS. 2013. The pseudokinase MLKL mediates necroptosis via a molecular
switch mechanism. Immunity 39:443-53.
53. Kung PL, Chou TW, Lindman M, Chang NP, Estevez I, Buckley BD, Atkins C, Daniels
BP. 2022. Zika virus-induced TNF-alpha signaling dysregulates expression of neurologic
genes associated with psychiatric disorders. J Neuroinflammation 19:100.
54. Daniels BP, Holman DW, Cruz-Orengo L, Jujjavarapu H, Durrant DM, Klein RS. 2014.
Viral pathogen-associated molecular patterns regulate blood-brain barrier integrity via
competing innate cytokine signals. mBio 5:e01476-14.
784