The effects of social experience on host gut microbiome in male zebrafish (*Danio rerio*)

Emily Scott, Michael S Brewer, Ariane L Peralta[*], Fadi A Issa[*]

Department of Biology, East Carolina University, Greenville, NC 27858

*Corresponding authors:

Dr. Fadi A. Issa, East Carolina University, Department of Biology, 101 E 10th St., Greenville, NC,

Email: issaf14@ecu.edu

Dr. Ariane Peralta, East Carolina University, Department of Biology, 101 E 10th St., Greenville,

NC, Email: peraltaa@ecu.edu

ABSTRACT

Although the gut and the brain vastly differ in physiological function, they have been interlinked in a variety of different neurological and behavioral disorders. The bacteria that comprise the gut microbiome communicate and influence the function of various physiological processes within the body including nervous system function. However, the effects of social experience in the context of dominance and chronic stress on gut microbiome remain poorly understood. Here, we examined whether social experience impacts the host zebrafish (*Danio rerio*) gut microbiome. We studied how social dominance during the first two weeks of social interactions changed the composition of zebrafish gut microbiome by comparing gut bacterial composition, diversity and relative abundance among socially dominant, submissive, social isolates, and control group-housed communal fish. Using amplicon sequencing of the 16S rRNA gene, we report that social dominance significantly affects host gut bacterial community composition but not bacterial diversity. At the genus-level, *Aeromonas* and unclassified Enterobacteriaceae relative abundance decreased in dominant individuals while commensal bacteria (e.g., *Exiguobacterium* and *Cetobacterium)* increased in relative abundance. Conversely, the relative abundance of *Psychrobacter* and *Acinetobacter* was increased in subordinates, isolates, and communal fish compared to control group-housed fish. The shift in commensal and pathogenic bacteria highlights the impact of social experience and the accompanying stress on gut microbiome with potentially similar effects in other social organisms.

IMPORTANCE

Disruptions in the gut microbiome negatively impact various systems in the body. Recently, the

gut microbiome has been associated with neurological deficits in both behavioral and

neurodegenerative disorders. Given the increasing prevalence in diagnosis of both neurological

disease and behavioral disorders, researching the link between social behaviors and the gut

microbiome is critical to better understand how the gut and the brain communicate during

healthy and diseased states. Our research findings demonstrate the effects of social dominance

and chronic stress on host gut microbiome composition. By identifying variations in bacterial

relative abundance based on social experience and associated stress, a broader understanding

of *pathogenic* (e.g., Enterobacteriaceae*, Aeromonas*) versus commensal communities (e.g.,

Exiguobacterium, Cetobacterium) and related host physiology can be inferred.

INTRODUCTION

The gut microbiome plays key roles in biochemical functions in vertebrates. Gut bacterial metabolism has been linked to many systemic diseases in humans and influences numerous immune system pathways (1–5). Additionally, commensal bacteria within the gut microbiome serves as a line of defense to viral infections through interferon signaling (6). Along with these known functions of the gut biome, it has recently been linked to affecting brain function in the context of neurodevelopmental (7, 8), behavioral (9–11), and neurodegenerative disorders (12–14). For instance, the gut microbiome produces certain biochemicals (i.e., dopamine and oxytocin) that regulate neural nuclei involved in social behavior (15, 16). This link is known as the "microbiota-gut-brain axis" and its dysfunction has been associated with many behavioral disorders such as autism spectrum disorder, depression, and anxiety with long-lasting negative impact on health and wellbeing (17). Thus, understanding the mechanisms of disease onset and progression related to the gut microbiome is critical.

The typical gut microbiome consists of both commensal and pathogenic bacteria (18–20). While it is known that pathogenic bacteria are more prevalent in gut microbiomes of diseased organisms, evidence from multiple animal models has shown that a lack of commensal bacteria can also contribute to the detrimental progression of various behavioral disorders (21, 22). In Autism Spectrum Disorder mouse models, the abundance of commensal bacteria species, *Lactobacillus reuteri* is significantly reduced in comparison to wild type mice (15). Further, *Lactobacillus plantarum* has been used as a probiotic in zebrafish models to reduce stress-related behaviors and prevent stress-induced microbiome dysbiosis (23). *L. plantarum* was discovered to modulate anxiety related behavior through the GABAergic and serotonergic pathways (23). Additionally, gut bacterial composition influences host behavior through the modulation of Brain-Derived Neurotrophic Factor (BDNF) levels and serotonin metabolism (24). For instance, probiotic treatment with *Lactobacillus rhamnosus* in zebrafish affects shoaling behavior and brain expression levels of *bdnf* and serotoninergic pathways both

of which have been implicated in aggression and multiple psychiatric disorders (25, 26). In

addition to influencing neurotransmitters levels, bacteria from the genera *Lactobacillus* can

activate afferent neurons in the intestine that modulate pain sensation and actions of defensive

behavior in response to stress (27). This evidence supports the notion that certain bacteria can

influence the activity of neuromodulatory pathways responsible for social behavior. Collectively,

these relationships between host gut bacteria and nervous system function support the notion

that the gut microbiome regulates social activity by influencing neuromodulatory pathways

responsible for social behavior.

Zebrafish social behavior has been studied extensively (28–32). When paired, zebrafish

interact agonistically with conspecifics and quickly form stable dominance relationships

consisting of dominant and submissive fish (28, 33). Once dominance forms, dominants have

priority to food, shelter, and mates while submissive fish experience chronic stress (34).

Although the neurobiological bases of zebrafish social aggression and their long-term impact on

social activity have been investigated; the effects of social dominance on gut microbiome are

less understood (35). This is of particular importance because the effects of chronic social

stress on organisms are detrimental and have been adversely linked to anxiety-like behavior

relevant for survival in many social species (2, 36–40). Specifically, reduction of glucocorticoid

signaling activity (caused by insufficient hormone availability) in chronic stress models has been

associated with stress-related disorders (41). Interestingly, the gut microbiota has recently been

thought to modulate downstream glucocorticoid receptors in the hippocampus resulting in

behavioral abnormalities highlighting the relationship between stress and the gut microbiome

(42). The stability of zebrafish social dominance and ease of quantifying their aggressive social

activity provides an opportunity to examine the brain-gut microbiome axes and the impact of

social stress on the gut biome as dominance relationships are established.

The zebrafish gut microbiome has been examined in embryos and adults. A past study

showed that the gut microbiome composition of host zebrafish changes throughout its

development (43). Specifically, as embryonic zebrafish develop, the diversity of gut bacteria

decreases significantly in terms of the number of operational taxonomic units (OTUs), which are

clusters of bacteria that exhibit high sequence similarity of the 16S rRNA gene (44, 45).

However, microbiome composition was similar between stages of adult fish, which indicates that

major gut microbial changes occur before and during major developmental changes such as

sexual differentiation (44). The adult zebrafish gut microbiome is composed primarily of bacteria

within the Proteobacteria, Firmicutes, and Fusobacteria phyla (46). Pathogenic bacteria within

the Proteobacteria phylum are found in significantly lower relative abundance of healthy human

gut microbiomes and higher relative abundance of Proteobacterial members can be used as a

diagnostic for disease and dysbiosis (47). Additionally, in zebrafish, Proteobacteria including

pathogenic genera *Vibrio* and *Plesiomona* were reduced in probiotic treated fish as opposed to

commensal Firmicutes bacterial members (24). Thus, the presence of pathogenic bacteria in

both zebrafish and humans can indicate microbial dysbiosis and potential health issues. In this

study, we examined whether social dominance has opposing effects on gut microbiome in

dominant versus chronically stressed submissive animals. We tested the hypothesis that the

relative abundance, diversity, and overall composition of zebrafish gut microbiome is socially

regulated. We compared gut microbiomes (based on amplicon sequencing of the 16S rRNA

gene) among socially dominant, submissive, socially isolated, and group-housed communal

fish. Results from this work improve our understanding of how social aggression and stress

impact host gut microbiome and may help future studies to understand the gut-brain axis and

how the onset of behavioral disorders affect gut dysbiosis.

Experimental Methods

Pairing and behavioral analysis:

Male zebrafish (n=24) were isolated in individual tanks for 7 days to minimize the effects of prior social experience. Then, 12 fish were randomly paired and placed into new tanks while 6 fish remained in isolation, and the remaining 6 fish were placed into one new tank to serve as communal controls. Over a period of 14 days, the 6 zebrafish pairs self-established rank and their behavior was monitored and recorded over a course of 14 days (2 weeks). Behavior was observed visually, and written observations were made regarding dominant and subordinate behavior among the paired and communal fish. Specifically, the number of attacks and retreats by each fish were recorded during a 5-minute observation period 3 times a week. Additionally, tank locations of each fish (top vs. bottom of tank) were monitored and recorded as stressed or subordinate fish tend to stay lower in the tank and more aggressive/dominant fish claim the top of the tank. All animal methods and protocols were approved by East Carolina University's Institutional Animal Care and Use Committee (IACUC) AUP# D320b.

Fecal sample collection:

Initial fecal samples were collected from each fish on day 3 of the isolation period. This fecal collection before pairing or establishment of behaviors served as a method of comparison for later fecal collections. After pairing of fish after the 7-day isolation period, fecal samples were collected on day 0 (day of pairing), day 7 (one week after pairing), and day 14 (2 weeks after pairing). Collections were made throughout the progression of social hierarchal development to determine whether the gut microbiome composition of host fish evolves as dominance was established.

On days of fecal collection, fish were removed from tanks and each fish was placed into individual containers to prevent any cross contamination of fecal matter. Before pairing, each fish was carefully observed and distinct markings, size differences, and color differences were

noted to ensure easy recognition of each fish upon isolation and fecal sample collection. 2-4

hours after separation and feeding each fish, fecal samples were collected. Fecal samples from

each fish were extracted using sterile micropipette tips and were placed into 500 microliters of

DNAse free water. Gloves were always worn when handling fecal samples and fish, to minimize

any cross contamination. After samples were collected, they were stored at -80 °C until DNA

extractions and processing were performed.

Microbiome Analysis:

To determine differences in microbial composition in zebrafish based on social status,

microbial communities in each fecal sample were characterized with Illumina sequencing of the

highly conserved 16S rRNA gene (48). Following fecal sample collection, DNA extractions were

performed on each sample using the PowerLyzer PowerSoil DNA protocol. Approximately 0.02g

of feces were collected from each fish for extractions. This extracted genomic DNA was used as

the template for PCR reactions, where barcoded primers (515F-806RB) were used to target the

V4 region of the 16S rRNA gene (48, 49). Each fecal sample was run in triplicate PCR reactions

and then combined and purified using the Axygen® AxyPrep magnetic bead purification kit

(Corning Life Sciences). After successful cleanup, DNA concentrations (ng/mL) from each

sample were quantified using the Quant-iT dsDNA HS (high sensitivity) assay (Thermo

Scientific, Waltham, Massachusetts, USA). Recorded DNA concentrations (ng/mL) using HS

measurements were converted to the final DNA concentration (ng/μl) and then dilutions using

PCR grade water were performed to ensure all samples added to the final pooled product were

the same mass (ng). PCR products were pooled in equimolar concentrations and sequenced on

an Illumina MiSeq platform using paired-end reads (Illumina Reagent Kit v2, 500 reaction kit) at

the Center for Genomics and Bioinformatics at Indiana University.

We processed raw 16S rRNA gene sequences using a standard mothur pipeline

(v1.40.1) (50, 51). We assembled contigs from paired end reads and quality trimmed using a

moving average quality score of 35 bp. We aligned sequences to the SILVA rRNA gene

database (v.128) (52), and we removed chimeric sequences using the VSEARCH algorithm

(53). We divided sequences based on taxonomic class and binned into operational taxonomic

units (OTUs) with a 97% sequence similarity level, and we classified OTUs using the SILVA

rRNA database (v128). We used an OTU-based 3% distance threshold to avoid splitting the

same bacterial genome into distinct clusters (using amplicon sequence variant (ASV) of a single

base difference). In addition, the sequencing was conducted using short-reads (250 bp) and not

assembled genomes, which introduces PCR and sequencing errors which can result in over-

inflated number of ASVs (54). The broadscale community composition patterns are robust when

employing an OTU-based approach (55).

Species Diversity and Community Compositional Analysis:

We examined the diversity and composition of bacterial communities in each fecal

sample. For each microbiome sample, we calculated the Chao1 OTU richness using the

estimateR() function and Shannon diversity using the diversity() function in the vegan package

(56), and also calculated Simpson's evenness using a custom function after we rarefied the

OTU table to 10,181 observations. We calculated the Bray-Curtis dissimilarity matrix and

visualized community composition according to 'Social Status' and 'Day' using principal

Coordinate Analysis (PCoA). Finally, genus level compositions with relative abundances greater

than 0.05 were plotted after OTU table was rarefied to 6,000 observations.

Statistical Analyses:

All data analyses were completed using the R statistical environment (R v4.2.0, R Core

Development Team 2022; R Studio v2022.07.1). Using the lmer() function from the lmerTest

package (57), a linear mixed effects model with 'Day' and 'Social Status' as fixed effects was

used to analyze bacterial diversity metrics Chao1 OTU richness, Shannon diversity (H'), and

Simpson's evenness. We ran a permutational multivariate analysis of variance (PERMANOVA)

based on the Bray-Curtis dissimilarity of bacterial community composition to determine the

extent that 'Social Status' and 'Day' and the interaction explained bacterial community

composition. We next identified bacterial species that represented each treatment ('Social

Status x 'Day') for bacterial taxa with a relative abundance greater than 0.05 when summed

across all samples. We performed the PERMANOVA using the adonis() function from the vegan

package (56) and the indval() function in the indicspecies package (58).

Data Availability

All code and data used in this study are in a public GitHub repository

(https://github.com/PeraltaLab/ZebrafishMicrobiomes_SocialStatus) and NCBI SRA BioProject

PRJNA925886.

RESULTS

Bacterial species diversity and community composition

We compared gut microbiomes of zebrafish of different social statuses. Bacterial OTU richness (Fig. 1) was similar across 'Social Status' and 'Day', while the interaction of 'Social Status' and 'Day' but not the main effects influenced Shannon diversity (Fig. 2) and Simpson's evenness increased from Day 7 to Day 14 (Fig. 3) (Table S1). While statistical significance was detected, the linear model explained a relatively low amount of variation (richness: adjusted R^2= 0.1049, p-value=0.0645; diversity: adjusted R^2=0.1089, p-value=0.0585; evenness: adjusted R^2=0.1547, p-value=0.0171) (Table S1). Subordinate data points plotted similarly to isolate data points during the isolation period (IP) before any pairing or rank establishment. Additionally, dominants data points trended similarly to those of communal animals during the isolation period. These trends were observed for richness (Fig. 1), diversity (Fig. 2), and evenness (Fig. 3).

In contrast to bacterial diversity, 'Social Status' and 'Day' influenced bacterial community composition. The interaction of 'Social Status' and 'Day' accounted for 14.4% of the variation in bacterial composition ($F_{9,92}$=1.9214, P=0.001), while the main effects of 'Day' accounted for 12.5% ($F_{3,92}$=5.035, P=0.001) and 'Social Status' accounted for 9.1% ($F_{3,92}$=3.671, P=0.001) of the bacterial community composition (Fig. 4, Table S2). The bacterial gut microbiomes of subordinate and communal fish tended to group together while the gut microbiomes of dominant fish were distinct from all other groups, except for Day 14 which grouped by the gut microbiomes of isolated fish on Day 0 (Fig. 4). The isolation period (IP) for subordinate, communal, and isolated fish resulted in similar gut microbiomes (diamond symbols). For subordinate and isolated fish (blue and gray symbols), the gut microbiomes changed over time more than the dominant and communal gut microbiomes (red and purple symbols) (Fig. 4).

Figure 1

Figure 1. Boxplots representing bacterial Chao 1 richness in individuals from dominant (red),

subordinate (blue), communal (purple), and isolate (gray) animals throughout the isolation

period (Day IP) representing collection before pairing and pairing period (Day 0, Day 7, Day 14).

The boxplot is a visual representation of 5 key summary statistics: the median, the 25% and

75% percentiles, and the whiskers which represent the feasible range of the data as determined

by 1.5 × the interquartile range. Symbols represent individual raw data points from six replicate

samples. Summary of statistical output in Appendix S1: Table S1A.

Figure 2

Figure 2. Boxplots representing bacterial Shannon diversity index in individuals from dominant

(red), subordinate (blue), communal (purple), and isolate (gray) animals throughout the isolation

period (Day IP) representing collection before pairing and pairing period (Day 0, Day 7, Day 14).

The boxplot is a visual representation of 5 key summary statistics: the median, the 25% and

75% percentiles, and the whiskers which represent the feasible range of the data as determined

by 1.5 × the interquartile range. Symbols represent individual raw data points from six replicate

samples. Summary of statistical output in Appendix S1: Table S1B.

Figure 3

Figure 3. Boxplots representing bacterial Simpson's diversity index in individuals from dominant

(red), subordinate (blue), communal (purple), and isolate (gray) animals throughout the isolation

period (Day IP) representing collection before pairing and pairing period (Day 0, Day 7, Day 14).

The boxplot is a visual representation of 5 key summary statistics: the median, the 25% and

75% percentiles, and the whiskers which represent the feasible range of the data as determined

by 1.5 × the interquartile range. Symbols represent individual raw data points from six replicate

samples. Summary of statistical output in Appendix S1: Table S1B.

Figure 4

Social Status Zebrafish Gut Microbiomes

Figure 4. Ordination based on a Principal Coordinates Analysis depicting bacterial community

composition according to fertilization and hydrology. Symbols are colored according to social

status (dominant=red, subordinate=blue, communal=purple, and isolate=gray) and shapes

represent day of pairing (Day 0=square, Day 7=circle, Day 14=triangle, Day IP=diamond). The

centroid and standard error bars (along axes 1 and 2) were calculated for six replicate plots.

Summary of statistical output in Appendix S1: Table S2.

    Taxa-level Shifts Across Social Status

We evaluated which bacteria taxa represented gut microbiomes by 'Social Status' and

'Day' using indicator species analysis. *Paracoccus spp.* and *Achromobacter spp.* represented

gut microbiomes of dominant fish during the isolation period, and unclassified taxa from the

families Planctomycetacea, Caldilineaceae, Comamonadaceae and indicators *Staphylococcus*

spp. and *Exiguobacterium spp.* represented gut microbiomes of dominant fish on Day 7 (Table

S3). *Acinetobacter spp.* and *Psychrobacter spp.* represented gut microbiomes of subordinate

fish during Day 0 and Day 7, respectively (Table S3). Unclassified taxa from the class

Betaproteobacteria and family Rhizobiaceae represented gut microbiomes of isolated fish on

Day 0 (Table S3). For communal fish, an unclassified taxon from the Rhodobacteraceae family,

Pseudomonas spp., and *Stappia spp.* represented gut microbiomes during the isolation period,

while *Rhizobiales spp., Arthrobacter spp.*, and *Acinetobacter spp.* represented Day 0,

Chryseobacterium spp. represented Day 7, and *Shewanella spp.* represented Day 14 (Table

S3). Unclassified taxa from the class Betaproteobacteria and family Rhizobiaceae represented

gut microbiomes of isolated fish on Day 0 (Table S3).

To further explore trends, we evaluated genus level composition for genera observed in >

5% relative abundance. While community composition varied between different social statuses

and day of pairing, we observed changes in dominant individuals on day 7 and 14 of pairing

(Fig. S1). We observed a reduction in pathogenic genera that were consistently high in relative

abundance in non-dominant social status groups and throughout the pairing period. Specifically,

Aeromonas and unclassified Enterobacteriaceae relative abundance was lower in dominant

individuals on day 7 and 14, whereas relative abundance of these genera remained higher in

other groups (Fig. S1). We also observed an increased relative abundance of *Chitinobacteria,*

Vibrio, and *Pseudomonas spp.* in subordinate, isolate, and communal animals and the presence

of these genera were maintained before and during pairing (Fig. S1). In addition, certain

commensal genera such as *Exiguobacterium* and *Cetobacterium* increased in relative

abundance in dominant animals on day 7 of pairing. These genera were present in dominant

animals throughout pairing, but on day 7, the relative abundance increased, especially for the

Cetobacterium genus (Fig. S1).

Discussion

A growing number of studies highlight the microbial role in the link between the gut and

the brain (59–61). Our results suggest a link between the host gut microbiome and social

experience in zebrafish building on prior research that gut microbiome may be associated with

social activity and behavioral disorders. Here, we studied whether social aggression and chronic

stress influence gut microbiome content in zebrafish. We showed that while there were no

significant differences in gut microbiome diversity or richness among different social states,

social dominance leads to compositional differences. This result complements prior work in a

mouse model that showed a causative relationship between the gut and anxiety/depression

phenotypes (15). These similarities along with findings in other studies examining how probiotic

treatment alters behavioral activity in socially stressed individuals lends further credence of the

strong relationship between the gut-brain axis. Specifically, chronically stressed zebrafish that

were treated with probiotics, experienced a complete reversal and rescue of their anxiety-like

behavior (23). This supports a microbiome mediated behavior hypothesis in which changes to

the gut microbiome affects associated behaviors related to stress and anxiety. However, in

those studies, both the composition and diversity of the gut microbiome directly affects behavior

whereas we found that only bacterial composition was affected by social status.

While we observed compositional differences between fish of different social rank, these

differences were more pronounced after 7-14 days of social interactions when social dominance

solidified. This signifies that the chronic stress that accompanies social submission or isolation

may directly impact bacterial composition. Additionally, it seems that social dominance and

aggression promote an increased presence of commensal bacterial genera. Previous research

in autism models show an inverse relationship between the gut and behavior, as individuals

born with gut microbiota dysbiosis are predisposed to social deficits and autism-like behaviors

(62).

One of the questions we sought to answer in this study was whether inherent differences

in gut microbiome prior to social dominance formation would predispose individuals to become

either dominant or submissive. Based on our results of fecal samples that were collected prior to

pairing during the isolation period (Day IP), this hypothesis is not supported in this model.

However, it would be interesting to determine in future studies whether repeated probiotic or

antibiotic treatments over extended periods are sufficient to modulate aggression levels and

reverse social dominance. Such results would be very exciting and would strongly indicate that

the microbiome could directly affect social aggression.

Based on previous research in zebrafish and other animal models, the genus,

Lactobacillus, from phylum Bacteroidetes, is a common protective bacterial species often found

in the guts of many animal models (23). Because of these protective qualities, the *Lactobacillus*

genus is often used as a probiotic treatment in both zebrafish and mouse models to rescue

stressed phenotypes (15, 23). Interestingly, the *Lactobacillus* genus was not present in the

taxonomic analysis of this experiment. This is likely due to *Lactobacillus* species being a

transient to the gut microbiome, as they primarily colonize the host GI tract from food intake and

the oral cavity (63–65). The only genus of bacteria found from the Bacteroidetes phylum was

*Chryseobacterium spp. a*nd was primarily found in communal fish on day 7 (Table S3). While

this genus doesn't come from the same family as *Lactobacillus*, both genera seem to have

similar protective qualities in zebrafish gut microbiota. In a study examining bacterial key players

in pathogenic infection in zebrafish, *Chryseobacterium massillae* was determined to be

important in protecting both larvae and adult zebrafish from pathogenic infection (66).

Interestingly, this bacterial genus was only present in low relative abundance in both communal

and dominant individuals (Table S3, Fig. S1).

Within the Firmicutes phylum, the bacterial genus *Exiguobacterium* was also determined

as a component of compositional analysis of the zebrafish gut in our study. This genus of

bacteria has been shown to produce cyclic dipeptides, which have many antimicrobial,

antifungal, antiviral, and anti-inflammatory properties in humans and other animals (67, 68).

Intriguingly, *Exiguobacterium acetylicum* has therapeutic properties in zebrafish colorectal

cancer models, which highlights the protective properties of this genus of bacteria in the gut of

both zebrafish and humans (69). While the bacterial genus *Exiguobacterium* was present in

socially submissive and isolated zebrafish, this OTU only increased in relative abundance in

dominant individuals after a week of pairing (Fig. S1). The absence of the bacterial genus

Exiguobacterium in communal animals should be noted, which may be due to the heterogeneity

among mid-ranked members living in a communal setting whereby individuals may dominate

some low-ranking animals while being submissive to high-ranking opponents.

Given the high percent of Proteobacteria communities in zebrafish gut microbiomes, the

presence of pathogenic bacteria is not surprising, particularly since a large portion of the

species in these pathogenic genera are waterborne. However, the varying relative abundance of

certain pathogenic bacteria like *Aeromonas,* unclassified Enterobacteriaceae*, Staphylococcus,*

Vibrio, and *Pseudomonas spp.* in the community composition is noteworthy. *Aeromonas and*

unclassified Enterobacteriaceae were the dominant pathogenic genera in all groups and at all

time points during pairing, but several pathogenic genera were primarily seen in subordinate,

isolate, and even communal individuals (Fig. S1). Many of these genera have been used in

zebrafish infection models. A model of *Staphylococcus* infection in zebrafish has been studied,

and the interactions between *Staphylococcus aureus* and other bacteria in the gut to create

infection (70). This study specifically examined the immune response of the fish in response to

this infection, but it is important to note the pathogenesis associated with *Staphylococcus*

infection. Interestingly, *Staphylococcus* relative abundance is increased in dominant individuals

even though other pathogenic genera are decreased. This could contribute to the increase in

Firmicutes relative abundance seen in dominant individuals. This finding is inconsistent with

other trends observed, as other pathogenic bacterial communities decreased in dominant

individuals; however, these observations still provide insight into gut microbiome composition

and associated social experience. In addition to the presence of *Staphylococcus* in dominant

individuals, indicator species analysis also revealed the presence of *Achromobacter* and

Paracoccus in dominant individuals during the isolation period (Table S3). *Achromobacter* and

Paracoccus are genera of bacteria with pathogenic properties in both mammals and fish (71–

73). While these genera are present in dominant individuals prior to pairing, they decrease in

relative abundance post-pairing which supports the hypothesis that social status may affect gut

bacterial composition. Prior studies did not describe the role for the unclassified genera

indicators observed for dominant day 7, which included OTUs in *Planctomycetaceae*,

Caldineaceae, and *Comamondaceae* genera. The *Aeromonas* genus, which has a high relative

abundance in almost all zebrafish individuals regardless of social status, has been observed in

several pathogenic models of zebrafish. Interestingly, specific species within the *Aeromonas*

genus have been seen to induce severe infection of zebrafish models by influencing the release

of reactive oxygen species (ROS) and nitrogen free radicals triggering an immune response

accompanied by massive inflammatory reaction and increased mortality rates (74). While our

results do not specify the species of *Aeromonas* present, higher relative abundances of this

genus may result in pathogenesis in chronically stressed animals as well (e.g., socially

submissive or isolates).

Collectively, our results show that gut microbiome composition is socially regulated. This

is illustrated by a partial shift in the composition from pathogenic to commensal bacteria in

dominant animals; while chronically stressed animals (e.g., submissive, and socially isolated)

showed an increase in pathogenic bacteria and a decline in commensal bacterial genera. The

interconnectivity between gut function and cognitive processes are likely to mediated bi-

directionally via afferent and efferent neural and neuroendocrine signaling pathways (75).

Although our results suggest that the physiological mechanisms underlying the shift in microbial

composition is slow acting and require one to two weeks for their biological manifestation, prior

reports have shown that acute exposure to stress can impact the gut microbiota community

composition by changing the relative proportions of the main microbiota phyla (76). Moreover,

experimental manipulations of gut microbiota affect stress levels and anxiety-like behavior by

influencing the function of the hypothalamic-pituitary-adrenal stress axis (77). Indeed, social

stress induced increase in blood cortisol levels correlates with changes in the diversity of the

mammalian gut microbiome, including the abundance of lactic acid and pathogenic bacteria (76,

78). In addition, chronic injections of glucocorticoids can have positive and negative impact on

the presence of specific microbial taxa, which consequently affect host metabolism (79–81).

Thus, it is likely that chronic stress induced by either prolonged social isolation or subordination

likely to have increased cortisol levels in isolated and submissive zebrafish with substantial

increase in expression of gut pathogenic bacteria.

How these status-dependent changes in gut bacterial composition feedback to influence

brain function remains poorly understood. Given that zebrafish experiencing induced stress

exhibit changes in gut microbiome composition and brain gene expression (24), future

experiments testing this relationship in zebrafish in the context of social dominance and stress

will be highly informative. More specifically, it would be of interest to determine whether

transcriptional changes in the brains of dominant and subordinate fish can be linked to changes

in bacterial abundance in the host gut. Indeed, a large body of work has shown that

dopaminergic, GABAergic, and serotoninergic pathways implicated in social behavior are highly

plastic, but whether the plasticity is mediated by feedback regulation of gut microbiome to

influence motivational brain centers involved in social aggression remains an open question (32,

82–86).

ACKNOWLEDGMENTS

We thank Dr. Katie Clements for her assistance in the early design of behavioral experiments.

This work was supported by the National Science Foundation grant (#1754513) to F.A.I. and

grant (#1845845) to A.L.P..

REFERENCES

1.   Kinross JM, Darzi AW, Nicholson JK. 2011. Gut microbiome-host interactions in health and

disease. Genome medicine 3:14.

2.   Foster JA, Rinaman L, Cryan JF. 2017. Stress & the gut-brain axis: Regulation by the

microbiome. Neurobiology of stress 7:124–136.

3.   Lazar V, Ditu L-M, Pircalabioru GG, Gheorghe I, Curutiu C, Holban AM, Picu A, Petcu L,

Chifiriuc MC. 2018. Aspects of Gut Microbiota and Immune System Interactions in

Infectious Diseases, Immunopathology, and Cancer. Frontiers in immunology 9:1830.

4.   Geuking MB, Köller Y, Rupp S, McCoy KD. 2014. The interplay between the gut microbiota

and the immune system. Gut microbes 5:411–418.

5.   van der Meulen TA, Harmsen H, Bootsma H, Spijkervet F, Kroese F, Vissink A. 2016. The

microbiome-systemic diseases connection. Oral diseases 22:719–734.

6.   Stefan KL, Kim M V, Iwasaki A, Kasper DL. 2020. Commensal Microbiota Modulation of

Natural Resistance to Virus Infection. Cell 183:1312-1324.e10.

7.   Tamana SK, Tun HM, Konya T, Chari RS, Field CJ, Guttman DS, Becker AB, Moraes TJ,

Turvey SE, Subbarao P, Sears MR, Pei J, Scott JA, Mandhane PJ, Kozyrskyj AL. 2021.

Bacteroides-dominant gut microbiome of late infancy is associated with enhanced

neurodevelopment. Gut microbes 13:1–17.

8.   Warner BB. 2019. The contribution of the gut microbiome to neurodevelopment and

neuropsychiatric disorders. Pediatric research 85:216–224.

9.   Ntranos A, Casaccia P. 2018. The Microbiome-Gut-Behavior Axis: Crosstalk Between the

Gut Microbiome and Oligodendrocytes Modulates Behavioral Responses.

Neurotherapeutics : the journal of the American Society for Experimental

NeuroTherapeutics 15:31–35.

10.   Sharon G, Sampson TR, Geschwind DH, Mazmanian SK. 2016. The Central Nervous

System and the Gut Microbiome. Cell 167:915–932.

11.   Christian LM, Galley JD, Hade EM, Schoppe-Sullivan S, Kamp Dush C, Bailey MT. 2015.

Gut microbiome composition is associated with temperament during early childhood. Brain,

behavior, and immunity 45:118–127.

12.   Tan LY, Yeo XY, Bae H-G, Lee DPS, Ho RC, Kim JE, Jo D-G, Jung S. 2021. Association

of Gut Microbiome Dysbiosis with Neurodegeneration: Can Gut Microbe-Modifying Diet

Prevent or Alleviate the Symptoms of Neurodegenerative Diseases? Life (Basel,

Switzerland) 11.

13.   Ghaisas S, Maher J, Kanthasamy A. 2016. Gut microbiome in health and disease: Linking

the microbiome-gut-brain axis and environmental factors in the pathogenesis of systemic

and neurodegenerative diseases. Pharmacology & therapeutics 158:52–62.

14.   Chandra S, Alam MT, Dey J, Sasidharan BCP, Ray U, Srivastava AK, Gandhi S, Tripathi

PP. 2020. Healthy Gut, Healthy Brain: The Gut Microbiome in Neurodegenerative

Disorders. Current topics in medicinal chemistry 20:1142–1153.

15. Sgritta M, Dooling SW, Buffington SA, Momin EN, Francis MB, Britton RA, Costa-Mattioli

M. 2019. Mechanisms Underlying Microbial-Mediated Changes in Social Behavior in

Mouse Models of Autism Spectrum Disorder. Neuron 101:246-259.e6.

16. Lynch JB, Hsiao EY. 2019. Microbiomes as sources of emergent host phenotypes.

Science (New York, NY) 365:1405–1409.

17. Hughes HK, Rose D, Ashwood P. 2018. The Gut Microbiota and Dysbiosis in Autism

Spectrum Disorders. Current neurology and neuroscience reports 18:81.

18. Cresci GA, Bawden E. 2015. Gut Microbiome: What We Do and Don't Know. Nutrition in

clinical practice : official publication of the American Society for Parenteral and Enteral

Nutrition 30:734–746.

19. Jandhyala SM, Talukdar R, Subramanyam C, Vuyyuru H, Sasikala M, Nageshwar Reddy

D. 2015. Role of the normal gut microbiota. World journal of gastroenterology 21:8787–

8803.

20. Kamada N, Chen GY, Inohara N, Núñez G. 2013. Control of pathogens and pathobionts by

the gut microbiota. Nature immunology 14:685–690.

21. Dowlati Y, Herrmann N, Swardfager W, Liu H, Sham L, Reim EK, Lanctôt KL. 2010. A

meta-analysis of cytokines in major depression. Biological psychiatry 67:446–457.

22. Ochoa-Reparáz J, Mielcarz DW, Begum-Haque S, Kasper LH. 2011. Gut, bugs, and brain:

role of commensal bacteria in the control of central nervous system disease. Annals of

neurology 69:240–247.

23.   Davis DJ, Doerr HM, Grzelak AK, Busi SB, Jasarevic E, Ericsson AC, Bryda EC. 2016.

Lactobacillus plantarum attenuates anxiety-related behavior and protects against stress-

induced dysbiosis in adult zebrafish. Scientific reports 6:33726.

24.   Borrelli L, Aceto S, Agnisola C, De Paolo S, Dipineto L, Stilling RM, Dinan TG, Cryan JF,

Menna LF, Fioretti A. 2016. Probiotic modulation of the microbiota-gut-brain axis and

behaviour in zebrafish. Scientific reports 6:30046.

25.   Ventriglia M, Zanardini R, Bonomini C, Zanetti O, Volpe D, Pasqualetti P, Gennarelli M,

Bocchio-Chiavetto L. 2013. Serum brain-derived neurotrophic factor levels in different

neurological diseases. BioMed research international 2013:901082.

26.   Mondelli V, Cattaneo A, Murri MB, Di Forti M, Handley R, Hepgul N, Miorelli A, Navari S,

Papadopoulos AS, Aitchison KJ, Morgan C, Murray RM, Dazzan P, Pariante CM. 2011.

Stress and inflammation reduce brain-derived neurotrophic factor expression in first-

episode psychosis: a pathway to smaller hippocampal volume. The Journal of clinical

psychiatry 72:1677–1684.

27.   Chiu IM, Heesters BA, Ghasemlou N, Von Hehn CA, Zhao F, Tran J, Wainger B,

Strominger A, Muralidharan S, Horswill AR, Bubeck Wardenburg J, Hwang SW, Carroll

MC, Woolf CJ. 2013. Bacteria activate sensory neurons that modulate pain and

inflammation. Nature 501:52–57.

28.   Oliveira RF, Silva JF, Simoes JM. 2011. Fighting zebrafish: characterization of aggressive

behavior and winner-loser effects. Zebrafish 8:73–81.

29.   Clements KN, Miller TH, Keever JM, Hall AM, Issa FA. 2018. Social Status-Related

Differences in Motor Activity Between Wild-Type and Mutant Zebrafish. The Biological

Bulletin 235:71–82.

30.   Orr SA, Ahn S, Park C, Miller TH, Kassai M, Issa FA. 2021. Social Experience Regulates

540   Endocannabinoids Modulation of Zebrafish Motor Behaviors. Frontiers in behavioral

541   neuroscience 15:668589.

542   31.   Carver JJ, Carrell SC, Chilton MW, Brown JN, Yong L, Zhu Y, Issa FA. 2021. Nuclear

543   androgen and progestin receptors inversely affect aggression and social dominance in

544   male zebrafish (Danio rerio). Hormones and behavior 134:105012.

545   32.   Teles MC, Dahlbom SJ, Winberg S, Oliveira RF. 2013. Social modulation of brain

546   monoamine levels in zebrafish. Behavioural brain research 253:17–24.

547   33.   Teles MC, Oliveira RF. 2016. Quantifying Aggressive Behavior in Zebrafish. Methods in

548   molecular biology (Clifton, NJ) 1451:293–305.

549   34.   Miller TH, Clements K, Ahn S, Park C, Hye Ji E, Issa FA. 2017. Social Status–Dependent

550   Shift in Neural Circuit Activation Affects Decision Making. The Journal of Neuroscience

551   37:2137–2148.

552   35.   Sylvia KE, Demas GE. 2018. A gut feeling: Microbiome-brain-immune interactions

553   modulate social and affective behaviors. Hormones and behavior 99:41–49.

554   36.   Chrousos GP, Gold PW. 1992. The concepts of stress and stress system disorders.

555   Overview of physical and behavioral homeostasis. JAMA 267:1244–1252.

556   37.   Blanchard RJ, McKittrick CR, Blanchard DC. 2001. Animal models of social stress: effects

557   on behavior and brain neurochemical systems. Physiology & behavior 73:261–271.

558   38.   Mudra Rakshasa A, Tong MT. 2020. Making "Good" Choices: Social Isolation in Mice

559   Exacerbates the Effects of Chronic Stress on Decision Making. Frontiers in behavioral

560   neuroscience 14:81.

39. Bartolomucci A, Palanza P, Sacerdote P, Panerai AE, Sgoifo A, Dantzer R, Parmigiani S.

2005. Social factors and individual vulnerability to chronic stress exposure. Neuroscience

and biobehavioral reviews 29:67–81.

40. Egan RJ, Bergner CL, Hart PC, Cachat JM, Canavello PR, Elegante MF, Elkhayat SI,

Bartels BK, Tien AK, Tien DH, Mohnot S, Beeson E, Glasgow E, Amri H, Zukowska Z,

Kalueff A V. 2009. Understanding behavioral and physiological phenotypes of stress and

anxiety in zebrafish. Behavioural brain research 205:38–44.

41. Raison CL, Miller AH. 2003. When not enough is too much: the role of insufficient

glucocorticoid signaling in the pathophysiology of stress-related disorders. The American

journal of psychiatry 160:1554–1565.

42. Luo Y, Zeng B, Zeng L, Du X, Li B, Huo R, Liu L, Wang H, Dong M, Pan J, Zheng P, Zhou

C, Wei H, Xie P. 2018. Gut microbiota regulates mouse behaviors through glucocorticoid

receptor pathway genes in the hippocampus. Translational psychiatry 8:187.

43. Stagaman K, Sharpton TJ, Guillemin K. 2020. Zebrafish microbiome studies make waves.

Lab animal 49:201–207.

44. Stephens WZ, Burns AR, Stagaman K, Wong S, Rawls JF, Guillemin K, Bohannan BJM.

2016. The composition of the zebrafish intestinal microbial community varies across

development. The ISME journal 10:644–654.

45. Chen W, Zhang CK, Cheng Y, Zhang S, Zhao H. 2013. A comparison of methods for

clustering 16S rRNA sequences into OTUs. PloS one 8:e70837.

46. Roeselers G, Mittge EK, Stephens WZ, Parichy DM, Cavanaugh CM, Guillemin K, Rawls

JF. 2011. Evidence for a core gut microbiota in the zebrafish. The ISME journal 5:1595–

1608.

47. Shin N-R, Whon TW, Bae J-W. 2015. Proteobacteria: microbial signature of dysbiosis in

gut microbiota. Trends in biotechnology 33:496–503.

48. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, Owens SM,

Betley J, Fraser L, Bauer M, Gormley N, Gilbert JA, Smith G, Knight R. 2012. Ultra-high-

throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. The

ISME journal 6:1621–1624.

49. Apprill A, McNally S, Parsons R, Weber L. 2015. Minor revision to V4 region SSU rRNA

806R gene primer greatly increases detection of SAR11 bacterioplankton . Aquatic

Microbial Ecology 75:129–137.

50. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013. Development of a

dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence

data on the MiSeq Illumina sequencing platform. Appl Environ Microbiol 79:5112–20.

51. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA,

Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Horn DJV, Weber

CF. 2009. Introducing Mothur: open-source, platform-independent, community-supported

software for describing and comparing microbial communities. Appl Environ Microbiol

75:7537–7541.

52. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.

2013. The SILVA ribosomal RNA gene database project: improved data processing and

web-based tools. Nucleic Acids Research 41:D590–D596.

53.   Rognes T, Flouri T, Nichols B, Quince C, Mahé F. 2016. VSEARCH: a versatile open

605   source tool for metagenomics. PeerJ 4:e2584.

606   54.   Schloss PD. 2021. Amplicon Sequence Variants Artificially Split Bacterial Genomes into

607   Separate Clusters. mSphere 6:e00191-21.

608   55.   Glassman SI, Martiny JBH. 2018. Broadscale Ecological Patterns Are Robust to Use of

609   Exact Sequence Variants versus Operational Taxonomic Units. mSphere 3:e00148-18.

610   56.   Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR,

611   O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H. 2019. vegan:

612   Community Ecology Package (2.5-5).

613   57.   Kuznetsova A, Brockhoff PB, Christensen RHB. 2017. lmerTest Package: Tests in Linear

614   Mixed Effects Models. Journal of Statistical Software 82:1–26.

615   58.   De Caceres, M.; Jansen F. 2016. Indispecies: Relationship between species and groups of

616   sites. R package version 1.7.6 R Found. Stat. Comput., Vienna. 1.7.6 R.

617   59.   Skonieczna-Żydecka K, Marlicz W, Misera A, Koulaouzidis A, Łoniewski I. 2018.

618   Microbiome-The Missing Link in the Gut-Brain Axis: Focus on Its Role in Gastrointestinal

619   and Mental Health. Journal of clinical medicine 7.

620   60.   Reardon S. 2014. Gut-brain link grabs neuroscientists. Nature. England

621   https://doi.org/10.1038/515175a.

622   61.   Flight MH. 2014. Neurodevelopmental disorders: the gut-microbiome-brain connection.

623   Nature reviews Neuroscience 15:65.

62. Sharon G, Cruz NJ, Kang D-W, Gandal MJ, Wang B, Kim Y-M, Zink EM, Casey CP, Taylor

BC, Lane CJ, Bramer LM, Isern NG, Hoyt DW, Noecker C, Sweredoski MJ, Moradian A,

Borenstein E, Jansson JK, Knight R, Metz TO, Lois C, Geschwind DH, Krajmalnik-Brown

R, Mazmanian SK. 2019. Human Gut Microbiota from Autism Spectrum Disorder Promote

Behavioral Symptoms in Mice. Cell 177:1600-1618.e17.

63. Walter J. 2008. Ecological role of lactobacilli in the gastrointestinal tract: implications for

fundamental and biomedical research. Applied and environmental microbiology 74:4985–

4996.

64. Dal Bello F, Hertel C. 2006. Oral cavity as natural reservoir for intestinal lactobacilli.

Systematic and applied microbiology 29:69–76.

65. Reuter G. 2001. The Lactobacillus and Bifidobacterium microflora of the human intestine:

composition and succession. Current issues in intestinal microbiology 2:43–53.

66. Stressmann FA, Bernal-Bayard J, Perez-Pascual D, Audrain B, Rendueles O, Briolat V,

Bruchmann S, Volant S, Ghozlane A, Häussler S, Duchaud E, Levraud J-P, Ghigo J-M.

2021. Mining zebrafish microbiota reveals key community-level resistance against fish

pathogen infection. The ISME journal 15:702–719.

67. Graz CJ, Grant GD, Brauns SC, Hunt A, Jamie H, Milne PJ. 2000. Cyclic dipeptides in the

induction of maturation for cancer therapy. The Journal of pharmacy and pharmacology

52:75–82.

68. Jinendiran S, Teng W, Dahms H-U, Liu W, Ponnusamy VK, Chiu CC-C, Kumar BSD,

Sivakumar N. 2020. Induction of mitochondria-mediated apoptosis and suppression of

tumor growth in zebrafish xenograft model by cyclic dipeptides identified from

Exiguobacterium acetylicum. Scientific reports 10:13721.

69.   Ström K, Sjögren J, Broberg A, Schnürer J. 2002. Lactobacillus plantarum MiLAB 393

648   produces the antifungal cyclic dipeptides cyclo(L-Phe-L-Pro) and cyclo(L-Phe-trans-4-OH-

649   L-Pro) and 3-phenyllactic acid. Applied and environmental microbiology 68:4322–4327.

650   70.   Clatworthy AE, Lee JS-W, Leibman M, Kostun Z, Davidson AJ, Hung DT. 2009.

651   Pseudomonas aeruginosa infection of zebrafish involves both host and pathogen

652   determinants. Infection and immunity 77:1293–1303.

653   71.   Menetrey Q, Sorlin P, Jumas-Bilak E, Chiron R, Dupont C, Marchandin H. 2021.

654   Achromobacter xylosoxidans and Stenotrophomonas maltophilia: Emerging Pathogens

655   Well-Armed for Life in the Cystic Fibrosis Patients' Lung. 5. Genes 12:610.

656   72.   Loftie-Eaton W, Crabtree A, Perry D, Millstein J, Baytosh J, Stalder T, Robison BD, Forney

657   LJ, Top EM. 2021. Contagious Antibiotic Resistance: Plasmid Transfer among Bacterial

658   Residents of the Zebrafish Gut. Applied and Environmental Microbiology 87:e02735-20.

659   73.   Decewicz P, Dziewit L, Golec P, Kozlowska P, Bartosik D, Radlinska M. 2019.

660   Characterization of the virome of Paracoccus spp. (Alphaproteobacteria) by combined in

661   silico and in vivo approaches. 1. Sci Rep 9:7899.

662   74.   Rodríguez I, Novoa B, Figueras A. 2008. Immune response of zebrafish (Danio rerio)

663   against a newly isolated bacterial pathogen Aeromonas hydrophila. Fish & shellfish

664   immunology 25:239–249.

665   75.   Mayer EA. 2011. Gut feelings: the emerging biology of gut-brain communication. Nature

666   reviews Neuroscience 12:453–466.

76.   Galley JD, Nelson MC, Yu Z, Dowd SE, Walter J, Kumar PS, Lyte M, Bailey MT. 2014.

Exposure to a social stressor disrupts the community structure of the colonic mucosa-

associated microbiota. BMC microbiology 14:189.

77.   Crumeyrolle-Arias M, Jaglin M, Bruneau A, Vancassel S, Cardona A, Daugé V, Naudon L,

Rabot S. 2014. Absence of the gut microbiota enhances anxiety-like behavior and

neuroendocrine response to acute stress in rats. Psychoneuroendocrinology 42:207–217.

78.   Mudd AT, Berding K, Wang M, Donovan SM, Dilger RN. 2017. Serum cortisol mediates the

relationship between fecal Ruminococcus and brain N-acetylaspartate in the young pig.

Gut microbes 8:589–600.

79.   Huang EY, Inoue T, Leone VA, Dalal S, Touw K, Wang Y, Musch MW, Theriault B, Higuchi

K, Donovan S, Gilbert J, Chang EB. 2015. Using corticosteroids to reshape the gut

microbiome: implications for inflammatory bowel diseases. Inflammatory bowel diseases

21:963–972.

80.   Petrosus E, Silva EB, Lay DJ, Eicher SD. 2018. Effects of orally administered cortisol and

norepinephrine on weanling piglet gut microbial populations and Salmonella passage.

Journal of animal science 96:4543–4551.

81.   Wu T, Yang L, Jiang J, Ni Y, Zhu J, Zheng X, Wang Q, Lu X, Fu Z. 2018. Chronic

glucocorticoid treatment induced circadian clock disorder leads to lipid metabolism and gut

microbiota alterations in rats. Life sciences 192:173–182.

82.   Weitekamp CA, Nguyen J, Hofmann HA. 2017. Neuromolecular Regulation of Aggression

Differs by Social Role during Joint Territory Defense. Integrative and comparative biology

57:631–639.

83.   Inoue K, Ford CL, Horie K, Young LJ. 2022. Oxytocin receptors are widely distributed in

690   the prairie vole (Microtus ochrogaster) brain: Relation to social behavior, genetic

691   polymorphisms, and the dopamine system. The Journal of comparative neurology

692   530:2881–2900.

693   84.   Korzan WJ, Summers CH. 2007. Behavioral diversity and neurochemical plasticity:

694   selection of stress coping strategies that define social status. Brain, behavior and

695   evolution 70:257–266.

696   85.   McDonald MM, Markham CM, Norvelle A, Albers HE, Huhman KL. 2012. GABAA receptor

697   activation in the lateral septum reduces the expression of conditioned defeat and

698   increases aggression in Syrian hamsters. Brain research 1439:27–33.

699   86.   Weitekamp CA, Nguyen J, Hofmann HA. 2017. Social context affects behavior, preoptic

700   area gene expression, and response to D2 receptor manipulation during territorial defense

701   in a cichlid fish. Genes, brain, and behavior 16:601–611.

Social Status Zebrafish Gut Microbiomes