Neo-natal castration leads to subtle differences in porcine anterior cruciate ligament

morphology and function in adolescence

Jacob D. Thompson[1,2], Danielle Howe[1,2], Emily H. Griffith[3], Matthew B. Fisher[1,2,4,*]

[1]Joint Department of Biomedical Engineering, North Carolina State University and the

University of North Carolina – Chapel Hill; Raleigh, NC 27695

[2]Comparative Medicine Institute, North Carolina State University; Raleigh, NC 27695

[3]Department of Statistics, North Carolina State University; Raleigh, NC 27695

[4]Department of Orthopaedics, University of North Carolina – Chapel Hill; Chapel Hill, NC

27599

*Corresponding Author Contact:

Address: 4130 Engineering Building III, 911 Oval Drive, CB 7115, Raleigh, NC, 27695

Telephone: 919-515-5242

Fax: 919-513-3814

Email: mbfisher@ncsu.edu

Author Contribution

Jacob D. Thompson contributed to the development of research design, acquisition, analysis, and interpretation of data, and drafting of the manuscript. Danielle Howe contributed to the development of research design, acquisition of data, and revision of the manuscript. Emily H. Griffith contributed to the interpretation of data and revision of the manuscript. Matthew B. Fisher contributed to the development of research design, acquisition, analysis, and interpretation of data, and revision of the manuscript. All authors have read and approved the final submitted manuscript.

Abstract

Female adolescent athletes are at a higher risk of tearing their anterior cruciate ligament (ACL) than male counterparts. While most work related to hormones has focused on the effects of estrogen to understand the increased risk of ACL injury, there are other understudied factors, including testosterone. The purpose of this study was to determine how surgical castration in the male porcine model influences ACL size and function across skeletal growth. Thirty-six male Yorkshire crossbreed pigs were raised to 3 (juvenile), 4.5 (early adolescent), and 6 months (adolescent) of age. Animals were either castrated (barrows) within 1-2 weeks after birth or were left intact (boars). Post-euthanasia, joint and ACL size were assessed via MRI, and biomechanics were assessed via a robotic testing system. Joint size increased throughout age, yet barrows had smaller joints than boars ($p<0.001$ for all measures). ACL cross-sectional area (CSA), length, volume, and stiffness increased with age ($p<0.0001$), as did ACL anteromedial (AM) bundle percent contribution to resisting loads ($p=0.012$). Boar ACL, AM bundle, and PL bundle volumes were 19% ($p=0.002$), 25% ($p=0.003$), and 15% ($p=0.04$) larger than barrows across ages. However, CSA, stiffness, and bundle contribution were similar between boars and barrows ($p>0.05$). The barrows had smaller temporal increases in AM bundle percent function than boars, but these data were highly variable. Thus, early and sustained loss in testosterone leads to subtle differences in ACL morphology, but may not influence measures associated with increased injury risk, such as CSA or bundle forces in response to applied loads.

Key Words: anterior cruciate ligament; sex hormones; castration; porcine model

Introduction

There are an ever-increasing number of pediatric athletes participating in middle and high school sports[1], accompanied by an increase in pediatric anterior cruciate ligament (ACL) injuries[2–4]. Furthermore, during adolescence there is a difference in the rate of ACL injuries between males and females, with females being 1.4 times more likely to tear their ACL overall[5]. In specific sports, they are over 3 times more likely to tear their ACL, like basketball[6–8] or soccer[9]. Several groups attribute this difference to aspects of sexual maturity[10,11] or gendered expectations in training, sport selection, and coaching[12,13]. A large focus has been on the onset of spikes and cycles of sex hormones[14–17]. The female sex hormones estradiol and progesterone have gathered the most attention with some groups showing associations between peaks of estradiol and progesterone to knee joint laxity and time of ACL injury[13,14,16,18]. However, strong evidence for a direct correlation between estradiol levels and injury risk does not yet exist[16,17]. While a major focus has been on uncovering theorized catabolic effects of female sex hormones to understand the increased risk of ACL injury, an understudied factor is the influence of testosterone, the primary male sex hormone, on developing male ACLs.

It is known that there are androgen receptors present within the human male ACL[19], suggesting that the ACL is an androgen-responsive tissue. In humans, males who tore their ACLs have higher levels of testosterone, indicating that testosterone may play some direct or indirect role in increasing ACL injury risk[20]. To study testosterone in animal models, surgical castration has been performed to drastically decrease total serum testosterone by over 95%[21,22]. For example, adult rats castrated at 12 weeks of age (post-puberty) had a 25-fold lower testosterone at 5 weeks post-castration than age-matched controls. Although the ACLs in the castrated rat group had a similar size, they had 15% lower load-to-failure values and 18% lower ultimate

stresses than intact males[22]. This study provides some evidence that testosterone may play a

protective role in ligament function in adults, especially failure properties of the ACL[22]. Thus, it

may be possible that both high and low levels of testosterone may impact the ACL, yet the

evidence is still sparse, particularly during skeletal growth.

Large animal models provide a useful tool to study the ACL. Our group, along with

others, have used a porcine model to understand structural and functional changes of the ACL

throughout skeletal growth, including progression through adolescence[23–28]. We have shown

similar size and functional changes within the human and porcine ACL, making this a good

model to study changes during adolescence[23,29]. The size of the porcine ACL diverges in males

and females around the point of sexual maturity, mainly within the two distinct bundles of the

ACL, the anteromedial (AM) and posterolateral (PL) bundles[30]. The AM bundle in males and

females increase in size throughout adolescence, yet the PL bundle size plateaus near the onset of

puberty. Functionally, there is also a shift in bundle function when resisting anterior loads, where

the AM bundle becomes the primary stabilizer over the PL bundle in both males and females[30].

The time of these changes is sex-dependent, with the plateau in PL bundle size occurring earlier

and functional changes occurring later in females. These sex specific changes, and their timing

around adolescence, motivate the examination of the role of sex hormones as a potential cause of

these changes. It is still unclear, though, which sex hormones drive changes in males and

females. Therefore, the objective of this study was to determine how neo-natal castration

influences the structure and function of the male porcine ACL during youth and early

adolescence.

Methods

Specimen Collection and Structural Imaging

Thirty-six male Yorkshire crossbreed pigs were raised to 3 months (juvenile), 4.5 months
(pre-puberty), or 6 months (early puberty). Six of each age group were left sexually intact
(boars), while the other six were castrated (barrows) at an early age (n=6 animals/age/treatment).
The boar data presented in this study has been re-analyzed from a previous study[30] to be
compared directly with barrow data. All animals were bred at the North Carolina State
University Swine Education Unit, and all experimental protocols were approved by the NC State
University IACUC. Following euthanasia, hind limbs were removed, and the stifle (knee) joints
were stored at -20°C in saline-soaked gauze.

To determine structural data, stifle joints were thawed and imaged using a 7.0-T Siemens
Magnetom MRI scanner (Siemens Healthineers, Erlangen, Germany) using a double-echo
steady-state sequence (DESS, flip angle: 25°, TR: 17ms, TE: 6s, voxel size: 0.42 x 0.42 x
0.4mm). Joint size data, including bicondylar width, tibial CSA, anterior-posterior (AP) tibial
width, as well as ACL, AM and PL bundle CSAs, orientations, lengths, and volumes were
recorded after segmentation using a commercial software (Simpleware ScanIP, Synopsys,
Mountain View, CA, USA) and processing using MATLAB (MathWorks, Natick, MA, USA)
scripts. Morphological ACL characteristics, including CSA, length, and volume, were
normalized to bicondylar width to account for overall joint size.

Biomechanical Data Collection and Processing

Stifle joints were thawed, trimmed, and the femur and tibia mounted in custom molds
using a fiberglass reinforced epoxy (Everglass, Evercoat, Cincinnati, OH, USA) to prepare for
robotic biomechanical testing. Stifle joints were then mounted to a robotic testing system
(KR300 R2500, KRC4, Kuka, Shelby Charter Township, MI, USA) with a universal force-

moment sensor (Omega160 IP65, ATI Industrial Automatic, Apex, NC, USA), following

previously described methods[30]. All systems were controlled through the simVITRO software

knee module (Cleveland Clinic, Cleveland, OH, USA).

During each test, the femur was fixed to the testing platform and the tibia was mounted to

the robotic system. Anatomical points on each leg were recorded using a 3D digitizer (G2X

MicroScribe, Amherst, VA, USA) to establish an anatomic coordinate system. The joints were

then flexed from 40° (approximately full extension in pigs) to 90° while minimizing joint forces

to establish a passive path during the test. AP forces were then applied to the joints at 60° of

flexion, while minimizing forces in other directions and recording kinematic data. Maximum

loads were scaled across age (40 N, 80 N, and 100 N for 3-month, 4.5-month, and 6-month,

respectively) based on the femoral cortical CSA. The recorded kinematics were then repeated

and the forces were measured. Joint tissues were then removed sequentially (capsule, AM

bundle, then PL bundle), and the intact kinematics were repeated to determine the force

contribution of each tissue through the principle of superposition[31,32]. After the AM bundle was

transected, the same loads were applied to the joint while new kinematics were recorded to fully

engage the remaining PL bundle.

Force-translation curves were then plotted across AP loading. The engagement point of

the ACL was determined by the point of inflection in the force-translation curves, which

separated the toe region and the linear region in anterior loading[33,34]. The toe region was fit using

an exponential curve, and the higher loads above the inflection point were fit to a line. *In situ*

joint slack and stiffnesses were then calculated using Matlab scripts, as previously described[25,33]

(Figure S1). The stiffness values for the ACL and AM bundle were calculated under the intact

kinematics, where the PL bundle stiffness was calculated under the AM deficient state to ensure

PL engagement. AM and PL contribution percentages to resisting anterior loading were

calculated by dividing the anterior force for each bundle by the total ACL contribution to

anterior loading and multiplying by 100. The difference in AM bundle contribution and PL

bundle contribution were plotted to directly compare the boars and barrows.

Statistical Analysis

Overall effects of age and surgical castration on bicondylar width, CSA, length, volume,

stiffness, slack, and difference in bundle percent contribution were assessed using two-way

ANOVAs for each variable. No animals were excluded from this work. Multiple comparisons

made between ages and boars and barrows at each age were run and the Type I error rate was

controlled using Holm-Sidak's post hoc method. The difference in AM and PL bundle percent

contribution was analyzed using one sample t-tests comparing the means to zero. Overall

significance was set to $\alpha=0.05$ for all tests.

Results

Joint Size Decreases following Surgical Castration

From MRI scans, joint bicondylar width, tibial plateau CSA, and AP tibial width were

recorded as metrics of joint size between the boars and barrows (Fig 1, Table S1). Across all

three measurements of joint size, age had significant effects on overall knee joint size ($p<0.0001$

for all measures). Specifically, bicondylar width increased by 20% and 21% from 3-months to 6-

months for the boars and the barrows, respectively (Fig 1A,D). Tibial plateau CSA increased by

64% and 63% from 3-months to 6-months for boars and barrows, respectively (Fig 1B,E).

Similar changes were seen for AP tibial width, increasing by 25% and 32% for boars and

barrows between the 3-month and 6-month age groups (Fig 1C,F). Surgical castration had

significant effects on joint size (bicondylar width: p=0.001, tibial plateau CSA: p=0.0007, AP

tibial width: p<0.0001). Barrow bicondylar width was 8% and 7% smaller at 3-months (adj

p=0.03) and 6-months (adj p=0.02) but was not statistically significantly smaller at 4.5 months

(adj p=0.40). There were statistically significant (adj p=0.003) differences at 6 months in tibial

plateau CSA between boars and barrows with the barrow plateau CSA being 13% smaller on

average. All barrow AP tibial widths at each age group were statistically significantly smaller

than the boars (3-month: 10% smaller, adj p=0.002; 4.5-month: 5% smaller, adj p=0.05; 6-

month: 6% smaller, adj p=0.02).

Figure 1. Barrows have smaller joints than boars throughout growth. Joint size measurements

taken from the (A) coronal, (B) transverse, and (C) sagittal planes from the MRI scans.

Quantitative comparison of (D) bicondylar width, (E) tibial plateau CSA, and (F) AP tibial width

measurements between boars and barrows across three ages tested, where each point represents

independent specimens. Data presented as mean ± 95% confidence interval with main effects

from two-way ANOVA shown in graph corner. Bars indicate p<0.05 between boars and barrows

within ages.

ACL Structure and Morphology Between Castrated and Intact Animals

In addition to overall joint size, the morphology of the ACL and its corresponding AM

and PL bundles were recorded. CSAs, lengths, and volumes for the ACL and the bundles were

segmented from the MRI scans. For both boars and barrows, there were significant age effects

seen for ACL, AM, and PL CSAs ($p<0.0001$ for all) (Fig 2A, Table S2). When comparing boars

and barrows, there was a minor, but not statistically significant, effect of castration on ACL and

AM CSA ($p=0.12$, $p=0.20$), with boars having slightly higher CSA values at 4.5 and 6-months

(12% larger for both ages). At 3 months, boars and barrows had similarly-sized ACLs, but

barrows had slightly larger AM bundles (3-months: 3% larger). There was no significant effect

of castration on raw PL bundle CSA size ($p=0.66$). ACL and bundle orientation were also

recorded in the sagittal plane, and the orientation of the ACL and the bundles increased across

age but were not statistically significant between boars and barrows (Fig S2). All CSA values

were normalized to the bicondylar width of the joint to account for overall joint size (Fig 2B).

After normalizing, the effect of age persisted for the ACL, AM, and PL CSAs ($p<0.0001$ for all),

but the lack of substantial effect due to castration stayed the same ($p=0.81$, $p=0.70$, $p=0.27$).

Figure 2. Boar and barrow ACL and bundle CSA increased throughout skeletal growth and adolescence in similar ways. (A) CSAs of the ACL, AM bundle, and PL bundle increased steadily across adolescence, and increases were similar between boars and barrows. (B) Normalizing by bicondylar width to account for overall joint size diminished any minor differences between boars and barrows. Data presented as mean ± 95% confidence interval with main effects from two-way ANOVA shown in graph corner.

Age had similar effects on ACL and bundle lengths, steadily increasing from 3 months to 6 months (p<0.0001 for all) (Fig 3A, Table S3). There was also an effect of castration on the ACL length (p=0.0018) and PL bundle length (p=0.0002), but not the AM bundle length. The boars had on average 8% longer ACLs across all age groups tested. The boars also had longer PL bundles at each age (3-month: 17% greater, adj p=0.005; 4.5-month: 10% greater, adj p=0.07; 6-month: 6% greater, adj p=0.12). After normalizing to joint size, the statistically significant

difference in ACL lengths across ages and between boars and barrows disappeared (Fig 3B). The

age effect also disappeared for the AM bundle and PL bundle lengths. However, when

comparing normalized boar and barrow ACL and bundle lengths, there was still a statistically

significant difference in the PL bundle length (p=0.012), with barrows having a statistically

significant shorter PL bundle at 4.5 months (11% shorter, adj p=0.02).

Figure 3. Boar and barrow ACL and bundle lengths all increased throughout skeletal growth and

adolescence but had differing lengths. (A) Lengths of the ACL, AM bundle, and PL bundle

increased steadily across adolescence, but barrows had shorter ACLs and PL bundles than boars.

(B) Normalizing by bicondylar width to account for overall joint size diminished the differences

in ACL length, but not PL bundle length, between boars and barrows. Data presented as mean ±

95% confidence interval with main effects from two-way ANOVA shown in graph corner. Bars

indicate p<0.05 between boars and barrows within ages.

While ACL and bundle CSA and length showed some minor differences between boars

and barrows across the ages tested, the ACL and bundle volumes showed more substantial

changes between boars and barrows. A similar age effect was observed in ACL, AM bundle, and

PL bundle volumes (p<0.0001 for all), which all increased during growth (Fig 4A, Table S4).

Castration also had a statistically significant impact on ACL (p=0.002), AM bundle (p=0.003),

and PL bundle (p=0.04) volume. Comparing boars and barrows across ages, the boars had on

average 19% larger ACLs, 25% larger AM bundles, and 15% larger PL bundles than the

barrows. The age effect is still seen after normalizing to the bicondylar width of the joints, but

the effects of castration are no longer statistically significant for the ACL (p=0.06) and PL

bundle (p=0.11), but persists in the AM bundle (p=0.04) (Fig 4B).

Figure 4. Boar and barrow ACL and bundle volumes all increased throughout skeletal growth

and adolescence with barrows having smaller ACL and AM bundle. (A) Volumes of the ACL,

AM bundle, and PL bundle increased steadily across adolescence, but barrows had smaller ACLs

and AM bundles, but not PL bundles than boars. (B) Normalizing by bicondylar width to account

for overall joint size diminished the differences in ACL and AM bundle volume between boars

and barrows. Data presented as mean ± 95% confidence interval with main effects from two-way

ANOVA shown in graph corner. Bars indicate $p<0.05$ between boars and barrows within ages.

Minor Differences in ACL Mechanics in Castrated Animals

Using a 6-degree of freedom robotic testing system, force translation curves showed

similar trends in both boars and barrows, where the slopes increased with age, indicating an

increase in overall joint stiffness ($p<0.0001$) (Fig 5A-B). There were no statistically significant

differences in overall joint stiffness, however, between boars and barrows ($p=0.10$) (Fig 5C).

Joint slack was calculated as the distance between engagement points of the ACL and the PCL.

Both age ($p=0.03$) and castration ($p=0.009$) had overall significant effects on joint slack (Fig

5D). The barrows, in general, had shorter slack lengths than the boars, and slack increased

throughout growth. Age did not seem to have an effect on the boar group, however, with slacks

remaining somewhat similar across ages. Normalized slack values showed no statistically

significant effects of age ($p=0.40$) or castration ($p=0.17$) (Table S5). Similar changes were seen

in the joint laxity, reported as APTT (Fig 5E, Table S6). Both age ($p=0.0008$) and castration

($p=0.008$) had pronounced impacts on APTT, with barrows having lower APTT values than the

boars, specifically 19% shorter APTT at 3 months ($p=0.03$). Both groups had increased APTT

values across the age groups tested. However, once normalizing to joint size, these age and

castration effects disappeared (Fig 5F).

A

Boars

B

Barrows

C

In Situ Joint Stiffness

D

In Situ Joint Slack

E

APTT

F

Normalized APTT

Figure 5. *In situ* joint stiffness, slack, and APTT all change throughout skeletal growth with

barrows having less slack and raw laxity than boars. Force translation curves are similar between

(A) boars and (B) barrows, with increasing slopes, indicating (C) more stiff joints throughout

skeletal growth. (D) *In situ* joint slack slowly increases across age in boars and barrows, but

more prominently in the barrow group. (E) Castration and age had significant impacts on joint

APTT yet had (F) no significant impact after normalizing to joint size. Force translation data

presented as mean forces and mean translations. All other data presented as mean ± 95%

confidence interval with main effects from two-way ANOVA shown in graph corner. Bars

indicate $p < 0.05$ between boars and barrows within ages.

Similar effects were seen in terms of ACL and bundle function. Stiffness increased with

age (ACL: $p < 0.0001$, AM bundle: $p < 0.0001$, PL bundle: $p = 0.003$), but did not change due to

castration (ACL: $p = 0.11$, AM bundle: $p = 0.76$, PL bundle: $p = 0.15$) (Fig 6A-C, Table S7). ACL

stiffness values for the boars increased by 111% from 3 to 6 months, and the barrow stiffness

values increased by 91%. The ACL was the primary contributor to resisting anterior drawer in

both the boars and barrows, but the two bundles of the ACL had distinct contributions to

resisting load that varied across age (Fig 6D, Table S8). At 3-months, the AM bundle contributed

slightly more to resisting anterior drawer than the PL bundle for both boars and barrows. From 3-

months to 4.5-months, the AM bundle in the boars became the primary stabilizer in the ACL,

taking on 85% of the total force applied to the ACL during anterior drawer. This continued to

increase up to 91% in the 6-month boars. In the barrows, the AM bundle contributed 70% and

77% at 4.5 and 6-months, respectively. To compare boar and barrow bundle contributions across

age, the difference in AM and PL percent contribution was plotted over time, with negative

values indicating greater PL bundle forces and positive values indicating greater AM bundle

forces (Fig 6E). There was a statistically significant age effect seen (p=0.012), with AM bundles

becoming more dominant across growth for both boars and barrows. Furthermore, all 4.5-month

and 6-month differences in percent bundle contribution were statistically greater than 0,

indicating higher AM bundle contribution. There was no statistical significance between boars

and barrows (p=0.24), yet there was a more gradual shift in barrows towards higher AM bundle

contribution, whereas in boars, the contribution of the AM bundle increased sharply between 3

and 4.5 months.

Figure 6. *In situ* ACL, AM bundle, and PL bundle stiffnesses increase throughout skeletal

growth and are similar between boars and barrows. (A) ACL, (B) AM bundle, and (C) PL bundle

stiffnesses all increase from 3-month to 6-months. ACL and AM bundle stiffnesses calculated

during intact kinematics, and PL bundle stiffnesses calculated during the AM bundle deficient

kinematic state. (D) Tissue contributions varying across age, with the AM bundle become more

dominant in boars and slowly increasing in barrows. (E) Difference in bundle contribution

similar in boars and barrows, with both increasing toward AM bundle dominance. Data

presented as mean ± 95% confidence interval with main effects from two-way ANOVA shown

in graph corner. + indicates $p<0.05$ relative to zero (indicating AM≠PL) within age groups.

Discussion

This work explored the influence of neo-natal surgical castration on ACL morphology

and function later in youth and early adolescence in a male porcine model. Surgical castration

lowers serum testosterone levels by over 95%[21] and is a standard practice in the swine industry to

control for meat quality[35]. The ACL increased considerably in size and stiffness across skeletal

growth for both intact males (boars) and castrated males (barrows). Relative to changes in

growth, however, there were only subtle differences between the two boars and barrows.

Barrows had, on average, smaller overall knee joints along with smaller absolute ACL, AM

bundle, and PL bundle volumes, but no difference in ACL or bundle CSA. When accounting for

joint size, any differences disappeared. Functionally, the AM bundle percent contribution

increased across growth for both groups, yet the boars had a more dramatic shift from equal

bundle percent contribution to a large increase in AM bundle percent contribution.

While joint size has not been previously characterized between boars and barrows, it has

been shown that barrows have lighter tibias than boars despite similar density and length

measurements[36], indicating greater bone thickness in boars. Differences in body composition

between boars and barrows have been well-characterized, with barrows having more fat and less

bone and skin weight than boars[36–38]. This finding is similar in humans, where

supraphysiological doses of testosterone lead to more fat-free mass and muscle size in men[39].

When looking at changes across skeletal growth, the increase in ACL length, CSA, and volume

matches previous findings in both pigs[24,30] and humans[40–42]. The minor changes in ACL

morphology observed in this study, especially after normalizing to the overall joint size are

similar to other findings directly relating intact and castrated animals[22]. A previous study

measured ACL CSA after 12-weeks in rats castrated as adults and intact male rats and found that

the ACL CSA did not differ between the two groups. The volume of the ACL and the AM and

PL bundles had more substantial changes between boars and barrows, with barrows having

smaller ACLs, AM bundles, and PL bundles than boars. The more substantial changes in ACL,

AM bundle, and PL bundle volume could simply be a result of magnifying smaller changes seen

in the CSA and length or could be of a result of larger differences near the entheses of the ACL.

However, normalizing to joint size also diminished these differences in volume, indicating that

surgical castration may influence absolute size, but may not influence normalized joint properties

that can be used to compare individuals.

Regarding mechanics, both boars and barrows had increasing ACL, AM bundle, and PL

bundle stiffnesses across skeletal growth, yet no significant differences were observed between

boars and barrows. While the stiffness of the joint and the tissues increased throughout skeletal

growth, there were also minor increases across growth for boars and barrows in absolute joint

laxity as measured by APTT, but similar laxity values once normalized to joint size.

Furthermore, the barrows had on average, less lax joints and slack, especially at 3 months, yet

this disappeared after normalizing to joint size. The normalized laxity values for the boars and

barrows only varied 5.8% in the boars and 5.1% in the barrows across ages. Human studies often

report raw values of APTT at static loads across ages, such as one study that showed that

skeletally mature men have a decreased knee laxity compared to immature men where female

knee laxity does not change[43]. This differs in this study where scaled loads were applied and

laxity values were similar within boars and barrows. If static loads were applied, then there

might have been similar decreases in absolute laxity across skeletal growth.

Previously, the pig model has been used to detect differences in bundle function across

skeletal growth and found that the AM bundle takes on more of a dominant load (92% by 6-

months in males and 76% by 6-months in females) over the PL bundle[30]. The AM bundle

contribution in the barrows matches the 76% AM bundle contribution in female pigs at 6-months

yet is higher than boars or females at 3 months (11% and 44%, respectively). Temporally, the

female AM bundle contribution increases from 3- to 4.5- to 6-months, boars have a rapid rise

between 3- and 4.5-months then plateau, and barrows have higher AM bundle contribution from

3-months and gradually rise over time. Because there are no significant differences in the

normalized morphological or stiffness values of the ACL or its bundles, it is interesting that there

is a slight, yet not statistically significant, difference in the bundle contribution to anterior

loading, where the bundles in the barrow behave more similar to female pigs than to intact male

pigs (Figure S3). When looking at the bundle function across skeletal growth and overall joint

size, the timing seems to support an earlier maturation in boars than in barrows, yet CSA and

stiffness values seems to indicate no difference in ACL maturation. Perhaps testosterone has

greater influence on the microstructural properties of the collagen fibers in the ACL, as shown by

a decrease in collagen content and increase in collagen fiber diameter in castrated male rabbits[44].

Given that the rise of testosterone in boars starts around 4-months[45], the slight difference in

timing of functional changes could be influenced more by testosterone especially between the 3-

month and 4.5-month range. Additional variables could also be at play when making

comparisons between gilts, boars, and barrows, such as fluctuations in other female sex

hormones[14–18].

Results from this work suggest that surgical castration at an early age has minor impacts

on the ACL structure and function several months later into early adolescence. The current

results cannot explain the continued rise in ACL, AM bundle, and PL bundle CSA in intact male

pigs in early adolescence compared to a plateau in ACL and PL bundle CSA in female pigs over

the same age period. Other factors exist, such as the equilibrium of the conversion between

androgens and estrogens via the aromatase enzyme[46]. With this conversion, castration can also

decrease serum estrogens in males. With other studies showing that females have higher joint

laxity than males, which is a known risk factor for ACL injury[47,48], the previous work in this pig

model showing a 38% greater APTT for female pigs at 18-months[30], and this work showing

similar normalized laxity values for boars and barrows, the future work should focus on the need

to directly measure multiple hormonal factors in females.

There are some limitations to the present study that may provide new pathways to

understanding the role of testosterone in the developing male ACL. The primary limitation is that

while the pig is a good model for understanding human knee function due to its similar

anatomical structures and size, no animal model completely replicates the human condition.

Another limitation of this study is the loss of stabilizing muscles surrounding the knee during

applied loads. Testosterone has been shown to have more drastic impacts on muscle mass and

strength[49,50], so there still could be a greater impact of testosterone on *in vivo* joint stability

between boars and barrows due to greater muscle stabilization. When looking at the functional

characteristics of the ACL and its bundles, only sub-failure properties were assessed. A previous

study in rats discovered that castrated rats had lower load-to-failure forces and ultimate tensile

stresses than intact males, yet these failure properties were not assessed in this current study[22].

Furthermore, there was a high variability in the barrow bundle percent contributions. Due to

limitations of specimen collection, there was also a lack of serum testosterone measurements in

the boars, so associative analyses could not be done. Longer-term data in older animals would

perhaps show a long-term impact of testosterone after its normal rise at 4 months. Sex hormones

are generally complex in nature, with independent hormones rarely impacting macroscopic

properties, and studying the interaction between fluctuating levels of serum sex hormones will

prove useful in attempting to understand sex-specific differences throughout skeletal growth and

ACL injury risk.

In conclusion, surgical castration had subtle impacts on joint and ACL morphology and

function across skeletal growth. Across skeletal growth, the boars and barrows had increasing

joint sizes, ACL CSA values and volumes, and stiffnesses. Between boars and barrows, the joint

size, ACL, AM bundle, and PL bundle volumes were smaller in the barrow group than the boars

across ages, but not the ACL CSA, stiffness, or bundle contribution. The barrows had a more

gradual shift in bundle function toward AM-dominance in resisting load than boars, although this

was not statistically significant due to high variability in the barrows. Thus, these results show

that early and sustained loss in testosterone leads to some differences in ACL morphology but

may not influence measures that may indicate increased injury risk, such as CSA or force in the

bundles in response to applied loads. This work is beneficial in understanding how hormones

might impact ACL performance and injury risk throughout male development.

Acknowledgments

We would like to thank the Swine Education Unit at North Carolina State University,

Laboratory Animal Resources at North Carolina State University, and the Biomedical Research

Imaging Center at the University of North Carolina – Chapel Hill for their contributions to this

work. Research in the publication was supported by the National Institute of Arthritis and

Musculoskeletal and Skin Diseases of the US National Institutes of Health (R01 AR071985 and

F31AR077997). The content is solely the responsibility of the authors and does not necessarily

represent the official views of the National Institutes of Health. Additional funding for this study

was provide in part by the National Science Foundation (DGE-1746939).

References

1. Smucny M, Parikh SN, Pandya NK. 2015. Consequences of Single Sport Specialization in the

Pediatric and Adolescent Athlete. Orthop. Clin. North Am. 46(2):249–258.

2. Beck NA, Lawrence JTR, Nordin JD, et al. 2017. ACL tears in school-aged children and

adolescents over 20 years. Pediatrics 139(3).

3. Weitz FK, Sillanpää PJ, Mattila VM. 2020. The incidence of paediatric ACL injury is

increasing in Finland. Knee Surgery, Sport. Traumatol. Arthrosc. 28(2):363–368.

4. Werner BC, Yang S, Looney AM, Gwathmey FW. 2016. Trends in pediatric and adolescent

anterior cruciate ligament injury and reconstruction. J. Pediatr. Orthop. 36(5):447–452.

5. Bram JT, Magee LC, Mehta NN, et al. 2021. Anterior Cruciate Ligament Injury Incidence in

Adolescent Athletes: A Systematic Review and Meta-analysis. Am. J. Sports Med.

49(7):1962–1972.

6. Joseph AM, Collins CL, Henke NM, et al. 2013. A multisport epidemiologic comparison of

anterior cruciate ligament injuries in high school athletics. J. Athl. Train. 48(6):810–817.

7. Messina DF, Farney WC, DeLee JC. 1999. The incidence of injury in Texas high school

basketball. A prospective study among male and female athletes. Am. J. Sports Med.

27(3):294–299.

8. Stanley LE, Kerr ZY, Dompier TP, Padua DA. 2016. Sex Differences in the Incidence of

Anterior Cruciate Ligament, Medial Collateral Ligament, and Meniscal Injuries in

Collegiate and High School Sports: 2009-2010 Through 2013-2014. Am. J. Sports Med.

44(6):1565–1572.

9. Gupta AS, Pierpoint LA, Comstock RD, Saper MG. 2020. Sex-Based Differences in Anterior

Cruciate Ligament Injuries Among United States High School Soccer Players: An

Epidemiological Study. Orthop. J. Sport. Med. 8(5):2325967120919178.

10. Cook CJ, Kilduff LP, Crewther BT. 2018. Basal and stress-induced salivary testosterone

variation across the menstrual cycle and linkage to motivation and muscle power. Scand.

J. Med. Sci. Sports 28(4):1345–1353.

11. Cook CJ, Crewther BT. 2019. Within- and between-person variation in morning testosterone

is associated with economic risk-related decisions in athletic women across the menstrual

cycle. Horm. Behav. 112:77–80.

12. Parsons JL, Coen SE, Bekker S. 2021. Anterior cruciate ligament injury: towards a gendered

environmental approach. Br. J. Sports Med. 55(17):984–990.

13. Legerlotz K, Nobis T. 2022. Insights in the Effect of Fluctuating Female Hormones on Injury

Risk—Challenge and Chance. Front. Physiol. 13:827726.

14. Wojtys EM, Huston LJ, Lindenfeld TN, et al. 1998. Association between the menstrual cycle

and anterior cruciate ligament injuries in female athletes. Am. J. Sports Med. 26(5):614–

619.

15. Slauterbeck JR, Fuzie SF, Smith MP, et al. 2002. The menstrual cycle, sex hormones, and

anterior cruciate ligament injury. J. Athl. Train. 37(3):275–278.

16. Wojtys EM, Huston LJ, Boynton MD, et al. 2002. The effect of the menstrual cycle on

anterior cruciate ligament injuries in women as determined by hormone levels. Am. J.

Sports Med. 30(2):182–188.

17. Shultz SJ, Kirk SE, Johnson ML, et al. 2004. Relationship between sex hormones and

anterior knee laxity across the menstrual cycle. Med. Sci. Sports Exerc. 36(7):1165–1174.

18. Adachi N, Nawata K, Maeta M, Kurozawa Y. 2008. Relationship of the menstrual cycle

phase to anterior cruciate ligament injuries in teenaged female athletes. Arch. Orthop.

Trauma Surg. 128(5):473–478.

19. Hamlet WP, Liu SH, Panossian V, Finerman GAM. 1997. Primary immunolocalization of

androgen target cells in the human anterior cruciate ligament. J. Orthop. Res. 15(5):657–

663.

20. Stijak L, Kadija M, Djulejić V, et al. 2015. The influence of sex hormones on anterior

cruciate ligament rupture: female study. Knee Surgery, Sport. Traumatol. Arthrosc.

23(9):2742–2749.

21. Liu C, Ma J, Zhang J, et al. 2019. Testosterone deficiency caused by castration modulates

mitochondrial biogenesis through the AR/PGC1α/TFAM pathway. Front. Genet.

10(MAY):505.

22. Romani WA, Belkoff SM, Elisseeff JH. 2016. Testosterone may increase rat anterior cruciate

ligament strength. Knee 23(6):1069–1073.

23. Cone SG, Simpson SG, Piedrahita JA, et al. 2017. Orientation changes in the cruciate

ligaments of the knee during skeletal growth: A porcine model. J. Orthop. Res.

35(12):2725–2732.

24. Cone SG, Lambeth EP, Ru H, et al. 2019. Biomechanical Function and Size of the

Anteromedial and Posterolateral Bundles of the ACL Change Differently with Skeletal

Growth in the Pig Model. Clin. Orthop. Relat. Res. 477(9):2161–2174.

25. Cone SG, Piedrahita JA, Spang JT, Fisher MB. 2019. In situ joint stiffness increases during

skeletal growth but decreases following partial and complete anterior cruciate ligament

injury. J. Biomech. Eng. 141(12).

26. Cone SG, Lambeth EP, Piedrahita JA, et al. 2020. Joint laxity varies in response to partial

and complete anterior cruciate ligament injuries throughout skeletal growth. J. Biomech.

101:109636.

27. Murray MM, Magarian EM, Harrison SL, et al. 2010. The effect of skeletal maturity on

functional healing of the anterior cruciate ligament. J. Bone Jt. Surg. 92(11):2039–2049.

28. Mastrangelo AN, Haus BM, Vavken P, et al. 2010. Immature animals have higher cellular

density in the healing anterior cruciate ligament than adolescent or adult animals. J.

Orthop. Res. 28(8):1100–1106.

29. Cone SG, Barnes RH, Howe D, et al. 2022. Age- and sex-specific differences in ACL and

ACL bundle size during adolescent growth. J. Orthop. Res. 40(7):1613–1620.

30. Howe D, Cone SG, Piedrahita JA, et al. 2022. Sex-specific biomechanics and morphology of

the anterior cruciate ligament during skeletal growth in a porcine model. J. Orthop. Res.

40(8):1853–1864.

31. Debski RE, Yamakawa S, Musahl V, Fujie H. 2017. Use of Robotic Manipulators to Study

Diarthrodial Joint Function. J. Biomech. Eng. 139(2).

32. Fujie H, Livesay GA, Woo SLY, et al. 1995. The use of a universal force-moment sensor to

determine in-Situ forces in ligaments: A new Methodology. J. Biomech. Eng. 117(1):1–7.

33. Imhauser CW, Kent RN, Boorman-Padgett J, et al. 2017. New parameters describing how

knee ligaments carry force in situ predict interspecimen variations in laxity during

simulated clinical exams. J. Biomech. 64:212–218.

34. Satopää V, Albrecht J, Irwin D, Raghavan B. 2011. Finding a "kneedle" in a haystack:

Detecting knee points in system behavior. Proc. - Int. Conf. Distrib. Comput. Syst. :166–

171.

35. Bañón S, Andreu C, Laencina J, Garrido MD. 2004. Fresh and eating pork quality from

entire versus castrate heavy males. Food Qual. Prefer. 15(3):293–300.

36. Knudson BK, Hogberg MG, Merkel RA, et al. 1985. Developmental comparisons of boars

and barrows: II. Body composition and bone development. J. Anim. Sci. 61(4):797–801.

37. Knudson BK, Hogberg MG, Merkel RA, et al. 1985. Developmental comparisons of boars

and barrows: I. Growth rate, carcass and muscle characteristics. J. Anim. Sci. 61(4):789–

796.

38. Castell AG, Strain JH. 1985. INFLUENCE OF DIET AND SEX-TYPE (BOAR,

CASTRATE OR GILT) ON LIVE AND CARCASS MEASUREMENTS OF SELF-FED

PIGS FROM TWO BREED LINES DIFFERING IN GROWTH RATES. undefined

65(1):185–195.

39. Bhasin S, Storer TW, Berman N, et al. 1996. The effects of supraphysiologic doses of

testosterone on muscle size and strength in normal men. N. Engl. J. Med. 335(1):1–7.

40. Edmonds EW, Bathen M, Bastrom TP. 2015. Normal Parameters of the Skeletally Immature

Knee: Developmental Changes on Magnetic Resonance Imaging. J. Pediatr. Orthop.

35(7):712–720.

41. Hosseinzadeh S, Kiapour AM. 2021. Age-related changes in ACL morphology during

skeletal growth and maturation are different between females and males. J. Orthop. Res.

39(4):841–849.

42. Putur DE, Slaven SE, Niu EL. 2020. ACL Growth With Age in Pediatric Patients: An MRI

Study. J. Pediatr. Orthop. 40(8):438–447.

43. Ahmad CS, Clark AM, Heilmann N, et al. 2006. Effect of gender and maturity on

quadriceps-to-hamstring strength ratio and anterior cruciate ligament laxity. Am. J. Sports

Med. 34(3):370–374.

44. Light VA, Montgomery RD, Akingbemi BT. 2012. Sex hormone regulation of collagen

concentrations in cranial cruciate ligaments of sexually immature male rabbits. Am. J.

Vet. Res. 73(8):1186–1193.

45. FlorCruz S V., Lapwood KR. 1978. A Longitudinal Study of Pubertal Development in Boars

Investigation of the Relationships Between Gonadal and Epididymal Development and

Plasma Luteinizing Hormone and Testosterone Profiles. Int. J. Androl. 1(1–6):317–330.

46. Capellino S, Straub RH, Cutolo M. 2014. Aromatase and regulation of the estrogen-to-

androgen ratio in synovial tissue inflammation: common pathway in both sexes. Ann. N.

Y. Acad. Sci. 1317(1):24–31.

47. Bayer S, Meredith SJ, Wilson K, et al. 2020. Knee Morphological Risk Factors for Anterior

Cruciate Ligament Injury: A Systematic Review. J. Bone Joint Surg. Am. 102(8):703–

718.

48. Alentorn-Geli E, Myer GD, Silvers HJ, et al. 2009. Prevention of non-contact anterior

cruciate ligament injuries in soccer players. Part 1: Mechanisms of injury and underlying

risk factors. Knee Surg. Sports Traumatol. Arthrosc. 17(7):705–729.

49. Griggs RC, Kingston W, Jozefowicz RF, et al. 1989. Effect of testosterone on muscle mass

and muscle protein synthesis. J. Appl. Physiol. 66(1):498–503.

50. Badash I, Wu B, Berger M, et al. 2017. Testosterone Supplementation Increases Lean Mass

in Men Undergoing Anterior Cruciate Ligament Reconstruction: A Randomized

Controlled Trial. Orthop. J. Sport. Med. 5(7 suppl6):2325967117S0026.