ciRS-7-miR7 regulate ischemia induced neuronal death via glutamatergic signaling

Flavia Scoyni[1], Valeriia Sitnikova[1*], Luca Giudice[1*], Paula Korhonen[1], Davide M Trevisan[2], Ana Hernandez de Sande[3], Mireia Gomez-Budia[1], Raisa Giniatullina[1], Irene F Ugidos[1], Hiramani Dhungana[4], Cristiana Pistono[1], Nea Korvenlaita[1], Nelli-Noora Välimäki[1], Salla M Kangas[5], Anniina E Hiltunen[6], Minna U Kaikkonen-Määttä[1], Jari Koistinaho[1,4], Seppo Ylä-Herttuala[1], Reetta Hinttala[5,6], Morten T Venø[7,8], Junyi Su[8], Markus Stoffel[9], Anne Schaefer[10,11], Nikolaus Rajewsky[12], Jørgen Kjems[8], Mary P LaPierre[9], Monika Piwecka[13], Jukka Jolkkonen[1], Rashid Giniatullin[1], Thomas B Hansen[8,14], Malm Tarja[1]

Affiliations:
[1]A.I.Virtanen Institute for Molecular Sciences, University of Eastern Finland, Kuopio, Finland
[2]Department of Biosciences and Nutrition, Karolinska Institute, Stockholm, Sweden
[3]School of Medicine, University of Eastern Finland, Kuopio, Finland
[4]Neuroscience Center, University of Helsinki, Finland
[5]Biocenter Oulu, University of Oulu, Oulu, Finland
[6]Medical Research Center Oulu and Research Unit of Clinical Medicine, University of Oulu and Oulu University Hospital, Oulu, Finland
[7]Omiics ApS, Aarhus, Denmark
[8]Interdisciplinary Nanoscience Center, Department of Molecular Biology and Genetics, Aarhus University, Aarhus, Denmark
[9]Institute of Molecular Health Sciences, ETH Zurich, Zürich, Switzerland
[10]Departments of Neuroscience and Psychiatry, Icahn School of Medicine at Mount Sinai, New York, USA
[11]Max Planck Institute, Biology of Ageing, Cologne, Germany
[12]Systems Biology of Gene Regulatory Elements, Max-Delbrück-Center for Molecular Medicine in the Helmholtz Association (MDC), Berlin Institute for Medical Systems Biology (BIMSB), Berlin, Germany
[13]Institute of Bioorganic Chemistry, Polish Academy of Sciences, Poznan, Poland
[14]Present address: Targovax ASA, Lysaker, Norway

*These authors contributed equally to this work

ABSTRACT

Brain functionality resides on finely tuned regulation of gene expression by networks of non-coding

RNAs (ncRNAs) such as the one composed by the circular RNA ciRS-7, the microRNA miR-7 and

the long non-coding RNA Cyrano. However, very little is known on how this network regulates stress

responses in neurodegeneration. Here we describe ischemia induced alterations in the ncRNA

network both *in vitro* and *in vivo* and in transgenic mice lacking ciRS-7 or miR-7. Our data show that

cortical neurons downregulate ciRS-7 and Cyrano and upregulate miR-7 expression upon ischemic

insults. Mice lacking ciRS-7 show reduced lesion size and motor impairment, whilst the absence of

miR-7 alone leads to an increase in the ischemia induced neuronal death. Moreover, miR-7 levels in

pyramidal excitatory neurons regulate dendrite morphology and glutamatergic signaling suggesting

a potential molecular link to the *in vivo* phenotype. Our data reveal a new endogenous mechanism by

which ciRS-7 and miR-7 regulate the outcome of ischemic stroke and shed new light into the

pathophysiological roles of intracellular networks of non-coding RNAs in the brain.

INTRODUCTION

The specialized, dynamic, and complex functionality of the brain relies on accurate regulation of gene expression, also mediated by non-coding RNAs (ncRNAs) which are particularly overrepresented in the brain and have been strongly connected with the increase of cognitive complexity in human[1]. MicroRNAs (miRNAs) are short (~22 nucleotides) ncRNAs known to post-transcriptionally regulate the expression of messenger RNA (mRNA) by binding short complementary sequences (seed)[2] and triggering mRNA decay or inhibition of translation[3]. Long non-coding RNAs (lncRNAs) are molecules of regulatory RNA longer than 200 nucleotides capable of regulating gene expression by interacting with other ncRNAs such as miRNAs, or proteins through secondary structures[4]. Circular RNAs (circRNAs) belong to a novel class of ncRNAs that, among other functions, are able to modulate gene expression by interacting with miRNAs, hence affecting miRNAs activity on mRNA targets[5]. CircRNAs are generated by backsplicing of linear transcripts, a specific splicing event in which the 5' splice site of the upstream exon binds to the 3' splice site of the downstream exon[6]. Thus, in RNA-sequencing data, the reads on the backsplice junction identify the circular specific form of the transcript and the circRNA abundance is defined by the ratio of these reads to the canonical splicing reads.

These ncRNAs independently control cellular function by regulating the expression of protein coding genes, but also interact with each other. An excellent example of ncRNAs regulatory network is the brain specific network composed by two miRNAs (miR-7, miR-671), a lncRNA (Cyrano) and a circRNA (ciRS-7)[5,7–9]. ciRS-7 is suggested to stabilize and promote miR-7 targeting through delocalization of miR-7 in different neuronal compartments[8,9]. On the other end, Cyrano has a single site of almost perfect complementarity to miR-7 which induces the degradation of miR-7 through target RNA–directed miRNA degradation (TDMD)[8]. The regulation of miR-7 mediated by Cyrano affect ciRS-7 abundancy and localization creating a network of interactions between these molecules that at the same time indirectly affect the expression of miR-7 target genes[5,8–10]. Gain and loss of

function experiments and the development of knock-out animals for ciRS-7, miR-7 and Cyrano have

revealed that Cyrano and ciRS-7 bind miR-7 and regulate its expression[5,8,9]. Nonetheless, no

information is available on how this network regulates gene expression in pathophysiological

conditions.

Ischemic stroke, induced by occlusion of one of the main cerebral arteries, leads to lack of oxygen

and nutrients in the brain parenchyma and subsequent cell death. Lack of energy and disruption in

the ion balance cause an uncontrolled release of glutamate in excitatory neurons and deficient uptake

of glutamate by astrocytes induces glutamate excitotoxity[11] and subsequent oxidative stress, necrosis

and apoptosis[12]. Due to the complexity of the ischemic stroke pathophysiology, it is not surprising

that numerous miRNAs and lncRNAs have been identified to regulate oxidative stress response and

glutamate excitotoxicity contributing to the outcome of stroke[13]. In fact, when the stress is sustained,

miRNAs are known to be involved in adaptative sudden switches of gene expression programs[14]. The

efficiency of miRNA-mediated target repression depends on the availability of miRNAs to interact

(expression, localization, activity) and the amount of mRNA targets that have a seed Mirna

Recognition Element (MRE), which confer robustness to crucial cellular functions creating a miRNA-

specific threshold[15]. In this context, sudden changes in MRE containing sequences can lead to an

imbalance in miRNA targeting and de-repression of specific targets. The crosstalk between miRNA

and all the expressed sequences containing MRE at any given moment suggested a theory that in

physiological conditions, a large number of MRE sequences compete for the binding of the miRNA[16].

Moreover, the expression of specific transcript with different binding strength or MRE abundance

during stress is known to be used by organisms to regulate miRNA effects in a process called target

mimicry[17].

Recent studies have also highlighted a strong connection between circRNAs and the production of

reactive oxygen species[18]. However, even though single ncRNAs have been studied in the context of

ischemic stroke[13], a comprehensive analysis of the regulatory role of ncRNAs network in stroke is

still lacking. In this study, we identified the brain highly expressed circRNA, ciRS-7, as one of the

seven circRNAs deregulated in ischemia-related conditions *in vitro* and *in vivo*. We expanded our

analysis to other ncRNAs species and discovered additional changes in miR-7 and Cyrano which

belong to the same regulatory circuitry of ciRS-7. Using ciRS-7 and miR-7 knock-out mouse models,

we revealed the potential role of ciRS-7 in preventing miR-7-mediated regulation of the target

mRNAs and demonstrate that ciRS-7 molecular network is involved in the regulation of stroke-

induced cell death.

RESULTS

Oxygen and glucose deprivation induces changes in the expression of circular RNAs

As circRNAs were previously connected to oxidative stress response[18] we evaluated the potential

deregulation of circRNAs in conditions mimicking ischemic stroke *in vitro*. We cultured murine

cortical neurons isolated from embryonic day fifteen (E15) cortices, subjected them to oxygen and

glucose deprivation (OGD) (Supplementary Figure S1A) and performed total RNA sequencing. To

assess whether our neuronal culture is composed of cell types representative of the murine cortex,

we inferred the presence of individual cell populations in our bulk RNA-seq samples through a

deconvolution analysis on single-cell RNA-seq dataset of embryonic mouse brain (E14.5[19]). The

results confirmed that interneurons (Int) and pyramidal neurons (Layer V-VI) are the most abundant

cell populations in our culture (Figure 1A). To validate that the culture is effectively enriched in

functional Interneurons (*GABAergic*) and Pyramidal neurons (*Glutamatergic*), we stimulated our

neuronal culture with GABA and glutamate and recorded the consequent induction of intracellular

calcium using calcium imaging. Our data confirmed that 98% of cells in our culture responded to

glutamate and 36% to GABA, providing functional proof that our culture mostly consists of glutamatergic excitatory neurons (Figure 1B).

Ingenuity Pathway Analysis (IPA) of the differentially expressed transcripts between normoxic and OGD neurons confirmed that OGD treatment induced the activation of Hypoxia-inducible factor 1-alpha (HIF-1alpha) (Figure 1C) and downstream upregulation of glycolysis, a characteristic hallmark of OGD[20] (Supplementary Table S1). Moreover, colorimetric cell viability assay indicated 30% decrease in neuronal viability upon the exposure (Supplementary Figure S1B), validating that our experimental setting represents a population of glutamatergic excitatory neurons that are vulnerable to ischemia-induced cell death.

CircRNAs are generated by back-splicing which produces transcripts that can be detected with back-splice junction reads (BSJ) spanning regions that are not present in regularly spliced transcripts (forward splice junction reads, FSJ) (Figure 1D). In order to identify OGD-induced alterations in circRNAs, we used *CIRIquant* algorithm[21], which considers circular specific reads (BSJ) and their ratio with linear transcript reads (FSJ), and identified 27 circRNAs with high confidence of which 7 were significantly differentially expressed (DE) between normoxic (control) and OGD conditions (Figure 1E, Supplementary Table S2). Among these DE circRNAs, ciRS-7 was the most highly expressed yielding the highest number of back-splice junctions (Figure 1F, left) and the highest circular to linear ratio (Junction ratio) (Figure 1F, right). In fact, ciRS-7 obtained a junction ratio score above 0.8, meaning that more than 80% of the transcripts originating from this locus are circular, in contrast with the other circRNAs which showed about 40% circular expression. Specifically, ciRS-7 was one of the three downregulated circRNAs upon OGD.

ciRS-7 network is altered in conditions mimicking ischemic stroke in vitro

ciRS-7 has been shown to be part of a feedback loop with miR-671 and miR-7 microRNA[5,7] and indirectly, through miR-7, with the long non-coding RNA Cyrano[8] (Figure 2A). To investigate the

relative relationship between these players in *in vitro* OGD conditions, we performed small RNA

sequencing of cortical neurons exposed to OGD. Together with the previously presented dataset, we

were able to capture circRNAs, mRNAs, long non-coding RNAs and small RNA transcripts

(Supplementary Table S2).

Small RNA-sequencing identified 333 upregulated and 88 downregulated miRNAs following OGD

(Figure 2B). The levels of miR-671-5p remained unaltered (Figure 2B) which was confirmed

independently by RT-qPCR (Supplementary Figure S2A), in contrast to previous findings of

diverging miR-671-5p and ciRS-7 expression in different contexts[7–9]. Instead, we detected a

significant upregulation in miR-7a-5p, but not of the variant miR-7b-5p (Figure 2B, Supplementary

Table S2). Similarly to human, in mouse miR-7 is redundantly encoded by three different loci (*miR-*

7a-1, *miR-7a-2*, *miR-7b*), each is produced from different primary transcript (pri-miRNA) and

precursor (pre-miRNA). Further processing, in all cases from -5p arm, give rise to two mature miR-

7 sequences (miR-7a and miR-7b), differing only by a single nucleotide in the non-seed position

number 10 (Figure 2C). To study the expression of these three miR-7 loci in our system, we took

advantage of Global run-on sequencing (GRO-seq), which allows the capture of nuclear nascent RNA

primary molecules, and compared our murine culture expression[22] with different mouse tissues and

cell lines[22–32] (Supplementary Table S3). In physiological conditions, only cortical neurons and the

pituitary gland actively transcribe all three independently regulated *miR-7* loci (Figure 2D). The

active transcription from three different loci in neuronal cells suggests a higher order of regulation of

this miRNA which influences the mature forms. Under OGD conditions, only the pri-miRNAs

contributing to the expression of miR-7a (miR-7a-1 and miR-7a-2) were significantly upregulated

(Figure 2E), indicating that the observed upregulation occurs already at the level of transcription.

None of miR-7 passenger strands generated from the three precursors (miR-7a-1-3p, miR-7a-2-3p,

and miR-7b-3p) were altered in our sequencing data (Supplementary Table S2). Moreover, we

detected no changes in the host gene in which miR-7a-1 is embedded *(Hnrnpk)*, and from which the

most abundant miR-7 primary transcript is generated (Supplementary Table S2). Taken together these

data show that OGD specifically regulates miR-7a-5p variant at the transcriptional and/or post-

transcriptional level.

To test the possible functional relevance of the upregulation of miR-7a-5p, we acquired predicted and

validated targets of miR-7a-5p using *miRWalk* algorithm analysis[33,34] and compared their overall

expression in conditions of OGD. In accordance with the upregulation of miR-7, miR-7 targets

showed a significant overall downregulation (Figure 2F, Supplementary Table S4), suggesting a

canonical functional repressive role of this miRNA.

The role of miR-7 interaction with ciRS-7 remains controversial, and it is thought to set the balance

between a positive and negative feedback[8,9] (Figure 2A). Additionally, Cyrano promotes template

mediated degradation (TMD) of miR-7 through a site of almost perfect complementarity, which was

shown to indirectly regulate ciRS-7 levels and localization[8] (Figure 2A). In our dataset the lncRNA

Cyrano was significantly downregulated upon OGD (Figure 2G), in line with the observed

upregulation in miR-7a-5p (Figure 2B).

To identify a first responder to OGD in the context of this molecular circuitry we evaluate the time-

dependent changes in the levels of these molecules. We subjected cortical neurons to OGD for 1, 6,

12 and 24 hours and assessed the expression level of ciRS-7, miR-7 and Cyrano. We detected a

significant downregulation of ciRS-7 and Cyrano already at 6 hours after OGD, prior to the

upregulation of miR-7 at 12 hours (Figure 2H). Interestingly, our data revealed a significant

downregulation of miR-7 at 6h of OGD concomitantly with downregulation in ciRS-7 and Cyrano,

which is in line with the previously suggested role of ciRS-7 in stabilizing miR-7[9].

ciRS-7 KO neurons exhibit differential OGD response

The absence of ciRS-7 alters synaptic transmission in excitatory neurons and produces schizophrenia-

like phenotype *in vivo*[9], connecting this molecule to glutamatergic transmission. This is of particular

interest in conditions of ischemic stroke, where glutamate mediated excitotoxicity is a critical

contributor to neuronal cell death[12]. Our cortical neuron culture enriched in glutamatergic neurons

(Figure 1B) provides a system in which these events can be studied *in vitro*. Neurons cultured from

ciRS-7 KO mice showed higher Ca^{2+}-responses to the stimulation with glutamate compared to their

wild-type (WT) counterparts (Figure 3A). Consequently, ciRS-7 KO neurons also exhibited a

significantly higher sensitivity to excitotoxicity upon exposure to high concentration of glutamate

(Figure 3B).

To test if this increased sensitivity to glutamate is affecting gene expression changes in OGD, we

subjected the ciRS-7 KO cortical cultures to OGD following the previous experimental design

(Supplementary Figure S1A) and performed total RNA-seq. OGD induced deregulation in total of

5767 genes in ciRS-7 KO neurons compared to normoxic conditions, of which the 85% were shared

with WT neurons (Figure 3C). In response to OGD, 871 genes were differentially expressed only in

ciRS-7 KO neurons, while the expression 1260 genes was exclusively altered in WT neurons

subjected to OGD. ciRS-7 WT specific genes altered in OGD were functionally enriched in

mitochondrial processes, DNA metabolism, regulation of cell-cycle and oxidative stress-induced

apoptosis (Figure 3D). The 871 KO-specific genes were instead involved in developmental and pro-

regenerative processes (*progenitor migration to the cortex, DNA replication, transcription,*

cerebellar cortex development) (Figure 3E). Thus, even though the genetic predisposition to

glutamate sensitivity of ciRS-7 KO would suggest a deleterious outcome in the OGD response, ciRS-

7 KO neurons in OGD regulate pathways of resilience. In support of this, ciRS-7 KO neurons did not

show increased cell death compared to WT in response of OGD treatment (Supplementary Figure

S2B). Interestingly, the lack of ciRS-7 did not affect OGD induced changes in the expression of miR-

7 and Cyrano (Figure 3F). Similar to WT neurons, Cyrano lncRNA was downregulated in ciRS-7 KO

neurons responding to OGD (Figure 3F, Supplementary Table S5). Moreover, even though ciRS-7

KO neurons stably express lower levels of miR-7a-5p[9] (Figure 3F, Supplementary Table S5), small-

RNA sequencing revealed a significant upregulation of miR-7a-5p in response to OGD also in these

neurons (Figure 3F, Supplementary Table S5). In line with our previous findings, we detected no

changes in miR-671-5p, miR-7b-5p and miR-7 passenger strands expression (Supplementary Table

S5). Nonetheless, whilst the expression in miR-7a-5p in OGD remained lower in ciRS-7 KO neurons

compared to WT neurons (Figure 3F), the targets of miR-7a-5p, both predicted and validated, showed

a significant global repression in these conditions (Figure 3G).

215
216
Alteration of ciRS-7 network affects ischemic stroke outcome in vivo

To evaluate the involvement of ciRS-7 network in ischemic stroke *in vivo*, *BALB/c* mice were

subjected to permanent middle cerebral artery occlusion (pMCAo) and levels of ciRS-7, Cyrano and

miR-7a-5p were evaluated by qPCR at six hours, one day and five days post-ischemia. In line with

our *in vitro* results, we detected a significant downregulation of ciRS-7 and Cyrano and upregulation

of miR-7a-5p in the peri-ischemic cortex (PI) at one day post-ischemia (dpi) compared to contralateral

cortex (CL) (Figure 4A). Moreover, similar to OGD, we failed to detect alterations in miR-671-5p

levels (Supplementary Figure S2C).

To evaluate the functional relevance of the ambiguous behavior of ciRS-7 KO neurons which showed

glutamate sensitivity but lack of increased cell death in OGD, we assessed whether ciRS-7 KO mice

subjected to transient ischemic stroke would show altered vulnerability to ischemic damage and

associated sensorimotor impairments. The size of the ischemic lesion was measured using Magnetic

Resonance Imaging (MRI) and behavioral deficits were assessed using Neurological Severity Scores

(NSS) at acute (1dpi), subacute (3dpi) and chronic (7dpi) timepoints after transient middle cerebral

artery occlusion (tMCAo) (Figure 4B). At 1dpi, ciRS-7 KO mice showed a significant reduction in

the ischemic lesion volume compared to their WT controls (Figure 4C) and significantly ameliorated

motor deficits (Figure 4D). Interestingly, ciRS-7 KO animals exhibited a reduced lesion size only

transiently at the acute timepoint (1dpi), showing no delay in the progression of the damage

(Supplementary Figure S3A,B).

To identify the molecular changes associated with the acute reduction in the lesion size and motor

deficits of the ciRS-7 KO mice, mice were sacrificed at 1dpi and peri-ischemic and the contralateral

cortex was used for mRNA sequencing. IPA analysis of the differentially expressed genes revealed

that ciRS-7 KO animals exhibited substantial inhibition in the pathways related to neuronal and tissue

damage, cytokine and interleukin release and calcium influx, in their peri-ischemic cortex compared

to WT controls (Figure 4E, Supplementary Table S6). Moreover, the differentially expressed genes

were mostly contributing to pathways of *necrosis* (205 genes) and *quantity of cytokine* (41 genes),

suggesting that the decreased lesion size in the ciRS-7 KO mice may be due to inhibition of necrotic

pathways and reduction in cytokine release.

In order to identify the major cellular responders mediating the observed alterations in gene

expression profiles, we carried out deconvolution analysis by using adult mouse brain scRNA-seq

data[35] to define the contribution of different cell populations to our bulk mRNA sequencing dataset.

This analysis identified neurons and astrocytes as major cellular responders contributing to the

differences between the WT and ciRS-7 KO gene expression profiles (Figure 4F). ciRS-7 KO mice

showed increased neuronal involvement in the peri-infarct cortex compared to the WT mice in line

with the significantly decreased tissue death in ciRS-7 KO mice, as measured by MRI (Figure 4C).

At the same time, ciRS-7 KO animals exhibited a decrease in the contribution of astrocytes in the

peri-ischemic cortex.

We then collected brain slices of ischemic ciRS-7 KO mice at 7dpi, timepoint of maximum immune

response[36], for staining with the astrocytic glial fibrillary acidic protein (GFAP). As expected,

ischemic stroke induced the expression of GFAP in the peri-infarct cortex in both WT and ciRS-7

KO mice. However, ciRS-7 KO mice showed significantly reduced astrogliosis in this area compered

to WT animals (Figure 4G).

At molecular level, tMCAo induced a significant downregulation of ciRS-7 in WT mice at 1dpi in

the peri-ischemic cortex, whereas the levels of miR-7-5p and Cyrano remained unaltered (Figure 4H,

Supplementary Figure S3C). However, ciRS-7 KO mice showed a significant upregulation of miR-

7-5p and unaltered levels of Cyrano (Figure 4H). These data suggest a correlation between the

absence of ciRS-7 and the upregulation of miR-7 in response to transient ischemic stroke.

ciRS-7 prevents miR-7 effects on glutamatergic signaling in excitatory neurons

Previous studies have identified the involvement of miR-7 in response to oxidative stress[37], in

particular to low glucose[38]. To establish the possible functional link between the upregulation of miR-

7 in ciRS-7 KO mice in conditions of ischemic stroke, we subjected cre-loxP inducible miR-7 KO

mice[39] to tMCAo (Figure 5A, Supplementary Figure S4A,B). We failed to detect changes in the lesion

volume as measured by MRI between miR-7 KO and WT mice at acute timepoint (1dpi)

(Supplementary Figure S4C). In line with this data, miR-7 KO mice did not show any differences in

motor deficits compared to their WT controls (Supplementary Figure S4D). However, we detected

an increase in the lesion volume in miR-7 KO mice (Figure 5B) compared to their WT controls at

later timepoint (7 dpi), which was not associated with changes in Neuroscore (Supplementary Figure

S4E), suggesting a potential role of miR-7 in preventing the ischemia induced cell death.

To understand the functional role of miR-7 in ischemic stroke, we first validated that miR-7 has

functional target sites in our system through the analysis of miR-7 target binding sites using Ago2

HITS-CLIP sequencing data in pyramidal excitatory neurons[40] (Figure 5C). We identified RISC-

associated binding sites on expressed transcripts in mature projection neurons and the analysis of

minimal seed site hexamer of miR-7 revealed 645 potential targets in the 3′ UTRs (Figure 5D). As

expected, ciRS-7 obtained the highest number of functional binding sites (n=139) and accounted for

more than 11% of the total reads (Supplementary Table S7). The overlap between Ago2 HITS-CLIP

physical targets and the downregulated genes in the peri-ischemic cortex of ciRS-7 KO mice, where

miR-7 is upregulated, highlighted 276 common genes (Figure 5E). These genes were functionally

enriched in pathways related to glutamatergic synaptic transmission and morphological changes of

neuronal projections (Figure 5F, Supplementary Figure S5).

Given the central role of glutamatergic signaling in ischemia, these miR-7 potential targets in

excitatory neurons might relate the improved ischemic outcome and the increased levels of miR-7 in

ciRS-7 KO mice. To test this connection, we transduced ciRS-7 WT and KO neurons with lentivirus

overexpressing GFP and miR-7a or GFP only, under the neuronal specific human Synapsin 1 (hSyn)

promoter. Efficiency of infection was assessed through fluorescence microscopy (Supplementary

Figure S6A) and we validated with RT-qPCR that miR-7a-5p levels resembled the endogenous levels

in conditions of OGD (Supplementary Figure S6B).

Whilst overexpression of miR-7 in WT neurons did not affect the overall neuronal glutamate

excitability, increased levels of miR-7 in ciRS-7 KO neurons significantly reduced their response to

glutamate (Figure 5G), thus reverting the genotype of glutamate sensibility of the ciRS-7 KO neurons.

This appeared to be a glutamate specific effect, as the overexpression of miR-7 did not affect

GABAergic response (Supplementary Figure S6C).

Finally, to evaluate the functional impact of the potential targets of miR-7 involved in morphological

changes of neuronal projections (Figure 5F), we performed live imaging analysis of neurite outgrowth

of ciRS-7 KO and WT neurons infected with miR-7 lentivirus. In agreement with the calcium imaging

experiment, we were able to detect a significant increase in the neurite length when over expressing

miR-7 in ciRS-7 KO cells, but not in WT neurons (Figure 5H). This result is in line with a previous

study demonstrating a direct effect of miR-7 overexpression in controlling neurite length in a

neuroblastoma cell line[41]. However, circRNAs are known to be poorly expressed in cell lines due of

their high proliferation rate[42]. So, we wondered whether this miR-7 direct effect on neurite length

was due to lack of ciRS-7 interference. A study identifying circRNAs in neuronal cell lines and tissue

discovered that human neuroblastoma cell line expresses an exiguous amount of ciRS-7[42]. In addition,

we validated that murine neuroblastoma cell line N2A also express negligible levels of ciRS-7 even

when differentiated using retinoid acid, albeit the levels of Cyrano and miR-7 are more abundant

(Supplementary Figure S7A,B,C).

DISCUSSION

Here we identify how the ncRNAs network composed of ciRS-7 – miR-7 – Cyrano regulates cellular

responses to OGD *in vitro* and ischemic stroke *in vivo*. In OGD *in vitro* model, we identified a new

endogenous relationship between ciRS-7 and miR-7 than previously shown as ciRS-7 and Cyrano

downregulation occurred ahead of miR-7 upregulation which was not dependent on ciRS-7 presence.

In *in vivo* conditions, the absence of ciRS-7 alleviated cellular death and subsequent sensorimotor

deficits in response to ischemic stroke whereas the lack of miR-7 led to a more extensive tissue

damage. These effects were partially executed via regulation of glutamatergic signaling by miR-7.

Together these data highlight an endogenous regulatory role for the ciRS-7 network in mediating

cellular stress responses under pathophysiological conditions.

We hypothesize that our results on the ncRNAs network composed by ciRS-7, miR-7 and Cyrano

reflects a stress response mediated by miR-7 that is modulated by ciRS-7 and Cyrano expression. In

mouse, ciRS-7 harbors 130 MREs (binding sites) for miR-7[9] and, because of its extremely high

expression in neuron[5,9], the low levels of miR-7 in physiological conditions[5] are likely to be bound

to ciRS-7, as suggested by our pyramidal excitatory neurons CLIP-seq data and others[9,10]. Moreover,

Cyrano, which is also very abundant in neurons, harbors a single almost complementary site for miR-

7 which mediates miR-7 degradation[8]. Our data show that in physiological conditions strong and high

abundant miR-7 binding competitors (ciRS-7/Cyrano) interact with limited regulator (miR-7) whereas in conditions of ischemic stroke, both *in vitro* and *in vivo*, there is a concomitant removal of ciRS-7 and Cyrano and subsequent upregulation of miR-7. Our data showing a downregulation in Cyrano prior to the upregulation of miR-7 is in line with previously known dynamics between these two molecules as Cyrano triggers miR-7 degradation through TDMD[8]. On the other hand, here we show for the first time the induction of miR-7 in the absence of ciRS-7 suggesting that expression level of miR-7 is not dependent on ciRS-7 in conditions of ischemic stroke. This data, together with the lack of involvement of miR-7 levels in ciRS-7 downregulation identified new endogenous relationship induced by stress between miR-7 and ciRS-7 in respect of what it has been previously shown in physiological conditions[8,9].

The role of miR-7 in ischemic stroke has been studied *in vivo* with controversial results[43,44]. Our large cohort of animals revealed a high biological variability in expression levels of miR-7 in WT animals subjected to transient ischemic stroke. This can be explained by the presence of feedback loops between ciRS-7, Cyrano and miR-7 and suggests the idea that ciRS-7 and Cyrano may behave as a buffer to regulate miR-7 expression. In line with this idea, ciRS-7 KO mice subjected to tMCAo significantly overexpress miR-7 one day post-ischemia and show a reduced stroke lesion size and motor deficits. Furthermore, at 7 dpi, miR-7 inducible KO mice showed an exaggerated ischemic lesion size compared to WT animals, reinforcing the idea of a role of miR-7 in regulating ischemic damage. Moreover, our data is in line with literature showing that the levels of miR-7 are downregulated at early timepoints in ischemic stroke and that upregulation in miR-7 is necessary for stroke recovery[45]. These data together propose a model in which ciRS-7 and Cyrano regulate miR-7 and possibly influence its upregulation acting as a "buffering" system.

Due to the lack of extensive changes in miR-7 targets in both ciRS-7, Cyrano and miR-7 KO animals

in physiological conditions[8,9], the exact role of ciRS-7 as an influencer of miR-7 target gene

expression has remained controversial. Our data show a significant shift in the overall expression of

miR-7 targets in conditions of OGD (Figure 2F), where a concerted downregulation of two strong

miR-7 regulators, ciRS-7 and Cyrano, occurs. Our data suggests that the inducible nature of this

network may explain the reported lack of extreme effects on miR-7 targets in physiological

conditions. This is supported by studies showing that single miRNA mutants exhibit a phenotype

solely under stress conditions[46,47]. Moreover, removal of a single competing target for endogenously

low expressed miRNAs, such as miR-7, is not expected to induce a relevant change in gene expression

in physiological conditions[15]. Finally, particularly for non-coding RNAs, cells are prone to adaptation

upon knock-out, generating an imbalance in output between knock-out and knock-down experiments,

as previously experienced[9].

We used the analysis of Ago2 CLIP-seq dataset to discover physiological interactions of miR-7 with

the downregulated genes in ciRS-7 KO tMCAo animals in which miR-7 is upregulated. Our data

suggest a role of miR-7 in regulating glutamatergic signaling and neuronal outgrowth. The central

role of glutamate in ischemic stroke pathophysiology is well established[12], contributing to the

neuronal loss upon exaggerated release and impaired clearance of glutamate in the synaptic cleft[11].

Overexpression of miR-7 failed to induce changes in glutamatergic signaling and neuronal outgrowth

in wild-type cortical neurons. However, in the absence of ciRS-7, miR-7 overexpression dampened

the neuronal glutamatergic response which likely contributes to the reduced cellular damage in

conditions of ischemic stroke.

Our hypothesis of ciRS-7 acting as a buffer on miR-7 mediated effects is strengthened further by our

data showing that miR-7 overexpression promotes neurite outgrowth only in ciRS-7 KO neurons,

which lack the ciRS-7 buffering effect. In line with our results, previously published data indicate

that miR-7 overexpression regulates neurite length in a neuroblastoma cell line[41]. Of note is that

murine neuroblastoma cell lines express negligible amounts of ciRS-7 even when differentiated with

retinoic acid, suggesting a more direct effect of miR-7 in the absence of its regulator. A recent study

has highlighted a potential role of miR-7 in controlling energy homeostasis in hypothalamic

neurons[39], linking this miRNA to conditions in which the energetic balance is strongly altered, such

as in ischemic stroke. However, alteration in the levels of miR-7 in hypothalamic neurons were not

accompanied by changes in the expression of ciRS-7 or Cyrano[39]. Together with our data showing

how the type of ischemic insult activates ciRS-7 network with different dynamics, this suggests that

the network responds in a stress type, strength, and duration dependent manner.

In summary, here we identify a role of ciRS-7 as a regulator of cellular stress response under OGD

in vitro and ischemic stroke *in vivo*. Upregulation of miR-7 following cerebral ischemia is critical for

the control of ischemic stroke induced damage and post-ischemia plasticity and depends on the levels

of ciRS-7. Our data support the role for ciRS-7 as a buffer of miR-7 effects against unwanted changes,

thus behaving as a "safe-guide" system. Finally, we propose a regulatory role for ciRS-7-miR-7

network in glutamatergic signaling through miR-7 target genes, hence providing a novel mechanism

which controls post-ischemia neuronal damage. More broadly, this study suggests a role of

intracellular network of non-coding RNAs in regulating pathophysiological processes in the brain.

MATERIAL AND METHODS

Primary cortical neuron culture preparation and OGD treatment

Primary cortical neurons were prepared from C57BL/6J and C57BL/6N-Cdr1asem1Nikr (Cdr1as KO and their WT counterpart) embryonic day 15 embryonal cortices. After dissection and removal of the meninges, cortices were incubated 15 minutes at 37 °C in a solution of Krebs buffer (0.126 M NaCl, 2.5 mM KCl, 25 mM NaHCO3, 1.2 mM NaH2PO4, 1.2 mM MgCl2, 2.5 mM CaCl2, supplemented with 45mM BSA, 0.8% of 3.85% MgSO4 and 1% Pen/Strep, pH 7.4) and 0.025% (w/v) trypsin (Sigma-Aldrich, T 9201). Tissue was then treated with 0.008% w/v DNaseI (Sigma-Aldrich, DN25) and 0.026% w/v trypsin inhibitor (Sigma-Aldrich, T9003) and centrifuged at 300 x g for 3 minutes. Cell pellet was resuspended in 3ml of DNaseI/Trypsin solution and then diluted in 7ml of Krebs. After centrifugation at 300 x g for 3 minutes the pellet containing embryonic neurons was resuspended in cortical neurons growth media: Neurobasal (Gibco 21103049), B27 Supplement (Gibco, 17504044), 0.2 mM L-glutamine (Lonza, BE17-605E), 0.01 mg/ml Penicillin/Streptomycin (Gibco, 15140122). Cells were plated on Poly-D lysine (Sigma-Aldrich, P6407) freshly precoated plates (50 μg/ml in sterile water plates for 1h at 37 °C and washed in sterile water prior use). Different density was used for 6-well plates (1.8 million cells per well), 48-well plates (125.000 cells per well), 13mm plastic coverslips (30.000 cells per coverslip). After 5 days in culture half of the in cortical neurons growth media was changed to fresh. Experiments were performed 7 days after the isolation day. Cells were maintained in the incubator 37 °C, 5% CO2. For OGD treatment experiments, before hypoxia induction the media was changed to Normoxic (cortical neurons growth media with Neurobasal changed to Gibco A2477501 supplemented with D-Glucose 25mM and Sodium Pyruvate 0.2mM) or OGD (cortical neurons growth media with Neurobasal changed to Gibco A2477501). Normoxic cells were then put back in the incubator, OGD cells were incubated in hypoxic chamber 37 °C, 5%CO_2, 1%O_2 (SCI-tive N, Ruskinn Technology). After the OGD timepoint cells were harvested for RNA extraction or subjected to MTT colorimetric test.

MTT test

Cell viability was measured from 48-well plates treating the cells with (3-(4, 5-dimethylthiazolyl-2)-2, 5-diphenyltetrazolium bromide), MTT reagent (Sigma-Aldrich, TOX1) diluted with culture media at a final concentration of 120 μM. Triton-X 100 1% v/v (Sigma-Aldrich, X100) treated wells were used as a positive control for this assay. Plates were incubated for 3-5h at +37°C. After, medium was discarded, formazan crystals were dissolved with dimethyl sulfoxide (DMSO) (Sigma-Aldrich, D2650) for 30min at RT in the dark. Absorbance was read at 585nm using Wallac 1420 Victor2 microplate reader (Perkin Elmer). Wells without cells were used as background and subtracted from the absorbance data, all the six technical replicates were plotted for each of the three biological replicate and the data was normalized on normoxic WT neurons or WT neurons infected with GFP only.

Neurite length measurement

Neurons were seeded at a density of 125.000 cells/well in 48-well plates. After OGD exposure or lentiviral infection, neurons were imaged at day 7 for 48 hours with IncuCyte® S3 Live Cell Analysis System (Essen BioScience Ltd.) in bright field and green channel live cell images (two 10x magnification images per well). Acquired data were analysed with the Incucyte® Neurotrack Analysis Software Module (Sartorius) considering the average value of the two images taken for each of the nine technical replicates for all the three biological replicates.

RNA isolation and qRT-PCR

Total RNA was extracted from primary cells and ipsilateral/contralateral animal cortex using TRIzol™ Reagent (Invitrogen) following the manufacturer's instructions. 1μl of GlycoBlue™ Coprecipitant (Ambion) was added at the isopropanol step in each sample. RNA was quantified using a Nanodrop 2000 spectrophotometer. 1 μg of total RNA was used for reverse-transcription

ofmRNAs, circRNA, lncRNAs and pri-miRNA species with High-Capacity cDNA Reverse

Transcription Kit (Applied Biosystems) following the manufacturer's protocol. SYBR Green qPCR

Master Mix (High ROX) (Bimake) and custom designed oligos were used to quantify mRNAs,

circRNAs, and lncRNAs following the manufacturer's indications. Mature microRNAs were reverse-

transcribed with TaqMan™ MicroRNA Reverse Transcription Kit (Invitrogen) using miRNA-

specific primers supplied in the TaqMan® probe kit for qPCR following the manufacturer's protocol.

Mature microRNAs and pri-microRNA PCRs were performed using Maxima Probe/ROX qPCR

Master Mix (Invitrogen) and TaqMan® specific probes (Thermofisher) following the manufacturer's

protocol. The result was analyzed with the ΔΔCT method and normalized geometric mean of 2

internal normalization controls (Gapdh and Rplp0) for SYBR green qPCRs and U6 expression for

TaqMan® miRNA and pri-miRNA. TaqMan® probes and sequence of the SYBR Green

oligonucleotide primers is available in the Supplementary Table S8.

Lentivirus vectors and virus generation

Lentiviral vectors LV1-eGFP (control) or LV1-eGFP-miR-7 were generated by subcloning inserts

from pAAV_hSYN1-eGFP-miR-7 and pAAV_hSYN1-eGFP (provided by Thomas B. Hansen)

inside LV1 (immunodeficiency virus 1 (HIV-1)-based LV-PGK-GFP) backbone by GenScript

Biotech Corporation. The generated construct contained HIV-1-LV backbone with hSYN1-eGFP-

miR-7 insert instead of PGK-GFPN inserted creating a terminal SmaI (CCCGGG) and a C-terminal

ApaI (GGGCCC) flanking restriction sites. The same was performed for the control vector with

hSYN1-eGFP insert only. 3rd generation lentiviral particles were produced by the BioCenter Kuopio

National Virus Vector Laboratory in Kuopio, Finland. The viral titer was assessed through qPCR

serial dilution quantification using eGFP ReadyMade™ Primers (IDT) (Supplementary Table S8).

Work with the virus vectors was carried out under permission from Finnish National Supervisory

Authority for Welfare and Health, Valvira. Cells were infected at day 2 post isolation with MOI 0.5

achieving 80% positive infected cells assessed by GFP expression in fluorescent microscope

(Supplementary Figure 5A) at day 7. The amount of overexpression of miR-7 was assessed by qPCR

as above specified (Supplementary Figure 5B).

Library preparation

All the samples RNA were isolated with TRIzol™ Reagent (Invitrogen) as specified above. RNA

samples were treated with TURBO DNA-free™ Kit (Ambion) following the manufacture's

instruction. RNA integrity was assessed through Agilent Bioanalyzer 2100 system with the Agilent

RNA 6000 Nano. The concentration of the samples was established with Qubit™ RNA Extended

range kit (Invitrogen). From cortical neurons subjected to OGD we generated 1) library to detect

circularRNAs and mRNA and lncRNAs using SMARTer® Stranded Total RNA Sample Prep Kit

(Takara Bio USA, Inc.) and 2) library for miRNAs detection utilizing NEBNext® Small RNA Library

Prep Set for Illumina (New England Biolabs (UK) Ltd) quality checked and size selected using pippin

prep method following manufacturer's protocol. We generated a library from ciRS-7 WT and KO

tMCAo animals contralateral and peri-ischemic cortices to detect mRNAs changes using CORALL

total RNA-Seq Library Prep Kit (Lexogen GmbH) after ribosomal RNA depletion of 600ng of RNA

with RiboCop rRNA Depletion Kit V1.2 (Lexogen GmbH). All the libraries were generated following

the manufacturer's protocol. After generation, the libraries were quantified with Qubit™ High

Sensitivity DNA kit (Invitrogen) and by qPCR using KAPA Library Quantification Kit for Illumina®

Platforms (Roche). Library size was determined with Agilent Bioanalyzer 2100 system using the

Agilent High Sensitivity DNA Kit. Cortical neurons samples to detect circularRNAs, mRNA, and

lncRNAs were sequenced paired-end 100 cycles on NovaSeq™ 6000 platform (Illumina) and single-

read 75 cycles on NextSeq™ 500 system (Illumina) to detect microRNAs. Animal samples were

sequenced single-read 75 cycles on NextSeq™ 550.

Calcium imaging

Calcium imaging was performed with murine E15 cortical neurons. The neurons were plated onto PDL-coated circular plastic coverslips (13 mm diameter) in 12-well plate at a density of 30,000 cells/coverslip and kept for 5 days in Neurobasal medium supplemented with B27. To quantify and compare the functional expression of glutamate and GABA receptors in neuronal cultures, we used calcium-imaging technique as previously described[48]. Briefly, neuronal cultures were loaded with the cell-permeable indicator Fluo-4am (Life Technologies, F10471) for 30 min at 37 °C, followed by 10 min washout, and placed in the perfusion chamber mounted on the stage of Olympus IX7010 microscope. Neurons were continuously perfused by basic salt solution (BSS) 3 ml/min containing in mM: 152 NaCl, 10 HEPES, 10 glucose, 2.5 KCl, 2 CaCl2, 1 MgCl2 and pH adjusted to 7.4. Test compounds diluted in the BSS to final concentrations were applied through fast perfusion system (Rapid Solution Changer RSC-200, BioLogic Science Instruments). Cells were imaged with 10x objective using Olympus IX-7010. Excitation wavelength was set as 494 nm, sampling frequency 2 FPS. Glutamate (100 µM with the co-agonist glycine 10 µM promoting activation of NMDA receptors subtype) or GABA (100 µM) were applied for 2 s. Finally, KCl (50 mM) application for 2 s was used to distinguish exciTable neurons from possible non-neuronal cells. Fluorescence was detected with the Till Photonics imaging system (FEI GmbH) equipped with a 12-bit CCD Camera (SensiCam) with a light excitation wavelength of 494 nm. Calcium responses to neurotransmitters were evaluated from changes in fluorescence intensity of individual neurons. To this end, regions of interest (ROI) of round shape around the cell body were selected from the whole image with the TILL vision Imaging Software (TILL Photonics GmbH). To distinguish from non-neuronal cells, ROI was taken at the time point corresponding to KCl-induced activation of neurons. The intensity values in each ROI were averaged at each time point to form a fluorescence signal for each neuron and normalized to the baseline level. Signals above 5% of the baseline were included in the analysis.

N2a Cell culture

Immortalized murine neuroblastoma cells (N2a) were maintained in DMEM, high glucose, GlutaMAX™ Supplement (Gibco, 31966021) media supplemented with supplemented with 10% Fetal Bovine Serum (Gibco, 10270106) and 1% penicillin and streptomycin in temperature controlled humidified incubator (37 °C, 5%CO_2). As previously published[49], for differentiation induction 100.000 cells were plated in a 6-well plate and maintained in complete media supplemented with 20μM retinoic acid (Sigma-Aldrich, R2625) in dimethyl sulfoxide (DMSO) with or without serum starvation (2% FBS instead of 10%) changing half of the media every second day for 5 days before collection.

MOUSE STRAINS AND ANIMAL PROCEDURES

All experiments follow the Helsinki Declaration and guidelines set by the European Commission (European Communities Council Directive, 86/609/EEC) and were approved by the National Animal Experiment Board of Finland.

Animals were housed with same sex siblings, in controlled temperature, humidity and light (12 hours light/dark cycle) conditions. Animals had access to ad libitum food. Before the beginning of the animal study mice were divided in single cages. Mice were randomized and all the participants were blinded to the genotype (surgery, MRI acquisition, behavioral test, sample collection and data analysis). Prior to the surgery, we use random number generator (GraphPad Prism quick calcs https://www.graphpad.com/quickcalcs/randomN1/) to randomize the mice into treatment groups. This numbering has been used in crescent order on 3-4 months old male mice that were divided into groups. pMCAo study was performed on 3-4 months old Balb/cOlaHsd male mice (n = 6-8 per timepoint). Cdr1as (ciRS-7) KO tMCAo study was performed on 3-4 months old C57BL/6N-Cdr1asem1Nikr[9] (Cdr1as KO mice and the WT counterpart), provided by Prof. Dr. Nikolaus Rajewsky, MDC, Berlin, Germany (n = 22-24 per genotype). miR-7 inducible KO tMCAo study was

performed on 3-4 months old B6.Cg-miR7a1tm1(fl/fl)ms miR7a2tm1(fl/fl)ms miR7btm1(fl/fl)ms

Ndor1Tg(UBC-cre/ERT2)1Ejb/Stf and UBC-cre negative counterpart[39] provided by Prof. Dr.

Markus Stoffel, ETH, Zurich, Switzerland (n = 20-26 per genotype).

Genotyping

Before the animal studies, ciRS-7 KO and miR-7 KO mice were genotyped as before[9,39] except for

the tissue dissociation step. Briefly, ear puncture samples were digested in 50 mM Sodium Hydroxide

solution for one hour at 95 °C. Once equilibrated to room temperature the samples were neutralized

with 1M Tris Hydrochloride pH 8. Two microliters of digested samples were used to run the reaction

of PCR with Taq 2X Master Mix (New England Biolabs (UK) Ltd) following manufacturer protocol

and using the oligonucleotide primers as specified in Supplementary Table S8. The PCR products

were separated on a stained 2.5% or 1.5% Agarose gel for ciRS-7 and miR-7 KO respectively and

run in an electrophoresis chamber. Images were acquired with ChemiDoc Imaging Systems (Bio-Rad

Laboratories, Inc).

Permanent middle cerebral artery surgery (pMCAo)

Permanent middle cerebral artery occlusion (pMCAo) was performed in 3-4 months old

Balb/cOlaHsd male mice as previously described[50]. Briefly, mice were anesthetized using 5%

isoflurane and anesthesia was maintained with 2% isoflurane. Temperature of the animals was

controlled during the surgery with heating blanket and rectal probe (Harvard apparatus). After the

skin incision, the left temporal bone was exposed under the temporal muscle, and 1 mm hole was

drilled on top of bifurcation in the middle cerebral artery (MCA). The dura was gently removed, and

the MCA was lifted and permanently occluded under the bifurcation using a thermocoagulator (Aaron

Medical Industries Inc.). Success of the occlusion was confirmed by cutting the MCA above the

occlusion site. After the occlusion, the temporal muscle was replaced, and the wound was sutured.

The surgery was performed on a total of 20 mice, 3 mice (one per timepoint) died during the surgery

and was then not included.

miR-7 KO induction with tamoxifen

Before tMCAo surgery, miR-7 inducible KO mice were injected with 2mg of Tamoxifen

intraperitoneally once per day for 5 consecutive days at the age of 3 to 4 weeks as before[39].

Recombination was induced in both Cre+ and Cre- littermates. Tamoxifen (Sigma-Aldrich, T5648)

was resuspended in 90% corn oil (Sigma-Aldrich, C8267) and 10% pure ethanol and dissolved at

56°C for 30 minutes in the dark. After injection the mice were put on soft food diet and their weight

was monitored daily. Mice showing a successful genomic recombination in ear puncture samples two

weeks after injection were included in the study (Supplementary Figure S4A). At the end of the study

the animals were analyzed for miR-7 abrogation in the brain through RT-qPCR (Supplementary

Figure S4B).

Transient middle cerebral artery surgery (tMCAo)

Transient middle cerebral artery occlusion (tMCAo) was performed in 3-4 months old male Cdr1as

and miR-7 KO and WT mice as previously described[51]. Briefly, induction of anesthesia was

performed with 3.5-4% isoflurane in 0.5 L/min of 100% O2-enriched air and maintained at 1-1.5%

isoflurane during the surgery. The body temperature maintained at 36 ± 1°C during surgery with a

heating pad attached to a rectal probe. A Doppler probe was fixed on the temporal bone over the

MCA to monitor blood flow. The right common, external, and internal carotid arteries (CCA, ECA,

and ICA) were exposed through an incision in the neck. The arteries were carefully dissected, and a

6-0 silicon coated monofilament of 0.21 mm of diameter (Doccol Corporation) was inserted into the

right CCA or left ECA (Cdr1as and mir-7 KO respectively) and led through the ICA until blocking

the origin of the MCA. Blood flow blockage was monitored with the Doppler probe, and animals

with less than 80% of blood flow decrease were discarded. After one hour of occlusion, the

monofilament was withdrawn allowing blood reperfusion. Animals that did not show a correct

reperfusion measured by laser doppler were discarded from the study. Incisions were permanently

sutured, and the animals were allowed to recover in a temperature-controlled environment for 24

hours. The surgery was performed on a total of 32 WT and 25 Cdr1as KO mice, and 36 WT and 26

miR-7 KO. In the ciRS-7 animal study a total of 7 mice was excluded as: 2 KO and 2 WT mice lacked

reperfusion upon surgery, 2 WT and 1 KO mice died during the study. In the miR-7 animal study a

total of 12 mice was excluded as: 3 WT and 2 KO lacked reperfusion upon surgery, 4 WT and 3 KO

died during the study.

Magnetic resonance imaging (MRI)

MRI was performed at 1, 3, and 7dpi using a vertical 9.4 T/89 mm magnet (Oxford instrument

PLC) upon anesthesia with 1.8% isoflurane in 30% O2/70% N2O. We acquired twelve slices of

0.8mm thickness per mouse (echo time 40ms, repetition time of 3000ms, matrix size of 128×256

and field of view $19.2 \times 19.2 \text{ mm}^2$) and analyzed the first seven images using the Aedes software

(http://aedes.uef.fi/) for MatLab program (Math-works). Upon definition of the region of interest

(ROI) of contralateral, ipsilateral, and lesion, the lesion volume normalized on oedema was

calculated on the first 7 section as: $Corrected\ lesion\ size\ =\ lesion\ size\ \times\ \Big(1 -$

$\frac{[ipsilateral\ size\ -\ contralateral\ size]}{contralateral\ size}\Big)$. The lesion size is expressed has percentage of lesion on the total

brain size.

Neurological severity score (NSS)

Mice were examined for neurological deficits at baseline and dpi 1, 3, and 7 using a severity scale

comprising the following tests: postural reflex, circling, falling to contralateral side, placement of the

contralateral forelimb during motion, and general state of alertness or consciousness. Deficits were

graded from 0 (normal) to 2 or 3 (severe)[52]. A sum of these scores were used for statistics. Behavioral

assessment was performed by an experimenter blinded to the genotype of mice.

Transcardiac perfusion and sample collection

After anesthesia, mice were perfused transcardially with cold saline solution with heparin 2500 IU/l

(Leo Pharma A/S). In the pMCAo study, the brains were collected and dissected into contralateral

and peri-ischemic cortical regions. In the tMCAo study the brains were cut in 6 coronal sections and

stained with 1% 2,3,5-Triphenyltetrazolium Chloride (TTC) (Sigma-Aldrich) in PBS solution for 5

minutes at 37 °C in agitation before dissection of contralateral and peri-ischemic cortical regions. For

immunohistochemistry staining, brains were collected and fixed in 4% paraformaldehyde solution in

0.1 M phosphate buffer (PB) pH 7.4. After 22 hours of fixation, the brains were transferred in 30%

sucrose in PB buffer solution for 48 hours and then frozen in liquid nitrogen before being stored in -

70 °C.

GFAP immunostaining

Each brain was then cut using a cryostat (Leica Microsystems) into six 20 μm coronal sections 400 μm

apart, collected on Superfrost™ Plus Microscope Slides (ThermoFisher Scientific) and stored in -

70°C until immunostaining. Sections were then rehydrated with phosphate-buffered saline (PBS) pH

7.4 for 10 min and PBS with 0.05% Tween-20 (PBST) (Sigma-Aldrich) for 5 min. Endogenous

peroxidase was blocked by using 0.3 % hydrogen peroxide (H_2O_2) in MeOH for 30 min after which

sections were washed 3 x 5 min in PBST. Non-specific binding was blocked with 10 % normal goat

serum (NGS)(Vector, S-1000) in PBST for 1 h at RT Sections were incubated overnight at RT in

primary antibody rabbit anti-GFAP (Agilent, Dako Z0334,1:500 in 5 % NGS-PBST) and then with

biotinylated secondary antibody anti-rabbit IgG (H+L) (Vector, BA-1000, 1:200 in 5 % NGS-PBST)

for 2 h at RT followed by incubation in ABC reagent (Vector Elite Kit) for 2 h at RT. Sections were

washed 3 x 5 min in PBST before and after the incubations. Nickel-3,3'-diaminobenzidine (Ni-DAB)

solution (0.175M Sodium acetate, 1% Nickel ammonium sulphate, 50mg DAB (Sigma-Aldrich, D-

5905)) with 0.075 % H_2O_2 was used to develop the colour for 6 min stopping the reaction by washing

the sections 2 x 5 min in dH_2O. Sections were then dehydrated in 50 % EtOH, 70 % EtOH, 95 %

EtOH, 100 % EtOH for 2 min in each and 3 x 5 min in xylene followed by mounting the coverslips

with Depex.

Image acquisition & analysis

Six GFAP immunoreactivity light microscope images per mouse were acquired by Leica DM6B-Z

Thunder Imager microscope (Leica Microsystems CMS GmbH) equipped with DMC2900 camera

using 10x magnification. Images were captured using LAS X software (Leica Microsystems CMS

GmbH) with exposure time 5 ms, color-gain mode: R:0-G:0-B:25. The images were quantified from

the peri-ischemic cortex next to the lesion border (1mm) and the corresponding area of the healthy

contralateral hemisphere from 10x images using ImageJ software[53] function "Measure particles". The

results were presented as relative immunoreactive area to the total area analyzed.

RNA-seq data analysis

Bulk RNA sequences of mRNA and miRNA have been aligned and quantified to the mouse genome

of reference mm10 with the Cdr1as annotation of "chrX:61183248|61186174|.|+" using the nf-core

workflow[54] (3.0 version of "rnaseq" and 1.1.0 version of "smrnaseq" applied without clipping and

the three prime adapter "AGATCGGAAGAGCACACGTCT"), while circular sequences using 1.1

CIRIquant[55] set to Read 1 matching the sense strand. Count data were prepared following the

workflow defined by Law et al[56]. We filtered out lowly expressed molecules in any condition using

the function "filterByExpr" to increase the reliability of the mean-variance relationship. We removed

the differences between samples due to the sequencing depth normalizing the count using the trimmed

mean of M-values (TMM)[57] method and applied a log transformation minimizing sum of sample-

specific squared difference to enhance the true positive and negative ratio in the downstream

analysis[58]. We checked for batch effect due to different timings in biological replicates preparation

by performing a principal component analysis and unsupervised consensus clustering with Cola[59].

We identified a batch effect due to different timings in biological replicates preparation influencing

the grouping of the normoxic and OGD samples in the dataset ciRS-7 KO/WT cortical neurons

(Supplementary Table S5). We corrected the batch effect of this dataset using the negative binomial

regression from Combat[60]. We adjusted the variance between the samples as before[61] through

winsorization[62]. We finally created the design matrix for each pair of conditions to compare (contrast)

and performed the differential expression analysis using limma/edgeR model[56] controlling for the

false discovery rate with Benjamini-Hochberg Procedure[63].

Ingenuity Pathway Analysis (IPA) and Functional enrichment analysis

We uploaded the differentially expressed genes of each contrast to QIAGEN IPA (QIAGEN Inc.,

https://digitalinsights.qiagen.com/IPA)[64] and Metascape[65] for Ingenuity pathway analysis and

functional enrichment analysis, respectively. The analysis has been performed with default

parameters and IPA's background was composed of non-differentially expressed genes.

Deconvolution analysis with scRNAseq

We exploited cell-type specific gene expression from external single-cell RNA sequencing (scRNA-

seq) data to define the cell subpopulations composing our cortical neuron culture (Supplementary

Table 2) and ischemic stroke animal tissue (Supplementary Table 6) bulk RNA sequencing datasets.

The scRNA-seq dataset of Loo et al.[66] of the mouse cerebral cortex at embryonic day 14.5 was

provided to SCDC[67] for performing the deconvolution of our gene expression matrix with cortical

neuron samples in normoxic conditions. The scRNA-seq data of Zeisel at al.[68] composed of murine

cerebral cortex samples from p25 to p60 was instead provided to MuSiC[69] to perform   the

deconvolution of our gene expression matrix with contralateral and peri-ischemic samples of wild-

type and ciRS-7 KO ischemic stroke animals (Supplementary Table 6) as this method is designed to

work with multi-subject scRNA-seq dataset.

GRO-seq data analysis

A summary of all GRO-seq samples used to quantify pri-miRNA expression levels are presented in

Supplementary Table S3. Raw reads for public GRO-seq data were acquired from the GEO database.

GRO-Seq reads were trimmed using the HOMER v4.3 (http://homer.salk.edu/homer)[70] software to

remove A-stretches originating from the library preparation. From the resulting sequences, those

shorter than 25 bp were discarded. The quality of raw sequencing reads was controlled using the

FastQC tool (http://www.bioinformatics.babraham.ac.uk/projects/fastqc)[71] and bases with poor

quality scores were trimmed using the FastX toolkit (http://hannonlab.cshl.edu/fastx_toolkit/). Reads

were aligned to mouse mm9 reference genome using the Bowtie[72] version bowtie-0.12.7. Up to two

mismatches and up to three locations were accepted per read and the best alignment was reported.

The data was used to create Tag Directories using HOMER. To optimize coverage, a combined tag

directory representing all samples of a given cell type (under one or several GSE numbers) was

created and used for pri-miRNA quantification using the 'analyzeRepeats.pl' command and '-rpkm -

strand +' options". The dataset used in this analysis are reported in Supplementary Table S3.

miRWalk analysis

We associated the significantly deregulated miRNAs and differentially expressed genes of wild-type

and ciRS-7 KO cortical neurons subjected to OGD (Supplementary Table 2, Supplementary Table

4). We filtered miRNA-Target interactions obtained from 3.0 miRWalk[73,74] database with 99%

probability and located in the 3-UTR region and included Oip5os1 (Cyrano) as target of mmu-miR-

7a-5p[8]. We linked the differentially expressed miRNAs and genes of each contrast by anticorrelation

(e.g. miRNA with positive log fold change and significant adjusted probability value is associated to

genes with negative log fold change, significant adjusted probability value and targets in miRWalk

of the miRNA) focusing on mmu-miR-7a-5p.

CDF generation

We tested the assumption of anticorrelation between mmu-miR-7a-5p and its miRWalk gene targets

using the empirical Cumulative Distribution Function (eCDF). For each contrast, we compared the

eCDF (control function) obtained from the values of log fold change of the non-target genes against

the one of the miRNA's targets and tested their equality with the Kolmogorov–Smirnov test[75]. To

determine the magnitude and direction of the shift of the eCDF of the targets in respect of the non-

targets, we measured the area between the two curves following this formula: $\int \{F_Y(t) - F_X(t)\}\, dt$

and the Wasserstein distance[76]. Both the considered targets and non-targets genes passed the filtering

by expression, count normalization and participated at the differential expression analysis.

CLIP-seq analysis

We collected the HITS-CLIP dataset of Argonate 2 in pyramidal excitatory neurons produced by Tan

et al.[40] (Supplementary Table S9) and replicated the original analysis to map and annotate the

genomic regions with a significant read cluster (peak). Briefly, we performed a quality control with

FastQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc)[71] and applied trimming to

remove bases after 50 base pairs due to low quality. We applied Cutadapt[77] to remove adapters and

over-represented sequences. We selected reads of at least 24 nucleotides in length and quality score

higher than 20 for each nucleotide. Reads were aligned to the mouse reference genome mm10 using

Burrows-Wheeler aligner[78] with default parameters, then the mapped reads were expanded by 30

nucleotides on each side. We considered only reads present in at least 2 experimental replicates and

clusters composed of at least 10 overlapping reads. Bowtie[79] was used to map all occurrences of

sequences complementary to the 6mer seed sequence (positions 2-7 of the mature miRNA) of mouse

miRNAs from miRBase V21 (https://www.mirbase.org/) to the mouse genome mm10, allowing no

mismatches. Strand specific intersection was done with the filtered clusters in the 3'UTR, 5'UTR,

coding sequence (CDS) and intron regions from RefSeq gene definitions, as well as antisense matches

to RefSeq genes[80]. Strand specific intersection was also done to circRNAs from circBase[81]. This

generated a genome-wide list with predicted miRNA target sites backed by the detected Ago2 HITS-

CLIP clusters for all mouse miRNAs. A subset including all miR-7a-5p targets is supplied as

Supplementary Table S7. Only 3'UTR target sites were used to define miRNA mediated regulation.

The full list of all predicted miRNA target sites was uploaded to GEO, as described in the Data

Availability section.

Statistical analysis

Graphs and statistical analysis were performed in GraphPad Prism 9. Every Figure legend reports

parameters of replicates (n), statistical test and p-value obtained. Where not specified, p-value was

not statistically significant (p-value > 0.05). We refer to n in animal study as single biological

replicate (mouse) and in cortical neuron as technical replicates in the same batch, the experiment has

been performed in three independent biological replicates.

Data availability

The data produced and analyzed in this publication have been deposited in NCBI's Gene Expression

Omnibus[82] and are accessible through GEO Series accession number GSE213179, GSE213067,

GSE213177, GSE211552 and GSE215210 as presented in the Supplementary Table S9. The code

developed to analyze the data produced in this study is available upon request at
https://doi.org/10.5281/zenodo.7233809.

ACKNOWLEDGEMENTS

The authors acknowledge Nicholas Downes for the fruitful discussions, Mirka Tikkanen for her technical assistance with immunohistochemistry staining, Reetta Vuolteenaho for mouse colony maintenance and Miia Salo for assistance in the preparation of primary neuronal cultures. Moreover, the authors acknowledge the Biocenter Kuopio National Virus Vector Laboratory technology service, Cell and Tissue Imaging Unit of UEF, The Lab Animal Centre at the University of Eastern Finland in particular Nina Liimatainen and Hanna Miettinen, for the support. The Biocenter Oulu Transgenic and Tissue Phenotyping core facility, Oulu Laboratory Animal Centre Research Infrastructure, Translational cell biology core facility at Kontinkangas campus and the Faculty of Biochemistry and Molecular Medicine of the University of Oulu are also acknowledged. This project is co-founded by Horizon 2020 Framework Programme of the European Union (Marie Skłodowska Curie grant agreement No 740264). This study was supported by the Academy of Finland, The Finnish Cultural Foundation, Finnish Foundation for Cardiovascular Research, Instrumentarium Science Foundation, Inkeri and Mauri Vänskä Foundation, Saastamoinen Foundation and Kuopio University Foundation. The illustrations present in this study were created with BioRender.com.

AUTHOR CONTRIBUTIONS

F.S. and T.M. conceived and planned the study; F.S., T.M., T.H., J.J., R. Giniatullin, and D.T. provided intellectual contribution and contributed to the interpretation of the results; F.S., V.S., P.K., I.U., J.J, H.D., C.P., N.V and J. Koistinaho contributed to the *in vivo* experiments; F.S., V.S., D.M.T., M.G.B., R. Giniatullina, N.K., S.YH. contributed to the *in vitro* experiments; F.S., A. H.S., J.S., and J. Kjems planned and performed the libraries for the RNA-seq experiments; L.G. planned and

generated the code for the bioinformatic analysis and analyzed RNA-seq data; M.L., M.P., M.S., N.R.
provided the transgenic animals and assisted in the animal study on these models; M.V. and A.S.
produced and analyzed the HITS-CLIP-seq data; R.H. coordinated and supported the sample
preparation in Oulu University by F.S., V.S., S.K. and A.H; M.K. analyzed the GRO-seq data; F.S.
and T.M. wrote the manuscript in consultation with T.H., J.J. R. Giniatullin, J. Kjems with input from
all authors. All authors provided critical feedback and helped shape the research, analysis and
manuscript.

REFERENCES

1. Mehler, M. F. & Mattick, J. S. Noncoding RNAs and RNA editing in brain development, functional diversification, and neurological disease. *Physiol. Rev.* **87**, 799–823 (2007).

2. Lewis, B. P., Burge, C. B. & Bartel, D. P. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* **120**, 15–20 (2005).

3. Mattick, J. S. & Makunin, I. V. Small regulatory RNAs in mammals. *Hum. Mol. Genet.* **14 Spec No 1**, R121-132 (2005).

4. Fatica, A. & Bozzoni, I. Long non-coding RNAs: new players in cell differentiation and development. *Nat. Rev. Genet.* **15**, 7–21 (2014).

5. Hansen, T. B. *et al.* Natural RNA circles function as efficient microRNA sponges. *Nature* **495**, 384–388 (2013).

6. Kristensen, L. S. *et al.* The biogenesis, biology and characterization of circular RNAs. *Nat. Rev. Genet.* **20**, 675–691 (2019).

7. Hansen, T. B. *et al.* miRNA-dependent gene silencing involving Ago2-mediated cleavage of a circular antisense RNA: miRNA mediated cleavage of circular antisense RNA. *EMBO J.* **30**, 4414–4422 (2011).

8.   Kleaveland, B., Shi, C. Y., Stefano, J. & Bartel, D. P. A Network of Noncoding Regulatory

RNAs Acts in the Mammalian Brain. *Cell* **174**, 350-362.e17 (2018).

9.   Piwecka, M. *et al.* Loss of a mammalian circular RNA locus causes miRNA deregulation and

affects brain function. *Science* **357**, eaam8526 (2017).

10. Memczak, S. *et al.* Circular RNAs are a large class of animal RNAs with regulatory potency.

Nature **495**, 333–338 (2013).

11. Olney, J. W., Rhee, V. & Ho, O. L. Kainic acid: a powerful neurotoxic analogue of glutamate.

Brain Res. **77**, 507–512 (1974).

12. Dirnagl, U., Iadecola, C. & Moskowitz, M. A. Pathobiology of ischaemic stroke: an integrated

view. *Trends Neurosci.* **22**, 391–397 (1999).

13. Tiedt, S. & Dichgans, M. Role of Non-Coding RNAs in Stroke. *Stroke* **49**, 3098–3106 (2018).

14. Leung, A. K. L. & Sharp, P. A. MicroRNA Functions in Stress Responses. *Mol. Cell* **40**, 205–

215 (2010).

15. Denzler, R. *et al.* Impact of MicroRNA Levels, Target-Site Complementarity, and

Cooperativity on Competing Endogenous RNA-Regulated Gene Expression. *Mol. Cell* **64**, 565–

579 (2016).

16. Salmena, L., Poliseno, L., Tay, Y., Kats, L. & Pandolfi, P. P. A ceRNA hypothesis: the Rosetta

Stone of a hidden RNA language? *Cell* **146**, 353–358 (2011).

17. Franco-Zorrilla, J. M. *et al.* Target mimicry provides a new mechanism for regulation of

microRNA activity. *Nat. Genet.* **39**, 1033–1037 (2007).

18. Zhang, Y. *et al.* Circular RNAs in the Regulation of Oxidative Stress. *Front. Pharmacol.* **12**,

(2021).

19. Loo, L. *et al.* Single-cell transcriptomic analysis of mouse neocortical development. *Nat.*

Commun. **10**, 134 (2019).

20. Semenza, G. L. Hypoxia-inducible factor 1: control of oxygen homeostasis in health and

disease. *Pediatr. Res.* **49**, 614–617 (2001).

21. Zhang, J., Chen, S., Yang, J. & Zhao, F. Accurate quantification of circular RNAs identifies

extensive circular isoform switching events. *Nat. Commun.* **11**, 90 (2020).

22. Loppi, S. *et al.* Peripheral inflammation preceeding ischemia impairs neuronal survival through

mechanisms involving miR-127 in aged animals. *Aging Cell* **20**, e13287 (2021).

23. Step, S. E. *et al.* Anti-diabetic rosiglitazone remodels the adipocyte transcriptome by

redistributing transcription to PPARγ-driven enhancers. *Genes Dev.* **28**, 1018–1028 (2014).

24. Meng, F.-L. *et al.* Convergent transcription at intragenic super-enhancers targets AID-initiated

genomic instability. *Cell* **159**, 1538–1548 (2014).

25. Min, I. M. *et al.* Regulating RNA polymerase pausing and transcription elongation in

embryonic stem cells. *Genes Dev.* **25**, 742 (2011).

26. Fang, B. *et al.* Circadian enhancers coordinate multiple phases of rhythmic gene transcription in

vivo. *Cell* **159**, 1140–1152 (2014).

27. Hah, N. *et al.* Inflammation-sensitive super enhancers form domains of coordinately regulated

enhancer RNAs. *Proc. Natl. Acad. Sci.* **112**, E297–E302 (2015).

28. Heinz, S. *et al.* Effect of natural genetic variation on enhancer selection and function. *Nature*

503, 487–492 (2013).

29. Escoubet-Lozach, L. *et al.* Mechanisms Establishing TLR4-Responsive Activation States of

Inflammatory Response Genes. *PLOS Genet.* **7**, e1002401 (2011).

30. Ji, Z. *et al.* Transcriptional activity regulates alternative cleavage and polyadenylation. *Mol.*

Syst. Biol. **7**, 534 (2011).

31. Zhang, F. *et al.* Enhancer-bound LDB1 regulates a corticotrope promoter-pausing repression

program. *Proc. Natl. Acad. Sci.* **112**, 1380–1385 (2015).

32. Telese, F. *et al.* LRP8-Reelin-regulated Neuronal (LRN) Enhancer Signature Underlying

Learning and Memory Formation. *Neuron* **86**, 696–710 (2015).

33. Sticht, C., Torre, C. D. L., Parveen, A. & Gretz, N. miRWalk: An online resource for prediction

of microRNA binding sites. *PLOS ONE* **13**, e0206239 (2018).

34. Dweep, H. & Gretz, N. miRWalk2.0: a comprehensive atlas of microRNA-target interactions.

Nat. Methods **12**, 697–697 (2015).

35. Zeisel, A. *et al.* Molecular Architecture of the Mouse Nervous System. *Cell* **174**, 999-1014.e22

(2018).

36. Yoon, J. S. *et al.* Spatiotemporal Protein Atlas of Cell Death-Related Molecules in the Rat

MCAO Stroke Model. *Exp. Neurobiol.* **27**, 287–298 (2018).

37. Kabaria, S. *et al.* MicroRNA-7 activates Nrf2 pathway by targeting Keap1 expression. *Free*

Radic. Biol. Med. **89**, 548–556 (2015).

38. Chaudhuri, A. D., Kabaria, S., Choi, D. C., Mouradian, M. M. & Junn, E. MicroRNA-7

Promotes Glycolysis to Protect against 1-Methyl-4-phenylpyridinium-induced Cell Death. *J.*

Biol. Chem. **290**, 12425–12434 (2015).

39. LaPierre, M. P., Lawler, K., Godbersen, S., Farooqi, I. S. & Stoffel, M. MicroRNA-7 regulates

melanocortin circuits involved in mammalian energy homeostasis. *Nat. Commun.* **13**, 5733

(2022).

40. Tan, C. L. *et al.* MicroRNA-128 Governs Neuronal Excitability and Motor Behavior in Mice.

Science **342**, 1254–1258 (2013).

41. Chen, H. *et al.* miR-7 and miR-214 are specifically expressed during neuroblastoma

differentiation, cortical development and embryonic stem cells differentiation, and control

neurite outgrowth in vitro. *Biochem. Biophys. Res. Commun.* **394**, 921–927 (2010).

42. Rybak-Wolf, A. *et al.* Circular RNAs in the Mammalian Brain Are Highly Abundant,

Conserved, and Dynamically Expressed. *Mol. Cell* **58**, 870–885 (2015).

43. Yao, G.-Y. *et al.* Ischemic postconditioning confers cerebroprotection by stabilizing VDACs

after brain ischemia. *Cell Death Dis.* **9**, 1–15 (2018).

44. Zhao, J. & Wang, B. MiR-7-5p Enhances Cerebral Ischemia-Reperfusion Injury by Degrading

sirt1 mRNA. *J. Cardiovasc. Pharmacol.* **76**, 227–236 (2020).

45. Mehta, S. L. *et al.* CDR1as regulates α-synuclein-mediated ischemic brain damage by

controlling miR-7 availability. *Mol. Ther. Nucleic Acids* **31**, 57–67 (2023).

46. Bushati, N. & Cohen, S. M. microRNA functions. *Annu. Rev. Cell Dev. Biol.* **23**, 175–205

(2007).

47. Miska, E. A. *et al.* Most Caenorhabditis elegans microRNAs are individually not essential for

development or viability. *PLoS Genet.* **3**, e215 (2007).

48. Tiihonen, J. *et al.* Sex-specific transcriptional and proteomic signatures in schizophrenia. *Nat.*

Commun. **10**, 3933 (2019).

49. Kumar, M. & Katyal, A. Data on retinoic acid and reduced serum concentration induced

differentiation of Neuro-2a neuroblastoma cells. *Data Brief* **21**, 2435–2440 (2018).

50. Korhonen, P. *et al.* Immunomodulation by interleukin-33 is protective in stroke through

modulation of inflammation. *Brain. Behav. Immun.* **49**, 322–336 (2015).

51. Shahjouei, S. *et al.* Middle Cerebral Artery Occlusion Model of Stroke in Rodents: A Step-by-

Step Approach. *J. Vasc. Interv. Neurol.* **8**, 1–8 (2016).

52. Senda, D. M., Franzin, S., Mori, M. A., de Oliveira, R. M. W. & Milani, H. Acute, post-

ischemic sensorimotor deficits correlate positively with infarct size but fail to predict its

occurrence and magnitude after middle cerebral artery occlusion in rats. *Behav. Brain Res.* **216**,

29–35 (2011).

53. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image

analysis. *Nat. Methods* **9**, 671–675 (2012).

54. Ewels, P. A. *et al.* The nf-core framework for community-curated bioinformatics pipelines. *Nat.*

Biotechnol. **38**, 276–278 (2020).

55. Zhang, J., Chen, S., Yang, J. & Zhao, F. Accurate quantification of circular RNAs identifies

extensive circular isoform switching events. *Nat. Commun.* **11**, 90 (2020).

56. Law, C. W. *et al.* RNA-seq analysis is easy as 1-2-3 with limma, Glimma and edgeR.

F1000Research **5**, (2016).

57. Robinson, M. D. & Oshlack, A. A scaling normalization method for differential expression

analysis of RNA-seq data. *Genome Biol.* **11**, R25 (2010).

58. Zhang, Z. *et al.* Novel Data Transformations for RNA-seq Differential Expression Analysis.

Sci. Rep. **9**, 4820 (2019).

59. Gu, Z., Schlesner, M. & Hübschmann, D. cola: an R/Bioconductor package for consensus

partitioning through a general framework. *Nucleic Acids Res.* **49**, e15 (2021).

60. Zhang, Y., Parmigiani, G. & Johnson, W. E. ComBat-seq: batch effect adjustment for RNA-seq

count data. *NAR Genomics Bioinforma.* **2**, lqaa078 (2020).

61. Allswede, D. M. *et al.* Complement Gene Expression Correlates with Superior Frontal Cortical

Thickness in Humans. *Neuropsychopharmacology* **43**, 525–533 (2018).

62. Winsor, C. P., Hastings, C., Mosteller, F. & Tukey, J. W. Low Moments for Small Samples: A

Comparative Study of Order Statistics. *Ann. Math. Stat.* (1947) doi:10.1214/aoms/1177730388.

63. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B Methodol.* **57**, 289–300 (1995).

64. Krämer, A., Green, J., Pollard, J. & Tugendreich, S. Causal analysis approaches in Ingenuity

Pathway Analysis. *Bioinforma. Oxf. Engl.* **30**, 523–530 (2014).

65. Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of systems-

level datasets. *Nat. Commun.* **10**, 1523 (2019).

66. Loo, L. *et al.* Single-cell transcriptomic analysis of mouse neocortical development. *Nat.*

Commun. **10**, 134 (2019).

67. Dong, M. *et al.* SCDC: bulk gene expression deconvolution by multiple single-cell RNA

sequencing references. *Brief. Bioinform.* **22**, 416–427 (2021).

68. Zeisel, A. *et al.* Molecular Architecture of the Mouse Nervous System. *Cell* **174**, 999-1014.e22

(2018).

69. Wang, X., Park, J., Susztak, K., Zhang, N. R. & Li, M. Bulk tissue cell type deconvolution with

multi-subject single-cell expression reference. *Nat. Commun.* **10**, 380 (2019).

70. Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-

regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38**, 576–589

(2010).

71. Blankenberg, D. *et al.* Manipulation of FASTQ data with Galaxy. *Bioinformatics* **26**, 1783

(2010).

72. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25 (2009).

73. Dweep, H. & Gretz, N. miRWalk2.0: a comprehensive atlas of microRNA-target interactions.

Nat. Methods **12**, 697–697 (2015).

74. Sticht, C., Torre, C. D. L., Parveen, A. & Gretz, N. miRWalk: An online resource for prediction

of microRNA binding sites. *PLOS ONE* **13**, e0206239 (2018).

75. Smirnov, N. Table for Estimating the Goodness of Fit of Empirical Distributions. *Ann. Math.*

Stat. **19**, 279–281 (1948).

76. Kantorovich, L. V. Mathematical Methods of Organizing and Planning Production. *Manag. Sci.*

6, 366–422 (1960).

77. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal **17**, 10–12 (2011).

78. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform.

Bioinformatics **25**, 1754–1760 (2009).

79. Trapnell, C., Pachter, L. & Salzberg, S. L. TopHat: discovering splice junctions with RNA-Seq.

Bioinforma. Oxf. Engl. **25**, 1105–1111 (2009).

80. Kd, P., T, T. & Dr, M. NCBI Reference Sequence (RefSeq): a curated non-redundant sequence

database of genomes, transcripts and proteins. *Nucleic Acids Res.* **33**, (2005).

81. Glažar, P., Papavasileiou, P. & Rajewsky, N. circBase: a database for circular RNAs. *RNA N. Y.*

N **20**, 1666–1670 (2014).

82. Edgar, R., Domrachev, M. & Lash, A. E. Gene Expression Omnibus: NCBI gene expression

and hybridization array data repository. *Nucleic Acids Res.* **30**, 207–210 (2002).

FIGURES

Figure 1. ciRS-7 is the most abundant circRNA in mouse cortical neuron cultures and downregulated by in vitro OGD.

(A) Box plot of bulk RNA-seq deconvolution scores. Cell types are sorted based on the median of the universal semantic groups which belong to (e.g. Int1, Int2 in Interneurons). (B) Calcium imaging analysis of the cellular response to glutamate and GABA with representative fluorescence images (n = 11 biological replicates; signal included > 5% of baseline; data as mean ± SD). (C) Bar plot of

Ingenuity pathway analysis (IPA) conducted on bulk RNA-seq data of cortical neurons treated with OGD. Z-score indicates IPA prediction of transcription factor activation (positive value) or inhibition (negative value). Green color highlights HIF1α activation ($|z| > 2$ is considered significant). **(D)** Schematic representation of CIRIquant algorithm. Gene composed by three exons (dark blue, light blue, purple) generating a circular molecule through back-splicing or a linear transcript. In green specific circular reads on the back-splice junction (BSJ), in gray reads generated from forward splicing junctions (FSJ) in the same back-splicing region. Junction ratio score formula calculates a circularization score, higher the score and higher percentage of circular form of the gene of interest. **(E)** Heat map of logCPM of detected circularRNAs between normoxic (N) and oxygen and glucose deprived (OGD) murine cortical neuron culture. Blue color represents low expression and purple color represents high expression. Significant differentially expressed circularRNA grouped with a bar (* = adj. p-value < 0.05, n = 5 biological replicates). **(F)** (*left*) bar plot of CPM of the BSJ and FSJ of differentially expressed circularRNAs between normoxic and oxygen and glucose deprivation conditions, BSJ represents circular specific reads. (*right*) bar plot of the Junction ratio score calculated by CIRIquant algorithm. The score is a ratio of BSJ reads and FSJ reads mapped to BSJ site and represents percentage of circularization in respect of the linear transcript.

1043
1044

Figure 2. ciRS-7 network is dynamically altered in ischemic stroke like conditions.

(A) Schematic illustration of the relationships between the molecules of the regulatory network composed by ciRS-7, miR-7, Cyrano and miR-671. (1) Hansen et al. EMBO J, 2011; Hansen et al. Nature, 2013; (2) Piwecka et al. Science, 2017; (3) Kleaveland et al. Cell, 2018. **(B)** Volcano plot of differentially expressed microRNAs in cortical neuron cultures treated with OGD. Highlighted significantly upregulated (pink) and downregulated (green) microRNAs, miR-7 variants (red), miR-7 passenger strands (white), miR-671 (black) (n = 5 biological replicates; p-adj. <0.05 and |log2FC|> 0.3). **(C)** Schematic illustration of the murine genomic loci from which miR-7 primary transcripts

(pri-miR) are transcribed, processed in precursor molecules (pre-miR) and finally into mature miR-7. Highlighted in red the seed sequence shared between miR-7a-5p and miR-7b-5p, underlined in black the non-seed mismatch in position 10 between the variants. **(D)** Bar plot of reads per kilobase million (RPKM) of miR-7 genomic loci in different cell type obtained from GRO-seq datasets. **(E)** Bar plot of the expression of miR-7 pri-mirna in cortical neuron after OGD treatment quantified by quantitative PCR. Relative expression normalized on normoxic condition (n = 4; * = p-value < 0.05, ** = p-value < 0.01 calculated with paired t-test; data as mean ± SD). **(F)** Plot of the cumulative distribution functions (CDFs) of the log fold changes of genes compared between OGD treated wild-type cortical neurons for 12h and normoxic conditions. Red curve is the CDF obtained with only miR-7 targets, while black curve with non-targets (n = 5 biological replicates; p-value calculated with Kolmogorov-Smirnov test). **(G)** Volcano plot of differentially expressed transcripts in cortical neuron cultures treated with OGD. Highlighted significantly upregulated (pink) and downregulated (green) transcripts, Cyrano (purple), miR-7 hosting gene HnrnpK (black) (n = 5 biological replicates; p-adj. <0.05 and |log2FC|> 0.3). **(H)** Bar plot of the quantification by quantitative PCR of ciRS-7 (blue), miR-7 (magenta), Cyrano (purple) in cortical neurons OGD treated at different timepoints (1 hour, 6 hours, 12 hours and 24 hours). Relative expression normalized on normoxic condition (normoxia n = 2-4, OGD n=5-6; * = p-value < 0.05, *** = p-value < 0.001, **** = p-value < 0.0001 calculated with unpaired t-test; data as mean ± SD).

Figure 3. ciRS-7 knock-out genotype exhibit differential OGD response without affecting the OGD induced changes in miR-7 and Cyrano.

(A) (*left*) Representative trace of calcium induced fluorescence in response to glutamate treatment in wild-type and ciRS-7 KO cortical neurons. (*right*) Scatter dot plot of calcium imaging analysis of the cellular response to glutamate of wild-type and ciRS-7 KO cortical neurons. (n = 8 biological replicates; signal included > 5% of baseline; * = p-value < 0.05 calculated with unpaired t-test; central

bar represents median value). **(B)** Bar plot of relative absorbance in MTT viability assay of wild-type

and ciRS-7 KO cortical neurons treated with vehicle or glutamate 250μM. Data expressed as survival

% normalized on wild-type vehicle (n = 4 biological replicates; * = p-value < 0.05, **** = p-value <

0.0001 calculated with one-way ANOVA corrected with Tukey's post-hoc test; data as mean ± SD).

(C) Venn diagram of the differentially expressed genes in normoxic versus OGD condition of wild-

type and ciRS-7 KO cortical neurons (n = 5 biological replicates; p-adj. <0.05 and |log2FC|> 0.3).

(D) Bar plot of functional enrichment analysis top 20 significant Metascape clusters performed on

differentially expressed genes in wild-type normoxic versus OGD cortical neurons. **(E)** Bar plot of

functional enrichment analysis top 20 significant Metascape clusters performed on differentially

expressed genes in ciRS-7 KO normoxic versus OGD cortical neurons. **(F)** (*left*) Bar plot of log2FC

of ciRS-7 (blue), miR-7 (magenta) and Cyrano (purple) obtained from RNA-seq of wild-type and

ciRS-7 KO cortical neurons in normoxic conditions versus OGD. (*right*) Bar plot of log2FC of miR-

7 (magenta) obtained from RNA-seq showing differences in the expression in normoxic conditions

(N) or OGD between the wild-type and ciRS-7 KO genotype (n = 5 biological replicates; p-adj. <0.05

and |log2FC|> 0.3). **(G)** Plot of the cumulative distribution functions (CDFs) of the log fold changes

of genes compared between OGD treated ciRS-7 KO cortical neurons for 12h and normoxic

conditions. Red curve is the CDF obtained with only miR-7 targets, while black curve with non-

targets (n = 5 biological replicates; p-value calculated with Kolmogorov-Smirnov test).

1107

Figure 4. Lack of ciRS-7 ameliorates ischemic stroke outcome in vivo.

(A) Bar plot of the quantification by quantitative PCR of ciRS-7 (blue), miR-7 (magenta), Cyrano (purple) in 3-4 months old BALB/c mice subjected to pMCAo surgery. Peri-ischemic (PI) and contralateral (CL) cortices were collected at different timepoints (6 hours, 1 day and 5 days). Relative expression normalized on the average expression of CL (n = 6-7; * = p-value < 0.05, *** = p-value < 0.001, **** = p-value < 0.0001 calculated with paired t-test; data as mean ± SD). **(B)** Schematic illustration of the experimental design: wild-type and ciRS-7 KO mice were subjected to tMCAo

surgery and monitored with neuroscore testing and MRI at 1 day, 3 days and 7 days post-surgery (dpi). Six mice were used at 1dpi to obtain peri-ischemic and contralateral cortices to perform RNA-sequencing. Eight to nine mice per group were used at dpi post ischemia for immunohistochemistry analysis. **(C)** (*left*) Bar plot of MRI quantification of the lesion size of wild-type and ciRS-7 KO mice subjected of tMCAo 1 day post-surgery. Data expressed as lesion percentage on total brain size adjusted for oedema (n = 25-28 per group; ** = p-value < 0.01 calculated with Mann–Whitney test; data as median with 95% confidence interval). (*right*) Representative image of lesion perimeter (blue) changes measured by MRI between wild-type (WT) and ciRS-7 KO animals. **(D)** Scatter dot plot of sensorimotor deficits measured by neuroscore in tMCAo wild-type and ciRS-7 KO mice at 1dpi (data expressed as total neuroscore; n = 22-25 per group; * = p-value < 0.05 calculated with Mann–Whitney test; data as median with 95% confidence interval). **(E)** Bubble plot of Ingenuity Pathway Analysis of differentially expressed genes between wild-type and ciRS-7 KO peri-ischemic cortices obtained through RNA-seq. Z-score indicates IPA prediction of pathway activation (positive value) or inhibition (negative value) in ciRS-7 KO. Color bar expresses p-value significancy (yellow to purple), size expresses the number of genes contained in the pathway ($|z| > 2$ is considered significant, pathway containing >10 genes were included). **(F)** (*left*) Spiderweb plot of wild-type (gray) and ciRS-7 KO (blue) contralateral and peri-ischemic cortices bulk RNA-seq deconvolution scores. (*right*) Bar plot of neurons and astrocyte contribution based on deconvolution scores in wild-type (gray) and ciRS-7 KO (blue) contralateral and peri-ischemic regions. **(G)** (*left*) Bar plot of quantification of GFAP DAB (3,3′-Diaminobenzidine) immunostaining of wild-type and ciRS-7 KO tMCAo mice at 7dpi. (*right*) representative images of the staining highlighting GFAP+ cells. Relative expression normalized on the average expression of contralateral of wild-type mice (n = 8; * = p-value < 0.05, ** = p-value < 0.01, **** = p-value < 0.0001 calculated with one-way ANOVA corrected with Tukey's post-hoc test; data as mean ± SD). **(H)** Bar plot of the quantification by quantitative PCR of ciRS-7 (blue), miR-7 (magenta), Cyrano (purple) in contralateral (CL) and peri-ischemic (PI) cortices of ischemic wild-type and ciRS-7 KO mice at 1dpi. Schematic representation correlating it to MRI lesion size at the same timepoint. Data are normalized on the average expression of contralateral of wild-type mice (n = 5 per group; * = p-value < 0.05, ** = p-value < 0.01 calculated with paired t-test; data as mean ± SD).

1146
1147
Figure 5. Lack of miR-7 exaggerates ischemic stroke outcome regulating glutamatergic

response.

(A) Schematic illustration of Cre-LoxP miR-7 KO animal model. Mice transgenic genome with LoxP

sequences (blue) flanking all the three miR-7 loci (magenta) and tamoxifen (orange) inducible Cre

recombinase (purple). Tamoxifen activation of the Cre recombinase results in miR-7 loci deletion.

Control mice are not transgenic for Cre recombinase. **(B)** (*left*) Bar plot of MRI quantification of the

lesion size of wild-type and miR-7 KO mice subjected of tMCAo one day post-surgery. Data

expressed as lesion percentage on total brain size adjusted for oedema (n = 8-11 per group; * = p-

value < 0.05 calculated with Mann–Whitney test; data as median with 95% confidence interval). (*right*) Representative image of lesion perimeter (blue) changes measured by MRI between wild-type (WT) and miR-7 KO animals. **(C)** Schematic illustration of recombinant mice utilized in to generate Ago2 CLIP-seq data in excitatory neurons. FLAG-Ago2 (blue) is translated only in Cam2Ka+ (red) excitatory neurons in which Cre recombinase (yellow) expression lead to FLAG-Ago2 translation due to the removal of a stop codon (gray). After cross-linking (thunder) FLAG-Ago2-miRNA-target complex is immunoprecipitated with anti-FLAG antibody (purple) and the miRNA-target complex output is sequenced. **(D)** Pie chart of the distribution of miR-7 binding sites obtained by CLIP-seq analysis. The 3779 identified binding sites are divided into protein coding sequences (CDS, purple), 5' untranslated regions (5' UTR, blue), 3' untranslated regions (3' UTR, dark green), intronic sequences (Intron, light green). **(E)** Venn diagram of the miR-7 binding sites in the 3' UTR identified by CLIP-seq in excitatory neurons (645) and the downregulated genes in ciRS-7 KO tMCAo animals peri-ischemic region (3286), in which miR-7 is upregulated. The genes resulting from the overlap of these datasets (276) represent miR-7 physical targets potentially affected in ischemic stroke. **(F)** Bar plot of functional enrichment analysis top 20 significant Metascape clusters performed of the genes identified as miR-7 physical targets potentially affected in ischemic stroke. **(G)** (*left*) Representative trace of calcium induced fluorescence in response to glutamate treatment in wild-type and ciRS-7 KO neurons infected with miR-7 or control lentivirus. (*right*) Scatter dot plot of the calcium imaging analysis quantifying the cellular response to glutamate of wild-type and ciRS-7 KO cortical neurons infected with control (GFP) or miR-7 overexpressing (GFP + miR-7) virus (n = 5 biological replicates; signal included > 5% of baseline; * = p-value < 0.05, **** = p-value < 0.0001 calculated with Kruskal–Wallis test; central bar represent median value). **(H)** (*left*) Bar plot of neurite length quantified through live imaging of wild-type type and ciRS-7 KO cortical neurons infected with control (GFP) or miR-7 overexpressing lentivirus (GFP + miR-7). Data normalized on average neurite length of wild-type GFP infected neurons (n = 3 biological replicates, n = 8 technical replicates; ** = p-value < 0.01 calculated with one-way ANOVA test corrected with Tukey's post-hoc test; data as mean ± SD). (*right*) Representative image of ciRS-7 KO cortical neurons infected with control (GFP) or miR-7 overexpressing virus (GFP + miR-7).