ciRS-7 and miR-7 regulate ischemia induced neuronal death via glutamatergic signaling

Flavia Scoyni[1]*, Valeriia Sitnikova[1]†, Luca Giudice[1]†, Paula Korhonen[1], Davide M Trevisan[2], Ana Hernandez de Sande[3], Mireia Gomez-Budia[1], Raisa Giniatullina[1], Irene F Ugidos[1], Hiramani Dhungana[4], Cristiana Pistono[1], Nea Korvenlaita[1], Nelli-Noora Välimäki[1], Salla M Kangas[5], Anniina E Hiltunen[6], Emma Gribchenko[1], Minna U Kaikkonen-Määttä[1], Jari Koistinaho[1,4], Seppo Ylä-Herttuala[1], Reetta Hinttala[5,6], Morten T Venø[7,8], Junyi Su[8], Markus Stoffel[9], Anne Schaefer[10,11], Nikolaus Rajewsky[12], Jørgen Kjems[8], Mary P LaPierre[9], Monika Piwecka[13], Jukka Jolkkonen[1], Rashid Giniatullin[1], Thomas B Hansen[8, ‡], Malm Tarja[1]*

Affiliations:

[1]A.I.Virtanen Institute for Molecular Sciences, University of Eastern Finland, Kuopio, 70221, Finland

[2]Department of Biosciences and Nutrition, Karolinska Institute, Stockholm, 17177, Sweden

[3]School of Medicine, University of Eastern Finland, Kuopio, 70221, Finland

[4]Neuroscience Center, University of Helsinki, Helsinki, 00290, Finland

[5]Biocenter Oulu, University of Oulu, Oulu, 90014, Finland

[6]Medical Research Center Oulu and Research Unit of Clinical Medicine, University of Oulu and Oulu University Hospital, Oulu, 90014, Finland

[7]Omiics ApS, Aarhus, 8200, Denmark

[8]Interdisciplinary Nanoscience Center, Department of Molecular Biology and Genetics, Aarhus University, Aarhus, 8000, Denmark

[9]Institute of Molecular Health Sciences, ETH Zurich, Zürich, 8093, Switzerland

[10]Departments of Neuroscience and Psychiatry, Icahn School of Medicine at Mount Sinai, New York, 10029-6504, USA

[11]Max Planck Institute, Biology of Ageing, Cologne, 50931, Germany

[12]Systems Biology of Gene Regulatory Elements, Max-Delbrück-Center for Molecular Medicine in the Helmholtz Association (MDC), Berlin Institute for Medical Systems Biology (BIMSB), Berlin, 10115, Germany

[13]Institute of Bioorganic Chemistry, Polish Academy of Sciences, Poznan, 61-704, Poland

*Correspondence: flavia.scoyni@uef.fi and tarja.malm@uef.fi;

† These authors contributed equally; ‡ Present address: Targovax ASA, Lysaker, 1366, Norway

ABSTRACT

Brain functionality relies on finely tuned regulation of gene expression by networks of non-coding RNAs (ncRNAs) such as the one composed by the circular RNA ciRS-7 (also known as CDR1as), the microRNA miR-7 and the long non-coding RNA Cyrano. Here we describe ischemia induced alterations in the ncRNA network both *in vitro* and *in vivo* and in transgenic mice lacking ciRS-7 or miR-7. Our data show that cortical neurons downregulate ciRS-7 and Cyrano and upregulate miR-7 expression upon ischemic insults. Mice lacking ciRS-7 show reduced lesion size and motor impairment, whilst the absence of miR-7 alone leads to an increase in the ischemia induced neuronal death. Moreover, miR-7 levels in pyramidal excitatory neurons regulate dendrite morphology and glutamatergic signaling suggesting a potential molecular link to the *in vivo* phenotype. Our data reveal that ciRS-7 and miR-7 contribute to the outcome of ischemic stroke and shed new light into the pathophysiological roles of intracellular networks of non-coding RNAs in the brain.

INTRODUCTION

The intricate functionality of the brain relies on precisely regulated gene expression, also mediated by non-coding RNAs (ncRNAs), molecules abundant in the brain linked to the increased cognitive complexity in human[1]. MicroRNAs (miRNAs), short (~22 nucleotides) ncRNAs, post-transcriptionally regulate messenger RNA (mRNA) expression by binding short complementary sequences (seed)[2] and triggering mRNA decay or inhibition of translation[3]. Long non-coding RNAs (lncRNAs), RNA molecules longer than 200 nucleotides, regulate gene expression by interacting with other ncRNAs, including miRNAs, or proteins[4]. Circular RNAs (circRNAs), a novel class of ncRNAs, modulate gene expression also by interacting with miRNAs, hence affecting miRNAs activity on mRNA targets[5]. CircRNAs result from backsplicing of linear transcripts, an uncanonical splicing event in which a 5' splice site is spliced with the 3' splice site of the upstream exon[6], forming a circular molecule.

These ncRNAs independently control cellular function by regulating the expression of protein-coding genes, but also interact with each other. In the brain-specific ncRNA network involving miR-7, miR-671, the lncRNA Cyrano and the circRNA ciRS-7 (also known as CDR1as)[5,7–9], ciRS-7 is suggested to stabilize and promote miR-7 targeting[8,9], while Cyrano triggers miR-7 degradation via target RNA–directed miRNA degradation (TDMD)[8] with a nearly perfectly complementary binding site. Analysis through gain and loss of function experiments and knock-out animals revealed that Cyrano and ciRS-7 bind miR-7, regulating its expression[5,8,9] and that Cyrano TDMD effect on miR-7 affects ciRS-7 abundance and localization, indirectly affecting the gene expression of miR-7 targets[5,8–10]. Nonetheless, the physiological purpose of this network in the brain remains unknown, including lack of information on its role in pathophysiological conditions.

Ischemic stroke, induced by occlusion of one of the major cerebral arteries, leads nutrients and oxygen deprivation in the brain parenchyma inducing cell death. Lack of energy and disruption in the ion balance cause an uncontrolled release of glutamate in excitatory neurons, leading to excitotoxicity[11], oxidative stress, necrosis, and apoptosis[12]. Various miRNAs and lncRNAs have been implicated in regulating oxidative stress response and glutamate excitotoxicity, impacting stroke outcomes[13]. Sustained stress prompts miRNAs to facilitate adaptive switches in gene expression program[14]. The efficiency of the miRNA-mediated response depends on miRNAs availability to interact (expression, localization, activity) and the amount of mRNA targets possessing a Mirna Recognition Element (MRE), creating a miRNA-specific threshold[15]. Sudden changes in MRE-containing sequences can disrupt miRNA targeting and derepress specific targets. This crosstalk led to theorize that in physiological conditions a large number of MRE sequences compete for the binding of the miRNA[16]. Organisms exploit this phenomenon during stress by regulating miRNA activity through transcripts with varying binding strength or MRE abundance following a process called target mimicry[17].

Recent studies emphasize a strong connection between circRNAs and reactive oxygen species production[18]. Despite individual ncRNAs being studied in the context of ischemic stroke[13], a comprehensive analysis of the regulatory role of ncRNAs network in stroke is still lacking. In this study, we identified ciRS-7, brain enriched circRNA, ciRS-7, among seven circRNAs deregulated in ischemia-related conditions *in vitro* and *in vivo*. Expanding our analysis to other ncRNAs species, we identified changes in miR-7 and Cyrano, part of the same regulatory circuitry as ciRS-7. Using ciRS-7 and miR-7 knock-out mouse models, we revealed the potential role of ciRS-7 in preventing miR-7-mediated regulation of the target mRNAs, unveiling the contribution of ciRS-7 molecular network in preventing stroke-induced cell death.

RESULTS

Oxygen and glucose deprivation induces changes in the expression of circular RNAs

Given the association of circRNAs with oxidative stress response[18] we evaluated their potential deregulation in conditions mimicking ischemic stroke *in vitro*. We cultured murine cortical neurons isolated from embryonic day fifteen (E15) cortices, subjected them to oxygen and glucose deprivation (OGD) for 12 hours (Supplementary Figure S1A) and performed total RNA sequencing.

A deconvolution analysis on single-cell RNA-seq dataset of embryonic mouse brain (E14.5[19]) confirmed that our culture is representative of the murine cortex, identifying interneurons (Int) and pyramidal neurons (Layer V-VI) as the predominant cell populations (Figure 1A). Functional validation using calcium imaging recording upon GABA and glutamate stimulation revealed that our culture primarily consists of glutamatergic excitatory neurons, as 98% of cells in our culture responded to glutamate and 36% to GABA (Figure 1B).

Ingenuity Pathway Analysis (IPA) of the differentially expressed genes between normoxic and OGD neurons confirmed the activation of Hypoxia-inducible factor 1-alpha (HIF-1alpha) (Figure 1C) and downstream upregulation of glycolysis, a characteristic hallmark of OGD[20] (Supplementary Table S1, Supplementary Figure S1B). Moreover, colorimetric cell viability assay indicated 30% decrease in neuronal viability post-exposure (Supplementary Figure S1C), confirming vulnerability to ischemia-induced cell death.

CircRNAs, generated by back-splicing, are identified through back-splice junction reads (BSJ) spanning regions that are not present in regularly spliced transcripts (forward splice junction reads, FSJ) (Figure 1D). By using *CIRIquant* algorithm[21], which comprises several circRNAs identification tools, we identified 27 circRNAs with high confidence, of which 7 were significantly differentially expressed (DE) between normoxic (control) and OGD conditions (Figure 1E, Supplementary Table

S2). Notably, ciRS-7 was the most expressed circRNA yielding the highest number of back-splice

junctions (Figure 1F, left) and the highest circular to linear ratio (Junction ratio) (Figure 1F, right).

The obtained junction ratio score of above 0.88, indicate that the transcript generated by the ciRS-7

locus are over 88% in circular form, while other circRNAs exhibited only 40% of circular expression.

Specifically, ciRS-7 was one of the three downregulated circRNAs upon OGD with a $\log_2$FC of -

0.552.

Figure 1. ciRS-7 is the most abundant circRNA in mouse cortical neuron cultures and

downregulated by in vitro OGD.

(A) Box plot of bulk RNA-seq deconvolution scores, indicating neuronal type contributions to overall

gene expression profile. X-axis labels derive from initial categorization by the authors of the dataset

(GSE123335), sorted by median of universal semantic groups (e.g., Int1, Int2 in Interneurons). **(B)**

Calcium imaging analysis of cellular response to glutamate and GABA with representative trace (n = 11; signal > 5% of baseline; mean ± SD). **(C)** Bar plot of Ingenuity Pathway Analysis (IPA) of bulk RNA-seq data from OGD-treated cortical neurons. Z-score denotes prediction of transcription factor activation (positive) or inhibition (negative); significant HIF1α activation is highlighted in green (|z| > 2 is considered significant). **(D)** Schematic of CIRIquant algorithm: Gene with three exons (dark blue, light blue, purple) produces a circular molecule via back-splicing or a linear transcript. Green bars denotes specific circular reads on back-splice junction (BSJ), the reads from forward splicing junctions (FSJ) in the same region are in gray. The Junction ratio score formula calculates a circularization score, indicating the percentage of the gene in circular form. **(E)** Heat map of logCPM for detected circRNAs in normoxic (N) and oxygen/glucose-deprived (OGD) murine cortical neurons. Green indicates low expression, while purple indicates high expression. Significantly differentially expressed circRNAs are grouped with a bar (* = adj. p-value < 0.05, n = 5). **(F)** (*left*) Bar plot of CPM of the BSJ and FSJ of differentially expressed circularRNAs between normoxic and OGD conditions, BSJ represents circular specific reads. (*right*) Bar plot of the Junction ratio score bar plot from CIRIquant algorithm, showing the ratio of BSJ to FSJ reads mapped to the BSJ site and indicating the percentage of circularization relative to the linear transcript.

ciRS-7 network is altered in conditions mimicking ischemic stroke in vitro

ciRS-7 is part of a feedback loop with miR-671 and miR-7 microRNA[5,7] and indirectly, through miR-7, with the long non-coding RNA Cyrano[8] (Figure 2A). To investigate the relative expression of these players in *in vitro* OGD conditions, we performed small RNA sequencing of cortical neurons exposed to OGD for 12h. Together with the previously presented dataset, we were able to capture circRNAs, mRNAs, long non-coding RNAs and small RNA transcripts (Supplementary Table S2).

Small RNA-sequencing identified 333 upregulated and 88 downregulated miRNAs following OGD (Figure 2B). Notably, the levels of miR-671-5p remained unaltered (Figure 2B) which was confirmed

independently by RT-qPCR (Supplementary Figure S2A), in contrast to previous reports of divergent miR-671-5p and ciRS-7 expression in different contexts[7–9]. Instead, we detected a significant upregulation in miR-7a-5p, but not of the variant miR-7b-5p (Figure 2B, Supplementary Table S2). Similarly to human, in mouse miR-7 is redundantly encoded by three different loci (*miR-7a-1*, *miR-7a-2*, *miR-7b*), each produced from different primary transcript (pri-miRNA) and precursor (pre-miRNA). Further processing, in all cases from -5p arm, give rise to two mature miR-7 sequences (miR-7a and miR-7b), differing only by a single nucleotide in the non-seed position number 10 (Figure 2C). To study the expression of the three miR-7 loci in our system, we utilized Global run-on sequencing (GRO-seq), which enable the capture of nuclear nascent RNA primary molecules, and compared the expression in our murine culture[22] with different mouse tissues and cell lines[22–32] (Supplementary Table S3). In physiological conditions, only cortical neurons and the pituitary gland actively transcribe all three independently regulated *miR-7* loci (Figure 2D, Supplementary figure S3). The active transcription from three different loci in neuronal cells suggests a higher order of regulation of this miRNA which influences the mature forms. Under OGD conditions, only the pri-miRNAs contributing to the expression of miR-7a (miR-7a-1 and miR-7a-2) were significantly upregulated (Figure 2E), indicating that the observed upregulation occurs already at the level of transcription. None of miR-7 passenger strands generated from the three precursors (miR-7a-1-3p, miR-7a-2-3p, and miR-7b-3p) were altered in our sequencing data (Supplementary Table S2). Moreover, we detected no changes in the host gene in which miR-7a-1 is embedded *(Hnrnpk)*, and from which the most abundant miR-7 primary transcript is generated (Supplementary Table S2). Taken together these data show that OGD specifically regulates miR-7a-5p variant at the transcriptional and/or post-transcriptional level.

To test the possible functional relevance of the upregulation of miR-7a-5p, we acquired predicted and validated targets of miR-7a-5p using *miRWalk* algorithm analysis[33,34] and compared their overall

expression in conditions of OGD. In accordance with the upregulation of miR-7, miR-7 targets

showed a significant overall downregulation (Figure 2F, Supplementary Table S4, Supplementary

Figure S2B), suggesting a canonical functional repressive role of this miRNA.

The role of miR-7 interaction with ciRS-7 remains controversial, and it is thought to set the balance

between a positive and negative feedback[8,9] (Figure 2A). Additionally, Cyrano promotes target-

directed miRNA degradation (TDMD) of miR-7 through a site of almost perfect complementarity,

indirectly regulating ciRS-7 levels and localization[8] (Figure 2A). In our dataset the lncRNA Cyrano

was significantly downregulated upon OGD (Figure 2G), in line with the observed upregulation in

miR-7a-5p (Figure 2B). To identify a first responder to OGD in the context of this molecular circuitry

we evaluate the time-dependent changes in the levels of these molecules. We subjected cortical

neurons to OGD for 1, 6, 12 and 24 hours and assessed the expression level of ciRS-7, miR-7 and

Cyrano. We detected a significant downregulation of ciRS-7 and Cyrano already at 6 hours after

OGD, prior to the upregulation of miR-7 at 12 hours (Figure 2H). Interestingly, our data revealed a

significant downregulation of miR-7 at 6h of OGD concomitantly with downregulation in ciRS-7 and

Cyrano, which is in line with the previously suggested role of ciRS-7 in stabilizing miR-7[9].

Figure 2. ciRS-7 network is dynamically altered in ischemic stroke like conditions.

(A) Schematic of the regulatory network involving ciRS-7, miR-7, Cyrano, and miR-671. References:
(1) Hansen et al., EMBO J, 2011; Hansen et al., Nature, 2013; (2) Piwecka et al., Science, 2017; (3)
Kleaveland et al., Cell, 2018. **(B)** Volcano plot of differentially expressed microRNAs in OGD-
treated cortical neuron cultures. Significantly upregulated (pink) and downregulated (green)
microRNAs, miR-7 variants (red), miR-7 passenger strands (white), and miR-671 (black) are

highlighted (n = 5; p-adj. < 0.05 and |log2FC| > 0.3). **(C)** Schematic of murine genomic loci illustrating the transcription of miR-7 primary transcripts (pri-miR), their processing into precursor molecules (pre-miR), and maturation into mature miR-7. The shared seed sequence between miR-7a-5p and miR-7b-5p is highlighted in red, and the non-seed mismatch at position 10 between the variants is underlined in black. **(D)** Bar plot of miR-7 loci reads per kilobase million (RPKM) in different cell type obtained from GRO-seq datasets. **(E)** Bar plot of miR-7 pri-miRNA expression in cortical neurons post-OGD treatment quantified by qPCR. Relative expression normalized to normoxic condition (n = 4; * $p < 0.05$, ** $p < 0.01$, paired t-test; mean ± SD).**(F)** Cumulative distribution functions (CDFs) plot of log fold changes of genes in OGD-treated wild-type cortical neurons (12h) compared to normoxic conditions. Red curve represents CDF for miR-7 targets, and black curve for non-targets (n = 5; p-value from Kolmogorov-Smirnov test). **(G)** Volcano plot of differentially expressed transcripts in OGD-treated cortical neuron cultures. Significantly upregulated (pink) and downregulated (green) transcripts, Cyrano (purple), and miR-7 hosting gene HnrnpK (black) are highlighted (n = 5; p-adj. < 0.05 and |log2FC| > 0.3). **(H)** Bar plot of ciRS-7 (blue), miR-7 (magenta), and Cyrano (purple) quantification by qPCR in OGD-treated cortical neurons at various timepoints (1h, 6h, 12h, and 24h). Relative expression normalized to normoxic condition (normoxia n = 2-4, OGD n = 5-6; * $p < 0.05$, *** $p < 0.001$, **** $p < 0.0001$, unpaired t-test; mean ± SD).

ciRS-7 KO neurons exhibit a distinct outcome after OGD without altering the overall OGD response

The absence of ciRS-7 alters synaptic transmission in excitatory neurons and produces schizophrenia-like phenotype *in vivo*[9], connecting this molecule to glutamatergic transmission. This is of particular interest in conditions of ischemic stroke, where glutamate mediated excitotoxicity is a critical contributor to neuronal cell death[12]. Our cortical neuron culture enriched in glutamatergic neurons

(Figure 1B) provides a system in which these events can be studied *in vitro*. Neurons cultured from

ciRS-7 KO mice showed higher Ca^{2+}-responses to the stimulation with glutamate compared to their

wild-type (WT) counterparts (Figure 3A). Consequently, ciRS-7 KO neurons also exhibited a

significantly higher sensitivity to excitotoxicity upon exposure to high concentration of glutamate

(Figure 3B).

To test if this increased sensitivity to glutamate is affecting gene expression changes in OGD, we

subjected the ciRS-7 KO cortical cultures to OGD for 12 hours, following the previous experimental

design (Supplementary Figure S1A) and performed total RNA-seq. OGD induced deregulation in

total of 5767 genes in ciRS-7 KO neurons compared to normoxic conditions, of which the 85% were

shared with WT neurons (Figure 3C, Supplementary Table S5). Consistent with this, the analysis of

interaction terms in the differential expression analysis, aimed at distinguishing differences in the

OGD response between WT and ciRS-7 KO, revealed no significant difference in the overall OGD

response between the two genotypes (Supplementary Table S6). However, 871 genes were

differentially expressed only in ciRS-7 KO neurons, while the expression of 1260 genes was

exclusively altered in WT neurons subjected to OGD. ciRS-7 WT specific genes altered in OGD were

functionally enriched in mitochondrial processes, DNA metabolism, regulation of cell-cycle and

oxidative stress-induced apoptosis (Figure 3D). The 871 KO-specific genes were instead involved in

developmental and pro-regenerative processes (*progenitor migration to the cortex, DNA replication,*

transcription, cerebellar cortex development) (Figure 3E). Despite the genetic predisposition to

glutamate sensitivity of the ciRS-7 KO, which would suggest a deleterious outcome in the response

to OGD, ciRS-7 KO neurons regulate pathways of resilience during OGD. In support of this, ciRS-7

KO neurons did not show increased cell death compared to WT in response of OGD treatment

(Supplementary Figure S2C). This result implies a distinction in OGD outcome not attributed to a

change in the OGD response itself, but rather to the regulation of the cascade of processes downstream

of the response.

Interestingly, the lack of ciRS-7 did not affect OGD induced changes in the expression of miR-7 and

Cyrano (Figure 3F). Similar to WT neurons, Cyrano lncRNA was downregulated in ciRS-7 KO

neurons responding to OGD (Figure 3F, Supplementary Table S5). Moreover, even though ciRS-7

KO neurons stably express lower levels of miR-7a-5p[9] (Figure 3F, Supplementary Table S5), small-

RNA sequencing revealed a significant upregulation of miR-7a-5p in response to OGD also in these

neurons (Figure 3F, Supplementary Table S5). In line with our previous findings, we detected no

changes in miR-671-5p, miR-7b-5p and miR-7 passenger strand expression (Supplementary Table

S5). Nonetheless, whilst the expression in miR-7a-5p in OGD remained lower in ciRS-7 KO neurons

compared to WT neurons (Figure 3F), the targets of miR-7a-5p, both predicted and validated, showed

a significant global repression in these conditions (Figure 3G).

Figure 3. ciRS-7 KO genotype exhibits differential OGD phenotypical output without affecting the OGD induced changes in miR-7 and Cyrano. (A) (Left) Representative trace of calcium-induced fluorescence in response to glutamate treatment in wild-type and ciRS-7 KO cortical neurons. (Right) Scatter dot plot of calcium imaging analysis showing the cellular response to glutamate in wild-type and ciRS-7 KO cortical neurons (n = 8; * p < 0.05, unpaired t-test; central bar represents

median value). **(B)** Bar plot of relative absorbance in MTT viability assay for wild-type and ciRS-7 KO cortical neurons treated with vehicle or 250μM glutamate. Data presented as survival % normalized to wild-type vehicle ($n = 4$; * $p < 0.05$, **** $p < 0.0001$, one-way ANOVA corrected with Tukey's post-hoc test; mean $\pm$ SD). **(C)** Venn diagram of differentially expressed genes in normoxic versus OGD conditions of wild-type and ciRS-7 KO cortical neurons ($n = 5$; p-adj. < 0.05 and $|log2FC| > 0.3$). **(D)** Bar plot of functional enrichment analysis top 20 significant Metascape clusters performed on differentially expressed genes in wild-type normoxic versus OGD cortical neurons. **(E)** Bar plot of top 20 significant Metascape clusters from functional enrichment analysis on differentially expressed genes in wild-type normoxic versus OGD cortical neurons. **(F)** (Left) Bar plot of log2FC for ciRS-7 (blue), miR-7 (magenta), and Cyrano (purple) from RNA-seq of wild-type and ciRS-7 KO cortical neurons in normoxic conditions versus after 12h of OGD treatment. (Right) Bar plot of log2FC for miR-7 (magenta) from RNA-seq, highlighting expression differences in normoxic conditions (N) or OGD between wild-type and ciRS-7 KO genotypes ($n = 5$; p-adj. < 0.05 and $|log2FC| > 0.3$). **(G)** Cumulative distribution functions (CDFs) plot of log fold changes of genes in OGD-treated ciRS-7 KO cortical neurons (12h) compared to normoxic conditions. The red curve represents CDF for miR-7 targets, while the black curve represents non-targets ($n = 5$; p-value from Kolmogorov-Smirnov test).

Alteration of ciRS-7 affects ischemic stroke outcome in vivo

To assess the reproducibility of our findings in ischemic stroke *in vivo*, *BALB/c* mice were subjected to permanent middle cerebral artery occlusion (pMCAO) and levels of ciRS-7, Cyrano and miR-7a-5p were evaluated by qPCR at six hours, one day and five days post-ischemia. In line with our *in vitro* results, we detected a significant downregulation of ciRS-7 and Cyrano and upregulation of miR-7a-5p in the peri-ischemic cortex (PI) at one day post-ischemia (dpi) compared to contralateral cortex (CL) (Figure 4A). Moreover, similar to OGD, we failed to detect alterations in miR-671-5p levels (Supplementary Figure S2D).

To explore the functional implications of ciRS-7 KO neurons ambiguous behavior involving glutamate sensitivity without increased cell death during OGD, we tested whether ciRS-7 KO mice exposed to transient ischemic stroke would show altered vulnerability to ischemic damage and associated sensorimotor impairments. The ischemic lesion size was measured using Magnetic Resonance Imaging (MRI) and behavioral deficits were assessed using Neurological Severity Scores (NSS) at acute (1dpi), subacute (3dpi) and chronic (7dpi) timepoints after transient middle cerebral artery occlusion (tMCAO) (Figure 4B). At 1dpi, ciRS-7 KO mice showed a significant reduction in the ischemic lesion volume compared to their WT controls (Figure 4C) and significantly ameliorated motor deficits (Figure 4D). Interestingly, ciRS-7 KO animals lesion size changes occurred transiently at the acute timepoint (1dpi), without delaying damage progression (Supplementary Figure S4A,B). To identify the molecular changes associated with acute reduction in the lesion size and motor deficits in ciRS-7 KO mice, mice were sacrificed at 1dpi and peri-ischemic and the contralateral cortex was used for mRNA sequencing. IPA analysis of differentially expressed genes in the peri-ischemic cortex of ciRS-7 KO animals indicated substantial inhibition in pathways related to neuronal and tissue damage, cytokine and interleukin release, and calcium influx compared to WT controls (Figure 4E, Supplementary Table S7). Moreover, most differentially expressed genes were mostly contributing

to pathways of *necrosis* (205 genes) and *quantity of cytokine* (41 genes), suggesting that the decreased

lesion size in the ciRS-7 KO mice may be due to inhibition of necrotic pathways and reduction in

cytokine release.

Aiming to identify crucial cellular responders mediating the observed gene expression changes, we

carried out deconvolution analysis by using adult mouse brain scRNA-seq data[35] to assess the

contribution of different cell populations in our bulk mRNA sequencing dataset. This analysis

identified neurons and astrocytes as major cellular responders contributing to WT and ciRS-7 KO

gene expression profiles (Figure 4F). In line with the significantly reduced tissue death measured by

MRI (Figure 4C), ciRS-7 KO mice exhibited increased neuronal involvement in the peri-infarct

cortex compared to WT mice, accompanied by a decrease in astrocyte contribution. To confirm this,

brain slices from ischemic ciRS-7 KO mice were collected at the peak of the immune response (7dpi)

[36] and stained for astrocytic glial fibrillary acidic protein (GFAP). Although ischemic stroke induced

GFAP expression in the peri-infarct cortex for both WT and ciRS-7 KO mice, ciRS-7 KO mice

exhibited significantly reduced astrogliosis compared to WT animals (Figure 4G).

At molecular level, tMCAO significantly downregulated ciRS-7 in WT mice at 1dpi in the peri-

ischemic cortex, while miR-7-5p and Cyrano levels remained unaltered (Figure 4H, Supplementary

Figure S4C). However, ciRS-7 KO mice showed a significant upregulation of miR-7-5p and unaltered

Cyrano levels (Figure 4H). These data suggest a correlation between the absence of ciRS-7 and a

faster upregulation of miR-7 in response to transient ischemic stroke and a marginal role of Cyrano

in this system at this timepoint.

Figure 4. Lack of ciRS-7 ameliorates ischemic stroke outcome in vivo.

(A) Bar plot of qPCR quantification for ciRS-7 (blue), miR-7 (magenta), and Cyrano (purple) in 3-4

months old BALB/c mice after pMCAo surgery. Peri-ischemic (PI) and contralateral (CL) cortices

were collected at different timepoints (6h, 1d, and 5d). Relative expression normalized to the average

CL expression (n = 6-7; * p < 0.05, *** p < 0.001, **** p < 0.0001, paired t-test; mean ± SD). **(B)**

Schematic of the experimental design: Wild-type and ciRS-7 KO mice underwent tMCAO surgery,

followed by neuroscore testing and MRI monitoring at 1, 3, and 7 days post-surgery (dpi). At 1dpi,

six mice were utilized to obtain peri-ischemic and contralateral cortices for RNA-sequencing.

Immunohistochemistry analysis was performed at later timepoints using eight to nine mice per group.

(C) (Left) Bar plot of MRI quantification of the lesion size in wild-type and ciRS-7 KO mice 1 day

post-tMCAO surgery. Data presented as lesion percentage on total brain size adjusted for edema (n =

25-28 per group; ** p < 0.01, Mann–Whitney test; median with 95% confidence interval). (Right)

Representative MRI image illustrating changes in lesion perimeter (blue) between wild-type (WT)

and ciRS-7 KO animals. **(D)** Scatter dot plot of sensorimotor deficits assessed by neuroscore in

tMCAO wild-type and ciRS-7 KO mice at 1dpi. Data presented as total neuroscore (n = 22-25 per

group; * p < 0.05, Mann–Whitney test; median with 95% confidence interval). **(E)** Bubble plot of

Ingenuity Pathway Analysis (IPA) for differentially expressed genes between wild-type and ciRS-7

KO peri-ischemic cortices from RNA-seq. Z-score reflects IPA prediction of pathway activation

(positive) or inhibition (negative) in ciRS-7 KO. Color bar indicates p-value significance (yellow to

purple), and bubble size represents the number of genes in the pathway ($|z| > 2$ considered significant;

pathways with >10 genes included). **(F)** (Left) Spiderweb plot of bulk RNA-seq deconvolution scores

of wild-type (gray) and ciRS-7 KO (blue) contralateral and peri-ischemic cortices. (Right) Bar plot

illustrating neurons and astrocyte contributions based on deconvolution scores in wild-type (gray)

and ciRS-7 KO (blue) contralateral and peri-ischemic regions. **(G)** (Left) Bar plot of quantification

for GFAP DAB (3,3′-Diaminobenzidine) immunostaining in wild-type and ciRS-7 KO tMCAO mice

at 7dpi. (Right) Representative images of the staining highlighting GFAP[+] cells. Relative expression

normalized to the average contralateral expression of wild-type mice (n = 8; * p < 0.05, ** p < 0.01,

**** p < 0.0001, one-way ANOVA corrected with Tukey's post-hoc test; mean ± SD). **(H)** Bar plot

of qPCR quantification for ciRS-7 (blue), miR-7 (magenta), and Cyrano (purple) in contralateral (CL)

and peri-ischemic (PI) cortices of ischemic wild-type and ciRS-7 KO mice at 1dpi. Schematic

representation correlates with MRI lesion size at the same timepoint. Data normalized to the average

contralateral expression of wild-type mice (n = 5 per group; * p < 0.05, ** p < 0.01, paired t-test;

mean ± SD).

ciRS-7 prevents miR-7 effects on dampening glutamatergic signaling in excitatory neurons

Previous studies linked miR-7 to oxidative stress response[37], in particular in low glucose conditions[38].

To establish the possible functional link between the upregulation of miR-7 in ciRS-7 KO mice in

conditions of ischemic stroke, we subjected cre-loxP inducible miR-7 KO mice[39] to tMCAO (Figure

5A, Supplementary Figure S5A,B). At the acute timepoint (1dpi), no changes in lesion volume, as

measured by MRI, were detected between miR-7 KO and WT mice (Supplementary Figure S5C).

Accordingly, miR-7 KO mice did not show any differences in motor deficits compared to their WT

controls (Supplementary Figure S5D). However, miR-7 KO mice (Figure 5B) exhibited an increase

in the lesion volume compared to their WT controls at later timepoint (7 dpi), unrelated to changes in

Neuroscore (Supplementary Figure S5E), indicating a potential role of miR-7 in regulating ischemia-

induced secondary cell death.

To understand the functional role of miR-7 in ischemic stroke, we first validated miR-7 functional

target sites in our system by analyzing miR-7 target binding sites in an Ago2 HITS-CLIP sequencing

dataset of pyramidal excitatory neurons[40] (Figure 5C, Supplementary Table S8). By identifying

RISC-associated binding sites on expressed transcripts in mature pyramidal neurons, our analysis of

the minimal seed site hexamer of miR-7 revealed 645 potential targets in the 3' UTRs (Figure 5D).

As expected, ciRS-7 had the highest number of functional binding sites (n=139), representing over

11% of the total reads (Supplementary Table S8). The intersection of Ago2 HITS-CLIP physical

targets and the downregulated genes in the peri-ischemic cortex of ciRS-7 KO mice, where miR-7 is

upregulated, highlighted 276 common genes (Figure 5E). These genes were functionally enriched in pathways related to glutamatergic synaptic transmission and morphological changes of neuronal projections (Figure 5F, Supplementary Figure S6).

To discern the specific contributions of ciRS-7 and miR-7 to glutamatergic signaling and neuronal morphological changes, we opted to replicate the molecular changes associated with low ciRS-7 and high miR-7 *in vitro*. We transduced ciRS-7 WT and KO neurons with lentivirus overexpressing GFP and miR-7a or GFP only, under the neuronal specific human Synapsin 1 (hSYN) promoter. Infection efficiency was evaluated through fluorescence microscopy (Supplementary Figure S7A) and we confirmed by RT-qPCR that miR-7a-5p levels resembled the endogenous levels under OGD condition (Supplementary Figure S7B). Whilst overexpression of miR-7 in WT neurons did not affect the overall neuronal glutamate excitability, increased levels of miR-7 in ciRS-7 KO neurons significantly reduced their response to glutamate (Figure 5G), thus reverting the genotype of glutamate sensibility of the ciRS-7 KO neurons. This appeared to be a glutamate specific effect, as the overexpression of miR-7 did not affect GABAergic response (Supplementary Figure S7C).

Finally, to evaluate the functional impact of the potential targets of miR-7 involved in morphological changes of neuronal projections (Figure 5F), we performed live imaging analysis of neurite outgrowth of ciRS-7 KO and WT neurons infected with miR-7 lentivirus. In agreement with the calcium imaging experiment, we were able to detect a significant increase in the neurite length when over expressing miR-7 in ciRS-7 KO cells, but not in WT neurons (Figure 5H). This aligns with a previous study showing a direct impact of miR-7 overexpression on neurite length in a neuroblastoma cell line[41]. Considering circRNAs are typically low in cell lines due to their rapid proliferation[42], we questioned whether the effect of miR-7 on neurite length was due to the absence of ciRS-7 interference.

372 However, circRNAs are known to be poorly expressed in cell lines due of their high proliferation

373 rate[42]. So, we wondered whether this miR-7 direct effect on neurite length was due to lack of ciRS-7

374 interference. We confirmed that the murine neuroblastoma cell line N2A expresses negligible ciRS-

375 7 levels even when differentiated with retinoid acid, despite higher levels of Cyrano and miR-7

376 (Supplementary Figure S8A,B,C).

Figure 5. Lack of miR-7 exaggerates ischemic stroke outcome regulating glutamatergic response.

(A) Schematic of Cre-LoxP miR-7 KO animal model: mice have a transgenic genome with LoxP sequences (blue) flanking all three miR-7 loci (magenta), and tamoxifen (orange)-inducible Cre recombinase (purple). Tamoxifen activation of Cre recombinase leads to miR-7 loci deletion. Control mice lack the Cre recombinase transgene. **(B)** (Left) Bar plot of MRI quantification for lesion size in wild-type and miR-7 KO mice one day post-tMCAO surgery. Data presented as lesion percentage on total brain size adjusted for edema (n = 8-11 per group; * p < 0.05, Mann–Whitney test; median with 95% confidence interval). (Right) Representative MRI image illustrating changes in lesion perimeter (blue) between wild-type (WT) and miR-7 KO animals. **(C)** Schematic of recombinant mice for generating Ago2 CLIP-seq data in excitatory neurons: FLAG-Ago2 (blue) is selectively translated in Cam2Ka$^+$ (red) excitatory neurons, where Cre recombinase (yellow) expression removes a stop codon (gray) leading to FLAG-Ago2 translation. Following cross-linking (thunder), the FLAG-Ago2-miRNA-target complex is immunoprecipitated with an anti-FLAG antibody (purple), and the sequenced output represents the miRNA-target complex. **(D)** Pie chart illustrating the distribution of miR-7 binding sites from CLIP-seq analysis. Among the 3779 identified binding sites, categories include protein coding sequences (CDS, purple), 5' untranslated regions (5' UTR, blue), 3' untranslated regions (3' UTR, dark green), and intronic sequences (Intron, light green). **(E)** Venn diagram depicting the miR-7 binding sites in the 3' UTR identified by CLIP-seq in excitatory neurons (645) and the downregulated genes in the peri-ischemic region of ciRS-7 KO tMCAO animals (3286), where miR-7 is upregulated. The overlap (276 genes) represents potential miR-7 physical targets affected in ischemic stroke. **(F)** Bar plot of the top 20 significant Metascape clusters from functional enrichment analysis on genes identified as miR-7 physical targets potentially affected in ischemic stroke. **(G)** (Left) Representative trace of calcium-induced fluorescence in response to glutamate treatment in wild-type and ciRS-7 KO neurons infected with miR-7 or control lentivirus. (Right)

Scatter dot plot of calcium imaging analysis quantifying the cellular response to glutamate in wild-

type and ciRS-7 KO cortical neurons infected with control (GFP) or miR-7 overexpressing (GFP +

miR-7) virus (n = 5; * $p < 0.05$, **** $p < 0.0001$, Kruskal–Wallis test; central bar represents median

value). **(H)** (Left) Bar plot of neurite length quantified through live imaging of wild-type and ciRS-7

KO cortical neurons infected with control (GFP) or miR-7 overexpressing lentivirus (GFP + miR-7).

Data normalized to the average neurite length of wild-type GFP-infected neurons (n = 3 biological

replicates, n = 8 technical replicates; ** $p < 0.01$, one-way ANOVA test corrected with Tukey's post-

hoc test; mean ± SD). (Right) Representative image of ciRS-7 KO cortical neurons infected with

control (GFP) or miR-7 overexpressing virus (GFP + miR-7).

miR-7 modulates AMPA-mediated glutamatergic signaling via ciRS-7/Cyrano network in ischemic

conditions

To further unravel the intricate interactions within the miR-7/ciRS-7/Cyrano network, we conducted

gain and loss of function experiments to modulate miR-7 expression in the context of ischemic stroke.

We utilized a lentivirus expressing the Cyrano CR1 region, known for its role in mediating miR-7

degradation through TDMD[8]. Wild-type cortical neurons were transduced with lentivirus carrying

CR1 or a control region featuring a mutated miR-7 binding site before exposure to a 12-hour period

of OGD. As expected, overexpression of Cyrano CR1 region led to a decrease in miR-7 expression

in normoxic conditions. Additionally, it prevented the OGD-induced increase in miR-7, in stark

contrast to the mutated CR1, which had no impact on OGD-induced upregulation of miR-7 (Fig. 6A).

Remarkably, preventing the increase in miR-7 expression preserved the levels of ciRS-7 during OGD,

disclosing a previously proposed antagonistic relationship between these molecules[8]. The

overexpression of the synthetic construct had no influence on the OGD-induced downregulation of

the endogenous full-length Cyrano. Conversely, lentiviral induced overexpression of miR-7 induced

an amplified miR-7 response during OGD (Fig. 6B). This molecular phenotype was consistently

associated with a downregulation of ciRS-7, even under physiological conditions, further suggesting

a potential involvement of miR-7 expression in the downregulation of ciRS-7. As expected, miR-7

overexpression had no effect on Cyrano basal expression or OGD-induced Cyrano downregulation

(Fig. 6B).

By leveraging miR-7 targets validated through CLIP-seq that were dysregulated in ischemic mice

(Fig. 5E, Supplementary Table S7), we pinpointed Gria1, a gene encoding a subunit of the AMPA

receptor, critical for mediating glutamate excitotoxicity in ischemic stroke pathophysiology[43]. The 3'

UTR of Gria1 contains two potential 6mer binding sites for miR-7a-5p (Fig. 6C). To validate the

physical interaction of these binding sites with miR-7a, we co-expressed a luciferase vector

containing the 3' UTR of Gria1 and miR-7a-5p in the HEK293T cell line. We observed a significant

post-transcriptional canonical effect of miR-7 on the 3'UTR which was absent with an unrelated

control miRNA (Fig. 6C). To further confirm the significance of this relationship in conditions of

ischemic stroke, we measured the levels of Gria1 in cortical neurons transduced with miR-7 lentivirus

prior OGD for 12h. In line with the RNA-seq data (Supplementary Figure S2B), OGD treatment led

to a decrease in Gria1 expression in neurons transduced with the control vector. However,

overexpression of miR-7 resulted in a 50% reduction in Gria1 mRNA levels under normal oxygen

conditions, with no further alterations during OGD (Fig 6D). Overexpression of Cyrano CR1 region

in neurons subjected to OGD prevented the OGD-induced downregulation of Gria1 (Fig. 6E). To

demonstrate that the miR-7-mediated effect on Gria1 lead to a decrease response to AMPA, we

carried out calcium imaging in ciRS-7 KO neurons that were lentivirally transduced to overexpress

miR-7. In these conditions, ciRS-7 buffering effect on miR-7 is disrupted. As expected, the

upregulation of miR-7 had no impact on wild-type neurons. However, in ciRS-7 KO neurons, we

observed a significant decrease in AMPA-induced Ca^{2+} responses in miR-7 overexpressing neurons

(Fig. 6F). This finding reinforces the link between miR-7 and glutamatergic signaling, indicating that

the diminished glutamatergic response attributed to miR-7 is, to some extent, a result of suppression

of AMPA signaling achieved through the downregulation of Gria1.

Figure 6. miR-7 regulation in OGD depends on ciRS-7 and Cyrano and modulate AMPA

response. (A) (Left) Schematic of the wild-type Cyrano CR1 region (CR1) and its miR-7 binding

sites mutated version (MUT). Magenta indicates pairing with miR-7a-5p, encompassing both seed

region and a more extensive one. (Right) Bar plot of qPCR quantification for miR-7 (magenta), ciRS-

7 (blue), and Cyrano (purple) in cortical neurons overexpressing Cyrano CR1 region (Cyrano CR1)

or its mutated version (MUT) along with mCherry, treated with OGD for 12 hours. Relative

expression normalized to normoxic condition of the mutated lentivirus (n = 5; * p < 0.05, *** p <

0.001, **** p < 0.0001, one-way ANOVA corrected with Tukey's post-hoc test; mean ± SD). **(B)** Bar

plot of qPCR quantification for miR-7 (magenta), ciRS-7 (blue), and Cyrano (purple) in cortical

neurons overexpressing miR-7 GFP or a control vector with only GFP (Ctrl), treated with OGD for

12 hours. Relative expression normalized to normoxic condition of the control lentivirus (n = 5; * p

< 0.05, ** p < 0.01, *** p < 0.001, one-way ANOVA corrected with Tukey's post-hoc test; mean ±

SD). **(C)** (Left) Schematic of the heteromeric AMPA receptor with its four subunits highlighted in

different colors. Gria1, encoding the GluR1 subunit, is highlighted in magenta. Two 6mer sites on Gria1 3' UTR and their potential pairing with miR-7a-5p in the seed region are shown (underlined). (Right) Bar plot of luciferase assay fluorescence quantification in HEKT293 cells transfected with control vector (Ctrl vector), Gria1 3' UTR, and miR-532-5p (lacking binding sites) or miR-7a-5p (magenta). Relative expression normalized to Gria1 3' UTR vector alone (n = 3; **** $p < 0.0001$, one-way ANOVA corrected with Tukey's post-hoc test; mean ± SD). **(D)** Bar plot of qPCR quantification for Gria1 (magenta) in cortical neurons overexpressing miR-7 GFP or a control vector with only GFP (Ctrl), treated with OGD for 12 hours. Relative expression normalized to normoxic condition of the control lentivirus (n = 5; * $p < 0.05$, *** $p < 0.001$, one-way ANOVA corrected with Tukey's post-hoc test; mean ± SD). **(E)** Bar plot of qPCR quantification for Gria1 (magenta) in cortical neurons overexpressing Cyrano CR1 region (Cyrano CR1) or its mutated version (MUT) along with mCherry, treated with OGD for 12 hours. Relative expression normalized to normoxic condition of the mutated lentivirus (n = 5; **** $p < 0.0001$, one-way ANOVA corrected with Tukey's post-hoc test; mean ± SD). **(F)** (Left) Representative trace of calcium-induced fluorescence in response to AMPA treatment in wild-type and ciRS-7 KO cortical neurons transduced with miR-7 or control lentivirus. (Right) Scatter dot plot of calcium imaging analysis of the cellular response to AMPA in wild-type and ciRS-7 KO cortical neurons infected with control (GFP) or miR-7 overexpressing (GFP + miR-7) virus (n = 6-7; **** $p < 0.0001$, Kruskal–Wallis test; central bar represents median value).

DISCUSSION

Here we identify oxygen and glucose deprivation as a metabolic stressor triggering endogenous changes in the ncRNA network of ciRS-7 – miR-7 – Cyrano both *in vitro* and *in vivo* during permanent ischemic stroke. In OGD *in vitro* model, we demonstrate dynamic changes in gene expression for these molecules independent of ciRS-7. In *in vivo* condition of transient ischemic stroke, we found no changes in the lncRNA Cyrano. However, the absence of ciRS-7 reduced cellular death and sensorimotor deficits, while the lack of miR-7 resulted in more extensive tissue damage. This suggests that both ciRS-7 and miR-7 may play a more crucial role in this system. In an *in vitro* model recapitulating the molecular changes occurring during ischemic stroke (low ciRS-7 and high miR-7), these effects were partly executed through miR-7-mediated regulation of glutamatergic signaling, contingent on the absence of ciRS-7. These data suggest that ciRS-7 may regulate miR-7 targeting, highlighting an endogenous regulatory role for the ciRS-7/miR-7 network in mediating cellular stress responses under pathophysiological conditions.

In mice, ciRS-7 harbors 130 binding sites for miR-7[9] and, because of its remarkably high expression in neurons[5,9], the limited molecules of miR-7 in physiological conditions[5] are likely associated to ciRS-7, as indicated by our pyramidal excitatory neurons CLIP-seq data and by others[9,10]. Moreover, Cyrano, highly abundant in neurons, harbors a single nearly complementary site for miR-7 which mediates miR-7 degradation[8]. Our data show that OGD induces changes in the neuronal physiological landscape by concomitant decrease in ciRS-7 and Cyrano and subsequent upregulation of miR-7. These changes seem to be independently regulated, as the absence of ciRS-7 did not influence OGD-induced alterations in the gene expression in miR-7 and Cyrano. While the anticorrelation between Cyrano and miR-7 follow established dynamics, with Cyrano triggering miR-7 degradation through TDMD[8], our study report the induction of miR-7 in the absence of ciRS-7, suggesting that during ischemic stroke the expression of miR-7 does not depend on ciRS-7. Consistently, miR-7

overexpression induces ciRS-7 downregulation and this effect is rescued by overexpressing the Cyrano site that mediates miR-7 degradation through TDMD. Moreover, in instances where miR-7 suppresses ciRS-7 levels, miR-7 upregulation during OGD becomes even more pronounced.

The role of miR-7 in ischemic stroke has been studied *in vivo* with controversial results[44,45] and our extensive animal cohort revealed substantial biological variability in the expression levels of miR-7 in a mouse model of tMCAo. This outcome is potentially tied to established feedback loops among ciRS-7, Cyrano, and miR-7, as tMCAo in ciRS-7 knockout mice results in a significant overexpression of miR-7 one day post-ischemia, accompanied by a reduced stroke lesion size and motor deficits. Additionally, at 7 dpi, inducible miR-7 knockout mice exhibited an exacerbated ischemic lesion size compared to WT animals, which aligns with literature reporting that miR-7 is necessary for stroke recovery[46] , further suggesting a role for miR-7 in regulating ischemic damage. The transient ciRS-7 KO phenotype is explained by an initial advantage provided by higher miR-7 levels, while subsequent ciRS-7 downregulation in WT animals minimizes genotypical differences at later timepoints. Furthermore, our in vitro data, mimicking molecular changes in ischemic conditions (low ciRS-7, high miR-7), indicate that under physiological conditions highly abundant ciRS-7 may function as a buffering system to regulate miR-7 targeting.

Given the lack of extensive changes in miR-7 targets in both ciRS-7, Cyrano and miR-7 KO animals in physiological conditions[8,9], the exact role of ciRS-7 as an influencer of miR-7 targeting has remained controversial. Our data show a significant shift in the overall expression of miR-7 targets during OGD, where a concerted downregulation of two known miR-7 regulators, ciRS-7 and Cyrano, occurs. Our data suggest that the inducible nature for this network may explain the reported absence of drastic effects on miR-7 targets in physiological conditions. Consistent with this concept, numerous studies show that single miRNA mutants exhibit a phenotype solely under stress conditions[47,48].

Moreover, unlike experimental settings involving acute treatment, knock-out models are subjected to

the buffering effect of cellular adaptation, as previously experienced[9].

The analysis of Ago2 CLIP-seq neuronal dataset revealed the lead miR-7 targets downregulated in

ciRS-7 KO ischemic mice belong to the functional class of glutamatergic signaling and neuronal

outgrowth. Glutamate plays a well-established role in ischemic stroke pathophysiology[12],

contributing to neuronal loss through exaggerated release and impaired clearance of glutamate in the

synaptic cleft[11]. Intriguingly, ciRS-7 KO neurons exhibit a pronounced sensitivity to glutamate in

physiological conditions, which represent a disadvantage in ischemic conditions, due to the

susceptibility to glutamate-mediated excitotoxicity. However, ciRS-7 KO neurons do not exhibit

increase cellular death upon OGD, but rather present a distinct regulation of processes following

OGD response, both *in vitro* and *in vivo*. We propose that the ameliorative effect during ischemic

stroke observed ciRS-7 KO mice may be linked to the levels of miR-7 and its unregulated accessibility

to targets. In line with this, overexpressing miR-7 dampened the neural glutamatergic response and

increased neuronal outgrowth only in ciRS-7 KO neurons, with no effect on the wild-type neurons.

A mechanism through which miR-7 partially exerts this role involves targeting the subunit 1 of

AMPA receptors (GluR1) by interacting with the 3' UTR of its mRNA, Gria1, thereby dampening

the AMPA response. Our data confirms that miR-7 effects on both glutamatergic signaling and neurite

outgrowth are dependent on ciRS-7 levels. In line with our results, the effect of miR-7 on neurite

outgrowth was previously reported in a neuroblastoma cell line[41] that expresses a negligible amount

of ciRS-7 even upon differentiation, suggesting a more direct effect of miR-7 in the absence of its

targeting regulator.

A recent study associates miR-7 with energy homeostasis in hypothalamic neurons[39] during

challenging energetic conditions similar to ischemic stroke. However, in this context, alterations in

miR-7 were not accompanied by changes in ciRS-7 and Cyrano[39]. Combined with our data revealing

diverse dynamics in the activation of the ciRS-7 network by various ischemic insults, this indicates

that the network response depends on the type, strength, and duration of the stress and the specific

cell type.

In summary, we propose a regulatory role for ciRS-7-miR-7 in glutamatergic signaling through miR-7 target genes, hence contributing to the control of post-ischemia neuronal damage. Our data support the hypothesis of a role for ciRS-7 in buffering miR-7 effects against unwanted changes, thus behaving as a "safe-guide" system. This study suggests a role of intracellular network of non-coding RNAs in regulating pathophysiological processes in the brain.

LIMITATIONS OF THE STUDY

The dynamic nature of multicellular systems, unlike in vitro cultures, results in data incongruencies across different timepoints and stressors. The intrinsic ability of regulatory molecules to induce phenotypic changes even at low doses complicates causality identification in processes involving multiple cellular events, such as disease onset. Growing number of studies revealing non-coding RNA alterations in pathophysiological in vitro systems underscores the necessity for in vivo research to understand the biological relevance of these complex regulatory networks. For a comprehensive analysis of ciRS-7 in ischemic stroke, future studies may benefit from in vivo investigations using Cyrano KO animals and direct rescue experiments on ciRS-7, although these remain technically challenging and time-consuming.

MATERIAL AND METHODS

PRIMARY AND IMMORTALIZED CELLS

Primary cortical neuron culture preparation and OGD treatment

Primary cortical neurons were prepared from C57BL/6J and C57BL/6N-Cdr1asem1Nikr (Cdr1as KO and their WT counterpart) embryonic day 15 embryonal cortices. After dissection and removal of the meninges, cortices were incubated 15 minutes at 37 °C in a solution of Krebs buffer (0.126 M NaCl, 2.5 mM KCl, 25 mM NaHCO3, 1.2 mM NaH2PO4, 1.2 mM MgCl2, 2.5 mM CaCl2, supplemented with 45mM BSA, 0.8% of 3.85% MgSO4 and 1% Pen/Strep, pH 7.4) and 0.025% (w/v) trypsin (Sigma-Aldrich, T 9201). Tissue was then treated with 0.008% w/v DNaseI (Sigma-Aldrich, DN25) and 0.026% w/v trypsin inhibitor (Sigma-Aldrich, T9003) and centrifuged at 300 x g for 3 minutes. Cell pellet was resuspended in 3ml of DNaseI/Trypsin solution and then diluted in 7ml of Krebs. After centrifugation at 300 x g for 3 minutes the pellet containing embryonic neurons was resuspended in cortical neurons growth media: Neurobasal (Gibco 21103049), B27 Supplement (Gibco, 17504044), 0.2 mM L-glutamine (Lonza, BE17-605E), 0.01 mg/ml Penicillin/Streptomycin (Gibco, 15140122). Cells were plated on Poly-D lysine (Sigma-Aldrich, P6407) freshly precoated plates (50 µg/ml in sterile water plates for 1h at 37 °C and washed in sterile water prior use). Different density was used for 6-well plates (1.8 million cells per well), 48-well plates (125.000 cells per well), 13mm plastic coverslips (30.000 cells per coverslip). After 5 days in culture half of the in cortical neurons growth media was changed to fresh. Experiments were performed 7 days after the isolation day. Cells were maintained in the incubator 37 °C, 5% CO2. For OGD treatment experiments, before hypoxia induction the media was changed to Normoxic (cortical neurons growth media with Neurobasal changed to Gibco A2477501 supplemented with D-Glucose 25mM and Sodium Pyruvate 0.2mM) or OGD (cortical neurons growth media with Neurobasal changed to Gibco A2477501). Normoxic cells were then put back in the incubator, OGD cells were incubated in hypoxic chamber

37 °C, 5%CO_2, 1%O_2 (SCI-tive N, Ruskinn Technology). After the OGD timepoint cells were

harvested for RNA extraction or subjected to MTT colorimetric test.

MTT test

Cell viability was measured from 48-well plates treating the cells with (3-(4, 5-dimethylthiazolyl-2)-

2, 5-diphenyltetrazolium bromide), MTT reagent (Sigma-Aldrich, TOX1) diluted with culture media

at a final concentration of 120 μM. Triton-X 100 1% v/v (Sigma-Aldrich, X100) treated wells were

used as a positive control for this assay. Plates were incubated for 3-5h at +37°C. After, medium was

discarded, formazan crystals were dissolved with dimethyl sulfoxide (DMSO) (Sigma-Aldrich,

D2650) for 30min at RT in the dark. Absorbance was read at 585nm using Wallac 1420 Victor2

microplate reader (Perkin Elmer). Wells without cells were used as background and subtracted from

the absorbance data, all the six technical replicates were plotted for each of the three biological

replicate and the data was normalized on normoxic WT neurons or WT neurons infected with GFP

only.

Neurite length measurement

Neurons were seeded at a density of 125.000 cells/well in 48-well plates. After OGD exposure or

lentiviral infection, neurons were imaged at day 7 for 48 hours with IncuCyte® S3 Live Cell Analysis

System (Essen BioScience Ltd.) in bright field and green channel live cell images (two 10x

magnification images per well). Acquired data were analysed with the Incucyte® Neurotrack

Analysis Software Module (Sartorius) considering the average value of the two images taken for each

of the nine technical replicates for all the three biological replicates.

RNA isolation and qRT-PCR

Total RNA was extracted from primary cells and ipsilateral/contralateral animal cortex using

TRIzol™ Reagent (Invitrogen) following the manufacturer's instructions. 1μl of GlycoBlue™

Coprecipitant (Ambion) was added at the isopropanol step in each sample. RNA was quantified using

a Nanodrop 2000 spectrophotometer. 1 μg of total RNA was used for reverse-transcription of

mRNAs, circRNA, lncRNAs and pri-miRNA species with High-Capacity cDNA Reverse

Transcription Kit (Applied Biosystems) following the manufacturer's protocol. SYBR Green qPCR

Master Mix (High ROX) (Bimake) and custom designed oligos were used to quantify mRNAs,

circRNAs, and lncRNAs following the manufacturer's indications. Mature microRNAs were reverse-

transcribed with TaqMan™ MicroRNA Reverse Transcription Kit (Invitrogen) using miRNA-

specific primers supplied in the TaqMan® probe kit for qPCR following the manufacturer's protocol.

Mature microRNAs and pri-microRNA PCRs were performed using Maxima Probe/ROX qPCR

Master Mix (Invitrogen) and TaqMan® specific probes (Thermofisher) following the manufacturer's

protocol. The result was analyzed with the ΔΔCT method and normalized geometric mean of 2

internal normalization controls (Gapdh and Rplp0) for SYBR green qPCRs and U6 expression for

TaqMan® miRNA and pri-miRNA. TaqMan® probes and sequence of the SYBR Green

oligonucleotide primers is available in the Supplementary Table S9.

Lentivirus vectors and virus generation

Lentiviral vectors LV1-eGFP (control) or LV1-eGFP-miR-7 were generated by subcloning inserts

from pAAV_hSYN1-eGFP-miR-7 and pAAV_hSYN1-eGFP (provided by Thomas B. Hansen)

inside LV1 (immunodeficiency virus 1 (HIV-1)-based LV-PGK-GFP) backbone by GenScript

Biotech Corporation. The generated construct contained HIV-1-LV backbone with hSYN1-eGFP-

miR-7 insert instead of PGK-GFPN inserted creating a terminal SmaI (CCCGGG) and a C-terminal

ApaI (GGGCCC) flanking restriction sites. The same was performed for the control vector with

hSYN1-eGFP insert only. In addition, GenScript Biotech Corporation generated lentiviral vectors,

namely LV1-mCherry-CR1 and LV1-mCherry-MUT, for the overexpression of the Cyrano CR1

region involved in miR-7 degradation through TDMD or a mutated form of it. The sequences for

these inserts were obtained from the prior publication by Kleaveland et al (2018)[8] and are accessible

on Addgene (https://www.addgene.org/128768/, https://www.addgene.org/128748/). These

sequences were then cloned into the previously established LV1-hSYN backbone, with the creation

of terminal AgeI (ACCGGT) and C-terminal SalI (GTCGAC) sites.

3rd generation lentiviral particles were produced by the BioCenter Kuopio National Virus Vector

Laboratory in Kuopio, Finland. The viral titer was assessed through qPCR serial dilution

quantification using eGFP or mCherry ReadyMade™ Primers (IDT) (Supplementary Table S9).

Work with the virus vectors was carried out under permission from Finnish National Supervisory

Authority for Welfare and Health, Valvira. Cells were infected at day 2 post isolation with MOI 0.5

achieving 80% positive infected cells assessed by GFP or mCherry expression in fluorescent

microscope (Supplementary Figure 7A,D) at day 7. The amount of overexpression of miR-7 or

Cyrano CR1 region was assessed by qPCR as above specified (Supplementary Figure 7B,E).

Library preparation OGD cortical neurons

All the samples RNA were isolated with TRIzol™ Reagent (Invitrogen) as specified above. RNA

samples were treated with TURBO DNA-free™ Kit (Ambion) following the manufacture's

instruction. RNA integrity was assessed through Agilent Bioanalyzer 2100 system with the Agilent

RNA 6000 Nano. The concentration of the samples was established with Qubit™ RNA Extended

range kit (Invitrogen). From cortical neurons subjected to OGD we generated 1) library to detect

circularRNAs and mRNA and lncRNAs using SMARTer® Stranded Total RNA Sample Prep Kit

(Takara Bio USA, Inc.) and 2) library for miRNAs detection utilizing NEBNext® Small RNA Library

Prep Set for Illumina (New England Biolabs (UK) Ltd) quality checked and size selected using pippin

prep method following manufacturer's protocol. All the libraries were generated following the

manufacturer's protocol. After generation, the libraries were quantified with Qubit™ High Sensitivity

DNA kit (Invitrogen) and by qPCR using KAPA Library Quantification Kit for Illumina® Platforms

(Roche). Library size was determined with Agilent Bioanalyzer 2100 system using the Agilent High

Sensitivity DNA Kit. Cortical neurons samples to detect circularRNAs, mRNA, and lncRNAs were

sequenced paired-end 100 cycles on NovaSeq™ 6000 platform (Illumina) and single-read 75 cycles

on NextSeq™ 500 system (Illumina) to detect microRNAs.

Calcium imaging

Calcium imaging was performed with murine E15 cortical neurons. The neurons were plated onto

PDL-coated circular plastic coverslips (13 mm diameter) in 12-well plate at a density of 30,000

cells/coverslip and kept for 5 days in Neurobasal medium supplemented with B27. To quantify and

compare the functional expression of glutamate and GABA receptors in neuronal cultures, we used

calcium-imaging technique as previously described[49]. Briefly, neuronal cultures were loaded with the

cell-permeable indicator Fluo-4am (Life Technologies, F10471) for 30 min at 37 °C, followed by 10

min washout, and placed in the perfusion chamber mounted on the stage of Olympus IX7010

microscope. Neurons were continuously perfused by basic salt solution (BSS) 3 ml/min containing

in mM: 152 NaCl, 10 HEPES, 10 glucose, 2.5 KCl, 2 CaCl2, 1 MgCl2 and pH adjusted to 7.4. Test

compounds diluted in the BSS to final concentrations were applied through fast perfusion system

(Rapid Solution Changer RSC-200, BioLogic Science Instruments). Cells were imaged with 10x

objective using Olympus IX-7010. Excitation wavelength was set as 494 nm, sampling frequency 2

FPS. Glutamate (100 µM with the co-agonist glycine 10 µM promoting activation of NMDA

receptors subtype) or GABA (100 µM) were applied for 2 s. Finally, KCl (50 mM) application for 2 s

was used to distinguish excitable neurons from possible non-neuronal cells. Fluorescence was

detected with the Till Photonics imaging system (FEI GmbH) equipped with a 12-bit CCD Camera

(SensiCam) with a light excitation wavelength of 494 nm. Calcium responses to neurotransmitters

were evaluated from changes in fluorescence intensity of individual neurons. To this end, regions of

interest (ROI) of round shape around the cell body were selected from the whole image with the TILL

vision Imaging Software (TILL Photonics GmbH). To distinguish from non-neuronal cells, ROI was

taken at the time point corresponding to KCl-induced activation of neurons. The intensity values in

each ROI were averaged at each time point to form a fluorescence signal for each neuron and

normalized to the baseline level. Signals above 5% of the baseline were included in the analysis.

N2a Cell culture

Immortalized murine neuroblastoma cells (N2a) were maintained in DMEM, high glucose,

GlutaMAX™ Supplement (Gibco, 31966021) media supplemented with supplemented with 10%

Fetal Bovine Serum (Gibco, 10270106) and 1% penicillin and streptomycin in temperature controlled

humidified incubator (37 °C, 5%CO_2). As previously published[50], for differentiation induction

100.000 cells were plated in a 6-well plate and maintained in complete media supplemented with

20μM retinoic acid (Sigma-Aldrich, R2625) in dimethyl sulfoxide (DMSO) with or without serum

starvation (2% FBS instead of 10%) changing half of the media every second day for 5 days before

collection.

Luciferase assay

Human embryonic kidney 293 cells containing the SV40 T-antigen (HEK293T) were cultured in

DMEM, high glucose, GlutaMAX™ Supplement (Gibco, 31966021) medium, supplemented with

10% Fetal Bovine Serum (Gibco, 10270106), and 1% penicillin and streptomycin in a temperature-

controlled humidified incubator (37 °C, 5%CO2). For the luciferase assay, 20,000 cells were seeded

per well in a 96-well plate. The following day, cells were co-transfected with 100ng of murine Gria1

3' UTR renilla/luciferase plasmid (MmiT076857-MT06-GC, GeneCopoeia) or miRNA 3' UTR target

control plasmid (CmiT000001-MT06-GC, GeneCopoeia) and 100nM of either miRIDIAN miRNA

mimic of mmu-miR-532 (C-310769-01-0002, Dharmacon) as a negative control or mmu-miR-7a-5p

(C-310591-07-0002, Dharmacon) using Lipofectamine™ 2000 (11668030, Invitrogen) according to

the manufacturer's protocol. After 48 hours, the luciferase assay was conducted with the Luc-Pair

Luciferase Assay Kit 2.0 (LF001-GC, GeneCopoeia), following the manufacturer's instructions,

except for the cell lysis step, which was performed directly on the plate in lysis buffer using three

freeze and thaw cycles with dry ice. Luciferase data were normalized to Renilla expression and

presented as the Fold Change of Relative Fluorescence between the samples with respect to 3' UTR

vector overexpression alone.

MOUSE STRAINS AND ANIMAL PROCEDURES

All experiments follow the Helsinki Declaration and guidelines set by the European Commission

(European Communities Council Directive, 86/609/EEC) and were approved by the National Animal

Experiment Board of Finland. Animals were housed with same sex siblings, in controlled

temperature, humidity and light (12 hours light/dark cycle) conditions. Animals had access to ad

libitum food. Before the beginning of the animal study mice were divided in single cages. Mice were

randomized and all the participants were blinded to the genotype (surgery, MRI acquisition,

behavioral test, sample collection and data analysis). Prior to the surgery, we use random number

generator (GraphPad Prism quick calcs https://www.graphpad.com/quickcalcs/randomN1/) to

randomize the mice into treatment groups. This numbering has been used in crescent order on 3-4

months old male mice that were divided into groups. pMCAO study was performed on 3-4 months

old BALB/cOlaHsd male mice (n = 6-8 per timepoint). Cdr1as (ciRS-7) KO tMCAO study was

performed on 3-4 months old C57BL/6N-Cdr1asem1Nikr[9] (Cdr1as KO mice and the WT

counterpart), provided by Prof. Dr. Nikolaus Rajewsky, MDC, Berlin, Germany (n = 22-24 per

genotype). miR-7 inducible KO tMCAO study was performed on 3-4 months old B6.Cg-

miR7a1tm1(fl/fl)ms miR7a2tm1(fl/fl)ms miR7btm1(fl/fl)ms Ndor1Tg(UBC-cre/ERT2)1Ejb/Stf and

UBC-cre negative counterpart[39] provided by Prof. Dr. Markus Stoffel, ETH, Zurich, Switzerland (n

= 20-26 per genotype).

Genotyping

Before the animal studies, ciRS-7 KO and miR-7 KO mice were genotyped as before[9,39] except for

the tissue dissociation step. Briefly, ear puncture samples were digested in 50 mM Sodium Hydroxide

solution for one hour at 95 °C. Once equilibrated to room temperature the samples were neutralized

with 1M Tris Hydrochloride pH 8. Two microliters of digested samples were used to run the reaction

of PCR with Taq 2X Master Mix (New England Biolabs (UK) Ltd) following manufacturer protocol

and using the oligonucleotide primers as specified in Supplementary Table S9. The PCR products

were separated on a stained 2.5% or 1.5% Agarose gel for ciRS-7 and miR-7 KO respectively and

run in an electrophoresis chamber. Images were acquired with ChemiDoc Imaging Systems (Bio-Rad

Laboratories, Inc).

Permanent middle cerebral artery surgery (pMCAo)

Permanent middle cerebral artery occlusion (pMCAo) was performed in 3-4 months old

BALB/cOlaHsd male mice as previously described[51]. Briefly, mice were anesthetized using 5%

isoflurane and anesthesia was maintained with 2% isoflurane. Temperature of the animals was

controlled during the surgery with heating blanket and rectal probe (Harvard apparatus). After the

skin incision, the left temporal bone was exposed under the temporal muscle, and 1 mm hole was

drilled on top of bifurcation in the middle cerebral artery (MCA). The dura was gently removed, and

the MCA was lifted and permanently occluded under the bifurcation using a thermocoagulator (Aaron

Medical Industries Inc.). Success of the occlusion was confirmed by cutting the MCA above the

occlusion site. After the occlusion, the temporal muscle was replaced, and the wound was sutured.

The surgery was performed on a total of 20 mice, 3 mice (one per timepoint) died during the surgery

and was then not included.

miR-7 KO induction with tamoxifen

Before tMCAO surgery, miR-7 inducible KO mice were injected with 2mg of Tamoxifen intraperitoneally once per day for 5 consecutive days at the age of 3 to 4 weeks as before[39]. Recombination was induced in both Cre+ and Cre- littermates. Tamoxifen (Sigma-Aldrich, T5648) was resuspended in 90% corn oil (Sigma-Aldrich, C8267) and 10% pure ethanol and dissolved at 56°C for 30 minutes in the dark. After injection the mice were put on soft food diet and their weight was monitored daily. Mice showing a successful genomic recombination in ear puncture samples two weeks after injection were included in the study (Supplementary Figure S5A). At the end of the study the animals were analyzed for miR-7 abrogation in the brain through RT-qPCR (Supplementary Figure S5B).

Transient middle cerebral artery surgery (tMCAO)

Transient middle cerebral artery occlusion (tMCAO) was chosen to continue the in vivo stroke studies as pMCAO in C57BL/6J produces considerably smaller infarcted regions than in the BALB/c background[52,53]. For this reason tMCAO was performed in 3-4 months old male Cdr1as and miR-7 KO and WT mice as previously described[54]. Briefly, induction of anesthesia was performed with 3.5-4% isoflurane in 0.5 L/min of 100% O2-enriched air and maintained at 1-1.5% isoflurane during the surgery. The body temperature maintained at 36 ± 1°C during surgery with a heating pad attached to a rectal probe. A Doppler probe was fixed on the temporal bone over the MCA to monitor blood flow. The right common, external, and internal carotid arteries (CCA, ECA, and ICA) were exposed through an incision in the neck. The arteries were carefully dissected, and a 6-0 silicon coated monofilament of 0.21 mm of diameter (Doccol Corporation) was inserted into the right CCA or left ECA (Cdr1as and mir-7 KO respectively) and led through the ICA until blocking the origin of the MCA. Blood flow blockage was monitored with the Doppler probe, and animals with less than 80%

of blood flow decrease were discarded. After one hour of occlusion, the monofilament was withdrawn

allowing blood reperfusion. Animals that did not show a correct reperfusion measured by laser

doppler were discarded from the study. Incisions were permanently sutured, and the animals were

allowed to recover in a temperature-controlled environment for 24 hours. The surgery was performed

on a total of 32 WT and 25 Cdr1as KO mice, and 36 WT and 26 miR-7 KO. In the ciRS-7 animal

study a total of 7 mice was excluded as: 2 KO and 2 WT mice lacked reperfusion upon surgery, 2

WT and 1 KO mice died during the study. In the miR-7 animal study a total of 12 mice was excluded

as: 3 WT and 2 KO lacked reperfusion upon surgery, 4 WT and 3 KO died during the study.

Magnetic resonance imaging (MRI)

MRI was performed at 1, 3, and 7dpi using a vertical 9.4 T/89 mm magnet (Oxford instrument PLC)

upon anesthesia with 1.8% isoflurane in 30% O_2/70% N_2O. We acquired twelve slices of 0.8mm

thickness per mouse (echo time 40ms, repetition time of 3000ms, matrix size of 128×256 and field

of view $19.2 \times 19.2\,mm^2$) and analyzed the first seven images using the Aedes software

(http://aedes.uef.fi/) for MatLab program (Math-works). Upon definition of the region of interest

(ROI) of contralateral, ipsilateral, and lesion, the lesion volume normalized on oedema was calculated

on the first 7 section as: $Corrected\ lesion\ size\ =\ lesion\ size\ \times\ \Big(1 -$

$\frac{[ipsilateral\ size\ -\ contralateral\ size]}{contralateral\ size}\Big)$. The lesion size is expressed has percentage of lesion on the total

brain size.

Neurological severity score (NSS)

Mice were examined for neurological deficits at baseline and dpi 1, 3, and 7 using a severity scale

comprising the following tests: postural reflex, circling, falling to contralateral side, placement of the

contralateral forelimb during motion, and general state of alertness or consciousness. Deficits were

graded from 0 (normal) to 2 or 3 (severe)[55]. A sum of these scores were used for statistics. Behavioral

assessment was performed by an experimenter blinded to the genotype of mice.

Transcardiac perfusion and sample collection

After anesthesia, mice were perfused transcardially with cold saline solution with heparin 2500 IU/l

(Leo Pharma A/S). In the pMCAo study, the brains were collected and dissected into contralateral

and peri-ischemic cortical regions. In the tMCAO study the brains were cut in 6 coronal sections and

stained with 1% 2,3,5-Triphenyltetrazolium Chloride (TTC) (Sigma-Aldrich) in PBS solution for 5

minutes at 37 °C in agitation before dissection of contralateral and peri-ischemic cortical regions. For

immunohistochemistry staining, brains were collected and fixed in 4% paraformaldehyde solution in

0.1 M phosphate buffer (PB) pH 7.4. After 22 hours of fixation, the brains were transferred in 30%

sucrose in PB buffer solution for 48 hours and then frozen in liquid nitrogen before being stored in -

70 °C.

GFAP immunostaining

Each brain was then cut using a cryostat (Leica Microsystems) into six 20 μm coronal sections 400 μm

apart, collected on Superfrost™ Plus Microscope Slides (ThermoFisher Scientific) and stored in -

70°C until immunostaining. Sections were then rehydrated with phosphate-buffered saline (PBS) pH

7.4 for 10 min and PBS with 0.05% Tween-20 (PBST) (Sigma-Aldrich) for 5 min. Endogenous

peroxidase was blocked by using 0.3 % hydrogen peroxide (H_2O_2) in MeOH for 30 min after which

sections were washed 3 x 5 min in PBST. Non-specific binding was blocked with 10 % normal goat

serum (NGS)(Vector, S-1000) in PBST for 1 h at RT Sections were incubated overnight at RT in

primary antibody rabbit anti-GFAP (Agilent, Dako Z0334,1:500 in 5 % NGS-PBST) and then with

biotinylated secondary antibody anti-rabbit IgG (H+L) (Vector, BA-1000, 1:200 in 5 % NGS-PBST)

for 2 h at RT followed by incubation in ABC reagent (Vector Elite Kit) for 2 h at RT. Sections were

washed 3 x 5 min in PBST before and after the incubations. Nickel-3,3'-diaminobenzidine (Ni-DAB)

solution (0.175M Sodium acetate, 1% Nickel ammonium sulphate, 50mg DAB (Sigma-Aldrich, D-

5905)) with 0.075 % H_2O_2 was used to develop the colour for 6 min stopping the reaction by washing

the sections 2 x 5 min in dH_2O. Sections were then dehydrated in 50 % EtOH, 70 % EtOH, 95 %

EtOH, 100 % EtOH for 2 min in each and 3 x 5 min in xylene followed by mounting the coverslips

with Depex.

Image acquisition & analysis

Six GFAP immunoreactivity light microscope images per mouse were acquired by Leica DM6B-Z

Thunder Imager microscope (Leica Microsystems CMS GmbH) equipped with DMC2900 camera

using 10x magnification. Images were captured using LAS X software (Leica Microsystems CMS

GmbH) with exposure time 5 ms, color-gain mode: R:0-G:0-B:25. The images were quantified from

the peri-ischemic cortex next to the lesion border (1mm) and the corresponding area of the healthy

contralateral hemisphere from 10x images using ImageJ software[56] function "Measure particles". The

results were presented as relative immunoreactive area to the total area analyzed. This part of the

work was carried out with the support of UEF Cell and Tissue Imaging Unit, University of Eastern

Finland, Biocenter Kuopio and Biocenter Finland.

Library preparation tMCAO tissue samples

All the samples RNA were isolated with TRIzol™ Reagent (Invitrogen) as specified above. RNA

samples were treated with TURBO DNA-free™ Kit (Ambion) following the manufacture's

instruction. RNA integrity was assessed through Agilent Bioanalyzer 2100 system with the Agilent

RNA 6000 Nano. The concentration of the samples was established with Qubit™ RNA Extended

range kit (Invitrogen). We generated a library from ciRS-7 WT and KO tMCAO animals contralateral

and peri-ischemic cortices to detect mRNAs changes using CORALL total RNA-Seq Library Prep

Kit (Lexogen GmbH) after ribosomal RNA depletion of 600ng of RNA with RiboCop rRNA

Depletion Kit V1.2 (Lexogen GmbH). All the libraries were generated following the manufacturer's

protocol. After generation, the libraries were quantified with Qubit™ High Sensitivity DNA kit

(Invitrogen) and by qPCR using KAPA Library Quantification Kit for Illumina® Platforms (Roche).

Library size was determined with Agilent Bioanalyzer 2100 system using the Agilent High Sensitivity

DNA Kit. Animal samples were sequenced single-read 75 cycles on NextSeq™ 550.

DATA ANALYSIS

RNA-seq data analysis

Bulk RNA sequences of mRNA and miRNA have been aligned and quantified to the mouse genome

of reference mm10 with the Cdr1as annotation of "chrX:61183248|61186174|.|+" using the and 1.1.0

version of "smrnaseq" applied without clipping and the three prime adapter

"AGATCGGAAGAGCACACGTCT"), while circular sequences using 1.1 CIRIquant[57] set to Read

1 matching the sense strand. Count data were prepared following the workflow defined by Law et

al[58]. We filtered out lowly expressed molecules in any condition using the function "filterByExpr" to

increase the reliability of the mean-variance relationship. We removed the differences between

samples due to the sequencing depth normalizing the count using the trimmed mean of M-values

(TMM)[59] method and applied a log transformation minimizing sum of sample-specific squared

difference to enhance the true positive and negative ratio in the downstream analysis[60]. We checked

for batch effect due to different timings in biological replicates preparation by performing a principal

component analysis and unsupervised consensus clustering with Cola[61]. We identified a batch effect

due to different timings in biological replicates preparation influencing the grouping of the normoxic

and OGD samples in the dataset ciRS-7 KO/WT cortical neurons (Supplementary Table S5). We

corrected the batch effect of this dataset using the negative binomial regression from Combat[62]. We

adjusted the variance between the samples as before[63] through winsorization[64]. We finally created

the design matrix for each pair of conditions to compare (contrast) and performed the differential

expression analysis using limma/edgeR model[58] controlling for the false discovery rate with

Benjamini-Hochberg Procedure[65]. We employed an interaction term in differential expression

analysis to assess whether the OGD response varied between ciRS-7 KO and WT backgrounds. Our

analysis using the interaction term identified only one significant DE gene, Gm2004, with a logFC

change of 9.631571, aligning with our expectations (Supplementary Table S6). This outcome

emphasized that the lack of ciRS-7 did not modulate the OGD response per se and that our

backgrounds remained consistent.

Ingenuity Pathway Analysis (IPA) and Functional enrichment analysis

We uploaded the differentially expressed genes of each contrast to QIAGEN IPA (QIAGEN Inc.,

https://digitalinsights.qiagen.com/IPA)[66] and Metascape[67] for Ingenuity pathway analysis and

functional enrichment analysis, respectively. The analysis has been performed with default

parameters and IPA's background was composed of non-differentially expressed genes.

Deconvolution analysis with scRNAseq

Deconvolution aims to estimate the proportions of different cell types within a mixed population in

bulk RNA sequencing data using expression profiles from individual cells of scRNA-seq data and

represents them in a composite expression profile. The term "contribution" is used to describe the

proportion of gene expression attributed to a specific cell type, such as astrocytes or neurons, within

the composite profile. We exploited cell-type specific gene expression from external single-cell RNA

sequencing (scRNA-seq) data to define the cell subpopulations composing our cortical neuron culture

(Supplementary Table S2) and ischemic stroke animal tissue (Supplementary Table S7) bulk RNA

sequencing datasets. The scRNA-seq dataset of Loo et al.[68] of the mouse cerebral cortex at embryonic

day 14.5 was provided to SCDC[69] for performing the deconvolution of our gene expression matrix

with cortical neuron samples in normoxic conditions. The scRNA-seq data of Zeisel at al.[70] composed

of murine cerebral cortex samples from p25 to p60 was instead provided to MuSiC[71] to perform the

deconvolution of our gene expression matrix with contralateral and peri-ischemic samples of wild-

type and ciRS-7 KO ischemic stroke animals (Supplementary Table S7) as this method is designed

to work with multi-subject scRNA-seq dataset.

GRO-seq data analysis

A summary of all GRO-seq samples used to quantify pri-miRNA expression levels are presented in

Supplementary Table S3. Raw reads for public GRO-seq data were acquired from the GEO database.

GRO-Seq reads were trimmed using the HOMER v4.3 (http://homer.salk.edu/homer)[72] software to

remove A-stretches originating from the library preparation. From the resulting sequences, those

shorter than 25 bp were discarded. The quality of raw sequencing reads was controlled using the

FastQC tool (http://www.bioinformatics.babraham.ac.uk/projects/fastqc)[73] and bases with poor

quality scores were trimmed using the FastX toolkit (http://hannonlab.cshl.edu/fastx_toolkit/). Reads

were aligned to mouse mm9 reference genome using the Bowtie[74] version bowtie-0.12.7. Up to two

mismatches and up to three locations were accepted per read and the best alignment was reported.

The data was used to create Tag Directories using HOMER. To optimize coverage, a combined tag

directory representing all samples of a given cell type (under one or several GSE numbers) was

created and used for pri-miRNA quantification using the 'analyzeRepeats.pl' command and '-rpkm -

strand +' options".

miRWalk analysis

We associated the significantly deregulated miRNAs and differentially expressed genes of wild-type

and ciRS-7 KO cortical neurons subjected to OGD (Supplementary Table S2, Supplementary Table

S5). We filtered miRNA-Target interactions obtained from 3.0 miRWalk[75,76] database with 99%

probability and located in the 3' UTR region and included Oip5os1 (Cyrano) as target of mmu-miR-

7a-5p[8]. We linked the differentially expressed miRNAs and genes of each contrast by anticorrelation

(e.g. miRNA with positive log fold change and significant adjusted probability value is associated to

genes with negative log fold change, significant adjusted probability value and targets in miRWalk

of the miRNA) focusing on mmu-miR-7a-5p.

CDF generation

We tested the assumption of anticorrelation between mmu-miR-7a-5p and its miRWalk gene targets

using the empirical Cumulative Distribution Function (eCDF). For each contrast, we compared the

eCDF (control function) obtained from the values of log fold change of the non-target genes against

the one of the miRNA's targets and tested their equality with the Kolmogorov–Smirnov test[77]. Genes

characterized by longer 3' UTRs often possess a greater number of potential miRNA binding sites

and may display distinct expression characteristics owing to their intrinsically less stable nature. To

address this concern, we have incorporated a normalization approach inspired by Kleaveland *et al.*

(2018)[8] in which the relationship between fold change and 3' UTR length is fitted to a linear function

using the 3' UTR length annotated by Eichhorn *et al.* (2014)[78]. The original fold changes are adjusted

by subtracting predicted values, assuming these values exclusively represent expected fold changes

due to 3' UTR length. This subtraction is done as absolute values to adjust the magnitude of the

original fold change, based on the assumption that 3' UTR length, as a gene-specific attribute, is

independent of class order. The subtraction aims to eliminate the fold change component attributed

to 3' UTR length and retain the component associated with miRNA targeting. To determine the

magnitude and direction of the shift of the eCDF of the targets in respect of the non-targets, we

measured the area between the two curves following this formula: $\int \{F_Y(t) - F_X(t)\}\, dt$ and the

Wasserstein distance[79]. Both the considered targets and non-targets genes passed the filtering by

expression, count normalization and participated at the differential expression analysis.

CLIP-seq analysis

We collected the HITS-CLIP dataset of Argonaute 2 in pyramidal excitatory neurons produced by Tan et al.[40] (GSE211552) and replicated the original analysis to map and annotate the genomic regions with a significant read cluster (peak) (Supplementary Table S8). Briefly, we performed a quality control with FastQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc)[73] and applied trimming to remove bases after 50 base pairs due to low quality. We applied Cutadapt[80] to remove adapters and over-represented sequences. We selected reads of at least 24 nucleotides in length and quality score higher than 20 for each nucleotide. Reads were aligned to the mouse reference genome mm10 using Burrows-Wheeler aligner[81] with default parameters, then the mapped reads were expanded by 30 nucleotides on each side. We considered only reads present in at least 2 experimental replicates and clusters composed of at least 10 overlapping reads. Bowtie[82] was used to map all occurrences of sequences complementary to the 6mer seed sequence (positions 2-7 of the mature miRNA) of mouse miRNAs from miRBase V21 (https://www.mirbase.org/) to the mouse genome mm10, allowing no mismatches. Strand specific intersection was done with the filtered clusters in the 3' UTR, 5' UTR, coding sequence (CDS) and intron regions from RefSeq gene definitions, as well as antisense matches to RefSeq genes[83]. Strand specific intersection was also done to circRNAs from circBase[84]. This generated a genome-wide list with predicted miRNA target sites backed by the detected Ago2 HITS-CLIP clusters for all mouse miRNAs. A subset including all miR-7a-5p targets is supplied as Supplementary Table S8. Only 3' UTR target sites were used to define miRNA mediated regulation. The full list of all predicted miRNA target sites was uploaded to GEO, as described in the Data Availability section. We conducted quality control on the CLIP-seq output by using MEME-ChIP[85] (https://meme-suite.org/meme/tools/meme-chip), a dedicated tool for motif enrichment analysis. This tool allowed us to systematically explore the enrichment of miRNA binding motifs within all identified CLIP-seq 3' UTR clusters. The results of this analysis are provided in Supplementary Table S8.

1007

Statistical analysis

Graphs and statistical analysis were performed in GraphPad Prism 9. Every Figure legend reports

parameters of replicates (n), statistical test and p-value obtained. Where not specified, p-value was

not statistically significant (p-value > 0.05). We refer to n in animal study as single biological

replicate (mouse) and in cortical neuron as technical replicates in the same batch, the experiment has

been performed in three independent biological replicates.

1014

Data availability

The data produced and analyzed in this publication have been deposited in NCBI's Gene Expression

Omnibus[86] and are accessible through GEO Series accession number GSE213179, GSE213067,

GSE213177, GSE211552 and GSE215210. The code developed to analyze the data produced in this

study is deposited in Zenodo at https://zenodo.org/records/10489728.

ACKNOWLEDGEMENTS

We thank Nicholas Downes, Mirka Tikkanen, Reetta Vuolteenaho, Miia Salo, Irina Belaia, Nikita

Mikhailov, and Gianluca Como for their support. We acknowledge UEF Biocenter Kuopio In Vitro

and Ex Vivo Electrophysiology Core Facility, National Virus Vector Laboratory Service, Cell and

Tissue Imaging Unit, Lab Animal Centre (Nina Liimatainen and Hanna Miettinen), Biocenter Oulu

Transgenic and Tissue Phenotyping Core, Oulu Laboratory Animal Centre Research Infrastructure,

Translational Cell Biology Core at Kontinkangas Campus, and the Faculty of Biochemistry and

Molecular Medicine of the University of Oulu. This project received co-funding from the Horizon

2020 Framework Programme of the European Union (Marie Skłodowska Curie grant agreement No

740264), as well as the Academy of Finland, Finnish Cultural Foundation, Finnish Foundation for

Cardiovascular Research, Instrumentarium Science Foundation, Inkeri and Mauri Vänskä

AUTHOR CONTRIBUTIONS

F.S. and T.M. conceived and planned the study, with intellectual contributions from T.H., J.J., R. Giniatullin, and D.T. In vivo experiments involved F.S., V.S., P.K., I.U., J.J., H.D., C.P., N.V, and J. Koistinaho. In vitro experiments involved F.S., V.S., D.M.T., M.G.B., R. Giniatullina, N.K., E.G., S.YH. RNA-seq experiments involved F.S., A.H.S., J.S., and J. Kjems, with analysis performed by L.G. Transgenic animals and study assistance were provided by M.L., M.P., M.S., N.R. HITS-CLIP-seq was generated and analyzed by M.V. and A.S. GRO-seq was analyzed by M.K. Sample preparation at Oulu University was done by R.H., F.S., V.S., S.K., and A.H. F.S. and T.M., wrote the manuscript with input from T.H., J.J., R. Giniatullin, J. Kjems, and critical feedback from all authors.

CONFLICT OF INTEREST

The authors declare no competing interests.

REFERENCES

1. Mehler, M. F. & Mattick, J. S. Noncoding RNAs and RNA editing in brain development, functional diversification, and neurological disease. *Physiol Rev* **87**, 799–823 (2007).

2. Lewis, B. P., Burge, C. B. & Bartel, D. P. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* **120**, 15–20 (2005).

3. Mattick, J. S. & Makunin, I. V. Small regulatory RNAs in mammals. *Hum Mol Genet* **14 Spec No 1**, R121-132 (2005).

4.   Fatica, A. & Bozzoni, I. Long non-coding RNAs: new players in cell differentiation and

development. *Nat Rev Genet* **15**, 7–21 (2014).

5.   Hansen, T. B. *et al.* Natural RNA circles function as efficient microRNA sponges. *Nature* **495**,

384–388 (2013).

6.   Kristensen, L. S. *et al.* The biogenesis, biology and characterization of circular RNAs. *Nat Rev*

Genet **20**, 675–691 (2019).

7.   Hansen, T. B. *et al.* miRNA-dependent gene silencing involving Ago2-mediated cleavage of a

circular antisense RNA: miRNA mediated cleavage of circular antisense RNA. *The EMBO*

Journal **30**, 4414–4422 (2011).

8.   Kleaveland, B., Shi, C. Y., Stefano, J. & Bartel, D. P. A Network of Noncoding Regulatory

RNAs Acts in the Mammalian Brain. *Cell* **174**, 350-362.e17 (2018).

9.   Piwecka, M. *et al.* Loss of a mammalian circular RNA locus causes miRNA deregulation and

affects brain function. *Science* **357**, eaam8526 (2017).

10. Memczak, S. *et al.* Circular RNAs are a large class of animal RNAs with regulatory potency.

Nature **495**, 333–338 (2013).

11. Olney, J. W., Rhee, V. & Ho, O. L. Kainic acid: a powerful neurotoxic analogue of glutamate.

Brain Res **77**, 507–512 (1974).

12. Dirnagl, U., Iadecola, C. & Moskowitz, M. A. Pathobiology of ischaemic stroke: an integrated

view. *Trends Neurosci* **22**, 391–397 (1999).

13. Tiedt, S. & Dichgans, M. Role of Non-Coding RNAs in Stroke. *Stroke* **49**, 3098–3106 (2018).

14. Leung, A. K. L. & Sharp, P. A. MicroRNA Functions in Stress Responses. *Mol Cell* **40**, 205–

215 (2010).

15. Denzler, R. *et al.* Impact of MicroRNA Levels, Target-Site Complementarity, and

Cooperativity on Competing Endogenous RNA-Regulated Gene Expression. *Mol Cell* **64**, 565–

579 (2016).

16. Salmena, L., Poliseno, L., Tay, Y., Kats, L. & Pandolfi, P. P. A ceRNA hypothesis: the Rosetta

Stone of a hidden RNA language? *Cell* **146**, 353–358 (2011).

17. Franco-Zorrilla, J. M. *et al.* Target mimicry provides a new mechanism for regulation of

microRNA activity. *Nat Genet* **39**, 1033–1037 (2007).

18. Zhang, Y. *et al.* Circular RNAs in the Regulation of Oxidative Stress. *Frontiers in*

Pharmacology **12**, (2021).

19. Loo, L. *et al.* Single-cell transcriptomic analysis of mouse neocortical development. *Nat*

Commun **10**, 134 (2019).

20. Semenza, G. L. Hypoxia-inducible factor 1: control of oxygen homeostasis in health and

disease. *Pediatr Res* **49**, 614–617 (2001).

21. Zhang, J., Chen, S., Yang, J. & Zhao, F. Accurate quantification of circular RNAs identifies

extensive circular isoform switching events. *Nat Commun* **11**, 90 (2020).

22. Loppi, S. *et al.* Peripheral inflammation preceeding ischemia impairs neuronal survival through

mechanisms involving miR-127 in aged animals. *Aging Cell* **20**, e13287 (2021).

23. Step, S. E. *et al.* Anti-diabetic rosiglitazone remodels the adipocyte transcriptome by

redistributing transcription to PPARγ-driven enhancers. *Genes Dev* **28**, 1018–1028 (2014).

24. Meng, F.-L. *et al.* Convergent transcription at intragenic super-enhancers targets AID-initiated

genomic instability. *Cell* **159**, 1538–1548 (2014).

25. Min, I. M. *et al.* Regulating RNA polymerase pausing and transcription elongation in

embryonic stem cells. *Genes & Development* **25**, 742 (2011).

26. Fang, B. *et al.* Circadian enhancers coordinate multiple phases of rhythmic gene transcription in

vivo. *Cell* **159**, 1140–1152 (2014).

27. Hah, N. *et al.* Inflammation-sensitive super enhancers form domains of coordinately regulated

enhancer RNAs. *Proceedings of the National Academy of Sciences* **112**, E297–E302 (2015).

28. Heinz, S. *et al.* Effect of natural genetic variation on enhancer selection and function. *Nature*
503, 487–492 (2013).

29. Escoubet-Lozach, L. *et al.* Mechanisms Establishing TLR4-Responsive Activation States of
Inflammatory Response Genes. *PLOS Genetics* **7**, e1002401 (2011).

30. Ji, Z. *et al.* Transcriptional activity regulates alternative cleavage and polyadenylation. *Mol Syst*
Biol **7**, 534 (2011).

31. Zhang, F. *et al.* Enhancer-bound LDB1 regulates a corticotrope promoter-pausing repression
program. *Proceedings of the National Academy of Sciences* **112**, 1380–1385 (2015).

32. Telese, F. *et al.* LRP8-Reelin-regulated Neuronal (LRN) Enhancer Signature Underlying
Learning and Memory Formation. *Neuron* **86**, 696–710 (2015).

33. Sticht, C., Torre, C. D. L., Parveen, A. & Gretz, N. miRWalk: An online resource for prediction
of microRNA binding sites. *PLOS ONE* **13**, e0206239 (2018).

34. Dweep, H. & Gretz, N. miRWalk2.0: a comprehensive atlas of microRNA-target interactions.
Nat Methods **12**, 697–697 (2015).

35. Zeisel, A. *et al.* Molecular Architecture of the Mouse Nervous System. *Cell* **174**, 999-1014.e22
(2018).

36. Yoon, J. S. *et al.* Spatiotemporal Protein Atlas of Cell Death-Related Molecules in the Rat
MCAO Stroke Model. *Exp Neurobiol* **27**, 287–298 (2018).

37. Kabaria, S. *et al.* MicroRNA-7 activates Nrf2 pathway by targeting Keap1 expression. *Free*
Radic Biol Med **89**, 548–556 (2015).

38. Chaudhuri, A. D., Kabaria, S., Choi, D. C., Mouradian, M. M. & Junn, E. MicroRNA-7
Promotes Glycolysis to Protect against 1-Methyl-4-phenylpyridinium-induced Cell Death. *J*
Biol Chem **290**, 12425–12434 (2015).

39. LaPierre, M. P., Lawler, K., Godbersen, S., Farooqi, I. S. & Stoffel, M. MicroRNA-7 regulates

melanocortin circuits involved in mammalian energy homeostasis. *Nat Commun* **13**, 5733

(2022).

40. Tan, C. L. *et al.* MicroRNA-128 Governs Neuronal Excitability and Motor Behavior in Mice.

Science **342**, 1254–1258 (2013).

41. Chen, H. *et al.* miR-7 and miR-214 are specifically expressed during neuroblastoma

differentiation, cortical development and embryonic stem cells differentiation, and control

neurite outgrowth in vitro. *Biochemical and Biophysical Research Communications* **394**, 921–

927 (2010).

42. Rybak-Wolf, A. *et al.* Circular RNAs in the Mammalian Brain Are Highly Abundant,

Conserved, and Dynamically Expressed. *Mol Cell* **58**, 870–885 (2015).

43. Noh, K.-M. *et al.* Blockade of calcium-permeable AMPA receptors protects hippocampal

neurons against global ischemia-induced death. *Proc Natl Acad Sci U S A* **102**, 12230–12235

(2005).

44. Yao, G.-Y. *et al.* Ischemic postconditioning confers cerebroprotection by stabilizing VDACs

after brain ischemia. *Cell Death Dis* **9**, 1–15 (2018).

45. Zhao, J. & Wang, B. MiR-7-5p Enhances Cerebral Ischemia-Reperfusion Injury by Degrading

sirt1 mRNA. *J Cardiovasc Pharmacol* **76**, 227–236 (2020).

46. Mehta, S. L. *et al.* CDR1as regulates α-synuclein-mediated ischemic brain damage by

controlling miR-7 availability. *Mol Ther Nucleic Acids* **31**, 57–67 (2023).

47. Bushati, N. & Cohen, S. M. microRNA functions. *Annu Rev Cell Dev Biol* **23**, 175–205 (2007).

48. Miska, E. A. *et al.* Most Caenorhabditis elegans microRNAs are individually not essential for

development or viability. *PLoS Genet* **3**, e215 (2007).

49. Korvenlaita, N. *et al.* Dynamic release of neuronal extracellular vesicles containing miR-21a-5p

is induced by hypoxia. *Journal of Extracellular Vesicles* **12**, 12297 (2023).

50. Tiihonen, J. *et al.* Sex-specific transcriptional and proteomic signatures in schizophrenia. *Nat*

Commun **10**, 3933 (2019).

51. Kumar, M. & Katyal, A. Data on retinoic acid and reduced serum concentration induced

differentiation of Neuro-2a neuroblastoma cells. *Data in Brief* **21**, 2435–2440 (2018).

52. Korhonen, P. *et al.* Immunomodulation by interleukin-33 is protective in stroke through

modulation of inflammation. *Brain, Behavior, and Immunity* **49**, 322–336 (2015).

53. Majid, A. *et al.* Differences in vulnerability to permanent focal cerebral ischemia among 3

common mouse strains. *Stroke* **31**, 2707–2714 (2000).

54. Marques, S. M. *et al.* Genetic background determines mouse strain differences in inflammatory

angiogenesis. *Microvasc Res* **82**, 246–252 (2011).

55. Shahjouei, S. *et al.* Middle Cerebral Artery Occlusion Model of Stroke in Rodents: A Step-by-

Step Approach. *J Vasc Interv Neurol* **8**, 1–8 (2016).

56. Senda, D. M., Franzin, S., Mori, M. A., de Oliveira, R. M. W. & Milani, H. Acute, post-

ischemic sensorimotor deficits correlate positively with infarct size but fail to predict its

occurrence and magnitude after middle cerebral artery occlusion in rats. *Behav Brain Res* **216**,

29–35 (2011).

57. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image

analysis. *Nat Methods* **9**, 671–675 (2012).

58. Zhang, J., Chen, S., Yang, J. & Zhao, F. Accurate quantification of circular RNAs identifies

extensive circular isoform switching events. *Nat Commun* **11**, 90 (2020).

59. Law, C. W. *et al.* RNA-seq analysis is easy as 1-2-3 with limma, Glimma and edgeR. *F1000Res*

5, (2016).

60. Robinson, M. D. & Oshlack, A. A scaling normalization method for differential expression

analysis of RNA-seq data. *Genome Biol* **11**, R25 (2010).

61. Zhang, Z. *et al.* Novel Data Transformations for RNA-seq Differential Expression Analysis. *Sci*

Rep **9**, 4820 (2019).

62. Gu, Z., Schlesner, M. & Hübschmann, D. cola: an R/Bioconductor package for consensus

partitioning through a general framework. *Nucleic Acids Research* **49**, e15 (2021).

63. Zhang, Y., Parmigiani, G. & Johnson, W. E. ComBat-seq: batch effect adjustment for RNA-seq

count data. *NAR Genomics and Bioinformatics* **2**, lqaa078 (2020).

64. Allswede, D. M. *et al.* Complement Gene Expression Correlates with Superior Frontal Cortical

Thickness in Humans. *Neuropsychopharmacol.* **43**, 525–533 (2018).

65. Winsor, C. P., Hastings, C., Mosteller, F. & Tukey, J. W. Low Moments for Small Samples: A

Comparative Study of Order Statistics. *The Annals of Mathematical Statistics* (1947)

doi:10.1214/aoms/1177730388.

66. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*

(Methodological) **57**, 289–300 (1995).

67. Krämer, A., Green, J., Pollard, J. & Tugendreich, S. Causal analysis approaches in Ingenuity

Pathway Analysis. *Bioinformatics* **30**, 523–530 (2014).

68. Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of systems-

level datasets. *Nat Commun* **10**, 1523 (2019).

69. Loo, L. *et al.* Single-cell transcriptomic analysis of mouse neocortical development. *Nat*

Commun **10**, 134 (2019).

70. Dong, M. *et al.* SCDC: bulk gene expression deconvolution by multiple single-cell RNA

sequencing references. *Briefings in Bioinformatics* **22**, 416–427 (2021).

71. Zeisel, A. *et al.* Molecular Architecture of the Mouse Nervous System. *Cell* **174**, 999-1014.e22

(2018).

72. Wang, X., Park, J., Susztak, K., Zhang, N. R. & Li, M. Bulk tissue cell type deconvolution with

multi-subject single-cell expression reference. *Nat Commun* **10**, 380 (2019).

73. Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-

regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576–589

(2010).

74. Blankenberg, D. *et al.* Manipulation of FASTQ data with Galaxy. *Bioinformatics* **26**, 1783

(2010).

75. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. *Genome Biology* **10**, R25 (2009).

76. Dweep, H. & Gretz, N. miRWalk2.0: a comprehensive atlas of microRNA-target interactions.

Nat Methods **12**, 697–697 (2015).

77. Sticht, C., Torre, C. D. L., Parveen, A. & Gretz, N. miRWalk: An online resource for prediction

of microRNA binding sites. *PLOS ONE* **13**, e0206239 (2018).

78. Smirnov, N. Table for Estimating the Goodness of Fit of Empirical Distributions. *The Annals of*

Mathematical Statistics **19**, 279–281 (1948).

79. Eichhorn, S. W. *et al.* mRNA destabilization is the dominant effect of mammalian microRNAs

by the time substantial repression ensues. *Mol Cell* **56**, 104–115 (2014).

80. Kantorovich, L. V. Mathematical Methods of Organizing and Planning Production.

Management Science **6**, 366–422 (1960).

81. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal **17**, 10–12 (2011).

82. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform.

Bioinformatics **25**, 1754–1760 (2009).

83. Trapnell, C., Pachter, L. & Salzberg, S. L. TopHat: discovering splice junctions with RNA-Seq.

Bioinformatics **25**, 1105–1111 (2009).

84. Kd, P., T, T. & Dr, M. NCBI Reference Sequence (RefSeq): a curated non-redundant sequence

database of genomes, transcripts and proteins. *Nucleic acids research* **33**, (2005).

85. Glažar, P., Papavasileiou, P. & Rajewsky, N. circBase: a database for circular RNAs. *RNA* **20**,

1666–1670 (2014).

86. Machanick, P. & Bailey, T. L. MEME-ChIP: motif analysis of large DNA datasets.

Bioinformatics **27**, 1696–1697 (2011).

1228