Sugar restriction and blood ingestion shape divergent immune defense trajectories in the mosquito *Aedes*

*aegypti*

Dom Magistrado[1], Noha K. El-Dougdoug[1,2], Sarah M. Short[1*]

[1]Department of Entomology, College of Food, Agricultural, and Environmental Sciences, The Ohio State

University, Columbus, Ohio, United States of America

[2]Department of Botany and Microbiology, Faculty of Science, Benha University, Benha, Egypt

* Corresponding author

Email: short.343@osu.edu (SMS)

# Abstract

Immune defense is comprised of 1) resistance: the ability to reduce pathogen load, and 2) tolerance: the ability to limit the disease severity induced by a given pathogen load. The study of tolerance in the field of animal immunity is fairly nascent in comparison to resistance. Consequently, studies which examine immune defense comprehensively (i.e., considering both resistance and tolerance in conjunction) are uncommon, despite their exigency in achieving a thorough understanding of immune defense. Furthermore, understanding tolerance in arthropod disease vectors is uniquely relevant, as tolerance is essential to the cyclical transmission of pathogens by arthropods. Here, we tested the effect(s) of dietary sucrose concentration (high or low) and blood meal (present or absent) on resistance and tolerance to *Escherichia coli* infection in the yellow fever mosquito *Aedes aegypti*. Resistance and tolerance were measured concurrently and at multiple timepoints. We found that both blood and sucrose affected resistance. Mosquitoes from the low sugar treatment displayed enhanced resistance at all timepoints post-infection compared to those from the high sugar treatment. Additionally, blood-fed mosquitoes showed enhanced resistance compared to non-blood-fed mosquitoes, but only on day 1 post-infection. Sucrose had no effect on tolerance, but the effect of blood was significant and dynamic across time. Specifically, we show that consuming blood prior to infection ameliorates a temporal decline in tolerance that mosquitoes experience when provided with only sugar meals. Taken together, our findings indicate that different dietary components can have unique and sometimes temporally dynamic impacts on resistance and tolerance. Finally, our findings 1) highlight the value of experimental and analytical frameworks which consider the explicit testing of effects on both resistance and tolerance as separate, but equally important, components of immune defense, and 2) underscore the importance of including a temporal component in studies of immune defense.

# Introduction

An organism's response to infection (i.e., immune defense) is comprised of both resistance and tolerance (1,2). Resistance is defined as the ability to reduce the number of pathogens inside the body, while tolerance is defined as the ability to limit the impact of the infection on host fitness. Both strategies are critical to surviving infection. Resistance and tolerance were first conceptualized by botanists in the late 1800s (3), and plant biologists have obtained major insights into both resistance and tolerance in years since. In contrast, scientists studying animal hosts have focused disproportionately on resistance until recently (2,4,5). As a result, immune tolerance in animals is a nascent, but quickly developing, field of study. Tolerance strategies likely include tactics such as repair of pathogen-induced tissue damage, detoxification of pathogen by-products, limitation of immune response-mediated self-injury (i.e. immunopathology), and general homeostasis promotion during and following infection – all strategies that promote the health of the host without necessarily affecting pathogen levels (2,6–9). Ecological immunology posits that resistance and tolerance, as components of host defense, are costly and should only be employed when the benefits outweigh the costs (10). Therefore, a host's balance between resistance and tolerance is likely important from a resource limitation perspective, as an inappropriate balance between resistance and tolerance may lead to an undesirable infection outcome. For example, a strong resistance response may result in complete pathogen elimination, but such an outcome could leave inadequate energy reserves for repairing infection-induced damages. If this results in reduced lifetime reproductive fitness, it would not be a successful strategy.

While an appropriate equilibrium between resistance and tolerance is important in all host-pathogen systems, it has unique relevance in hematophagous arthropods that transmit pathogens. Arthropod-borne pathogens are highly diverse, and include viruses, bacteria, protozoan parasites, and filarial worms. Diseases caused by these pathogens comprise more than 17% of all global infectious diseases (11), and thus impose a massive public and veterinary health burden. Arthropod-borne pathogens are ingested by the arthropod when it takes a blood meal from an infected vertebrate host. Because transmission is

categorically dependent upon the arthropod's ability to tolerate infection long enough to spread the

pathogen to a subsequent vertebrate host, studying immune tolerance is critical to the ability to understand

and address arthropod-borne pathogen transmission. We therefore explored the effect of diet on resistance

and tolerance to infection concurrently in the yellow fever mosquito, *Ae. aegypti*, which transmits

multiple human pathogens including dengue virus and Zika virus.

Female adult mosquitoes have evolved to consume two meal types: 1) nectar, which is rich in sugar, and

2) blood, which is rich in protein and necessary for egg production in most species (12,13). Consumption,

storage, and digestion differ appreciably between the two meal types. For example, sugar meals are stored

in the ventral diverticulum (crop) while blood meals bypass the crop and are directly sent to the midgut.

Sugar meals can be stored in the crop until needed for energy-intensive activities such as flight while

blood digestion typically begins within a few hours of feeding (14). Blood digestion, unlike sucrose

digestion, is also associated with significant physiological stress due to rapid shifts in temperature and

pH, gut distension, heme toxicity resulting from hemoglobin digestion, and midgut redox stress (15–21).

Diet has previously been implicated in immune defense in arthropods. For example, lower dietary sugar

concentrations have been shown to enhance resistance to bacterial infection in *Drosophila melanogaster*

and resistance to *Plasmodium* infection in *Anopheles stephensi* (21,22). Effects of diet on tolerance have

been explored as well. For example, genotype interacts with diet to impact tolerance to bacterial infection

in *D. melanogaster* (22). Additionally, a study that investigated the effects of a blood meal on bacterial

infection in *Ae. aegypti* found that blood impacts resistance and tolerance early in infection, but they also

found that the effects were dose-dependent (23). In the present study, we tested the effects of two adult

female *Ae. aegypti* diet components, 1) low/high sucrose diet concentration and 2) the ingestion of a

blood meal, on resistance and tolerance to bacterial infection. We measured resistance and tolerance

concurrently at multiple timepoints across a 5-day infection timecourse, allowing us to examine each

treatment group's resistance/tolerance trajectory, as well as test for any potential interactions between the

two components of the mosquito's diet in affecting immune defense. Our results contribute to the

understanding of how arthropods of medical importance withstand pathogen infection, and more

specifically, elucidate effects of blood feeding and sugar feeding on immune defense that may be

conserved amongst other hematophagous arthropod vectors.

# Results

We investigated the effect of dietary sucrose concentration and blood feeding on resistance and tolerance

to infection over time. To accomplish this, we exposed female *Ae. aegypti* mosquitoes to four different

diet treatments: 10% sucrose alone, 10% sucrose + blood meal, 1% sucrose alone, and 1% sucrose +

blood meal. We then infected them with *E. coli* (S17) and measured bacterial load and survival at

multiple timepoints post-infection (Fig 1). The resultant dataset comprises seven variables (Table 1) and

was used to build multiple models describing the effect(s) of both Blood and Sucrose on resistance and

tolerance to bacterial infection across time.

107
**Fig 1. Experimental Design Schematic.** We reared adult female *Ae. aegypti* on four experimental diets: 10%
sucrose alone, 10% sucrose + blood meal, 1% sucrose alone, and 1% sucrose + blood meal. We then infected
individuals with fluorescent *E. coli* (S17) via intrathoracic microinjection. Infected mosquitoes were split into
monitoring groups for survival and bacterial load. Survival and bacterial load measurements were obtained in
parallel at 1, 3, and 5 days post-infection (dpi). The resultant dataset was used to build models describing the
effect(s) of blood and/or sucrose diets on resistance and tolerance simultaneously (Table 1).

**Table 1. Description of terms included in models.**

| Variable | Type | Description |
|---|---|---|
| Survival (Alive, Dead) | Weighted proportion following a binomial distribution | Proportion of mosquitoes alive, weighted for sample size |
| Blood | Categorical – binary | Mosquitoes either received or did not receive a blood meal |
| Sucrose | Categorical – binary | Mosquitoes received either 1% sucrose or 10% sucrose |
| Day | Discrete | Day (1, 3, or 5) post-infection at which measurements were taken |
| Bacterial Load | Continuous | Median Colony Forming Units per mosquito at time of sampling |

# Dietary sucrose and blood affect resistance

## Lower dietary sucrose increases resistance

We measured resistance to bacterial infection by testing the effects of Sucrose and Blood on Bacterial Load at 1, 3, and 5 days post-infection (dpi). We first built a model using Sucrose, Blood, and Day as predictor variables and Bacterial Load as the response variable, then assessed main effects as well as all potential interactions. We performed backward elimination to identify significant model terms and found that both Sucrose (Table 2: Model A, $p_{Sucrose} = 4.57 \times 10^{-5}$) and Day (Table 2: Model A, $p_{Day} = 2.92 \times 10^{-4}$) significantly affect resistance to infection. Females fed 1% sucrose had significantly lower bacterial loads (and therefore higher resistance) than females fed 10% sucrose, and bacterial load decreased over time for both treatment groups (Fig 2). Furthermore, pairwise comparisons between sucrose treatments on each individual day showed that 1% sucrose exposure significantly enhanced resistance to infection at all days post-infection. (Table 3: Model B (Day 1), $p_{Sucrose} = 0.008$; Model C (Day 3), $p_{Sucrose} = 0.005$; Model D (Day 5), $p_{Sucrose} = 0.048$).

**Table 2. Across-Days Resistance Model A and Summary Output.**

| Resistance Model A (Across Days): $Bacterial\ Load = \beta_0 + \beta_1(Day) + \beta_2(Sucrose) + \epsilon$ | | | | |
|---|---|---|---|---|
| Summary Output: | | | | |
| | Bacterial Load | | | |
| *Predictors* | *Estimate* | *SE* | *Statistic* | *p-value* |
| (Intercept) | 4.01 | 0.98 | 4.10 | 1.42e-4 |
| Day | -1.04 | 0.27 | -3.88 | **2.92e-4** |
| Sucrose | 3.81 | 0.86 | 4.44 | **4.57e-5** |
| Observations | 56 | | | |

Resistance Model A is a linear model testing the effect(s) of Blood, Sucrose, and Day on Bacterial Load.

**Fig 2. Mosquitoes fed 1% sucrose had significantly lower bacterial loads compared to mosquitoes fed 10%**
**sucrose.** Boxplots were constructed using median bacterial loads for females fed 1% sucrose and 10% sucrose at 1,
3, and 5 dpi. Each median value was calculated from four individuals, and point error bars show the interquartile
range. Asterisks represent significant differences in bacterial load between sucrose treatments at each time post-
infection (** $p < 0.01$, * $p < 0.05$). Data were collected from a total of five replicate experiments.

## Blood increases resistance at 1 dpi only

Blood feeding also significantly affected resistance to infection, but only at 1 dpi, at which blood-fed

females had significantly lower bacterial loads compared to non-blood-fed females (Table 3: Model B, p

$_{Blood}$ = 0.038; Fig 3).

**Table 3. Day-Specific Resistance Models B-D and Summary Outputs.**

Resistance Model B (Day 1): $\boldsymbol{Bacterial\ Load = \beta_0 + \beta_1(Sucrose) + \beta_2(Blood) + \epsilon}$

Resistance Model C (Day 3): $\boldsymbol{Bacterial\ Load = \beta_0 + \beta_1(Sucrose) + \epsilon}$

Resistance Model D (Day 5): $\boldsymbol{Bacterial\ Load = \beta_0 + \beta_1(Sucrose) + \epsilon}$

Summary Outputs:

| | Resistance Model B (Day 1) | | | | Resistance Model C (Day 3) | | | | Resistance Model D (Day 5) | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | Bacterial Load | | | | Bacterial Load | | | | Bacterial Load | | | |
| *Predictors* | *Est.* | *SE* | *Stat.* | *p-value* | *Est.* | *SE* | *Stat.* | *p-value* | *Est.* | *SE* | *Stat.* | *p-value* |
| (Intercept) | 4.40 | 1.41 | 3.11 | 0.006 | 0.22 | 1.04 | 0.21 | 0.836 | 0.15 | 0.45 | 0.34 | 0.742 |
| Sucrose | 4.86 | 1.63 | 2.98 | **0.008** | 4.70 | 1.47 | 3.19 | **0.005** | 1.39 | 0.64 | 2.17 | **0.048** |
| Blood | -3.67 | 1.63 | -2.25 | **0.038** | | | | | | | | |
| Observations | 20 | | | | 20 | | | | 16 | | | |

Resistance Models B-D are linear models testing the effect(s) of Blood and Sucrose on Bacterial Load at each dpi.

**Fig 3. Blood-fed mosquitoes had significantly lower bacterial loads compared to non-blood-fed mosquitoes at 1 dpi.** Boxplots were constructed using plotted points showing median (n = 4) bacterial load for blood-fed and non-blood-fed females at 1, 3, and 5 dpi. Each median value was calculated from four individuals, and point error bars show the interquartile range. Asterisks represent significant differences in bacterial load between sucrose treatments at each time post-infection (* p < 0.05, ns = no significant difference). Data were collected from a total of five replicate experiments.

## The effect of a blood meal on tolerance is dynamic across infection timecourse

We also tested the effects of Blood and Sucrose on tolerance to infection, where tolerance was measured

as the slope of population survival regressed on pathogen load (24). We first built a model with Survival

as the response variable and Bacterial Load, Blood, Sucrose, and Day as the predictor variables. We

assessed main effects as well as all potential interactions and performed backward elimination to achieve

the best fit model. The final model revealed a significant three-way interaction between Blood, Bacterial

Load, and Day (Table 4: Model E; $p_{\text{Bacterial Load} \times \text{Blood} \times \text{Day}} = 0.006$), indicating that the effect of blood on

tolerance changes across time. In order to examine the nature of blood's time-dependent effect on

tolerance, we parsed our data by day and used the resulting three datasets to build a tolerance model for

each day (Fig 4). These day-specific models reveal that blood-fed mosquitoes had significantly lower

tolerance at 1 dpi (Table 5: Model F, p $_{\text{Bacterial Load} \times \text{Blood}}$ = 0.002), no significant difference at 3 dpi, and

significantly higher tolerance at 5 dpi (Table 5: Model G, p $_{\text{Bacterial Load} \times \text{Blood}}$ = 0.040) when compared to

non-blood-fed mosquitoes. In parallel, we also tested whether tolerance changes across time for blood-fed

and/or non-blood-fed treatment groups. To achieve this, we parsed our dataset by blood feeding status and

used the resulting two datasets to build a tolerance model for each blood feeding treatment group. These

models revealed that tolerance significantly decreased over time in non-blood-fed mosquitoes (Table 6:

Model H, p $_{\text{Bacterial Load} \times \text{Day}}$ = $6.70 \times 10^{-6}$) but did not significantly change in blood-fed mosquitoes (Table

6: Model I).

**Table 4. Across-Days Tolerance Model E and Summary Output.**

| Tolerance Model E (Across Days): | $E(Alive, Dead) = \varphi[\beta_0 + \beta_1(Day) + \beta_2(Blood) + \beta_3(Bacterial\ Load)$ $+ \beta_4(Day \times Blood) + \beta_5(Day \times Bacterial\ Load)$ $+ \beta_6(Blood \times Bacterial\ Load)$ $+ \beta_7(Day \times Blood \times Bacterial\ Load)]$ |
|---|---|

Summary Output:

| | Survival (Alive, Dead) | | | |
|---|---|---|---|---|
| *Predictors* | *Estimate* | *SE* | *Statistic* | *p-value* |
| (Intercept) | 1.34 | 0.33 | 4.08 | 1.67e-4 |
| Day | -0.17 | 0.08 | -2.05 | **0.046** |
| Blood | 0.93 | 0.49 | 1.91 | 0.062 |
| Bacterial Load | 0.14 | 0.06 | 2.37 | **0.022** |
| Blood : Bacterial Load | -0.24 | 0.09 | -2.59 | **0.013** |
| Day : Bacterial Load | -0.07 | 0.02 | -3.42 | **0.001** |
| Blood : Day | -0.22 | 0.12 | -1.79 | 0.080 |
| Day : Blood : Bacterial Load | 0.09 | 0.03 | 2.86 | **0.006** |
| Observations | 56 | | | |

169 Tolerance Model E is a quasibinomial generalized linear model testing the effect(s) of Blood, Sucrose, Day, and
170 Bacterial Load on Survival (Alive, Dead).

**Fig 4. The effect of blood feeding on tolerance varies across time.** Points show population survival plotted against median bacterial load at each timepoint post-infection. Error bars around points show the interquartile range of the four values used to calculate bacterial load median values. Plotted lines are derived from interaction plots of estimated marginal means calculated from day-specific binomial tolerance models (Table 5: Model F (Day 1), p $_{\text{Bacterial Load} \times \text{Blood}}$ = 0.002; Day 3, no significant predictors; Table 5: Model G (Day 5), p $_{\text{Bacterial Load} \times \text{Blood}}$ = 0.040) plotted on a response scale rather than a linear scale. The presence of solid lines indicates significantly different slopes, and therefore a significant difference in tolerance, between blood treatment groups at that timepoint. Dashed lines indicate no significant difference in tolerance at that timepoint. Data were collected from a total of five replicate experiments.

**Table 5. Day-Specific Tolerance Models F-G and Summary Outputs.**

| | |
|---|---|
| Tolerance Model F (Day 1): | $P(Alive, Dead) = \varphi[\beta_0 + \beta_1(Blood) + \beta_2(Bacterial\ Load) + \beta_3(Bacterial\ Load \times Blood)]$ |
| Tolerance Model G (Day 5): | $E(Alive, Dead) = \varphi[\beta_0 + \beta_1(Blood) + \beta_2(Bacterial\ Load) + \beta_3(Bacterial\ Load \times Blood)]$ |

Summary Outputs:

| | Tolerance Model F (Day 1) | | | | Tolerance Model G (Day 5) | | | |
|---|---|---|---|---|---|---|---|---|
| | Survival (Alive, Dead) | | | | Survival (Alive, Dead) | | | |
| *Predictors* | *Estimate* | *SE* | *Statistic* | *p-value* | *Estimate* | *SE* | *Statistic* | *p-value* |
| (Intercept) | 0.93 | 0.23 | 4.10 | 4.19e-5 | 0.48 | 0.20 | 2.36 | 0.036 |
| Blood | 1.20 | 0.39 | 3.10 | **0.002** | -0.21 | 0.30 | -0.71 | 0.492 |
| Bacterial Load | 0.08 | 0.03 | 2.36 | **0.019** | -0.33 | 0.13 | -2.53 | **0.026** |
| Blood : Bacterial Load | -0.17 | 0.06 | -3.08 | **0.002** | 0.52 | 0.23 | 2.31 | **0.040** |
| Observations | 20 | | | | 16 | | | |

Tolerance Model F is a binomial generalized linear model testing the effect(s) of Blood, Sucrose, and Bacterial Load on Survival (Alive, Dead) at 1 dpi. Tolerance Model G is a quasibinomial generalized linear model testing the effect(s) of Blood, Sucrose, and Bacterial Load on Survival (Alive, Dead) at 5 dpi.

**Table 6. Blood Feeding Status-Specific Tolerance Models H-I and Summary Outputs.**

| Tolerance Model H (Non-Blood-Fed): | $P(Alive, Dead) = \varphi[\beta_0 + \beta_1(Day) + \beta_2(Bacterial\ Load) + \beta_3(Bacterial\ Load \times Day)]$ |
|---|---|
| Tolerance Model I (Blood-Fed): | $E(Alive, Dead) = \varphi[\beta_0 + \beta_1(Day) + \beta_2(Bacterial\ Load)]$ |

Summary Outputs:

| | Tolerance Model H (Non-blood-fed) | | | | Tolerance Model I (Blood-fed) | | | |
|---|---|---|---|---|---|---|---|---|
| | Survival (Alive, Dead) | | | | Survival (Alive, Dead) | | | |
| *Predictors* | *Estimate* | *SE* | *Statistic* | *p-value* | *Estimate* | *SE* | *Statistic* | *p-value* |
| (Intercept) | 1.34 | 0.25 | 5.38 | 7.38e-8 | 2.13 | 0.35 | 6.11 | 2.18e-6 |
| Day | -0.17 | 0.06 | -2.70 | **0.007** | -0.35 | 0.08 | -4.17 | **3.23e-4** |
| Bacterial Load | 0.14 | 0.04 | 3.12 | **0.002** | -0.04 | 0.03 | -1.46 | 0.158 |
| Day : Bacterial Load | -0.07 | 0.02 | -4.50 | **6.70e-6** | | | | |
| Observations | 28 | | | | 28 | | | |

Tolerance Model H is a binomial generalized linear model testing the effect(s) of Sucrose, Day, and Bacterial Load on survival (Alive, Dead) in non-blood-fed mosquitoes. Tolerance Model I is a quasibinomial generalized linear model testing the effect(s) of Sucrose, Day, and Bacterial Load on survival (Alive, Dead) in blood-fed mosquitoes.

# A blood meal alters the shape of a mosquito's immune defense trajectory through time

Blood altered resistance early in infection and tolerance dynamically across the infection timecourse. Therefore, we compared the relative utilization of each component of immune defense by quantitatively plotting resistance and tolerance for each blood feeding status group at all three timepoints (Fig 5). Blood-fed mosquitoes displayed an increase in resistance (Table 2: Model A, $p_{Day} = 2.92 \times 10^{-4}$) and a non-significant trend toward increased tolerance (Table 6: Model I) from 1 dpi to 5 dpi, while non-blood-fed mosquitoes displayed increased resistance (Table 2: Model A, $p_{Day} = 2.92 \times 10^{-4}$) but decreased tolerance (Table 6: Model H, $p_{Bacterial\ Load \times Day} = 6.70 \times 10^{-6}$) from 1 dpi to 5 dpi (Fig 5).

**Fig 5. Blood feeding alters the mosquito's trajectory of resistance and tolerance over time.** Plotted points
display the balance of tolerance and resistance through an infection timecourse for blood-fed (left panel) and non-
blood-fed mosquitoes (right panel). To estimate tolerance, we regressed Bacterial load on Survival separately for
each blood feeding status/day combination and found the estimated marginal mean and standard error of Bacterial
Load for each model using the emmeans package (25). Resistance is the population median CFU with interquartile
ranges for each group plotted on a reversed x-axis scale (as resistance is inversely related to CFU). Data were
collected from a total of five replicate experiments.

# Discussion

## Sucrose restriction enhances resistance throughout infection timecourse

We have shown that a 1% sucrose diet confers higher resistance to intrathoracic infection by the gram-

negative bacterium *E. coli* S17 in adult female *Ae. aegypti* (Thai strain) compared to a 10% sucrose diet.

In mosquitoes, effects of dietary sugars on infection response have primarily been studied using oral

infection models. In one study, pre-infection sugar feeding enhanced *Ae. aegypti* resistance to Zika by

increasing the expression of antiviral genes (26), while in *An. stephensi*, lower dietary glucose abundance

correlated with higher resistance to *Plasmodium berghei* (21). In *D. melanogaster*, lower dietary sugar

concentration increased resistance to bacterial infection in two separate studies (22,27). Altogether, the

current body of work suggests lower dietary sugar levels usually correlate with increased resistance to

infection, which is consistent with our findings.

While dietary sucrose significantly affected mosquito resistance in our study, it did not affect survival (S1

Fig). Conversely, effects of dietary sugar on survival have been observed in *D. melanogaster*, although

the directionality of the effect is not consistent. Howick & Lazzaro (22) showed that higher dietary sugars

were associated with lower survival (and lower resistance) post-infection. However, other studies have

shown a positive relationship between dietary sugar and infection outcome (although resistance was not

directly measured). For example, in one study, flies that consumed a relatively low-protein, high-

carbohydrate diet displayed increased survival to bacterial infection and higher constitutive expression of

antimicrobial peptides (AMPs) (28). In another study, dietary glucose supplementation led to significantly

improved *D. melanogaster* longevity and survival following bacterial infection (29). Multiple

dissimilarities between these studies and our own, including experimental design and pathogen type,

could underlie the different findings. Overall, relationships between dietary sugars and survival after

infection are not fully consistent and warrant further study.

The mechanisms by which sugar affects resistance are not known. As diet is known to affect hemolymph

composition in insects (30,31), higher dietary sugar may provide a more suitable environment for bacteria

to proliferate. Alternatively, but not mutually exclusively, a lower concentration of dietary sucrose may

directly or indirectly affect signaling of the mosquito's immune pathways, thereby inhibiting resistance

mechanisms such as melanization, AMP production, and/or hemocyte activity. Higher dietary sugar levels

are associated with a stronger melanization response in mosquitoes (32–34), which may initially appear

antithetical to our results. However, *E. coli* is preferentially phagocytized rather than melanized by

mosquitoes (35–37), suggesting that any reduction in melanization that may have occurred in the 1%

sucrose treatment is unlikely to impact infection outcomes in our system.  Sucrose may have a different

effect after challenge with a pathogen that is primarily melanized. In *D. melanogaster*, differences in

dietary sugar levels are associated with physiological changes that may explain effects on resistance. For

example, flies reared on high sugar diets display abnormal hemocyte morphology, defects in phagocytosis

ability, and excessive activation of Toll/JNK in the fat body (38). Other work has also implicated high-

sugar diets in inhibition of FOXO signaling in *D. melanogaster* (39), which has been shown to be critical

in resistance to bacterial infection (40).

While the differences in immune resistance between our sucrose treatments may be explained by the

effect of dietary sucrose availability *per se*, it is also feasible that the effect we observe is the result of

general caloric restriction. Insects subject to food restriction and insects subject to infection undergo

many of the same physiological changes, including reconfiguration of intermediate metabolism, reduced

energy storage, release of glucose and fatty acids from existing energy stores, and inhibition of the

insulin-like signaling pathway (41,42). By virtue of this commonality, food restriction prior to infection

may induce a metabolic switch that functionally primes the mosquito to resist pathogen invasion. Indeed,

positive effects of starvation or dietary restriction on resistance and/or infection outcome more generally

have been observed in multiple insect orders (43–48).  Imposed starvation does not always lead to

increased resistance, however, and in many cases has been shown to cause decreased resistance and/or

adverse infection outcomes (45,49–53). For example, starved *Leptinotarsa decemlineata* beetles display

increased susceptibility to *Beauveria bassiana* and heightened mortality post-infection compared to their

non-starved counterparts (50). Similarly, the tsetse fly *Glossina morsitans morsitans* displays increased

susceptibility to both *Trypanosoma congolense* or *Trypanosoma brucei brucei* under starvation conditions

(51).

## Blood improves resistance early post-infection

Our data also reveal that a blood meal enhances resistance at 1 dpi. This differs from the effect of sucrose,

which remains significant at every timepoint measured. Previous work reported similar effects of blood

feeding on resistance to *E. coli*, in association with insulin signaling in one instance (23) and 20-

hydroxyecdysone (20E) in another (54). In addition to supporting these findings, our data contribute an

enhanced understanding of the dynamic nature of blood's effect over time. Specifically, the effect of

blood on resistance is relatively transient, especially in comparison to the effect of dietary sucrose on

bacterial resistance which is consistent for at least five days post-infection. As the effect of blood on 20E

titers generally does not extend past 48 hours post-blood meal (55,56) and 20E can enhance resistance in

mosquitoes (Reynolds et al., 2020; but see Wang et al., 2022; Werling et al., 2019), it is possible that the

dynamic effect of blood feeding on resistance we observed is due to changing 20E titers over time. Titers

of another hormone, Juvenile Hormone (JH), rapidly and drastically decline in blood-fed *Ae. aegypti*,

reaching minimums between 24 and 48 hours post-blood meal (59,60). JH and JH analogs are associated

with various immunosuppressive effects, including the downregulation of AMP genes (61–63), decreased

activity of hemocyte activator molecules (64), and reductions in phenoloxidase activity (65,66).

Therefore, it is also feasible that improved resistance after blood feeding is mediated via declines in JH.

In addition to the potential regulatory role of hormones, investigations into the mosquito's transcriptional

profile post-blood feeding indicate altered expression of immune-related genes within the 24 hours

following a blood meal (67–70). Such genes may affect resistance to bacterial infection and may or may

not be subject to regulation by insulin signaling and/or hormones such as JH and 20E.

## A blood meal, but not sucrose concentration, affects tolerance to infection

While the effect of blood on resistance is limited to 1 dpi, we observe a dynamic effect of blood feeding

on tolerance across the 5-day infection timecourse. Blood-fed mosquitoes have significantly lower

tolerance compared to non-blood-fed mosquitoes at 1 dpi, and significantly higher tolerance at 5 dpi.

Further, non-blood-fed mosquitoes show a significant decrease in tolerance across a 5-day infection

timecourse while blood-fed mosquitoes show static tolerance across a 5-day infection timecourse. This

suggests that the blood meal ameliorates a decline in tolerance that mosquitoes experience when provided

only sugar meals. Blood is a unique nutritional resource – its digestion is associated with various

physiological stressors, including rapid shifts in temperature and pH (15,71) as well as heme toxicity and

oxidative stress (72–74). Despite the challenges posed by blood digestion, mosquitoes are well adapted to

overcome the associated stressors (15,17,71,75,76). Because many stressors associated with blood feeding

are also associated with infection (20,21,77,78) it is possible that the homeostasis-promoting processes

that occur during or after blood feeding have the additional effect of promoting host health during

infection (i.e. tolerance).

Blood feeding induces multiple signaling cascades that may promote tolerance to infection. For example,

heat shock proteins (HSP) are widely conserved in both eukaryotes and prokaryotes (79) and are

implicated in a variety of processes generally related to the reduction of stress (80). They are rapidly

induced upon blood feeding in multiple arthropods and are critical to surviving the stress of a blood meal

(15,19,81). Host HSPs have also been implicated in host response to bacterial, viral, and fungal infections

alike, playing roles in homeostasis via maintenance of protein stability and functionality, reductions in

inflammation, and attenuation of autoimmunity (82,83). HSPs that protect arthropods from blood-induced

damage may also promote host health and survival during infection. Additionally, the unfolded protein

response (UPR) is a similarly highly conserved, stress-mitigating, pathway that has been heavily

implicated in defense against pathogens (reviewed by Rosche et al., 2021). Transient UPR upregulation

occurs after blood feeding in *Ae. aegypti* (85). Further, UPR upregulation promotes tolerance to

*Pseudomonas aeruginosa* in the nematode *Caenorhabditis elegans* (86). It also promotes survival in

mosquito cells infected with dengue virus in vitro by ameliorating infection-induced endoplasmic

reticulum stress (87), a function which is characteristic of tolerance.  Importantly, it is unlikely that any

single signaling pathway or molecular cascade regulates tolerance to infection, as immunopathology

affects various physiological processes (1) that require repair or protection. Thus, if the aforementioned

processes do indeed affect tolerance, they likely do so in tandem with other mechanisms.

It is also possible that the tolerance benefit of a blood meal is explained by the influx of nutrients (e.g.,

protein, lipid) associated with a blood meal. However, we observed no effect of sucrose on tolerance, and

higher dietary sucrose concentrations are also strongly associated with higher nutrient reserves. Whole-

body homogenates of mosquitoes fed 10% sucrose have significantly higher levels of sugar, glycogen,

and lipids compared to mosquitoes fed 2% sucrose (88). This indicates that if tolerance is indeed

regulated nutritionally in our system, this regulation may be specific to the nutrients provided by a blood

meal.

## Blood-fed and non-blood-fed mosquitoes employ different, but equally effective, immune defense strategies

Blood-fed and non-blood-fed mosquitoes showed markedly different resistance-tolerance strategies for

surviving infection with a non-coevolved bacterial pathogen (Figure 5). In the absence of a blood meal, a

mosquito displays an increase in resistance and a decrease in tolerance across a 5-day infection

timecourse. But when provided with a blood meal, early infection is characterized by a significantly

stronger resistance response and significantly weaker tolerance response (possibly the result of a trade-

off). Across time, blood-fed mosquitoes show no change in tolerance, and an increase in resistance that is

similar to that of non-blood-fed mosquitoes. In comparing the groups at the end of infection timecourse,

blood-fed mosquitoes show equal resistance and higher tolerance compared to non-blood-fed mosquitoes.

Each diet clearly induced a unique immune defense strategy. Interestingly, we observed no significant

difference in survival (S1 Fig) indicating that the two strategies employed by the groups are equally

effective in surviving infection. A blood meal is not only physiologically stressful and energetically costly

to digest on its own (89), it also catalyzes reproductive processes (i.e. vitellogenesis) that are energetically

costly (90). Immune defense is also a costly process  (10,91–93). Understanding how organisms partition

limited resources amongst energy-intensive processes represents an ongoing challenge in biology.

Reproduction, immunity, and digestion have been shown to physiologically trade off in multiple insect

systems (90,94–100). In light of this, it may be somewhat surprising that mosquitoes under the combined

stress of blood digestion, reproduction, and infection in parallel show no difference in survival compared

to non-blood-fed mosquitoes. Overall, our results therefore suggest that the immune defense benefits of a

blood meal are great enough to mitigate the stressors and resource usage associated with this meal.

# Conclusions

In the present study, we show the effects of blood ingestion and two dietary sucrose concentrations on both resistance and tolerance to the non-coevolved bacterium *E. coli* in the adult female yellow fever mosquito *Ae. aegypti*. Our results indicate that dietary sucrose concentration and blood ingestion both affect resistance, while only blood ingestion affects tolerance. The effect of blood on tolerance was dynamic across time, significantly worsening tolerance at the start of the infection timecourse and significantly enhancing tolerance by the end. Mosquitoes are one of many arthropods that transmit pathogens by virtue of a hematophagous lifestyle: mosquitoes, ticks, biting midges, triatomine bugs, fleas, black flies, and sand flies alike all transmit pathogens when consuming vertebrate blood (101). Likewise, they all share the ability to tolerate the pathogens they transmit, which is critical for successful transmission. In light of this, motivation for exploring the relationship between hematophagy and tolerance is not limited to mosquitoes, but rather, is broadly relevant in all arthropod-borne pathogen systems. Further, shared characteristics of tolerance biology could potentially be leveraged as novel targets for effective and sustainable vector control (102). Because blood feeding is already one of the best-described areas of mosquito biology, has been investigated (103–106) and implemented (107) as a vector control target, and affects tolerance in *Ae. aegypti*, targeting blood feeding in the context of tolerance-focused vector control holds excellent potential.

# Methods

## Mosquitoes

Throughout the duration of the experiment, *Ae. aegypti* Thai strain (Laura C. Harrington, Cornell University) mosquitoes were reared in a chamber maintained at 27°C and 80% relative humidity under a 14h:10h light:dark cycle. First, eggs were hatched in RO water placed in a vacuum chamber. Upon hatching, larvae were reared in trays containing RO water at a density of 200-300 larvae per tray and given one pinch of Tetramin fish flakes as well as cat food ad libitum until pupation. Upon pupation, pupal cups were split into four 8"x8" mesh treatment cages (Bioquip, Rancho Dominguez, CA, USA):

10% sucrose only, 10% sucrose + blood, 1% sucrose only, and 1% sucrose + blood. Each cage received

the appropriate sucrose meal concurrent with the addition of pupal cups. Pupae were allotted 48 hours to

eclose before pupal cup removal. Post-eclosion, all adults were left undisturbed for 48 hours, then starved

for 24 hours. Next, mosquitoes from cages containing blood feeding treatments were provided with a

blood meal maintained at 37 °C via a membrane feeding system (Hemotek, Blackburn, UK) for 1-2 hours.

Sucrose meals were then returned to all mosquitoes. Infections were performed on females at 24 hours

post-blood feeding. At this time, blood feeding status was confirmed visually under a microscope by the

presence of a blood bolus.

## Sucrose and blood diets

392 Mosquitoes were maintained on one of four experimental diets: 10% sucrose alone, 10% sucrose + blood,

393 1% sucrose alone, and 1% sucrose + blood. Sucrose meals were created by passing a solution of DI water

394 and UltraPure sucrose (Life Technologies, Carlsbad, CA) through a 0.2 µm filter. Blood meals consisted

395 of defibrinated rabbit blood (Hemostat, Dixon, CA, USA) supplemented with $Na_2ATP$ to a 1 mM

396 concentration.

## Bacterial culture and mosquito infections

The bacteria used for infections, *E. coli* S17 pPROBE-mCherry (Dimopoulos Lab, unpublished data),

contains a fluorescent mCherry plasmid and a kanamycin resistance cassette. Bacteria were grown in

Luria broth (LB) supplemented with kanamycin (50 µg/mL) overnight at 30°C with shaking. Cultures

were washed thrice in sterile 1X PBS, then pelleted and resuspended to OD600 = 1 ± 0.1 (mean of $1.23\times^9$

CFU/mL across replicates) . At the time of infection, mosquitoes were anesthetized on ice and females

were injected with 69 nanoliters of a $1 \times 10^{-2}$ dilution of this culture by piercing the soft tissue of the

anepisternal cleft of the mesothorax using a Nanoject II Auto-Nanoliter Injector (Drummond, Broomall,

PA, USA). Fresh injection needles were prepared for each replicate by manually pulling a borosilicate

glass capillary tube (Drummond, Broomall, PA, USA) over a flame to achieve a tip with an outer

diameter no greater than 500 microns as measured using a stage micrometer. Preliminary experiments

indicated that Nanoject II delivery of bacteria was adequately precise, delivering a mean of $312 \pm 13$ CFU

per mosquito (n = 6 mosquitoes were injected, then each homogenate was immediately cultured as

described in the subsequent section) (S1 File). Following infection, mosquitoes from each diet group were

randomly allocated into two groups and placed in separate cages for survival and bacterial load

measurements.

## Monitoring survival and bacterial load

Survival and bacterial load were measured on days 1 (24 hours +/- 2 hours), 3 (72 hours +/- 2 hours), and

5 (120 hours +/- 2 hours) post-infection. Immediately after data collection at each timepoint, dead

individuals were removed from survival and bacterial load cages and discarded. Survival was monitored

by counting the number of mosquitoes dead and alive. To measure bacterial load, four living mosquitoes

were sampled using an InsectaVac Aspirator (Bioquip, Rancho Dominguez, CA, USA) and individually

homogenized in 150 µL sterile 1X PBS. Serial dilutions were performed at $10^{-2}$ and $10^{-4}$ and cultured

alongside undiluted homogenate on LB supplemented with kanamycin (50 µg /mL) at 30°C for 24-48

hours. Resulting fluorescent colonies were counted using a stereo microscope fluorescence adapter

system (Nightsea, Lexington, MA). For each individual, the least dilute plate with countable CFUs was

used to obtain a representative value. After obtaining a CFU count for each individual mosquito, the

median of four mosquitoes was calculated and used in the data set. Each row in the data set (S1 File)

contains the median CFU of four  mosquitoes paired with the accompanying survival value for that

treatment group.

## Statistical analysis

All analyses were performed using R statistical software (108) and RStudio (109).

To measure resistance, we built linear models testing the effect(s) of Blood, Sucrose, Day, and Replicate

on Bacterial Load (Table 1). Backward elimination was used to assess all possible interactions as well as

main effects. The presence of any main effect indicates that variable significantly affects resistance.

To measure tolerance, we used a reaction norm approach (24) to test for variation in health across a range

of real-time pathogen loads between diet groups at multiple time points. Our approach is thus categorized

as range tolerance rather than point tolerance (110). We built binomial generalized linear models testing

the effect(s) of Blood, Sucrose, Day, and Bacterial Load on Survival (Table 1). Backward elimination was

used to assess all possible interactions as well as main effects. Further, we tested all tolerance models for

overdispersion (111) by performing residual deviance goodness-of-fit tests and Pearson goodness-of-fit

tests. When detected, we adjusted for overdispersion by correcting the standard errors in those models

through incorporation of a dispersion parameter when calculating the variance. Models with

overdispersion parameters are defined as quasi-generalized linear models using a binomial distribution

where the variance is given by $\varphi \times \mu$, in which $\mu$ is the mean and $\varphi$ the dispersion parameter. We used

these final models to interpret any effect(s) of our predictor variables on tolerance. Specifically, a

significant interaction between Bacterial Load and another variable in predicting Survival indicates that

variable significantly affects tolerance.

To obtain interaction plots of estimated marginal means for each day-specific binomial tolerance model

(Figure 4), we used the emmip function from the emmeans package (25). To obtain representative values

for tolerance for each level of Blood on Days 1, 3, and 5 (Figure 5), we used the emtrends function from

the emmeans package, a function that creates a reference grid for each model of interest, then calculates

difference quotients of predictions from those reference grids and computes the marginal averages and

standard errors for those averages. We used the emtrends function to obtain these values for each day

using Model F (Day 1), Model G (Day 5), and a model with a non-significant Bacterial Load × Blood

term for Day 3 (model not shown).

# **Acknowledgements**

454 We thank Ellen Klinger and Megan Meuti (The Ohio State University) for statistical advice and helpful

455 experimental design comments, George Dimopoulos and Yuemei Dong (Johns Hopkins University) for

456   providing the mCherry-transformed strain of *E. coli*, as well as Laura C. Harrington (Cornell University)

457   for the gift of *Ae. aegypti* (Thai strain).

# Supporting information

**S1 Fig. Survival across a 5-day infection timecourse is affected by neither blood nor sucrose.**

Kaplan-Meier curves are displayed to compare the effects of (A) blood treatment and (B) sucrose

treatment on survival in mosquitoes infected with *E. coli*. Corresponding log-rank test p-values are

displayed on each panel.

**S1 File. Selected R code used for statistical analysis and figures.**

**S1 Dataset. Data used in analyses.** Data are divided into three tabs: final dataset (df_final), survival

analysis dataset (df_surv), and raw CFU dilution counts (raw CFU counts) used to calculate medians used

in df_final.

# References

1.   Medzhitov R, Schneider DS, Soares MP. Disease Tolerance as a Defense Strategy. Science. 2012 Feb
24;335(6071):936–41.

2.   Schneider DS, Ayres JS. Two ways to survive infection: what resistance and tolerance can teach us
about treating infectious diseases. Nat Rev Immunol. 2008 Nov;8(11):889–95.

3.   Cobb NA. Contributions to an economic knowledge of Australian rusts (Uredineae). [Chapter X].
Improving wheat by selection [Internet]. Departmet of Agriculture, New South Wales; 1894 Apr
[cited 2022 Sep 26]. (Contributions to an economic knowledge of Australian rusts (Uredineae).).
Report No.: 18. Available from: https://catalog.hathitrust.org/Record/008904165

4.   Corby-Harris V, Habel KE, Ali FG, Promislow DEL. Alternative measures of response to
Pseudomonas aeruginosa infection in Drosophila melanogaster. J Evol Biol. 2007;20(2):526–33.

5.   Råberg L, Sim D, Read AF. Disentangling Genetic Variation for Resistance and Tolerance to
Infectious Diseases in Animals. Science. 2007 Nov;318(5851):812–4.

6.   McCarville JL, Ayres JS. Disease tolerance: concept and mechanisms. Curr Opin Immunol. 2018 Feb
1;50:88–93.

7.   Råberg L, Graham AL, Read AF. Decomposing health: tolerance and resistance to parasites in
animals. Philos Trans R Soc B Biol Sci. 2009 Jan 12;364(1513):37–49.

8.    Read AF, Graham AL, Råberg L. Animal Defenses against Infectious Agents: Is Damage Control
485            More Important Than Pathogen Control. PLOS Biol. 2008 Dec 23;6(12):e1000004.

486    9.    Soares MP, Teixeira L, Moita LF. Disease tolerance and immunity in host protection against
487            infection. Nat Rev Immunol. 2017 Feb;17(2):83–96.

488    10.   Baucom RS, de Roode JC. Ecological immunology and tolerance in plants and animals. Funct Ecol.
489            2011;25(1):18–28.

490    11.   Vector-borne diseases [Internet]. World Health Organization; 2022 Mar [cited 2022 Oct 17].
491            Available from: https://www.who.int/news-room/fact-sheets/detail/vector-borne-diseases

492    12.   Barredo E, DeGennaro M. Not Just from Blood: Mosquito Nutrient Acquisition from Nectar
493            Sources. Trends Parasitol. 2020 May 1;36(5):473–84.

494    13.   Wolff GH, Riffell JA. Olfaction, experience and neural mechanisms underlying mosquito host
495            preference. J Exp Biol. 2018 Feb 27;221(4):jeb157131.

496    14.   Foster WA. Mosquito Sugar Feeding and Reproductive Energetics. Annu Rev Entomol.
497            1995;40(1):443–74.

498    15.   Benoit JB, Lopez-Martinez G, Patrick KR, Phillips ZP, Krause TB, Denlinger DL. Drinking a hot blood
499            meal elicits a protective heat shock response in mosquitoes. Proc Natl Acad Sci. 2011 May
500            10;108(19):8026–9.

501    16.   Champion CJ, Xu J. The impact of metagenomic interplay on the mosquito redox homeostasis. Free
502            Radic Biol Med. 2017 Apr 1;105:79–85.

503    17.   Lahondère C, Lazzari CR. Mosquitoes Cool Down during Blood Feeding to Avoid Overheating. Curr
504            Biol. 2012 Jan 10;22(1):40–5.

505    18.   Oliveira MF, Silva JR, Dansa-Petretski M, de Souza W, Lins U, Braga CMS, et al. Haem detoxification
506            by an insect. Nature. 1999 Aug;400(6744):517–8.

507    19.   Pereira MH, Paim RMM, Lahondère C, Lazzari CR. Heat Shock Proteins and Blood-Feeding in
508            Arthropods. In: Asea AAA, Kaur P, editors. Heat Shock Proteins in Veterinary Medicine and
509            Sciences [Internet]. Cham: Springer International Publishing; 2017 [cited 2022 Sep 16]. p. 349–59.
510            (Heat Shock Proteins). Available from: https://doi.org/10.1007/978-3-319-73377-7_13

511    20.   Peterson TML, Luckhart S. A mosquito 2-Cys peroxiredoxin protects against nitrosative and
512            oxidative stresses associated with malaria parasite infection. Free Radic Biol Med. 2006 Mar
513            15;40(6):1067–82.

514    21.   Wang M, An Y, Gao L, Dong S, Zhou X, Feng Y, et al. Glucose-mediated proliferation of a gut
515            commensal bacterium promotes Plasmodium infection by increasing mosquito midgut pH. Cell
516            Rep. 2021 Apr 20;35(3):108992.

517    22.   Howick VM, Lazzaro BP. Genotype and diet shape resistance and tolerance across distinct phases
518            of bacterial infection. BMC Evol Biol. 2014 Mar 22;14(1):56.

519    23.    Castillo J, Brown MR, Strand MR. Blood Feeding and Insulin-like Peptide 3 Stimulate Proliferation
520           of Hemocytes in the Mosquito Aedes aegypti. PLOS Pathog. 2011 Oct 6;7(10):e1002274.

521    24.    Simms EL. Defining tolerance as a norm of reaction. Evol Ecol. 2000 Jul;14(4–6):563–70.

522    25.    Lenth RV, Buerkner P, Herve M, Jung M, Love J, Miguez F, et al. emmeans: Estimated Marginal
523           Means, aka Least-Squares Means [Internet]. 2022 [cited 2022 Sep 19]. Available from:
524           https://CRAN.R-project.org/package=emmeans

525    26.    Almire F, Terhzaz S, Terry S, McFarlane M, Gestuveo RJ, Szemiel AM, et al. Sugar feeding protects
526           against arboviral infection by enhancing gut immunity in the mosquito vector Aedes aegypti. PLOS
527           Pathog. 2021 Sep 2;17(9):e1009870.

528    27.    Unckless RL, Rottschaefer SM, Lazzaro BP. The Complex Contributions of Genetics and Nutrition to
529           Immunity in Drosophila melanogaster. PLOS Genet. 2015 Mar 12;11(3):e1005030.

530    28.    Ponton F, Morimoto J, Robinson K, Kumar SS, Cotter SC, Wilson K, et al. Macronutrients modulate
531           survival to infection and immunity in Drosophila. J Anim Ecol. 2020;89(2):460–70.

532    29.    Galenza A, Hutchinson J, Campbell SD, Hazes B, Foley E. Glucose modulates Drosophila longevity
533           and immunity independent of the microbiota. Biol Open. 2016 Jan 21;5(2):165–73.

534    30.    Mullins DE. Chemistry and Physiology of the Hemolymph. In: Comprehensive Insect Physiology,
535           Volume 3. 1st ed. 1985. p. 355–400.

536    31.    Thompson SN. Trehalose – The Insect 'Blood' Sugar. In: Advances in Insect Physiology [Internet].
537           Elsevier; 2003 [cited 2023 Jan 3]. p. 205–85. Available from:
538           https://linkinghub.elsevier.com/retrieve/pii/S0065280603310045

539    32.    Ferguson LV, Beckett NH, French MC, Campbell MJ, Smith TG, Adamo SA. Sugar intake interacts
540           with temperature to influence reproduction and immunity in adult Culex pipiens mosquitoes. Can J
541           Zool. 2019 May;97(5):424–8.

542    33.    Koella JC, Sørensen FL. Effect of adult nutrition on the melanization immune response of the
543           malaria vector Anopheles stephensi. Med Vet Entomol. 2002;16(3):316–20.

544    34.    Schwartz A, Koella JC. Melanization of Plasmodium falciparum and C-25 Sephadex Beads by Field-
545           Caught Anopheles gambiae (Diptera: Culicidae) from Southern Tanzania. J Med Entomol. 2002 Jan
546           1;39(1):84–8.

547    35.    Hernández-Martínez S, Lanz H, Rodrguez MH, González-Ceron L, Tsutsumi V. Cellular-Mediated
548           Reactions to Foreign Organisms Inoculated into the Hemocoel of Anopheles albimanus (Diptera:
549           Culicidae). J Med Entomol. 2002 Jan 1;39(1):61–9.

550    36.    Hillyer JF, Schmidt SL, Christensen BM. Hemocyte-mediated phagocytosis and melanization in the
551           mosquito Armigeres subalbatus following immune challenge by bacteria. Cell Tissue Res. 2003 Jul
552           1;313(1):117–27.

37.   Hillyer JF, Schmidt SL, Christensen BM. RAPID PHAGOCYTOSIS AND MELANIZATION OF BACTERIA
AND PLASMODIUM SPOROZOITES BY HEMOCYTES OF THE MOSQUITO AEDES AEGYPTI. J Parasitol.
2003 Feb 1;89(1):62–9.

38.   Yu S, Zhang G, Jin LH. A high-sugar diet affects cellular and humoral immune responses in
Drosophila. Exp Cell Res. 2018 Jul 15;368(2):215–24.

39.   Dobson AJ, Ezcurra M, Flanagan CE, Summerfield AC, Piper MDW, Gems D, et al. Nutritional
Programming of Lifespan by FOXO Inhibition on Sugar-Rich Diets. Cell Rep. 2017 Jan 10;18(2):299–
306.

40.   Fink C, Hoffmann J, Knop M, Li Y, Isermann K, Roeder T. Intestinal FoxO signaling is required to
survive oral infection in Drosophila. Mucosal Immunol. 2016 Jul;9(4):927–36.

41.   Adamo SA. How insects protect themselves against combined starvation and pathogen challenges,
and the implications for reductionism. Comp Biochem Physiol B Biochem Mol Biol. 2021 Aug
1;255:110564.

42.   Adamo SA, Miyashita A. Stayin' Alive: Endocrinological Stress Responses in Insects. In: Advances in
Invertebrate (Neuro)Endocrinology: A Collection of Reviews in the Post-Genomic Era, Volume 2:
Arthropoda. 1st ed. 2020. p. 283–323.

43.   Brown AE, Baumbach J, Cook PE, Ligoxygakis P. Short-Term Starvation of Immune Deficient
Drosophila Improves Survival to Gram-Negative Bacterial Infections. PLOS ONE. 2009 Feb
16;4(2):e4490.

44.   Burger JMS, Hwangbo DS, Corby-Harris V, Promislow DEL. The functional costs and benefits of
dietary restriction in Drosophila. Aging Cell. 2007;6(1):63–71.

45.   Conroy TJ, Palmer-Young EC, Irwin RE, Adler LS. Food Limitation Affects Parasite Load and Survival
of Bombus impatiens (Hymenoptera: Apidae) Infected With Crithidia (Trypanosomatida:
Trypanosomatidae). Environ Entomol. 2016 Oct 1;45(5):1212–9.

46.   Kang KD, Kamita SG, Suzuki K, Seong SI. Effect of starvation upon baculovirus replication in larval
Bombyx mori and Heliothis virescens. J Invertebr Pathol. 2011 Feb 1;106(2):205–10.

47.   Logan A, Ruiz-González MX, Brown MJF. The impact of host starvation on parasite development
and population dynamics in an intestinal trypanosome parasite of bumble bees. Parasitology. 2005
Jun;130(6):637–42.

48.   Moller-Jacobs LL, Murdock CC, Thomas MB. Capacity of mosquitoes to transmit malaria depends
on larval environment. Parasit Vectors. 2014 Dec 14;7(1):593.

49.   Akoda K, Van den Bossche P, Lyaruu EA, De Deken R, Marcotty T, Coosemans M, et al. Maturation
of a Trypanosoma Brucei Infection to the Infectious Metacyclic Stage Is Enhanced in Nutritionally
Stressed Tsetse Flies. J Med Entomol. 2009 Nov 1;46(6):1446–9.

50.    Furlong MJ, Groden E. Starvation induced stress and the susceptibility of the Colorado potato
588           beetle, Leptinotarsa decemlineata, to infection by Beauveria bassiana. J Invertebr Pathol. 2003 Jun
589           1;83(2):127–38.

590    51.    Kubi C, Van Den Abbeele J, De Deken R, Marcotty T, Dorny P, Van Den Bossche P. The effect of
591           starvation on the susceptibility of teneral and non-teneral tsetse flies to trypanosome infection.
592           Med Vet Entomol. 2006;20(4):388–92.

593    52.    Le Rohellec M, Le Bourg É. Contrasted effects of suppressing live yeast from food on longevity,
594           aging and resistance to several stresses in Drosophila melanogaster. Exp Gerontol. 2009 Nov
595           1;44(11):695–707.

596    53.    Pavlushin SV, Belousova IA, Chertkova EA, Akhanaev YuB, Martemyanov VV, Glupov VV. Effect of
597           Starvation as a Population Stress-Factor on the Activation of Covert Baculovirus Infection in the
598           Gypsy Moth. Biol Bull Rev. 2021 Jan 1;11(1):86–91.

599    54.    Reynolds RA, Kwon H, Smith RC. 20-Hydroxyecdysone Primes Innate Immune Responses That Limit
600           Bacterial and Malarial Parasite Survival in Anopheles gambiae. mSphere. 2020 Apr 15;5(2):e00983-
601           19.

602    55.    Baldridge GD, Feyereisen R. Ecdysteroid titer and oocyte growth in the northern house mosquito,
603           Culex pipiens L. Comp Biochem Physiol A Physiol. 1986 Jan 1;83(2):325–9.

604    56.    Martín D, Wang SF, Raikhel AS. The vitellogenin gene of the mosquito Aedes aegypti is a direct
605           target of ecdysteroid receptor. Mol Cell Endocrinol. 2001 Feb 22;173(1):75–86.

606    57.    Wang M, Wang Y, Chang M, Wang X, Shi Z, Raikhel AS, et al. Ecdysone signaling mediates the
607           trade-off between immunity and reproduction via suppression of amyloids in the mosquito Aedes
608           aegypti. PLOS Pathog. 2022 Sep 22;18(9):e1010837.

609    58.    Werling K, Shaw WR, Itoe MA, Westervelt KA, Marcenac P, Paton DG, et al. Steroid hormone
610           function controls non-competitive Plasmodium development in Anopheles. Cell. 2019 Apr
611           4;177(2):315-325.e14.

612    59.    Hernández-Martínez S, Rivera-Perez C, Nouzova M, Noriega FG. Coordinated changes in JH
613           biosynthesis and JH hemolymph titers in Aedes aegypti mosquitoes. J Insect Physiol. 2015 Jan
614           1;72:22–7.

615    60.    Shapiro AB, Wheelock GD, Hagedorn HH, Baker FC, Tsai LW, Schooley DA. Juvenile hormone and
616           juvenile hormone esterase in adult females of the mosquito Aedes aegypti. J Insect Physiol. 1986
617           Jan 1;32(10):867–77.

618    61.    Chang MM, Wang YH, Yang QT, Wang XL, Wang M, Raikhel AS, et al. Regulation of antimicrobial
619           peptides by juvenile hormone and its receptor, Methoprene-tolerant, in the mosquito Aedes
620           aegypti. Insect Biochem Mol Biol. 2021 Jan 1;128:103509.

621    62.    Flatt T, Heyland A, Rus F, Porpiglia E, Sherlock C, Yamamoto R, et al. Hormonal regulation of the
622           humoral innate immune response in Drosophila melanogaster. J Exp Biol. 2008 Aug
623           15;211(16):2712–24.

63. Schwenke RA, Lazzaro BP. Juvenile Hormone Suppresses Resistance to Infection in Mated Female
Drosophila melanogaster. Curr Biol. 2017 Feb 20;27(4):596–601.

64. Clark KD, Kim Y, Strand MR. Plasmatocyte sensitivity to plasmatocyte spreading peptide (PSP)
fluctuates with the larval molting cycle. J Insect Physiol. 2005 May 1;51(5):587–96.

65. Contreras-Garduño J, Córdoba-Aguilar A, Lanz-Mendoza H, Cordero Rivera A. Territorial behaviour
and immunity are mediated by juvenile hormone: the physiological basis of honest signalling?
Funct Ecol. 2009;23(1):157–63.

66. Rolff J, Siva-Jothy MT. Copulation corrupts immunity: A mechanism for a cost of mating in insects.
Proc Natl Acad Sci. 2002 Jul 23;99(15):9916–8.

67. Bonizzoni M, Dunn WA, Campbell CL, Olson KE, Dimon MT, Marinotti O, et al. RNA-seq analyses of
blood-induced changes in gene expression in the mosquito vector species, Aedes aegypti. BMC
Genomics. 2011 Jan 28;12(1):82.

68. Das S, Radtke A, Choi YJ, Mendes AM, Valenzuela JG, Dimopoulos G. Transcriptomic and functional
analysis of the Anopheles gambiae salivary gland in relation to blood feeding. BMC Genomics.
2010 Oct 14;11(1):566.

69. Hixson B, Bing XL, Yang X, Bonfini A, Nagy P, Buchon N. A transcriptomic atlas of Aedes aegypti
reveals detailed functional organization of major body parts and gut regional specializations in
sugar-fed and blood-fed adult females. Lemaître B, Banerjee U, Pondeville E, editors. eLife. 2022
Apr 26;11:e76132.

70. Price DP, Nagarajan V, Churbanov A, Houde P, Milligan B, Drake LL, et al. The Fat Body
Transcriptomes of the Yellow Fever Mosquito Aedes aegypti, Pre- and Post- Blood Meal. PLOS
ONE. 2011 Jul 27;6(7):e22573.

71. del Pilar Corena M, VanEkeris L, Salazar MaI, Bowers D, Fiedler MM, Silverman D, et al. Carbonic
anhydrase in the adult mosquito midgut. J Exp Biol. 2005 Sep 1;208(17):3263–73.

72. Graça-Souza AV, Maya-Monteiro C, Paiva-Silva GO, Braz GRC, Paes MC, Sorgine MHF, et al.
Adaptations against heme toxicity in blood-feeding arthropods. Insect Biochem Mol Biol. 2006 Apr
1;36(4):322–35.

73. Gutteridge JMC, Smith A. Antioxidant protection by haemopexin of haem-stimulated lipid
peroxidation. Biochem J. 1988 Dec 15;256(3):861–5.

74. Jeney V, Balla J, Yachie A, Varga Z, Vercellotti GM, Eaton JW, et al. Pro-oxidant and cytotoxic
effects of circulating heme. Blood. 2002 Aug 1;100(3):879–87.

75. Oliveira JHM, Gonçalves RLS, Lara FA, Dias FA, Gandara ACP, Menna-Barreto RFS, et al. Blood Meal-
Derived Heme Decreases ROS Levels in the Midgut of Aedes aegypti and Allows Proliferation of
Intestinal Microbiota. PLOS Pathog. 2011 Mar 17;7(3):e1001320.

76.    Saeaue L, Morales NP, Komalamisra N, Morales Vargas RE. Antioxidative systems defense against
659              oxidative stress induced by blood meal in Aedes aegypti. Southeast Asian J Trop Med Public
660              Health. 2011 May;42(3):542–9.

661     77.    Ahmed AM. Lipid peroxidation and oxidative protein products as biomarkers of oxidative stress in
662              the autogenous mosquito, Aedes caspius, upon infection with the mosquitocidal bacterium,
663              Bacillus thuringiensis Kurstaki. Pak J Zool Pak [Internet]. 2012 [cited 2022 Dec 8]; Available from:
664              https://scholar.google.com/scholar_lookup?title=Lipid+peroxidation+and+oxidative+protein+prod
665              ucts+as+biomarkers+of+oxidative+stress+in+the+autogenous+mosquito%2C+Aedes+caspius%2C+
666              upon+infection+with+the+mosquitocidal+bacterium%2C+Bacillus+thuringiensis+Kurstaki&author=
667              Ahmed%2C+A.M.+%28Kind+Saud+Univ.%2C+Riyadh+%28Saudi+Arabia%29.+Zoology+Dept.%29&p
668              ublication_year=2012

669     78.    Chen TH, Chiang YH, Hou JN, Cheng CC, Sofiyatun E, Chiu CH, et al. XBP1-Mediated BiP/GRP78
670              Upregulation Copes with Oxidative Stress in Mosquito Cells during Dengue 2 Virus Infection.
671              BioMed Res Int. 2017 Oct 1;2017:3519158.

672     79.    Lindquist S, Craig EA. The heat-shock proteins. Annu Rev Genet. 1988;22:631–77.

673     80.    Santoro MG. Heat shock factors and the control of the stress response. Biochem Pharmacol. 2000
674              Jan 1;59(1):55–63.

675     81.    Paim RMM, Araujo RN, Leis M, Sant'anna MRV, Gontijo NF, Lazzari CR, et al. Functional evaluation
676              of Heat Shock Proteins 70 (HSP70/HSC70) on Rhodnius prolixus (Hemiptera, Reduviidae)
677              physiological responses associated with feeding and starvation. Insect Biochem Mol Biol. 2016
678              Oct;77:10–20.

679     82.    Bolhassani A, Agi E. Heat shock proteins in infection. Clin Chim Acta. 2019 Nov 1;498:90–100.

680     83.    Pockley AG. Heat shock proteins as regulators of the immune response. The Lancet. 2003 Aug
681              9;362(9382):469–76.

682     84.    Rosche KL, Sidak-Loftis LC, Hurtado J, Fisk EA, Shaw DK. Arthropods Under Pressure: Stress
683              Responses and Immunity at the Pathogen-Vector Interface. Front Immunol [Internet]. 2021 [cited
684              2022 Jul 6];11. Available from: https://www.frontiersin.org/articles/10.3389/fimmu.2020.629777

685     85.    Weng SC, Shiao SH. The unfolded protein response modulates the autophagy-mediated egg
686              production in the mosquito Aedes aegypti. Insect Mol Biol. 2020;29(4):404–16.

687     86.    Richardson CE, Kooistra T, Kim DH. An essential role for XBP-1 in host protection against immune
688              activation in C. elegans. Nature. 2010 Feb;463(7284):1092–5.

689     87.    Hou JN, Chen TH, Chiang YH, Peng JY, Yang TH, Cheng CC, et al. PERK Signal-Modulated Protein
690              Translation Promotes the Survivability of Dengue 2 Virus-Infected Mosquito Cells and Extends Viral
691              Replication. Viruses [Internet]. 2017 Sep 20 [cited 2021 Jan 1];9(9). Available from:
692              https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5618028/

693    88.    Vaidyanathan R, Fleisher AE, Minnick SL, Simmons KA, Scott TW. Nutritional Stress Affects
694           Mosquito Survival and Vector Competence for West Nile Virus. Vector-Borne Zoonotic Dis. 2008
695           Dec;8(6):727–32.

696    89.    Sarfati M, Krasnov BR, Ghazaryan L, Khokhlova IS, Fielden LJ, Degen AA. Energy costs of blood
697           digestion in a host-specific haematophagous parasite. J Exp Biol. 2005 Jul 1;208(13):2489–96.

698    90.    Schwenke RA, Lazzaro BP, Wolfner MF. Reproduction–Immunity Trade-Offs in Insects. Annu Rev
699           Entomol. 2016;61:239–56.

700    91.    Buttgereit F, Burmester GR, Brand MD. Bioenergetics of immune functions: fundamental and
701           therapeutic aspects. Immunol Today. 2000 Apr 1;21(4):194–9.

702    92.    Lazzaro BP. Adenosine Signaling and the Energetic Costs of Induced Immunity. PLOS Biol. 2015 Apr
703           27;13(4):e1002136.

704    93.    Sheldon BC, Verhulst S. Ecological immunology: costly parasite defences and trade-offs in
705           evolutionary ecology. Trends Ecol Evol. 1996 Aug 1;11(8):317–21.

706    94.    Adamo SA, Bartlett A, Le J, Spencer N, Sullivan K. Illness-induced anorexia may reduce trade-offs
707           between digestion and immune function. Anim Behav. 2010 Jan 1;79(1):3–10.

708    95.    Ahmed AM, Baggott SL, Maingon R, Hurd H. The costs of mounting an immune response are
709           reflected in the reproductive fitness of the mosquito Anopheles gambiae. Oikos. 2002;97(3):371–7.

710    96.    Ahmed AM, Hurd H. Immune stimulation and malaria infection impose reproductive costs in
711           Anopheles gambiae via follicular apoptosis. Microbes Infect. 2006 Feb 1;8(2):308–15.

712    97.    Andrews KB, Kemper D, Powell KS, Cooper PD. Spatial trade-offs in the digestive and reproductive
713           systems of grape phylloxera. Aust J Zool. 2011;59(6):392.

714    98.    Nordling D, Andersson M, Zohari S, Lars G. Reproductive effort reduces specific immune response
715           and parasite resistance. Proc R Soc Lond B Biol Sci. 1998 Jul 22;265(1403):1291–8.

716    99.    Schwartz A, Koella JC. The cost of immunity in the yellow fever mosquito, Aedes aegypti depends
717           on immune activation. J Evol Biol. 2004;17(4):834–40.

718    100.   Tripet F, Aboagye-Antwi F, Hurd H. Ecological immunology of mosquito–malaria interactions.
719           Trends Parasitol. 2008 May 1;24(5):219–27.

720    101.   Berenger JM, Parola P. 12 - Arthropod Vectors of Medical Importance. In: Cohen J, Powderly WG,
721           Opal SM, editors. Infectious Diseases (Fourth Edition) [Internet]. Elsevier; 2017 [cited 2022 Dec
722           14]. p. 104-112.e1. Available from:
723           https://www.sciencedirect.com/science/article/pii/B9780702062858000125

724    102.   Lambrechts L, Saleh MC. Manipulating Mosquito Tolerance for Arbovirus Control. Cell Host
725           Microbe. 2019 Sep;26(3):309–13.

103. Chen Q, Pei D, Li J, Jing C, Wu W, Man Y. The antenna transcriptome changes in mosquito
Anopheles sinensis, pre- and post- blood meal. PLOS ONE. 2017 Jul 17;12(7):e0181399.

104. Das De T, Thomas T, Verma S, Singla D, Chauhan C, Srivastava V, et al. A Synergistic Transcriptional
Regulation of Olfactory Genes Drives Blood-Feeding Associated Complex Behavioral Responses in
the Mosquito Anopheles culicifacies. Front Physiol. 2018;9:577.

105. Ni M, Zhao T, Lv H xin, Li M jin, Xing D, Zhao T yan, et al. Screening for odorant receptor genes
expressed in Aedes aegypti involved in host-seeking, blood-feeding and oviposition behaviors.
Parasit Vectors. 2022 Mar 4;15(1):71.

106. Wu Q, Li C xiao, Liu Q mei, Guo X xia, Shi Q ming, Zhang H duan, et al. RNA interference of odorant
receptor CquiOR114/117 affects blood-feeding behavior in Culex quinquefasciatus. Acta Trop.
2020 Apr 1;204:105343.

107. Lyimo IN, Kessy ST, Mbina KF, Daraja AA, Mnyone LL. Ivermectin-treated cattle reduces blood
digestion, egg production and survival of a free-living population of Anopheles arabiensis under
semi-field condition in south-eastern Tanzania. Malar J. 2017 Jun 6;16(1):239.

108. R Core Team. R: A language and environment for statistical computing [Internet]. Vienna, Austria:
R Foundation for Statistical Computing; 2022. Available from: https://www.R-project.org/

109. RStudio Team. RStudio: Integrated Development Environment for R [Internet]. Boston, MA:
RStudio, PBC; 2022. Available from: http://www.rstudio.com/

110. Ayres JS, Schneider DS. Tolerance of Infections. Annu Rev Immunol. 2012 Apr 23;30(1):271–94.

111. Dunn PK, Smyth GK. Chapter 9: Models for Proportions: Binomial GLMs. In: Generalized Linear
Models With Examples in R [Internet]. New York, NY: Springer New York; 2018 [cited 2023 Jan 13].
p. 333–69. (Springer Texts in Statistics). Available from: http://link.springer.com/10.1007/978-1-
4419-0118-7_9

749