Title

Circadian oscillation in primary cilium length by clock genes

regulate fibroblast cell migration

Ryota Nakazato, Yuki Matsuda, Koji Ikegami

Department of Anatomy and Developmental Biology, Graduate School of Biomedical

and Health Sciences, Hiroshima University, Hiroshima, Japan

Abstract

Various mammalian cells have autonomous cellular clocks that are produced by the transcriptional cycle of clock genes. Cellular clocks provide circadian rhythms for cellular functions via transcriptional and cytoskeletal regulation. The vast majority of mammalian cells possess a primary cilium, an organelle protruding from the cell surface. Here, we investigated the little-known relationship between circadian rhythm and primary cilia. The length and number of primary cilia showed circadian dynamics both in vitro and in vivo. The circadian rhythm of primary cilia morphology was abolished by SR9011, a clock genes suppressor. A centrosomal protein, pericentrin, transiently accumulates in centriolar satellites, the base of primary cilia at the shortest cilia phase, and induces elongation of primary cilia at the longest cilia phase in the circadian rhythm of primary cilia. In addition, rhythmic cell migration during wound healing depends on the length of primary cilia and affects the rate of wound healing. Our findings demonstrate that the circadian dynamics of primary cilia length by clock genes control fibroblast migration and could provide new insights into chronobiology.

Introduction

Circadian rhythmicity is observed in various biological phenomena in organisms,

including the sleep/wake cycle, ranging from bacteria to mammals. This system adjusts

physiological processes to an approximately 24-h cycle, and adapts it to the external

environment of the day–night cycle. The suprachiasmatic nucleus (SCN) of

hypothalamus is believed to be a "central clock" that synchronizes the clock in every

cell and tissues in vivo (Yamazaki *et al*, 2000). Disruption of the central clock causes

various diseases, including abnormal sleep behaviors, obesity, and vascular diseases

(Nakazato *et al*, 2017; Turek *et al*, 2005; Zee & Vitiello, 2009). Circadian rhythms are

generated by a transcriptional feedback loop of "clock gene" expression. The core loop

consisted of two positive and two negative regulators. The brain and muscle aryl

hydrocarbon receptor nuclear translocator-like protein-1 (BMAL1; also known as

ARNTL) and circadian locomotor output cycles protein kaput (CLOCK) function as

two positive regulators (Gekakis *et al*, 1998). The BMAL1/CLOCK complex binds to

the E-box motif and drives the transcription of various genes, including period (PER)

and cryptochrome (CRY), which inhibit the transcriptional activity of the

BMAL1/CLOCK complex as negative regulators of the loop (Griffin *et al*, 1999;

Yamaguchi *et al*, 2000). This core feedback loop, which takes place in a 24-h cycle

generates circadian rhythmicity (Kume *et al*, 1999).

Most organs, tissues, and cells in the non-SCN also have an intrinsic circadian

clock, while the center of the circadian clock is the SCN (Shimba *et al*, 2005; Storch *et*

al, 2002; Young *et al*, 2001). Negative feedback loops by clock genes exist in individual

cells, and oscillations are observed in each single cell. The oscillations are also called an

autonomous cellular clock. The autonomous cellular clock provides circadian rhythms

for cell-specific functions in an SCN-independent manner. This clock regulates the cytoskeleton and organelles, and provides rhythmic changes in cellular functions (Mofatteh *et al*, 2021). For example, the autonomous cellular clock regulates fibroblast migration by controlling actin dynamics. Human studies have shown that burns sustained at night take longer to heal than those sustained during the day, possibly because fibroblast migration to the wound is regulated by clock genes (Hoyle *et al*, 2017). The accumulation of key mitochondrial enzymes is regulated by Per1/2 and exhibits diurnal oscillations, which are thought to adapt mitochondrial activities to changes in body activity and rest, feeding, and fasting throughout the day (Neufeld-Cohen *et al*, 2016). The formation of daughter centrioles has been reported to exhibit diurnal oscillations due to Plk4 oscillations in centrioles and centriolar duplicating independent of the cell cycle (Aydogan *et al*, 2020).

Most cells possess a primary cilium, which is a small organelle protruding from the cell surface that functions as a sensory antenna for cells to detect extracellular signals (Phua *et al*, 2015; Singla & Reiter, 2006). The primary cilium is recognized as a crucial organelle for various cellular processes including proliferation, differentiation, and migration (Ezratty *et al*, 2011; Higginbotham *et al*, 2012; Irigoín & Badano, 2011; Rohatgi *et al*, 2007). Defects in primary cilia due to congenital factors induce several diseases "ciliopathy", including obesity, polycystic kidney, and retinopathy (Reiter & Leroux, 2017). Primary cilia are immotile, but not static. In the cell cycle, primary cilia assemble in non-dividing cells in the G_0/G_1 phase and disassemble with cell cycle entry (Plotnikova *et al*, 2009). We previously reported that the decapitation of primary cilia and release of vesicles are signals that induce primary cilia disassembly and drive the cell cycle (Phua *et al*, 2017). Cilia decapitation is caused by the accumulation of

phosphatidylinositol 4,5-bisphosphate (PI(4,5)P2) on the primary cilia membrane, leading to actin polymerization at the tip of the primary cilia (Phua *et al.*, 2017; Stilling *et al*, 2022). G protein-coupled receptors (GPCRs) are localized in primary cilia and are receptive to the extracellular environment, such as growth factors. When ligands bind to GPCRs, activated GPCRs accumulate at the tip of cilia and are disposed of extracellularly by vesicle release, thereby regulating receptor signaling (Nager *et al*, 2017).

Numerous studies on the effects of loss or mutation of primary cilia on cellular function in ciliopathy have revealed much regarding the importance of the "presence" of primary cilia. Morphological dynamics (including length changes) in primary cilia may also contribute to the regulation of cellular functions and extracellular signal reception. Cell sensitivity to mechanical stimuli increases depending on the length of primary cilia (Spasic & Jacobs, 2017). Aldosterone-stimulated increase in cell size of renal collecting duct epithelial cells may be dependent on the length of primary cilia (Komarynets *et al*, 2020). Despite of accumulation of these evidence, the mechanisms regulating primary cilium morphology and the physiological significance of alterations in primary cilium morphology have not been fully elucidated. Here, we investigated the relationship between the circadian rhythm and primary cilia. We demonstrated that the length of primary cilia in cultured murine fibroblasts, neurons, and astrocytes of the murine brain exhibits circadian dynamics similar to the rhythmic oscillations of clock genes. Wound healing assay revealed that the circadian dynamics of primary cilium length regulate fibroblast migration. Our data indicate that circadian rhythms in primary cilium length produced by clock genes regulate cell migration. The findings provide new insights into chronobiology.

Results

Ciliogenesis and primary cilium length in cultured cells exhibit circadian rhythms

To investigate the relationship between primary cilia and circadian rhythm, we

analyzed changes in primary cilia morphology and clock gene expression in NIH/3T3

mouse embryonic fibroblasts. In cultured cells, primary cilia assembly is induced by

serum starvation-induced cell cycle arrest (R W Tucker, 1979). Cultured cells also have

autonomous clock gene expression rhythms, but their individual phases are disparate

(Nagoshi *et al*, 2004). Dexamethasone (DEX), a synthetic corticosteroid, induces Per1

expression and synchronizes the clock between cells (Nagoshi *et al.*, 2004). In this study,

ciliogenesis was induced by incubation in culture medium for 24 h with serum

concentration decreased from 10% to 1%, followed by 2-h exposure to DEX to

synchronize clock gene expression (Fig. 1A). The DEX-induced synchronization of

clock genes expression was verified by using Per1::Luc-expressing NIH/3T3 cells. The

cells were exposed to DEX for 2 h to synchronize the expression of clock genes.

Bioluminescence was measured every 15 min for 96 h using photomultiplier tube

detector assemblies. The oscillatory pattern of luciferase activity was observed in cells

exposed to DEX (Fig. 1B). Based on these results, we used this synchronization

regimen in subsequent experiments.

To investigate the influence of the cell intrinsic clock on primary cilia, we

performed immunostaining of primary cilia in synchronized NIH/3T3 cells. NIH/3T3

cells were synchronized by DEX, then fixed at 24, 30, 36, 42, and 48 h after switching

the medium to DEX-free one. Primary cilia were visualized by staining Arl13b, a

marker of primary cilia, and the base of primary cilia by staining γ-tubulin, a

peri-centrosome component (Fig. 1C). In NIH/3T3 cells treated with vehicle, where

conditions were unsynchronized, Arl13b-positive primary cilia were observed in approximately 40% of the total cells at all times (Fig. 1C and 1D). In DEX-treated synchronized conditions, cells with Arl13b-positive primary cilia were minimum 30 h after synchronization (32.31 ± 1.73%, Fig. 1D), and maximum 42 h after synchronization (60.70 ± 2.26%, Fig. 1D). At other time points, the proportion of ciliated cells was approximately 40% (Fig. 1D). We further examined the effect of the circadian rhythms on the length of primary cilia. The average length of the primary cilia in unsynchronized cells was approximately 3 µm (Fig. 1E). Primary cilia length in synchronized cells was shortest 30 h after synchronization (2.64 ± 0.05 µm, Fig. 1E) and longest 42 h after synchronization (3.47 ± 0.03 µm, Fig. 1E). At other time points, the length of primary cilia was approximately 3 µm (Fig. 1E). These data indicate that both the number and length of primary cilia exhibit circadian oscillations.

Circadian changes of primary cilia morphology is regulated by clock genes

To examine whether the circadian oscillation of primary cilia length is mediated by clock gene expression, NIH/3T3 cells synchronized by DEX treatment were exposed to SR9011, a synthetic agonist for Rev-Erb, which suppresses Bmal1 and Clock expression (Sulli *et al*, 2018) (Fig. 2A). SR9011 also inhibits autophagy and lipogenesis in cancer cells at 20 µM (Sulli *et al*, 2018). The blocking of autophagy is reported to cause primary cilia elongation (Pampliega & Cuervo, 2016; Pampliega *et al*, 2013). Indeed, 10 µM SR9011 induced the elongation of primary cilia (average of primary cilium length 4.17 ± 0.18 µm for 10 µM SR9011 versus 2.81 ± 0.20 µm for control) in unsynchronized cells (Fig. 2B and 2C). We thus sought out an appropriate concentration of SR9011 that did not induce primary cilia elongation, but inhibited the

circadian dynamics of primary cilia. To this end, we treated NIH/3T3 cells with 2-to-10-µM SR9011 at 30 h after synchronization, at which the shortest cilia length and maximal Bmal/Clock activity were observed. Total RNA was collected at 30 h after 2-h exposure to DEX, cDNA was synthesized by reverse transcription, and the expression level of the clock gene Bmal1 mRNA (gene symbol is *Arntl*) was measured. Two micromolar SR9011 was enough to significantly suppress Bmal1 expression (Fig. 2D).

SR9011 at 2 µM did not induce the abnormal primary cilia elongation at 24 h after synchronization (Fig. 2E). In contrast, the primary cilia shortening at 30 h after synchronization was canceled by the same concentration of SR9011 (average primary cilia length 2.15 ± 0.03 µm with DEX versus 2.82 ± 0.02 µm with DEX and SR9011) (Fig. 2E). At 1 µM, SR9011 failed to block the shortening of primary cilia 30 h after synchronization (Fig. 2E). We then examined whether the circadian oscillation of primary cilia length was abolished with SR9011 at this concentration. The elongation of primary cilia at 42 h after synchronization was also abolished by 2 µM SR9011 (3.13 ± 0.07 µm with DEX versus 2.64 ± 0.11 µm with DEX and SR9011) (Fig. 2F and 2G). Hence, SR9011 significantly suppressed the circadian oscillation of primary cilium length in NIH/3T3 cells (Fig. 2G). These results indicate that the circadian oscillation of primary cilia length is mediated by rhythmic expression of clock genes.

Primary cilium length in murine brain also exhibits circadian rhythms

Next, to assess whether primary cilium length also shows circadian dynamics *in vivo*, we performed immunostaining of murine brains collected at different time points of day and night. In the adult murine brain, marker proteins for primary cilia differ among cell types (Sterpka & Chen, 2018). Hippocampal dentate gyrus of the

murine brain was used for immunohistochemistry targeting adenylate cyclase 3 (AC3) as a primary ciliary neuronal marker, NeuN (neuronal marker), glial fibrillary acidic protein (GFAP, astroglial marker), and Arl13b (Fig. 3A). AC3-positive primary cilia were observed in NeuN-positive cells but not in GFAP-positive cells (Fig. 3A; left and middle panels). Arl13b-positive primary cilia were observed in GFAP-positive cells (Fig. 3A; right panel). Perfusion fixation was performed in adult mice using 4% paraformaldehyde (PFA) at Zeitgeber Time (ZT)5 and ZT17, followed by immunohistochemical staining. The length of primary cilia in neurons in the murine hippocampus was significantly longer at ZT17 than at ZT5 (Fig. 3B and 3C). The primary cilia length in astrocytes was also markedly longer at ZT17 than at ZT5 (Fig. 3D and 3E). These results indicate that primary cilium length *in vivo* also exhibits circadian oscillations.

Transient increase in centriolar satellites regulates circadian rhythm of primary cilium length

To investigate the mechanism by which the cell intrinsic clock regulates the length of the primary cilia, we focused on centriolar satellites. Centriolar satellites are an array of 70-100-nm granules at the base of the primary cilia that regulate the transport of centrosomal and ciliary proteins. They are also involved in the formation of primary cilia (Odabasi *et al*, 2019; Quarantotti *et al*, 2019). Pericentrin (PCNT) is a centrosomal protein that localizes to centriolar satellites through binding to pericentriolar material 1 (PCM1), and is involved in primary cilia assembly (Jurczyk *et al*, 2004). In this study, immunostaining for PCNT was performed in NIH/3T3 cells synchronized with DEX treatment (Fig. 4A). PCNT-positive centriolar satellites

transiently increased in NIH/3T3 cells 30 h after synchronization with primary cilium length minimum. The average area of PCNT-positive granules was 11.95 ± 0.55 μm^2 24 h, 29.59 ± 3.33 μm^2 30 h, 12.64 ± 0.87 μm^2 36 h, 12.02 ± 1.38 μm^2 42 h, and 13.78 ± 1.32 μm^2 48 h after synchronization (Fig. 4B). The increase in PCNT-positive centriolar satellites at 30 h was suppressed by 2 μM SR9011, with an average area of PCNT-positive granules of 17.49 ± 1.35 μm^2 in cells exposed to DEX and 2 μM SR9011 (Fig. 4A and 4B). We also performed a western blotting assay using whole cell lysate of NIH/3T3 cells synchronized with DEX to test whether the oscillation of PCNT-positive granules occurs through changes in the amount of PCNT protein. No significant difference in the amount of PCNT protein was observed at any time point after synchronization (Fig. 4C). These results indicate that the transient increase in PCNT-positive centriolar satellites observed in NIH/3T3 cells 30 h after synchronization is caused by changes in subcellular localization.

It has been reported that correct subcellular localization of centriolar satellites is necessary for the formation of primary cilia (Aydin *et al*, 2020). The elongation of primary cilia is known to take place over several hours (Westlake *et al*, 2011). To investigate the effect of centriolar satellites increasing at the base of primary cilia on the circadian dynamics of primary cilium length, we inhibited transport of centriolar satellites to the base of primary cilia. Centriolar satellites are transported to the base of primary cilia via microtubule transport (Dammermann & Merdes, 2002). We used colchicine, an inhibitor of microtubule polymerization, to block the accumulation of PCNT-positive granules at the base of the primary cilia. The effect of colchicine was confirmed by immunostaining the microtubules with anti-α-tubulin antibody. Microtubules visible in control NIH/3T3 cells (Fig. 4D; left panel) were disrupted by

200 nM colchicine (Fig. 4D; right panel). Colchicine inhibited the transient increase in PCNT-positive centriolar satellites at 30 h after synchronization in a concentration-dependent manner, to a level comparable to that of unsynchronized control cells (Fig. 4E and 4F). We then examined whether the blockade of PCNT-positive centriolar satellites accumulation at 30 h after synchronization impaired later primary cilia elongation. To this end, we exposed DEX-synchronized NIH/3T3 cells to 200 nM colchicine for 2 h at 28 h after synchronization, and subsequently measured the length of primary cilia visualized by immunostaining at 30 h and 42 h after synchronization (Fig. 4G). At 30 h after synchronization, the administration of colchicine made no significant difference to the length of primary cilia compared to control cells. However, at 42 h after synchronization (the longest cilia phase), 200 nM colchicine markedly suppressed the length of primary cilia compared to control cells (average primary cilia length 3.11 ± 0.10 µm for DEX versus 2.28 ± 0.09 µm for DEX with colchicine) (Fig 4H, 4I and 4J). These results indicate that the circadian clock controls the circadian regulation of primary cilium length by regulating the transport of centriolar satellites to the base of primary cilia.

Wound healing depends on the circadian phase of wounding, and its differences are eliminated by loss of primary cilia in NIH/3T3 cells

To investigate the physiological significance of circadian rhythms in primary cilium length, we performed a wound healing assay on synchronized NIH/3T3 cells. The cells were wounded when the primary cilium length was minimal (30 h after synchronization) or maximal (18 h or 42 h after synchronization), and the area of recovery was measured up to 24 h after wounding (Fig. 5A). We generated NIH/3T3

cells with deficient primary cilia, Kif3a knockout (KO) NIH/3T3 cells (Kif3a KO), as control references to evaluate the effect of primary cilia, and performed a wound healing assay with wild-type (WT) NIH/3T3 cells. The area of recovery in WT cells at 24 h after wounding was significantly larger for cells wounded when primary cilium length was minimal, rather than maximal (average recovery area 24.98 ± 0.66 μm^2 at 30 h, 18.46 ± 1.56 μm^2 at 18 h, and 16.16 ± 1.83 μm^2 at 42 h) (Fig. 5B and 5C). In contrast, these differences were abolished by the loss of primary cilia in Kif3a KO (average recovery area 13.17 ± 2.56 μm^2 at 30 h, 15.29 ± 2.89 μm^2 at 18 h, and 8.39 ± 0.75 μm^2 at 42 h) (Fig. 5B and 5C). These results indicate that primary cilium length exhibiting circadian rhythms is involved in the circadian dynamics of cell motility.

To investigate in detail how the length of primary cilia exhibiting circadian rhythms affects cell motility, we performed time-lapse analysis. NIH/3T3 WT cells were wounded 18 or 30 h after synchronization and were photographed every 10 min. The coordinate migration and migration speed of these cells was determined. Cells with wounds at the time when the primary cilia length is minimal (30 h after synchronization) migrated almost linearly toward the wound (Fig. 5D, upper panels, 5E, and Movie EV1). In contrast, cells with wounds at the time when the primary cilia length was maximal (18 h after synchronization) did not move much and movement was not toward the wound (Fig. 5D, lower panels, 5E, and Movie EV2). The migration rate was also faster for cells wounded 30 h after synchronization than at 18 h (Fig. 5F). These results indicate that NIH/3T3 cells with shorter primary cilium lengths migrate faster than cells with longer primary cilium lengths and move in a straight line in the direction of the wound, leading to faster wound healing.

Rhythmic oscillations of primary cilium length of NIH/3T3 cells toward wound repair is arrested

To further assess primary cilia in the wound healing assay, wounds were created in lawns of NIH/3T3 cell growth 18 or 30 h after synchronization, followed by immunostaining 12 h after wounding (Fig. 6A). At 12 h after wounding, three cell populations were observed based on differences in cell density: cells not facing the wound, cells facing the wound, and cells invading the wound area (Fig. 6B). The primary cilium length in NIH/3T3 cells wounded 30 h after synchronization was longer in cells facing the wound than in cells wounded at 18 h. The results were similar to the comparative lengths of primary cilia 42 h and 30 h after synchronization (average length: 3.67 ± 0.11 µm versus 2.27 ± 0.06 µm) (Fig. 6C and 6D). In cells wounded 30 h after synchronization, primary cilia in cells facing the wound were shorter (average length 2.31 ± 0.09 µm) than those in cells not facing the wound (Fig. 6C and 6D). The length of primary cilia in cells wounded 18 h after synchronization were a mixture of longer and shorter cilia in cells facing the wound (average length 3.18 ± 0.10 µm) (Fig. 6C and 6D). In cells invading the wound, primary cilia in cells wounded at 30 and 18 h after synchronization were absent or very short (Fig. 6C). These results indicate that the circadian rhythm of primary cilium length is maintained in cells not facing the wound, similar to the results of Figure 1E. In contrast, in cells facing the wound, the length of primary cilia was maintained at the time of wounding, indicating a lack of connection with the circadian rhythm. Furthermore, the absence or very short primary cilia in cells invading the wound area indicates that resorption of primary cilia is necessary for cell migration to repair the wound. Thus, wound healing may have been faster in cells wounded 30 h after synchronization than in those wounded 42 h after synchronization.

To further investigate the effect of primary cilium length on cell migration in

the wound healing assay, we analyzed the structure of actin filaments by fluorescent

staining of F-actin with phalloidin. One hour after wounding, no significant change in

actin polymerization was observed at either 18 or 30 h after synchronization (Fig. 6E).

However, the distance from the cell nucleus to the tip of the cell in the direction of the

wound was greater at 30 h than at 18 h after synchronization (Fig. 6E and 6F). No

significant change was observed in the angle of the direction of the wound relative to

the orientation of the primary cilia among these cells (Fig. 6E and Expanded View Fig.

1. These results indicate that short primary cilia can facilitate cell body changes in the

direction of migration in the wound healing assay.

The collective findings of the wound healing assay can be summarized as

follows. When a wound is introduced in a monolayer of cells, circadian changes in the

length of primary cilia in cells contacting the wound are arrested. In addition, cell

morphology changes in response to the wound and cell migration increase progressively

as cilia length decreases. These observations suggest that wounds heal faster at the

shortest cilia phase (30 h after synchronization) than at the longest cilia phase (18 or 42

h after synchronization) (Fig. 6G).

Discussion

The most important finding of this study is that primary cilium length exhibits

an oscillatory rhythm that coordinates with rhythmic clock gene expression. Ciliary

structure regulates cell cycle progression (Plotnikova *et al.*, 2009). Several clock genes

act as gatekeepers in the cell cycle (Matsuo *et al*, 2003). BMAL1/CLOCK controls the

cell cycle by directly regulating the expression of WEE1, an inhibitor of CDC2/cyclin

B1 (Farshadi *et al*, 2019; Obi-Ioka *et al*, 2013). PER1 inhibits Cyclin D1 (Fu *et al*, 2016). In this study, we performed all experiments under conditions in which the concentration of serum in the culture medium was reduced from 10% to 1% to induce ciliogenesis by arresting the cells in the G0 phase of the cell cycle. Circadian rhythms in primary cilium length have also been observed in neurons, which are non-dividing cells. The circadian clock oscillates independently of the cell cycle (Matsuo *et al.*, 2003). The formation of centrioles, which play a major role during cell division as the center of spindle formation, also exhibit a circadian rhythm independent of the cell cycle (Aydogan *et al.*, 2020). Our current findings indicate that circadian rhythm or primary cilium length, regulated by clock genes, is a phenomenon that is independent of the cell cycle.

In this study, we observed oscillatory rhythms in the primary cilium length in cultured cells and in the murine brain. A previous study reported a relationship between circadian rhythms and ciliary protein expression in the brain (Baldi *et al*, 2021). Further analysis is needed to elucidate whether the mechanism of primary cilia *in vivo*, as in cultured cells, is mediated by the regulation of PCNT localization to centriolar satellites by clock genes. Whether primary cilia also exhibit circadian oscillations in tissues other than the brain, such as the kidneys, requires further investigation.

SR9011 is a synthetic agonist of REV-ERB, the nuclear receptor involved in the cell-autonomous circadian transcriptional feedback loop as a transcriptional repressor. REV-ERB alters circadian behavior and the circadian pattern of core clock gene expression in the hypothalamus of mice (Solt *et al*, 2012). SR9011 applied at 20 µM is shown to inhibit autophagy in cancer cells (Sulli *et al.*, 2018). The functional interaction between primary cilia and autophagy, and the autophagy-induced

degradation of proteins essential for ciliogenesis have been described (Pampliega *et al.*, 2013). In this study, the elongation of primary cilia by high concentrations of SR9011 thus could result from both inhibition of clock genes and blockade of the autophagy process. To eliminate the possibility of autophagy involvement, we tried lower concentrations of SR9011 and found that Bmal1 mRNA expression oscillations were suppressed, even with 2 µM SR9011, especially at 30 h after synchronization. SR9011 at 2 µM did not induce excessive elongation of primary cilia. Thus, the low concentration of SR9011 could inhibit circadian oscillations of primary cilia length by inhibiting only the expression of clock genes, not autophagy. The transient increase in PCNT at the centriolar satellites 30 h after synchronization was also inhibited with 2 µM SR9011. These results suggest that oscillations in clock gene expression produce circadian rhythms in the length of primary cilia through direct regulation of PCNT expression in centriolar satellites. Further detailed analysis is required to confirm this suggestion.

PCNT is a component of centriolar satellites and contributes to centrosome stability (Quarantotti *et al.*, 2019). PCNT forms a complex with intraflagellar transport (IFT) proteins and polycystin-2 (PC2), which is reportedly required for the assembly of primary cilia (Jurczyk *et al.*, 2004). In contrast, in trisomy 21, the most prevalent human chromosomal disorder, an excessive increase in PCNT inhibits the transport of centrosome and cilia proteins, resulting in defects in cilia formation (Galati *et al*, 2018). In this study, PCNT-positive centriolar satellites transiently increased in NIH/3T3 cells 30 h after synchronization when the primary cilium length was minimal. In addition, inhibition of microtubule polymerization with colchicine suppressed the transient increase in PCNT-positive centriolar satellites and significantly suppressed the primary

cilium length maxima observed in NIH/3T3 cells 42 h after synchronization. The centriolar satellites are reported to be required for the process of primary cilium formation, but do not contribute much to the stability of primary cilia (Aydin *et al.*, 2020). Serum starvation-induced ciliogenesis is considered to begin within minutes of the onset of serum starvation, but the length of primary cilia becomes maximal 24 to 48 hours after the onset of serum starvation, indicating a time lag (Kukic *et al*, 2016; Westlake *et al.*, 2011; Zhang *et al*, 2020). The increase in the number of ciliated cells by serum starvation is suppressed in centriolar satellites-deficient cells compared to normal cells, but the difference is observed at 18 h after the onset of serum starvation (Aydin *et al.*, 2020). Although it is not reported how long it takes for the primary cilia to reach their maximum length from the formation of centriolar satellites, a time lag of about 12 h, as in the results of this study, is expected to exist. Thus, our observations indicate that clock genes confer circadian rhythms on primary cilium length through the subcellular localization of PCNT-positive centriolar satellites. Further experiments should be performed to reveal the mechanisms by which clock genes regulate the subcellular localization of centriolar satellites.

The physiological significance of the circadian rhythm of primary cilium length requires further investigation. It is possible that changes in the length of primary cilia could produce circadian rhythms in their sensitivity as antennae to extracellular signals. No such changes were observed in this study. Primary cilia reportedly play important roles in development. However, circadian rhythms in the SCN have not been observed in prenatal mice or rats (Sumová *et al*, 2006). Oscillations in clock gene expression are also not observed in embryonic stem cells and occur with cell differentiation (Yagita *et al*, 2010). Therefore, it is necessary to focus on primary cilia

and the developmental stages of immature cells to elucidate the physiological significance of primary cilia that exhibit circadian rhythms.

A previous study reported that the cellular clock in fibroblasts modulates cell migration, which affects the efficacy of wound healing (Hoyle *et al.*, 2017). Our observations from the wound healing assay also suggest a circadian rhythm in the mobilization of NIH/3T3 cells. In addition, the circadian rhythm of cell migration is absent in NIH/3T3 cells that have lost their primary cilia (Kif3a KO), suggesting that rhythmic cell mobilization depends on the circadian rhythm of primary cilium length. Previous studies have reported that primary cilia regulate cell migration as antennae by detecting chemokines, such as platelet-derived growth factor (Christensen *et al*, 2013; Schneider *et al*, 2010). Previous studies have also reported that the tips of primary cilia orient in the direction of travel of the migrating cell (Albrecht-Buehler, 1977). In this study, no correlation was observed between the orientation of primary cilia and the direction of the wound site at 1 h in the wound healing assay. Other authors reported that primary cilia regulate neural migration through centrosomal dynamics (Tsai & Gleeson, 2005). Neural migration is essential for centrosome movement toward the leading edge and cell nuclei. AC3 in primary cilia produces cAMP, which activates protein kinase A to regulate centrosome dynamics and subsequent nuclear movement (Stoufflet *et al*, 2020). In this study, we observed that NIH/3T3 cells with short primary cilia migrated faster and in a straight line into wounds than did cells with long primary cilia, resulting in faster wound repair. Previous studies have not reported a relationship between primary cilium length and cell mobilization.

In the wound healing assay that used cells synchronized by DEX, circadian rhythms of primary cilium length were maintained in cells not facing the wound. In

contrast, the length of primary cilia in cells facing the wound was not associated with the circadian rhythm and maintained their length at the time of wounding (Fig. 6G). A previous study reported that the rhythmic oscillation of clock gene expression depends on cell density and is altered in conditions of low cell density (Noguchi *et al*, 2013). Cell-to-cell communication has also been suggested to affect circadian rhythms (Fuhr *et al*, 2019). In addition, in the present study, cells that invaded wounds were devoid of primary cilia or had very short primary cilia, suggesting that cells migrating to wounds have short primary cilia. These findings further suggest that wound healing is faster in cells that are wounded when primary cilia are short, because the primary cilia remain short and fixed, allowing for rapid conversion to a cell type that can invade the wound (i.e., cells with no or very short primary cilia). In contrast, cells that are wounded when primary cilia are long take longer to convert to the cell type that invades the wound. Thus, cells facing the wound have a mixture of long and short primary cilia, and the wound appears to heal more slowly (Fig. 6G). Further experiments are needed to determine why cells that invade wounds must eliminate or shorten their primary cilia.

We hypothesize that primary cilia anchor for the centrosome. During cell division, disassembly of primary cilia could free the centrosome and allow it to move dynamically. This phenomenon is thought to occur during cell migration. Indeed, in this study, cells in the wound area had no or very short primary cilia. Our prediction from the results we obtained is that the centrosome can be easily released from short primary cilia more than from long primary cilia, and that centrosome dynamics and cell migration can commence sooner (Fig. 6G).

In conclusion, the autonomous cellular clock provides rhythmic oscillations in primary cilium length by regulating PCNT accumulation in centriolar satellites. Primary

cilia contribute to the circadian dynamics of cell migration during wound repair. Our findings contribute to an improved understanding of the functional role of the non-SCN autonomous cellular clock system and primary cilium length alterations in a variety of physiological and pathological processes involving cell migration in wound repair and ciliopathies. Thus, clock genes may represent novel targets for the discovery and development of therapies for ciliopathies.

Materials and Methods

Cell culture and treatments

NIH/3T3 cells purchased from American Type Culture Collection (CRL-1658) were cultured in Dulbecco's Modified Eagle's Medium (DMEM)-High Glucose (044-29765, Wako) supplemented with 10% fetal bovine serum (FBS; Gibco) at 37°C in an atmosphere of 5% CO_2. The workflow for the induction of ciliogenesis and clock gene synchronization is shown in Fig. 1. Ciliogenesis was induced by changing the culture medium to DMEM with 1% FBS for 24 h. Next, the cells were exposed to DEX (Sigma-Aldrich) for 2 h to synchronize the expression of clock genes. DEX was then washed out and various experiments were performed. SR9011 (Sigma-Aldrich) was added to the culture medium simultaneously with DEX and continued to be added to the medium after DEX had been washed out. Colchicine (Sigma-Aldrich) was added to the culture medium 28 h after synchronization, incubated for 2 h, and then washed out.

Generation of Kif3a KO cell line

Detailed methods for the generation of KO cell lines have been previously described (Ijaz & Ikegami, 2021; Ijaz *et al*, 2022; Phua *et al.*, 2017). Kif3a-KO cells were

generated using CRISPR/Cas9-based genome-editing techniques. To construct an all-in-one expression plasmid for KO, a 20-bp target sequence (Kif3a: CTATAGACAGGCCGTCAGCG) was subcloned into the U6-gRNA/CMV-Cas9-2A-GFP plasmid backbone (ATUM). Cells were plated at a density of 2×10^4 per cm^2 and were transfected with the all-in-one plasmid U6-gRNA/CMV-Cas9-2A-GFP using polyethylenimine. Three days after transfection, green fluorescent protein (GFP)-positive cells were sorted by fluorescence-activated cell sorting (FACS) using a FACS Aria2 SORP cell sorter (Becton Dickinson Biosciences). Single cell clones were generated by the limiting dilution culture method. Mutant cells were screened by double-strand break site-targeted PCR genotyping (Ijaz *et al.*, 2022). KO cells were identified by DNA sequencing.

Real-time qPCR

Total RNA was extracted from cells and reverse-transcribed to cDNA using the PrimeScript 1st Strand cDNA Synthesis Kit (TaKaRa Bio). cDNA was used as a template for real-time qPCR analysis on CFX96 Touch (Bio-Rad) using specific primers targeting *glyceraldehyde 3-phosphate dehydrogenase* (*Gapdh*; 5'-AGGTCGGTGTGAACGGATTTG-3' as the forward primer and 5'-TGTAGACCATGTAGTTGAGGTCA-3' as the reverse primer), *Arntl* (5'-TGACCCTCATGGAAGGTTAGAA-3' as the forward primer and 5'-GGACATTGCATTGCATGTTGG-3'), and TB Green Premix Ex Taq II (TaKaRa Bio). In each sample, expression levels of the *Arntl* (BMAL1 gene symbol) were normalized to *Gapdh* expression as an internal control.

Mice

C57BL/6J mice purchased from Japan SLC, Inc. were maintained under conditions of controlled temperature and humidity with a 12-h light/dark cycle and *ad libitum* access to standard laboratory food and water. The experiments involving mice were approved by the Natural Science Center for Basic Research and Development at Hiroshima University.

Immunocytochemistry

Cells plated on cover glass (MATSUNAMI) were fixed with 4% PFA for 15 min at 37°C or 100% MeOH for 10 min at 4°C. The cells were blocked with 5% normal goat serum containing 0.1% Triton X-100 in PBS (blocking buffer) for 1 h at room temperature (22 to 28°C). The primary antibodies were diluted in blocking buffer and incubated with cells overnight at 4°C. The cells were immunostained with anti-Arl13b (Proteintech or Abcam; both 1:1000), anti-pericentrin (Abcam; 1:400), anti-γ-tubulin (Sigma-Aldrich; GTU88; 1:400), anti-α-tubulin (Sigma-Aldrich; 1:400), and anti-5-HT6 receptor (Abcam; 1:400). The cells were washed with PBS to remove unbound antibody and incubated for 2 h with Alexa Fluor-conjugated secondary antibodies (Invitrogen; 1:500), 4′,6-diamidino-2-phenylindole (DAPI; Wako; 1:1000), and Alexa488-conjugated phalloidin (1:1000) at room temperature. After washing with PBS, the cover glasses were mounted on glass slides with VECTASHIELD mounting medium (Vector Laboratories). Images were acquired using a model DMI3000B epifluorescence microscope (Leica) and model FV1000D confocal microscope (Olympus). Quantitative image analyses were performed using ImageJ software (NIH).

Immunohistochemistry

Adult 8–12 week old male mice were deeply anesthetized with isoflurane before intracardial perfusion with PBS and 4% PFA. The brains were post-fixed overnight and cryoprotected in 30% sucrose. The brains were embedded in OCT tissue TEC (Sakura) and 30 µm-thick frozen coronal sections were cut using a model CM 3050 cryostat (Leica). The sections were blocked with blocking buffer for 1 h and incubated with primary antibodies overnight at 4°C. The antibodies used for the brain sections in this study were anti-Arl13b (Abcam; 1:400), anti-ACIII (Novus; 1:1000), anti-NeuN (Abcam; 1:400), and anti-GFAP (Abcam; 1:400). After washing with PBS, the sections were incubated for 2 h with Alexa Fluor-conjugated secondary antibody (Invitrogen, 1:400) and DAPI (Wako, 1:1000). Confocal images were acquired using a model FV1000 microscope (Olympus). Quantitative image analysis was performed using ImageJ software (NIH).

Western blotting

Cells were solubilized in 1× sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) sample buffer (Wako). Lysate samples were subjected to 6% SDS-PAGE and transferred to polyvinylidene difluoride membranes. Membranes were blocked with 5% bovine serum albumin (BSA) in Tris buffered saline containing 0.1% Tween 20 (TBST) and immunoblotted using primary antibody to pericentrin (Abcam) diluted 1:1000 in 1% BSA in TBST overnight at 4°C. After washing with TBST, the membranes were incubated for 1 h with horseradish peroxidase-conjugated secondary antibody (1:5000) and developed using ECL Prime (GE).

Wound healing assay

A scratch was made in a cell monolayer using a sterile 200 µL plastic pipette tip at the indicated time after removing DEX. Images were acquired every 6 h for 24 h after wounding. Cell closure of the wound was quantitated using ImageJ software. Live cell imaging by time-lapse analysis was performed using a model LCV110 microscope (OLYMPUS) every 10 min for 12 h at 37°C. Quantification of the coordinates of the cellular centroid and cellular moving distance was performed using ImageJ software.

Statistical analysis

All experiments and quantitative analyses were blinded and randomized to remove bias. The results are presented as the mean ± SEM. Group means were compared using Student's t-test. Statistical significance was set at $P < 0.05$.

Acknowledgments

This work was supported in part by JSPS Grants-in-Aids for Young Scientists (20K15792) to RN and for Scientific Research (C) (21K06172) to KI. We thank Madoka Hamada, Mimoko Katoh, and Yoko Hayashi for technical assistance. Experiments were performed at the Analysis Center of Life Science, Natural Science Center for Basic Research and Development, Hiroshima University, and the Research Institute for Radiation Biology and Medicine, Hiroshima University.

Author contributions

RN and KI designed experiments and wrote the manuscript. RN performed all the experiments. YM quantified the length of primary cilia in NIH/3T3 cells, immunostaining of primary cilia in murine brains, and quantification of the length of primary cilia in neurons and astrocytes.

Conflict of interest

The authors declare that there are no conflicts of interest.

References

Albrecht-Buehler G (1977) The phagokinetic tracks of 3T3 cells. *Cell* 11: 395-404

Aydin OZ, Taflan SO, Gurkaslar C, Firat-Karalar EN (2020) Acute inhibition of centriolar satellite function and positioning reveals their functions at the primary cilium. *PLoS Biol* 18: e3000679

Aydogan MG, Steinacker TL, Mofatteh M, Wilmott ZM, Zhou FY, Gartenmann L, Wainman A, Saurya S, Novak ZA, Wong SS *et al* (2020) An Autonomous Oscillation Times and Executes Centriole Biogenesis. *Cell* 181: 1566-1581 e1527

Baldi P, Alhassen W, Chen S, Nguyen H, Khoudari M, Alachkar A (2021) Large-scale analysis reveals spatiotemporal circadian patterns of cilia transcriptomes in the primate brain. *J Neurosci Res* 99: 2610-2624

Christensen ST, Veland IR, Schwab A, Cammer M, Satir P (2013) Analysis of primary cilia in directional cell migration in fibroblasts. *Methods Enzymol* 525: 45-58

Dammermann A, Merdes A (2002) Assembly of centrosomal proteins and microtubule organization depends on PCM-1. *J Cell Biol* 159: 255-266

Ezratty EJ, Stokes N, Chai S, Shah AS, Williams SE, Fuchs E (2011) A role for the primary cilium in Notch signaling and epidermal differentiation during skin development. *Cell* 145: 1129-1141

Farshadi E, Yan J, Leclere P, Goldbeter A, Chaves I, van der Horst GTJ (2019) The positive circadian regulators CLOCK and BMAL1 control G2/M cell cycle transition through Cyclin B1. *Cell Cycle* 18: 16-33

Fu XJ, Li HX, Yang K, Chen D, Tang H (2016) The important tumor suppressor role of PER1 in regulating the cyclin-CDK-CKI network in SCC15 human oral squamous cell carcinoma cells. *Onco Targets Ther* 9: 2237-2245

Fuhr L, Abreu M, Carbone A, El-Athman R, Bianchi F, Laukkanen MO, Mazzoccoli G, Relógio A (2019) The Interplay between Colon Cancer Cells and Tumour-Associated Stromal Cells Impacts the Biological Clock and Enhances Malignant Phenotypes. *Cancers (Basel)* 11

Galati DF, Sullivan KD, Pham AT, Espinosa JM, Pearson CG (2018) Trisomy 21

Represses Cilia Formation and Function. *Dev Cell* 46: 641-650 e646

Gekakis N, Staknis D, Nguyen HB, Davis FC, Wilsbacher LD, King DP, Takahashi JS, Weitz CJ (1998) Role of the CLOCK protein in the mammalian circadian mechanism. *Science* 280: 1564-1569

Griffin EA, Jr., Staknis D, Weitz CJ (1999) Light-independent role of CRY1 and CRY2 in the mammalian circadian clock. *Science* 286: 768-771

Higginbotham H, Eom TY, Mariani LE, Bachleda A, Hirt J, Gukassyan V, Cusack CL, Lai C, Caspary T, Anton ES (2012) Arl13b in primary cilia regulates the migration and placement of interneurons in the developing cerebral cortex. *Dev Cell* 23: 925-938

Hoyle NP, Seinkmane E, Putker M, Feeney KA, Krogager TP, Chesham JE, Bray LK, Thomas JM, Dunn K, Blaikley J *et al* (2017) Circadian actin dynamics drive rhythmic fibroblast mobilization during wound healing. *Sci Transl Med* 9

Ijaz F, Ikegami K (2021) Knock-in of Labeled Proteins into 5'UTR Enables Highly Efficient Generation of Stable Cell Lines. *Cell Struct Funct* 46: 21-35

Ijaz F, Nakazato R, Setou M, Ikegami K (2022) A pair of primers facing at the double-strand break site enables to detect NHEJ-mediated indel mutations at a 1-bp resolution. *Sci Rep* 12: 11681

Irigoín F, Badano JL (2011) Keeping the balance between proliferation and differentiation: the primary cilium. *Curr Genomics* 12: 285-297

Jurczyk A, Gromley A, Redick S, San Agustin J, Witman G, Pazour GJ, Peters DJ, Doxsey S (2004) Pericentrin forms a complex with intraflagellar transport proteins and polycystin-2 and is required for primary cilia assembly. *J Cell Biol* 166: 637-643

Komarynets O, Chassot A, Bernabeu E, Czogalla J, Roth I, Liaudet N, Prodon F, Loffing J, Feraille E (2020) Aldosterone controls primary cilium length and cell size in renal collecting duct principal cells. *FASEB J* 34: 2625-2640

Kukic I, Rivera-Molina F, Toomre D (2016) The IN/OUT assay: a new tool to study ciliogenesis. *Cilia* 5: 23

Kume K, Zylka MJ, Sriram S, Shearman LP, Weaver DR, Jin X, Maywood ES, Hastings MH, Reppert SM (1999) mCRY1 and mCRY2 are essential components of the negative

limb of the circadian clock feedback loop. *Cell* 98: 193-205

Matsuo T, Yamaguchi S, Mitsui S, Emi A, Shimoda F, Okamura H (2003) Control mechanism of the circadian clock for timing of cell division in vivo. *Science* 302: 255-259

Mofatteh M, Echegaray-Iturra F, Alamban A, Dalla Ricca F, Bakshi A, Aydogan MG (2021) Autonomous clocks that regulate organelle biogenesis, cytoskeletal organization, and intracellular dynamics. *Elife* 10

Nager AR, Goldstein JS, Herranz-Perez V, Portran D, Ye F, Garcia-Verdugo JM, Nachury MV (2017) An Actin Network Dispatches Ciliary GPCRs into Extracellular Vesicles to Modulate Signaling. *Cell* 168: 252-263 e214

Nagoshi E, Saini C, Bauer C, Laroche T, Naef F, Schibler U (2004) Circadian gene expression in individual fibroblasts: cell-autonomous and self-sustained oscillators pass time to daughter cells. *Cell* 119: 693-705

Nakazato R, Kawabe K, Yamada D, Ikeno S, Mieda M, Shimba S, Hinoi E, Yoneda Y, Takarada T (2017) Disruption of Bmal1 Impairs Blood-Brain Barrier Integrity via Pericyte Dysfunction. *J Neurosci* 37: 10052-10062

Neufeld-Cohen A, Robles MS, Aviram R, Manella G, Adamovich Y, Ladeuix B, Nir D, Rousso-Noori L, Kuperman Y, Golik M *et al* (2016) Circadian control of oscillations in mitochondrial rate-limiting enzymes and nutrient utilization by PERIOD proteins. *Proc Natl Acad Sci U S A* 113: E1673-1682

Noguchi T, Wang LL, Welsh DK (2013) Fibroblast PER2 circadian rhythmicity depends on cell density. *J Biol Rhythms* 28: 183-192

Obi-Ioka Y, Ushijima K, Kusama M, Ishikawa-Kobayashi E, Fujimura A (2013) Involvement of Wee1 in the circadian rhythm-dependent intestinal damage induced by docetaxel. *J Pharmacol Exp Ther* 347: 242-248

Odabasi E, Gul S, Kavakli IH, Firat-Karalar EN (2019) Centriolar satellites are required for efficient ciliogenesis and ciliary content regulation. *EMBO Rep* 20

Pampliega O, Cuervo AM (2016) Autophagy and primary cilia: dual interplay. *Curr Opin Cell Biol* 39: 1-7

Pampliega O, Orhon I, Patel B, Sridhar S, Díaz-Carretero A, Beau I, Codogno P, Satir BH, Satir P, Cuervo AM (2013) Functional interaction between autophagy and ciliogenesis. *Nature* 502: 194-200

Phua SC, Chiba S, Suzuki M, Su E, Roberson EC, Pusapati GV, Schurmans S, Setou M, Rohatgi R, Reiter JF *et al* (2017) Dynamic Remodeling of Membrane Composition Drives Cell Cycle through Primary Cilia Excision. *Cell* 168: 264-279 e215

Phua SC, Lin YC, Inoue T (2015) An intelligent nano-antenna: Primary cilium harnesses TRP channels to decode polymodal stimuli. *Cell Calcium* 58: 415-422

Plotnikova OV, Pugacheva EN, Golemis EA (2009) Chapter 7 - Primary Cilia and the Cell Cycle. In: *Methods in Cell Biology*, Sloboda R.D. (ed.) pp. 137-160. Academic Press:

Quarantotti V, Chen JX, Tischer J, Gonzalez Tejedo C, Papachristou EK, D'Santos CS, Kilmartin JV, Miller ML, Gergely F (2019) Centriolar satellites are acentriolar assemblies of centrosomal proteins. *Embo j* 38: e101082

R W Tucker ABP, K Fujiwara (1979) Centriole ciliation is related to quiescence and DNA synthesis in 3T3 cells. *Cell* 17: 527-535

Reiter JF, Leroux MR (2017) Genes and molecular pathways underpinning ciliopathies. *Nat Rev Mol Cell Biol* 18: 533-547

Rohatgi R, Milenkovic L, Scott MP (2007) Patched1 regulates hedgehog signaling at the primary cilium. *Science* 317: 372-376

Schneider L, Cammer M, Lehman J, Nielsen SK, Guerra CF, Veland IR, Stock C, Hoffmann EK, Yoder BK, Schwab A *et al* (2010) Directional cell migration and chemotaxis in wound healing response to PDGF-AA are coordinated by the primary cilium in fibroblasts. *Cell Physiol Biochem* 25: 279-292

Shimba S, Ishii N, Ohta Y, Ohno T, Watabe Y, Hayashi M, Wada T, Aoyagi T, Tezuka M (2005) Brain and muscle Arnt-like protein-1 (BMAL1), a component of the molecular clock, regulates adipogenesis. *Proc Natl Acad Sci U S A* 102: 12071-12076

Singla V, Reiter JF (2006) The primary cilium as the cell's antenna: signaling at a sensory organelle. *Science* 313: 629-633

Solt LA, Wang Y, Banerjee S, Hughes T, Kojetin DJ, Lundasen T, Shin Y, Liu J, Cameron MD, Noel R *et al* (2012) Regulation of circadian behaviour and metabolism by synthetic REV-ERB agonists. *Nature* 485: 62-68

Spasic M, Jacobs CR (2017) Lengthening primary cilia enhances cellular mechanosensitivity. *Eur Cell Mater* 33: 158-168

Sterpka A, Chen X (2018) Neuronal and astrocytic primary cilia in the mature brain. *Pharmacol Res* 137: 114-121

Stilling S, Kalliakoudas T, Benninghoven-Frey H, Inoue T, Falkenburger BH (2022) PIP(2) determines length and stability of primary cilia by balancing membrane turnovers. *Commun Biol* 5: 93

Storch KF, Lipan O, Leykin I, Viswanathan N, Davis FC, Wong WH, Weitz CJ (2002) Extensive and divergent circadian gene expression in liver and heart. *Nature* 417: 78-83

Stoufflet J, Chaulet M, Doulazmi M, Fouquet C, Dubacq C, Métin C, Schneider-Maunoury S, Trembleau A, Vincent P, Caillé I (2020) Primary cilium-dependent cAMP/PKA signaling at the centrosome regulates neuronal migration. *Sci Adv* 6

Sulli G, Rommel A, Wang X, Kolar MJ, Puca F, Saghatelian A, Plikus MV, Verma IM, Panda S (2018) Pharmacological activation of REV-ERBs is lethal in cancer and oncogene-induced senescence. *Nature* 553: 351-355

Sumová A, Bendová Z, Sládek M, El-Hennamy R, Laurinová K, Jindráková Z, Illnerová H (2006) Setting the biological time in central and peripheral clocks during ontogenesis. *FEBS Lett* 580: 2836-2842

Tsai LH, Gleeson JG (2005) Nucleokinesis in neuronal migration. *Neuron* 46: 383-388

Turek FW, Joshu C, Kohsaka A, Lin E, Ivanova G, McDearmon E, Laposky A, Losee-Olson S, Easton A, Jensen DR *et al* (2005) Obesity and metabolic syndrome in circadian Clock mutant mice. *Science* 308: 1043-1045

Westlake CJ, Baye LM, Nachury MV, Wright KJ, Ervin KE, Phu L, Chalouni C, Beck JS, Kirkpatrick DS, Slusarski DC *et al* (2011) Primary cilia membrane assembly is initiated by Rab11 and transport protein particle II (TRAPPII) complex-dependent

trafficking of Rabin8 to the centrosome. *Proc Natl Acad Sci U S A* 108: 2759-2764

Yagita K, Horie K, Koinuma S, Nakamura W, Yamanaka I, Urasaki A, Shigeyoshi Y, Kawakami K, Shimada S, Takeda J *et al* (2010) Development of the circadian oscillator during differentiation of mouse embryonic stem cells in vitro. *Proc Natl Acad Sci U S A* 107: 3846-3851

Yamaguchi S, Mitsui S, Miyake S, Yan L, Onishi H, Yagita K, Suzuki M, Shibata S, Kobayashi M, Okamura H (2000) The 5' upstream region of mPer1 gene contains two promoters and is responsible for circadian oscillation. *Curr Biol* 10: 873-876

Yamazaki S, Numano R, Abe M, Hida A, Takahashi R, Ueda M, Block GD, Sakaki Y, Menaker M, Tei H (2000) Resetting central and peripheral circadian oscillators in transgenic rats. *Science* 288: 682-685

Young ME, Razeghi P, Cedars AM, Guthrie PH, Taegtmeyer H (2001) Intrinsic diurnal variations in cardiac metabolism and contractile function. *Circ Res* 89: 1199-1208

Zee PC, Vitiello MV (2009) Circadian Rhythm Sleep Disorder: Irregular Sleep Wake Rhythm Type. *Sleep Med Clin* 4: 213-218

Zhang C, Li C, Siu GKY, Luo X, Yu S (2020) Distinct Roles of TRAPPC8 and TRAPPC12 in Ciliogenesis via Their Interactions With OFD1. *Front Cell Dev Biol* 8: 148

Figure legends

Figure 1 – Length of primary cilia in cultured cells exhibits circadian rhythm.

A) Schematic diagram of the protocol for induction of ciliogenesis by serum starvation

and synchronization of clock gene expression by exposure of cultured NIH/3T3 cells to

dexamethasone (DEX).

B) Real-time luciferase bioluminescence assay for Per1 promoter activity levels in

Per1::Luc-expressing NIH/3T3 cells exposed to DEX as in A.

C) Immunostaining for primary cilia in NIH/3T3 cells at the indicated time points

following treatment with the protocol used in panel A. Red denotes labeling of cilia by

anti-Arl13b antibody, green denotes labeling of centrosomes by anti-γ-tubulin antibody,

and blue denotes labeling of DNA by DAPI. White arrows point to primary cilia. Scale

bar, 20 µm.

D) Quantitative analysis of the percentage of Arl13b$^+$ NIH/3T3 cells at each indicated

time point in panel C. The data are from three independent experiments.

E) Quantitative analysis of the primary cilium length in NIH/3T3 cells at each indicated

time point in panel C. The data are from three independent experiments. Representative

views of DEX-synchronized primary cilia are provided on the graph. Scale bar, 2 µm.

Data in panels D and E are presented as mean ± SEM. *P≤0.05 and **P≤0.01 (Student's

t-test).

Figure 2 – SR9011 inhibits the circadian rhythms of primary cilium length.

A) Schematic diagram of molecular core clock mechanism and SR9011 pharmacology.

B) Immunostaining of primary cilia in NIH/3T3 cells treated with 10 µM SR9011. Scale

bar, 5 µm.

C) Quantitative analysis of length of primary cilia in panel B. The data are from three independent experiments.

D) Real-time qPCR analysis of Bmal1 mRNA expression levels in the NIH/3T3 cells incubated with SR9011 at 30 h after release from DEX.

E) Quantitative analysis of primary cilium length in NIH/3T3 cells treated with the indicated concentration of SR9011 24 or 30 h after release from DEX, as in panel A. The data are from over 50 cells per condition.

F) Immunostaining for primary cilia in NIH/3T3 cells treated with 2 µM SR9011 30 or 42 h after release from DEX. Scale bar, 5 µm.

G) Quantitative analysis of primary cilium length in NIH/3T3 cells at each indicated time point after release from DEX in panel F. The data are from three independent experiments.

Data in panels C, D and G are presented as mean ± SEM. *P≤0.05 and **P≤0.01 (Student's t-test).

Figure 3 – Length of primary cilia also exhibits circadian rhythm in murine brain.

A) Immunostaining for primary cilia on neurons (green; labeled by anti-AC3 antibody) and neural markers (red; labeled with anti-NeuN antibody) or primary cilia on astrocytes (green; labeled by anti-Arl13b antibody) and astroglial markers (red; labeled by anti-GFAP antibody) in murine brain hippocampus. Scale bar, 20 µm.

B) Immunostaining for primary cilia in neurons in the hippocampus of perfusion-fixed murine brains at ZT5 or ZT17. Scale bar, 20 µm.

C) Quantitative analysis of primary cilium length in panel B (n = 50 cilia per condition).

D) Immunostaining for primary cilia on astrocytes in the hippocampus of

perfusion-fixed murine brains at ZT5 or ZT17. Scale bar, 5 µm.

E) Quantitative analysis of primary cilium length from panel D (n = 15 cilia per

condition).

Data in panels C and E are presented as mean ± SEM. **P≤0.01 (Student's t-test).

Figure 4 – Pericentrin accumulates around the base of shortest primary cilia.

A) Immunostaining for primary cilia and pericentrin. Arl13b is stained red, pericentrin

(PCNT) green, and DNA blue (by DAPI) in NIH/3T3 cells at 30 and 42 h after release

from DEX. Scale bar, 5 µm. White dotted lines enclose PCNT-positive granules.

B) Quantitative analysis of areas positive for PCNT in NIH/3T3 cells treated with

SR9011 or dimethyl sulfoxide (DMSO) as vehicle at the indicated time points after

release from DEX (N = 3 independent experiments).

C) Western blot analyses of expression of PCNT and GAPDH (as an internal standard)

in NIH/3T3 cells at each indicated time point after release from DEX.

D) Immunostaining for microtubules (green; labeled with anti-α-tubulin antibody) and

DNA (blue; DAPI) in NIH/3T3 cells incubated with 200 nM colchicine or DMSO as

vehicle for 2 h. Scale bar, 10 µm.

E) Immunostaining for primary cilia (red, Arl13b; green, PCNT) in NIH3T3 cells

treated with 200 nM colchicine or DMSO 30 h after release from DEX. White dotted

lines enclose PCNT-positive granules. Scale bar, 20 µm.

F) Quantitative analysis of the areas positive for PCNT in NIH/3T3 cells with the

indicated concentration of colchicine 30 h after release from DEX (N = 3 independent

experiments).

G) Schematic diagram of the protocol for NIH/3T3 cells synchronized with DEX and

exposure to colchicine.

H) Immunostaining of primary cilia (red; Arl13b) and centrosomes (green; γ-tubulin) in NIH/3T3 cells treated with 200 nM colchicine or DMSO 42 h after release from DEX. Scale bar, 20 µm.

I) Representative views of primary cilia for panel J. Scale bar, 2 µm

J) Quantitative analysis of primary cilium length in NIH/3T3 cells treated with 200 nM colchicine or DMSO 30 or 42 h after release from DEX (N = four independent experiments).

In panels B, F, and J, data are presented as mean ± SEM. *P≤0.05 **P≤0.01 #P≤0.05 ## P≤0.01 (Student's t-test).

Figure 5 – Primary cilia influence fibroblast mobilization with circadian changes in wound healing.

A) Schematic diagram of the protocol for wound healing assay using NIH/3T3 cells exposed to DEX.

B) Representative images of NIH/3T3 WT and Kif3a KO 0 or 24 h after wounding in the wound healing assay with the protocol illustrated in panel A. Scale bar, 200 µm.

C) Quantitative analysis of the area of wound healing from panel B at the indicated time points after wounding. The data are from three independent experiments. Data are presented as the mean ± SEM. *P≤0.05; n.s. indicates no significant difference (Student's t-test).

D) Representative time-lapse images of NIH/3T3 WT at the indicated time after wounding in the wound healing assay with the protocol illustrated in panel A. Scale bar, 50 µm.

E and F) Quantitative analysis of the coordinates of the cellular centroid (E) and the moving distance of cells (F) from panel D (The data represent three cells per condition from three individual experiments).

Figure 6 – Rhythmic dynamics of primary cilia in wound healing.

A) Schematic diagram of experimental protocol after the wound healing assay for NIH/3T3 cells exposed to DEX.

B) Representative fluorescence image of DAPI-stained NIH/3T3 12 h after wounding using the protocol illustrated in panel A. Scale bar, 200 μm.

C) Immunostaining for primary cilia (red; Arl13b) in NIH/3T3 cells at the indicated area (I-III as in B) with the protocol illustrated in panel A.

D) Quantitative analysis of primary cilium length from panel C. The data are from over 60 cells per condition from two individual experiments.

E) Immunostaining for primary cilia (red; Arl13b), F-actin (green; phalloidin), and DNA (blue; DAPI) in NIH/3T3 cells 1 h after wounding with the protocol illustrated in panel A. Arrows indicate the direction and length between the nucleus and the migrating edge of cell. Scale bar, 20 μm.

F) Quantitative analysis of the distance from the nucleus to the migrating edge of cell from panel E (white arrows in panel E). The data are from 27 cells per condition.

G) Hypothetical schematic diagram of the relationship between primary cilia and cell migration showing circadian rhythms.

Figure EV1 – Primary cilia angle in wound healing.

A) Schematic diagram of the angle of primary cilia relative to the direction of the

wound.

B) Quantitative analysis of the angle of primary cilia from panel E in Figure 6. The data are from 19 cells at 18 h and 14 cells at 30 h.

Figure 1

Figure 2

Figure 3

(A)

NeuN / AC3 / DAPI   GFAP / AC3 / DAPI   GFAP / Arl13b / DAPI

(B)

NeuN / AC3 / DAPI

ZT 17 (2:00 a.m.)   ZT 5 (2:00 p.m.)

(C)

(D)

GFAP / Arl13b / DAPI

ZT 17 (2:00 a.m.)   ZT 5 (2:00 p.m.)

(E)

Figure 4

Figure 5

Figure 6

(A)

(B) 12 h after wound

(C) Arl13b

(D)

(E) 1 h after wounding
Arl13b / F-actin / DAPI

(F)

(G) [Wound lock cilia oscillation]

[Cilia need to be shortened to migrate]