Two avian *Plasmodium* species trigger different transcriptional responses on their
vector *Culex pipiens*

Running title: *Cx. pipiens genic response to Plasmodium*

Garrigós M[1,2*], Ylla G[3], Martínez-de la Puente J[2,4], Figuerola J[1,4], Ruiz-López MJ[1,4].

[1] Department of Wetland Ecology, Estación Biológica de Doñana, CSIC, Sevilla
(Spain).

[2] Department of Parasitology, University of Granada, Granada (Spain).

[3] Bioinformatics and Genome Biology Lab, Faculty of Biochemistry, Biophysics and
Biotechnology, Jagiellonian University, Krakow (Poland).

[4] CIBER Epidemiologia y Salud Pública (CIBERESP), Madrid (Spain).

*Corresponding author: Marta Garrigós garrigosm@ugr.es

Abstract

Malaria is a mosquito-borne disease caused by protozoans of the genus

Plasmodium that affects both humans and wildlife. The fitness consequences of

infections by avian malaria are well known in birds, however, little information exists

on its impact on mosquitoes. Here we study how *Culex pipiens* mosquitoes

transcriptionally respond to infection by two different *Plasmodium* species, *P. relictum*

and *P. cathemerium*, differing in their virulence (mortality rate) and transmissibility

(parasite presence in exposed mosquitoes' saliva). We study the mosquito response to

the infection at three critical stages of parasite development: formation of ookinetes at

24 hours post-infection (hpi), the release of sporozoites into the hemocoel at 10 days

post-infection (dpi), and storage of sporozoites in the salivary glands at 21dpi. For each

time point, we characterized the gene expression of mosquitoes infected with each *P.*

relictum and *P. cathemerium* and mosquitoes fed on an uninfected bird and,

subsequently, compared their transcriptomic responses. Differential gene expression

analysis showed most of the transcriptomic changes occurred during the early infection

stage (24 hpi), especially when comparing *P. relictum* and *P. cathemerium* infected

mosquitoes. Differentially expressed genes in mosquitoes infected with each species

were related mainly to the immune response, trypsin, and other serine-proteases

metabolism. We conclude that these differences in response may partly play a role in

the differential virulence and transmissibility previously observed in *P. relictum* and *P.*

cathemerium in *Cx. pipiens*.

Keywords: avian malaria, *Plasmodium relictum*, *Plasmodium cathemerium*, vector-

borne parasites, mosquito transcriptome, RNAseq.

1. Introduction

Vector-borne diseases are a major challenge for both human and animal health, representing more than 17% of emerging infectious diseases (WHO 2021). Mosquitoes are vectors of relevant pathogens including viruses that cause yellow fever, dengue, or West Nile fever and parasites such as nematode worms which cause lymphatic filariasis, and haemosporidians, like *Plasmodium*, which cause malaria (Lehane 2010). Malaria is one of the most important vector-borne diseases for humans, and only in 2020, it caused around 602,000 deaths (WHO 2021). Malaria parasite species affect humans and related primates as well as other mammals, reptiles, or birds, driving some populations to extinction (van Riper et al. 1986).

Avian malaria is a worldwide distributed mosquito-borne disease caused by *Plasmodium* parasites which use birds as obligate hosts (Valkiūnas 2005). Avian *Plasmodium* is transmitted by mosquitoes, mainly of the genus *Culex,* which are the definitive hosts and where *Plasmodium* reproduces sexually (Valkiūnas 2005). In birds, as a part of their life cycle, *Plasmodium* merozoites invade the host erythrocytes and can differentiate into mature gametocytes (Sinden 1983). When a female mosquito feeds on a *Plasmodium*-infected bird, gametocytes are released from the erythrocytes, and male and female gametes unite resulting first in the zygote, and then in the motile ookinete (Sinden 2002). About 24 hours post-infection (hpi) the ookinete crosses the midgut epithelium and remains located between the epithelial surface and the basal lamina, where it transforms into a sessile oocyst. After about 10 days post-infection (dpi), mature oocysts liberate thousands of sporozoites into the hemocoel. Sporozoites that survive the immune system of the mosquito eventually invade the salivary glands (Sinden 1983; Sinden 2002; Abraham and Jacobs-Lorena 2004) from where they can be then transmitted to a new avian host upon a mosquito bite.

For an appropriate development and transmission of the parasites, mosquitoes

susceptible to the infection need to survive it, allowing parasites to complete their life

cycle. Thus, the nature of the interaction between pathogens and mosquitoes determines

the ability of a vector to acquire, maintain and transmit parasites to a new host, i.e. the

vector competence (Beerntsen et al. 2000; Bonizzoni et al. 2013). Therefore, the vector

competence is conditioned on the mosquito response against the pathogen (Higgs and

Beaty 2005), which includes a number of defense mechanisms. After a blood meal,

mosquitoes synthesize and release serine proteases that constitute a chemical barrier

against pathogens (Vizioli et al. 2001; Molina-Cruz et al. 2005; Muller et al. 1995) and

form a chitin-containing peritrophic matrix around the blood. This matrix constitutes a

physical barrier for harmful food particles, digestive enzymes, and pathogens (Lehane

1997). Despite the existence of these and other barriers, some pathogens manage to

reach the mosquito midgut, hemocoel, or internal organs and activate the immune

response, which may be categorized into cellular and humoral immune responses

(Hillyer 2016). The cellular response includes mechanisms such as phagocytosis,

cellular encapsulation, autophagy, melanization, and induction of apoptosis, while the

humoral response consists of the activation of signaling pathways that eventually result

in the synthesis of factors with antimicrobial activity (Michel and Kafatos 2005; Hillyer

2016). In insects, one of the main immune signaling pathways is the Toll pathway,

which is activated by the Spätzle cytokine and results in the activation of the

transcription of antimicrobial peptides (AMPs) and other immune effectors that may

combat pathogens (Kumar et al. 2018). In addition, two other pathways have been

shown to be involved in the response to *Plasmodium*, including avian malaria parasites,

the immune deficiency (Imd) and the Janus Kinase signal transducer of activation (JAK-

STAT) (Clayton et al. 2014, García-Longoria et al. 2022).

The infection by *Plasmodium* and the mosquito response against infection

ultimately result in a cost to vectors (Ahmed et al. 2002), which might depend, among

other factors, on which *Plasmodium* species infects the mosquito. For example,

Gutiérrez-López et al. (2020), found a higher survival rate and transmissibility

(measured as the presence of parasite DNA in the saliva of mosquitoes) in *Culex pipiens*

infected with *Plasmodium cathemerium* compared to those infected with *Plasmodium*

relictum. However, the genetic mechanisms that underlay these phenotypic differences

are unknown. Part of the heterogeneity in parasite virulence and therefore in the fitness

consequences in their hosts (Gutiérrez-López et al. 2020) is due to the remarkable

cellular plasticity and transcriptional variation of *Plasmodium* (García-Longoria et al.

2020). Avian *Plasmodium* is an extremely diverse clade with at least 55 species

(Valkiūnas and Iezhova 2018) divided into more than 1,446 unique genetic lineages

(Bensch et al. 2009). This extensive inter and intra-specific genomic variation will

translate into different phenotypic characters, including differences of virulence, which

will also determine how the mosquitoes respond to an infection.

Although studies on avian malaria that use transcriptomic approaches are

increasing in the last few years, they have mainly focused on the avian host, addressing

either the bird response to infection (e.g. Videvall et al. 2015) or the gene expression of

Plasmodium infecting birds (e.g. Videvall et al. 2017; García-Longoria et al. 2020;

Videvall et al. 2021). In contrast, to the best of our knowledge, only three studies have

analysed the gene expression of mosquitoes infected by avian *Plasmodium* (Zou et al.

2011; Ferreira et al. 2022; García-Longoria et al. 2022). However, none of them focus

on *Cx. pipiens*, which is the main vector of avian *Plasmodium* in Europe. Another

significant limitation is that all of these studies focus on *P. relictum* and consequently

do not take into consideration the effect of the differences in virulence and

transmissibility between *Plasmodium* lineages.

Here, we analyse the gene expression of *Cx. pipiens* infected with two widely

distributed species of avian *Plasmodium* with different characteristics namely, *P.*

relictum (lineage SGS1) and *P. cathemerium* (lineage PADOM02). To that end, we

obtained RNA-seq data of mosquitoes infected with the two *Plasmodium* species at

three time points corresponding with key stages of parasite development in the vector.

These stages are (1) during the formation of ookinetes, (2) during the release of

sporozoites into the hemocoel, and (3) after sporozoites invade and are stored in the

salivary glands. This study uses a natural avian malaria system that will help to address

the current knowledge gaps on molecular mechanisms occurring during *Plasmodium*

infections in mosquitoes. In addition, these results will allow us to further understand

the differences in *Cx. pipiens* response against different *Plasmodium* species and how

this might affect vector competence.

2. Material and Methods
130

Sampling and experimental conditions

We captured juvenile house sparrows (*Passer domesticus*) with mist nets in

September 2020 at Granja Escuela de Trigueros (Huelva province, Spain). We ringed,

weighted and measured the individuals before bringing them into captivity at the animal

facilities of the Doñana Biological Station, following the ethical guidelines (article 34

RD 53/2013).

In the field, we took blood samples from each bird jugular vein using a sterile

syringe, taking both blood samples and blood smears. Blood samples were used to

visually and molecularly identify the blood parasite infections and the parasite lineage

identity. To do that, we extracted genomic DNA from blood samples using a Lithium

Chloride protocol (Gemmell and Akiyama 1996) and detected parasite infections

following Hellgren et al. (2004). We sequenced the amplified products for positive

samples on both strands using Capillary Electrophoresis Sequencing by Macrogen

(Madrid, Spain). We analysed the sequences using Geneious v. 2020.0.3 (Kearse et al.

2012) and identified lineages in MalAvi (Bensch et al. 2009). After molecular analyses,

we chose three birds for further analyses, namely: A bird that was not infected by

Plasmodium, *Haemoproteus* nor *Leucocytozoon* (control), a bird infected with *P.*

relictum lineage SGS1, and a bird infected with *P. cathemerium* lineage PADOM02. To

visually assess the intensity of parasitemia we analyzed the blood smears of the selected

birds infected by *P. cathemerium* and *P. relictum* using a light microscope Olympus

CX33. The intensity of parasitemia was less than 1 parasite cell/15.000 erytrocites in

both cases.

We collected mosquito larvae on October 2020 in Aljaraque (Huelva province)

and reared them following Gutiérrez-López et al. (2020). We maintained larvae in

dechlorinated water and fed *ad libitum* with Hobby-Mikrozell 20 ml/22 g (Dohse

Aquaristik GmbH & Co.101 KG, D-53501, Gelsdorf, Germany) and Hobby-Liquizell

50 ml (Dohse Aquaristik GmbH & Co.101 KG, D-53501, Gelsdorf, Germany). After

emergence, we identified adults to the species level and sexed them following Gunay et

al. (2018). We kept adult *Cx. pipiens* females in separate cages of 50 individuals

maximum and fed them with a 10% sugar solution. We maintained both larvae and

adult mosquitoes under controlled conditions (26ºC ± 1, 55-60% relative humidity (RH)

and 12:12 light:dark photoperiod cycle).

We divided the 11-day-old adults (± 1 day) into 3 groups mixing individuals

originating from the different cages and allowed them to feed overnight on a *P.*

relictum-infected bird, a *P. cathemerium*-infected bird and an uninfected control bird.

Only one individual of each category was used in this experiment and they were

exposed to mosquitoes only one time. In the morning after exposure, we separated the

fed females of each group into three different cages and maintained them under the

conditions described above.

For transcriptome analyses, we processed mosquitoes at three time points after

exposure (24 hpi, 10 dpi and 21 dpi). At each time point, we created pools of 5

mosquitoes of each infection status capturing the mosquitoes alive and immediately

transferring them to dry ice. To ensure that 24h had passed since feeding, the degree of

blood digestion was assessed according to the Sella scale, following Detinova et al.

1962. , and selecting mosquitos classified as Sella stage IV. We preserved the

mosquitoes at -80C until RNA extractions were carried out. We collected a total of 36

samples including controls (4 pools x 3 time-points x 3 conditions).

RNA extraction, library preparation, and sequencing

We extracted RNA and DNA from pools of 5 mosquitoes using TRIzol®

(Invitrogen, Carlsbad, CA, USA) followed by column purification using RNeasy mini

kit® (QIAGEN, Hilden, Germany) following Ferreira et al. (2022). We tested the

remaining DNA for the presence of *Plasmodium* following Hellgreen et al. (2004)

confirming that parasite DNA was present in all positive samples and not present in

negative samples. We prepared RNAseq libraries at the Polo d'Innovazione di

Genomica, Genetica e Biologia, Siena (Italy) using and Illumina library. Then, we

quantified samples using Qubit® 4.0 Fluorometer and checked RNA integrity using the

Fragment Analyzer to measure RNA Quality Number (RQN) specific for mosquitoes.

Finally, we prepared the libraries following the QIAseqTM Stranded mRNA Selected

Kit Handbook for Illumina Paired-End Indexed Sequencing. Indexed DNA libraries

were sequenced in an Illumina NextSeq550 Flowcell, using the Illumina chemistry

V2.5, 2x75bp run.

Data analysis
193

To check the quality of the reads we used FastQC (ver. 0.11.9; Andrews et al.

2010) and MultiQC (Ewels et al. 2016). Then, we filtered low quality and under 36 bp

reads using Trimmomatic (Bolger et al. 2014). Because the reference genome and

annotations of *Cx. pipiens* are not published yet, we used the reference genome and

annotations of phylogenetically closest species that were available in Ensembl, *Cx.*

quinquefasciatus (https://metazoa.ensembl.org/Culex_quinquefasciatus/Info/Index). We

used STAR (Dobin et al. 2012) to map the short reads to the reference genome of *Cx.*

quinquefasciatus and RSEM (Li and Dewey 2011) to quantify gene abundances.

Following steps were carried out in R (R Core Team 2021) using Bioconductor

packages (Huber et al. 2015). We carried out a Variance Stabilizing Transformation

(VST) of the counts to represent the samples on a PCA plot. Then, we used the DESeq2

package (Love et al. 2014) to perform the differential gene expression analysis

comparing: i) *P. cathemerium* infected mosquitoes vs controls, ii) *P. relictum* infected

mosquitoes vs controls, and iii) *P. relictum* infected mosquitoes vs *P. cathemerium*

infected mosquitoes. We kept those genes with an adjusted p-value < 0.01 and sorted

them by the log2 fold changeestimations, considering genes with a LFC>1 or LFC<-1,

excepting one gene specified above. We finished the differential gene expression

analysis visualizing differentially expressed genes by MA plots and an UpSet plot.

Finally, we performed a Gene Ontology (GO) enrichment analysis for the up-

and down-regulated genes using the topGO package (Alexa and Rahnenfuhrer 2010)

including the "biological processes" and "molecular functions" categories from

VectorBase (Giraldo-Calderón et al. 2015). For the Enrichment analysis we used

classical algorithm and Fisher's exact test and considered enriched the GO terms with p-

value < 0,01. We used the ggplot2 (Wickham 2016) package to visualize the enriched

GO terms as described by Bonnot et al. (2019).

3. Results

Transcriptomics data description

We obtained 36 mRNA-seq libraries: four replicates for three infection status at

three time points. The mean number of reads per library was 33,177,082, ranging from

15,406,828 to 45,332,626 reads for raw samples. Raw data has been made publicly

available through to the European Nucleotide Archive ENA database

(https://www.ebi.ac.uk/ena/browser/home) under project accession number

PRJEB1609, Study ERP125411. The percentage of reads kept after trimming ranged

from 80.56% to 99.52%, (12,411,723 to 44,887,231 reads see Table S1). The MultiQC

report showed a mean quality score above q30 in all base calls across the read. On

average, 80.87% of reads mapped to the genome of *Cx. quinquefasciatus*. For

downstream analyses, we removed two samples taken at 24 hpi that had barcoding

errors due to lab processing.

Sample clustering reveals early transcriptomic response to infection

The principal component analysis (PCA) results show that most of the transcriptome

variation is contained in the PC1 (85% var.) driven by the time post-bloodmeal, under

both *Plasmodium* species infections. At 24 hpi, there were clear differences (PC2, 3%

var.) between control samples, *P. cathemerium*, and *P. relictum* infected mosquitoes. By

contrast, there were no clear transcriptomic differences at later stages, such as between

10 dpi and at 21 dpi, nor between infection statuses (Figure 1).

Figure 1. Principal component analysis (PCA) of transcriptome variation *in Cx. pipiens*
at three time points after feeding on either an uninfected bird (control), a *P.*
cathemerium-infected bird and a *P. relictum*-infected bird. Time points analysed were
24 hours post-infection (hpi), 10 days post-infection (dpi) and 21 dpi. The x-axis shows
the first principal component score, which captures 84% of variation and the y-axis
shows the second principal component score, which captures 3% of variation.

Differential gene expression analysis and enrichment analysis unveil the clues of the

mosquitoe transcriptomic response to Plasmodium infections

Overall, 2,051 genes were differentially expressed in *Cx. pipiens,* from which

590 showed a log2 fold change (LFC) higher than 1 or lower than -1 (see Table 1 for

differentially expressed genes distribution among comparisons). Most of the

transcriptomic differences were found at an early infection stage (24 hpi), especially

when comparing *P. relictum*-infected mosquitoes vs *P. cathemerium*-infected

mosquitoes, and *P. relictum*-infected mosquitoes vs controls. No differences were found

between mosquitoes infected by *P. cathemerium* and those infected by *P. relictum* at 10

dpi and between mosquitoes infected with *P. relictum* and controls at 21 dpi. At 24 hpi,

the comparisons *P. relictum*-infected mosquitoes vs controls and *P. relictum* vs *P.*

cathemerium-infected mosquitoes shared 460 differentially expressed genes (Figure 2;

see Supplementary Material for MA plots).

Table 1. Number of differentially expressed genes (DEG) with a p<0.01 for all
comparisons at 24 hours post-infection (24 hpi), 10 days post-infection (10 dpi), and 21
days post-infection (21 dpi). In brackets, DEG filtered by log2 fold change (LFC) > 1
for up- regulated genes and LFC < -1 for down-regulated genes. No DEG were found
for the comparisons *P. relictum* vs *P. cathemerium* infected mosquitoes at 10 dpi and *P.*
cathemerium infected mosquitoes vs controls at 21 dpi.

Time-point	Comparisons	Total DGE	Up-regulated	Down-regulated
24 hpi	*P. relictum* vs controls	645 (115)	107 (16)	538 (99)
	P. cathemerium vs controls	83 (18)	79 (19)	4 (2)
	P. relictum vs *P. cathemerium*	1256 (406)	330 (49)	926 (357)
10 dpi	*P. relictum* vs controls	62 (26)	46 (24)	16 (2)
	P. cathemerium vs controls	1 (0)	1 (0)	0 (0)
21 dpi	*P. relictum* vs controls	2 (2)	1 (1)	1 (1)
	P. relictum vs *P. cathemerium*	2 (2)	1 (1)	1 (1)

268
269   **Figure 2.** UpSet plot showing overlap size of sets of differentially expressed genes for
270   i) *P. cathemerium* infected mosquitoes vs controls, ii) *P. relictum infected* mosquitoes
      vs controls, and iii) *P. relictum* infected mosquitoes vs *P. cathemerium* infected

272　mosquitoes at three time points (24 hpi, 10 dpi and 21 dpi). The top vertical bar plot

273　shows the number of genes (y-axis) contained in each intersection (x-axis). The

274　horizontal bar plot at the bottom shows the number of differentially expressed genes for

each comparison.

　　　　a) Cx. pipiens gene expression response to P. relictum infection

At 24 hpi, up-regulated genes in *P. relictum*-infected *Cx. pipiens* compared to

controls included a cecropin gene (CPIJ005108), which is part of the Toll pathway in

the immune response, and a mitochondrial NADH-ubiquinone oxidoreductase gene

(CPIJ009076). Another gene involved in the Toll pathaway, *spätzle 1B* (CPIJ006792),

was also up-regulated in *P.relictum*-infected mosquitoes, although with a LFC = 0.3984.

Within the down-regulated genes there were four multicopper oxidase genes

(CPIJ016802, CPIJ010466, CPIJ012244, and CPIJ010465), four chymotrypsins

(CPIJ003915, CPIJ018205, CPIJ007838, and CPIJ006568) and two serine proteases

genes (CPIJ004984 and CPIJ002112) (Figure S1A).

At 10 dpi, when compared to controls, mosquitoes infected by *P. relictum* two

up-regulated genes were ribosomal genes (CPIJ040837 and CPIJ009519). At 21 dpi, the

aromatic amino acid decarboxylase gene (CPIJ010562) was down-regulated (Figure

S1B).

At 24 hpi, the GO categories within biological processes related to cellular

mechanisms and mitochondrial chain complex assemblies presented the greatest number

of up-regulated genes. Within down-regulated genes, the enriched biological processes

included molecules metabolism, transport and localization (Figure 3). The molecular

functions enriched for up-regulated genes were cation binding, metal ion binding and

ion binding. Protein, ATP, and nucleic acids binding molecular functions (including

purine ribonucleoside triphosphate, purine nucleotide, purine ribonucleotide, adenyl

nucleotide and adenyl ribonucleotide, among others) were enriched for down-regulated

genes (Figure 4).

300
Figure 3. Dot plot of significantly (adjusted p-value <0.01) enriched GO biological
processes for differentially expressed genes (up and down-regulated) at 24 hpi for i) *P.*
cathemerium infected mosquitoes vs controls, ii) *P. relictum* infected mosquitoes vs
controls, and iii) *P. relictum* infected mosquitoes vs *P. cathemerium* infected
mosquitoes. We did not find enriched GO terms for down-regulated genes for *P.*
cathemerium infected mosquitoes vs controls. Larger dots correspond to a higher
number of significant genes, and the color gradient goes from green for the least
significant terms to red for the most significant terms, measured as False Discovery
        Rate (FDR).

Figure 4. Dot plot of significantly (adjusted p-value <0.01) enriched GO molecular functions for differentially expressed genes (up and down-regulated) at 24 hpi for i) *P. cathemerium* infected mosquitoes vs controls, ii) *P. relictum* infected mosquitoes vs controls, and iii) *P. relictum* infected mosquitoes vs *P. cathemerium* infected mosquitoes. We did not find enriched GO terms for down-regulated genes for *P. cathemerium* infected mosquitoes vs controls.

At 10 dpi enriched biological processes were related to amino acid biosynthesis for up-regulated genes and to protein geranylgeranylation for the down-regulated genes (Figure S2). Molecular functions included actin monomer binding and GTPase activity for up-regulated genes (Figure S3).

b) *Cx. pipiens* gene expression during response to *P. cathemerium* infection

In *P. cathemerium*-infected mosquitoes compared to controls at 24 hpi (Figure

S4), we found up-regulated genes related to digestive enzymes like *trypsin 2 precursor*

(CPIJ005273), two protein G12 precursors (CPIJ012846 and CPIJ012848), and genes

related to the mitochondrial electron chain including genes for an ADP-ATP carrier

protein (CPIJ005941), a mitochondrial phosphate carrier protein (CPIJ013141), and the

cytochrome P450 4C51 (CPIJ018944). The *pom1* gene (CPIJ019938) was down-

regulated. At both 10 dpi and 21dpi no genes had differential gene expression with

LFC > 1.

At 24 hpi, GO terms associated with biological processes were mostly related to

the generation of precursor metabolites and energy, ATP generation, and nucleic acids

metabolic processes, including nucleoside phosphate, ribose phosphate, nucleotide,

ribonucleotide, purine nucleotide and purine ribonucleotide metabolic processes (Figure

3). GO terms associated with molecular functions included ion and transmembrane

transporter, oxidoreductase and proton-transporting ATP synthase activities (Figure 4).

c) *Cx. pipiens* respond differently to *P. relictum* and *P. cathemerium* infections

At 24 hpi, the comparison *P. relictum* vs *P. cathemerium*- infected mosquitoes

showed the larger number of differentially expressed genes (Table 1; Figure S5). The

up-regulated genes included four zinc finger proteins genes (CPIJ015530, CPIJ012592,

CPIJ008782, and CPIJ009347). On the other hand, mosquitoes infected by *P. relictum*

compared to those infected by *P. cathemerium* showed down-regulated genes of five

protein G12 precursors (CPIJ005176, CPIJ012848, CPIJ012844, CPIJ012845, and

CPIJ012846), four multicopper oxidases (CPIJ016802, CPIJ010466, CPIJ010465, and

CPIJ000864), two chitin synthases (CPIJ014268 and CPIJ014269), a peritrophic

membrane chitin binding protein (CPIJ007042), and a number of serine proteases

including five chymotrypsin genes (CPIJ003915, CPIJ018205, CPIJ007838,

CPIJ006542, and CPIJ006568) and three trypsin precursors (CPIJ005273, CPIJ006019,

and CPIJ004660). From the down-regulated genes, three multicopper oxidases and three

chymotrypsin precursors were also down-regulated in *P. relictum*-infected mosquitoes

compared to controls and two G12 protein precursors and one trypsin precursor were

also up-regulated in *P. cathemerium*-infected mosquitoes.

No differentially expressed genes were found at 10 dpi and only 2 genes at 21

dpi, a down-regulated testicular acid phosphatase precursor (CPIJ007697) and an up-

regulated uncharacterized protein.

At 24 hpi, biological processes related to metabolic regulation were enriched for up-

regulated genes, while down-regulated genes included nucleic acid nucleotide metabolic

processes (Figure 3), some of them also enriched for up-regulated genes *in P.*

cathemerium-infected mosquitoes compared to controls. Molecular functions like

microtubule and nucleic acid binding were enriched in up-regulated genes in

mosquitoes infected by *P. relictum* compared to those infected by *P. cathemerium*. Most

down-regulated enriched molecular functions were related to protein and nucleotides

binding (Figure 4), as observed in mosquitoes infected by *P. relictum* compared to

controls.

4. Discussion

How mosquitoes respond to infection and the impact of such infection will

ultimately influence the transmission of different *Plasmodium* species. For example, *P.*

relictum and *P. cathemerium* infections in *Cx. pipiens* have a different impact on the

mosquitoes' fitness. In particular, *P. relictum* causes higher mortality than *P.*

cathemerium infections in *Cx. pipiens*, and the transmissibility of *P. relictum* is lower

than that of *P. cathemerium* (Gutiérrez-López et al. 2020). But little is known about the

genetic underpinnings of mosquito response to *Plasmodium* infections. Here, we

compare for the first time the transcriptional response of *Cx. pipiens* infected by these

two avian *Plasmodium* species. Our results show that although responses of infected *Cx.*

pipiens by these *Plasmodium* species share some common pathways, there are key

differences in gene expression associated with the immune system and serine-protease

synthesis. These information provides new insights into the molecular mechanism

underlying *Culex pipiens* immune response against *Plasmodium*. Furthermore, by

comparing two *Plasmodium* species we show that there is variation in the immune

response of *Culex pipiens* to different avian malaria species. This information will in the

long term help to understand the complex interaction between the mosquitos and

different malaria parasites and potentially uncover the key to understanding differences

on transmission and fitness consequences for the mosquitoes.

Early and different response to the infection

When a mosquito feeds on a vertebrate host, a number of processes involving

changes in gene expression are triggered (Dana et al. 2005). We found that the vast

majority of the differentially expressed genes between infected and uninfected

mosquitoes were found at 24 hpi. Between 18-24 h after blood feeding ookinetes form,

invade the peritrophic matrix leaving behind the blood bolus and start the midgut

epithelium cell invasion (Cirimotich et al. 2010; Valkiūnas et al. 2015; Baia-da-Silva et

al. 2018). Ookinete formation and invasion of the midgut epithelium are considered

critical steps that will determine the success of the infection. Parasite abundance drops

drastically during this step due to lumenal and epithelial immune responses mounted by

the mosquito (Cirimotich et al. 2010), which may explain the significant differences in

gene expression between uninfected and infected mosquitoes at 24 hpi (Ferreira et al.

2022). In addition, Vlachou et al. (2005) using the *Plasmodium berghei - Anopheles*

gambiae model found that 7% of the mosquito transcriptome was differentially

regulated during ookinete invasion of the midgut.

At 24 hpi, for both mosquitoes infected with *P. cathemerium* and *P. relictum*, we

found an increase in differentially expressed genes and enriched GO terms related with

the mitochondrial respiratory chain activity that ultimately produces reactive oxygen

species (ROS) (Kowaltowski et al. 2009). Small regulatory changes in the

mitochondrial respiratory chain can drastically affect ROS generation (Korshunov et al.

1997; Kowaltowski et al. 2009). For example, *Anopheles stephensi* and *An. gambiae*

increased the levels of ROS in response to *Plasmodium* infection (Han 2000; Kumar et

al. 2003), and higher levels of ROS improve mosquito survival after a bacterial

infection (Molina-Cruz et al. 2008). At 24 hpi we also found most of the differences in

gene expression between mosquitoes infected by each of these parasites.

Differential immune response between mosquitoes exposed to *P. relictum* or *P.*

cathemerium

We found important differences in the expression of genes associated with the

immune response between *P. relictum-* and *P. cathemerium*-infected mosquitoes. At 24

hpi, in *Cx. pipiens* infected by *P. cathemerium* compared to controls, we found that two

of the most up-regulated genes were protein G12 precursors. These two genes and other

three protein G12 precursor genes were down-regulated in mosquitoes infected by *P.*

relictum compared to those infected by *P. cathemerium*. G12 transcripts accumulate in

the midgut of mosquitoes after blood feeding (Shao et al. 2005; Bonizzoni et al. 2012),

and may aid in erythrocyte digestion given their hemolytic activity (Foo et al. 2020).

G12 transcripts have been suggested to play a role in immune function because: i) G12

protein in Ae. *Aegypti* has a high level of identity with cockroach allergens (Morlais et

al. 2003), ii) it is up-regulated after flavivirus infection via the JAK-STAT pathway

(Etebari et al. 2017), and iii) it has a cytolytic effect on flaviviruses and several types of

eukaryotic cells (Foo et al. 2020). In fact, a G12 protein gene was found up-regulated in

Ae. aegypti 12 hours after feeding on blood infected with *Plasmodium gallinaceum*

(Morlais et al. 2003), suggesting a potential role in the immune response against these

parasites as well.

In *P. relictum*-infected mosquitoes two of the up-regulated genes at 24 hpi were

related to the Toll pathway of the innate immune response when compared to the

controls: the Spätzle and Cecropin N proteins. In insects, when a pathogen is

recognized, the extracellular Spätzle cytokine activates the Toll receptors, which

regulate the antimicrobial peptides (AMPs), an essential innate immune response (De

Gregorio 2002; Shia et al. 2009). The AMPs eventually kill pathogens by a number of

strategies including disrupting the microbial membrane (Shen et al. 2018). Spätzle

protein activates a Toll receptor in *Ae. aegypti* mosquitoes when infected with the

fungus *Beauveria bassiana* (Shin et al. 2006) and in other insects after bacterial and

fungal exposure (Bae et al. 2021). Cecropins are one of the largest groups of insect

AMPs found in the orders Diptera, Lepidoptera, and Coleoptera among others (Vizioli

et al. 2000; An et al. 2009; Memarpoor-Yazdi et al. 2013). The activation of the Toll

pathway and specifically cecropin-analogs may kill *Plasmodium* parasites (Jaynes et al.

1988; Frolet et al. 2006) and disrupt sporogonic development by aborting the normal

development of oocysts (Gwadz et al. 1989; Kim et al. 2004), which could reduce the

parasite load in posterior infection stages. Our results showing the role of the Toll

pathway are consistent with a recent study addressing the effect of *P. relictum* (lineage

SGS1) on the immune system of *Cx. quinquefasciatus* (García-Longoria et al. 2022).

They found that over 50% of immune genes identified as being part of the Toll pathway

in *Cx. quinquefasciatus* were up-regulated after exposure to *P. relictum.*

Differential expression of genes associated with trypsin and serine metabolism

Serine proteases have several functions in insects including blood digestion

(Borovsky and Schlein, 1988) and mediation in the immune response (melanization,

cytokine activation, and antimicrobial peptides; Jiang et al. 2010). Trypsins and

chymotrypsins, two types of serine proteases, are two essential digestive enzymes in

mosquitoes (Molina-Cruz et al. 2005; Borges-Veloso et al. 2012). The synthesis of

trypsin and chymotrypsin is triggered after mosquito blood feeding to digest the chitin-

containing peritrophic matrix (Vizioli et al. 2001; Muller et al. 1995) which is fully

formed at 24 h post feeding (Hegedus et al. 2009). However, the production of trypsin

and chymotrypsin may be down or up-regulated by the infection with different parasites

(Borovsky and Schlein 1987; Shahabuddin et al. 1996; Serrano-Pinto et al. 2010).

Mosquito trypsin may be a signal for *Plasmodium* ookinetes to cross the peritrophic

matrix at the right time for proper *Plasmodium* development (Shahabuddin et al. 1996).

In particular, mosquito trypsin proteases play a fundamental role in allowing *P.*

gallinaceum to cross the peritrophic matrix by activating a *Plasmodium* prochitinase

enzyme (Shahabuddin et al. 1996). In fact, trypsin inhibitors block the development of

Plasmodium oocysts (Shahabuddin et al. 1996), suggesting that mosquito trypsin is a

key molecule for pathogen infection.

When compared to controls, mosquitoes exposed to *P. cathemerium* up-

regulated the trypsin 2 precursor gene, while mosquitoes exposed to *P. relictum*

strongly down-regulated genes ofchymotrypsins and serine protease precursors .

Furthermore, when comparing gene expression between mosquitoes exposed to each

species, we found that mosquitoes exposed to *P. relictum* also down-regulated the

chymotrypsins and serine protease precursors that when compared to control, as well as

others chymotrypsins, serine protease precursors, and three trypsin precursos. This

differential expression was found at 24 hpi, therefore coinciding with the moment when

the peritrophic matrix is fully formed and the ookinetes are crossing it. Agreeing with

the findings of Shahabuddin et al. (1996) for the infection by *P. gallinaceum*, we also

found that trypsin (*trypsin 2 precursor*) was up-regulated in mosquitoes infected with *P.*

cathemerium when compared with uninfected mosquitoes. This suggests the potential

role of the trypsin proteases to allow *P. cathemerium* to cross the peritrophic matrix. In

contrast, mosquitoes infected with *P. relictum* had a number of down-regulated genes

characterized as serine proteases, including several chymotrypsin and one trypsin.

Ferreira et al. (2022) also found serine-type endopeptidase activity enriched for down-

regulated genes when studying the expression of *Cx. quinquefasciatus* infected by *P.*

relictum. We hypothesize that the lower levels of trypsin in response to *P. relictum*

infection might lead to a lower number of oocysts and would explain why the

transmissibility of *P. relictum* is lower than that of *P. cathemerium* (Gutiérrez-López et

al. 2020).

In addition, an under-expression of serine-proteases related to blood digestion

such as chymotrypsins should partly block the digestion of the blood, which is very rich

in proteins (Kumar et al. 2017). This would make it difficult to obtain nutrients needed

for various processes like egg formation and could lead to digestive dysregulation in the

mosquito which might increase mortality. Here mosquitoes seem to be down-regulating

important processes due to the infection with *P. relictum* which might have a cost for

the mosquito, induce damage and decrease the tolerance to infection.

Reduced differential gene expression at 10 and 21 days post-infection

An unexpected result was the strong decrease of differential gene expression

between infected and uninfected mosquitoes at 10 dpi. We chose this time point because

at about 10 dpi *Plasmodium* mature oocysts release the sporozoites into the mosquito

hemocoel (Aly et al. 2020). However, at this time point, only *P. relictum*-infected

mosquitoes compared to controls showed differences in gene expression and the number

of differentially expressed genes decreased significantly compared to 24 hpi. Similar

results were obtained by Ferreira et al. (2022) when studying gene expression of *Cx.*

quinqueafasciatus exposed to *P. relictum*. In addition, the absence of differential

expression in mosquitoes exposed to *P. relictum* compared to those exposed to *P.*

cathemerium may be due to differences in the developmental time of different species

of *Plasmodium* (Sinden 1983). *P. cathemerium* produces sporozoites faster than *P.*

relictum when infecting *Cx. pipiens* mosquitoes (Kazlauskienė et al. 2013; Aly et al.

2020), and therefore they may not be exactly at the same point of development within

the mosquito at 10 dpi. In addition, when the sporozoites are realesed, part of them are

destroyed by a rapid immune response (Hillyer et al. 2007), which may also result in a

lower genetic response by the mosquito.

By 21 dpi, the differences in gene expression were drastically reduced. With only

two genes differentially expressed in mosquitoes infected with *P. relictum*, and none in

those infected with *P. cathemerium*. This decrease in differential expression towards the

end of the infection has been found before in *Cx. quinqueafasciatus* infected with *P.*

relictum (Ferreira et al. 2022, García-Longoria et al. 2022). Although the causes behind

this decrease are not clear, we hypothesize that the response towards the end of the

infection might be localized to the salivary glands, and by analysing whole mosquitoes

we are diluting any potential effect.

Comparisons across avian malaria studies

Although studies analysing the response of mosquitoes to avian malaria are still

scarce we can already see some common patterns and differences. Two other recent

studies have analysed the response to *P. relictum* infection in *Cx. quinquefasciatus*, a

species closely related to *Cx. pipiens*. A common pattern found in all studies is the

progressive reduction in the differential gene expression at 10 and 21 dpi with respect to

24 hpi. In addition, like García-Longoria et al. (2022), we found the activation of the

Toll-like receptor pathway in response to *P. relictum SGS1* infection. Interestingly,

Ferreira et al. (2022) did not find this pattern. Although different factors may be

influencing this result, like experimental procedures or the genetic differences between

the populations of *Cx. quinquefasciatus* used, one of the main factors might have been

the lineage used for the infection. Ferreira et al. (2022) infected the mosquitoes with

another lineage, *P. relictum* GRW4, which dominates in America. Like Ferreira et al.

(2022), the response to infection we found for mosquitoes infected with *P.*

cathemerium, was different. But, a similar pattern has been found before. Shahabuddin

et al. (1996) reported the important role of a mosquito trypsin in the passage of *P.*

gallinaceum through the peritrophic matrix, as our results indicate for *P. cathemerium*.

Altogether these result suggests that different *Plasmodium* species or lineages may

trigger a differential immune response in mosquitoes.

Concluding remarks

Because we used naturally infected wild birds as *P. cathemerium* and *P. relictum*

donors and wild collected mosquitoes, the results obtained here represent a good

example of a natural system. In this respect, we found a different transcriptomic

response to infections, especially at 24 hpi. This time point coincides with one of the

key stages of *Plasmodium* development in mosquitos, when the ookinetes form, cross

the peritrophic matrix and start to invade the midgut epithelium. Both elicit an innate

immune response, with mosquitoes exposed to *P. relictum* activating the Toll Pathway

(Spätzle and Cecropin N proteins) which may result in a a reduction of *Plasmodium*

development (Frolet et al. 2006) and *P. cathemerium*- infected mosquitoes up-

regulating the antimicrobial G12 proteins. On the other hand, the lower levels of trypsin

in mosquitoes exposed to *P. relictum* compared to those expose to *P. cathemerium* may

affect the parasite development within the mosquito, which may affect parasite

transmission (Gutiérrez-López et al. 2020). If the cost of this response is high, this can

also potentially lead to higher mosquito mortality. In particular, the proteases and

trypsins are necessary to digest the blood meal, and if levels are too low this might

increase mosquito mortality. Future studies are necessary to better understand how the

observe differences are linked to differences in transmission. In addition, it will also be

important to understand how these differences may be related to the different ecology

and incidence of *Plasmodium* lineages/species in the wild.

5. Acknowledgments

We thank Francisco Ferreira for his help optimizing the RNA extraction protocol,

Juan Pascual, and Alazne Diez for their help during the experiments, and Cristina Pérez

for her help with the laboratory work. This publication was supported by the project

Research Infrastructures for the control of vector-borne diseases (Infravec2), which has

received funding from the European Union's Horizon 2020 research and innovation

program, under grant agreement No 6738; project PGC2018-095704-B-I00 from

Agencia Española de Investigación supported by FEDER Funds from the European

Union and the computing infrastructure provided by ICTS-RBD-CSIC. This study was

also partially financed by the PID2020-118205GB-I00 grant funded by MCIN/AEI/

10.13039/501100011033 and by "ERDF A way of making Europe". MG was supported

by a FPI grant (PRE2021-098544). GY contributions were supported by the Faculty of

Biochemistry, Biophysics and Biotechnology at Jagiellonian University (Poland), under

the Strategic Programme Excellence Initiative.

6. References
574

- Abraham E, Jacobs-Lorena M. 2004. Mosquito midgut barriers to malaria parasite

development. *Insect Biochem. Mol Biol* **34**(7): 667-671.

- Ahmed AM, Baggott SL, Maingon R, Hurd H. 2002. The costs of mounting an

immune response are reflected in the reproductive fitness of the mosquito *Anopheles*

gambiae. *Oikos (Copenhagen, Denmark)*, **97**(3): 371-377.

https://doi.org/10.1034/j.1600-0706.2002.970307.x

- Alexa A, Rahnenfuhrer J. 2010. topGO: enrichment analysis for gene ontology. R

package version, **2**(0), 2010.

- Aly MZY, Mohamed III, Sebak SI, Vanstreels RET, El Gendy AM. 2020.

Morphological and molecular characterization of *Plasmodium cathemerium* (lineage

PADOM02) from the sparrow *Passer domesticus* with complete sporogony in *Culex*

pipiens complex. *Parasitology* **147**(9): 985-993.

https://doi.org/10.1017/S0031182020000566

- An C, Jiang H, Kanost M. 2009. Proteolytic activation and function of the cytokine

Spätzle in the innate immune response of a lepidopteran insect, *Manduca sexta*. *FEBS*

J **277**(1): 148-162.

- Andrews S. 2010. FastQC: A Quality Control Tool for High Throughput Sequence

Data [Online].

- Baia-da-Silva DC, Alvarez LCS, Lizcano OV, Costa FTM, Lopes SCP, et al. 2018.

The role of the peritrophic matrix and red blood cell concentration in *Plasmodium vivax*

infection of *Anopheles aquasalis*. *Parasit Vectors* **11**(1).

https://doi.org/10.1186/s13071-018-2752-5

- Bae Y, Jo Y, Patnaik B, Kim B, Park K, Edosa T, et al. 2021. *Tenebrio molitor* Spätzle

1b Is Required to Confer Antibacterial Defense Against Gram-Negative Bacteria by

Regulation of Antimicrobial Peptides. *Front Physiol* **12**. doi:

10.3389/fphys.2021.758859

- Beerntsen B, James A, Christensen B. 2000. Genetics of Mosquito Vector

Competence. *Microbiol. Mol Biol Rev* **64**(1): 115-137.

- Bensch S, Hellgren O, Pérez-Tris J. 2009. MalAvi: a public database of malaria

parasites and related haemosporidians in avian hosts based on mitochondrial

cytochrome b lineages. *Mol Ecol Resour* **9**(5): 1353-1358.

- Bolger A, Lohse M, Usadel B. (2014). Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**(15): 2114-2120.

- Bonizzoni M, Gasperi G, Chen X, James A. 2013. The invasive mosquito species

Aedes albopictus: current knowledge and future perspectives. *Trends Parasitol* **29**(9):

460-468.

- Bonizzoni M, Dunn W, Campbell C, Olson K, Marinotti O, et al. 2012. Strain

variation in the transcriptome of the dengue fever vector, *Aedes aegypti. G3 (Bethesda,*

Md) **2**(1): 103-114. https://doi.org/10.1534/g3.111.001107

- Bonnot T, Gillard M, Nagel D. 2019. A Simple Protocol for Informative Visualization

of Enriched Gene Ontology Terms. *Bio Protoc* **9**(22). doi: 10.21769/bioprotoc.3429

- Borges-Veloso A, Saboia-Vahia L, Cuervo P, Pires R, Britto C, et al. 2012. Proteolytic

profiling and comparative analyses of active trypsin-like serine peptidases in

preimaginal stages of *Culex quinquefasciatus*. *Parasite Vector* **5**(1). doi: 10.1186/1756-

3305-5-123

- Borovsky D, Schlein Y. 1988. Quantitative determination of trypsinlike and

chymotrypsinlike enzymes in insects. *Arch Insect Biochem Physiol* **8**(4): 249-260.

- Borovsky D, Schlein Y. 1987. Trypsin and chymotrypsin-Iike enzymes of the sandfly

Phlebotomus papatasi infected with *Leishmania* and their possible role in vector

competence. *Med Vet Entomol* **1**(3): 235-242.

- Cirimotich C, Dong Y, Garver L, Sim S, Dimopoulos G. 2010. Mosquito immune

defenses against *Plasmodium* infection. *Dev Comp Immunol* **34**(4): 387-395.

- Clayton AM, Dong Y, Dimopoulos G. 2014. The Anopheles innate immune system in

the defense against malaria infection. J Innate Immun **6**: 169-181.

- Dana AN, Hong Y, Kern MK, Hillenmeyer ME, Harker B, et al. 2005. Gene

expression patterns associated with blood-feeding in the malaria mosquito *Anopheles*

gambiae. *BMC Genet* **6**(1). doi: 10.1186/1471-2164-6-5

- De Gregorio E. 2002. The Toll and Imd pathways are the major regulators of the

immune response in *Drosophila*. *EMBO J* **21**(11): 2568-2579.

-Detinova TS, Bertram DS, World Health Organization. 1962. Age-grouping methods in

diptera of medical importance, with special reference to some vectors of malaria / T. S.

Detinova ; [with] an Annex on the ovary and ovarioles of mosquitos (with glossary) by

D. S. Bertram. World Health Organization.

- Dobin A, Davis C, Schlesinger F, Drenkow J, Zaleski C, et al. 2012. STAR: ultrafast

universal RNA-seq aligner. *Bioinformatics* **29**(1): 15-21.

- Etebari K, Hegde S, Saldaña M, Widen S, Wood T, et al. 2017. Global transcriptome

analysis of *Aedes aegypti* mosquitoes in response to Zika virus infection. *MSphere* **2**(6).

https://doi.org/10.1128/mSphere.00456-17

- Ewels P, Magnusson M, Lundin S, Käller M. 2016. MultiQC: summarize analysis

results for multiple tools and samples in a single report. *Bioinformatics* **32**(19): 3047-

3048.

- Ferreira FC, Videvall E, Seidl CM, Wagner NE, Kilpatrick AM, et al. 2022.

Transcriptional response of individual Hawaiian *Culex quinquefasciatus* mosquitoes to

the avian malaria *parasite Plasmodium relictum*. *Malar J* **21**(1): 249.

https://doi.org/10.1186/s12936-022-04271-x

- Foo A., Thompson P, Chen S-H, Jadi R, Lupo B, et al. 2021. The mosquito protein

AEG12 displays both cytolytic and antiviral properties via a common lipid transfer

mechanism. *Proc Natl Acad Sci USA* **118**(11): e2019251118

- Frolet C, Thoma M, Blandin S, Hoffmann JA, Levashina EA. 2006. Boosting NF-

kappaB-dependent basal immunity of *Anopheles gambiae* aborts development of

Plasmodium berghei. Immunity **25**(4): 677685.

https://doi.org/10.1016/j.immuni.2006.08.019

- García-Longoria L, Palinauskas V, Ilgūnas M, Valkiūnas G, Hellgren O. 2020.

Differential gene expression of *Plasmodium homocircumflexum* (lineage pCOLL4)

across two experimentally infected passerine bird species. *Genomics* **112**(4): 2857-

2865.

- García-Longoria L, Ahrén D, Berthomieu A, Kalbskopf V, Rivero A, et al. 2022.

Immune gene expression in the mosquito vector *Culex quinquefasciatus* during an avian

malaria infection. *Mol Ecol* https://doi.org/10.1111/mec.16799

- Gemmell N, Akiyama S. 1996. An efficient method for the extraction of DNA from

vertebrate tissues. *Trends Genet* **12**(9): 338-339.

- Giraldo-Calderón G, Emrich S, MacCallum R, Maslen G, Dialynas E, et al. 2015.

VectorBase: an updated bioinformatics resource for invertebrate vectors and other

organisms related with human diseases. *Nucleic Acids Res* **43**(D1): D707-D713.

- Gunay F, Picard M, Robert V. 2018. MosKeyTool, an interactive identification key for

mosquitoes of Euro-Mediterranean. Version 2.1. 2018. Available from:

http://www.medilabsecure.com/moskeytool.

- Gutiérrez-López R, Martínez-de la Puente J, Gangoso L, Soriguer R, Figuerola J.

2020. *Plasmodium* transmission differs between mosquito species and parasite lineages.

Parasitology **147**(4): 441-447.

- Gwadz R, Kaslow D, Lee J, Maloy W, Zasloff M, et al. 1989. Effects of magainins

and cecropins on the sporogonic development of malaria parasites in mosquitoes. *Infect*

Immun **57**(9): 2628-2633.

- Han YS. 2000. Molecular interactions between Anopheles stephensi midgut cells and

Plasmodium berghei: the time bomb theory of ookinete invasion of mosquitoes. *EMBO*

J **19**(22): 6030-6040.

- Hegedus D, Erlandson M, Gillott C, Toprak U. 2009. New Insights into Peritrophic

Matrix Synthesis, Architecture, and Function. *Annu Rev Entomol* **54**(1): 285-302.

- Hellgren O, Waldenström J, Bensch S. 2004. A new PCR assay for simultaneous

studies of leucocytozoon, *Plasmodium*, and *Haemoproteus* from avian blood. *J*

Parasitol **90**(4): 797-802.

- Higgs S, Beaty BJ. 2005. Natural cycles of vector-borne pathogens. Biology of disease

vectors. In Biology of disease vectors, 2nd ed. (ed. Marquardt WC, *et al*.). Elsevier

Academic Press, New York.

- Hillyer J. 2016. Insect immunology and hematopoiesis. *Dev Comp Immunol* **58**: 102-

118.

- Huber W, Carey V, Gentleman R, Anders S, Carlson M, et al. 2015. Orchestrating

high-throughput genomic analysis with Bioconductor. *Nat Methods* **12**(2): 115-121.

- Janeh M, Osman D, Kambris Z. 2019. Comparative Analysis of Midgut Regeneration

Capacity and Resistance to Oral Infection in Three Disease-Vector Mosquitoes. *Sci Rep*

9(1). doi: 10.1038/s41598-019-50994-4

- Jaynes J, Burton C, Barr S, Jeffers G, Julian G, et al. 1988. In vitro cytocidal effect of

novel lytic peptides on *Plasmodium* falciparum and *Trypanosoma cruzi. FASEB J*

2(13): 2878-2883.

- Jiang H, Vilcinskas A, Kanost MR. 2010. Immunity in lepidopteran insects. *Adv Exp*

Med Biol **708**:181-204. https://doi.org/10.1007/978-1-4419-8059-5_10

- Kazlauskienė R, Bernotienė R, Palinauskas V, Iezhova TA, Valkiūnas G. 2013.

Plasmodium relictum (lineages pSGS1 and pGRW11): complete synchronous

sporogony in mosquitoes *Culex pipiens pipiens. Exp Parasitol* **133**(4): 454-461.

https://doi.org/10.1016/j.exppara.2013.01.008

- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, et al. 2012. Geneious

Basic: An integrated and extendable desktop software platform for the organization and

analysis of sequence data. *Bioinformatics* **28**(12): 1647-1649.

- Kim W, Koo H, Richman A, Seeley D, Vizioli J, et al. 2004. Ectopic Expression of a

Cecropin Transgene in the Human Malaria Vector Mosquito *Anopheles gambiae*

(Diptera: Culicidae): Effects on Susceptibility to *Plasmodium. J Med Entomol* **41**(3):

447-455.

- Korshunov SS, Skulachev VP, Starkov AA. 1997. High protonic potential actuates a

mechanism of production of reactive oxygen species in mitochondria. *FEBS Lett* **416**:

15–18

- Kowaltowski A, de Souza-Pinto N, Castilho, Vercesi A. 2009. Mitochondria and

reactive oxygen species. *Free Radic Biol Med* **47**(4): 333-343.

- Kumar A, Srivastava P, Sirisena P, Dubey S, Kumar R, et al. 2018. Mosquito Innate

Immunity. *Insects* **9**(3): 95.

- Kumar M, Mohanty A, Sreenivasamurthy S, Dey G, Advani J, et al. 2017. Response to

Blood Meal in the Fat Body of *Anopheles stephensi* Using Quantitative Proteomics:

Toward New Vector Control Strategies Against Malaria. *OMICS J Integr Biol* **21**(9):

520-530.

- Kumar S, Christophides GK, Cantera R, Charles B, Han YS, et al. 2003. The role of

reactive oxygen species on *Plasmodium* melanotic encapsulation in Anopheles gambiae.

Biol Sci **100**(24): 14139-14144.

- Lehane MJ. 2010. *The biology of blood-sucking in insects,* 2nd ed. Cambridge

University Press, Cambridge.

- Lehane M. 1997. Peritrophic matrix structure and function. *Annu Rev Entomol* **42**(1):

525-550.

- Li B, Dewey C. 2011. RSEM: accurate transcript quantification from RNA-Seq data

with or without a reference genome. *BMC Bioinform* **12**(1). doi: 10.1186/1471 2105-12-

323

- Love M, Huber W, Anders S. 2014. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**(12). doi: 10.1186/s13059-

014- 0550-8

- Memarpoor-Yazdi M, Zare-Zardini H, Asoodeh A. 2013. A Novel Antimicrobial

Peptide Derived from the Insect *Paederus dermatitis*. *Int J Pept Res Ther* **19**(2): 99-108.

- Michel K, Kafatos F. 2005. Mosquito immunity against *Plasmodium*. *Insect Biochem*

Mol Biol **35**(7): 677-689.

- Molina-Cruz A, Bennett K, Barillas-Mury C, Richardson J, Black W, et al. 2005.

Effect of mosquito midgut trypsin activity on dengue-2 virus infection and

dissemination in *Aedes aegypti*. *Am J Trop Med* Hyg **72**(5): 631-637.

- Molina-Cruz A, DeJong RJ, Charles B, Gupta L, Kumar S, et al. 2008 Reactive

oxygen species modulate Anopheles gambiae immunity against bacteria and

Plasmodium. *J Biol Chem* **283**: 3217–3223.

- Morlais I, Mori A, Schneider J, Severson D. 2003. A targeted approach to the

identification of candidate genes determining susceptibility to *Plasmodium gallinaceum*

in *Aedes aegypti*. *Mol Genet Genomics* **269**(6): 753-764.

- Muller H, Catteruccia F, Vizioli J, Dellatorre A, Crisanti A. 1995. Constitutive and

Blood Meal-Induced Trypsin Genes in *Anopheles gambiae*. *Exp Parasitol* **81**(3): 371-

385.

- Nászai M, Carroll L, Cordero J. 2015. Intestinal stem cell proliferation and epithelial

homeostasis in the adult *Drosophila* midgut. *Insect Biochem Mol Biol* **67**: 9-14.

- R Core Team. 2021. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-

project.org/.

- Serrano-Pinto V, Acosta-Pérez M, Luviano-Bazán D, Hurtado-Sil G, Batista CVF, et

al. 2010. Differential expression of proteins in the midgut of *Anopheles albimanus*

infected with *Plasmodium berghei*. *Insect Biochem. Mol Biol* **40**(10): 752-758.

https://doi.org/10.1016/j.ibmb.2010.07.011

- Shahabuddin M, Lemos F, Kaslow D, Jacobs-Lorena M. 1996. Antibody mediated

inhibition of *Aedes aegypti* midgut trypsins blocks sporogonic development of

Plasmodium gallinaceum. *Infect Immun* **64**(3): 739-743.

- Shao L. Devenport M, Fujioka H, Ghosh A, Jacobs-Lorena M. 2005. Identification

and characterization of a novel peritrophic matrix protein, Ae-Aper50, and the

microvillar membrane protein, AEG12, from the mosquito, *Aedes aegypti*. *Insect*

Biochem Mol Biol **35**(9): 947-959.

- Shen W, He P, Xiao C, Chen X. 2018. From Antimicrobial Peptides to Antimicrobial

Poly(α-amino acid)s. *Adv Healthc Mater* **7**(20): 1800354. doi:

10.1002/adhm.201800354

- Shia A, Glittenberg M, Thompson G, Weber A, Reichhart J, et al. 2009. Toll-

dependent antimicrobial responses in *Drosophila* larval fat body require Spa□tzle

secreted by haemocytes. *J Cell Sci* **122**(24): 4505-4515. doi: 10.1242/jcs.049155

- Shin S, Bian G, Raikhel A. 2006. A Toll Receptor and a Cytokine, Toll5A and Spz1C,

Are Involved in Toll Antifungal Immune Signaling in the Mosquito *Aedes aegypti. J*

Biol Chem **281**(51): 39388-39395.

- Sinden R. 2002. Molecular interactions between *Plasmodium* and its insect vectors.

Cell Microbiol **4**(11): 713-724.

- Sinden R. 1983. Sexual Development of Malarial Parasites. *Adv Parasitol* **22**: 153-

216.

- Taracena M, Bottino-Rojas V, Talyuli O, Walter-Nuno A, Oliveira J, et al. 2018.

Regulation of midgut cell proliferation impacts *Aedes aegypti* susceptibility to dengue

virus. *PLoS Negl Trop. Dis*. **12**(5): e0006498. doi: 10.1371/journal.pntd.0006498

- Valkiūnas G. 2005. Avian Malaria Parasites and Other Haemosporidia. CRC Press,

New York.

-Valkiūnas G, Iezhova TA. 2018. Keys to the avian malaria parasites. *Malar J* **17**(1).

https://doi.org/10.1186/s12936-018-2359-5

- Valkiūnas G, Žiegytė R, Palinauskas V, Bernotienė R, Bukauskaitė D, et al. 2015.

Complete sporogony of *Plasmodium relictum* (lineage pGRW4) in mosquitoes *Culex*

pipiens pipiens, with implications on avian malaria epidemiology. *Parasitol Res* **114**(8):

3075-3085. https://doi.org/10.1007/s00436-015-4510-3

- van Riper C, van Riper S, Goff M, Laird M. 1986. The Epizootiology and Ecological

Significance of Malaria in Hawaiian Land Birds. *Ecol Monogr* **56**(4): 327-344.

- Vizioli J, Catteruccia F, della Torre A, Reckmann I, Müller H. 2001. Blood digestion

in the malaria mosquito *Anopheles gambiae*. *Eur J Biochem* **268**(14): 4027-4035.

- Vizioli J, Bulet P, Charlet M, Lowenberger C, Blass C, et al. 2000. Cloning and

analysis of a cecropin gene from the malaria vector mosquito, *Anopheles gambiae*.

Insect Mol Biol **9**(1): 75-84.

- Videvall E, Cornwallis CK, Palinauskas V, Valkiūnas G, Hellgren O. 2015. The avian

transcriptome response to malaria infection. *Mol Biol Evol* **32**(5): 1255-1267.

https://doi.org/10.1093/molbev/msv016

- Videvall E, Paxton K, Campana M, Cassin☐Sackett L, Atkinson C, et al. 2021.

Transcriptome assembly and differential gene expression of the invasive avian malaria

parasite *Plasmodium relictum* in Hawai☐i. *Ecol Evol* **11**(9): 4935-4944.

- Videvall E, Cornwallis C, Ahrén D, Palinauskas V, Valkiūnas G, et al. 2017. The

transcriptome of the avian malaria parasite *Plasmodium ashfordi* displays host-specific

gene expression. *Mol Ecol* **26**(11): 2939-2958.

- Vlachou D, Schlegelmilch T, Christophides G, Kafatos F. 2005. Functional Genomic

Analysis of Midgut Epithelial Responses in *Anopheles* during *Plasmodium* Invasion.

Curr Biol **15**(13): 1185-1195.

- Wickham H. 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag,

New York.

- WHO. 2021. World malaria report 2021: regional data and trends. In World Health:

Vol. WHO/UCN/GMP/2021.09 (Issue December).

https://www.who.int/publications/m/item/WHO-UCN-GMP-2021.09 (Accessed May

18, 2022)

- Zou Z, Souza-Neto J, Xi Z, Kokoza V, Shin S, Dimopoulos G, et al. 2011.

Transcriptome Analysis of *Aedes aegypti* Transgenic Mosquitoes with Altered

Immunity. *Plos Pathog* **7**(11): e1002394. doi: 10.1371/journal.ppat.1002394

7. Data Availability Statement

Raw sequences generated in this study have been submitted to the European Nucleotide

Archive ENA database (https://www.ebi.ac.uk/ena/browser/home) under project

accession number PRJEB1609, Study ERP125411.

8. Author Contributions

María José Ruiz-López, Josué Martínez-de la Puente and Jordi Figuerola Borras

designed the original study. María José Ruiz-López developed the experimental assay

and the laboratory work. Marta Garrigós did the bioinformatic work and interpreted the

results with María José Ruiz-López and Guillem Ylla. Marta Garrigós wrote the first

original draft of the manuscript and subsequent versions with considerable assistance

from the rest of the authors.