Yersinia Type III-Secreted Effectors Subvert Caspase-4-dependent Inflammasome Activation in

Human Cells

Jenna Zhang[a], Igor E. Brodsky[b,#], and Sunny Shin[a,#]

[a]Department of Microbiology, University of Pennsylvania Perelman School of Medicine,

Philadelphia, Pennsylvania 19104

[b]Department of Pathobiology, University of Pennsylvania School of Veterinary Medicine,

Philadelphia, Pennsylvania 19104

Running title: *Yersinia* suppresses human inflammasome activation

#Co-corresponding authors: Igor E. Brodsky ibrodsky@vet.upenn.edu and Sunny Shin

sunshin@pennmedicine.upenn.edu

Abstract word count: 191

Text word count: 4927

Abstract

Yersinia are gram-negative zoonotic bacteria that use a type three secretion system (T3SS) to

inject *Yersinia* outer proteins (Yops) into the host cytosol in order to subvert essential

components of innate immune signaling. However, *Yersinia* virulence can elicit activation of

immune complexes known as inflammasomes, which lead to inflammatory cell death and

cytokine release aimed at containing infection. *Yersinia* activation and evasion of

inflammasomes have been characterized in macrophages but remain poorly defined in intestinal

epithelial cells (IECs), the primary site of gastrointestinal *Yersinia* infection. In contrast to murine

macrophages, we find that in human IECs, *Yersinia pseudotuberculosis* T3SS effectors fully

suppress activation of the caspase-4 inflammasome, which senses cytosolic lipopolysaccharide

(LPS). The antiphagocytic Yops YopE and YopH, as well as YopK, were entirely responsible for

inhibiting inflammasome activation, in part by inhibiting *Yersinia* internalization into IECs.

Surprisingly, Yops E, H and K also suppressed inflammasome activation in human

macrophages, which, like human IECs, failed to undergo cell death in response to wild-type

Yersinia. These data suggest species-specific differences underlying inflammasome activation

in response to *Yersinia*, and provide insight into the mechanisms of *Yersinia*-mediated

inflammasome activation and suppression in human cells.

Importance *Yersinia* are responsible for significant disease burdens in humans, ranging from

recurrent disease outbreaks (yersiniosis) to pandemics (*Yersinia pestis* plague). Together with

rising antibiotic resistance rates, there is a critical need to better understand *Yersinia*

pathogenesis and host immune mechanisms, as this information will aid in developing improved

immunomodulatory therapeutics. Intestinal epithelial cells are a critical component of intestinal

immunity, yet their inflammasome responses are understudied compared to those of innate

immune cells. Furthermore, inflammasome responses in human models are less studied relative

to murine models of infection although key innate immune differences exist between mice and

humans. Here, we dissect human intestinal epithelial cell and macrophage inflammasome

responses to *Yersinia pseudotuberculosis.* Our findings provide fundamental insight into

species- and cell-type specific differences in inflammasome responses to *Yersinia.*

50
Introduction

Innate immune responses are a critical component of host defense and employ pattern

recognition receptors (PRRs) to detect pathogen-associated molecular patterns (PAMPs) (1, 2).

Specific cytosolic PRRs in the Nucleotide binding domain and Leucine-rich Repeat-containing

protein (NLR) family induce the formation and activation of multimeric immune complexes

known as inflammasomes in response to cytosolic PAMPs and distinct stimuli downstream of

pathogen virulence-associated activity,(3–5). The NLRP3 inflammasome is activated by a

variety of stimuli including potassium efflux downstream of bacterial-induced pore formation (5–

9), whereas the NAIP-NLRC4 inflammasome responds to bacterial flagellin and components of

bacterial type III secretion systems (T3SS) (10–18). Inflammasomes recruit and activate

caspase-1, which in turn processes members of the IL-1 family of cytokines and the pore-

forming protein GSDMD into their active forms (19–23). In addition to these canonical

inflammasome pathways, a non-canonical caspase-4 inflammasome in humans, and

orthologous caspase-11 inflammasome in mice, responds to cytosolic lipopolysaccharide (LPS)

during gram-negative bacterial infection to initiate GSDMD pore formation and cytokine release

(24–31). Consequently, inflammasome activation leads to release of active IL-1 family

cytokines, specifically IL-1β and IL18, and an inflammatory form of cell death known as

pyroptosis, that collectively amplify immune signaling and promote anti-bacterial defense (32).

In parallel, bacterial infection triggers other programmed cell death pathways, including

apoptosis and necroptosis, thereby providing multiple layers of defense against bacterial

pathogens (33, 34).

The enteric pathogenic Yersiniae, *Yersinia pseudotuberculosis* and *Y. enterocolitica,*

express a conserved T3SS that injects *Yersinia* outer proteins (Yops) into target cells (35).

T3SS-injected Yops manipulate host cellular pathways to promote infection, but also activate

effector-triggered cell death responses. For example, in murine macrophages, YopJ-mediated

suppression of NF-κB signaling induces apoptosis whereas YopE-mediated disruption of

cytoskeletal dynamics and hyper-translocation of T3SS components both induce inflammasome

activation and subsequent pyroptosis (36–45). Notably, *Yersinia* utilizes two other effectors,

YopM and YopK, to evade YopE-triggered pyrin and T3SS-triggered NLRP3 and caspase-11-

inflammasome activation respectively and promote infection (42, 43, 45, 46), highlighting

competing necessities between host signaling manipulation to facilitate colonization and evasion

of ensuing effector-triggered immunity (47).

Inflammasome responses to *Yersinia* have primarily been studied in murine

macrophages. However, inflammasomes are expressed in multiple cell types, including

intestinal epithelial cells (IECs) (48), which are the first colonization barrier and the primary site

of infection of many gastrointestinal pathogens, including *Y. enterocolitica* and *Y.*

pseudotuberculosis. In mice, intestinal epithelial-intrinsic NAIP/NLRC4 and caspase-11

inflammasome responses restrict intraepithelial bacterial burdens during infection with enteric

pathogens, including *Salmonella*, *Shigella*, enteropathogenic *Escherichia coli* (EPEC), and

Citrobacter rodentium (49–55). The NAIP/NLRC4 inflammasome is not present in human IECs,

wherein the human homolog of caspase-11, caspase-4, instead plays a critical role in controlling

bacterial burdens and driving expulsion of infected cells (49, 56, 57), highlighting species-

specific differences as well as cell type-specific differences in inflammasome responses.

Importantly, many enteric bacterial pathogens use secreted virulence factors to inhibit IEC death

in order to preserve their replicative niche during early infection (58, 59). *Y. enterocolitica* uses

two anti-phagocytic Yops, YopE and YopH, to suppress caspase-1 and NLRP3-dependent

inflammasome activation in human IECs (60). However, the role of other Yops or other

inflammasomes such as the caspase-4 inflammasome, is unknown.

Here, we find that in contrast to mouse macrophages, the *Y. pseudotuberculosis* (*Yptb*)

effector YopJ does not induce cell death in human IECs or macrophages. In contrast, during

human IEC infection, *Yptb* T3SS-secreted effectors suppress caspase-4 inflammasome and

GSDMD-dependent IL-18 release. Surprisingly, NLRP3 was dispensable for inflammasome

activation induced by effector-less *Yptb*, as was NAIP/NLRC4, consistent with lack of

expression of both inflammasomes in human IECs (56). Both caspase-1 and caspase-8 partially

contributed to inflammasome activation, although neither was absolutely required. Instead,

inflammasome activation was entirely dependent on caspase-4. Three of the six injected Yops,

YopE, YopH, and YopK, collectively suppress activation of caspase-4. Mechanistically, YopE

and YopH blockade of bacterial phagocytosis prevented accumulation of intracellular bacteria in

IECs and correlated with reduced inflammasome activation. Like in human IECs, we found that

YopE, YopH, and YopK also synergistically suppress inflammasome activation in human

macrophages, a departure from our current understanding of murine macrophage

inflammasome responses to *Yersinia*. These findings demonstrate a key role for disruption of

actin-mediated phagocytosis in *Yersinia* evasion of the noncanonical inflammasome in human

intestinal epithelial cells and macrophages, thus uncovering important species and cell-type

specific differences in inflammasome responses to *Yersinia* infection.

Results

Yptb effectors suppress T3SS-dependent inflammasome activation in human IECs

During infection of murine macrophages, *Yersinia* injects Yops to suppress

inflammasome activation (42, 43, 45, 46). However, wild type (WT) *Yersinia* induces rapid cell

death (Fig. 1A) and caspase-1 activation due to YopJ-mediated inhibition of NF-κB and MAPK

signaling, enabling antibacterial defense despite effector-mediated immune modulation (38, 44,

61, 62). In agreement with previous studies on *Y. enterocolitica* infection of human IECs (60),

infection of the human colorectal cell line Caco-2 with WT *Y. pseudotuberculosis* failed to induce

release of lactate dehydrogenase (LDH), a marker of dying cells, into the cell supernatant (Fig.

1B), suggesting that human IECs do not undergo cell death in response to WT *Yptb* infection.

Consistently, WT *Yptb*-infected cells contained high levels of ATP, indicating that these cells are

viable and that WT *Yptb* does not induce cell death in human IECs, in contrast to murine

macrophages (Fig. 1A-C).

Lack of cell death in *Yptb*-infected human IECs could be due to a failure to activate

programmed cell death pathways, or to suppression of programmed cell death by WT *Yersinia*.

During infection of human IECs, *Y. enterocolitica* deploys several injected effectors, specifically

YopH and YopE, to suppress inflammasome-dependent cell death (60). To determine if *Yptb*

similarly modulates cell death in human IECs using its injected effectors, we infected Caco-2

cells with a strain of *Yptb* lacking all six of its injected effectors *(Δ6 Yptb)* and measured

induction of cell death by LDH release and ATP viability assay. Notably, Δ*6 Yptb* triggered

significantly higher levels of cell death compared to either mock or WT *Yptb*-infected cells (Fig.

1B and 1C). Further, Δ*6 Yptb*-infected Caco-2 cells robustly cleaved and secreted the

inflammasome-dependent cytokine IL-18 compared to mock or WT *Yptb*-infected cells (Fig. 1D

and 1E). These data indicate that in the absence of its injected effectors, *Yersinia* triggers

inflammasome activation in human IECs, in agreement with prior studies (60). IL-18 release

following Δ*6 Yptb* infection was dose-dependent, whereas WT *Yptb* failed to induce IL-18

release across a range of increasing MOIs (Fig. S1A). Consistent with our and others' findings

that expression and release of the inflammasome-dependent cytokine IL-1β is very low in

human IECs (49, 56, 60), we did not detect IL-1β release during Δ*6 Yptb* infection (Fig. S1B).

Inflammasome activation in response to Δ*6 Yptb* infection is T3SS-dependent, as an isogenic

Yptb strain cured of its virulence plasmid encoding the T3SS did not induce IL-18 release in

human IECs (Fig. 1D). Intriguingly, coinfection of Caco-2 cells with Δ*6 Yptb* and increasing

doses of *WT Yptb* resulted in a dose-dependent decrease in inflammasome activation (Fig. 1F

S1C), indicating that the effectors from WT *Yptb* block inflammasome activation in trans, further

supporting a role for *Yersinia*-injected effectors in suppressing inflammasome activation.

Collectively, these results indicate that like *Y. enterocolitica*, *Yptb* Yops suppress inflammasome

activation in human IECs, a fundamentally distinct outcome from the induction of apoptosis and

pyroptosis triggered in murine macrophages during WT *Yersinia* infection (38, 44, 61, 62).

Caspase-4 is required for Δ*6 Yptb*-induced inflammasome activation in human IECs

Caspases play important roles in cleaving and activating inflammasome-dependent

cytokines and executing cell death (34). Pretreatment of Caco-2 cells with the pan-caspase

inhibitor ZVAD prior to infection with Δ6 Yptb completely abrogated IL-18 release and cell death

compared to the vehicle control (Fig. 2A, B), indicating that caspases likely mediate

inflammasome activation downstream of Δ6 Yptb infection in human IECs. As ZVAD broadly

inhibits multiple caspases, we next sought to determine which specific caspases are required for

Δ6 Yptb-induced inflammasome activation in human IECs. Caspase-1 is critical for

inflammasome-dependent cytokine release and pyroptosis during bacterial infection (34, 44, 52,

63), and is activated during Y. enterocolitica infection of Caco-2 cells (60). To evaluate the role

for caspase-1 in inflammasome activation during Δ6 Yptb infection, we took parallel genetic and

pharmacologic approaches by using $CASP1^{-/-}$ Caco-2 cells (56) and the caspase-1 inhibitor

YVAD respectively. Δ6 Yptb infection of two independent clones of $CASP1^{-/-}$ Caco-2 cells

resulted in a partial loss of IL-18 release compared to WT Caco-2 cells (Fig. S2A). Additionally,

WT Caco-2s pretreated with YVAD exhibited significant but incomplete loss of IL-18 release

after Δ6 Yptb infection as compared to DMSO vehicle-treated Caco-2 (Fig. S2B), suggesting

that caspase-1 contributes to, but is not absolutely required for, Δ6 Yptb-induced inflammasome

activation. Caspase-8 is activated in response to infection by multiple pathogens, including

Yersinia (44, 64, 65), and can process caspase-1 substrates such as IL-1β and GSDMD to

mediate pyroptosis in the absence of caspase-1 (52, 64, 66). We therefore performed siRNA

knockdown of CASP8 in WT Caco-2, which resulted in 70-80% knockdown of caspase-8 (Fig.

S2C). CASP8 knockdown resulted in a partial reduction in IL-18 release following Δ6 Yptb

infection (Fig. S2D). We observed a similar reduction when pretreating WT Caco-2 cells with the

caspase-8 inhibitor IETD (Fig. S2E), suggesting that like caspase-1, caspase-8 is partially

required for inflammasome activation during Δ6 Yptb infection. CASP8 siRNA knockdown in

$CASP1^{-/-}$ Caco-2 cells also failed to completely abrogate inflammasome activation during Δ6

Yptb infection (Fig. S2F, S2G). Together, these results suggest that while both caspase-1 and

caspase-8 are contributing to inflammasome activation during Δ6 Yptb infection of human IECs,

neither is absolutely required and additional caspases likely contribute.

Caspase-4 plays a critical role in human IECs in response to a variety of enteric

pathogens (49, 55–57), and its activation triggers both cell death and IL-18 release in intestinal

epithelial cells(49). To test whether caspase-4 contributes to inflammasome activation during Δ6

Yptb infection, we infected two independent single-cell clones of $CASP4^{-/-}$ Caco-2 cells (56) with

either WT or Δ6 Yptb. As expected, WT Yptb infection did not elicit inflammasome activation in

either WT or $CASP4^{-/-}$ Caco-2 cells. Δ6 Yptb infection of WT Caco-2 cells resulted in robust

inflammasome activation, leading to release of cleaved IL-18 and cell death. Notably, CASP4

deficiency in Caco-2 cells abrogated cleavage and release of active IL-18 and cell death in

response to Δ6 Yptb infection, indicating that caspase-4 is absolutely required for

inflammasome activation induced by Δ6 Yptb infection in human IECs (Fig. 2C-E). Caspase-5

contributes to Salmonella-induced inflammasome activation in Caco-2 cells (56). To test

whether caspase-5 also contributes to Δ6 Yptb-induced inflammasome activation, we treated

WT Caco-2 cells with either a control scramble siRNA or CASP5 siRNA. Knockdown of CASP5

resulted in a partial but significant decrease in IL-18 secretion and cell death (Fig. S3B and

S3C), suggesting that while caspase-5 plays a contributing role in inflammasome activation, it

may not be absolutely required. Collectively, these data indicate that caspase-4 is essential for

inflammasome activation in response to Δ6 Yptb infection, with caspase-1, -8, and -5 playing

contributory roles as well.

GSDMD pore formation downstream of caspase-4 activation is required for Δ6 Yptb-

induced IL-18 release and cell death in human IECs

Inflammasome activation leads to cleavage of the pore-forming protein GSDMD into its

active transmembrane-inserting N-terminal domain, leading to its oligomerization into a large

ungated pore (19, 25, 67). Formation of the GSDMD pore in the plasma membrane leads to

release of IL-1 family cytokines as well as osmotic cell lysis and death, collectively termed

"pyroptosis" (68–71). Caspase-4 cleaves and activates GSDMD via release of its N-terminal

domain (25, 72). Notably, Δ6 Yptb infection led to robust GSDMD cleavage in WT Caco-2 cells,

which was completely absent in CASP4$^{-/-}$ Caco-2 cells, indicating that caspase-4 is required for

GSDMD cleavage in human IECs in response to Yersinia lacking its secreted effectors (Fig.

3A). In contrast, consistent with a lack of observed cell death and IL-18 release, WT Yptb

infection did not elicit GSDMD cleavage in either WT or two independent CASP4$^{-/-}$ clones (Fig.

3A).

To test whether GSDMD is required for inflammasome activation in human IECs during

Δ6 Yptb infection, we pretreated Caco-2 cells with disulfiram, a chemical inhibitor of GSDMD

pore formation (73). Critically, disulfiram treatment completely abrogated IL-18 release and cell

death downstream of inflammasome activation in Δ6 Yptb-infected cells compared to infected

vehicle control treated cells (Fig. 3B and 3C).

The NLRP3 inflammasome can be activated a variety of stimuli during infection,

including potassium efflux downstream of caspase-4 and GSDMD activation (5–9). Previous

studies of human IEC responses during Y. enterocolitica infection identified a critical role for the

NLRP3 inflammasome (60). However, studies of human IECs during *Salmonella* infection found

that NLRP3 does not play a role in inflammasome activation, potentially due to very low levels of

NLRP3 expression in human IECs as compared to human macrophages (56). Interestingly, WT

Caco-2 cells pretreated with a chemical inhibitor of the NLRP3 inflammasome, MCC950,

underwent comparable levels of inflammasome activation in response to *Δ6 Yptb* infection as

infected vehicle control-treated Caco-2 cells (Fig. S4A). Stimulating Caco-2 cells with LPS and

nigericin, a known agonist of the NLRP3 inflammasome, also failed to induce IL-18 release,

further suggesting a lack of NLRP3 inflammasome activity in Caco-2 cells. The NAIP/NLRC4

inflammasome, which senses and responds to flagellin and type III secretion system ligands

(10–18) (Fig. S4B), and the inflammasome adaptor protein ASC (Fig. S4C) were also

dispensable for *Δ6 Yptb*-induced inflammasome activation, consistent with prior findings that

expression of these proteins is very low in human IECs (56). Collectively, these results indicate

that during *Δ6 Yptb* infection of human IECs, GSDMD cleavage and activation occurs

downstream of caspase-4 and is required for IL-18 release and cell death, but the canonical

NLRP3 and NAIP/NLRC4 inflammasomes, as well as broadly ASC-dependent inflammasomes,

are dispensable.

YopE, YopH, and YopK synergistically suppress inflammasome activation in human cells

Our findings demonstrate that *Yptb* lacking its entire repertoire of injected effectors

induce inflammasome activation in human IECs (Fig. 1). In contrast, Caco-2 cells infected with a

panel of *Yptb* mutant strains each lacking one of the six Yops failed to elicit IL-18 secretion (Fig.

4A), indicating that loss of any single secreted Yop was insufficient to alleviate inflammasome

suppression and that several Yops likely have overlapping functions in suppressing

inflammasome activation. Notably, single loss of YopK and YopM failed to induce

inflammasome activation (Fig. 4A), despite their roles in suppressing the NLRP3/caspase-11

and pyrin inflammasome respectively in murine macrophages (40, 43, 45, 46). Indeed, *Y.*

enterocolitica was previously proposed to regulate NLRP3 inflammasome activation in Caco-2

cells by a combination of YopE and YopH-mediated blockade of "outside-in" integrin signaling

(60). Consistently, *Yptb* lacking both YopE and YopH *(ΔyopEH Yptb)* elicited significantly

elevated IL-18 release downstream of inflammasome activation in Caco-2 cells, indicating that

combinatorial loss of both YopE and YopH was sufficient to alleviate inflammasome

suppression in human IECs (Fig. 4B). Nonetheless, IL-18 levels during *ΔyopEH Yptb* infection

were still significantly lower than IL-18 levels released during *Δ6 Yptb* infection (Fig. 4B),

suggesting that additional Yops contribute to inflammasome suppression during *Yersinia*

infection of human IECs.

YopK is a translocated effector that negatively regulates the translocation of other

effector proteins and T3SS components (42, 43, 45). In murine macrophages, YopK suppresses

inflammasome activation, whereas YopE and YopH do not contribute to inflammasome

suppression (42, 43, 45). Given that deletion of *yopK* alone failed to elicit IL-18 release in

human IECs (60) (Fig. 4A), we considered that YopK modulates inflammasome activation in

human IECs in a manner that is masked by YopE and YopH, perhaps because these effectors

are hypertranslocated in a *yopK* mutant. We therefore infected Caco-2 cells with a combined

mutant *Yptb* strain lacking *yopE*, *yopH* and *yopK (ΔyopEHK Yptb)* and assayed for IL-18

release downstream of inflammasome activation. Notably, IL-18 release during Δ*yopEHK Yptb*

infection was substantially elevated compared to Δ*yopEH Yptb,* and fully recapitulated levels of

IL-18 observed during Δ*6 Yptb* infection (Fig. 4B). These data indicate that YopE, YopH and

YopK function together to suppress inflammasome activation during infection. Importantly,

individual loss of YopK, dual loss of YopK and YopE *(ΔyopEK)* or dual loss of YopK and YopH

(ΔyopHK) all failed to induce inflammasome activation. Only in a *yopEH* mutant background did

additional deletion of YopK lead to an increase in IL-18 release (Fig. 4B). Further, as with Δ*6*

Yptb infection, Δ*yopEHK*-induced inflammasome activation was fully dependent on the

caspase-4 inflammasome (Fig. 4C, 4D). Taken together, these data suggest that YopE, YopH,

and YopK act to inhibit components of a shared inflammasome-activating pathway, a departure

from the murine macrophage literature where YopK and YopM but not YopE or H, suppress

inflammasome activation (42, 43, 45, 46)

Fundamental differences exist between human and murine macrophage inflammasome

responses to enteric pathogens such as *Salmonella* Typhimurium (74–76). As such, we

hypothesized that human macrophage responses to *Yptb* infection may similarly diverge from

those of murine macrophages. Infection of the human monocytic THP-1 cell line revealed that

Δ*yopEH* and Δ*yopEHK* infection induced IL-1β and pyroptosis downstream of inflammasome

activation (Fig. 4E and 4F). As was the case during infection of human IECs, Δ*yopEHK* infection

triggered higher levels of inflammasome activation than Δ*yopEH* infection in THP-1 cells (Fig.

4E and 4F). These data indicate that human macrophages behave similarly to human IECs, with

Yops E and H functioning together with YopK to suppress inflammasome activation. Intriguingly,

and in marked contrast to established findings in murine macrophages, WT *Yptb* also did not

induce cell death in human macrophages, implying that WT *Yptb* effectively suppresses cell

death in human IECs and macrophages (Fig. 4F vs 1A). These data suggest that species-

specific differences exist in the response to the activities of *Yersinia* effector proteins between

human and murine cells addition to cell-type specific differences. Overall, these results reveal a

role for Yops E, H and K in synergistically suppressing inflammasome activation in human IECs

and macrophages.

YopE and YopH inhibit caspase-4-dependent inflammasome activation by blocking actin-

dependent bacterial phagocytosis in human IECs

During *Y. enterocolitica* infection of human IECs, YopE and YopH suppress inflammasome

activation by disrupting signaling downstream of *Yersinia* invasin binding to host β1-integrin

receptor (60). Invasin-β1-integrin interactions initiate host cytoskeletal rearrangements to

facilitate bacterial internalization (77–80), and YopE and YopH disruption of focal adhesion

complexes and actin filamentation consequently inhibits *Yersinia* uptake into host cells (80–87).

Notably, consistent with both Δ*6* and Δ*yopEHK Yptb* infection, we found that caspase-4 was

absolutely required for inflammasome activation induced by Δ*yopEH Yptb* infection (Fig. 5A and

5B). Considering YopE and YopH's known role in inhibiting bacterial internalization, we

hypothesized that YopE and YopH suppression of inflammasome activation in human IECs is

linked to their antiphagocytic activity, which would limit bacterial internalization, subsequent

cytosolic delivery of LPS, and caspase-4 inflammasome activation. We therefore assessed *Yptb*

internalization into human IECs by measuring intracellular bacterial burdens at 2 hours post-

infection. As expected, intracellular bacterial burdens were lowest in WT *Yptb*-infected cells,

while Δ*yopEH*-infected cells had significantly elevated levels of intracellular bacteria, as did

Δ*yopEHK* and Δ*6*-infected cells (Fig. 5C and Fig. S5A). Δ*yopEHK* and Δ*6*-infected cells had

comparable levels of intracellular bacteria to Δ*yopEH*-infected cells at 2 hpi, indicating that

YopE and YopH regulate phagocytosis inhibition in human IECs, while YopK limits

inflammasome activation through a mechanism distinct from phagocytosis inhibition. To

corroborate our bacteriologic observation that YopE and YopH suppress IEC internalization of

Yersinia, we performed microscopic analysis on Caco-2 cells infected with GFP-expressing WT,

Δ*yopEH*, or Δ*6 Yptb*. In agreement with our CFU data, Δ*yopEH Yptb*-infected cells had a

significantly higher levels of intracellular bacteria (i.e. GFP only bacteria) than WT *Yptb*-infected

cells, and levels of intracellular bacteria in Δ*yopEH* and Δ*6 Yptb* infected cells were similar (Fig.

5D, 5E, S5B). As YopE and YopH block bacterial uptake via disruption of actin filamentation, we

thus asked whether pharmacological inhibition of the actin cytoskeleton would complement

inhibition of bacterial uptake during Δ*yopEH Yptb* infection of human IECs. Indeed, pretreating

Caco-2 cells with cytochalasin D, an inhibitor of actin polymerization, reduced levels of

intracellular bacteria following Δ*yopEH* infection to that of WT *Yptb* infection (Fig. 5F and S5C).

Furthermore, cytochalasin D treatment completely abrogated inflammasome activation during

Δ*yopEH* and Δ*6 Yptb* infection (Fig. 5G), suggesting that inhibition of phagocytosis restored

inflammasome suppression in Δ*yopEH Yptb*-infected human IECs. Collectively, these results

point to a link between YopE and YopH-mediated inhibition of phagocytosis and inflammasome

activation, offering a potential mechanism by which YopE and YopH mediate inflammasome

evasion by *Yptb* in human IECs.

Discussion

In this study, we demonstrate that *Y. pseudotuberculosis* injected effectors suppress

caspase-4-dependent inflammasome activation, GSDMD-mediated cell death, and IL-18

release in human intestinal epithelial cells (IECs) (Fig. 1-3, S1). Caspase-1, caspase-8 and

caspase-5 partially contribute, but none individually or in combination were required for this

response (Fig. S2, S3). Consistent with their low levels of expression in IECs, the NAIP/NLRC4

and NLRP3 inflammasomes did not contribute to this inflammasome activation, nor did the

adaptor protein ASC, which is utilized by multiple inflammasomes to mediate caspase

recruitment (Fig. S4). Collectively these data imply that canonical inflammasomes are not

activated during *Yptb* infection of IECs. Rather, we report here that YopE, YopH, and YopK act

in concert to suppress caspase-4 inflammasome activation in human IECs.

While loss of YopK alone had no effect on inflammasome suppression, YopK deletion in

the absence of YopE and YopH phenocopied Δ*6 Yptb* infection and resulted in maximum

inflammasome activation, suggesting that YopK suppresses a component of an inflammasome-

activating pathway masked by the activity of YopE and YopH (Fig. 4). Consistent with prior

findings in *Y. enterocolitica* infection (60), we found that WT *Yptb* does not induce YopJ-

dependent cell death in human IECs, in contrast to *Yersinia* interactions reported in murine

macrophages. Surprisingly, during infection of human macrophages, WT *Yptb* also fails to

induce cell death and deploys YopE, YopH and YopK to synergistically suppress inflammasome

activation (Fig. 4). Finally, we demonstrate, in agreement with previous findings (80–84), that

YopE and YopH block bacterial internalization into IECs. YopE and YopH are known to inhibit

cytoskeletal dynamics, and we found that chemical inhibition of actin polymerization mimicked

YopE and H by ablating both inflammasome activation and bacterial internalization (Fig. 5 and

S5).

Previous studies found that *Y. enterocolitica* infection of murine macrophages and

human IECs activates the NLRP3 inflammasome (45, 60). Surprisingly, we found that NLRP3

was dispensable for inflammasome activation during *Y. pseudotuberculosis* infection of Caco-2

cells (Fig. S4A). Furthermore, known activators of NLRP3 failed to induce inflammasome

activation in Caco-2 cells (56) (Fig. S4A). Consistently, NLRP3 expression is very low in Caco-2

cells and human intestinal enteroids and colonoids, and NLRP3 is dispensable for

inflammasome activation during *Salmonella* infection of human IECs (56). However, human

patient mutations in NLRP3 have been found to be associated with development of Crohn's

disease, suggesting that there may be a gastrointestinal role for this inflammasome in humans

despite its low expression (88). It is possible that differences in cell culture conditions or *Y.*

enterocolitica and *Y. pseudotuberculosis* infection properties may account for the differential

role of NLRP3 during respective infections of human IECs, and that under certain conditions,

low levels of NLRP3 are able to respond and execute inflammasome activation in human IECs.

We found instead that caspase-4 was absolutely required for inflammasome activation

during *Y. pseudotuberculosis* infection of human IECs (Fig. 2, 4C-D, 5A-B). Caspase-4 is more

highly expressed in human IECs than other inflammasome components (56), and a broad range

of intracellular enteric pathogens engage or inhibit caspase-4/11 in human and murine IECs,

respectively (49–59). In myeloid cells, activated caspase-4/11 directly executes cell death, but

requires NLRP3 and caspase-1 for downstream IL-18 maturation and secretion (24). However,

it has been reported that in IECs, caspase-1 is dispensable for IL-18 release, whereas caspase-

4 is absolutely required (49, 55, 57, 89). Indeed, we found that caspase-1 is not required for IL-

18 release or cell death during *Yptb* infection of human IECs, although it plays a partial role (Fig.

S2A and S2B). We similarly found a partial role for caspase-8 in cell death (Fig. S2D and S2E).

Caspase-8 and caspase-1 can be recruited to the same inflammasome complexes and can

have redundant or compensatory roles (52, 63). Moreover, caspase-8 cleaves and activates

caspase-1 in response to YopJ-dependent NF-κB blockade during *Yersinia* infection in murine

macrophages (44). However, we found that YopJ did not impact inflammasome activation

during *Y. pseudotuberculosis* infection of Caco-2 cells, indicating that fundamentally distinct

pathways mediate *Yptb*-induced cell death in IECs. Whether caspase-8 and capase-1 are

activated downstream of caspase-4 or whether they are acting in a parallel pathway during

Yersinia infection of human IECs is still an open question. Simultaneous knockdown of caspase-

1 and caspase-8 still allowed for release of IL-18 during Δ*6 Yptb* infection (Fig. S2F), suggesting

that caspase-1 and caspase-8 are unlikely to play overlapping roles and instead potentially act

sequentially. These results also raise the possibility that there is a role for other caspases, such

as caspase-10, in inflammasome activation in human IECs.

Our findings that YopE and YopH block bacterial internalization into human IECs are

consistent with known functions of YopE and H (80–84), and demonstrate that in the absence of

YopE and YopH, bacterial uptake and inflammasome activation are able to occur (Fig. 5 and

S4). We hypothesize that blocking uptake of the bacteria into the host cell enables *Yptb* to limit

delivery of bacterial LPS into the host cell, thereby limiting the activation of the noncanonical

inflammasome. These findings suggest a potential mechanism by which YopE and YopH enable

Yptb to evade caspase-4 inflammasome activation during infection. Intriguingly these results are

a departure from the paradigm of YopE-dependent activation of the pyrin inflammasome via

disruption of cytoskeletal dynamics in murine macrophages (40, 46). In the murine macrophage

system, this activation is suppressed by YopM, yet we found that during human IEC infection

YopM was dispensable for inflammasome suppression even in the presence of YopE (Fig. 4A).

A previous study proposed that during *Y. entercolitica* infection of IECs, YopE and YopH

suppress inflammasome activation by blocking a critical priming "first signal" downstream of

integrin signaling that upregulates IL-18 transcript levels (60). The downstream signaling

elements of integrin binding are crucial for initiation of phagocytosis as well, and thus it is

possible that YopE and YopH disruption of FAK and RhoGTPase signaling contributes to

inflammasome suppression both by limiting intracellular LPS and inhibition of the "first signal".

After being internalized, *Yersinia* enters a membrane-enclosed vacuole that it remodels to

facilitate intracellular survival (90–92). In murine macrophages, YopK prevents LPS-mediated

noncanonical inflammasome activation by limiting hypertranslocation of T3SS components

YopB and D, thus maintaining vacuolar integrity and preventing destabilization of the *Yersinia*-

containing vacuole (42, 43). Critically, in human IECs, we found that YopK contributes to

caspase-4-inflammasome suppression downstream of YopE and YopH and that inflammasome

activation in the absence of Yops E, H and K is T3SS-dependent (Fig. 4B and 1D). It is possible

that in the absence of YopE, YopH and YopK, *Yersinia* is more readily taken up into IECs

downstream of integrin binding, and *Yersinia*-derived LPS is somehow exposed to the cytosol

from within its vacuole, perhaps due to T3SS activity. Further studies are needed to determine

whether *Yersinia* LPS is exposed to the cytosol and binds to caspase-4, as well as the precise

contribution of YopE, YopH and YopK to inhibition of this process in human IECs.

The synergistic blockade of caspase-4 inflammasome activation by YopE, YopH, and

YopK, as well as absence of YopJ-induced cell death, in both human IECs and THP1

macrophages represents a fundamentally distinct model from the murine macrophage

response, suggesting that *Yersinia* infection may trigger both cell type-specific and species-

specific inflammasome responses. Varying cellular properties and general organ systems needs

may underly these differential responses. For example, as components of a physical as well as

innate immune barrier, it may be advantageous for IECs to modulate and execute tightly

coordinated responses downstream of virulence detection in order to avoid disruption to the

monolayer that bacteria can use as access points (84, 93). Whether inflammasome activation

induces changes in the intestinal epithelium during *Yersinia* infection is unknown and would be

worth exploring in the future. Furthermore, unlike macrophages, IECs are not phagocytic and as

such, cytosolic LPS may be a more specific indicator of virulence activity in IECs, leading to a

stronger reliance on caspase-4-mediated responses. Surprisingly, during human macrophage

infection *Yptb* also deployed Yops E, H and K to suppress inflammasome activation and failed

to induce YopJ-dependent cell death, suggesting that certain inflammasome dynamics may be

shared between multiple cell types within a given species and differ broadly between disparate

species like mice and humans. Further studies comparing inflammasome responses between

murine and human IECs and murine and human macrophages will provide important insight into

cell type-specific and species-specific differences in inflammasome pathways.

Overall, our data demonstrate that *Yersinia* employs three injected Yops, YopE, YopH,

and YopK, to inhibit caspase-4 and GSDMD-dependent inflammasome activation in human

IECs. These Yops also inhibit inflammasome activation in human macrophages, in a manner

distinct from what has been previously observed in murine macrophages. These findings lend

further support to the critical role for the caspase-4 inflammasome in human IECs during host

defense against enteric pathogens and provide new insight into innate immune interactions

between *Yersinia* and a critical component of intestinal immunity. Moreover, it highlights how

inflammasome responses can vary between different cell types and between mice and humans,

which provides crucial insight into how inflammasome responses contribute to human health

and disease.

Materials and Methods

Bacterial strains and growth conditions

Yersinia strains are described in Table S1 in the supplemental material. Δ*yopEHK*, Δ*yopEK*, and Δ*yopHK* were generated by introducing a frameshift mutation of the *yopK* open reading frame into the Δ*yopEH*, Δ*yopE* and Δ*yopH* backgrounds respectively using a plasmid provided by Dr. James Bliska and an allelic exchange method (94). Δ*yopM* was generated by introducing an unmarked deletion of the *yopM* open reading frame into IP2666 using a plasmid provided by Dr. James Bliska and the same allelic exchange method. Yersiniae were cultured overnight at 26°C with aeration in 2x yeast extract-tryptone (YT) broth. To induce T3SS expression, in the morning, the bacteria were diluted into fresh 2xYT containing 20 mM sodium oxalate and 20 mM MgCl$_2$. Bacteria were grown with aeration for 1 hour at 26°C followed by 2 hour at 37°C prior to infection. All cultures were pelleted at 6000 x *g* for 3 min and resuspended in phosphate-buffered saline (PBS). Cells were infected at an MOI of 60 unless otherwise indicated, centrifuged at 290 x *g* for 10 min and incubated at 37°C. At 1 hour post-infection, cells were treated with 20 ng/ml or 100 ng/ml of gentamicin for 6 hour or 2 hour time points respectively. Infections proceeded at 37°C for the indicated length of time for each experiment. In all experiments control cells were mock infected with PBS.

Cell culture of cell lines

Caco-2 cells (HTB-27; American Type Culture Collection) were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% (vol/vol) heat-inactivated fetal bovine serum (FBS), 100 IU/mL penicillin and 100 μg/mL streptomycin. THP-1 cells (TIB-202; American Type Culture Collection) were maintained in RPMI supplemented with 10% (vol/vol) heat-inactivated fetal bovine serum (FBS), 0.05 nM β-mercaptoethanol, 100 IU/mL penicillin and 100 μg/mL streptomycin. All cells were grown at 37°C in a humidified incubator with 5% CO$_2$. One day prior to infection, Caco-2 cells were incubated with 0.25% trypsin-EDTA (Gibco) diluted 1:1 with 1 x PBS at 37°C for 15 min to dissociate cells. Trypsin was neutralized with serum-containing medium. Cells were replated in medium without antibiotics in a 24-well plate at a density of 3 x 10^5 cells/well. Two days prior to infection, THP-1 cells were replated in medium without antibiotics in a 48-well plate at a density of 2 x 10^5 cells/well and incubated with phorbol 12-myristate 13-acetate (PMA) for 24 hours to allow differentiation into macrophages.

Macrophages were primed with 100 ng/mL Pam3CSK4 (Invivogen) for 16 hours prior to

bacterial infections.

ELISAs

Supernatants harvested from infected cells were assayed using enzyme-linked immunosorbent

assay (ELISA) kits for human IL-18 (R&D Systems) and IL-1β (BD Biosciences).

LDH cytotoxicity assays

Supernatants harvested from infected cells were assayed for cytotoxicity by measuring loss of

cellular membrane integrity via lactate dehydrogenase (LDH) assay. LDH release was

quantified using an LDH Cytotoxicity Detection Kit (Clontech) according to the manufacturer's

instructions and normalized to mock-infected (min cytotoxicity) and 2% triton-treated cells (max

cytotoxicity)

Cell viability assay

Viability of infected cells was assessed using the CellTiter-Glo 2.0 Assay Kit (CTG2.0;

Promega) according to manufacturer's instructions. Caco-2 cells were seeded in 96-well flat

bottom white polystyrene TC-treated microplates (Corning) at a density of 7.5 x 10^4 cells/well in

medium without antibiotics. Cells were infected at an MOI of 60 and treated with 20 ng/mL

gentamicin 1hpi. After 6 hours, cells were treated with CTG2.0 reagent mix and incubated in the

dark at 37°C for 30 min. Luminescence was read on a luminometer and values normalized to

cells treated with 1% TritonX-100 (min cell viability) and mock-transfected cells (max cell

viability)

Immunoblot analysis

Cells were replated and infected on serum-free medium to collect supernatant samples.

Supernatant samples were centrifuged at 200 x g to pellet any cell debris and treated with

trichloroacetic acid (TCA) (25 μL TCA per 500 μL supernatant) overnight at 4°C. The following

day, TCA-treated samples were centrifuged at max speed (15,871 x g) for 15 min at 4°C and

washed with ice-cold acetone. TCA-precipitated supernatant samples and cell lysates were

resuspended in 1 x SDS-PAGE sample buffer and boiled for 5 min. Samples were separated by

SDS-PAGE on a 12% (vol/vol) acrylamide gel and transferred to polyvinylidene difluoride

(PVDF) Immobilon-P membranes (Millipore). Primary antibodies specific for human IL-18 (MLB

International PM014), β-actin (4967L; Cell Signaling) and GSDMD (G7422 Sigma-Aldrich) and

horseradish peroxidase (HRP)-conjugated secondary antibody anti-rabbit IgG (7074S; Cell

Signaling) were used. Enhanced chemiluminescence (ECL) Western blotting substrate or

SuperSignal West Femto (Pierce Thermo Scientific) HRP substrate were used for detection.

Inhibitor experiments

Cells were treated 1 h prior to infection at the indicated concentrations of the following inhibitors:

10 μM MCC950 (Sigma-Aldrich; PZ0280), 20 μM pan-caspase inhibitor Z-VAD(Ome)-FMK (SM

Biochemicals; SMFMK001), 20 μM caspase-1 inhibitor Ac-YVAD-cmk (Sigma-Aldrich;

SML0429), and 30 μM disulfiram (Sigma).

siRNA-mediated gene knockdown

CASP5 (S2417), *CASP8* (S2427), and two Silencer Select negative-control siRNAs (Silencer

Select negative control no. 1 and no. 2 siRNA) were purchased from Ambion (Life

Technologies). Three days before infection, 30 nM siRNA was transfected into Caco-2 cells

using Lipofectamine RNAiMAX transfection reagent (Thermo Fisher Scientific) following the

manufacturer's protocol.

Quantitative RT-PCR analysis

RNA was isolated using the RNeasy Plus Mini Kit (Qiagen) following the manufacturer's

instructions. Cells were lysed in 350 μL RLT buffer with β-mercaptoethanol and centrifuged

through a QIAshredder spin column (Qiagen). cDNA was synthesized from isolated RNA using

SuperScript II Reverse Transcriptase (Invitrogen) following the manufacturer's instructions.

Quantitative PCR was conducted with the CFX96 real-time system from Bio-Rad using the

SsoFast EvaGreen Supermix with Low ROX (Bio-Rad). For analysis, mRNA levels of siRNA-

treated cells were normalized to housekeeping gene *HPRT* and control siRNA-treated cells

using the $2^{-\Delta\Delta CT}$ (cycle threshold) method to calculate knockdown efficiency (95). The following

primers were used:

CASP5 forward: TTCAACACCACATAACGTGTCC

CASP5 reverse: GTCAAGGTTGCTCGTTCTATGG

CASP8 forward: GTTGTGTGGGGTAATGACAATCT

Casp8 reverse: TCAAAGGTCGTGGTCAAAGCC

HPRT forward: CCTGGCGTCGTGATTAGTGAT

HPRT reverse: AGACGTTCAGTCCTGTCCATAA

Bacterial uptake enumeration with colony forming units (CFUs)

Cells were infected with indicated strains of *Yersinia* at an MOI of 20. 1 hpi, cells were treated with 100 μg/mL of gentamicin to kill extracellular bacteria. 2 hpi the supernatants were aspirated and cells were lysed with PBS containing 0.5% Triton to collect intracellular bacteria. Harvested bacteria were serially diluted in PBS and plated on LB agar plates containing 2 μg/mL Irgasan. Plates were incubated at 28°C for two days and CFUIs were counted.

Fluorescence microscopy of intracellular *Yersinia*

One day before infection 2 x 10^5 cells/well were plated on glass coverslips in a 24-well plate. Cells were infected with indicated strains of *Yersinia* constitutively expressing GFP at an MOI of 20. At 2hpi, cells were washed 2 times with PBS, fixed with 4% paraformaldehyde for 10 min at 37°C and stored overnight at 4°C in PBS. The following day, cells were blocked for 30 min at room temperature in blocking solution containing 1% BSA in PBS and incubated for 1 h at room temperature in blocking solution with the polyclonal anti-*Yersinia* antibody SB349 diluted 1:1000 (kindly provided by Dr. James Bliska) (82). AF594-conjugated goat anti-Rabbit IgG antibody (A-11012 Thermo Fisher Scientific) was diluted 1:500 in blocking solution was added to cells and incubated for 45 min at room temperature. Cells were mounted on glass slides with DAPI mounting medium (Sigma Fluoroshield). Coverslips were imaged on an inverted fluorescence microscope (IX81; Olympus) and images were collected using a high-resolution charge-coupled-devise camera (FAST1394; QImaging) at a magnification of 60x. Images were analyzed and presented using SlideBook (version 5.0) software (Intelligent Imaging Innovations, Inc.). Average intracellular bacteria/cell and %intracellular bacteria were scored by counting 20 captures per coverslip across triplicate coverslips.

Statistical analysis

Prism 9.4.1 (GraphPad Software) was utilized for the graphing of data and all statistical analyses. Statistical significance for experiments were determined using the appropriate test and are indicated in each figure legend. Differences were considered statistically significant if the *p* value was <0.05.

Data availability

All data are included in the manuscript and supplemental material. Bacterial strains available upon request

Acknowledgments

We thank members of the Shin and Brodsky laboratories for helpful scientific discussions. We thank Dr. James Bliska for generously providing plasmids and Yop mutant strains.

This work is supported by NIH/NIAID grants AI151476, AI118861, AI123243 (S.S.), AI128630, AI163596, and AI139102 (I.E.B). S.S. and I.E.B. are both recipients of the Burroughs-Welcome Fund Investigators in the Pathogenesis of Infectious Disease Award. J.Z. is a recipient of the NIH/NIAID Microbial Pathogenesis and Genomics training grant 5T32AI141393-03.

Figure 1: *Yptb* T3SS-injected effectors suppress inflammasome activation in human IECs.
BMDMs (A) or Caco-2 cells (B-F) were infected with PBS (Mock or MOI 0), WT *Yptb*, or *Δ6*
Yptb. (A, B) Cell death was measured as percent cytotoxicity normalized to cells treated with
2% Triton at 6hpi. (C) Cell death was measured as percent viability normalized to mock-infected
cells at 6hpi. (D) Release of IL-18 into the supernatant was measured by ELISA at 6hpi. (E)
Lysates and supernatants collected 6hpi were immunoblotted for IL-18 and β-actin. (F) Release
of IL-1i into the supernatant was measured by ELISA at 6hpi. ns — not significant, ** p < 0.01,
*** p < 0.001, **** p < 0.0001 by Welch's t test (A-C) or by one-way ANOVA (D, F). Error bars
represent the standard deviation of triplicate wells and are representative of two or three
independent experiments.
608
Figure 2: caspase-4 is required for *Δ6 Yptb*-induced inflammasome activation in human IECs.
(A, B) One hour prior to infection, WT Caco-2 cells were treated with 20 uM ZVAD or DMSO as
a vehicle control. Cells were then infected with PBS (Mock), WT *Yptb* or *Δ6 Yptb*. At 6hpi (A)
release of IL18 into the supernatant was measured by ELISA and (B) cell death was measured
as percent cytotoxicity normalized to cells treated with 2% triton. (C-E) WT or two independent
single cell clones of *CASP4*-/- Caco-2 cells were infected with PBS (Mock), WT *Yptb* or *Δ6 Yptb*.
(C) Release of IL-18 into the supernatant was measured by ELISA at 6hpi. (D) Lysates and
supernatants collected 6hpi were immunoblotted for IL-18 and β-actin. (E) Cell death was
measured at 6hpi as percent cytotoxicity normalized to cells treated with 2% triton. ** p < 0.01,
*** p < 0.001, **** p < 0.0001 by two-way ANOVA. Error bars represent the standard deviation
of triplicate wells and are representative of two or three independent experiments.
620
Figure 3: GSDMD pore formation downstream of caspase-4 activation is required for *Δ6 Yptb*-
induced inflammasome activation in human IECs. (A) WT or two independent single cell clones
of *CASP4*-/- Caco-2 cells were infected with PBS (Mock), WT *Yptb* or *Δ6 Yptb*. Lysates and
supernatants were collected at 6hpi and immunoblotted for GSDMD and β-actin. (B, C) One
hour prior to infection, WT Caco-2 cells were treated with 30 uM disulfiram or DMSO as a
vehicle control. Cells were then infected with PBS (Mock), WT *Yptb* or *Δ6 Yptb*. (B) Release of
IL-18 into the supernatant and (C) percent cytotoxicity normalized to cells treated with 2% triton
were measured at 6hpi. **** p < 0.0001 by two-way ANOVA. Error bars represent the standard
deviation of triplicate wells and are representative of three independent experiments.
630
Figure 4: YopE, YopH, and YopK synergistically suppress inflammasome activation in human
cells. (A, B) WT Caco-2 cells were infected with PBS (Mock) or indicated strain of *Yptb*. Release
of IL-18 into the supernatant was measured at 6hpi. (C, D) WT or two independent single cell
clones of *CASP4*-/- Caco-2 cells were infected with PBS (Mock) or the indicated strain of *Yptb*.
(C) Release of IL-18 into the supernatant and (D) percent cytotoxicity normalized to cells treated
with 2% triton were measured at 6hpi. (E, F) WT THP-1 monocyte-derived macrophages were
primed with 100 ng/ml Pam2CSK4 for 16 hours. Cells were infected with PBS (Mock) of the
indicated strain of *Yptb*. (E) Release of IL-1β into the supernatant and (F) percent cytotoxicity
normalized to cells treated with 2% triton were measured at 6hpi. * p < 0.05, *** p < 0.001, **** p
< 0.0001 by one-way ANOVA (A, B, E, F) or two-way ANOVA (C, D). Error bars represent the
standard deviation of triplicate wells and are representative of two or three independent
experiments.
643
Figure 5: YopE and YopH inhibit caspase-4-dependent inflammasome activation and actin-
dependent bacterial phagocytosis in human IECs. (A, B) WT or two independent single cell
clones of *CASP4*-/- Caco-2 cells were infected with PBS (Mock) or the indicated strain of *Yptb*.
(A) Release of IL-18 into the supernatant and (B) percent cytotoxicity normalized to cells treated

with 2% triton were measured at 6hpi. (C) WT Caco-2 cells were infected with the indicated
strain of *Yptb* at an MOI=20 and lysed at 2 hpi. Bacteria were plated on *Yersinia*-selective agar
to calculate CFUs. (D, E) WT Caco-2 cells were seeded on glass coverslips and infected with
indicated strain of *Yptb* expressing GFP at an MOI=20. Cells were fixed at 2hpi and stained for
extracellular *Yersinia* (Red) and DAPI to label DNA (blue). (D) Representative images are
shown. Scale bar represents 10 μm. (E) Proportion of total bacteria that was intracellular (green
only) was scored by fluorescence microscopy. Bars represent standard deviation of triplicate
coverslips with 20 fields scored per coverslip. (F, G) One hour prior to infection, WT Caco-2
cells were treated with 10 uM cytochalasin D or DMSO as a vehicle control. Cells were then
infected with PBS (Mock) or indicated strain of *Yptb*. (F) Cells were lysed at 2hpi and bacteria
were plated on *Yersinia*-selective agar to calculate CFUs. (G) Release of IL-18 into the
supernatant was measured at 6hpi. * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$ by two-way ANOVA
(A, B, F, G) or one-way ANOVA (C, E). Error bars represent the standard deviation of triplicate
wells and are representative of two or three independent experiments
662
663

664 Table s1: *Yersinia* strains used in this study

Strain name	Relevant characteristics	Reference or Source
IP2666 (WT *Yptb*)	Wild-type, pYV$^+$, naturally *yopT$^-$*	(94)
Δ6 Yptb	*yopEHJMKO$^-$*	(62)
ΔyopE Yptb	*yopE-*	(82)
ΔyopH Yptb	*yopH-*	(85)
ΔyopM Yptb	*yopM-*	This study and (40)
ΔyopJ Yptb	*yopJ-*	(62)
ΔyopK Yptb	*yopK-*	(45)
ΔyopO Yptb	*yopO-*	(96)
ΔyopEH Yptb	*yopEH-*	(85)
ΔyopEK Yptb	*yopEK-*	This study and (97)
ΔyopHK Yptb	*yopHK-*	This study and (97)
ΔyopEHK Yptb	*yopEHK-*	This study and (97)

667 Figure s1 (related to Fig 1): *Yersinia* Yops suppress inflammasome activation. (A) Caco-2 cells
668 were infected with PBS (Mock) or WT *Yptb* or *Δ6 Yptb*. Release of IL-1β into the supernatant
669 was measured by ELISA at 6hpi. (B, C) Caco-2 cells were infected with PBS or WT *Yptb* or *Δ6*
670 *Yptb* at the indicated MOI. (B) Release of IL-18 into the supernatant was measured by ELISA at
671 6hpi. (C) Percent cytotoxicity normalized to cells treated with 2% triton were measured at 6hpi. *
672 $p < 0.05$, ** $p < 0.01$ by one-way ANOVA. Error bars represent the standard deviation of
673 triplicate wells and are representative of two independent experiments.

675 Figure s2 (related to Fig 2): Caspase-1 and caspase-8 are involved but not absolutely required
676 for *Δ6 Yptb*-induced inflammasome activation in human IECs. (A) WT Caco-2 cells or two
677 independent single cell clones of *CASP1*-/- Caco-2 cells were infected with PBS (Mock), WT
678 *Yptb*, or *Δ6 Yptb*. (B) One hour prior to infection, WT Caco-2 cells were treated with 20 uM
679 YVAD or DMSO as a vehicle control. Cells were then infected with PBS (Mock), WT *Yptb* or *Δ6*
680 *Yptb*. (A, B) IL-18 release into the supernatant was measured by ELISA at 6hpi. (C) Knockdown
681 efficiency of siRNA targeting *CASP8* in WT Caco-2 cells was measured by qRT-PCR and
682 normalized to housekeeping gene *HPRT* and calculated relative to control-siRNA-treated cells.
683 (D) WT Caco-2 cells were treated with siRNA targeting a control scrambled siRNA or siRNA
684 targeting *CASP8* for 72 hours. (E) One hour prior to infection WT Caco-2 cells were treated with
685 20 uM IETD or DMSO as a vehicle control. (D, E) Cells were infected with PBS (Mock), WT
686 *Yptb* or *Δ6 Yptb*. IL-18 release into the supernatant was measured at 6hpi. (F-G) WT or two
687 independent single cell clones of *CASP1*-/- Caco-2 cells were treated with siRNA targeting a
688 control scrambled siRNA or siRNA targeting *CASP8* for 72 hours. (F) IL-18 release into the
689 supernatant was measured at 6hpi. (G) Knockdown efficiency of siRNA targeting *CASP8* in WT
690 Caco-2 cells was measured by qRT-PCR and normalized to housekeeping gene *HPRT* and
691 calculated relative to control-siRNA-treated cells. ** $p < 0.01$, **** $p < 0.0001$ by two-way
692 ANOVA. Error bars represent the standard deviation of triplicate wells and are representative of
693 two or three independent experiments.

695 Figure s3 (related to Fig 2.): Caspase-5 contributes to *Δ6 Yptb*-induced inflammasome
696 activation in human IECs (A) Knockdown efficiency of siRNA targeting *CASP5* in WT Caco-2
697 cells was measured by qRT-PCR, normalized to *HPRT* and calculated relative to control-siRNA-

698   treated cells. (B, C) WT Caco-2 cells were treated with siRNA targeting a control scrambled
699   siRNA or siRNA targeting *CASP5* for 72 hours. (B) IL-18 release and (C) % cytotoxicity
700   normalized to cells treated with 2% triton were measured at 6hpi. ** $p < 0.01$, **** $p < 0.0001$ by
701   two-way ANOVA. Error bars represent the standard deviation of triplicate wells and are
702   representative of three independent experiments.
703
Figure s4 (related to Fig 3.) NLRP3, NAIP/NLRC4 and ASC are dispensable for *Δ6 Yptb*-
induced inflammasome activation in human IECs. (A) One hour prior to infection cells were
treated with 10 uM MCC950 or DMSO as a vehicle control. Cells were then infected with PBS
(Mock) WT *Yptb* or *Δ6 Yptb.* (B, C) WT or two independent single cell clones of (B) *NAIP*-/- or
(C) *PYCARD*-/- Caco-2 cells were infected with PBS (Mock), WT *Yptb*, or *Δ6 Yptb*. (A, B, C) IL-
18 release was measured in supernatants at 6hpi. **** $p < 0.0001$ by two-way ANOVA. Error
bars represent the standard deviation of triplicate wells and are representative of three
independent experiments.
712
Figure s5 (related to Fig. 5): YopE and YopH block phagocytosis in human IECs. (A) WT Caco-
2 cells were infected with the indicated strain of *Yptb* at an MOI=20 and lysed at 2 hpi. Bacteria
were plated on *Yersinia*-selective agar to enumerate CFUs and %internalization was calculated
as 2hpi CFUs divided by bacterial inoculum. (B) WT Caco-2 cells were seeded on glass
coverslips and infected with indicated strain of *Yptb* expressing GFP at an MOI=20. Cells were
fixed at 2hpi and stained for extracellular *Yersinia* (Red) and DAPI to label DNA (blue). Number
of intracellular bacteria (green only) in a field and number of cells was scored by fluorescence
microscopy. Bars represent standard deviation of triplicate coverslips with 20 fields scored per
coverslip. (C) One hour prior to infection, WT Caco-2 cells were treated with 10 uM cytochalasin
D or DMSO as a vehicle control. Cells were then infected with PBS (Mock) or indicated strain of
Yptb and lysed at 2hpi. Bacteria were plated *Yersinia*-selective agar to enumerate CFUs
and %internalization was calculated as 2hpi CFUs divided by bacterial inoculum. * $p < 0.05$, ** p
< 0.01, **** $p < 0.0001$ by one-way ANOVA (A, B) or two-way ANOVA (C). Error bars represent
the standard deviation of triplicate wells and are representative of two or three independent
experiments.
728

1.   Janeway CA. 1989. Approaching the Asymptote? Evolution and Revolution in
Immunology. Cold Spring Harb Symp Quant Biol 54:1–13.
2.   Janeway CA, Medzhitov R. 2003. Innate Immune Recognition.
https://doi.org/101146/annurev.immunol20083001084359 20:197–216.
3.   Vance RE, Isberg RR, Portnoy DA. 2009. Patterns of Pathogenesis: Discrimination of
Pathogenic and Nonpathogenic Microbes by the Innate Immune System. Cell Host
Microbe 6:10–21.
4.   Brodsky IE, Monack D. 2009. NLR-mediated control of inflammasome assembly in the
host response against bacterial pathogens. Semin Immunol 21:199–207.
5.   Lamkanfi M, Dixit VM. 2009. Inflammasomes: guardians of cytosolic sanctity. Immunol
Rev 227:95–105.
6.   Mariathasan S, Weiss DS, Newton K, McBride J, O'Rourke K, Roose-Girma M, Lee WP,
Weinrauch Y, Monack DM, Dixit VM. 2006. Cryopyrin activates the inflammasome in
response to toxins and ATP. Nat 2006 4407081 440:228–232.
7.   Muñoz-Planillo R, Kuffa P, Martínez-Colón G, Smith BL, Rajendiran TM, Núñez G. 2013.
K+ Efflux Is the Common Trigger of NLRP3 Inflammasome Activation by Bacterial Toxins
and Particulate Matter. Immunity 38:1142–1153.
8.   Hornung V, Bauernfeind F, Halle A, Samstad EO, Kono H, Rock KL, Fitzgerald KA, Latz
E. 2008. Silica crystals and aluminum salts activate the NALP3 inflammasome through
phagosomal destabilization. Nat Immunol 2008 98 9:847–856.
9.   Franchi L, Kanneganti TD, Dubyak GR, Núñez G. 2007. Differential requirement of P2X7
receptor and intracellular K+ for caspase-1 activation induced by intracellular and
extracellular bacteria. J Biol Chem 282:18810–18818.
10.   Ren T, Zamboni DS, Roy CR, Dietrich WF, Vance RE. 2006. Flagellin-Deficient
Legionella Mutants Evade Caspase-1- and Naip5-Mediated Macrophage Immunity. PLOS
Pathog 2:e18.
11.   Zhao Y, Yang J, Shi J, Gong Y-N, Lu Q, Xu H, Liu L, Shao F. 2011. The NLRC4
inflammasome receptors for bacterial flagellin and type III secretion apparatus
https://doi.org/10.1038/nature10510.
12.   Kofoed EM, Vance RE. 2011. Innate immune recognition of bacterial ligands by NAIPs
determines inflammasome specificity https://doi.org/10.1038/nature10394.
13.   Miao EA, Mao DP, Yudkovsky N, Bonneau R, Lorang CG, Warren SE, Leaf IA, Aderem
A. Innate immune detection of the type III secretion apparatus through the NLRC4
inflammasome https://doi.org/10.1073/pnas.0913087107.
14.   Zhao Y, Shi J, Shi X, Wang Y, Wang F, Shao F. 2016. Genetic functions of the NAIP
family of inflammasome receptors for bacterial ligands in mice. J Exp Med 213:647–656.
15.   Rauch I, Tenthorey JL, Nichols RD, Al Moussawi K, Kang JJ, Kang C, Kazmierczak BI,
Vance RE. 2016. NAIP proteins are required for cytosolic detection of specific bacterial
ligands in vivo. J Exp Med 213:657–665.
16.   Yang J, Zhao Y, Shi J, Shao F. 2013. Human NAIP and mouse NAIP1 recognize
bacterial type III secretion needle protein for inflammasome activation. Proc Natl Acad Sci
U S A 110:14408–14413.
17.   Rayamajhi M, Zak DE, Chavarria-Smith J, Vance RE, Miao EA. 2013. Cutting Edge:
Mouse NAIP1 Detects the Type III Secretion System Needle Protein. J Immunol
191:3986–3989.
18.   Molofsky AB, Byrne BG, Whitfield NN, Madigan CA, Fuse ET, Tateda K, Swanson MS.
2006. Cytosolic recognition of flagellin by mouse macrophages restricts Legionella
pneumophila infection. J Exp Med 203:1093–1104.
19.   Shi J, Zhao Y, Wang K, Shi X, Wang Y, Huang H, Zhuang Y, Cai T, Wang F, Shao F.
2015. Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell death.
Nat 2015 5267575 526:660–665.

20.   Li P, Allen H, Banerjee S, Franklin S, Herzog L, Johnston C, Mcdowell J, Paskind M,
781   Rodman L, Salfeld J, Towne E, Tracey D, Wardwell S, Wei F-Y, Wong W, Kamen R,
782   Seshadri T. 1995. Mice Deficient in IL-lp-Converting Enzyme Are Defective in Production
783   of Mature IL-lp and Resistant to Endotoxic Shock. Cell 80:401–411.
784   21.   Kuida K, Lippke JA, Ku G, Harding MW, Livingston DJ, Su MSS, Flavell RA. 1995.
785   Altered Cytokine Export and Apoptosis in Mice Deficient in Interleukin-1β Converting
786   Enzyme. Science (80-) 267:2000–2003.
787   22.   Cookson BT, Brennan MA. 2001. Pro-inflammatory programmed cell death [2]. Trends
788   Microbiol 9:113–114.
789   23.   Mariathasan S, Monack DM. 2007. Inflammasome adaptors and sensors: intracellular
790   regulators of infection and inflammation. Nat Rev Immunol 2007 71 7:31–40.
791   24.   Kayagaki N, Warming S, Lamkanfi M, Walle L Vande, Louie S, Dong J, Newton K, Qu Y,
792   Liu J, Heldens S, Zhang J, Lee WP, Roose-Girma M, Dixit VM. 2011. Non-canonical
793   inflammasome activation targets caspase-11. Nature 479:117–121.
794   25.   Kayagaki N, Stowe IB, Lee BL, O'Rourke K, Anderson K, Warming S, Cuellar T, Haley B,
795   Roose-Girma M, Phung QT, Liu PS, Lill JR, Li H, Wu J, Kummerfeld S, Zhang J, Lee WP,
796   Snipas SJ, Salvesen GS, Morris LX, Fitzgerald L, Zhang Y, Bertram EM, Goodnow CC,
797   Dixit VM. 2015. Caspase-11 cleaves gasdermin D for non-canonical inflammasome
798   signalling. Nat 2015 5267575 526:666–671.
799   26.   Rathinam VAK, Vanaja SK, Waggoner L, Sokolovska A, Becker C, Stuart LM, Leong JM,
800   Fitzgerald KA. 2012. TRIF licenses caspase-11-dependent NLRP3 inflammasome
801   activation by gram-negative bacteria. Cell 150:606–619.
802   27.   Gurung P, Subbarao Malireddi RK, Anand PK, Demon D, Vande Walle L, Liu Z, Vogel P,
803   Lamkanfi M, Kanneganti TD. 2012. Toll or interleukin-1 receptor (TIR) domain-containing
804   adaptor inducing interferon-β (TRIF)-mediated caspase-11 protease production
805   integrates toll-like receptor 4 (TLR4) protein- and Nlrp3 inflammasome-mediated host
806   defense against enteropathogens. J Biol Chem 287:34474–34483.
807   28.   Broz P, Ruby T, Belhocine K, Bouley DM, Kayagaki N, Dixit VM, Monack DM. 2012.
808   Caspase-11 increases susceptibility to Salmonella infection in the absence of caspase-1.
809   Nat 2012 4907419 490:288–291.
810   29.   Kayagaki N, Wong MT, Stowe IB, Ramani SR, Gonzalez LC, Akashi-Takamura S, Miyake
811   K, Zhang J, Lee WP, Muszyński A, Forsberg LS, Carlson RW, Dixit VM. 2013.
812   Noncanonical inflammasome activation by intracellular LPS independent of TLR4.
813   Science 341:1246–1249.
814   30.   Shi J, Zhao Y, Wang Y, Gao W, Ding J, Li P, Hu L, Shao F. 2014. Inflammatory caspases
815   are innate immune receptors for intracellular LPS. Nature 514:187–192.
816   31.   Hagar JA, Powell DA, Aachoui Y, Ernst RK, Miao EA. 2013. Cytoplasmic LPS activates
817   caspase-11: implications in TLR4-independent endotoxic shock. Science 341:1250–
818   1253.
819   32.   Lamkanfi M, Dixit VM. 2014. Mechanisms and Functions of Inflammasomes. Cell
820   157:1013–1022.
821   33.   Lamkanfi M, Dixit VM. 2010. Manipulation of Host Cell Death Pathways during Microbial
822   Infections. Cell Host Microbe 8:44–54.
823   34.   Labbé K, Saleh M. 2008. Cell death in the host response to infection. Cell Death Differ
824   2008 159 15:1339–1349.
825   35.   Cornelis GR, Wolf-Watz H. 1997. The Yersinia Yop virulon: a bacterial system for
826   subverting eukaryotic cells. Mol Microbiol 23:861–867.
827   36.   Bliska JB, Wang X, Viboud GI, Brodsky IE. 2013. Modulation of innate immune
828   responses by Yersinia type III secretion system translocators and effectors. Cell Microbiol
829   15:1622–1631.
37.   Viboud GI, Bliska JB. 2005. YERSINIA OUTER PROTEINS: Role in Modulation of Host

Cell Signaling Responses and Pathogenesis.
https://doi.org/101146/annurev.micro59030804121320 59:69–89.

38. Palmer LE, Hobble S, Galán JE, Bliska JB. 1998. YopJ of Yersinia pseudotuberculosis is
required for the inhibition of macrophage TNF-α production and downregulation of the
MAP kinases p38 and JNK. Mol Microbiol 27:953–965.

39. Orth K, Palmer LE, Bao ZQ, Stewart S, Rudolph AE, Bliska JB, Dixon JE. 1999. Inhibition
of the Mitogen-Activated Protein Kinase Kinase Superfamily by a Yersinia Effector.
Science (80-) 285:1920–1923.

40. Medici NP, Rashid M, Bliska JB. 2019. Characterization of pyrin dephosphorylation and
inflammasome activation in macrophages as triggered by the Yersinia effectors YopE and
YopT. Infect Immun 87.

41. Wang X, Parashar K, Sitaram A, Bliska JB. 2014. The GAP Activity of Type III Effector
YopE Triggers Killing of Yersinia in Macrophages. PLoS Pathog 10:1004346.

42. Zwack EE, Feeley EM, Burton AR, Baofeng H, Yamamoto M, Kanneganti T-D, Bliska JB,
Coers J, Brodsky IE. 2017. Guanylate Binding Proteins Regulate Inflammasome
Activation in Response to. Infect Immun 85:1–16.

43. Zwack EE, Snyder AG, Wynosky-Dolfi MA, Ruthel G, Philip NH, Marketon MM, Francis
MS, Bliska JB, Brodsky IE. 2015. Inflammasome activation in response to the yersinia
type III secretion system requires hyperinjection of translocon proteins YopB and YopD.
MBio 6:1–11.

44. Philip NH, Dillon CP, Snyder AG, Fitzgerald P, Wynosky-Dolfi MA, Zwack EE, Hu B,
Fitzgerald L, Mauldin EA, Copenhaver AM, Shin S, Wei L, Parker M, Zhang J, Oberst A,
Green DR, Brodsky IE. 2014. Caspase-8 mediates caspase-1 processing and innate
immune defense in response to bacterial blockade of NF- B and MAPK signaling. Proc
Natl Acad Sci 111:7385–7390.

45. Brodsky IE, Palm NW, Sadanand S, Ryndak MB, Sutterwala FS, Flavell RA, Bliska JB,
Medzhitov R. 2010. A Yersinia effector protein promotes virulence by preventing
inflammasome recognition of the type III secretion system. Cell Host Microbe 7:376–387.

46. Chung LK, Park YH, Zheng Y, Brodsky IE, Hearing P, Kastner DL, Chae JJ, Bliska JB.
2016. The Yersinia Virulence Factor YopM Hijacks Host Kinases to Inhibit Type III
Effector-Triggered Activation of the Pyrin Inflammasome. Cell Host Microbe 20:296–306.

47. Lopes Fischer N, Naseer N, Shin S, Brodsky IE. 2020. Effector-triggered immunity and
pathogen sensing in metazoans. Nat Microbiol 5:14–26.

48. Lei-Leston AC, Murphy AG, Maloy KJ. 2017. Epithelial Cell Inflammasomes in Intestinal
Immunity and Inflammation. Front Immunol 8:1168.

49. Knodler LA, Crowley SM, Sham HP, Yang H, Wrande M, Ma C, Ernst RK, Steele-
Mortimer O, Celli J, Vallance BA. 2014. Noncanonical inflammasome activation of
caspase-4/caspase-11 mediates epithelial defenses against enteric bacterial pathogens.
Cell Host Microbe 16:249–256.

50. Sellin ME, Müller AA, Felmy B, Dolowschiak T, Diard M, Tardivel A, Maslowski KM, Hardt
WD. 2014. Epithelium-intrinsic NAIP/NLRC4 inflammasome drives infected enterocyte
expulsion to restrict salmonella replication in the intestinal mucosa. Cell Host Microbe
16:237–248.

51. Hausmann A, Böck D, Geiser P, Berthold DL, Fattinger SA, Furter M, Bouman JA,
Barthel-Scherrer M, Lang CM, Bakkeren E, Kolinko I, Diard M, Bumann D, Slack E,
Regoes RR, Pilhofer M, Sellin ME, Hardt W-D. 2020. Intestinal epithelial NAIP/NLRC4
restricts systemic dissemination of the adapted pathogen Salmonella Typhimurium due to
site-specific bacterial PAMP expression. Mucosal Immunol
https://doi.org/10.1038/s41385-019-0247-0.

52. Rauch I, Deets KA, Ji DX, von Moltke J, Tenthorey JL, Lee AY, Philip NH, Ayres JS,
Brodsky IE, Gronert K, Vance RE. 2017. NAIP-NLRC4 Inflammasomes Coordinate

882   Intestinal Epithelial Cell Expulsion with Eicosanoid and IL-18 Release via Activation of
883   Caspase-1 and -8. Immunity 46:649–659.

884 53.   Mitchell PS, Roncaioli JL, Turcotte EA, Goers L, Chavez RA, Lee AY, Lesser CF, Rauch
885   I, Vance RE. 2020. Naip–nlrc4-deficient mice are susceptible to shigellosis. Elife 9:1–25.

886 54.   Roncaioli JL, Babirye JP, Chavez RA, Liu FL, Turcotte EA, Lee AY, Lesser CF, Vance
887   RE. 2022. A hierarchy of cell death pathways confers layered resistance to shigellosis in
888   mice. bioRxiv 2022.09.21.508939.

889 55.   Pallett MA, Crepin VF, Serafini N, Habibzay M, Kotik O, Sanchez-Garrido J, Di Santo JP,
890   Shenoy AR, Berger CN, Frankel G. 2016. Bacterial virulence factor inhibits caspase-4/11
891   activation in intestinal epithelial cells. Mucosal Immunol 2017 103 10:602–612.

892 56.   Naseer N, Zhang J, Bauer R, Constant DA, Nice TJ, Brodsky IE, Rauch I, Shin S. 2022.
893   Salmonella enterica Serovar Typhimurium Induces NAIP/NLRC4- And NLRP3/ASC-
894   Independent, Caspase-4-Dependent Inflammasome Activation in Human Intestinal
895   Epithelial Cells. Infect Immun 90.

896 57.   Kobayashi T, Ogawa M, Sanada T, Mimuro H, Kim M, Ashida H, Akakura R, Yoshida M,
897   Kawalec M, Reichhart JM, Mizushima T, Sasakawa C. 2013. The Shigella OspC3 effector
898   inhibits caspase-4, antagonizes inflammatory cell death, and promotes epithelial
899   infection. Cell Host Microbe 13:570–583.

900 58.   Ashida H, Mimuro H, Ogawa M, Kobayashi T, Sanada T, Kim M, Sasakawa C. 2011. Cell
901   death and infection: A double-edged sword for host and pathogen survival. J Cell Biol
902   195:931–942.

903 59.   Fink SL, Cookson BT. 2007. Pyroptosis and host cell death responses during Salmonella
904   infection. Cell Microbiol 9:2562–2570.

905 60.   Thinwa J, Segovia JA, Bose S, Dube PH. 2014. Integrin-Mediated First Signal for
906   Inflammasome Activation in Intestinal Epithelial Cells. J Immunol 193:1373–1382.

907 61.   Denecker G, Declercq W, Geuijen CAW, Boland A, Benabdillah R, Van Gurp M, Sory
908   MP, Vandenabeele P, Cornelis GR. 2001. Yersinia enterocolitica YopP-induced
909   apoptosis of macrophages involves the apoptotic signaling cascade upstream of bid. J
910   Biol Chem 276:19706–19714.

911 62.   Lilo S, Zheng Y, Bliska JB. 2008. Caspase-1 activation in macrophages infected with
912   Yersinia pestis KIM requires the type III secretion system effector YopJ. Infect Immun
913   76:3911–3923.

914 63.   Man SM, Hopkins LJ, Nugent E, Cox S, Glück IM, Tourlomousis P, Wright JA, Cicuta P,
915   Monie TP, Bryant CE. 2014. Inflammasome activation causes dual recruitment of NLRC4
916   and NLRP3 to the same macromolecular complex. Proc Natl Acad Sci U S A 111:7403–
917   7408.

918 64.   Sarhan J, Liu BC, Muendlein HI, Li P, Nilson R, Tang AY, Rongvaux A, Bunnell SC, Shao
919   F, Green DR, Poltorak A. 2018. Caspase-8 induces cleavage of gasdermin D to elicit
920   pyroptosis during Yersinia infection. Proc Natl Acad Sci U S A 115:E10888–E10897.

921 65.   Weng D, Marty-Roix R, Ganesan S, Proulx MK, Vladimer GI, Kaiser WJ, Mocarski ES,
922   Pouliot K, Chan FKM, Kelliher MA, Harris PA, Bertin J, Gough PJ, Shayakhmetov DM,
923   Goguen JD, Fitzgerald KA, Silverman N, Lien E. 2014. Caspase-8 and RIP kinases
924   regulate bacteria-induced innate immune responses and cell death. Proc Natl Acad Sci U
925   S A 111:7391–7396.

926 66.   Antonopoulos C, Russo HM, El Sanadi C, Martin BN, Li X, Kaiser WJ, Mocarski ES,
927   Dubyak GR. 2015. Caspase-8 as an Effector and Regulator of NLRP3 Inflammasome
928   Signaling. J Biol Chem 290:20167–20184.

929 67.   Agard NJ, Maltby D, Wells JA. 2010. Inflammatory stimuli regulate caspase substrate
930   profiles. Mol Cell Proteomics 9:880–893.

931 68.   He WT, Wan H, Hu L, Chen P, Wang X, Huang Z, Yang ZH, Zhong CQ, Han J. 2015.
932   Gasdermin D is an executor of pyroptosis and required for interleukin-1β secretion. Cell

Res 2015 2512 25:1285–1298.

69.   Liu X, Zhang Z, Ruan J, Pan Y, Magupalli VG, Wu H, Lieberman J. 2016. Inflammasome-
activated gasdermin D causes pyroptosis by forming membrane pores. Nat 2016
5357610 535:153–158.

70.   Sborgi L, Rühl S, Mulvihill E, Pipercevic J, Heilig R, Stahlberg H, Farady CJ, Müller DJ,
Broz P, Hiller S. 2016. GSDMD membrane pore formation constitutes the mechanism of
pyroptotic cell death. EMBO J 35:1766–1778.

71.   Ding J, Wang K, Liu W, She Y, Sun Q, Shi J, Sun H, Wang DC, Shao F. 2016. Pore-
forming activity and structural autoinhibition of the gasdermin family. Nat 2016 5357610
535:111–116.

72.   Aglietti RA, Estevez A, Gupta A, Ramirez MG, Liu PS, Kayagaki N, Ciferri C, Dixit VM,
Dueber EC. 2016. GsdmD p30 elicited by caspase-11 during pyroptosis forms pores in
membranes. Proc Natl Acad Sci U S A 113:7858–7863.

73.   Hu JJ, Liu X, Xia S, Zhang Z, Zhang Y, Zhao J, Ruan J, Luo X, Lou X, Bai Y, Wang J,
Hollingsworth LR, Magupalli VG, Zhao L, Luo HR, Kim J, Lieberman J, Wu H. 2020. FDA-
approved disulfiram inhibits pyroptosis by blocking gasdermin D pore formation. Nat
Immunol 2020 217 21:736–745.

74.   Valeria MRR, Ramirez J, Naseer N, Palacio NM, Siddarthan IJ, Yan BM, Boyer MA,
Pensinger DA, Sauer JD, Shin S. 2017. Broad detection of bacterial type III secretion
system and flagellin proteins by the human NAIP/NLRC4 inflammasome. Proc Natl Acad
Sci U S A 114:13242–13247.

75.   Naseer N, Egan MS, Reyes Ruiz VM, Scott WP, Hunter EN, Demissie T, Rauch I,
Brodsky IE, Shin S. 2022. Human NAIP/NLRC4 and NLRP3 inflammasomes detect
Salmonella type III secretion system activities to restrict intracellular bacterial replication.
PLoS Pathog 18.

76.   Gram AM, Wright JA, Pickering RJ, Lam NL, Booty LM, Webster SJ, Bryant CE. 2021.
Salmonella Flagellin Activates NAIP/NLRC4 and Canonical NLRP3 Inflammasomes in
Human Macrophages. J Immunol 206:631–640.

77.   Dersch P, Isberg RR. 2000. An Immunoglobulin Superfamily-Like Domain Unique to the
Yersinia pseudotuberculosis Invasin Protein Is Required for Stimulation of Bacterial
Uptake via Integrin Receptors. Infect Immun 68:2930.

78.   Isberg RR. 1989. Determinants for thermoinducible cell binding and plasmid-encoded
cellular penetration detected in the absence of the Yersinia pseudotuberculosis invasin
protein. Infect Immun 57:1998–2005.

79.   Isberg RR, Tran Van Nhieu G. 1994. Binding and internalization of microorganisms by
integrin receptors. Trends Microbiol 2:10–14.

80.   Fallman M, Andersson K, Hakansson S, Magnusson KE, Stendahl O, Wolf- Watz H.
1995. Yersinia pseudotuberculosis inhibits Fc receptor-mediated phagocytosis in J774
cells. Infect Immun 63:3117–3124.

81.   Andersson K, Carballeira N, Magnusson KE, Persson C, Stendahl O, Wolf-Watz H,
Fällman M. 1996. YopH of Yersinia pseudotuberculosis interrupts early phosphotyrosine
signalling associated with phagocytosis. Mol Microbiol 20:1057–1069.

82.   Black DS, Bliska JB. 2000. The RhoGAP activity of the Yersinia pseudotuberculosis
cytotoxin YopE is required for antiphagocytic function and virulence. Mol Microbiol
37:515–527.

83.   Cornelis GR. 2002. The Yersinia Ysc–Yop "Type III" weaponry. Nat Rev Mol Cell Biol
2002 310 3:742–753.

84.   Fasciano AC, Dasanayake GS, Estes MK, Zachos NC, Breault DT, Isberg RR, Tan S,
Mecsas J. 2021. Yersinia pseudotuberculosis YopE prevents uptake by M cells and
instigates M cell extrusion in human ileal enteroid-derived monolayers. Gut Microbes 13.

85.   Black DS, Bliska JB. 1997. Identification of p130Cas as a substrate of Yersinia YopH

(Yop51), a bacterial protein tyrosine phosphatase that translocates into mammalian cells
and targets focal adhesions. EMBO J 16:2730–2744.

86. Persson C, Carballeira N, Wolf-Watz H, Fällman M. 1997. The PTPase YopH inhibits
uptake of Yersinia, tyrosine phosphorylation of p130Cas and FAK, and the associated
accumulation of these proteins in peripheral focal adhesions. EMBO J 16:2307–2318.

87. Hamid N, Gustavsson A, Andersson K, McGee K, Persson C, Rudd CE, Fällman M.
1999. YopH dephosphorylates Cas and Fyn-binding protein in macrophages. Microb
Pathog 27:231–242.

88. Zhen Y, Zhang H. 2019. NLRP3 inflammasome and inflammatory bowel disease. Front
Immunol 10:276.

89. Holly MK, Han X, Zhao EJ, Crowley SM, Allaire JM, Knodler LA, Vallance BA, Smith JG.
2020. Salmonella enterica infection of murine and human enteroid-derived monolayers
elicits differential activation of epithelium-intrinsic inflammasomes. Infect Immun 88.

90. Mills SD, Finlayl BB. 1998. Isolation and characterization of Salmonella typhimurium and
Yersinia pseudotubercu/osis-containing phagosomes from infected mouse macrophages:
Y; pseudotuberculosis traffics to terminal lysosomes where they are degraded. Eur J Cell
Biol 77:35–47.

91. Fallman M, Andersson K, Hakansson S, Magnusson KE, Stendahl O, Wolf- Watz H.
1995. Yersinia pseudotuberculosis inhibits Fc receptor-mediated phagocytosis in J774
cells. Infect Immun 63:3117–3124.

92. Feeley EM, Pilla-Moffett DM, Zwack EE, Piro AS, Finethy R, Kolb JP, Martinez J, Brodsky
IE, Coers J. 2017. Galectin-3 directs antimicrobial guanylate binding proteins to vacuoles
furnished with bacterial secretion systems. Proc Natl Acad Sci U S A 114:E1698–E1706.

93. Ventayol PS, Geiser P, Di Martino ML, Florbrant A, Fattinger SA, Walder N, Sima E,
Shao F, Gekara NO, Sundbom M, Hardt WD, Webb DL, Hellström PM, Eriksson J, Sellin
ME. 2021. Bacterial detection by NAIP/NLRC4 elicits prompt contractions of intestinal
epithelial cell layers. Proc Natl Acad Sci U S A 118:1–11.

94. Zhang Y, Murtha J, Roberts MA, Siegel RM, Bliska JB. 2008. Type III secretion
decreases bacterial and host survival following phagocytosis of Yersinia
pseudotuberculosis by macrophages. Infect Immun 76:4299–4310.

95. Livak KJ, Schmittgen TD. 2001. Analysis of Relative Gene Expression Data Using Real-
Time Quantitative PCR and the 2−ΔΔCT Method. Methods 25:402–408.

96. Viboud GI, So SSK, Ryndak MB, Bliska JB. 2003. Proinflammatory signalling stimulated
by the type III translocation factor YopB is counteracted by multiple effectors in epithelial
cells infected with Yersinia pseudotuberculosis. Mol Microbiol 47:1305–1315.

97. Ryndak MB, Chung H, London E, Bliska JB. 2005. Role of predicted transmembrane
domains for Type III translocation, pore formation, and signaling by the Yersinia
pseudotuberculosis YopB protein. Infect Immun 73:2433–2443.

1022

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure S1

Figure S2

A

Average knockdown efficiency

	CASP5
WT Caco-2s	75%

B

C

Figure S4

Figure S5