Title

Overexpression of NIMA-related kinase suppresses cell proliferation and tip growth in a liverwort

Marchantia polymorpha

Running head

Overexpression of MpNEK1 suppresses growth

Authors

Hikari Mase[1], Yoshihiro Yoshitake[2], Takayuki Kohchi[2], Taku Takahashi[1], Hiroyasu Motose[1,*]

[1]Department of Biological Science, Graduate School of Natural Science & Technology, Okayama

University, Tsushimanaka 3-1-1, Okayama 700-8530, Japan

[2]Graduate School of Biostudies, Kyoto University, Kyoto 606-8502, Japan

* corresponding author

Hiroyasu Motose: motose@cc.okayama-u.ac.jp

Subject area: growth and development

Number of black and white figures 0

Color figures 12

Tables 0

Type and number of supplementary material: 7 supplementary figures, 7 videos

Abstract

NIMA-related kinases (NEKs) regulate a series of mitotic events in fungi and animals, whereas plant NEKs regulate growth direction of cells and organs. The liverwort *Marchantia polymorpha* has a single functional Mp*NEK1* gene, whose knockout leads to twisted growth of rhizoids. Mp*NEK1* is also expressed in the meristem of vegetative flat organ, thallus, while its function remains unknown. Here, we generated transgenic lines for the inducible expression of Mp*NEK1* using an estrogen receptor mediated system. Estradiol treatment efficiently induced the accumulation of Mp*NEK1* mRNA and MpNEK1-Citrine fusion protein throughout plant body. Overexpression of MpNEK1 severely suppressed growth of rhizoids and thalli, eventually causing the lethality of juvenile plants. The effect of estradiol was reversible until 3 days, whereas 7-days treatment resulted in irreversible suppression of growth. This severe effect was observed even at the nanomolar level of estradiol. EdU staining and microtubule imaging clearly indicated the suppression of cell proliferation by estradiol-induced MpNEK1. Unexpectedly, the overexpression of kinase-deficient MpNEK1 also suppressed thallus growth and rhizoid formation, despite their slightly mild effect than the full length MpNEK1, indicating phosphorylation-independent mechanism of growth suppression. In conclusion, overexpression of MpNEK1 suppresses cell division and elongation, leading to growth cessation and lethality. Our results imply that the expression of MpNEK1 is tightly regulated and plant NEKs might control cell division as in fungi and animals.

Keywords

Estradiol, *Marchantia polymorpha*, NIMA-related kinase, rhizoid, tip growth, thallus

Abbreviations

51   EdU; 5-ethynyl-2-deoxyuridine, EF1α; elongation factor1α, NEK; NIMA-related kinase, NIMA;

52   never in mitosis A, WDL, WAVE DANPENED-LIKE, XVE: LexA-VP16-ER artificial transcription

53   activator, WT; wild type

Introduction

Never in mitosis A (NIMA)-related kinase (NEK) is a Ser/Thr protein kinase that controls a series of mitotic events in fungi and animals (O'Connell et al. 2003, Fry et al. 2012), while plant NEK members have been shown to regulate directional growth (Motose, et al. 2008; Sakai, et al. 2008). *Arabidopsis* NEK6 suppresses ectopic outgrowth of epidermal cells through microtubule organization and interaction with other NEK members (Motose, et al. 2011). NEK6 localizes to the shrinking ends of microtubules, phosphorylates five amino acid residues of β-tubulin, and depolymerizes aberrant cortical microtubules to promote epidermal cell elongation (Takatani, et al. 2017). Furthermore, NEK6 has been shown to stabilize growth direction of hypocotyls through selectively removing cortical microtubules that best align with maximal tensile stress (Takatani, et al. 2020). NEK6-mediated attenuation of tensile-stress response increases the noise in microtubule orientation to buffer mechanical conflicts and to stabilize growth direction.

The liverwort *Marchantia polymorpha* is an emerging model to study evolution of land plants (Kohchi, et al. 2021) because of its low genetic redundancy (Bowman, et al. 2017) and various genetic tools (Ishizaki, et al. 2015; Ishizaki, et al. 2016). Indeed, *Arabidopsis* has seven NEK members, whereas *M. polymorpha* has a single *NEK* gene, Mp*NEK1* (Takatani, et al. 2015; Otani, et al. 2018). Thanks to these advantages, MpNEK1 has been shown to direct tip growth of rhizoids through microtubule depolymerization in the apical region (Otani, et al. 2018). Rhizoids are filamentous rooting cells, mainly elongating from the ventral epidermis and are thought to be required for plant anchorage to the soil and water/nutrient uptake (Jones and Dolan 2012; Shimamura, 2016; Kohchi, et al. 2021; Kanda, et al. 2022). Mp*nek1* mutant rhizoids frequently change their growth direction, resulting in the zigzag morphology (Otani, et al. 2018) and the defect of invasive growth to enter the hard substratum (Mase, et al. 2022). MpNEK1 localization in the apical microtubule foci and the increased microtubule stability in Mp*nek1* mutants demonstrate that MpNEK1 reorganizes and destabilizes apical microtubules to maintain growth polarity of rhizoids

(Otani, et al. 2018). Similar phenotype was observed in the mutants of a microtubule-associated

protein, WAVE DANPENED-LIKE (MpWDL) (Honkanen, et al. 2016; Champion, et al. 2021).

MpWDL might stabilize longitudinal microtubules in the shank region of rhizoids. Thus,

microtubule organization and dynamics are essential for directional growth of rhizoids.

However, it is not clear how MpNEK1 regulates microtubules to stabilize growth direction.

Furthermore, functional significance of Mp*NEK1* expression in the meristem of thalli remains

unclear. The gain of function approach could be helpful to elucidate these problems and to identify

downstream proteins that are phosphorylated by MpNEK1. Here, we adopted estradiol-inducible

system to overdrive Mp*NEK1* and to estimate its effects on growth and development in *M.*

polymorpha.

Results

Establishment of MpNEK1 inducible lines

In this study, we generated Mp*NEK1* inducible transgenic lines employing the XVE chimeric

transcriptional activator (Zuo, et al. 2000). This activator is composed of the DNA-binding domain

of LexA, the transcriptional activation domain of VP16, and the carboxyl estradiol binding domain

of the human estrogen receptor. This system has no apparent toxic physiological effects and has

been effectively used in flowering plants. The XVE vector was optimized for the application to *M.*

polymorpha in several points. Because the endogenous elongation factor Mp*EF1α* promoter is more

active in the meristem and sexual organ than the cauliflower mosaic virus *35S* promoter (Althoff, et

al. 2014), the Mp*EF1α* promoter was utilized to drive XVE. However, only a few research has been

reported on the application of XVE system in *M. polymorpha* (e.g. Flores-Sandoval, et al. 2016;

Furuya, et al. 2022).

We isolated eight transgenic lines by transformation of wild type sporelings with

pMpGWB344-MpNEK1 (MpEF1α pro:XVE>>LexAop:MpNEK1), in which XVE is expressed

under the control of Mp*EF1α* promoter and activates transcription of Mp*NEK1* through the binding

to the LexA operator in the presence of estradiol. Among them, seven transgenic lines (line #2 to 8)

were most identical to the wild type in the growth and morphology of thalli. These transgenic lines

were examined for the efficiency of Mp*NEK1* induction by quantitative real-time PCR (RT-qPCR)

(Supplementary Fig. S1). Two-week-old thalli of the wild type (WT) and transgenic lines were

transferred to the medium with or without β-estradiol and incubated for 24 hours. The expression

levels of Mp*NEK1* were remarkably increased in most transgenic lines (#2, 4, 5, 7, and 8).

Mp*NEK1* expression was 4-15 fold higher in estradiol treatment than in the estradiol-free medium.

Estradiol did not induce Mp*NEK1* expression in the wild type and line #6. Mp*NEK1* induction was

well reproducible and reliable, especially in line #2, 4, and 8, which exhibit no obvious defects in

growth and morphology without estradiol. Thus, these were used for the further analyses.

MpNEK1 induction severely suppresses thallus growth

We conducted phenotypic analysis to access the effects of estradiol-induced MpNEK1 on growth

and development. The gemmae of the wild type and Mp*NEK1* inducible lines were grown in the

presence or absence of exogenous β-estradiol for 24 days. There was no obvious effect of

β-estradiol on the growth and morphology of the wild type thalli (Supplementary Fig. S2). In the

inducible lines, thallus growth was severely suppressed when cultured with β-estradiol (Fig. 1).

Most gemmalings stopped growing and died in the presence of estradiol (Fig. 1), whereas the line

#6 did not show growth retardation (Supplementary Fig. S3). It was noteworthy that Mp*NEK1* was

not induced in the line #6 (Supplementary Fig. S1). These results clearly demonstrate that induction

of Mp*NEK1* severely suppresses thallus growth and causes juvenile lethality.

We next examined whether effect of Mp*NEK1* induction is reversible or not. The gemmalings of

the wild type and Mp*NEK1* inducible lines were grown in the presence of estradiol, and then

transferred to the estradiol-free medium at various times after estradiol treatment (Fig. 2). After 1

and 3-day treatment with estradiol, thallus growth seemed to be recovered, whereas irreversible

growth suppression was observed after 7 and 14-day treatment. In the wild type, growth inhibitory

effect of estradiol was not observed. Therefore, one-week Mp*NEK1* induction (estradiol treatment)

confers irreversible effect on thallus growth.

In the XVE induction system in flowering plants, 2 to 10 µM concentrations of estradiol have

been commonly used. In the Mp*NEK1* inducible lines, 10 µM estradiol caused severe growth

suppression and lethality. However, more mild conditions would be desirable to further investigate

MpNEK1 function. To determine less-effective conditions and dose-dependent effect of estradiol,

thalli of the wild type and Mp*NEK1* inducible lines were incubated in the presence of estradiol at

various concentrations (Fig. 3). The wild-type thallus showed no obvious phenotype in all

concentration ranges, whereas severe growth suppression was observed in the Mp*NEK1* inducible

lines even at the 10 and 100 nM of estradiol. From this result, Mp*NEK1* induction has the potent

effect on thallus growth.

We noticed that some plants obtained the resistance to estradiol and survived in the agar medium

supplemented with estradiol (Fig. 4). The estrogen resistance was observed in the next generation

gemmae, suggesting cell-cell movement of siRNA from the thallus to gemma. These

estradiol-resistant plants appeared more frequently in the lower concentrations and exhibited no

induction of Mp*NEK1* (Supplementary Fig. S4). At the lower concentrations, larger numbers of

thallus cells could survive and acquire estradiol resistance by gene silencing mechanism. Thus,

lower concentrations of estradiol are not desirable conditions, which often induce resistant plants.

MpNEK1 induction suppresses cell proliferation

Previous studies show that plant NEKs mainly regulate growth direction of cells and organs.

However, severe suppression of thallus growth by estradiol-induced Mp*NEK1* suggests some roles

of plant NEK in cell division and proliferation. To determine whether estradiol-induced Mp*NEK1*

affects cell division, we monitored cell proliferation activity by staining with EdU

(5-ethynyl-2'-deoxyuridine) (Fig. 5). This thymidine analog can be incorporated into DNA during S

phase. The subsequent reaction of EdU with a fluorescent azide labels newly replicated DNA strands of the cells in S phase during the incubation with EdU.

In both the wild type and inducible lines, EdU-positive nuclei were observed in the meristematic notch region in thalli grown in the absence of estradiol (Fig. 5). The amount of EdU-positive nuclei was almost similar in the wild type with or without estradiol, whereas it was significantly reduced in the estradiol-treated inducible lines. This result demonstrates that estradiol-induced Mp*NEK1* suppresses cell proliferation.

MpNEK1 induction suppresses mitotic cell division

Because *Arabidopsis* NEK6 depolymerizes specific microtubules, which detached from plasma membrane or aligned to the maximal tensile stress, to regulate cell expansion and organ growth (Takatani, et al. 2017; Takatani, et al. 2020), we monitored the effect of MpNEK1 overexpression on microtubule organization and mitotic apparatus. MpNEK1 inducible constructs were introduced into the transgenic lines expressing a microtubule marker Citrine-MpTUB2. We obtained the MpNEK1 inducible lines expressing Citrine-MpTUB2. The gemmae of these transgenic plants were grown for 3 days in the estradiol-free medium and then applied with the liquid medium with or without 10 μM estradiol. After 24 and 48 hours of treatment, thallus cells were observed with confocal microscopy. In the estradiol-treated thalli, number of mitotic cells was significantly decreased compared with the mock-treated thalli (Fig. 6A, B). We classified mitotic cells according to mitotic structures, prospindle, spindle, and phragmoplast, since these structures are clearly distinguished under confocal microscopy (Buschmann et al. 2016). Despite the slight increase of cells with prospindle, there are no considerable differences in the ratio of mitotic structures (Fig. 6C), suggesting that MpNEK1 induction does not severely affect mitotic progression. We also found that cortical microtubules were not affected with or without estradiol (Fig. 6A). Because we could not detect microtubule signal in rhizoids of the obtained transgenic lines, it could not be determined whether MpNEK1 induction affects microtubule organization in rhizoids. In summary,

overexpression of MpNEK1 suppresses mitotic cell division.

The overaccumulation of MpNEK1-Citrine suppresses growth of thalli and rhizoids

We generated inducible lines of MpNEK1-Citrine fusion protein (Fig. 7), in which protein

induction and localization could be easily detected by confocal microscopy, facilitating functional

analysis of MpNEK1. Eight transgenic lines were isolated by transformation of wild type sporelings

with pMpGWB144-MpNEK1-Citrine. Among them, four transgenic lines (line #1, 5, 7, and 8) were

most identical to the wild type in the absence of estradiol. When grown in the agar medium

supplemented with estradiol, these transgenic lines exhibited severe growth retardation and juvenile

lethality (Supplementary Fig. S5).

We next analyzed expression and localization of MpNEK1-Citrine fusion protein (Fig. 7B, C).

The gemmae of MpNEK1-Citrine inducible lines were grown for 4 days in the medium with or

without 1 µM β-estradiol and observed under a confocal microscope (Fig. 7B). MpNEK1-Citrine

was not detected in the absence of estradiol, whereas it was accumulated throughout

estradiol-treated thalli (Fig. 7B, Video S1). MpNEK1-Citrine localized in the cytoplasmic particles.

In the 24-hour estradiol treatment (2-day old gemmalings grown without estradiol were treated with

estradiol solution for 24 hours), similar expression and localization pattern were observed (Fig. 7C).

From these results, MpNEK1-Citrine was induced throughout thalli within 24 hours after estradiol

treatment and severely suppressed thallus growth.

We noticed accumulation of MpNEK1-Citrine particles in the meristem in some transgenic plants

even in the absence of estradiol (Supplementary Fig. S6). This leaky expression levels were varied

in each individual and transgenic line. In this study, we used plants and transgenic lines (line #1, 5,

7, and 8), which showed less leaky expression of MpNEK1-Citrine.

Taking accounts that MpNEK1 localizes in the apical microtubule foci to direct tip growth of

rhizoids (Otani, et al. 2018), we analyzed effect and localization of estradiol-induced

MpNEK1-Citrine in tip-growing rhizoids (Fig. 8). Because rhizoids were not formed when gemmae

were grown with estradiol at the start of culture, we added estradiol solution on the 3-day-old

gemmalings grown without estradiol to examine localization and effect of MpNEK1-Citrine in the

time course. After 24-hour treatment, MpNEK1-Citrine accumulated in the apical region of rhizoids

(Fig. 8A, Video S2). The expression level of MpNEK1-Citrine was varied in each rhizoid. A few

rhizoids showed highly accumulated MpNEK1-Citrine, which formed large droplets, moved to the

apical direction, and fused with each other (Video S3). To determine when MpNEK1-Citrine starts

to accumulate, we observed rhizoids at the early periods of estradiol treatment (Fig. 8B).

MpNEK1-Citrine began to accumulate at about 1 hour of treatment and gradually increased in the

rhizoid apex. After 48 and 120 hours, MpNEK1-Citrine accumulated throughout the entire region of

rhizoids in the particle and filamentous patterns (Fig. 8C, Video S4-S7). Tip growth was ceased or

severely reduced after 48 and 120 hours of estradiol treatment (Fig. 8D, Video S4-6). After 48 hours,

movement of MpNEK1-Citrine particles was obvious (Fig. 9A, Video S4, S5). MpNEK1-Citrine

particles moved toward the apex (anterograde) and basal region (retrograde) at the same velocity

(Fig. 9B). These results suggest that overaccumulation of MpNEK1 suppresses rhizoid growth.

The overexpression of kinase-deficient MpNEK1 suppresses growth of thalli and rhizoids

To determine whether the kinase activity of MpNEK1 is required for growth suppression, we

generated the inducible lines of MpNEK1 without its kinase activity (Fig. 10). For this purpose,

lysine-34 essential for the kinase activity was substituted with glutamic acid (K37E, MpNEK1^{K37E})

or the N-terminal kinase domain was deleted (KD deletion, MpNEK1^{dKD}) (Fig. 10A). MpNEK1^{K37E}

and MpNEK1^{dKD} were fused with Citrine at the C-termini and constructed into the XVE system as

shown in Fig. 7A.

We compared the effects of MpNEK1^{K37E} and MpNEK1^{dKD} induction to that of the full-length

MpNEK1 induction (Fig. 10, 11, Supplementary Fig. S7). The gemmae of the wild type and

inducible lines of full length MpNEK1, MpNEK1^{K37E}, and MpNEK1^{dKD} were grown in the medium

with (+ER) or without 1 µM estradiol (Mock). In the presence of estradiol, thallus growth was

suppressed in all inducible lines (Fig. 10B). This finding indicates that the kinase activity of

MpNEK1 is not required for the suppression of thallus growth. However, close-up observation

showed that the thalli of MpNEK1^{K37E} and MpNEK1^{dKD} inducible lines were larger than that of

full-length MpNEK1 inducible line (Fig. 10C). Furthermore, rhizoids were scarcely formed in the

full length inducible lines, whereas MpNEK1^{K37E} and MpNEK1^{dKD} inducible lines developed

several rhizoids, which elongated as those of the wild type (Fig. 10C).

We quantified the number of rhizoids, which seems to be a good indicator of the difference

among the inducible lines (Fig. 11). The gemmae of the wild type and inducible lines of full length

MpNEK1, MpNEK1^{K37E}, and MpNEK1^{dKD} were grown in the medium with (+ER) or without 1 μM

estradiol (Mock) for 7 days (Fig. 11A). The number of rhizoids in the wild type was not affected by

estradiol. Although the number of rhizoids was significantly decreased in all of the inducible lines,

the inducible lines of MpNEK1^{K37E} and MpNEK1^{dKD} develop more rhizoids than the full length

inducible lines. This difference was confirmed in the normalized data by using the average number

of rhizoids in estradiol-free medium as a standard (Fig. 11B). The difference between kinase

deficient and full length inducible lines was more clear in the quantification of rhizoids in 14-d

treatment (Fig. 11C). Thus, the induction of kinase-deficient MpNEK1 is slightly milder than the

full length induction. The kinase activity of MpNEK1 may have a little contribution to growth

suppression.

To investigate the accumulation and subcellular localization of kinase-deficient MpNEK1, the

inducible lines were grown in the medium with or without 1 μM estradiol and observed under

confocal microscopy (Fig. 12). Because the inducible lines of kinase-deficient MpNEK1 develop

larger thalli and a lot of rhizoids than the full length inducible lines, they provide good opportunity

to analyze localization of MpNEK1 in the growing organs and more healthy cells. Estradiol

treatment induced accumulation of MpNEK1^{K37E} and MpNEK1^{dKD} in the granules throughout the

thallus as the full length MpNEK1 (Fig. 12A).

We observed localization of MpNEK1^{K37E} and MpNEK1^{dKD} in rhizoids (Fig. 12B). Some

rhizoids showed granular accumulation throughout rhizoids (whole localization), while others

showed preferential localization at the apex of rhizoids (tip localization). We quantified the number

of rhizoids with the whole localization or tip localization (Fig. 12C). Tip-localized rhizoids were

approximately 47% in MpNEK1^{K37E} while 94% in MpNEK1^{dKD}. Thus, MpNEK1^{dKD} exhibited

relatively normal localization compared with MpNEK1^{K37E} in the inducible condition.

Next, we measured the growth velocity of rhizoids in the MpNEK1^{K37E} inducible lines (Fig. 12D).

The rhizoid growth was almost stopped after 48 hours of estradiol treatment. This is comparable to

the effect of the full length inducible lines. Therefore, excess accumulation of kinase-dead

MpNEK1 also suppressed rhizoid elongation.

Discussion

Here, we established and characterized induction system of MpNEK1 as a new tool to analyze

plant NEK function. Estradiol-based XVE induction system has been widely used in angiosperms,

but its application and characterization in *M. polymorpha* still remain limited. The efficient

induction of Mp*NEK1* transcripts and overaccumulation of MpNEK1-Citrine fusion protein

throughout plant body indicate that this system is very helpful for the functional analysis of

MpNEK1 and other proteins in *M. polymorpha*.

We noticed several unexpected phenomena in which caution is required. MpNEK1-Citrine

showed leaky expression in the meristem without estradiol. This might be due to the basal

expression of artificial transcriptional activator XVE under the control of Mp*EF1α* promoter, that is

highly active in meristematic tissues (Althoff, et al. 2014). Nonetheless, this leaky expression did

not affect thallus growth, suggesting that effective threshold of MpNEK1. Since the leaky

expression was varied in plants and transgenic lines, estrogen-free XVE-mediated transactivation

might be stochastic and/or responsive to fluctuating physiological states.

Secondly, estrogen-resistant silenced plants emerged as survivors in the prolonged treatment

especially at the lower concentrations of estradiol. The estrogen resistance could be transferred to

the next generation gemmae, implying thallus-to-gemma movement of siRNA. Thus, it would be

useful for the analysis of RNA silencing in *M. polymorpha*, while invokes concerns about treatment

time and estradiol concentration to minimize the contamination of silenced plants.

MpNEK1 induction severely suppressed thallus growth probably through cell proliferation. There

are several possible mechanisms of this growth-suppressive effect. Firstly, MpNEK1 may

depolymerize microtubule-based mitotic apparatus including spindle and phragmoplast or affect

cortical microtubule organization, which is required for directional growth prior to cell division.

However, our microtubule imaging showed that overexpression of MpNEK1 did not severely affect

mitotic apparatus and cortical microtubules. Secondly, overexpression of MpNEK1 may activate

cell cycle checkpoints to prevent cell cycle progression. According to the experiment of EdU

labeling, the cells through the S phase were reduced under the MpNEK1 overexpression.

Furthermore, MpNEK1 induction reduced mitotic cells in the M phase. Thus, MpNEK1 induction

may stop cell cycle in the G1 phase and eventually reduce the number of dividing cells. The *nimA*

mutants of a filamentous fungus *Aspergillus nidulans* exhibit cell cycle arrest just before M phase,

and conversely, overexpression of *NimA* causes premature chromosome condensation and aberrant

spindle formation, resulting in cell cycle arrest (Oakley and Morris, 1983; Osmani, et al. 1988).

Taking into accounts of remarkable promoter activity of Mp*NEK1* in the meristem (Otani, et al.

2018), it would be likely that MpNEK1 controls cell cycle progression and checkpoints.

Overexpression of MpNEK1 suppressed tip growth in rhizoids. Furthermore, overexpression

altered localization of MpNEK1-Citrine from the rhizoid apex to the entire region as the time of

estradiol treatment increased. This may reflect the overflow of MpNEK1 from the transport and

localization machineries. As the consequence, tip growth was stopped in rhizoids with this

dispersed localization pattern. Over-accumulated MpNEK1 may affect microtubule organization

and/or microtubule-dependent processes during rhizoid growth. Because the longitudinal

microtubules promote rhizoid elongation through organelle transport (Kanda, et al. 2022), it is

plausible that MpNEK1 accumulation throughout rhizoids suppresses transport system (see the last paragraph).

Unexpectedly, we found growth suppression of thalli and rhizoids by kinase-deficient MpNEK1. This result demonstrates that growth suppression is independent of protein phosphorylation by MpNEK1. Therefore, the induction system is not suitable for the identification of target proteins phosphorylated by MpNEK1. Nevertheless, kinase-dead MpNEK1 is slightly milder than the wild type MpNEK1 in the estradiol induction system. The kinase activity of MpNEK1 is not essential for the growth suppression, but may have a minor contribution. Excessively accumulated MpNEK1 might trap interacting proteins and substrates in the granules to suppress thallus growth and rhizoid elongation in a kinase-independent manner. Animal NEK7 licenses the assembly and activation of NLRP3 inflammasome, a large protein complex for the activation of inflammatory caspases, independently of its kinase activity (He et al. 2016, Sharif et al. 2019). Furthermore, NEK7 is required for oligomerization of NLRP3 and its adaptor protein ASC, which form the granular structure in the cytosol (He et al. 2016, Sharif et al. 2019). Plant NEK may also function as the assembly factor of protein supercomplex and cytosolic granular structures.

In summary, MpNEK1 induction system is useful for the analyses of its function in cell proliferation. Furthermore, it would be helpful to visualize directional protein transport in rhizoids because we observed MpNEK1 particles moved to the acropetal and basipetal direction (Video S3-S5). The velocity of MpNEK1 particles is quite similar to that of armadillo-repeat kinesin (Kanda et al. 2022), about half of velocity of growing microtubule plus ends, suggesting this motility is not due to plus-end tracking but rather to the kinesin-dependent transport along microtubules. MpNEK1 overexpression has the potent growth effect than that of *Arabidopsis* NEK6, whose overexpression caused slight reduction of root growth (Takatani, et al. 2017). Comparing with MpNEK1, NEK6-Citrine fusion protein is not highly accumulated in the XVE system (H. Mo., unpublished result), implying the differential degradation efficiency of NEK proteins and the potential advantage of protein induction and analyses in *M. polymorpha*.

Materials and Methods

Plant materials and growth conditions

M. polymorpha accessions Takaragaike-1 (Tak-1, male) and Takaragaike-2 (Tak-2, female) were used as the wild type strains. Plants were grown on the half-strength Gamborg's B5 medium solidified with 1% agar at 22°C under the light cycles with 16 hours of light and 8 hours of darkness.

Plasmid construction and transformation

Plasmids were mainly constructed based on the Gateway technology (Life technologies). Primers used in the plasmid construction are listed in Table S1. To generate a plasmid of Mp*NEK1* induction by estradiol, the full-length MpNEK1 coding sequence in the pENTR/D-TOPO entry vector (Otani et al. 2018) was transferred by LR reaction (LR clonase □ enzyme mix, Life Technologies) into the Gateway binary vector pMpGWB344 (MpEF1α pro:XVE >>LexA operator:Gateway cassette).

To generate a plasmid of estradiol-inducible MpNEK1-mCitrine, *mCitrine* coding sequence was amplified by KOD One (Toyobo) and a primer set (Table S1) and was cloned into the *Asc*I site of the pENTR/D-TOPO entry vector containing full length MpNEK1 CDS (Otani et al., 2018) by In-fusion system (Takara). The resulting vector pENTR/D-TOPO-MpNEK1-mCitrine was subjected to LR reaction to transfer MpNEK1-Citrine fusion into the Gateway binary vector pMpGWB144 (MpEF1α pro:XVE >>LexA operator:Gateway cassette).

To generate a plasmid of estradiol-inducible kinase-deficient MpNEK1, pENTR/D-TOPO-MpNEK1-mCitrine was subjected to the inverse PCR-based mutagenesis using KOD mutagenesis kit (Toyobo) and primer sets (Table S1). The resulting vectors, pENTR/D-TOPO-MpNEK1^{K37E}-mCitrine and pENTR/D-TOPO-MpNEK1^{dKD}-mCitrine were subjected to LR reaction to transfer mutated MpNEK1-Citrine fusion into the Gateway binary vector pMpGWB144 (MpEF1α pro:XVE >>LexA operator:Gateway cassette).

The constructs, pMpGWB344-MpNEK1 (MpEF1α pro:XVE>>LexA operator:MpNEK1), pMpGWB144-MpNEK1-Citrine (MpEF1α pro:XVE>>LexA operator:MpNEK1-Citrine), pMpGWB144-MpNEK1^{K37E}-Citrine (MpEF1α pro:XVE>>LexA operator:MpNEK1^{K37E}-Citrine), and pMpGWB144-MpNEK1^{dKD}-Citrine (MpEF1α pro:XVE>>LexA operator:MpNEK1^{dKD}-Citrine) were transformed into F1 sporelings derived from sexual crosses between Tak-2 and Tak-1 by *Agrobacterium*-mediated method (Ishizaki et al., 2008). The pMpGWB344-MpNEK1 construct was transformed into the regenerating thalli of CaMV35Spro:Citrine-MpTUB2 plants harboring pMpGWB105-MpTUB2 construct (Otani et al. 2018) according to the method of Kubota et al., 2013. Chlorsulfuron and hygromycin was used for the selection of transformants in pMpGWB344 and pMpGWB144, respectively. In pMpGWB344 and pMpGWB144, the artificial transcriptional activator XVE is expressed under the control of Mp*EF1α* promoter. XVE translocates to the nucleus in the presence of estradiol and binds to the LexA operator to induce the expression of MpNEK1 and MpNEK1-Citrine, respectively.

RT-qPCR

Total RNA was isolated from the thalli of the wild type and transgenic plants. For each sample, 0.5 µg of total RNA was reverse transcribed to cDNA using ReverTra Ace reverse transcriptase (Toyobo) according to the manufacturer's protocol. Real-time PCR was performed on a thermal cycler Dice Real Time System (Takara) using KOD SYBR qPCR Kit (Toyobo) according to the manufacturer's protocol. Transcript levels of Mp*EF1α* and Mp*ACT* were used as a reference for normalization (Saint-Marcoux et al., 2015). Primers used in RT-qPCR are listed in Table S1. RT-qPCR experiments were performed using four biological replicates.

Microscopy

The morphology of gemmalings and rhizoids was observed under a stereo microscope S8APO0 (Leica Microsystems) or a light microscope DM5000B (Leica) equipped with a CCD camera

DFC500 (Leica). To analyze the localization of MpNEK1-Citrine, plants were grown on

half-strength B5 medium for 3 and 5 days under the cycles with 16 hours of light and 8 hours of

darkness. For live imaging, half-strength B5 agar medium was solidified in the glass bottom dish

(35 mm in diameter x 10 mm in thickness, Thickness No. 1, Matsunami) and then the central region

of agar medium on the bottom glass strip was removed by tweezers. This region was poured with

about 200 µL of melted half-strength B5 agar medium to generate a thin solidified medium. The

gemmae were planted and grown in the central thin medium for 2-7 days. The details of this method

will be described elsewhere by H.M. Gemmalings were observed under a FV1200 confocal

laser-scanning microscope (Olympus) equipped with a high-sensitivity GaAsP detector and silicone

oil objective lenses (30×, 60× Olympus) or FV3000 (Olympus) equipped with high-sensitivity

GaAsP detectors and oil immersion objective lenses (40x, NA = 1.4; 60x, NA = 1.42, Olympus).

Silicone oil (SIL300CS, Olympus) or immersion oil (F30CC, Olympus) was used as immersion

media for the objective lenses. The samples were excited at 473 nm and 559 nm (laser diode) and

the emission was separated using a FV12-MHSY SDM560 filter (490-540 nm, 575-675 nm,

Olympus) in FV1200. The images were analyzed using ImageJ (National Institutes of Health, USA).

For estradiol treatment, an aliquot (200□300 µL) of 1/2 B5 liquid medium supplemented with

estradiol at a concentration of 1 µM or 10 µM was added to the gemmalings for the induction of

Mp*NEK1*. As a control, same amount of 1/2 B5 liquid medium supplemented with same

concentration of solvent (DMSO, less than 0.1%) was applied.

EdU staining

S-phase cells were visualized using Click-iT EdU Imaging Kits (Life Technologies) according to

the manufacturer's instruction. Gemmae were incubated on the half-strength Gamborg's B5 agar

medium supplemented with or without 1 µM β-estradiol for 4 days. Alternatively, for 1-day

treatment, gemmae were incubated on the agar medium without 1 µM β-estradiol for 3 days and

then transferred to and incubated in the agar medium with 1 µM β-estradiol for 1 day. Gemmalings

were then transferred to a solution containing 10 μM 5-ethynyl-2-deoxyuridine (EdU) and

incubated for 1 hour. Plants were fixed with 3.7 % formaldehyde solution in phosphate-buffered

saline (PBS) for 20 min. EdU incorporated into DNA was labelled with Alexa Fluor

488-azide-containing Click-iT reaction cocktail in the dark for 30 min. EdU-labelled cells were

observed using a confocal laser scanning microscope, FV1200 (Olympus). The maximum

Z-projection images were created using the ImageJ software (Schneider et al. 2012).

Funding

This work was supported by Grants-in-Aid from the JSPS (JSPS KAKENHI Grant Numbers

16K07403, 19K06709, 21H00370, and 23H04708 to H.Mo.), Grant-in-Aid for JSPS Fellows to

H.Ma. (23KJ1608), the Sasakawa Scientific Research Grant from The Japan Science Society to

H.Ma., Ryobi Teien Memory Foundation to H.Mo., and Naito Foundation to H.Mo.

Acknowledgements

We thank Drs. Ryuichi Nishihama and Kimitsune Ishizaki for their advice on EdU staining and

transformation.

Author contributions

HMa and HMo designed the research; HMa performed experiments and analyzed the data; YY and

TK developed XVE system; HMa and HMo wrote the manuscript; HMa, YY, TK, TT and HMo

revised the manuscript.

Conflict of interest

The authors declare that they have no conflicts of interest.

Data availability

The data of this study are available upon request from the corresponding author, Hiroyasu Motose.

Supplementary data

Fig S1. Effects of estradiol on the accumulation of Mp*NEK1* transcripts in the wild type and

transgenic lines introduced with *XVE::*Mp*NEK1*.

Fig S2. Effect of estradiol on thallus growth of the wild type.

Fig S3. Effect of estradiol on thallus growth of the transgenic line #6 with *XVE::*Mp*NEK1*.

Fig S4. Effect of estradiol on the accumulation of Mp*NEK1* transcripts in the wild type and

estradiol-resistant transgenic plants harboring *XVE::*Mp*NEK1*.

Fig S5. Effect of estradiol on the growth of MpNEK1-Citrine inducible lines

Fig S6. Expression and localization of MpNEK1-Citrine in the meristem of thallus without

estradiol.

Fig S7. Effect of estradiol on the growth of the wild type and MpNEK1-Citrine inducible lines

Video S1. Localization of MpNEK1-Citrine in a gemmaling grown for 4 days in the medium

supplemented with 1µM estradiol (same plant shown in Fig. 6B right panels).

Video S2. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine

inducible line treated with estradiol for 24 hours.

Video S3. Localization and dynamics of over-accumulated MpNEK1-Citine in a rhizoid of

MpNEK1-Citrine inducible line treated with estradiol for 24 hours.

Video S4. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine

inducible line treated with estradiol for 48 hours.

Video S5. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine

inducible line treated with estradiol for 48 hours.

Video S6. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine

inducible line treated with estradiol for 120 hours.

Video S7. Localization of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine inducible line treated

with estradiol for 120 hours.

References

Althoff F, Kopischke S, Zobell O, Ide K, Ishizaki K, Kohchi T, Zachgo S. 2014. Comparison of the

Mp*EF1α* and *CaMV35* promoters for application in *Marchantia polymorpha* overexpression studies.

Transgenic Research 23, 235-244.

Bowman JL, Kohchi T, Yamato KT, et al. 2017. Insights into land plant evolution garnered from the

Marchantia polymorpha genome. Cell 171: 287-304.

Buschmann H, Holtmannspötter M, Borchers A, O'Donoghue MT, Zachgo S (2016) Microtubule

dynamics of the centrosome-like polar organizers from the basal land plant Marchantia polymorpha.

New Phytol. 209, 999-1013.

Champion C, Lamers J, Jones VAS, Morieri G, Honkanen S, Dolan L. 2021. Microtubule associated

protein WAVE DAMPENED2-LIKE (WDL) controls microtubule bundling and the stability of the

site of tip-growth in *Marchantia polymorpha* rhizoids. PLoS Genetics 17, e1009533.

Flores-Sandoval E, Dierschke T, Fisher TJ, Bowman JL. 2016. Efficient and inducible use of

artificial microRNAs in *Marchantia polymorpha*. Plant & Cell Physiolgy 57, 281-290.

Fry AM, O'Regan L, Sabir SR, Bayliss R. 2012. Cell cycle regulation by the NEK family of protein

kinases. Journal of Cell Science 125, 4423-4433.

Furuya T, Nishihama R, Ishizaki K, Kohchi T, Fukuda H, Kondo Y. 2022. A glycogen synthase

kinase 3-like kinase MpGSK regulates cell differentiation in *Marchantia polymorpha*. Plant

Biotechnology 39, 65-72.

He Y, Zeng MY, Yang D, Motro B, Núñez G (2016) NEK7 is an essential mediator of NLRP3 activation downstream of potassium efflux. Nature 530, 354-359.

Honkanen S, Jones VAS, Morieri G, Champion C, Hetherington AJ, Kelly S, Proust H, Saint-Marcoux D, Prescott H, Dolan L. 2016. The mechanism forming the cell surface of tip-growing rooting cells is conserved among land plants. Current Biology 26, 3238–3244.

Ishizaki K, Chiyoda S, Yamato KT, Kohchi T. 2008. *Agrobacterium*-mediated transformation of the haploid liverwort *Marchantia polymorpha* L., an emerging model for plant biology. Plant & Cell Physiology 49, 1084-1091.

Ishizaki K, Nishihama R, Ueda M, Inoue K, Ishida S, Nishimura Y, Shikanai T, Kohchi T. 2015. Development of gateway binary vector series with four different selection markers for the liverwort *Marchantia polymorpha*. PLoS One 10, e0138876.

Ishizaki K, Nishihama R, Yamato KT, Kohchi T. 2016. Molecular genetic tools and techniques for *Marchantia polymorpha* research. Plant & Cell Physiology 57, 262-270.

Jones VAS, Dolan L. 2012. The evolution of root hairs and rhizoids. Annals of Botany 110, 205-212.

Kanda A, Otani K, Takahashi T, Motose H. 2022. Plant specific armadillo repeat kinesin directs organelle transport and microtubule convergence to promote tip growth. BioRxiv doi: https://doi.org/10.1101/2022.07.08.499237. [Preprint]

Kohchi T, Yamato KT, Ishizaki K, Yamaoka S, Nishihama R. 2021. Development and molecular

genetics of *Marchantia polymorpha*. Annual Review of Plant Biology 72, 677-702.

Kubota A, Ishizaki K, Hosaka M, Kohchi T (2013) Efficient *Agrobacterium*-mediated

transformation of the liverwort *Marchantia polymorpha* using regenerating thalli. Biosci.

Biotechnol. Biochem. 77, 167-172.

Mase H, Nakagami H, Okamoto T, Takahashi T, Motose H. 2022. Establishment and application of

novel culture methods in *Marchantia polymorpha*: persistent tip growth is required for substrate

penetration by rhizoids. Communicative & Integrative Biology 15, 164-167.

Motose H, Tominaga R, Wada T, Sugiyama M, Watanabe Y. 2008. A NIMA-related protein kinase

suppresses ectopic outgrowth of epidermal cells through its kinase activity and the association with

microtubules. Plant Journal 54, 829-844.

Motose H, Hamada T, Yoshimoto K, Murata T, Hasebe M, Watanabe Y, Hashimoto T, Sakai T,

Takahashi T. 2011. NIMA-related kinases 6, 4, and 5 interact with each other to regulate

microtubule organization during epidermal cell expansion in *Arabidopsis thaliana*. Plant Journal 67,

993-1005.

Oakley BR, Morris NR. 1983. A mutation in *Aspergillus nidulans* that blocks the transition from

interphase to prophase. Journal of Cell Biology 96, 1155-1158.

O'Connell MJ, Krien MJ, Hunter T. 2003. Never say never. The NIMA-related protein kinases in

mitotic control. Trends in Cell Biology 13, 221-228.

Osmani SA, Pu RT, Morris NR. 1988. Mitotic induction and maintenance by overexpression of a G2-specific gene that encodes a potential protein kinase. Cell 53, 237-244.

Otani K, Ishizaki K, Nishihama R, Takatani S, Kohchi T, Takahashi T, Motose H. 2018. An evolutionarily conserved NIMA-related kinase directs rhizoid tip growth in the basal land plant *Marchantia polymorpha*. Development 145, pii: dev154617.

Saint-Marcoux D, Proust H, Dolan L, Langdale JA. 2015. Identification of reference genes for real-time quantitative PCR experiments in the liverwort *Marchantia polymorpha*. PLoS One 10, e0118678.

Sakai T, Honing H, Nishioka M, et al. 2008. Armadillo repeat-containing kinesins and a NIMA-related kinase are required for epidermal-cell morphogenesis in *Arabidopsis*. Plant Journal 53, 157-171.

Sharif H, Wang L, Wang WL, Magupalli VG, Andreeva L, Qiao Q, Hauenstein AV, Wu Z, Núñez G, Mao Y, Wu H (2019) Structural mechanism for NEK7-licensed activation of NLRP3 inflammasome. Nature 570, 338-347.

Shimamura M. 2016. *Marchantia polymorpha*: taxonomy, phylogeny and morphology of a model system. Plant & Cell Physiology 57, 230-256.

Takatani S, Otani K, Kanazawa M, Takahashi T, Motose H. 2015. Structure, function, and evolution of plant NIMA-related kinases: implication for phosphorylation-dependent microtubule regulation. Journal of Plant Research 128, 875-891.

Takatani S, Ozawa S, Yagi N, Hotta T, Hashimoto T, Takahashi Y, Takahashi T, Motose H. 2017.

Directional cell expansion requires NIMA-related kinase 6 (NEK6)-mediated cortical microtubule

destabilization. Scientific Reports 7, 7826.

Takatani S, Verger S, Okamoto T, Takahashi T, Hamant O, Motose H. 2020. Microtubule response

to tensile stress is curbed by NEK6 to buffer growth variation in the *Arabidopsis* hypocotyl. Current

Biology 30, 1491-1503.

Zuo J, Niu QW, Chua NH. 2000. An estrogen receptor-based transactivator XVE mediates highly

inducible gene expression in transgenic plants. Plant Journal 24, 265-273.

Figure legend

Fig1. Effect of estradiol on thallus growth of Mp*NEK1* inducible line.

(A) Construct for Mp*NEK1* induction. Mp*EF1a* promoter, terminator (T), LexA operator (LexA op),

minimal CaMV35S promoter (m35Spro). The diagram is not shown to scale.

(B) The gemmae of Mp*NEK1* inducible line (#2) were planted in the agar medium with (+ER) or

without 10 µM estradiol (Mock) and grown for 3, 11, and 20 days.

(C) Time course of growth of Mp*NEK1* inducible line (#2) with (+ER) or without 10 µM estradiol

(Mock) .

(D) Quantification of thallus growth of Mp*NEK1* inducible line (#2, 4, 8) with (+ER) or without 10

µM estradiol (Mock). The mean projection area of thalli (n = 8-11 plants) was quantified by

ImageJ. Circles and error bars indicate mean values and standard deviations, respectively.

Fig 2. Reversible or irreversible effect of estradiol on thallus growth

(A) Schematic diagram of treatments. Gemmae were planted in the agar medium with 10 µM

estradiol, grown in the same medium for 1, 3, 7, or14 days, and then transferred to and grown in

the estradiol-free medium for 23 days.

(B) Gemmae of the wild type (WT) and four independent inducible lines (#2, 3, 4, and 8) were

grown as shown in (A). The periods of estradiol treatment is shown above the photographs

taken at the 23 day after transfer to the estradiol-free medium.

Fig 3. Effect of various concentrations of estradiol on thallus growth.

The gemmae of the wild type and Mp*NEK1* inducible lines were planted in the agar medium

supplemented with or without estradiol at the concentration of 10 nM, 100 nM, and 1000 nM and

grown for 13 (A) and 16 days (B).

Fig. 4. Estradiol resistance and its transfer to the next generation.

(A) The gemmae of the estradiol resistant plants or sensitive plants of Mp*NEK1* inducible lines

were planted in the agar medium supplemented with 10 μM estradiol and grown for 11, 15, and

21 days.

(B) Quantification of growth of gemmalings derived from the estradiol resistant plants or sensitive

plants of Mp*NEK1* inducible lines. The mean projection area of thalli (n = 10 plants) was

quantified by ImageJ. Circles and error bars indicate mean values and standard deviations,

respectively.

Fig. 5. Effect of estradiol on cell proliferation

(A) EdU-labelled nuclei in the gemmalings treated with 1 μM estradiol for 4 days. The gemmae of

the wild type (WT) and Mp*NEK1* inducible line were planted in the medium supplemented

with or without 1 μM estradiol, grown for 4 days, and incubated with 10 μM EdU for 1 hour in

liquid medium of the same kinds. EdU-labelled nuclei were visualized according to the

manufacture's instruction as described in methods.

630 (B) EdU-labelled nuclei in the gemmalings treated with 1 µM estradiol for 1 day. The gemmae of

631 the wild type (WT) and Mp*NEK1* inducible line were planted in the medium without estradiol,

632 grown for 3 days, and then transferred to and grown in the medium supplemented with or

633 without 1 µM estradiol for 1 day. The gemmalings were incubated with 10 µM EdU for 1 hour

634 in liquid medium of the same kinds. EdU-labelled nuclei were visualized according to the

635 manufacture's instruction as described in methods.

636 (C) Quantification of EdU-labeled nuclei in the thalli grown as in (A). Data are shown by box plots

637 (n = 12-16 plants). An asterisk indicates significant difference from the control (-ER) (*t*-test, P

638 < 0.05).

639 (D) Quantification of EdU-labeled nuclei in the thalli grown as in (B). Data are shown by box plots

640 (n = 12-16 plants). An asterisk indicates significant difference from the control (-ER) (*t*-test, P

641 < 0.05).

643 Fig. 6. Effect of MpNEK1 induction on microtubules

644 (A) The gemmae of MpNEK1 inducible line with a microtubule marker

645 CaMV35S:Citrine-MpTUB2 were grown in the estradiol-free medium for 3 days and then

646 applied with the liquid medium with (+ER) or without 10 µM estradiol (Mock). Thalli were

647 observed under a confocal microscope after 24 h and 48 h of treatment. Green: Microtubule,

648 magenta: plastid autofluorescence.

649 (B) Number of mitotic cells per notch in the MpNEK1 inducible line with

650 CaMV35S:Citrine-MpTUB2 after 24 h and 48 h of treatment with or without estradiol. Data are

651 shown in the box plot (n = 10 notches, N > 6 plants). An asterisk indicates significant difference

652 (t-test, P < 0.01).

653 (C) The classification of mitotic cells with prospindle, spindle, or phragmoplast (n indicates the

654 number of mitotic cells used for the classification).

Fig. 7. Localization of estradiol-induced MpNEK1-Citrine in the thallus

(A) Construct for Mp*NEK1* induction. Mp*EF1a* promoter, terminator (T), LexA operator (LexA op),

minimal CaMV35S promoter (m35Spro). The diagram is not shown to scale.

(B) The gemmae of the Mp*NEK1* inducible line were planted in the medium with (+ER) or without

1 µM estradiol (Mock), grown for 4 days, and observed under a confocal microscope.

(C) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol,

grown for 2 days, and then added with (+ER) or without 1 µM estradiol (Mock) for 1 day.

Upper panels are light field images merged with confocal images shown in the below panels.

Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

Fig. 8. Localization of estradiol-induced MpNEK1-Citrine in rhizoids.

(A) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol,

grown for 3 days, and then applied with (+ER) or without 1 µM estradiol (Mock). Rhizoids

were observed under a confocal microscope. Green: MpNEK1-Citrine, magenta: plastid

autofluorescence.

(B, C) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol,

grown for 3 days, and then applied with 1 µM estradiol and observed at the time indicated in

each panel. The same rhizoid was observed under a confocal microscope in the time course.

Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

(D) Velocity of rhizoid growth in the MpNEK1-Citrine inducible line incubated for 48 hours with

(+ER) or without 1 µM estradiol (-ER). Data are shown by box plots (n = 32 rhizoids). An

asterisk indicates significant difference (*t*-test, $P < 0.01$).

Fig. 9. Movement of MpNEK1-Citrine particles in rhizoids.

(A) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol,

grown for 3 days, and then applied with 1 µM estradiol. Rhizoids were observed under a

confocal microscope after 48 hours. The lower panel indicates kymograph of MpNEK1-Citrine

particles. Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

(B) Velocity of MpNEK1-Citrine particles. Data are shown by box plots (n = 24 or 31 particles).

There was no significant difference (*t*-test, P > 0.05).

Fig. 10. Effect of kinase-deficient MpNEK1 induction on the growth of thalli and rhizoids.

(A) Full length and kinase-deficient MpNEK1 used for estradiol induction.

(B) The gemmae of the wild type and inducible lines of full length MpNEK1, MpNEK1^{K37E}, or

MpNEK1^{dKD} (KD deletion) were planted in the medium with (+ER) or without 1 µM estradiol

(Mock), and grown for 14 days.

(C) The gemmae of the wild type and inducible lines of full length MpNEK1, MpNEK1^{K37E}, or

MpNEK1^{dKD} (KD deletion) were planted in the medium with (+ER) or without 1 µM estradiol

(Mock), and grown for 7 days.

Fig. 11. Effect of MpNEK1 induction on the number of rhizoids.

(A) The number of rhizoids in the gemmae grown for 7 days in the medium with (+ER) or without

1 µM estradiol (Mock). The gemmae of the wild type (WT) and inducible lines of full length

MpNEK1-Citrine (FL), MpNEK1^{K37E}-Citrine (K37E), and MpNEK1^{dKD}-Citrine (dKD) were

used. Data are shown by box plots (n = 10 plants). The different letters indicate significant

differences by Tukey's HSD test (P < 0.03). Asterisks indicate significant differences from the

value of FL with estradiol (+ER) (*t*-test, P < 0.01).

(B) The relative number of rhizoids in (A). The average number of rhizoids in estradiol-free media

(ER -) in each genotype was used for the normalization. Data are shown by box plots (n = 10

plants). The different letters indicate significant differences by Tukey's HSD test (P < 0.03).

Asterisks indicate significant differences from the value of FL with estradiol (+ER) (*t*-test, P <

0.01).

(C) The number of rhizoids in the gemmae grown for 14 days in the medium supplemented with 1

μM estradiol. The gemmae of the inducible lines of full length MpNEK1-Citrine (FL),

MpNEK1^{K37E}-Citrine (K37E), and MpNEK1^{dKD}-Citrine (dKD) were used. Data are shown by

box plots   (n indicates the number of plants). The different letters indicate significant

differences by Tukey's HSD test (P < 0.03).

Fig. 12. Localization of kinase-deficient MpNEK1-Citrine

(A) The gemmae of the inducible lines of MpNEK1^{K37E}-Citrine (K37E) or MpNEK1^{dKD}-Citrine

(KD deletion) were planted in the medium with (+ER) or without 1 μM estradiol (Mock),

grown for 4 days, and observed under a confocal microscope. Green: MpNEK1-Citrine,

magenta: plastid autofluorescence.

(B) The gemmae of the inducible lines of MpNEK1^{K37E}-Citrine (K37E) or MpNEK1^{dKD}-Citrine

(KD deletion) were grown as in (A) and rhizoids were observed under a confocal microscope.

Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

(C) Ratio of rhizoids with tip-localized or whole localized MpNEK1^{dKD}-Citrine (KD deletion) or

MpNEK1^{K37E}-Citrine (K37E) after 120 h of estradiol treatment.

(D) Growth velocity of rhizoids in the MpNEK1^{K37E}-Citrine inducible lines incubated for 48 hours

with (+ER) or without 1 μM estradiol (-ER). Data are shown by box plots (n = 30 rhizoids). An

asterisk indicates significant difference (*t*-test, P < 0.01).

Fig S1. Effects of estradiol on the accumulation of Mp*NEK1* transcripts in the wild type and

transgenic lines introduced with pMpGWB344-Mp*NEK1* (*XVE::*Mp*NEK1*).

The 10-day old plants of the wild type (WT) and the transgenic lines were incubated for 1 day in the

medium with (+) or without 10 μM estradiol (-). Total RNA was extracted and subjected to

RT-qPCR. All transcript levels are relative to that of mock-treated wild type plants. Mp*EF1α* and

Mp*ACT* were used as the control genes. Columns and error bars indicate mean values and standard

errors, respectively (n indicates the number of biological replicates).

Fig S2. Effect of estradiol on thallus growth of the wild type.

(A) The gemmae of the wild type (Tak-1) were planted in the agar medium with (+ER) or without

10 µM estradiol (Mock) and grown for 3, 11, and 20 days.

(B) Time course of growth of the wild type with (+ER) or without 10 µM estradiol (Mock).

(C) Quantification of thallus growth of the wild type with (+ER) or without 10 µM estradiol

(Mock). The mean projection area of thalli (n = 10 plants) was quantified by ImageJ. Circles

and error bars indicate mean values and standard deviations, respectively.

Fig S3. Effect of estradiol on thallus growth of the transgenic line #6 with *XVE::*Mp*NEK1*.

(A) The gemmae of the line (#6) were planted in the agar medium with (+ER) or without 10 µM

estradiol (Mock) and grown for 3, 11, and 20 days.

(B) Time course of growth of the line (#6) with (+ER) or without 10 µM estradiol (Mock) .

(C) Quantification of thallus growth of the line (#6) with (+ER) or without 10 µM estradiol (Mock).

The mean projection area of thalli (n = 10 plants) was quantified by ImageJ. Circles and error

bars indicate mean values and standard deviations, respectively.

Fig S4. Effect of estradiol on the accumulation of Mp*NEK1* transcripts in the wild type and

estradiol-resistant transgenic plants harboring *XVE::*Mp*NEK1*.

The 10-day old plants of the wild type (WT) and estradiol-resistant plants derived from line #8 and

#2 were incubated for 1 day in the medium with (+) or without 10 µM estradiol (-). Total RNA was

extracted and subjected to RT-qPCR. All transcript levels are relative to that of mock-treated wild

type plants. Mp*EF1α*were used as the control genes. Data are shown by box plots (n = 2 biological

replicates).

Fig S5. Effect of estradiol on the growth of MpNEK1-Citrine inducible lines

The gammae of the wild type (WT) and the MpNEK1-Citrine inducible lines were planted in the

agar medium supplemented with (+ER) or without 1 µM estradiol (Mock) and grown for 15 days.

Fig S6. Expression and localization of MpNEK1-Citrine in the meristem of thallus without

estradiol.

Left panel shows a light field image merged with a confocal image in the right panel. Green;

MpNEK1-Citrine, magenta; plastid autofluorescence.

Fig S7. Effect of estradiol on the growth of the wild type and MpNEK1-Citrine inducible lines

The gammae of the wild type (WT) and the MpNEK1-Citrine inducible lines were planted in the

agar medium supplemented with (+ER) or without 1 µM estradiol (Mock) and grown for 7, 14, and

21 days.

Video S1. Localization of MpNEK1-Citrine in a gemmaling grown for 4 days in the medium

supplemented with 1µM estradiol (same plant shown in Fig. 6B right panels).

Video S2. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine

inducible line treated with estradiol for 24 hours.

MpNEK1-Citrine inducible line was grown without estradiol for 3 days and then applied with 1 µM

estradiol and grown for 24 hours. Green: MpNEK1-Citrine, magenta: plastid autofluorescence

(same rhizoid shown in Fig. 7C).

Video S3. Localization and dynamics of over-accumulated MpNEK1-Citine in a rhizoid of

MpNEK1-Citrine inducible line treated with estradiol for 24 hours.

MpNEK1-Citrine inducible line was grown without estradiol for 3 days and then applied with 1 μM

estradiol and grown for 24 hours. A rhizoid highly accumulating MpNEK1-Citrine was observed.

Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

Video S4. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine

inducible line treated with estradiol for 48 hours.

MpNEK1-Citrine inducible line was grown without estradiol for 3 days and then applied with 1 μM

estradiol and grown for 48 hours. Green: MpNEK1-Citrine, magenta: plastid autofluorescence

(same rhizoid shown in Fig. 7C).

Video S5. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine

inducible line treated with estradiol for 48 hours.

MpNEK1-Citrine inducible line was grown without estradiol for 3 days and then applied with 1 μM

estradiol and grown for 48 hours. Green: MpNEK1-Citrine, magenta: plastid autofluorescence

(same rhizoid shown in Fig. 8A).

Video S6. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine

inducible line treated with estradiol for 120 hours.

MpNEK1-Citrine inducible line was grown without estradiol for 3 days and then applied with 1 μM

estradiol and grown for 120 hours. Green: MpNEK1-Citrine, magenta: plastid autofluorescence

(same rhizoid shown in Fig. 7C).

Video S7. Localization of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine inducible line treated

with estradiol for 120 hours.

MpNEK1-Citrine inducible line was grown without estradiol for 3 days and then applied with 1 μM

estradiol and grown for 120 hours. Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

Fig1. Effect of estradiol on thallus growth of Mp*NEK1* inducible line.

(A) Construct for Mp*NEK1* induction. Mp*EF1α* promoter, terminator (T), LexA operator (LexA op), minimal CaMV35S promoter (m35Spro). The diagram is not shown to scale.

(B) The gemmae of Mp*NEK1* inducible line (#2) were planted in the agar medium with (+ER) or without 10 μM estradiol (Mock) and grown for 3, 11, and 20 days.

(C) Time course of growth of Mp*NEK1* inducible line (#2) with (+ER) or without 10 μM estradiol (Mock) .

(D) Quantification of thallus growth of Mp*NEK1* inducible line (#2, 4, 8) with (+ER) or without 10 μM estradiol (Mock). The mean projection area of thalli (n = 10 plants) was quantified by ImageJ. Circles and error bars indicate mean values and standard deviations, respectively.

Fig 2. Reversible or irreversible effect of estradiol on thallus growth

(A) Schematic diagram of treatments. Gemmae were planted in the agar medium with 10 μM estradiol, grown in the same medium for 1, 3, 7, or14 days, and then transferred to and grown in the estradiol-free medium for 23 days.

(B) Gemmae of the wild type (WT) and four independent inducible lines (#2, 3, 4, and 8) were grown as shown in (A). The periods of estradiol treatment is shown above the photographs taken at the 23 day after transfer to the estradiol-free medium.

Fig 3. Effect of various concentrations of estradiol on thallus growth.

The gemmae of the wild type and Mp*NEK1* inducible lines were planted in the agar medium supplemented with or without estradiol at the concentration of 10 nM, 100 nM, and 1000 nM and grown for 13 (A) and 16 days (B).

Fig. 4. Estradiol resistance and its transfer to the next generation.

(A) The gemmae of the estradiol resistant plants or sensitive plants of Mp*NEK1* inducible lines were planted in the agar medium supplemented with 10 μM estradiol and grown for 11, 15, and 21 days.

(B) Quantification of growth of gemmalings derived from the estradiol resistant plants or sensitive plants of Mp*NEK1* inducible lines. The mean projection area of thalli (n = 10 plants) was quantified by ImageJ. Circles and error bars indicate mean values and standard deviations, respectively.

Fig. 5. Effect of estradiol on cell proliferation

(A) EdU-labelled nuclei in the gemmalings treated with 1 μM estradiol for 4 days. The gemmae of the wild type (WT) and MpNEK1 inducible line were planted in the medium supplemented with or without 1 μM estradiol, grown for 4 days, and incubated with 10 μM EdU for 1 hour in liquid medium of the same kinds. EdU-labelled nuclei were visualized according to the manufacture's instruction as described in methods.

(B) EdU-labelled nuclei in the gemmalings treated with 1 μM estradiol for 1 day. The gemmae of the wild type (WT) and MpNEK1 inducible line were planted in the medium without estradiol, grown for 3 days, and then transferred to and grown in the medium supplemented with or without 1 μM estradiol for 1 day. The gemmalings were incubated with 10 μM EdU for 1 hour in liquid medium of the same kinds. EdU-labelled nuclei were visualized according to the manufacture's instruction as described in methods.

(C) Quantification of EdU-labeled nuclei in the thalli grown as in (A). Data are shown by box plots (n = 12-16 plants). An asterisk indicates significant difference from the control (-ER) (*t*-test, P < 0.05).

(D) Quantification of EdU-labeled nuclei in the thalli grown as in (B). Data are shown by box plots (n = 12-16 plants). An asterisk indicates significant difference from the control (-ER) (*t*-test, P < 0.05).

Fig. 6. Effect of MpNEK1 induction on microtubules

(A) The gemmae of MpNEK1 inducible line with a microtubule marker CaMV35S:Citrine-MpTUB2 were grown in the estradiol-free medium for 3 days and then applied with the liquid medium with (+ER) or without 10 μM estradiol (Mock). Thalli were observed under a confocal microscope after 24 h and 48 h of treatment. Green: Microtubule, magenta: plastid autofluorescence.

(B) Number of mitotic cells per notch in the MpNEK1 inducible line with CaMV35S:Citrine-MpTUB2 after 24 h and 48 h of treatment with or without estradiol. Data are shown in the box plot (n = 10 notches, N > 6 plants). Asterisks indicate significant difference (t-test, P < 0.01).

(C) The classification of mitotic cells with prospindle, spindle, or phragmoplast (n indicates the number of mitotic cells used for the classification).

A

B

Mock	+ER (4 days)

C

Mock	+ER (1 day)

Fig. 7. Localization of estradiol-induced MpNEK1-Citrine in the thallus

(A) Construct for Mp*NEK1* induction. Mp*EF1α* promoter, terminator (T), LexA operator (LexA op), minimal CaMV35S promoter (m35Spro). The diagram is not shown to scale.

(B) The gemmae of the Mp*NEK1* inducible line were planted in the medium with (+ER) or without 1 μM estradiol (Mock), grown for 4 days, and observed under a confocal microscope.

(C) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol, grown for 2 days, and then added with (+ER) or without 1 μM estradiol (Mock) for 1 day. Upper panels are light field images merged with confocal images shown in the below panels. Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

Fig. 8. Localization of estradiol-induced MpNEK1-Citrine in rhizoids.

(A) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol, grown for 3 days, and then applied with (+ER) or without 1 μM estradiol (Mock). Rhizoids were observed under a confocal microscope. Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

(B, C) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol, grown for 3 days, and then applied with 1 μM estradiol and observed at the time indicated in each panel. The same rhizoid was observed under a confocal microscope in the time course. Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

(D) Velocity of rhizoid growth in the MpNEK1-Citrine inducible line incubated for 48 hours with (+ER) or without 1 μM estradiol (-ER). Data are shown by box plots (n = 32 rhizoids). An asterisk indicates significant difference (*t*-test, P < 0.01).

Fig. 9. Movement of MpNEK1-Citrine particles in rhizoids.

(A) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol, grown for 3 days, and then applied with 1 μM estradiol. Rhizoids were observed under a confocal microscope after 48 hours. The lower panel indicates kymograph of MpNEK1-Citrine particles. Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

(B) Velocity of MpNEK1-Citrine particles. Data are shown by box plots (n = 24 or 31 particles). There was no significant difference (*t*-test, P > 0.05).

Fig. 10. Effect of kinase-deficient MpNEK1 induction on the growth of thalli and rhizoids.
(A) Full length and kinase-deficient MpNEK1 used for estradiol induction.
(B) The gemmae of the wild type and inducible lines of full length MpNEK1, MpNEK1^{K37E}, or MpNEK1^{dKD} (KD deletion) were planted in the medium with (+ER) or without 1 μM estradiol (Mock), and grown for 14 days.
(C) The gemmae of the wild type and inducible lines of full length MpNEK1, MpNEK1^{K37E}, or MpNEK1^{dKD} (KD deletion) were planted in the medium with (+ER) or without 1 μM estradiol (Mock), and grown for 7 days.

Fig. 11. Effect of MpNEK1 induction on the number of rhizoids.

(A) The number of rhizoids in the gemmae grown for 7 days in the medium with (+ER) or without 1 μM estradiol (Mock). The gemmae of the wild type (WT) and inducible lines of full length MpNEK1-Citrine (FL), MpNEK1^{K37E}-Citrine (K37E), and MpNEK1^{dKD}-Citrine (dKD) were used. Data are shown by box plots (n = 10 plants). The different letters indicate significant differences by Tukey's HSD test (P < 0.03). Asterisks indicate significant differences from the value of FL with estradiol (+ER) (*t*-test, P < 0.01).

(B) The relative number of rhizoids in (A). The average number of rhizoids in estradiol-free media (ER -) in each genotype was used for the normalization. Data are shown by box plots (n = 10 plants). The different letters indicate significant differences by Tukey's HSD test (P < 0.03). Asterisks indicate significant differences from the value of FL with estradiol (+ER) (*t*-test, P < 0.01).

(C) The number of rhizoids in the gemmae grown for 14 days in the medium supplemented with 1 μM estradiol. The gemmae of the inducible lines of full length MpNEK1-Citrine (FL), MpNEK1^{K37E}-Citrine (K37E), and MpNEK1^{dKD}-Citrine (dKD) were used. Data are shown by box plots (n indicates the number of plants). The different letters indicate significant differences by Tukey's HSD test (P < 0.03).

Fig. 12. Localization of kinase-deficient MpNEK1-Citrine

(A) The gemmae of the inducible lines of MpNEK1^{K37E}-Citrine (K37E) or MpNEK1^{dKD}-Citrine (KD deletion) were planted in the medium with (+ER) or without 1 μM estradiol (Mock), grown for 4 days, and observed under a confocal microscope. Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

(B) The gemmae of the inducible lines of MpNEK1^{K37E}-Citrine (K37E) or MpNEK1^{dKD}-Citrine (KD deletion) were grown as in (A) and rhizoids were observed under a confocal microscope. Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

(C) Ratio of rhizoids with tip-localized or whole localized MpNEK1^{dKD}-Citrine (KD deletion) or MpNEK1^{K37E}-Citrine (K37E) after 120 h of estradiol treatment.

(D) Growth velocity of rhizoids in the MpNEK1^{K37E}-Citrine inducible lines incubated for 48 hours with (+ER) or without 1 μM estradiol (-ER). Data are shown by box plots (n = 30 rhizoids). An asterisk indicates significant difference (*t*-test, P < 0.01).