Title

Overexpression of NIMA-related kinase suppresses cell proliferation and tip growth in a liverwort

Marchantia polymorpha

(119 letters with spaces)

Hikari Mase[1], Yoshihiro Yoshitake[2], Takayuki Kohchi[2], Taku Takahashi[1], Hiroyasu Motose[1]

[1]Department of Biological Science, Graduate School of Natural Science & Technology, Okayama

University, Tsushimanaka 3-1-1, Okayama 700-8530, Japan

[2]Graduate School of Biostudies, Kyoto University, Kyoto 606-8502, Japan

Hikari Mase, pxn56xng@s.okayama-u.ac.jp

Yoshihiro Yoshitake, yoshitake.yoshihiro.7v@kyoto-u.ac.jp

Takayuki Kohchi, tkohchi@lif.kyoto-u.ac.jp

Taku Takahashi, perfect@cc.okayama-u.ac.jp

Hiroyasu Motose, motose@cc.okayama-u.ac.jp

Date of submission: 25 Jan. 2023

Number of tables and figures: 8 figures

Supplementary data: 6 supplementary figures, 7 videos

Word count: 2813 (introduction to the start of the acknowledgements)

Running title: Overexpression of NIMA-related kinase in *M. polymorpha* (54 letters with spaces)

Highlight

Overexpression of NIMA-related kinase suppresses cell proliferation and tip growth in a liverwort *Marchantia polymorpha*. Mp*NEK1* expression might be tightly regulated to control cell division as fungi and animal NEKs.

(30 words)

Abstract

NIMA-related kinases (NEK) regulate various mitotic events in fungi and animals, whereas plant NEKs regulate growth direction of cells and organs. The liverwort *Marchantia polymorpha* has a single functional Mp*NEK1* gene, whose knockout leads to aberrant growth direction of rhizoids. Its gain-of-function phenotype, however, remains unknown. Here, we generated inducible transgenic lines of Mp*NEK1*, using an estrogen receptor mediated system. Estradiol treatment efficiently increased accumulation of Mp*NEK1* mRNA and overexpression at the protein level was confirmed in the inducible lines of MpNEK1-Citrine. Overexpression of MpNEK1 severely suppressed growth of rhizoids and thalli, eventually causing lethality of juvenile plants. The effect of estradiol was reversible until 3 days, whereas 7-days treatment resulted in irreversible suppression of growth. This severe growth suppression was observed even at the nanomolar level of estradiol. EdU staining clearly indicated elimination of cell proliferation by estradiol-induced MpNEK1. In conclusion, overexpression of MpNEK1 suppresses cell division and elongation, leading to severe growth defects and lethality. Our results imply that the expression of MpNEK1 is tightly regulated at the appropriate level for cell growth. Plant NEK might control cell division as in the case of fungi and animal NEKs.

(190 words)

Keywords

Estradiol, *Marchantia polymorpha*, NIMA-related kinase, rhizoid, tip growth, thallus

Abbreviations

EdU; 5-ethynyl-2-deoxyuridine, EF1α; elongation factor1α, NEK; NIMA-related kinase, NIMA;

never in mitosis A, WDL, WAVE DANPENED-LIKE, XVE: LexA-VP16-ER artificial transcription

activator, WT; wild type

Materials and Methods

Plant materials and growth conditions

M. polymorpha accessions Takaragaike-1 (Tak-1, male) and Takaragaike-2 (Tak-2, female) were used as the wild type strains. Plants were grown on the half-strength Gamborg's B5 medium solidified with 1% agar at 22°C under the light cycles with 16 hours of light and 8 hours of darkness.

Plasmid construction and transformation

Plasmids were mainly constructed based on the Gateway technology (Life technologies). Primers used in the plasmid construction are listed in Table S1. To generate a plasmid of Mp*NEK1* induction by estradiol, the full-length MpNEK1 coding sequence in the pENTR/D-TOPO entry vector (Otani et al. 2018) was transferred by LR reaction (LR clonase □ enzyme mix, Life Technologies) into the Gateway binary vector pMpGWB344 (MpEF1α pro:XVE >>LexA operator:Gateway cassette).

To generate a plasmid of estradiol-inducible MpNEK1-mCitrine, *mCitrine* coding sequence was amplified by KOD One (Toyobo) and a primer set (Table S1) and was cloned into the *Asc*I site of the pENTR/D-TOPO entry vector containing full length MpNEK1 CDS (Otani et al., 2018) by In-fusion system (Takara). The resulting vector pENTR/D-TOPO-MpNEK1-mCitrine was subjected to LR reaction to transfer MpNEK1-Citrine fusion into the Gateway binary vector pMpGWB144 (MpEF1α pro:XVE >>LexA operator:Gateway cassette).

The constructs, pMpGWB344-Mp*NEK1* (MpEF1α pro:XVE>>LexA operator:MpNEK1) and pMpGWB144-MpNEK1-Citrine (MpEF1α pro:XVE>>LexA operator:MpNEK1-Citrine) were transformed into F1 sporelings derived from sexual crosses between Tak-2 and Tak-1 by *Agrobacterium*-mediated method (Ishizaki et al., 2008). Chlorsulfuron and hygromycin was used for the selection of transformants in pMpGWB344 and pMpGWB144, respectively. In pMpGWB344 and pMpGWB144, the artificial transcriptional activator XVE is expressed under the

control of Mp*EF1α* promoter. XVE translocates to the nucleus in the presence of estradiol and binds

to the LexA operator to induce the expression of MpNEK1 and MpNEK1-Citrine, respectively.

RT-qPCR

Total RNA was isolated from the thalli of the wild type and transgenic plants. For each sample, 0.5

µg of total RNA was reverse transcribed to cDNA using ReverTra Ace reverse transcriptase

(Toyobo) according to the manufacturer's protocol. Real-time PCR was performed on a thermal

cycler Dice Real Time System (Takara) using KOD SYBR qPCR Kit (Toyobo) according to the

manufacturer's protocol. Transcript levels of Mp*EF1α* and Mp*ACT* were used as a reference for

normalization (Saint-Marcoux et al., 2015). Primers used in RT-qPCR are listed in Table S1.

RT-qPCR experiments were performed using four biological replicates.

Microscopy

The morphology of gemmalings and rhizoids was observed under a stereo microscope S8APO0

(Leica Microsystems) or a light microscope DM5000B (Leica) equipped with a CCD camera

DFC500 (Leica). To analyze the localization of MpNEK1-Citrine, plants were grown on

half-strength B5 medium for 3 and 5 days under the cycles with 16 hours of light and 8 hours of

darkness. For live imaging, half-strength B5 agar medium was solidified in the glass bottom dish

(35 mm in diameter x 10 mm in thickness, Thickness No. 1, Matsunami) and then the central region

of agar medium on the bottom glass strip was removed by tweezers. This region was poured with

about 200 µL of melted half-strength B5 agar medium to generate a thin solidified medium. The

gemmae were planted and grown in the central thin medium for 2-7 days. The details of this method

will be described elsewhere by H.M. Gemmalings were observed under a FV1200 confocal

laser-scanning microscope (Olympus) equipped with a high-sensitivity GaAsP detector and silicone

oil objective lenses (30×, 60× Olympus). Silicone oil (SIL300CS, Olympus) was used as immersion

media for the objective lenses. The images were analyzed using ImageJ (National Institutes of

Health, USA). For estradiol treatment, an aliquot (200□300 µL) of estradiol solution at a

concentration of 1µM was added to the gemmalings for the induction of Mp*NEK1*. As a control,

same amount of water supplemented with same concentration of solvent (DMSO, less than 0.1%)

was applied.

EdU staining

S-phase cells were visualized using Click-iT EdU Imaging Kits (Life Technologies) according to

the manufacturer's instruction. Gemmae were incubated on the half-strength Gamborg's B5 agar

medium supplemented with or without 1 µM β-estradiol for 4 days. Alternatively, for 1-day

treatment, gemmae were incubated on the agar medium without 1 µM β-estradiol for 3 days and

then transferred to and incubated in the agar medium with 1 µM β-estradiol for 1 day. Gemmalings

were then transferred to a solution containing 10 µM 5-ethynyl-2-deoxyuridine (EdU) and

incubated for 1 hour. Plants were fixed with 3.7 % formaldehyde solution in phosphate-buffered

saline (PBS) for 20 min. EdU incorporated into DNA was labelled with Alexa Fluor

488-azide-containing Click-iT reaction cocktail in the dark for 30 min. EdU-labelled cells were

observed using a confocal laser scanning microscope, FV1200 (Olympus). The maximum

Z-projection images were created using the ImageJ software (Schneider et al. 2012).

Introduction

Never in mitosis A (NIMA)-related kinase (NEK) is a Ser/Thr protein kinase that controls various mitotic events in fungi and animals (O'Connell et al. 2003, Fry et al. 2012), while plant NEK members have been shown to regulate directional growth (Motose, et al. 2008; Sakai, et al. 2008). *Arabidopsis* NEK6 suppresses ectopic outgrowth of epidermal cells through microtubule organization and interaction with other NEK members (Motose, et al. 2011). NEK6 localizes to the shrinking ends of microtubules, phosphorylates five amino acid residues of β-tubulin, and depolymerizes aberrant cortical microtubules to promote epidermal cell elongation (Takatani, et al. 2017). Furthermore, NEK6 has been shown to stabilize growth direction of hypocotyls through selectively removing cortical microtubules that best align with maximal tensile stress (Takatani, et al. 2020). NEK6-mediated attenuation of tensile-stress response increases the noise in microtubule orientation to buffer mechanical conflicts and to stabilize growth direction.

The liverwort *Marchantia polymorpha* is an emerging model to study evolution of land plants (Kohchi, et al. 2021) because of its low genetic redundancy (Bowman, et al. 2017) and various genetic tools (Ishizaki, et al. 2015; Ishizaki, et al. 2016). Indeed, *Arabidopsis* has seven NEK members, whereas *M. polymorpha* has a single *NEK* gene, Mp*NEK1* (Takatani, et al. 2015; Otani, et al. 2018). Thanks to these advantages, MpNEK1 has been shown to direct tip growth of rhizoids through microtubule depolymerization in the apical region (Otani, et al. 2018). Rhizoids are filamentous rooting cells, mainly elongating from the ventral epidermis and are thought to be required for plant anchorage to the soil and water/nutrient uptake (Jones and Dolan 2012; Shimamura, 2016; Kohchi, et al. 2021; Kanda, et al. 2022). Mp*nek1* mutant rhizoids frequently change their growth direction, resulting in the zigzag morphology (Otani, et al. 2018) and the defect of invasive growth to enter the hard substratum (Mase, et al. 2022). MpNEK1 localization in the apical microtubule foci and the increased microtubule stability in Mp*nek1* mutants demonstrate that MpNEK1 reorganizes and destabilizes apical microtubules to maintain growth polarity of rhizoids

(Otani, et al. 2018). Similar phenotype was observed in the mutants of a microtubule-associated

protein, WAVE DANPENED-LIKE (MpWDL) (Honkanen, et al. 2016; Champion, et al. 2021).

MpWDL might stabilize longitudinal microtubules in the shank region of rhizoids. Thus,

microtubule organization and dynamics are essential for directional growth of rhizoids.

However, it is not clear how MpNEK1 regulates microtubules to stabilize growth direction.

Furthermore, functional significance of Mp*NEK1* expression in the meristem remains unclear. The

gain of function approach could be helpful to elucidate these problems and to identify downstream

proteins that are phosphorylated by MpNEK1. Here, we adopted estradiol-inducible system to

overdrive Mp*NEK1* and to estimate its effects on growth and development in *M. polymorpha*.

Results

Establishment of MpNEK1 inducible lines

In this study, we generated Mp*NEK1* inducible transgenic lines employing the XVE chimeric

transcriptional activator (Zuo, et al. 2000). This activator is composed of the DNA-binding domain

of LexA, the transcriptional activation domain of VP16, and the carboxyl estradiol binding domain

of the human estrogen receptor. This system has no apparent toxic physiological effects and has

been effectively used in flowering plants. The XVE vector was optimized for the application to *M.*

polymorpha in several points. Because the endogenous elongation factor Mp*EF1α* promoter is more

active in the meristem and sexual organ than the cauliflower mosaic virus *35S* promoter (Althoff, et

al. 2014), the Mp*EF1α* promoter was utilized to drive XVE. However, only a few research has been

reported on the application in *M. polymorpha* (e.g. Flores-Sandoval, et al. 2016; Furuya, et al.

2022).

We isolated eight transgenic lines by transformation of wild type sporelings with

pMpGWB344-MpNEK1 (MpEF1α pro:XVE>>LexAop:MpNEK1), in which XVE is expressed

under the control of Mp*EF1α* promoter and activates transcription of Mp*NEK1* through the binding

to the LexA operator in the presence of estradiol. Among them, seven transgenic lines (line #2 to 8)

were most identical to the wild type in growth and morphology of thalli. These transgenic lines

were examined for the efficiency of Mp*NEK1* induction by quantitative real-time PCR (RT-qPCR)

(Fig. S1). Two-week-old thalli of the wild type (WT) and transgenic lines were transferred to the

medium with or without β-estradiol and incubated for 24 hours. The expression levels of Mp*NEK1*

were remarkably increased in most transgenic lines (#2, 4, 5, 7, and 8). Mp*NEK1* expression was

100 to 200 times higher in estradiol treatment than in the estradiol-free medium. Estradiol did not

induce Mp*NEK1* expression in the wild type and line #6. Mp*NEK1* induction was well reproducible

and reliable, especially in line #2, 4, and 8, which exhibit no obvious defects in growth and

morphology without estradiol. Thus, these were used for the further analyses described below.

MpNEK1 induction severely suppresses thallus growth

We conducted phenotypic analysis to access the effects of estradiol-induced MpNEK1 on growth

and development. The gemmae of wild type and Mp*NEK1* inducible lines were grown in the

presence or absence of exogenous β-estradiol for 24 days. There was no obvious effect of

β-estradiol on the growth and morphology of the wild type thalli (Fig. S2). In the inducible lines, on

the other hand, thallus growth was severely suppressed when cultured with β-estradiol (Fig. 1).

Most gemmalings stopped growing and died in the presence of estradiol (Fig. 1), whereas line #6

did not show growth retardation (Fig. S3). It was noteworthy that Mp*NEK1* was not induced in the

line #6 (Fig. S1). These results clearly demonstrate that induction of Mp*NEK1* severely suppresses

thallus growth and causes juvenile lethality.

We next examined whether effect of Mp*NEK1* induction is reversible or not. The gemmalings of

the wild type and Mp*NEK1* inducible lines were grown in the presence of estradiol, and then

transferred to the estradiol-free medium at various times after estradiol treatment (Fig. 2). After 1

and 3-day treatment with estradiol, thallus growth seemed to be recovered, whereas irreversible

growth suppression was observed after 7 and 14-day treatment. In the wild type, growth inhibitory

effect of estradiol was not observed. Therefore, one-week MpNEK1 induction (estradiol treatment)

confers irreversible effect on thallus growth.

In the XVE induction system in flowering plants, 2 to 10 µM concentrations of estradiol have

been commonly used. In the Mp*NEK1* inducible lines, 10 µM estradiol caused severe growth

suppression and lethality. However, more mild conditions would be desirable to further investigate

MpNEK1 function. To determine less-effective conditions and dose-dependent effect of estradiol,

thalli of the wild type and Mp*NEK1* inducible lines were incubated in the presence of estradiol at

various concentrations (Fig. 3). The wild-type thallus showed no obvious phenotype in all

concentration ranges, whereas severe growth suppression was observed in the Mp*NEK1* inducible

lines even at the 10 and 100 nM of estradiol. From this result, Mp*NEK1* induction has the potent

effect on thallus growth.

We noticed that some plants obtained the resistance to estradiol and survived in the agar medium

supplemented with estradiol (Fig. 4). The estrogen resistance was observed in the next generation

gemmae, suggesting cell-cell movement of siRNA from the thallus to gemma. These

estradiol-resistant plants appeared more frequently in the lower concentrations and exhibited no

induction of Mp*NEK1* (Fig. S4). At the lower concentrations, larger numbers of thallus cells could

survive and acquire estradiol resistance by gene silencing mechanism. Thus, lower concentrations

of estradiol are not desirable conditions, which often induce resistant plants.

MpNEK1 induction suppresses cell proliferation

Previous studies show that plant NEKs mainly regulate growth direction of cells and organs.

However, severe suppression of thallus growth by estradiol-induced Mp*NEK1* suggests some roles

of plant NEK in cell division and proliferation. To determine whether estradiol-induced Mp*NEK1*

affects cell division, we monitored cell proliferation activity by staining with EdU

(5-ethynil-2'-deoxyuridine) (Fig. 5). This thymidine analog can be incorporated into DNA during S

phase. The subsequent reaction of EdU with a fluorescent azide labels newly replicated DNA

strands of the cells in S phase during the incubation with EdU.

In both the wild type and inducible lines, EdU-positive nuclei were observed in the meristematic

notch region in thalli grown in the absence of estradiol (Fig. 5). The amount of EdU-positive nuclei

was almost similar in the wild type with or without estradiol, whereas it was significantly reduced

in the estradiol-treated inducible lines. This result demonstrates that estradiol-induced Mp*NEK1*

suppresses cell proliferation.

MpNEK1-Citrine overaccumulation suppresses growth of thalli and rhizoids

We generated inducible lines of MpNEK1-Citrine fusion protein, in which protein induction and

localization could be easily detected by confocal microscopy, facilitating functional analysis of

MpNEK1. Eight transgenic lines were isolated by transformation of wild type sporelings with

pMpGWB144-MpNEK1-Citrine. Among them, four transgenic lines (line #1, 5, 7, and 8) were

most identical to the wild type in the absence of estradiol. When grown in the agar medium

supplemented with estradiol, these transgenic lines exhibited severe growth retardation and juvenile

lethality (Fig. S5).

We next analyzed expression and localization of MpNEK1-Citrine fusion protein (Fig. 6A, B).

The gemmae of MpNEK1-Citrine inducible lines were grown for 4 days in the medium with or

without 1 µM β-estradiol and observed under a confocal microscope. MpNEK1-Citrine was not

detected in the absence of estradiol, whereas it was accumulated throughout estradiol-treated thalli

(Fig. 6A, Video S1). MpNEK1-Citrine localized in the cytoplasmic particles. In the 24-hour

estradiol treatment (2-day old gemmalings grown without estradiol were treated with estradiol

solution for 24 hours), similar expression and localization pattern were observed (Fig. 6B). From

these results, MpNEK1-Citrine was induced throughout thalli within 24 hours after estradiol

treatment and severely suppressed thallus growth.

We noticed accumulation of MpNEK1-Citrine particles in the meristem in some transgenic plants

even in the absence of estradiol (Fig. S6). This leaky expression levels were varied in the

individuals and transgenic lines. In this study, we used plants and transgenic lines (line #1, 5, 7, and

8), which showed less leaky expression of MpNEK1-Citrine.

Taking accounts that MpNEK1 localizes in the apical microtubule foci to direct tip growth of

rhizoids (Otani, et al. 2018), we analyzed effect and localization of estradiol-induced

MpNEK1-Citrine in tip-growing rhizoids (Fig. 7). Because rhizoids were not formed when gemmae

were treated with estradiol at the start of culture, we added estradiol solution on the 3-day-old

gemmalings grown without estradiol to examine localization and effect of MpNEK1-Citrine in the

time course. After 24-hour treatment, MpNEK1-Citrine accumulated in the apical region of rhizoids

(Fig. 7A, 7B, Video S2). The expression level of MpNEK1-Citrine was varied in each rhizoid. A

few rhizoids showed highly accumulated MpNEK1-Citrine, which formed large droplets, moved to

the apical direction, and fused with each other (Video S3). To determine when MpNEK1-Citrine

starts to accumulate, we observed rhizoids at the early periods of estradiol treatment (Fig. 7B).

MpNEK1-Citrine began to accumulate at about 1 hour of treatment and gradually increased in the

rhizoid apex. After 48 and 120 hours, MpNEK1-Citrine accumulated throughout the entire region of

rhizoids in the particle and filamentous patterns (Fig. 7C, Video S4-S7). Tip growth was ceased or

severely reduced after 48 and 120 hours of estradiol treatment (Fig. 7D, Video S4-6). After 48 hours,

movement of MpNEK1-Citrine particles was obvious (Fig. 8, Video S4, S5). MpNEK1-Citrine

particles moved toward the apex (anterograde) and basal region (retrograde) at the same velocity

(Fig. 8). These results suggest that overaccumulation of MpNEK1 suppresses rhizoid growth.

Discussion

Here, we established and characterized induction system of MpNEK1 as a new tool to analyze

plant NEK function. Estradiol-based XVE induction system has been widely used in angiosperms,

but its application and characterization in *M. polymorpha* still remain limited. The efficient

induction of Mp*NEK1* transcripts and overaccumulation of MpNEK1-Citrine fusion protein

throughout plant body indicate that this system is very helpful for the functional analysis of

MpNEK1.

We noticed several unexpected phenomena about which caution is required. MpNEK1-Citrine

showed leaky expression in the meristem without estradiol. This might be due to the artificial

transcription activator XVE under the control of Mp*EF1α* promoter, that is highly active in

meristematic tissues (Althoff, et al. 2014). Nonetheless, this leaky expression did not affect thallus

growth, suggesting that effective threshold of MpNEK1 could exist. Since the leaky expression was

varied in plants and transgenic lines, estrogen-free XVE-mediated transactivation might be

stochastic and/or responsive to fluctuating physiological states.

Secondly, estrogen-resistant silenced plants emerged as survivors in the prolonged treatment

especially at the lower concentrations of estradiol. The estrogen resistance could be transferred to

the next generation gemmae, implying thallus-to-gemma movement of siRNA. Thus, it would be

useful for the analysis of RNA silencing in *M. polymorpha*, while invokes concerns about treatment

time and estradiol concentration to minimize the contamination of silenced plants.

MpNEK1 induction severely suppressed thallus growth probably through cell proliferation. There

are several possible mechanisms of this growth-suppressive effect. Firstly, MpNEK1 may

depolymerize microtubule-based mitotic apparatus including preprophase band, spindle and

phragmoplast because MpNEK1 regulates tip growth of rhizoids through microtubule

depolymerization (Otani, et al. 2018). Secondly, overexpression of MpNEK1 may activate cell

cycle checkpoints to prevent cell cycle progression. The *nimA* mutants of a filamentous fungus

Aspergillus nidulans exhibit cell cycle arrest just before M phase, and conversely, overexpression of

NimA causes premature chromosome condensation and aberrant spindle formation, resulting in cell

cycle arrest (Oakley and Morris, 1983; Osmani, et al. 1988). Thirdly, MpNEK1 may affect cortical

microtubule organization, which is required for directional growth prior to cell division.

Arabidopsis NEK6 depolymerize specific microtubules, detached from plasma membrane or

aligned to the maximal tensile stress, to stabilize growth direction (Takatani, et al. 2017; Takatani, et

al. 2020). Plant cells need to elongate to some extent prior to cell division, whereas if microtubules

are depolymerized by MpNEK1, cell elongation could be suppressed, resulting in suppression of

cell division as well. However, we could not succeed to introduce microtubule markers in the

MpNEK1 inducible lines and to observe the effects of overexpression on microtubules. Taking into

accounts of remarkable promoter activity of MpNEK1 in the meristem (Otani, et al. 2018), it would

be likely that MpNEK1 regulates cell division in addition of rhizoid growth.

Overexpression of MpNEK1 suppressed tip growth in rhizoids. Furthermore, overexpression

altered localization of MpNEK1-Citrine from the rhizoid apex to the entire region as the time of

estradiol treatment increased. This may reflect the overflow of MpNEK1 from the transport and

localization machineries. As the consequence, tip growth was stopped in rhizoids with this

dispersed localization pattern. Over-accumulated MpNEK1 may destabilize microtubule foci at the

apex and/or longitudinal microtubules at the shank region. Because the longitudinal microtubules

might promote rhizoid elongation through organelle transport (Kanda, et al. 2022), it is plausible

that MpNEK1 depolymerizes longitudinal microtubules to suppress rhizoid growth.

In summary, MpNEK1 induction system is useful for the analyses of its function in cell division

and directional growth. Furthermore, it would be helpful to visualize protein transport toward the

rhizoid apex and shank region because we observed MpNEK1 particles moved to the acropetal and

basipetal direction (Video S3-S5). The velocity of MpNEK1 particles is quite similar to that of

armadillo-repeat kinesin, about half of velocity of growing microtubule plus ends (Kanda et al.

2022), suggesting this motility is not due to plus-end tracking but rather to the kinesin-dependent

transport along microtubules. Anyway, MpNEK1 overexpression has the potent growth effect than

that of *Arabidopsis* NEK6, whose overexpression caused slight reduction of root growth (Takatani,

et al. 2017). Comparing with MpNEK1, NEK6-Citrine fusion protein is not highly accumulated in

the XVE system (HMo, unpublished result), implying the differential degradation efficiency of

NEK proteins and the advantage of protein induction in *M. polymorpha*.

Supplementary data

Fig S1. Effects of estradiol on the accumulation of Mp*NEK1* transcripts in the wild type and transgenic lines introduced with *XVE::*Mp*NEK1*.

Fig S2. Effect of estradiol on thallus growth of the wild type.

Fig S3. Effect of estradiol on thallus growth of the transgenic line #6 with *XVE::*Mp*NEK1*.

Fig S4. Effect of estradiol on the accumulation of Mp*NEK1* transcripts in the wild type and estradiol-resistant transgenic plants harboring *XVE::*Mp*NEK1*.

Fig S5. Effect of estradiol on the growth of MpNEK1-Citrine inducible lines

Fig S6. Expression and localization of MpNEK1-Citrine in the meristem of thallus without estradiol.

Video S1. Localization of MpNEK1-Citrine in a gemmaling grown for 4 days in the medium supplemented with 1μM estradiol (same plant shown in Fig. 6B right panels).

Video S2. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine inducible line treated with estradiol for 24 hours.

Video S3. Localization and dynamics of over-accumulated MpNEK1-Citine in a rhizoid of MpNEK1-Citrine inducible line treated with estradiol for 24 hours.

Video S4. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine inducible line treated with estradiol for 48 hours.

Video S5. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine inducible line treated with estradiol for 48 hours.

Video S6. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine inducible line treated with estradiol for 120 hours.

Video S7. Localization of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine inducible line treated with estradiol for 120 hours.

Acknowledgements

We thank Drs. Ryuichi Nishihama and Kimitsune Ishizaki for their advice on EdU staining and

transformation.

Author contributions

HMa and HMo designed the research; HMa performed experiments and analyzed the data; YY and

TK developed XVE system; HMa and HMo wrote the manuscript; HMa, YY, TK, TT and HMo

revised the manuscript.

Conflict of interest

The authors declare that they have no conflicts of interest.

Funding

This work was supported by Grants-in-Aid from the JSPS (JSPS KAKENHI Grant Numbers

16K07403, 19K06709, and 21H00370 to H.M.), Ryobi Teien Memory Foundation, and Naito

Foundation to H.M.

Data availability

The data of this study are available upon request from the corresponding author, Hiroyasu Motose.

References

Althoff F, Kopischke S, Zobell O, Ide K, Ishizaki K, Kohchi T, Zachgo S. 2014. Comparison of the

Mp*EF1α* and *CaMV35* promoters for application in *Marchantia polymorpha* overexpression studies.

Transgenic Research 23, 235-244.

Bowman JL, Kohchi T, Yamato KT, et al. 2017. Insights into land plant evolution garnered from the

Marchantia polymorpha genome. Cell 171: 287-304.

Champion C, Lamers J, Jones VAS, Morieri G, Honkanen S, Dolan L. 2021. Microtubule associated

protein WAVE DAMPENED2-LIKE (WDL) controls microtubule bundling and the stability of the

site of tip-growth in *Marchantia polymorpha* rhizoids. PLoS Genetics 17, e1009533.

Flores-Sandoval E, Dierschke T, Fisher TJ, Bowman JL. 2016. Efficient and inducible use of

artificial microRNAs in *Marchantia polymorpha*. Plant & Cell Physiolgy 57, 281-290.

Fry AM, O'Regan L, Sabir SR, Bayliss R. 2012. Cell cycle regulation by the NEK family of protein

kinases. Journal of Cell Science 125, 4423-4433.

Furuya T, Nishihama R, Ishizaki K, Kohchi T, Fukuda H, Kondo Y. 2022. A glycogen synthase

kinase 3-like kinase MpGSK regulates cell differentiation in *Marchantia polymorpha*. Plant

Biotechnology 39, 65-72.

Honkanen S, Jones VAS, Morieri G, Champion C, Hetherington AJ, Kelly S, Proust H,

Saint-Marcoux D, Prescott H, Dolan L. 2016. The mechanism forming the cell surface of

tip-growing rooting cells is conserved among land plants. Current Biology 26, 3238–3244.

Ishizaki K, Chiyoda S, Yamato KT, Kohchi T. 2008. *Agrobacterium*-mediated transformation of the

haploid liverwort *Marchantia polymorpha* L., an emerging model for plant biology. Plant & Cell

Physiology 49, 1084-1091.

Ishizaki K, Nishihama R, Ueda M, Inoue K, Ishida S, Nishimura Y, Shikanai T, Kohchi T. 2015.

Development of gateway binary vector series with four different selection markers for the liverwort

Marchantia polymorpha. PLoS One 10, e0138876.

Ishizaki K, Nishihama R, Yamato KT, Kohchi T. 2016. Molecular genetic tools and techniques for *Marchantia polymorpha* research. Plant & Cell Physiology 57, 262-270.

Jones VAS, Dolan L. 2012. The evolution of root hairs and rhizoids. Annals of Botany 110, 205-212.

Kanda A, Otani K, Takahashi T, Motose H. 2022. Plant specific armadillo repeat kinesin directs organelle transport and microtubule convergence to promote tip growth. BioRxiv doi: https://doi.org/10.1101/2022.07.08.499237. [Preprint]

Kohchi T, Yamato KT, Ishizaki K, Yamaoka S, Nishihama R. 2021. Development and molecular genetics of *Marchantia polymorpha*. Annual Review of Plant Biology 72, 677-702.

Mase H, Nakagami H, Okamoto T, Takahashi T, Motose H. 2022. Establishment and application of novel culture methods in *Marchantia polymorpha*: persistent tip growth is required for substrate penetration by rhizoids. Communicative & Integrative Biology 15, 164-167.

Motose H, Tominaga R, Wada T, Sugiyama M, Watanabe Y. 2008. A NIMA-related protein kinase suppresses ectopic outgrowth of epidermal cells through its kinase activity and the association with microtubules. Plant Journal 54, 829-844.

Motose H, Hamada T, Yoshimoto K, Murata T, Hasebe M, Watanabe Y, Hashimoto T, Sakai T, Takahashi T. 2011. NIMA-related kinases 6, 4, and 5 interact with each other to regulate microtubule organization during epidermal cell expansion in *Arabidopsis thaliana*. Plant Journal 67,

993-1005.

Oakley BR, Morris NR. 1983. A mutation in *Aspergillus nidulans* that blocks the transition from

interphase to prophase. Journal of Cell Biology 96, 1155-1158.

O'Connell MJ, Krien MJ, Hunter T. 2003. Never say never. The NIMA-related protein kinases in

mitotic control. Trends in Cell Biology 13, 221-228.

Osmani SA, Pu RT, Morris NR. 1988. Mitotic induction and maintenance by overexpression of a

G2-specific gene that encodes a potential protein kinase. Cell 53, 237-244.

Otani K, Ishizaki K, Nishihama R, Takatani S, Kohchi T, Takahashi T, Motose H. 2018. An

evolutionarily conserved NIMA-related kinase directs rhizoid tip growth in the basal land plant

Marchantia polymorpha. Development 145, pii: dev154617.

Saint-Marcoux D, Proust H, Dolan L, Langdale JA. 2015. Identification of reference genes for

real-time quantitative PCR experiments in the liverwort *Marchantia polymorpha*. PLoS One 10,

e0118678.

Sakai T, Honing H, Nishioka M, et al. 2008. Armadillo repeat-containing kinesins and a

NIMA-related kinase are required for epidermal-cell morphogenesis in *Arabidopsis*. Plant Journal

53, 157-171.

Shimamura M. 2016. *Marchantia polymorpha*: taxonomy, phylogeny and morphology of a model

system. Plant & Cell Physiology 57, 230-256.

Takatani S, Otani K, Kanazawa M, Takahashi T, Motose H. 2015. Structure, function, and evolution

of plant NIMA-related kinases: implication for phosphorylation-dependent microtubule regulation.

Journal of Plant Research 128, 875-891.

Takatani S, Ozawa S, Yagi N, Hotta T, Hashimoto T, Takahashi Y, Takahashi T, Motose H. 2017.

Directional cell expansion requires NIMA-related kinase 6 (NEK6)-mediated cortical microtubule

destabilization. Scientific Reports 7, 7826.

Takatani S, Verger S, Okamoto T, Takahashi T, Hamant O, Motose H. 2020. Microtubule response

to tensile stress is curbed by NEK6 to buffer growth variation in the *Arabidopsis* hypocotyl. Current

Biology 30, 1491-1503.

Zuo J, Niu QW, Chua NH. 2000. An estrogen receptor-based transactivator XVE mediates highly

inducible gene expression in transgenic plants. Plant Journal 24, 265-273.

Figure legend

Fig1. Effect of estradiol on thallus growth of Mp*NEK1* inducible line.

(A) Construct for Mp*NEK1* induction. Mp*EF1a* promoter, terminator (T), LexA operator (LexA op),

minimal CaMV35S promoter (m35Spro). The diagram is not shown to scale.

(B) The gemmae of Mp*NEK1* inducible line (#2) were planted in the agar medium with (+ER) or

without 10 μM estradiol (Mock) and grown for 3, 11, and 20 days.

(C) Time course of growth of Mp*NEK1* inducible line (#2) with (+ER) or without 10 μM estradiol

(Mock) .

(D) Quantification of thallus growth of Mp*NEK1* inducible line (#2, 4, 8) with (+ER) or without 10

μM estradiol (Mock). The mean projection area of thalli (n = 10 plants) was quantified by

ImageJ.

Fig 2. Reversible or irreversible effect of estradiol on thallus growth

(A) Schematic diagram of treatments. Gemmae were planted in the agar medium with 10 μM

estradiol, grown in the same medium for 1, 3, 7, or14 days, and then transferred to and grown in

the estradiol-free medium for 23 days.

(B) Gemmae of the wild type (WT) and four independent inducible lines (#2, 3, 4, and 8) were

grown as shown in (A). The periods of estradiol treatment is shown above the photographs

taken at the 23 day after transfer to the estradiol-free medium.

Fig 3. Effect of various concentrations of estradiol on thallus growth.

The gemmae of the wild type and Mp*NEK1* inducible lines were planted in the agar medium

supplemented with or without estradiol at the concentration of 10 nM, 100 nM, and 1000 nM and

grown for 13 (A) and 16 days (B).

Fig. 4. Estradiol resistance and its transfer to the next generation.

(A)   The gemmae of the estradiol resistant plants or sensitive plants of Mp*NEK1* inducible lines

were planted in the agar medium supplemented with 10 μM estradiol and grown for 11, 15, and

21 days.

(B)   Quantification of growth of gemmalings derived from the estradiol resistant plants or sensitive

plants of Mp*NEK1* inducible lines. The mean projection area of thalli (n = 10 plants) was

quantified by ImageJ.

Fig. 5. Effect of estradiol on cell proliferation

(A) EdU-labelled nuclei in the gemmalings treated with 1 μM estradiol for 4 days. The gemmae of

the wild type (WT) and Mp*NEK1* inducible line were planted in the medium supplemented

with or without 1 μM estradiol, grown for 4 days, and incubated with 10 μM EdU for 1 hour in liquid medium of the same kinds. EdU-labelled nuclei were visualized according to the manufacture's instruction as described in methods.

(B) EdU-labelled nuclei in the gemmalings treated with 1 μM estradiol for 1 day. The gemmae of the wild type (WT) and Mp*NEK1* inducible line were planted in the medium without estradiol, grown for 3 days, and then transferred to and grown in the medium supplemented with or without 1 μM estradiol for 1 day. The gemmalings were incubated with 10 μM EdU for 1 hour in liquid medium of the same kinds. EdU-labelled nuclei were visualized according to the manufacture's instruction as described in methods.

(C) Quantification of EdU-labeled nuclei in (A). Data are shown by box plots (n = 12-16 plants). An asterisk indicates significant difference from the control (-ER) (*t*-test, P < 0.05).

(D) Quantification of EdU-labeled nuclei in (B). Data are shown by box plots (n = 12-16 plants). An asterisk indicates significant difference from the control (-ER) (*t*-test, P < 0.05).

Fig. 6. Localization of estradiol-induced MpNEK1-Citrine in the thallus

(A) Construct for Mp*NEK1* induction. Mp*EF1a* promoter, terminator (T), LexA operator (LexA op), minimal CaMV35S promoter (m35Spro). The diagram is not shown to scale.

(B) The gemmae of the Mp*NEK1* inducible line were planted in the medium with (+ER) or without 1 μM estradiol (Mock), grown for 4 days, and observed under a confocal microscope.

(C) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol, grown for 2 days, and then added with (+ER) or without 1 μM estradiol (Mock) for 1 day. Upper panels are light field images merged with confocal images shown in the below panels. Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

Fig. 7. Localization of estradiol-induced MpNEK1-Citrine in rhizoids.

(A) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol,

grown for 3 days, and then applied with (+ER) or without 1 µM estradiol (Mock). Rhizoids

were observed under a confocal microscope in the time course. Green: MpNEK1-Citrine,

magenta: plastid autofluorescence.

(B, C) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol,

grown for 3 days, and then applied with 1 µM estradiol and observed at the time indicated in

each panel. The same rhizoid was observed under a confocal microscope in the time course.

Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

(D) Velocity of rhizoid growth in the MpNEK1-Citrine inducible line incubated for 48 hours with

(+ER) or without 1 µM estradiol (-ER). Data are shown by box plots (n = 32 rhizoids). An

asterisk indicates significant difference (*t*-test, $P < 0.01$).

Fig. 8. Movement of MpNEK1-Citrine particles in rhizoids.

(A) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol,

grown for 3 days, and then applied with 1 µM estradiol. Rhizoids were observed under a

confocal microscope after 48 hours. The lower panel indicates kymograph of MpNEK1-Citrine

particles. Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

(B) Velocity of MpNEK1-Citrine particles. Data are shown by box plots (n = 24 or 31 particles).

There was no significant difference (*t*-test, $P > 0.05$).

Fig S1. Effects of estradiol on the accumulation of Mp*NEK1* transcripts in the wild type and

transgenic lines introduced with pMpGWB344-Mp*NEK1* (*XVE::*Mp*NEK1*).

The 10-day old plants of the wild type (WT) and the transgenic lines were incubated for 1 day in the

medium with (+) or without 10 µM estradiol (-). Total RNA was extracted and subjected to

RT-qPCR. All transcript levels are relative to that of mock-treated wild type plants. Mp*EF1α* and

Mp*ACT* were used as the control genes.

Fig S2. Effect of estradiol on thallus growth of the wild type.

(A) The gemmae of the wild type (Tak-1) were planted in the agar medium with (+ER) or without

10 µM estradiol (Mock) and grown for 3, 11, and 20 days.

(B) Time course of growth of the wild type with (+ER) or without 10 µM estradiol (Mock).

(C) Quantification of thallus growth of the wild type with (+ER) or without 10 µM estradiol

(Mock). The mean projection area of thalli (n = 10 plants) was quantified by ImageJ.

Fig S3. Effect of estradiol on thallus growth of the transgenic line #6 with *XVE::*Mp*NEK1*.

(A) The gemmae of the line (#6) were planted in the agar medium with (+ER) or without 10 µM

estradiol (Mock) and grown for 3, 11, and 20 days.

(B) Time course of growth of the line (#6) with (+ER) or without 10 µM estradiol (Mock) .

(C) Quantification of thallus growth of the line (#6) with (+ER) or without 10 µM estradiol (Mock).

The mean projection area of thalli (n = 10 plants) was quantified by ImageJ.

Fig S4. Effect of estradiol on the accumulation of Mp*NEK1* transcripts in the wild type and

estradiol-resistant transgenic plants harboring *XVE::*Mp*NEK1*.

The 10-day old plants of the wild type (WT) and estradiol-resistant plants derived from line #8 and

#2 were incubated for 1 day in the medium with (+) or without 10 µM estradiol (-). Total RNA was

extracted and subjected to RT-qPCR. All transcript levels are relative to that of mock-treated wild

type plants. Mp*EF1α* and Mp*ACT* were used as the control genes.

Fig S5. Effect of estradiol on the growth of MpNEK1-Citrine inducible lines

The gammae of the wild type (WT) and the MpNEK1-Citrine inducible lines were planted in the

agar medium supplemented with (+ER) or without 1 µM estradiol (Mock) and grown for 15 days.

Fig S6. Expression and localization of MpNEK1-Citrine in the meristem of thallus without

estradiol.

Left panel shows a light field image merged with a confocal image in the right panel. Green;

MpNEK1-Citrine, magenta; plastid autofluorescence.

Video S1. Localization of MpNEK1-Citrine in a gemmaling grown for 4 days in the medium

supplemented with 1μM estradiol (same plant shown in Fig. 6B right panels).

Video S2. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine

inducible line treated with estradiol for 24 hours.

MpNEK1-Citrine inducible line was grown without estradiol for 3 days and then applied with 1 μM

estradiol and grown for 24 hours. Green: MpNEK1-Citrine, magenta: plastid autofluorescence

(same rhizoid shown in Fig. 7C).

Video S3. Localization and dynamics of over-accumulated MpNEK1-Citine in a rhizoid of

MpNEK1-Citrine inducible line treated with estradiol for 24 hours.

MpNEK1-Citrine inducible line was grown without estradiol for 3 days and then applied with 1 μM

estradiol and grown for 24 hours. A rhizoid highly accumulating MpNEK1-Citrine was observed.

Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

Video S4. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine

inducible line treated with estradiol for 48 hours.

MpNEK1-Citrine inducible line was grown without estradiol for 3 days and then applied with 1 μM

estradiol and grown for 48 hours. Green: MpNEK1-Citrine, magenta: plastid autofluorescence

(same rhizoid shown in Fig. 7C).

Video S5. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine

inducible line treated with estradiol for 48 hours.

MpNEK1-Citrine inducible line was grown without estradiol for 3 days and then applied with 1 µM

estradiol and grown for 48 hours. Green: MpNEK1-Citrine, magenta: plastid autofluorescence

(same rhizoid shown in Fig. 8A).

Video S6. Localization and dynamics of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine

inducible line treated with estradiol for 120 hours.

MpNEK1-Citrine inducible line was grown without estradiol for 3 days and then applied with 1 µM

estradiol and grown for 120 hours. Green: MpNEK1-Citrine, magenta: plastid autofluorescence

(same rhizoid shown in Fig. 7C).

Video S7. Localization of MpNEK1-Citine in a rhizoid of MpNEK1-Citrine inducible line treated

with estradiol for 120 hours.

MpNEK1-Citrine inducible line was grown without estradiol for 3 days and then applied with 1 µM

estradiol and grown for 120 hours. Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

Fig1. Effect of estradiol on thallus growth of Mp*NEK1* inducible line.

(A) Construct for Mp*NEK1* induction. Mp*EF1α* promoter, terminator (T), LexA operator (LexA op), minimal CaMV35S promoter (m35Spro). The diagram is not shown to scale.

(B) The gemmae of Mp*NEK1* inducible line (#2) were planted in the agar medium with (+ER) or without 10 μM estradiol (Mock) and grown for 3, 11, and 20 days.

(C) Time course of growth of Mp*NEK1* inducible line (#2) with (+ER) or without 10 μM estradiol (Mock) .

(D) Quantification of thallus growth of Mp*NEK1* inducible line (#2, 4, 8) with (+ER) or without 10 μM estradiol (Mock). The mean projection area of thalli (n = 10 plants) was quantified by ImageJ.

Fig 2. Reversible or irreversible effect of estradiol on thallus growth

(A) Schematic diagram of treatments. Gemmae were planted in the agar medium with 10 µM estradiol, grown in the same medium for 1, 3, 7, or14 days, and then transferred to and grown in the estradiol-free medium for 23 days.

(B) Gemmae of the wild type (WT) and four independent inducible lines (#2, 3, 4, and 8) were grown as shown in (A). The periods of estradiol treatment is shown above the photographs taken at the 23 day after transfer to the estradiol-free medium.

Fig 3. Effect of various concentrations of estradiol on thallus growth.

The gemmae of the wild type and Mp*NEK1* inducible lines were planted in the agar medium supplemented with or without estradiol at the concentration of 10 nM, 100 nM, and 1000 nM and grown for 13 (A) and 16 days (B).

Fig. 4. Estradiol resistance and its transfer to the next generation.

(A) The gemmae of the estradiol resistant plants or sensitive plants of Mp*NEK1* inducible lines were planted in the agar medium supplemented with 10 μM estradiol and grown for 11, 15, and 21 days.

(B) Quantification of growth of gemmalings derived from the estradiol resistant plants or sensitive plants of Mp*NEK1* inducible lines. The mean projection area of thalli (n = 10 plants) was quantified by ImageJ.

Fig. 5. Effect of estradiol on cell proliferation

(A) EdU-labelled nuclei in the gemmalings treated with 1 µM estradiol for 4 days. The gemmae of the wild type (WT) and Mp*NEK1* inducible line were planted in the medium supplemented with or without 1 µM estradiol, grown for 4 days, and incubated with 10 µM EdU for 1 hour in liquid medium of the same kinds. EdU-labelled nuclei were visualized according to the manufacture's instruction as described in methods.

(B) EdU-labelled nuclei in the gemmalings treated with 1 µM estradiol for 1 day. The gemmae of the wild type (WT) and Mp*NEK1* inducible line were planted in the medium without estradiol, grown for 3 days, and then transferred to and grown in the medium supplemented with or without 1 µM estradiol for 1 day. The gemmalings were incubated with 10 µM EdU for 1 hour in liquid medium of the same kinds. EdU-labelled nuclei were visualized according to the manufacture's instruction as described in methods.

(C) Quantification of EdU-labeled nuclei in (A). Data are shown by box plots (n = 12-16 plants). An asterisk indicates significant difference from the control (-ER) (*t*-test, P < 0.05).

(D) Quantification of EdU-labeled nuclei in (B). Data are shown by box plots (n = 12-16 plants). An asterisk indicates significant difference from the control (-ER) (*t*-test, P < 0.05).

A

B

| Mock | +ER (4 days) |

C

| Mock | +ER (1 day) |

Fig. 6. Localization of estradiol-induced MpNEK1-Citrine in the thallus

(A) Construct for Mp*NEK1* induction. Mp*EF1α* promoter, terminator (T), LexA operator (LexA op), minimal CaMV35S promoter (m35Spro). The diagram is not shown to scale.

(B) The gemmae of the Mp*NEK1* inducible line were planted in the medium with (+ER) or without 1 µM estradiol (Mock), grown for 4 days, and observed under a confocal microscope.

(C) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol, grown for 2 days, and then added with (+ER) or without 1 µM estradiol (Mock) for 1 day. Upper panels are light field images merged with confocal images shown in the below panels. Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

Fig. 7. Localization of estradiol-induced MpNEK1-Citrine in rhizoids.
(A) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol, grown for 3 days, and then applied with (+ER) or without 1 μM estradiol (Mock). Rhizoids were observed under a confocal microscope in the time course. Green: MpNEK1-Citrine, magenta: plastid autofluorescence.
(B, C) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol, grown for 3 days, and then applied with 1 μM estradiol and observed at the time indicated in each panel. The same rhizoid was observed under a confocal microscope in the time course. Green: MpNEK1-Citrine, magenta: plastid autofluorescence.
(D) Velocity of rhizoid growth in the MpNEK1-Citrine inducible line incubated for 48 hours with (+ER) or without 1 μM estradiol (-ER). Data are shown by box plots (n = 32 rhizoids). An asterisk indicates significant difference (*t*-test, P < 0.01).

Fig. 8. Movement of MpNEK1-Citrine particles in rhizoids.

(A) The gemmae of the Mp*NEK1* inducible line were planted in the medium without estradiol, grown for 3 days, and then applied with 1 μM estradiol. Rhizoids were observed under a confocal microscope after 48 hours. The lower panel indicates kymograph of MpNEK1-Citrine particles. Green: MpNEK1-Citrine, magenta: plastid autofluorescence.

(B) Velocity of MpNEK1-Citrine particles. Data are shown by box plots (n = 24 or 31 particles). There was no significant difference (*t*-test, P > 0.05).

Fig S1. Effects of estradiol on the accumulation of Mp*NEK1* transcripts in the wild type and transgenic lines introduced with *XVE::*Mp*NEK1*.
The 10-day old plants of the wild type (WT) and the transgenic lines were incubated for 1 day in the medium with (+) or without 10 µM estradiol (-). Total RNA was extracted and subjected to RT-qPCR. All transcript levels are relative to that of mock-treated wild type plants. Mp*EF1α* and Mp*ACT* were used as the control genes.

Fig S2. Effect of estradiol on thallus growth of the wild type.

(A) The gemmae of the wild type (Tak-1) were planted in the agar medium with (+ER) or without 10 µM estradiol (Mock) and grown for 3, 11, and 20 days.

(B) Time course of growth of the wild type with (+ER) or without 10 µM estradiol (Mock) .

(C) Quantification of thallus growth of the wild type with (+ER) or without 10 µM estradiol (Mock). The mean projection area of thalli (n = 10 plants) was quantified by ImageJ.

Fig S3. Effect of estradiol on thallus growth of the transgenic line #6 with *XVE::*Mp*NEK1*.

(A) The gemmae of the line (#6) were planted in the agar medium with (+ER) or without 10 µM estradiol (Mock) and grown for 3, 11, and 20 days.

(B) Time course of growth of the line (#6) with (+ER) or without 10 µM estradiol (Mock) .

(C) Quantification of thallus growth of the line (#6) with (+ER) or without 10 µM estradiol (Mock). The mean projection area of thalli (n = 10 plants) was quantified by ImageJ.

Fig S4. Effect of estradiol on the accumulation of Mp*NEK1* transcripts in the wild type and estradiol-resistant transgenic plants harboring *XVE::*Mp*NEK1*.

The 10-day old plants of the wild type (WT) and estradiol-resistant plants derived from line #8 and #2 were incubated for 1 day in the medium with (+) or without 10 µM estradiol (-). Total RNA was extracted and subjected to RT-qPCR. All transcript levels are relative to that of mock-treated wild type plants. Mp*EF1α* and Mp*ACT* were used as the control genes.

Fig S5. Effect of estradiol on the growth of MpNEK1-Citrine inducible lines

The gammae of the wild type (WT) and the MpNEK1-Citrine inducible lines were planted in the agar medium supplemented with (+ER) or without 1 μM estradiol (Mock) and grown for 15 days.

Fig S6. Expression and localization of MpNEK1-Citrine in the meristem of thallus without estradiol.

Left panel shows a light field image merged with a confocal image in the right panel. Green; MpNEK1-Citrine, magenta; plastid autofluorescence.