Ageing impairs the regenerative capacity of regulatory T cells in central nervous system

remyelination

Alerie Guzman de la Fuente[1,2,3*], Marie Dittmer[1], Elise Heesbeen[1,4], Nira de la Vega Gallardo[1], Jessica

White[3], Andrew Young[1,5], Katie Mayne[1,6], John Falconer[1,7], Christopher E. McMurran[8,9], Mohammed

Innayatullah[1], Rebecca Ingram[1], Vijay Tiwari[1], Rosana Penalva[1], Yvonne Dombrowski[1] and Denise C.

Fitzgerald[1*]

[1]Wellcome-Wolfson Institute for Experimental Medicine, Queen's University Belfast, Belfast, BT9 7BL,

Northern Ireland, UK.

[2]Institute for Health and Biomedical Sciences of Alicante (ISABIAL), Alicante, 03010, Spain.

[3] Institute of Neuroscience of Alicante CSIC-UMH, San Juan de Alicante, 03550, Spain.

[4]Current address: Division of Pharmacology, Institute for Pharmaceutical Sciences, Utrecht University,

Utrecht, The Netherlands.

[5] Current address Patrick G Johnston Centre for Cancer Research, Queen's University Belfast, Belfast,

BT9 7BL, Northern Ireland, UK

[6]Current address: Department of Pathology and Laboratory Medicine, University of British Columbia,

G227-2211 Wesbrook Mall, Vancouver, BC, V6T 2B5, Canada.

[7] Current address: CRUK Beatson Institute, G61 1BD, Glasgow, UK

[8] Department of Clinical Neurosciences, University of Cambridge, Cambridge, UK

[9] Department of Medical Epidemiology and Biostatistics, Karolinska Institute, Stockholm, Sweden

*Corresponding authors:

Denise C. Fitzgerald, d.fitzgerald@qub.ac.uk

Alerie G. de la Fuente, guzman_ale@isabial.es

Abstract

Myelin regeneration (remyelination) is essential to prevent neurodegeneration in

demyelinating diseases such as Multiple Sclerosis but its efficiency declines with age. Regulatory T

cells (Treg) recently emerged as critical players in tissue regeneration, including remyelination.

However, the effect of ageing on Treg-mediated regenerative processes is poorly understood. Here,

we show that expansion of aged Treg does not rescue age-associated remyelination impairment due

to an intrinsically diminished capacity of aged Treg to promote oligodendrocyte differentiation and

myelination. This decline in regenerative Treg functions can be rescued by a young environment. We

identified Melanoma Cell Adhesion Molecule 1 (MCAM1) and Integrin alpha 2 (ITGA2) as novel

candidates of Treg-mediated oligodendrocyte differentiation that decrease with age. Our findings

demonstrate that ageing limits the neuroregenerative capacity of Treg, likely limiting their

remyelinating therapeutic potential in aged patients and describe two novel mechanisms implicated

in Treg-driven remyelination that may be targetable to overcome this limitation.

Introduction

Myelin is critical to the metabolic support and function of axons in the central nervous system

(CNS)[1]. CNS myelin is produced by oligodendrocytes and damage to oligodendrocytes or myelin causes

neurological impairment[2–4]. Myelin regeneration (remyelination) is highly neuroprotective and holds

potential to restore lost function for patients with demyelinating diseases such as Multiple Sclerosis

(MS)[5–7]. Like most regenerative processes, remyelination efficiency declines with age[8,9], contributing

to age-associated disease progression and the accumulation of irreversible disability[10]. Despite the

critical neuroprotective role of remyelination, there are currently no approved remyelination-

enhancing therapies, a key unmet medical need. Thus, understanding how ageing alters the different

cellular mechanisms governing myelin regeneration is key to overcoming age-associated

remyelination failure. This holds potential to prevent neuronal loss and the accumulation of

irreversible cognitive, sensory and motor disabilities.

Regulatory T cells (Treg) have recently emerged as key cellular players in tissue regeneration

due to both anti-inflammatory functions and capacity to modulate tissue-resident stem cell

differentiation[11,12]. Similar crosstalk has also been described in the CNS, where we and others showed

that Treg promote oligodendrocyte progenitor cell (OPC) differentiation and CNS remyelination[13–15].

The discovery of pro-remyelinating functions of Treg opened a new therapeutic avenue to prevent

remyelination decline and associated neurodegeneration. However, whether this important

neuroregenerative function of Treg is impaired with age remains unknown. Understanding how ageing

affects myelin-regenerative functions of Treg is essential to determine the relevance and suitability of

potential Treg-based pro-remyelination therapies in patients of advanced age in MS and other

demyelinating diseases.

Here, we show that ageing impairs the capacity of natural Treg to drive OPC differentiation *in*

vitro and myelination *ex vivo*. Surprisingly, this defect was reversible in a young remyelinating

environment *in vivo*. Transcriptomic comparison of young and aged Treg identified Melanoma Cell

Adhesion Molecule 1 (MCAM1) and Integrin alpha 2 (ITGA2) as novel candidates of Treg-mediated

OPC differentiation that decrease with age. This study identifies Treg-driven OPC differentiation as a

novel remyelination mechanism that is impaired by ageing and identifies potential therapeutic targets

to promote CNS remyelination.

Results

Expansion of Treg does not rescue impaired OPC differentiation in aged mice

We first examined murine Treg frequency with age, and in agreement with previous

studies[16,17], found increased numbers and proportions of Treg in aged mice compared to young mice

(**Fig. 1 A-D, Sup. Fig. 1 A**). We next verified that OPC differentiation into CC1$^+$ oligodendrocytes is

impaired with ageing *in vivo* as previously reported[9,18] (**Fig. 1 E**), despite the endogenous increase in

Treg proportions with age. The impairment in oligodendrocyte differentiation with age was not

rescued by further expanding Treg numbers in aged mice *in vivo* using IL-2/anti-IL-2 administration

(**Fig 1 F-I**). Hence, we hypothesised that the myelin-regenerative capacity of Treg may be intrinsically

impaired in aged Treg, irrespective of Treg frequency.

Aged Treg demonstrate an impaired intrinsic myelin-regenerative capacity

To compare the capacity of young and aged Treg to support the process of myelin

regeneration, we first investigated effects on oligodendrocyte development. To do this, we isolated

natural Treg from the spleens and lymph nodes of young (3-4 months) and aged (14-18 months) mice

(**Sup. Fig. 1 A,B**) and co-cultured these Treg with neonatal murine OPCs. While neither young nor aged

Treg modified OPC proliferation *in vitro* (**Fig. 2 A, B**), young Treg significantly increased OPC

differentiation into myelin basic protein- (MBP) expressing oligodendrocytes, in agreement with our

previous findings of Treg secretome[13,19]. Aged Treg however, did not enhance OPC differentiation

demonstrating an impairment in their capacity to support this vital step in the process of myelin

regeneration (**Fig. 2 C-E**).

To determine if the inability of aged Treg to support OPC differentiation was functionally

relevant to the process of myelin production, we compared the capacity of young and aged Treg to

drive myelin ensheathment of axons *ex vivo.* To do this, we co-cultured murine organotypic brain slices

with young or aged Treg for 7 days. Young Treg significantly enhanced axonal myelination compared

to control slices as described before[13]. However, aged Treg failed to significantly increase myelination

in brain slices compared to controls, demonstrating an impairment in the capacity of aged Treg to

drive myelin production (**Fig. 2 F, G**).

Myelin-regenerative capacity of aged Treg is restored in a young environment

We previously reported that Treg depletion using diphtheria toxin (DT) in B6.129(Cg)-

Foxp3^{tm3(DTR/GFP)Ayr}/J (FoxP3-DTR)[20] mice impaired OPC differentiation and remyelination, which was

rescued by the administration of wild type Treg[13]. Thus, we next compared the regenerative capacity

of young and aged Treg in a model of lysolecithin-induced spinal cord myelin damage *in vivo.* Following

endogenous GFP⁺ Treg depletion (**Sup. Fig. 1 C, D**), we administered wild type (GFP⁻) young or aged

Treg intraperitoneally and confirmed comparable detection of adoptively transferred Treg in each

group (**Sup. Fig. 1 E, F**). We next compared the neuroregenerative capacity of adoptively transferred

young and aged wild type Treg to enhance OPC differentiation during remyelination (**Fig. 3 A**). We did

not observe any differences in the burden of damage for each experimental group, indicated by the

lesion area (**Sup. Fig. 1 H**). There was also no difference in the number of oligodendrocyte lineage

cells, quantified by Olig2 immunostaining (**Fig. 3 B-D**), or proliferating OPCs (Olig2⁺Ki67⁺) at 14 days

post-lesion (dpl) (**Sup. Fig. 1 G, I, J**). As before[13], Treg depletion impaired OPC differentiation into CC1⁺

and ASPA⁺ oligodendrocytes, which was rescued by adoptive transfer of wild type young Treg.

Unexpectedly however, aged Treg injected into young Treg-depleted mice also rescued impaired OPC

differentiation, comparable to young Treg, (**Fig. 3 B, C, E, F**). To determine if aged Treg not only rescued

OPC differentiation but also restored remyelination, we examined axonal wrapping through the

quantification of circular myelin basic protein (MBP⁺) patterns around neurofilament-H⁺ (NFH⁺) axons

as an early indication of axonal remyelination. Both, young and aged Treg administered to Treg-

depleted mice, restored axonal wrapping to levels comparable to non-depleted controls (**Fig. 3 G-J**).

These findings suggest that the age-induced defect of neuroregenerative Treg functions observed *in*

vitro/ex vivo is reversible in a young systemic environment *in vivo*, and therefore, potentially

therapeutically targetable.

Ageing significantly alters Treg transcriptome

We next sought to identify how ageing impairs the myelin regenerative functions of Treg by

examining gene expression differences in natural Treg purified from young and aged female mice. RNA

sequencing (RNAseq) analysis showed that young and aged Treg clustered separately (**Fig. 4 A, C**) as

shown in the literature[21,22] and identified 1423 upregulated and 302 downregulated mRNA transcripts

in aged Treg compared to young Treg (**Fig. 4 B**). However, there was no difference in the expression of classic Treg markers such as *Foxp3*, *Cd4* or *Il2ra* (**Sup. Fig. 2 B**). Gene Ontology (GO) analysis identified that the differentially expressed genes were associated with pathways linked to the adaptive immune response, lymphocyte-mediated immunity, phagocytosis, cell-surface receptor signalling and membrane invagination (**Fig. 4 D)**. Genes that were upregulated with ageing were enriched for the following GO biological processes: extracellular matrix organisation, cell migration, synapse assembly, axonal guidance, cell adhesion and inflammatory and immune responses (**Fig. 4 E**). On the other hand, mRNA transcripts that were downregulated in aged Treg were associated with defence responses to virus, B cell proliferation and protein folding GO biological processes (**Fig. 4 E)** (GO cellular component, GO Molecular function and Kyoto Encyclopedia of Genes and Genome (KEGG) database pathways analysis is shown in **Sup. Fig. 2**). To identify potential mechanisms underlying how ageing may impair myelin regenerative functions of Treg, we examined expression of genes associated with the Treg tissue repair programme described recently for skin, colon, lung and adipose tisse[23]. Unexpectedly, most of the 31 Treg tissue repair programme hallmark/signature genes[23] were in fact upregulated with ageing, except for *Klrb1, Ccr5, Selplg, Bach2, Rad50, Tfrc, Cd84 and Cd164* (**Fig. 4 F, G**). No difference in expression was detected in these hallmark genes between male and female natural Treg, showing that gender does not alter the expression of genes associated with the tissue repair programme (**Sup. Fig. 2 C, D**). These data suggest that the mechanism(s) involved in Treg-driven myelin regeneration capacity that are lost with ageing, are likely different to mechanisms from the Treg tissue repair programme described by Delacher et al[23].

Mcam and Itga2 are associated with Treg capacity to drive oligodendrocyte formation

        To narrow down age-induced changes in Treg that diminish their capacity to induce OPC differentiation, we first investigated whether the putative mechanisms are soluble or contact-mediated; the latter being suggested by the close physical association of Treg with OPCs observed in co-cultures (**Fig. 5A**). Young natural Treg were co-cultured either directly in contact with OPC, or in

the same well but separated using a transwell, which allowed for the exchange of secreted molecules

but not direct OPC-Treg contact. While young Treg co-cultured directly with OPCs enhanced OPC

differentiation into MBP+ oligodendrocytes, this pro-differentiating effect was lost when Treg were

physically separated from OPCs (**Fig. 5 B, C**). To identify potential cell-to-cell contact mechanisms

responsible for Treg-driven OPC differentiation that may be lost with ageing, we combined our aged

Treg RNA sequencing data with a publicly available RNAseq dataset that compares between OPCs,

newly differentiated and differentiated oligodendrocytes[24]. We hypothesised that ligand-receptor

interactions between Treg and plasma membrane receptors enriched in OPCs underlie the pro-

differentiation effect of Treg on OPC. We prioritised ligand-receptor partnerships between Treg and

OPCs that were impacted by ageing, to identify mechanisms that could explain the lost myelin

regenerative capacity of aged Treg. We then performed a protein-protein interaction analysis using

String[25] (**Fig. 5 D**) and prioritised plasma membrane proteins that had the highest decrease in

expression for aged Treg and the highest number of potential binding partners amongst OPC-enriched

receptors compared to oligodendrocytes (**Fig. 5 E**). We identified the following key candidates: *Ccr7,*

Itga2 (CD49b), *Klrb1c* (NK1.1), *Ly6c1, Mcam* (CD146) and *Sell* (CD62L) (**Fig. 5 D-F**). Next, we used a loss-

of-function approach to investigate whether these candidates may confer the lost regenerative

capacity of aged Treg, namely, CD62L, NK1.1, LY6C, MCAM (CD146) and ITGA2 (CD49b). Of these five

targets, antibody-mediated blocking of MCAM (CD146) and ITGA2 (CD49b) on Treg prior to Treg-OPC

co-culture inhibited the capacity of young Treg to drive OPC differentiation *in vitro* as indicated by the

fold change in the percentage of MBP+ cells and the MBP+ area per well (**Fig. 5 G-I, Sup. Fig. 2 H, I**).

This suggests that MCAM and ITGA2 are previously unidentified mediators of pro-remyelinating

functions of Treg and are downregulated with age.

Discussion

Ageing is one of the strongest risk factors associated with the transition between relapsing-remitting to secondary progressive MS[10], a phase that is characterised by the accumulation of neurodegeneration and irreversible disability. This is partly due to impaired remyelination with age[8,9]. Several factors have been linked to age-associated myelin regeneration failure, such as impaired myelin debris phagocytosis by microglia and macrophages[26–28] and reduced capacity of OPCs to differentiate into myelin-forming oligodendrocytes[18,29]. Therapies that enhance remyelination and prevent age-related neurological decline are an urgent unmet need. Beyond their classic immune-modulatory and anti-inflammatory role, Treg have recently emerged as key regulators of tissue homeostasis and regeneration in several tissues[27], including the CNS, where we have previously identified Treg as key promoters of remyelination[13]. However, how ageing affects the capacity of Treg to enhance myelin regeneration had not been addressed.

Peripheral Treg increase with age[16,21,30], a somewhat paradoxical finding given the tissue-regenerative functions of Treg and the impairment of tissue regeneration with age. Here, we further expanded the murine Treg population *in vivo* and showed that despite this increased cell number, aged-associated remyelination failure was not rescued. These data suggest that with ageing, Treg have an intrinsically impaired myelin regenerative capacity. Ageing is known to enhance Treg senescence, limit Treg proliferation[16,21] and there are conflicting reports on the effect of ageing on Treg immunosuppressive capacity[16,21]. Additionally, previous work on muscle regeneration showed that aged Treg have an impaired migratory capacity that hinders muscle regeneration[31]. In a model of influenza infection, aged Treg demonstrated an impaired capacity to support lung repair, which was associated with a cell autonomous impairment in their regenerative programme[22]. Here, we sought to determine whether aged Treg have an intrinsically impaired regenerative capacity to drive CNS myelin regeneration combining *in vitro, ex vivo* and *in vivo* approaches.

Purified aged Treg failed to significantly enhance OPC differentiation into MBP-expressing oligodendrocytes or the myelination of cerebellar slices, which is in striking contrast to young Treg, which robustly drove both biological processes. These data suggest that ageing is associated with a

cell-intrinsic impairment of the regenerative capacity of Treg in CNS myelin regeneration. Given that

IL-2-mediated Treg expansion *in vivo* is in clinical trial for MS (clinical trial number NCT02424396), our

findings are of translational importance as they suggest that Treg expansion as a potential pro-

remyelination therapy may be limited in older populations.

We next investigated whether the regenerative capacity of aged Treg could be restored, and

therefore therapeutically targetable. Surprisingly, aged Treg rescued OPC differentiation and myelin

wrapping similarly to that of young Treg in an *in vivo* model of young spinal cord remyelination. In

contrast to findings in the lung[22], the CNS remyelinating capacity of Treg can be partially restored by

a young environment. Previous studies identified a tissue repair transcriptomic signature in the skin,

adipose tissue, and the lung[23]. Unexpectedly, very few of these hallmark Treg tissue repair genes were

downregulated in aged Treg in our transcriptomic profiling studies. These data indicate that although

some Treg regenerative mechanisms may be common across tissues, other mechanisms may be

tissue-specific or process- specific[32].

We recognised that the impairment of myelin regenerative Treg functions with age presented

a new avenue to discover novel mechanisms of how Tregs support CNS remyelination and the

potential to identify new therapeutic targets to boost remyelination. Indeed, using a combined

approach of Treg/OPC transcriptomic dataset cross-refencing with *in vitro* antibody blocking assays,

we identified MCAM and ITGA2 as novel candidate mediators of Treg-driven OPC differentiation.

These two candidates are downregulated in aged compared to young Treg, likely interact directly with

OPCs and inhibit young Treg-driven OPC differentiation when blocked *in vitro*.

MCAM has previously been described as a cell-adhesion molecule contributing to brain

inflammation by facilitating pathogenic T-cell extravasation into the CNS[33–35]. However, our study is,

to our knowledge, the first report showing that MCAM signalling can promote OPC differentiation.

Together with a recently published abstract examining Treg in MS patients (Poster 4[36]) these data also

represent the first evidence that Treg express MCAM. We have also identified ITGA2 as a novel

mediator of Treg-driven OPC differentiation that is decreased with age. This aligns with previous

reports showing that other integrins are involved in oligodendrocyte development[37] and shows that

Treg are additional cellular contributors to integrin-mediated OPC differentiation.

Collectively, these studies have identified that ageing is associated with a cell-intrinsic

impairment of the myelin-regenerative functions of Treg. This discovery increases the range of

remyelination-associated mechanisms that are impaired with age[8,9,18,26–29,38]. Encouragingly, we have

shown that this impairment can be rescued in a young environment *in vivo*. We have also identified

two novel mediators of Treg-driven OPC differentiation, MCAM and ITGA2. The findings from this

study identify novel mechanisms involved in CNS regeneration which may hold therapeutic potential

for patients with demyelinating diseases.

Methods

Animals

All mice were on a C57BL/6J background and were either bred in-house or purchased from Charles

River Laboratories, UK. Foxp3-DTR mice were a kindly provided by Prof. Alexander Rudensky

(Memorial Sloan Kettering Institute, New York). Neonatal C57BL6/J P3-P7 pups were used for OPC and

brain slice preparations, which were combined with natural Treg were isolated exclusively from young

(3-4m) and aged (15-18m) C57BL6/J male mice for use with neonatal mixed sex pups. For *in vivo* Treg

adoptive transfer studies, natural Treg were isolated from young (3-4m) and aged (15m) C57BL6/J

female mice, and were injected into both, female and male recipient Foxp3-DTR mice (2-4m). To

deplete Foxp3+ cells in Foxp3-DTR mice, diphtheria toxin was administered as described below. All

animal maintenance and experiments were done in accordance with the UK Home Office regulation

(Project Licences 2789 and 2894) and were approved by the Queen's University Belfast's Animal

Welfare and Ethical Review Committee.

Lysolecithin-induced spinal cord demyelination *in vivo*

Spinal cord demyelination was induced as described previously[13]. In brief, demyelination was induced

in the ventral white matter funiculus of the thoracic spinal cord (between vertebrae T11-12 or T12-

13) by the injection of 1.2 µL of 1% (w/v) L-α-Lysophosphatidylcholine (Lysolecithin; Sigma-Aldrich)

under general anaesthesia. At 10- or 14-days post lesion (dpl), mice were terminally anaesthetised

with intraperitoneal (i.p.) pentobarbital injections and transcardially perfused with ice-cold phosphate

buffered saline (PBS) followed by 4% paraformaldehyde (PFA) (Sigma-Aldrich). Spinal cords were

dissected and immersed overnight in 4% PFA at 4 °C. Next, spinal cords were cryoprotected with 30%

sucrose (Sigma-Aldrich) in PBS for 72h and snap-frozen in OCT (Tissue-Tek). Frozen spinal cords were

cryosectioned at 12 µm thickness and immunostained as described below.

In vivo **Treg expansion**

Aged mice (15m) were intraperitoneally injected with IL-2/anti-IL-2 antibody complexes to expand

endogenous Treg[39]. In brief, murine IL-2 (1 µg; Peprotech) was mixed with anti-mouse IL-2 (5 µg;

Bioxcell, Clone JES6-1A12) in PBS at 37 °C for 30 min. Each mouse was injected daily for 3 consecutive

days with IL-2 (1 µg) and anti-IL-2 (5 µg) in 200µl of PBS. Adequate Treg expansion was assessed by

flow cytometry as described below. Spinal cord demyelination was induced on the day after the third

injection of IL-2/anti-IL-2, by injecting lysolecithin into the ventrolateral white matter as described

above. Treg expansion was examined by flow cytometry as described below on day 10dpl, while OPC

differentiation was examined at 10 dpl.

In vivo **Treg depletion and adoptive transfer of wild type natural Treg**

Young (2-4m) male and female Foxp3-DTR mice were injected daily i.p. with diphtheria toxin (DT; 0.04

µg/g of body weight; Sigma, Cat. No. D0564) for 3 days prior to demyelination. To maintain

endogenous Treg depletion during remyelination, DT (0.04 µg/g of body weight) was injected i.p. every

fourth day. Control animals received 200 µl of saline i.p. Depletion was confirmed at the endpoint by

flow cytometric analysis of endogenous GFP⁺ natural Treg in blood, spleen, and lymph nodes (see

below). To adoptively transfer young (2-4m) and aged (15m) natural Treg to Treg-depleted mice, we

immunomagnetically isolated natural Treg from wild type female mice using a CD4$^+$ and CD25$^+$ Treg

isolation kit (STEMCELL Technologies) as described below. In the 24h prior to lysolecithin-induced

demyelination, mice were injected i.p. with 10^6 wild type natural Treg which are resistant to DT.

Adequate reconstitution was evaluated by flow cytometry for Treg in lymph nodes at the endpoint

(see below).

OPC isolation and culture

OPCs were isolated from P3-P7 mice based on A2B5 expression. Briefly, pups were culled by an

overdose of pentobarbital (200 mg/mL, 20µl per mouse) and whole brain was dissected and placed in

ice-cold Hibernate-A (ThermoFisher Scientific). Dissected brain was minced using scalpels, transferred

to a 15mL tube, and centrifuged for 1 min at 100 g and 4 °C. Hibernate-A medium was aspirated, and

the pellet was resuspended in 5 mL dissociation media containing 165U of Papain (Worthington) and

DNAse type I (40 µg/mL; Worthington) in hibernate A for 30 min at 37 °C. Upon digestion papain was

washed off with Hanks Buffered Salt Solution (HBSS) (ThermoFisher Scientific) and cells were

centrifuged for 5 min at 300 g and 4 °C. The pellet was resuspended in 4 mL trituration buffer

(Hibernate-A with 2% B27; ThermoFisher Scientific) and 2 mM sodium pyruvate (ThermoFisher

Scientific) and gently triturated ten times using a 5 mL pipette. Tissue was left to settle for 2 min at

room temperature (RT), and supernatant was transferred to a clean 50 mL tube through a 70 µm

strainer (Corning). Remaining tissue was exposed to another 2 mL of trituration buffer and gently

triturated using a glass polished pipette. After trituration, tissue was left to settle for 2 min at RT and

supernatant was transferred to the tube through the 70 µm strainer. This step was repeated another

time with a glass fire-polished pipette of decreased diameter and then with a 1 mL pipette. To further

remove debris from the cell suspension, 11.5 mL of 90% Percoll (GE Healthcare) diluted in 10X PBS

(ThermoFisher Scientific) was added and topped up to a final volume of 45 mL with Dulbecco's

Modified Eagle's medium (DMEM) (ThermoFisher Scientific). The cell suspension was then centrifuged

at 800 g for 20 min at 4 °C. Upon centrifugation, the pellet was washed with HBSS and then

resuspended in 10 mL magnetic-activated cell sorting (MACS) buffer (Hibernate-A with 2 mM sodium

pyruvate, 2% B27, 0.5% BSA; ThermoFisher Scientific), 2 mM EDTA (Sigma-Aldrich) and 10 µg/mL

insulin (Sigma-Aldrich), and incubated for 30 min at 37 °C in a 10 cm petri dish coated with BSC1

Griffonia Simplicifolia Lectin (BSL1, 5 µg/mL)(Vektor Labs) to remove microglia. Then, the supernatant

was collected, and cells were counted and centrifuged at 300 g for 5 min at 4 °C. The pellet was

resuspended and incubated with 2 µg of A2B5 antibody (Millipore, Clone A2B5-105) in 500 µL of MACS

buffer per 10 million cells for 25 min on ice, with gentle resuspensions every 10 min. Cells were washed

with HBSS and spun at 300 g for 5min at 4 °C (Miltenyi Biotec). The pellet was resuspended and

incubated with 80 µL of MACS buffer and 20 µL of anti-IgM microbeads (Miltenyi Biotec) for 15 min.

Then, cell suspensions were centrifuged at 300 g for 5 min at 4 °C, resuspended in 0.5 ml MACS buffer

and placed in a MACS mini column (Miltenyi Biotec) on a MiniMACS Separator (Miltenyi Biotec) and

washed with 1.5mL MACS buffer. Once the liquid passed completely through the column, the column

was removed from the stand and placed in a new 15 mL tube, where 1 mL of OPC media was added

to the column and plunged through to elute the OPCs. OPC media contained DMEM, 2% B27, sodium

pyruvate (2 mM), insulin (5 µg/mL), Trace Elements B (0.01%; Corning), Forskolin (5 µM; Sigma-

Aldrich), Biotin (10 ng/mL; Sigma-Aldrich), Penicillin-streptomycin-glutamine (1%; ThermoFisher

Scientific), N-acetyl cysteine (60 µg/mL; Sigma-Aldrich) and 1% SATO stock solution. SATO stock

solution contained bovine serum albumin (BSA) fraction V (0.1 mg/mL; ThermoFisher Scientific),

sodium selenite (4 µg/mL; Sigma-Aldrich), putrescine (1.61 mg/mL; Sigma-Aldrich), apo-transferrin

(0.1 mg/mL; Sigma-Aldrich) and progesterone (4 µg/mL; Sigma-Aldrich). The purified population was

counted and diluted in OPC media for plating. OPCs were plated at a density of 3,000 cells per well in

96-well plates (Falcon) previously coated with poly-L-lysine (10 µg/mL; Sigma-Aldrich) and laminin (10

µg/mL; Sigma-Aldrich) diluted in DMEM. OPCs were plated with PDGFaa (20 ng/mL; Peprotech) and

NT-3 (10 ng/mL; Peprotech). The following day two thirds of the media was removed (100 µl) and 100

µl of fresh OPC media with PDGFaa (20 ng/mL) and NT3 (10 ng/mL) were added per well. Cells were

incubated at 37°C and 5% CO_2 for two days. On the third day, media was completely removed and

20,000 Treg in fresh media were added as indicated below.

For transwell assays, OPCs were selected with anti-PDGFRα panning. After microglia depletion in BSL1-

coated petri dishes, the supernatant was transferred to PDGFRα panning plates and incubated at RT

for 1 h. PDGFRα panning plates were coated with goat-anti-rat IgG (H+L) (7.5 µg/mL; ThermoFisher

Scientific) in PBS with 0.2% BSA and rat anti-PDGFRα (1.5 µg/mL in PBS; BD Biosciences) with 0.2% BSA

for 3 h. Supernatants were removed, cells were washed 5-8 times with PBS and the attached OPCs

were scraped, counted, and replated in 24-well plates (Sarstedt) at 10,000 cells per well. On the third

day *in vitro* 50,000 Treg were added either directly into the well with OPCs or in a transwell (Millipore)

above the OPCs.

After 6 days of OPC-Treg co-culture cells were fixed for 15 min with 4% PFA and washed twice with

PBS prior to immunostaining. Cells were blocked with 5% donkey serum (Sigma-Aldrich) with 0.1%

Triton-X-100 (Sigma-Aldrich) in PBS for 1h at RT. OPCs were then stained overnight at 4 °C with primary

antibodies diluted in blocking solution against Olig2 (1:500, Bio-techne), CNPase (1:500, Sigma-Aldrich,

clone 11-5B), MBP (1:500, Millipore, clone 12), Ki67 (1:300, Abcam, clone SP6), NG2 (1:200, Millipore),

and CD3 (1:500, eBioscience). Cells were then washed three times with PBS and incubated for 1 h at

room temperature with secondary antibodies diluted in blocking buffer (Alexa fluor (AF) 488 donkey

anti- rabbit (1:500; ThermoFisher Scientific), AF568 donkey-anti- rat (1:500; Abcam), AF647 donkey

anti-mouse (1:500; Abcam), AF647 donkey anti-rabbit (1:500; ThermoFisher Scientific) and AF755

donkey anti-goat (1:500; ThermoFisher Scientific) and Höechst (1:10,000; Sigma-Aldrich). Cells were

then washed with PBS three times and stored in 150 µL PBS/well for cell imaging.

Cell imaging was performed using the CellInsight CX5 high content imaging system (ThermoFisher

Scientific) for 96-plates and EVOS for 24-well plates. Twenty-five separate fields of view of each well

in 96-well plates and the cell populations of interest were quantified using CellInsight CX5 analysis

software and the SpotCounts programme. For 24-well plates we imaged randomly 5 fields of view per

well at 20X magnification and images were analysed manually with Fiji.

Natural Treg isolation

Young (2-4 m) and aged (15-18 m) male mice were culled by CO_2 overdose. Spleens and lymph nodes were removed and mashed through a 70 µm strainer with a 2 ml syringe plunger. $CD4^+$ T cells were immunomagnetically purified by negative selection according to manufacturer's instructions (STEMCELL Technologies). Isolated $CD4^+$ T cells were then subjected to a $CD25^+$ cell isolation (STEMCELl Technologies) using releasable magnetic spheres (STEMCELL Technologies) according to the manufacturer's protocol. Purity was confirmed by flow cytometry for CD4, CD25 and Foxp3 (**Sup. Fig. 2A**). For i.p. injections in adoptive transfer experiments, cells were resuspended at 10^6 cells per 200 µL in saline. For OPC-natural Treg co-cultures 20,000 natural Treg were resuspended in 150 µl of OPC Brainphys-based media and added to OPCs in 96-well plates. For 24-well transwell plate experiments 50,000 natural Treg were resuspended in 250 µl OPC-BrainPhys media and added to transwells above cultured OPCs. OPC BrainPhys-based media contains BrainPhys (STEMCELL Technologies), B27 (2%; ThermoFisher Scientific), Glutamax (0.5 mM; ThermoFisher Scientific), N-acetyl-cysteine (60 µg/mL), SATO (1:100, as described above) and insulin (5 µg/mL). In the case of brain slices, 50,000 Treg were diluted in 5 µL of brain slice media and dropped directly onto brain slices.

For natural Treg mechanistic studies, OPCs and Treg were isolated as described above. Prior to OPC-natural Treg co-culture, 120,000 Treg were transferred to a 1.5 mL microcentrifuge tube and centrifuged at 400 g for 5 min at 4 °C. Treg were resuspended in 100 µL of PBS with 2% FCS and a blocking antibody (20 µg/mL) or corresponding isotype were added per tube. Natural Treg were incubated with the blocking antibody for 45 min on ice and then cells were washed with PBS and centrifuged for 10 min at 400 g and 4 °C. Cells were resuspended in 900 µl of OPC BrainPhys-based media (see above) and 20,000 Treg in 150 µL of media were added per well in 96-well plates. As previously, Treg were co-cultured with OPCs for 7 days *in* vitro (DIV) and then fixed and stained as described above. Blocking antibodies used were anti-NK1.1 PE (eBioscience, clone PK136), anti-CD62L-APC (eBioscience, clone Mel14), anti-Ly6c1-APC (eBioscience, clone RB6-BC5), rat-isotype-APC

(eBioscience), anti-Itga2 (Abcam, clone EPR5788), anti-MCAM (Abcam, clone EPR3208) and rabbit-

isotype (Vektor labs).

Organotypic brain slice cultures

Brainstem slices from male and female P3 C57BL/6J mice were prepared at 300 μm thickness using a

McIlwain Tissue Chopper as described previously[13]. Slices were separated and placed on Millicell

inserts in individual wells of a 24-well plate (Millipore) with 250 μl brain slice medium containing 46.6%

minimum essential medium (ThermoFisher Scientific), 25% Earls balanced salt solution (Sigma-

Aldrich), 25% heat-inactivated horse serum (Thermofisher Scientific), 1% penicillin-streptomycin

(Thermofisher Scientific), 1% glutamax (Thermofisher Scientific) and 1.4% D-glucose (Sigma). Brain

slices were incubated at 37 °C and 5% CO_2 overnight and then media was fully replaced. On day 3,

media was changed again and 50000 young or aged Treg were added directly on top of each slice in a

5 μl droplet. Brain slices were cultured for a further 7 DIV, with media changes every other day. On

day 10 (7 days post-treatment), brain slices were fixed with 4% PFA for 45 min and immunostained. In

brief, brain slices were blocked for 2 h in 10% normal donkey serum (Sigma-Aldrich), 1 mM Hepes

(Thermofisher Scientific), 1% BSA (ThermoFisher Scientific) and 0.5% Triton X-100 (Sigma-Aldrich) in

PBS. Slices were incubated with primary antibodies for rat anti-MBP (1:500; Millipore, clone 12) and

chicken anti-NFH (1:500; EncorBiotech, polyclonal) in blocking buffer at 4 °C for two overnights. Brain

slices were then washed with PBS-0.01% triton three times for 30 min. Slices were incubated with

secondary antibodies (AF488 donkey anti-chicken IgY (1:500; Abcam, polyclonal) and AF568 donkey

anti-rat IgG (1:500; Abcam, polyclonal) in blocking buffer overnight at 4 °C. Brain slices were washed

twice with PBS-0.01% triton and then incubated with Höechst stain (1:20000; Sigma-Aldrich) for 10

min. Slices were washed with PBS and mounted using fluoromount G (Thermofisher Scientific).

Imaging was performed using a Leica SP8 confocal microscope, using a 63X oil objective and the Leica

Navigator at 0.5 μm intervals over 10 μm. Four fields of view per brain slice were selected for imaging

based on NFH and Höechst staining but blinded to MBP. To quantify myelination index, the area

stained by NFH, MBP and the colocalising area of NFH$^+$MBP$^+$ were measured per stack in Fiji[40] using a

Fiji plug-in developed and kindly provided by the Williams laboratory at the University of Edinburgh[41].

Then, the ratio between total NFH$^+$ and NFH$^+$MBP$^+$ areas (myelination index) was calculated per z-

stack and the average of the myelination index per slice was calculated.

Flow cytometry

Spleens and lymph nodes were mashed through a 70 µm strainer. For splenocytes, cells were exposed

to red blood cell lysis buffer (STEMCELL Technologies) for 2 min at room temperature. Both, lymph

nodes and splenocytes were then washed with PBS and centrifuged at 300 g for 5 min at 4°C. Cells

were resuspended in 200 µL PBS and stained with a cell viability dye with eFluor 455-UV viability dye

(1:2000; ThermoFisher Scientific) and cell surface stained with antibodies for CD4 (1:500; eBioscience,

clone RM4.5) and CD25 (1:500; eBioscience, clone PC61.5) for 15 min at RT. Cells were washed with

flow cytometry staining buffer (FCSB) (2% FCS in PBS) and centrifuged at 300 g for 5 min at 4 °C. Cells

were then fixed with Fix & Perm A (ThermoFisher Scientific) for 10 min at RT. Fixative was washed off

with FCSB and centrifuged at 300 g for 5 min at 4 °C. The pellet was resuspended in PBS and data were

acquired on a FACSCanto II. In the natural Treg depletion experiment, presence of endogenous natural

Treg was determined by the expression of GFP. To determine the purity of natural Treg isolations or

the extent of natural Treg reconstitution by adoptive transfer, cells were washed and centrifuged for

5 min at 300 g and 4 °C. Cells were then resuspended in 100 µL Fix & Perm B (ThermoFisher Scientific)

with an anti-Foxp3 antibody (1:100; eBioscience, clone FJK-16S) overnight at 4 °C. Cells were then

washed with FCSB and centrifuged at 300 g and 4 °C for 5 min. Cells were then resuspended, data

were acquired on a FACSCanto II and analysed using FlowJo software version 9.0 (BD). To calculate

cell numbers, singlets were identified by FSC-H versus FSC-A and viable cells gated for CD3 and CD4,

and subsequently CD25 and Foxp3$^+$GFP$^+$ and Foxp3$^+$GFP$^-$ cells. To evaluate young and aged Treg

numbers in blood, 30 µL were incubated with 100 µL of the viability and cell surface antibody mix for

30 min at room temperature. Cells were then washed with FCSB and centrifuged at 300 g for 5 min at

4 °C. After resuspension, cells were fixed and lysed in 100 µL of OptiLyse B (Beckman Coulter) for 10

min at RT, washed with distilled water and then after 15 min, acquired on a FACSCanto II. After initial

data acquisition, cells were centrifuged at 300 g for 5 min at 4 °C and incubated overnight with anti-

Foxp3 antibody in Fix & Perm B, as described above. Cells were washed and data were acquired on a

FACSCanto II.

Immunofluorescence staining of CNS tissue

Spinal cord sections were dried for 30 min at RT and washed for 10 min in PBS. For Olig2, Ki67, CC1

and ASPA staining, spinal cord sections underwent antigen retrieval at 85 °C for 10 min with 1X citrate

buffer pH 6.0 (Sigma-Aldrich). Tissue sections were washed with PBS and permeabilised with 1%

Triton-X-100 in PBS for 30 min at RT. Sections were washed with PBS and incubated with blocking

solution (5% donkey serum; Sigma-Aldrich) diluted in TBS with 0.25% tween (Sigma-Aldrich) for 1h at

RT. If spinal cord sections were incubated with a primary antibody raised in mouse (e.g. anti-CC1)

sections underwent an additional 1 h blocking step at RT using Mouse on Mouse (M.O.M) blocking

reagent (Vector labs) in 5% donkey serum. Then spinal cord sections were incubated with primary

antibodies against Olig2 (1:500; Bio-Techne, polyclonal), anti-APC (1:400, Abcam, clone CC1), anti-

ASPA (1:300, Millipore, polyclonal), anti-NFH (1:500; Abcam, polyclonal), anti-MBP (1:500; Millipore,

clone 12) or anti-Ki67 (1:300; Abcam, clone SP6) overnight at 4 °C. Sections were washed with Tris-

Buffered-Saline (TBS) (ThermoFischer Scientific) with 0.25% Tween (Sigma-Aldrich). Secondary

antibodies including donkey anti-goat AF488, donkey anti-mouse AF568, donkey anti-rabbit AF647 (all

1:500; ThermoFisher Scientific), donkey anti-rat AF568 (1:500; Abcam), donkey anti-rabbit AF647

(1:500, Abcam) and Höechst (1:10000; Sigma-Aldrich) were added for 1 h at room temperature. Spinal

cord sections were then washed with TBS-0.25% Tween twice and mounted with fluoromount G.

Image acquisition was performed using the Leica TIRF and Leica DM5500 widefield fluorescent

microscopes. Further image processing was performed in Fiji[40] software and analysis was undertaken

by blinded manual counting. For NFH and MBP wrapping analysis, images were quantified using Cell

Profiler and Cell Profiler Analyst softwares[42,43].

RNA sequencing and analysis

Young (2-4m) and aged (16m-20m) natural Treg were isolated from spleen and lymph nodes by

magnetic-activated cell sorting and purity was checked by flow cytometry as described above (**Sup.**

Fig. 3A). Natural Treg were lysed in 0.5 mL of Trizol (Sigma-Aldrich) by vortexing and pipetting and

frozen at -80 °C. RNA was extracted combining Trizol and Chloform isolations with RNAeasy micro kit

columns (Qiagen). RNA concentration was determined using a Nanodrop, and RNA quality was

assessed by Qubit measurement and in an RNA nanochip Bioanalyzer. Sequencing libraries were

prepared using the Kapa Hyper Preparation kit with riboerase (Roche) following manufacturer's

instructions. Sequencing was performed on the NovaSeq 6000 in a pair-end 75 base pair format and

80 million reads.

The raw sequencing data was checked for lower quality bases and adaptor sequences with FastQC and

quality trimming was performed using CutAdapt[44]. The high-quality trimmed data were mapped on

mm10 mouse reference genome using STAR aligner (v 2.7.0a)[45]. The quantification of each gene

transcripts was performed with ensemble transcript annotation gtf file (GRCm38) using featureCounts

tool (v 2.0.0)[46]. The resultant read counts matrix was analyse using DESeq2 (v1.38.1)[47] to identify

differentially expressed genes between the young and aged natural Treg samples using Wald tests

with the Bonferroni correction for multiple testing. The differentially expressed genes (DEG) with a

multiple testing adjusted p<0.05 and fold change >2 were considered for further downstream analysis.

These genes were subjected to gene ontology enrichment analysis using enrichGO function in

clusterProfiler (v4.2) R package and DAVID (https://david.ncifcrf.gov). Over and underrepresented

ontology terms were identified (p value <= 0.05 and Bonferroni correction p adjustment method).

Next, enriched activated and suppressed functional pathways were identified using gseKEGG function

in clusterProfiler with p value<0.05.

Raw data files from RNAseq are available at the GEO database (accession number GSE218804).

Bioinformatic analysis to identify putative Treg mechanisms

To determine the molecular mechanisms associated with Treg-driven OPC differentiation that are

impaired with ageing, we combined our young and aged natural Treg RNA sequencing with data

obtained from Zhang *et al*. 2014 (GSE: 52564)[24]. We first extracted genes that were downregulated in

aged natural Treg and associated with the GO cellular component term "plasma membrane". We then

extracted the FPKM matrix and selected genes that were upregulated in OPC when compared with

newly myelinating oligodendrocytes and oligodendrocytes in the Zhang *et al*. database. We performed

GO cellular component analysis on those OPC-enriched genes and selected genes associated with the

GO term Apical Plasma Membrane. Then, both gene lists were subjected to a protein-protein

interaction analysis using the String open-source tool (www.string-db.org)[25]. Protein-protein

interactions within genes that were associated only with the natural Treg or only the OPC databases

were eliminated, and the remaining downregulated natural Treg membrane proteins were prioritised

considering their significance, fold change and the number of potential interactors enriched in OPCs.

Candidate genes were then selected based on their p-adjusted value, fold enrichment and the number

of interactions with OPC receptors.

Statistical analysis

All statistical analyses were performed in GraphPad Prism (GraphPad Software, Inc. version 9) or R.

First normality of datasets was assessed using Shapiro-Wilk and Kolmogorov-Smirnov tests. For

comparisons between 2 groups, such as young and aged Treg and OPC differentiation (Sup. Fig. 1),

unpaired two-tailed Student's t tests were used if data were normally distributed, and Mann-Whitney

U tests for non-parametric datasets. When more than two groups were analysed and datasets were

normally distributed, a one-way analysis of variance (ANOVA) was performed assuming equal

variances, followed by Sidak's multiple comparison test. When two factors were controlling the

outcome, such as replicates and treatments, 2-way ANOVA followed by Dunnet's comparison was

used. When datasets were was not normally distributed, a Kruskal-Wallis followed by Dunn's multiple

comparison test was used. For percentage data, arcsin conversion was performed to analyse the data

using parametric tests. For all statistical tests, differences were considered significant at $p < 0.05$.

Data and code availability

Young and aged natural Treg RNA sequencing data will also be deposited at GEO (NCBI) with the

revised version.

Figure legends

Figure 1: Expansion of Treg does not rescue impaired OPC differentiation in aged mice. A) Flow

cytometric plot and **B**) quantification of natural Treg proportion, identified by CD4 and Foxp3

expression in young and aged mouse spleens (n=15-32, unpaired two-tailed Student's t test after

arcsin conversion). **C**) Flow cytometric plot and **D**) quantification showing the proportions of natural

Treg (endogenous GFP reporting Foxp3 expression in Foxp3-DTR mice) in aged blood compared to

young circulation (n=30-38 mice, unpaired two-tailed Student's t test after *arcsin* conversion). **E**) Bar

graph showing the density of OLIG2$^+$CC1$^+$ oligodendrocytes in young and aged lysolecithin-induced

demyelinating lesions at 14dpl (n=4 mice, unpaired Student's t test). **F**) Flow cytometric plot showing

CD25 and Foxp3 expression and the quantification of CD25$^+$Foxp3$^+$ natural Treg proportion in a CD4$^+$

T cell population from the spleen (**G**) and blood (**H**) of control aged mice and aged mice treated with

intraperitoneal injection of IL-2/anti-IL-2 complexes (n=3, unpaired two-tailed Student's t test after

arcsin conversion). **I**) Bar graph showing the quantification of OLIG2$^+$CC1$^+$ oligodendrocyte density at

10dpl in aged control and Treg-expanded mice (n=6-7, unpaired Student's t test).

Figure 2: Aged natural Treg demonstrate impaired capacity to drive OPC differentiation and

myelination *in vitro*. Representative images of OPCs co-cultured with young and aged natural Treg

and immunostained for OLIG2 (cyan) as a pan oligodendrocyte lineage marker, the proliferation

marker Ki67 (magenta, **A**) and differentiation markers CNP (grey) and MBP (magenta) (**C**) (Scale bar =

100 μm). **B**) Quantification of OPC proliferation when co-cultured with young and aged natural Treg

(n=6, 1-way ANOVA after *arcsin* conversion) **D**) Quantification of the proportion of OPCs reaching early

stage-differentiation when exposed to young and aged natural Treg, as indicated by CNPase staining

(n=9, 1-way ANOVA after *arcsin* conversion). **E**) Quantification of the proportion of OPCs expressing

late-stage differentiation marker MBP in control and OPCs treated with young and aged natural Treg

(n=9, 1-way ANOVA, Sidak's multiple comparisons test). **F**) Immunohistochemistry of control and

young and aged Treg treated cerebellar slices (MBP, magenta and NFH, green, Scale bar = 100 μm).

(**G**) Quantification of myelination index (ratio between MBP and NFH colocalization area and NFH

area) in neonatal cerebellar slices (n=7, 1-way ANOVA after *arcsin* conversion, Sidak's multiple

comparison tests).

Figure 3: Myelin-regenerative capacity of aged Treg is restored in a young environment. A) Diagram

explaining the experimental design of *in vivo* Treg depletion, Treg adoptive transfer and spinal cord

demyelination. Representative images of immunostaining identifying oligodendrocytes by the co-

localisation of the pan-oligodendrocyte lineage marker OLIG2 (green) with CC1 (magenta, **B**) or ASPA

(magenta, **C**) at 14 dpl (scale bar = 100 μm, demyelination area is highlighted by the white line). Bar

graphs show the quantification of total number of oligodendrocyte lineage cells (**D**), as well as CC1-

expressing (**E**) or ASPA-expressing (**F**) oligodendrocytes in the demyelinated lesions of PBS control

mice, natural Treg-depleted mice and mice depleted of endogenous Treg that received young or aged

natural Treg by adoptive transfer (n=7-9, 1-way ANOVA, Sidak's multiple comparisons test). **G**)

Representative images of immunostaining for neurofilament-H (NFH, green) and MBP (magenta) to

quantify myelin wrapping as an early marker of remyelination at 14 dpl (scale bar = 100 μm,

demyelination area is highlighted by the white line). Quantification shows the total number of axons

(**H**), the density of MBP wrapped axons (**I**) and the percentage of MBP-wrapped axons from the total

number of axons (**J**) (n=6-9, 1-way ANOVA, Sidak's multiple comparisons test).

Figure 4: Ageing significantly alters natural Treg transcriptome. A) Principal component analysis

demonstrating the clustering differences between young and aged natural Treg. **B**) Volcano plot

demonstrating 1456 genes upregulated and 302 genes downregulated genes in aged natural Treg

relative to young Treg. **C**) Heatmap demonstrating hierarchical clustering of the top 50 differentially

expressed genes between young and aged Treg. **D**) Graph showing the pathways enriched amongst

differentially expressed genes. **E**) Bar graph highlighting the GO biological processes associated with

genes that are upregulated or downregulated in aged Treg. **F**) Heatmap showing the normalised count

values for the Treg tissue repair programme identified by Delacher *et al.*[23] **G**) Graph showing the Treg

tissue repair programme signature genes score in aged and young natural Treg.

Figure 5: *Mcam* and *Itga2* contribute to Treg-driven OPC differentiation and are downregulated in

aged Treg. A) Representative images of immunostaining showing cell-to-cell contact between OPCs

and Treg in OPC-Treg co-cultures *in vitro*. OPCs are identified by the co-staining of OLIG2 (cyan) and

NG2 (grey), while Treg are identified by CD3 (red) (scale bar = 50 μm). **B**) Representative images of

immunostaining and **C**) quantification of MBP-expressing oligodendrocytes in control OPCs, OPCs

directly co-cultured with young Treg and OPCs cultured with young Treg in a transwell (n=6, 1-way

ANOVA after arcsin conversion, Sidak's multiple comparison's test). **D**) Diagram summarising

bioinformatic approaches to identify protein-protein interactions between OPCs and Treg. **E**) Graph

showing 21 protein candidates expressed in the Treg plasma membrane, that are downregulated in

aged Treg and have potential bindings partners enriched in OPCs vs oligodendrocytes. Log_2 Change, -

Log_{10} (Padj) and the number of OPC binding partners are indicated (see legend). **F**) Bar graphs showing

RNAseq normalised count values for the top 6 candidates. **G**) Representative images of

immunostaining showing OPC differentiation in co-culture with young Treg in the presence or absence

of neutralising antibodies against candidate cell surface mediators (scale bar = 100µm). Bar graphs

showing the quantification of OPC differentiation measured by the fold change in percentage of MBP⁺

cells (**H**) and MBP⁺ area per well (**I**) (n=7, 2-way ANOVA, Dunnett's multiple comparison tests).

Supplementary information

Supplementary Figure 1: Endogenous Treg depletion and reconstitution during lysolecithin- induced

demyelination *in vivo*. **A**) Gating strategy followed to identify natural Treg. **B**) Bar graph showing

natural Treg isolation purity for *in vitro* experiments and *in vivo* adoptive transfer studies. Natural Treg

are identified by CD4 and CD25 expression (average 91% of the isolated cells for both young and aged

are CD4⁺CD25⁺) and by the expression of CD4 and Foxp3 (average 87% of the isolated cells are positive

for CD4 and Foxp3 in both young and aged mouse isolations) (n=14 independent isolations for

different adoptive transfer rounds, each pool comprises 2-8 different mice, Student's t test, after

arcsin conversion). **C**) Flow cytometric plot showing CD4 and GFP expression to identify endogenous

natural Treg in lymph nodes. **D**) Bar graph showing quantification for the proportion of endogenous

GFP⁺ Treg within the CD4⁺ T cell population in the lymph nodes of young Foxp3-DTR mice at 14dpl

(n=7-9, 1-way ANOVA, after *arcsin* conversion, Sidak's multiple comparisons test). **E**) Flow cytometric

plots showing CD25 and Foxp3 expression to detect Treg reconstitution in lymph nodes after adoptive

transfer. **F**) Bar graph quantification shows the proportion of CD25⁺Foxp3⁺ cells within the CD4⁺ T cell

population present in the lymph nodes of young Foxp3-DTR mice at 14dpl (n=7-12, Kruskal Wallis and

Dunn's multiple comparisons test). **G**) Representative images of immunostaining showing proliferating

OPCs at 14 dpl (OLIG2 (green) and Ki67 (magenta)) (scale bar = 100 µm, demyelinated area is indicated

by the white line). Bar graph showing the areas of demyelination (**H**), the density of proliferating OPCs

in the lesion (**I**) and the proportion of oligodendrocyte lineage cells proliferating in response to

damage (**J**) (n=7-8 mice, 1 way ANOVA).

Acknowledgements

We acknowledge extensive technical support from Carmel McVicar, Maria P Athanasios and the staff of the animal facility. We thank A. Rudensky (Memorial Sloan Kettering Cancer Centre) for providing Foxp3-DTR mice. This work was supported by the Wellcome Trust (110138/Z/15/Z to DCF), Biotechnology and Biological Sciences Research Council (BB/J01026X/1 and BB/N003721/1, to DCF), ECTRIMS postdoctoral fellowship (to AGF), Wellcome ISSF fellowship (to AGF), Miguel Servet Fellowship from the Spanish Institute of Health Carlos III (CP21/00032 to AGF), The Leverhulme Trust (ECF-2014-390, to YD) and postgraduate studentship support from the Dept. for the Economy (Northern Ireland). We thank the laboratory of Anna Williams (University of Edinburgh) for the Fiji plug-in to analyse myelination index in brain slice cultures.

Author contributions

Experiments were designed by A.G.F., D.C.F., Y.D. and V.T. Experiments were performed and/or

analysed by A.G.F., M.D., E.H., N.V.G., J.W., A.Y., K.M., J.F., C.E.M., M.I., R.P., Y.D. and D.C.F.

R.J.I. and C.E.M. provided advice on experimental design and interpretation. Manuscript was written

by A.G.F. and D.C.F. with contributions from all authors. D.C.F and A.G.F. oversaw the study.

References

1. Nave, K.-A. & Werner, H. B. Ensheathment and Myelination of Axons: Evolution of Glial Functions. *Annu Rev Neurosci* **44**, 1–23 (2021).

2. Griffiths, I. *et al.* Axonal swellings and degeneration in mice lacking the major proteolipid of myelin. *Science* **280**, 1610–1613 (1998).

3. Lappe-Siefke, C. *et al.* Disruption of Cnp1 uncouples oligodendroglial functions in axonal support and myelination. *Nat Genet* **33**, 366–374 (2003).

4. Love, S. Demyelinating diseases. *J Clin Pathol* **59**, 1151 (2006).

5. Bodini, B. *et al.* Dynamic Imaging of Individual Remyelination Profiles in Multiple Sclerosis. *Ann Neurol* **79**, 726–738 (2016).

6. Franklin, R. J. M. & ffrench-Constant, C. Regenerating CNS myelin — from mechanisms to experimental medicines. *Nat Rev Neurosci* **18**, 753–769 (2017).

7. Lubetzki C. *et al*. Remyelination in multiple sclerosis: from basic science to clinical translation. *The Lancet Neurol* **19**, 678–688 (2020).

8. Shields, S., Gilson, J., Blakemore, W. & Franklin, R. Remyelination occurs as extensively but more slowly in old rats compared to young rats following fliotoxin-induced CNS demyelination. *Glia* **29**, 102 (2000).

9. Sim, F. J., Zhao, C., Penderis, J. & Franklin, R. J. M. The age-related decrease in CNS remyelination efficiency is attributable to an impairment of both oligodendrocyte progenitor recruitment and differentiation. *J Neurosci* **22**, 2451–2459 (2002).

10. Confavreux, C. & Vukusic, S. Age at disability milestones in multiple sclerosis. *Brain* **129**, 595–605 (2006).

11. Burzyn, D. *et al.* A special population of regulatory T cells potentiates muscle repair. *Cell* **155**, 1282–1295 (2013).

12. Ali, N. *et al.* Regulatory T Cells in Skin Facilitate Epithelial Stem Cell Differentiation. *Cell* **169**,
1119-1123.e11 (2017).

13. Dombrowski, Y. *et al.* Regulatory T cells promote myelin regeneration in the central nervous
system. *Nat Neurosci* **20**, 674–680 (2017).

14. McIntyre, L. L. *et al.* Regulatory T cells promote remyelination in the murine experimental
autoimmune encephalomyelitis model of multiple sclerosis following human neural stem cell
transplant. *Neurobiol Dis* **140**, 104868 (2020).

15. Shi, L. *et al.* Treg cell-derived osteopontin promotes microglia-mediated white matter repair
after ischemic stroke. *Immunity* **54**, 1527-1542.e8 (2021).

16. Garg, S. K. *et al.* Aging is associated with increased regulatory T-cell function. *Aging Cell* **13**, 441–
448 (2014).

17. Elyahu, Y. *et al.* Aging promotes reorganization of the CD4 T cell landscape toward extreme
regulatory and effector phenotypes. *Sci adv* **5**, eaaw8330 (2019).

18. Neumann, B. *et al.* Metformin Restores CNS Remyelination Capacity by Rejuvenating Aged Stem
Cells. *Cell Stem Cell* **25**, 473-485.e8 (2019).

19. Dittmer, M. *et al.* Characterization of a murine mixed neuron-glia model and cellular responses
to regulatory T cell-derived factors. *Mol brain* **11**, 25 (2018).

20. Kim, J. M., Rasmussen, J. P. & Rudensky, A. Y. Regulatory T cells prevent catastrophic
autoimmunity throughout the lifespan of mice. *Nat Immunol* **8**, 191–197 (2007).

21. Guo, Z. *et al.* DCAF1 regulates Treg senescence via the ROS axis during immunological ageing. *J*
Clin Invest **130**, 5893–5908 (2020).

22. Morales-Nebreda, L. *et al.* Aging imparts cell-autonomous dysfunction to regulatory T cells
during recovery from influenza pneumonia. *JCI Insight* **6**, e141690 (2021).

23. Delacher, M. *et al.* Single-cell chromatin accessibility landscape identifies tissue repair program in
human regulatory T cells. *Immunity* **54**, 702-720 (2021).

24. Zhang, Y. *et al.* An RNA-sequencing transcriptome and splicing database of glia, neurons, and
vascular cells of the cerebral cortex. *J Neurosci* **34**, 11929–11947 (2014).

25. Jensen, L. J. *et al.* STRING 8--a global view on proteins and their functional interactions in 630
organisms. *Nucleic Acids Research* **37**, D412-6 (2009).

26. Natrajan, M. S. *et al.* Retinoid X receptor activation reverses age-related deficiencies in myelin
debris phagocytosis and remyelination. *Brain* **138**, 3581–3597 (2015).

27. Linehan, E. *et al.* Aging impairs peritoneal but not bone marrow-derived macrophage
phagocytosis. *Aging cell* **13**, 699–708 (2014).

28. Cantuti-Castelvetri, L. *et al.* Defective cholesterol clearance limits remyelination in the aged
central nervous system. *Science* **359***, 684-688* (2018).

29. Segel, M. *et al.* Niche stiffness underlies the ageing of central nervous system progenitor cells.
Nature **350**, 1199–134 (2019).

30. Wang, W., Thomas, R., Oh, J. & Su, D. Accumulation of pTreg cells is detrimental in late-onset
(aged) mouse model of multiple sclerosis. *Aging Cell* **21**, e13630 (2022).

31. Kuswanto, W. *et al.* Poor Repair of Skeletal Muscle in Aging Mice Reflects a Defect in Local,
Interleukin-33-Dependent Accumulation of Regulatory T Cells. *Immunity* **44**, 355–367 (2016).

32. Muñoz-Rojas, A. R. & Mathis, D. Tissue regulatory T cells: regulatory chameleons. *Nat Rev*
Immunol **21**, 597-611 (2021).

33. Breuer, J. *et al.* Blockade of MCAM/CD146 impedes CNS infiltration of T cells over the choroid
plexus. *J Neuroinflamm* **15**, 236 (2018).

34. Zondler, L. *et al.* MCAM/CD146 Signaling via PLCγ1 Leads to Activation of β1-Integrins in Memory
T-Cells Resulting in Increased Brain Infiltration. *Front Immunol* **11**, 599936 (2020).

35. Larochelle, C. *et al.* Melanoma cell adhesion molecule–positive CD8 T lymphocytes mediate
central nervous system inflammation. *Ann Neurol.* **78**, 39–53 (2015).

36. ACTRIMS Forum 2020 - Poster Session 1. Poster 4. Zandee, S.E. *et al.* Can T Cell Migratory
Signature Predict MS Disease Progression? *Mult Scler J* **26**, 16–89 (2020).

37. Laursen, L. S., Chan, C. W. & ffrench-Constant, C. An Integrin-Contactin Complex Regulates CNS
Myelination by Differential Fyn Phosphorylation. *J Neurosci* **29**, 9174–9185 (2009).

38. Ruckh, J. M. *et al.* Rejuvenation of regeneration in the aging central nervous system. *Cell stem*
cell **10**, 96–103 (2012).

39. Webster, K. E. *et al.* In vivo expansion of T reg cells with IL-2–mAb complexes: induction of
resistance to EAE and long-term acceptance of islet allografts without immunosuppression. *J Exp*
Medicine **206**, 751–760 (2009).

40. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**,
676–682 (2012).

41. Zhang, H., Jarjour, A. A., Boyd, A. & Williams, A. Central nervous system remyelination in culture
— A tool for multiple sclerosis research. *Experimental Neurology* **230**, 138–148 (2011).

42. Stirling, D. R. *et al.* CellProfiler 4: improvements in speed, utility and usability. *Bmc Bioinformatics*
22, 433 (2021).

43. Dao, D. *et al.* CellProfiler Analyst: interactive data exploration, analysis and classification of large
biological image sets. *Bioinformatics* **32**, 3210–3212 (2016).

44. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
Embnet J **17**, 10–12 (2011).

45. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

46. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2013).

47. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

Figure 1: Expansion of Treg does not rescue impaired OPC differentiation in aged mice. A) Flow cytometric plot and **B**) quantification of natural Treg proportion, identified by CD4 and Foxp3 expression in young and aged mouse spleens (n=15-32, unpaired two-tailed Student's t test after *arcsin* conversion). **C**) Flow cytometric plot and **D**) quantification showing the proportions of natural Treg (endogenous GFP reporting Foxp3 expresion in Foxp3-DTR mice) in aged blood compared to young circulation (n=30-38, unpaired two-tailed Student's t test). **E**) Bar graph showing the density of OLIG2$^+$CC1$^+$ oligodendrocytes in young and aged lysolecithin-induced demyelinating lesions at 14dpl (n=4 mice, unpaired Student's t test). **F**) Flow cytometric plot showing CD25 and Foxp3 expression and the quantification of CD25$^+$Foxp3$^+$ natural Treg proportion in a CD4$^+$ T cell population from the spleen (**G**) and blood (**H**) of control aged mice and aged mice treated with intraperitoneal injection of IL-2/anti-IL-2 complexes (n=3, unpaired two-tailed Student's t test after arcsin c conversion). **I**) Bar graph showing the quantification of OLIG2$^+$CC1$^+$ oligodendrocyte density at 10dpl in aged control and Treg expanded mice (n=6-7, unpaired Student's t test).

Fig. 1

Figure 2: Aged natural Treg demonstrate impaired capacity to drive OPC differentiation and myelination *in vitro*. Representative images of OPCs co-cultured with young and aged nTreg and immunostained for OLIG2 (cyan) as a pan oligodendrocyte lineage marker, the proliferation marker Ki67 (magenta, **A**) and differentiation markers CNP (grey) and MBP (magenta) (**C**) (Scale bar = 100 µm). **B**) Quantification of OPC proliferation when co-cultured with young and aged nTreg (n=6, 1-way ANOVA after arcsin conversion) **D**) Quantification of the proportion of OPCs reaching early stage-differentiation when exposed to young and aged nTreg, as indicated by CNPase staining (n=9, 1-way ANOVA after arcsin conversion). **E**) Quantification of the proportion of OPCs expressing late-stage differentiation marker MBP in control and OPCs treated with young and aged nTreg (n=9, 1-way ANOVA, Sidak's multiple comparisons test). **F**) Immunohistochemistry of control and young and aged nTreg treated cerebellar slices (MBP, magenta and NFH, green, Scale bar = 100 µm). (**G**) Quantification of myelination index (ratio between MBP and NFH colocalization area and NFH area) in neonatal cerebellar slices (n=7, 1-way ANOVA after arcsin conversion, Sidak's multiple comparison tests). .

Fig. 2

Fig. 3

Figure 3: Myelin-regenerative capacity of aged Treg is restored in a young environment. A) Diagram explaining the experimenta design of *in vivo* Treg depletion, Treg adoptive transfer and spinal cord demyelination. Representative iamges of immunostaining identifying oligodendrocytes by the co-localisation of the pan-oligodendrocyte lineage marker OLIG2 (green) with CC1 (magenta, **B**) or ASPA (magenta, **C**) at 14dpl (scale bar=100µm, demyelinated area is highlighted by the white line). Bar graphs show the quantification of total number of oligodendrocyte lineage cells (**D**), as well as CC1-expressing (**E**) or ASPA-expressing (**F**) oligodendrocytes in the demyelinated lesions of PBS control mice, natural Treg-depleted mice and mice depleted of endogenous Treg that received young or aged natural Treg by adoptive transfer (n=7-9, 1-way ANOVA, Sidak's multiple comparisons test). **G**) Representative images of immunostaining for neurofilament-H (NFH, green) and MBP (magenta) to quantifiy myelin wrapping as an early marker of remyelinationg at 14dpl (scale bar = 100µm, demyelination area is highlighted by the white line). Quantification shows the total number of axons (**H**), the density of MBP wrapped axons (**I**) and the percentage of MBP-wrapped axons from the total number of axons (**J**) (n=6-9, 1-way ANOVA, Sidak's multiple comparisons test).

Fig. 4

Figure 4: Ageing significantly alters natural Treg transcriptome. A) Principal component analysis demostrating the clustering differences bedtween young and aged natural Treg. **B**) Volcano plot demostrating 1456 genes upregulated and 302 genes genes downregulated genes natural Treg relative to young Treg. **C**) Heatmap demonstrating hierarchical clustering of the top 50 differentially expressed genes between young and aged Treg. **D**) Graph showing the pathways enriched amongst differntially expressed genes. **E**) Bar graph highlighting the GO biological processes associated with genes that are upregulated or downregulated in aged Treg. **F**) Heatmap showing the normalised count values for the Treg tissue repair programme identified by Delacher *et al*.[23]. **G**) Graph showing the Treg tissue repair programme signature genes score in aged and young natural Treg.

Fig. 5

Figure 5: Mcam and Itga2 contribute to Treg-driven OPC differentiation and are downregulated in aged Treg.
A) Representative images of immunostaining showing cell-to-cell contact between OPCs and Treg in OPC-Treg co-cultures *in vitro*. OPCs are identified by the co-staining of OLIG2 (cyan) and NG2 (grey), while Treg are identified by CD3 (red) (scale bar = 50 μm). **B**) Representative images of immunostaining and **C**) quantification of MBP-expressing oligodendrocytes in control OPCs, OPCs directly co-cultured with young Treg and OPCs cultured with young Treg in a transwell (n=6, 1-way ANOVA after arcsin conversion, Sidak's multiple comparison's test). **D**) Diagram summarising bioinformatic approaches to identify protein-protein interactions between OPCs and Treg. **E**) Graph showing 21 protein candidates expressed in the nTreg plasma membrane, that are downregulated in aged nTreg and have potential bindings partners enriched in OPCs vs oligodendrocytes. Log_2 Change, $-Log_{10}$ (Padj) and the number of OPC binding partners are indicated (see legend). **F**) Bar graphs showing RNAseq normalised count values for the top 6 candidates. **G**) Representative images of immunostaining showing OPC differentiation in co-culture with young Treg in the presence ofr absence of neutralising antibodies against candidate cell surface mediators (scale bar=100μm). Bar graphs showing the quantification of OPC differentiation measured by the fold change in percentage of MBP$^+$ cells (**H**) and MBP$^+$ area per well (**I**) (n=7, 2-way ANOVA, Dunnett{s multiple comparsion tests).