Ageing impairs the regenerative capacity of regulatory T cells in central nervous system

remyelination

Alerie Guzman de la Fuente[1,2,3*], Marie Dittmer[1], Elise Heesbeen[1,4], Nira de la Vega Gallardo[1], Jessica

White[3], Andrew Young[1,5], Tiree McColgan[1], Amy Dashwood[6,7], Katie Mayne[1,8], Sonia Cabeza-

Fernández[2,3], John Falconer[1,9], Francisco Javier Rodriguez-Baena[3], Christopher E. McMurran[10,11],

Mohammed Innayatullah[12,13], Khalil S. Rawji[14], Robin J.M. Franklin[14], James Dooley[6,7], Adrian Liston[6,7],

Rebecca Ingram[1], Vijay K. Tiwari[1,12,13,15], Rosana Penalva[1], Yvonne Dombrowski[1] and Denise C.

Fitzgerald[1*]

[1]Wellcome-Wolfson Institute for Experimental Medicine, Queen's University Belfast, Belfast, BT9

7BL, Northern Ireland, UK.

[2]Institute for Health and Biomedical Sciences of Alicante (ISABIAL), Alicante, 03010, Spain.

[3]Instituto de Neurosciencias CSIC-UMH, San Juan de Alicante, 03550, Spain.

[4]Current address: Division of Pharmacology, Institute for Pharmaceutical Sciences, Utrecht

University, Utrecht, The Netherlands.

[5]Current address Patrick G Johnston Centre for Cancer Research, Queen's University Belfast, Belfast,

BT9 7BL, Northern Ireland, UK

[6]Department of Pathology, University of Cambridge, CB2 1QP, Cambridge, UK

[7]Babraham Institute, CB22 3AT, Cambridge, UK

[8]Current address: Department of Pathology and Laboratory Medicine, University of British Columbia,

G227-2211 Wesbrook Mall, Vancouver, BC, V6T 2B5, Canada.

[9]Current address: CRUK Beatson Institute, G61 1BD, Glasgow, UK

[10]Department of Clinical Neurosciences, University of Cambridge, Cambridge, UK

[11]Department of Medical Epidemiology and Biostatistics, Karolinska Institute, Stockholm, Sweden

[12]Institute of Molecular Medicine, University of Southern Denmark, 5000 Odense, Denmark

[13]Danish Institute for Advanced Study (DIAS), 5230 Odense, Denmark

[14]Altos Labs - Cambridge Institute of Science, Granta Park, Cambridge CB21 6GP, UK

[15] Department of Clinical Genetics, Odense University Hospital, 5000 Odense, Denmark

*Corresponding authors:

Denise C. Fitzgerald, d.fitzgerald@qub.ac.uk

Alerie G. de la Fuente, guzman_ale@isabial.es

Abstract

Myelin regeneration (remyelination) is essential to prevent neurodegeneration in

demyelinating diseases such as Multiple Sclerosis, however, its efficiency declines with age.

Regulatory T cells (Treg) recently emerged as critical players in tissue regeneration, including

remyelination. However, the effect of ageing on Treg-mediated regenerative processes is poorly

understood. Here, we show that expansion of aged Treg does not rescue age-associated

remyelination impairment due to an intrinsically diminished capacity of aged Treg to promote

oligodendrocyte differentiation and myelination. This decline in regenerative Treg functions can be

rescued by a young environment. We identified Melanoma Cell Adhesion Molecule 1 (MCAM1) and

Integrin alpha 2 (ITGA2) as novel candidates of Treg-mediated oligodendrocyte differentiation that

decrease with age. Our findings demonstrate that ageing limits the neuroregenerative capacity of

Treg, likely limiting their remyelinating therapeutic potential in aged patients, and describe two

novel mechanisms implicated in Treg-driven remyelination that may be targetable to overcome this

limitation.

Introduction

Myelin is critical to the metabolic support and function of axons in the central nervous

system (CNS)[1]. CNS myelin is produced by oligodendrocytes and damage to oligodendrocytes or

myelin causes neurological impairment[2-4]. Myelin regeneration (remyelination) is highly

neuroprotective and holds potential to restore lost function for patients with demyelinating diseases

such as Multiple Sclerosis (MS)[5-8]. Like most regenerative processes, remyelination efficiency

declines with age[9-11], contributing to age-associated disease progression and the accumulation of

irreversible disability[12]. Age-associated remyelination decline has been linked to intrinsic changes in

the capacity of oligodendrocyte progenitor cells (OPC) to respond to damage in the aged CNS[13-15],

changes in physical properties and the microenvironment (e.g extracellular matrix) of the aged CNS

which makes it refractory to remyelination[16,17] and impaired capacity of other cells involved in the

process of myelin regeneration (e.g macrophages, microglia, astrocytes) to drive this process [18-23].

Despite the critical neuroprotective role of remyelination[8,24], there are currently no approved

remyelination-enhancing therapies, a key unmet medical need. Thus, understanding how ageing

alters the different cellular mechanisms governing myelin regeneration is key to overcoming age-

associated remyelination failure. This holds potential to prevent neuronal loss and the accumulation

of irreversible cognitive, sensory and motor disabilities.

Regulatory T cells (Treg) have recently emerged as key cellular players in tissue regeneration

due to both anti-inflammatory functions and capacity to modulate tissue-resident stem cell

differentiation[25,26]. Similar crosstalk has also been described in the CNS, where we and others

showed that Treg promote oligodendrocyte progenitor cell (OPC) differentiation and CNS

remyelination[27-29]. The discovery of pro-remyelinating functions of Treg opened a new therapeutic

avenue to prevent remyelination decline and associated neurodegeneration. However, whether this

important neuroregenerative function of Treg is impaired with age remains unknown.

Understanding how ageing affects myelin-regenerative functions of Treg is essential to determine

the relevance and suitability of potential Treg-based pro-remyelination therapies in patients of

advanced age in MS and other demyelinating diseases.

Here, we show that ageing impairs the capacity of natural Treg to drive OPC differentiation

in vitro and myelination *ex vivo*. Surprisingly, this defect was reversible in a young remyelinating

environment *in vivo*. Transcriptomic comparison of young and aged Treg identified Melanoma Cell

Adhesion Molecule 1 (MCAM1) and Integrin alpha 2 (ITGA2) as novel candidates of Treg-mediated

OPC differentiation that decrease with age. This study identifies Treg-driven OPC differentiation as a

remyelination mechanism that is impaired by ageing and identifies novel Treg-expressed candidates

(ITGA2, MCAM), which enhance OPC differentiation and should be further investigated to determine

its potential relevance as therapeutic targets to promote CNS remyelination.

Results

Tissue-resident Treg are increased in the aged CNS

We first examined the frequency of murine Treg with age, and in agreement with previous studies[30,31], found increased numbers and proportions of Treg in aged mice compared to young mice in spleen and blood (**Fig. 1 A-E**). Next, we investigated whether ageing impacted T cell populations in the CNS. As described previously[32–34], we observed an increase in the proportion of CD8[+] and CD4[+] T cells in aged murine CNS tissue with significant differences observed more prominently in spinal cord than brain (**Sup. Fig. 1A-H**). We also observed a significant increase in the proportion of Treg in the spinal cord (**Fig. 1 F, G**) and a non-significant trend in the brain (**Fig. 1 I, J**). Proportions of tissue-resident (CD69[+]) Treg[35] were increased both in the aged brain and spinal cord (**Fig. 1 H, K**). Previous studies showed that ageing impairs migration of Treg to damaged muscle[36]. To determine if ageing affects the capacity of Treg to accumulate in demyelinated lesions in the CNS we used two models of demyelination, lysolecithin-induced demyelination, and experimental autoimmune encephalomyelitis (EAE), a model of immune-mediated demyelination. Despite the low frequency of Treg in demyelinated areas, we did not observe any differences in Treg density in either of the two models at the time points investigated (**Fig. 1L-N**). This suggests that aged Treg have a similar capacity to accumulate in demyelinated CNS tissue when compared to young Treg although differences at other phases of CNS damage cannot be conclusively ruled out.

Expansion of Treg does not rescue impaired OPC differentiation in aged mice

We next verified that oligodendrocytes numbers were comparable in the young and aged healthy CNS (**Sup. Fig. 2 A-D**)[37,38], but that OPC differentiation into CC1[+] oligodendrocytes was impaired in aged demyelinated lesions *in vivo* as previously reported[10,13] (**Sup. Fig. 2 E, F**). We next investigated whether expanding Treg populations in aged mice would recover oligodendrocyte differentiation. Administration of IL-2/anti-IL-2 *in vivo* significantly expanded systemic CD25[+]Foxp3[+] Treg (~3-5-fold increase in blood and spleen) (**Fig 2 A-C**) with a trend of higher densities of Treg in

demyelinated lesions of Treg-expanded mice (**Sup. Fig. 2 G-J**) which was not statistically significant

although numbers were low. We first examined whether Treg expansion enhanced the

remyelination response in young mice. As myelin regeneration is highly efficient in young rodents[9,10],

and our previous findings showed that Treg accelerated oligodendrocyte differentiation[27], we

examined oligodendrocyte differentiation at a comparatively early time point of 7 days-post-lesion

(dpl), when OPC differentiation is just starting in this model[39]. While there were no changes in

overall Olig2$^+$ cell density, Treg expansion significantly increased the number of Olig2$^+$CC1$^+$

differentiated oligodendrocytes in young mice (**Fig. 2D-F**). Next, we investigated whether Treg

expansion accelerate OPC differentiation in aged mice. Taking into account that remyelination is

delayed in aged mice, we first examined oligodendrocyte differentiation at 10dpl as a relevant time

point at which OPC differentiation has started in aged mice. Contrary to young mice, Treg expansion

did not significantly change the density of Olig2$^+$ cells or Olig2$^+$CC1$^+$ differentiated oligodendrocytes

in aged mice at 10dpl (**Fig. 2 G-I**). For a direct comparison to the timepoint in young mice, we also

examined oligodendrocyte differentiation at 7dpl in aged mice and observed no differences (data

not shown). We also evaluated neurofilament (NFH) and myelin basic protein (MBP) colocalization as

a surrogate measure of remyelination. As for OPC differentiation, Treg expansion did not enhance

axon wrapping in aged mice (**Fig. 2 J-L**). Hence, we hypothesised that the myelin-regenerative

capacity of Treg may be intrinsically impaired in aged Treg, irrespective of Treg population

frequency.

Aged Treg demonstrate an impaired intrinsic capacity to enhance myelin production

To compare the capacity of young and aged Treg to enhance OPC differentiation and

myelination, we first investigated effects on oligodendrocyte development. To do this, we isolated

natural Treg from the spleens and lymph nodes of young (3-4 months) and aged (14-18 months)

mice (**Sup. Fig. 3 A, B**) and co-cultured these Treg with neonatal murine OPCs. While neither young

nor aged Treg modified OPC proliferation *in vitro* (**Fig. 3 A, B**), young Treg significantly increased OPC

differentiation into MBP-expressing oligodendrocytes, in agreement with our previous findings of

Treg secretome[27,40]. Aged Treg however, did not enhance OPC differentiation demonstrating an

impairment in their capacity to support this necessary step in the process of myelin formation (**Fig. 3**

C-E).

146   To determine if the inability of aged Treg to support OPC differentiation was functionally

relevant to the process of myelin production, we compared the capacity of young and aged Treg to

drive myelin ensheathment of axons *ex vivo*. To do this, we co-cultured murine organotypic brain

slices with young or aged Treg for 7 days. Young Treg significantly enhanced axonal myelination

compared to control slices as described before[27]. However, aged Treg failed to significantly increase

myelination compared to controls (**Fig. 3 G**). In contrast to our observations in OPC differentiation,

we did not detect significant differences between young and aged Treg when evaluating myelination

index (**Fig. 3G**) but on balance, these *in vitro* and *ex vivo* data suggest that aged Treg fail to

significantly drive myelin production.

Myelin-regenerative capacity of aged Treg is restored in a young environment

157   We previously reported that Treg depletion using diphtheria toxin (DT) in B6.129(Cg)-

Foxp3[tm3(DTR/GFP)Ayr]/J (FoxP3-DTR)[41] mice impaired OPC differentiation and remyelination, which was

rescued by the administration of wild type Treg[27]. Thus, we next compared the regenerative capacity

of young and aged Treg in a model of lysolecithin-induced spinal cord myelin damage *in vivo*.

Following endogenous GFP[+] Treg depletion (**Fig. 4 A**, **Sup. Fig. 3 C, D**), we administered wild type

(GFP[-]) young or aged Treg intraperitoneally and confirmed comparable detection of adoptively

transferred Treg in each group (**Sup. Fig. 3 E, F**). We next compared the neuroregenerative capacity

of adoptively transferred young and aged wild type Treg to enhance OPC differentiation during

remyelination (**Fig. 4 A**). We did not observe any differences in the burden of damage for each

experimental group, indicated by the lesion area (**Sup. Fig. 3 H**). There was also no difference in the

number of oligodendrocyte lineage cells, quantified by Olig2 immunostaining (**Fig. 4 B-D**), or

proliferating OPCs (Olig2$^+$Ki67$^+$) at 14 days post-lesion (dpl) (**Sup. Fig. 3 G, I, J**). As before[27], Treg

depletion impaired OPC differentiation into CC1$^+$ and ASPA$^+$ oligodendrocytes, which was rescued by

adoptive transfer of wild type young Treg. Unexpectedly however, aged Treg injected into young

Treg-depleted mice also rescued impaired OPC differentiation, comparably to young Treg, (**Fig. 4 B,**

C, E, F). To determine if aged Treg not only rescued OPC differentiation but also restored

remyelination, we examined axonal wrapping through the quantification of circular MBP$^+$ patterns

around NFH$^+$ axons as an early indication of axonal remyelination. Both, young and aged Treg

administered to Treg-depleted mice, restored axonal wrapping to levels comparable to non-depleted

controls (**Fig. 4 G-J**). These findings suggest that the age-induced defect of neuroregenerative Treg

functions observed *in vitro/ex vivo* is reversible in a young systemic environment *in vivo*, and

therefore, potentially therapeutically targetable.

Young Treg partially rescue OPC differentiation during remyelination in aged mice

Next, we investigated whether young Treg could rescue impaired remyelination in aged mice

using Foxp3-DTR mice in which GFP$^-$ young and aged Treg were adoptively transfer following DT-

mediated Treg depletion. In contrast to our findings in young Treg-depleted mice, aged mice without

Treg showed decreased Olig2$^+$ cell density in the demyelinating area which was not rescued neither

by young nor by aged Treg (**Fig. 5 A, B**). As per young mice, Treg depletion did not impair OPC

proliferation (**Fig. 5 C**), suggesting that the decrease in Olig2$^+$ cells likely reflects either an impaired

survival or migration of oligodendrocyte lineage cells to the damaged area. The decreased density of

Olig2$^+$CC1$^+$ oligodendrocytes in the absence of Treg was partially rescued by the transfer of young

Treg into these animals but not by transferred aged Treg, which were significantly different to the

non-depleted control (**Fig. 5 A, C**). However, it is important to highlight that we did not detect

statistically significant differences between aged Treg depleted mice receiving young or aged Treg

(**Fig. 5 A, D**) and thus, we interpret these findings with caution. A further reason to be cautious in

interpreting these results is that contrary to young recipients, adoptively transferred Treg were not

detectable in the classic lymphoid organs (spleen, lymph nodes) of aged Foxp3-DTR mice (**Sup**. **Fig. 4**

E, F), suggesting that the adoptively transferred cells have a different migration pattern in aged

recipients, and thus we cannot be sure that young and aged cells have the same reconstitution

capacity or distribution. Additionally, since young and aged Treg failed to restore Olig2$^+$ cell density

in aged mice (**Fig. 5B**), the age-associated remyelination impairment, even when Treg are expanded,

is associated not only with impaired Treg regenerative capacity but also with a refractory aged CNS

environment that may limit OPC responsiveness to Treg-derived cues, as shown for other factors

previously[13].

Ageing significantly alters Treg transcriptome

We next sought to identify how ageing impairs the myelin regenerative functions of Treg by

examining gene expression differences in natural Treg purified from young and aged female mice.

RNA sequencing (RNAseq) analysis showed that young and aged Treg clustered separately (**Fig. 6 A,**

C) as shown in the literature[42,43] and identified 1423 upregulated and 302 downregulated mRNA

transcripts in aged Treg compared to young Treg (**Fig. 6 B**). However, there was no difference in the

expression of classic Treg markers such as *Foxp3*, *Cd4* or *Il2ra* (**Sup. Fig. 5 B**). Pathway analysis

identified that the differentially expressed genes were associated with pathways linked to the

adaptive immune response, lymphocyte-mediated immunity, phagocytosis, cell-surface receptor

signalling and membrane invagination (**Fig. 6 D**). Genes that were upregulated with ageing were

enriched for the following GO biological processes: extracellular matrix organisation, cell migration,

synapse assembly, axonal guidance, cell adhesion and inflammatory and immune responses (**Fig. 6**

E). On the other hand, mRNA transcripts that were downregulated in aged Treg were associated with

defence responses to virus, B cell proliferation and protein folding GO biological processes (**Fig. 6 E**)

(GO cellular component, GO Molecular function and Kyoto Encyclopedia of Genes and Genome

(KEGG) database pathways analysis is shown in **Sup. Fig. 5**). To identify potential mechanisms

underlying how ageing may impair myelin regenerative functions of Treg, we examined expression

of genes associated with the Treg tissue repair programme described recently for skin, colon, lung

and adipose tisse[44]. Unexpectedly, most of the 31 Treg tissue repair programme hallmark/signature

genes[44] were in fact upregulated with ageing, except for *Klrb1, Ccr5, Selplg, Bach2, Rad50, Tfrc, Cd84*

and Cd164 (**Fig. 6 F, G**). No difference in expression was detected in these hallmark genes between

male and female natural Treg, showing that sex does not alter the expression of genes associated

with the tissue repair programme (**Sup. Fig. 5 C, D**). These data suggest that the mechanism(s)

involved in Treg-driven myelin regeneration capacity that are impaired with ageing, are likely

different to mechanisms from the Treg tissue repair programme described by Delacher et al[44].

Mcam and Itga2 are associated with Treg capacity to drive oligodendrocyte formation

To narrow down age-induced changes in Treg that diminish their capacity to induce OPC

differentiation, we first investigated whether the putative mechanisms are soluble or contact-

mediated; the latter being suggested by the close physical association of Treg with OPCs observed in

co-cultures (**Fig. 7 A**) and in EAE-induced, demyelinated lesions *in vivo* (**Fig. 7 B**). Young natural Treg

were co-cultured either directly in contact with OPC, or in the same well but separated using a

transwell, which allowed for the exchange of secreted molecules but not direct OPC-Treg contact.

While young Treg co-cultured directly with OPCs enhanced OPC differentiation into MBP$^+$

oligodendrocytes, this pro-differentiating effect was lost when Treg were physically separated from

OPCs (**Fig. 7 C, D**). To identify potential cell-to-cell contact mechanisms responsible for Treg-driven

OPC differentiation that may be lost with ageing, we combined our aged Treg RNA sequencing data

with a publicly available RNAseq dataset that compares between OPCs, newly differentiated and

differentiated oligodendrocytes[45]. We hypothesised that ligand-receptor interactions between Treg

and plasma membrane receptors enriched in OPCs underlie the pro-differentiation effect of Treg on

OPC. We prioritised ligand-receptor partnerships between Treg and OPCs that were impacted by

ageing, to identify mechanisms that could explain the lost myelin regenerative capacity of aged Treg.

We then performed a protein-protein interaction analysis using String[46] (**Fig. 7 F**) and prioritised

plasma membrane proteins that had the highest decrease in expression for aged Treg and the

highest number of potential binding partners amongst OPC-enriched receptors compared to

oligodendrocytes (**Fig. 7 G**). We identified the following key candidates: *Ccr7, Itga2* (CD49b), *Klrb1c*

(NK1.1), *Ly6c1, Mcam* (CD146) and *Sell* (CD62L) (**Fig. 5 F-H**). Next, we used a loss-of-function

approach to investigate whether these candidates may confer the lost regenerative capacity of aged

Treg, namely, CD62L, NK1.1, LY6C, MCAM (CD146) and ITGA2 (CD49b). Of these five targets,

antibody-mediated blocking of MCAM (CD146) and ITGA2 (CD49b) on Treg prior to Treg-OPC co-

culture inhibited the capacity of young Treg to drive OPC differentiation *in vitro* as indicated by the

fold change in the percentage of MBP$^+$ cells and the MBP$^+$ area per well (**Fig. 8 A-C, Sup. Fig. 6 A, B**).

This suggests that MCAM and ITGA2 are previously unidentified mediators of pro-remyelinating

functions of Treg and are downregulated with age.

Next, we sought to determine whether MCAM and ITGA2 would directly promote OPC

differentiation by treating neonatal OPCs with recombinant ITGA2 and MCAM. While treatment with

rMCAM for up to 6 days *in vitro* did not affect OPC proliferation or differentiation (**Sup. Fig. 6C, E-G**),

rITGA2 directly enhanced OPC differentiation by day 3 (**Fig. 8 D-G**). rITGA2 did not significantly

change the proportion of proliferating OPC (**Sup. Fig. 6 D**) or the proportion of MBP-expressing

oligodendrocytes (**Fig. 8 D, E**). However, we observed an increased complexity of oligodendrocyte

morphology in rITGA2-treated cultures as indicated by the enhanced number of intersections

detected by Sholl analysis on ITGA2-treated OPCs (**Fig. 8 F, G**). This suggests that oligodendrocytes in

rITGA2 treated cultures are at later stage of development which aligns with an acceleration of

differentiation. Therefore, while rITGA2 directly promoted OPC differentiation, rMCAM did not,

suggesting that the function of this latter cell adhesion molecule in Treg-mediated OPC

differentiation may be to facilitate OPC-Treg interaction and potentially to activate other pro-

differentiative mechanisms rather than itself directly driving OPC differentiation.

To determine whether a young environment could boost MCAM and ITGA2 expression by

aged Treg, we adoptively transferred young and aged Treg expressing red fluorescent protein (RFP$^+$)

obtained from Foxp3-RFP mice (C57BL/6-*Foxp3*[tm1Flv]/J Foxp3-IRES-mRFP) into young, Treg-depleted

mice and assessed ITGA2 and MCAM expression in spleen and lymph node after 11 days by flow

cytometry (**Sup. Fig. 7 A, B**). Due to technical challenges and a lack of further aged FoxP3 reporter

mice, only n=2-3 was possible in this study which limits interpretation of statistical significance of

our findings. The proportions of MCAM$^+$ or ITGA2$^+$ cells within total Foxp3-RFP$^+$ populations were

low in basal conditions and were significantly increased in both aged and young Treg upon adoptive

transfer (**Sup. Fig. 7 C, F**). To examine the extent of expression of each marker on a per cell basis, we

quantified median fluorescent intensity of MCAM$^+$RFP$^+$ and ITGA2$^+$RFP$^+$ Treg and observed

decreased expression of both markers in aged Treg compared to young Treg, in agreement with our

RNAseq findings. Following adoptive transfer into young mice, expression of ITGA2 was fully

restored to levels comparable to that of young Treg (**Sup. Fig. 7 C-E**), while MCAM in aged Treg was

partially restored (**Sup. Fig. 7 F-H**). These findings suggest that ITGA2 and MCAM are Treg-expressed

proteins that directly or indirectly support OPC differentiation, are decreased with age and

expression of which can be restored when aged Treg are adoptively transferred into a young

environment.

Discussion

Ageing is one of the strongest risk factors associated with the transition from relapsing-

remitting to secondary progressive MS[12], a phase that is characterised by the accumulation of

neurodegeneration and irreversible disability. This is partly due to impaired remyelination with age,

which was first described in animal models of remyelination[9,10] but has now also been observed in

MS patients[8,11]. Several factors have been linked to age-associated myelin regeneration failure, such

as impaired myelin debris phagocytosis by microglia and macrophages[19,20,47], reduced intrinsic

capacity of OPCs to differentiate into myelin-forming oligodendrocytes[13,17] and a refractory CNS

environment with increased stiffness and changes in extracellular matrix[16,17]. Therapies that enhance

remyelination and prevent age-related neurological decline are an urgent unmet need. Beyond their

classic immune-modulatory and anti-inflammatory role, Treg have recently emerged as key

regulators of tissue homeostasis and regeneration in several tissues[27], including the CNS, where we

have previously identified Treg as key promoters of remyelination[27]. However, how ageing affects

the capacity of Treg to enhance myelin regeneration had not been addressed.

Peripheral Treg[30,42,48], as well as Treg and tissue-resident Treg increase with age, a somewhat

paradoxical finding given the tissue-regenerative functions of Treg and the impairment of tissue

regeneration with age. Here, we further expanded the murine Treg population *in vivo* and showed

that despite this increased cell number, aged-associated remyelination failure was not rescued.

These data suggest that with ageing, Treg have an intrinsically impaired myelin-regenerative

capacity. Ageing is known to enhance Treg senescence, limit Treg proliferation[30,42] and there are

conflicting reports on the effect of ageing on Treg immunosuppressive capacity[30,42]. Previous work

on muscle regeneration showed that aged Treg have an impaired migratory capacity that hinders

muscle regeneration[36]. Additionally, in a model of influenza infection, aged Treg demonstrated an

impaired capacity to support lung repair, which was associated with a cell autonomous impairment

in their regenerative programme[43]. Here, we sought to determine whether aged Treg have an

intrinsically impaired capacity to drive CNS myelin regeneration combining *in vitro, ex vivo* and *in*

vivo approaches.

Purified aged Treg failed to significantly drive OPC differentiation into MBP-expressing

oligodendrocytes or the myelination of organotypic cerebellar slices, which was in striking contrast

to young Treg, which robustly drove both biological processes. These data suggest that ageing is

associated with a cell-intrinsic impairment of the regenerative capacity of Treg in CNS myelin

regeneration. Given that IL-2-mediated Treg expansion *in vivo* is in clinical trial for MS (clinical trial

number NCT02424396)[49], our findings are of translational importance as they suggest that Treg

expansion as a potential pro-remyelination therapy may be limited in older populations.

We next investigated whether the regenerative capacity of aged Treg could be restored, and

therefore therapeutically targetable. Surprisingly, aged Treg rescued OPC differentiation and myelin

wrapping similarly to that of young Treg in an *in vivo* model of young spinal cord remyelination. In

contrast to findings in the lung[43], the CNS remyelinating capacity of Treg can be partially restored by

a young environment. Interestingly, in lesions of aged mice lacking Treg, young Treg partially

restored Olig2$^+$CC1$^+$ oligodendrocyte density, while aged Treg did not, suggesting that indeed young

Treg have a more potent regenerative capacity than aged Treg. However, neither young nor aged

Treg restored Olig2$^+$ density or remyelination caused by depletion of endogenous Treg, suggesting

that with age, the failure of Treg to drive remyelination is a combination of both an intrinsically

impaired regenerative capacity of Treg and an aged CNS that is refractory to Treg-derived pro-

differentiation signals. Previous studies identified a tissue repair transcriptomic signature in the skin,

adipose tissue, and the lung[44]. Unexpectedly, very few of these hallmark Treg tissue repair genes

were downregulated in aged Treg in our transcriptomic profiling studies. These data indicate that

although some Treg regenerative mechanisms may be common across tissues, other mechanisms

may be tissue-specific or process- specific[50].

We recognised that the impairment of myelin regenerative Treg functions with age

presented a new avenue to discover novel mechanisms of how Tregs support CNS remyelination and

the potential to identify new targets that may be of therapeutic relevance to boost remyelination.

Indeed, using a combined approach of Treg/OPC transcriptomic dataset cross-referencing with *in*

vitro antibody blocking assays, we identified MCAM and ITGA2 as novel candidate mediators of Treg-

driven OPC differentiation. These two candidates were downregulated in aged compared to young

Treg, likely interact directly with OPCs and inhibit young Treg-driven OPC differentiation when

blocked *in vitro*. Transfer of aged Treg into young mice boosted expression of ITGA2 and to a lesser

extent of MCAM by Treg in preliminary studies, suggesting that these are modifiable factors that can

be restored in aged Treg. Furthermore, recombinant ITGA2, but not MCAM, promoted OPC

differentiation *in vitro* suggesting that these proteins have direct and indirect roles in Treg-driven

OPC differentiation.

To our knowledge this is the first report identifying ITGA2 as a novel mediator of Treg-driven

OPC differentiation, that is decreased with age, and can directly promote OPC differentiation. This

aligns with previous reports showing that other integrins are involved in oligodendrocyte

development[51] and shows that Treg are additional cellular contributors to integrin-mediated OPC

differentiation. MCAM has previously been described as a cell-adhesion molecule contributing to

brain inflammation by facilitating pathogenic T cell extravasation into the CNS[52–54] and here, we

identify a putative regenerative function of MCAM expressed by Treg. Together these data

contribute to the growing evidence that Treg express MCAM[55,56] and suggests a potential

regenerative role mediating Treg-stem cell interactions. Even though our data show that, *in vitro*,

non-activated natural Treg require direct OPC-Treg contact to drive OPC differentiation, previous

work from our laboratory showed that activated Treg secretome also enhanced OPC

differentiation[27]. Thus, Treg-mediated regenerative capacity likely involves a range of signalling

molecules that regulate OPC-Treg interaction as well as subsequent OPC differentiation and

remyelination, providing a range of potential novel mechanisms to explore as therapeutic targets for

remyelination therapy in the future.

Collectively, these studies have identified that ageing is associated with a cell-intrinsic

impairment of the myelin-regenerative functions of Treg. This discovery increases the range of

remyelination-associated mechanisms that are impaired with age[9,10,13,17–20,47]. Encouragingly, we

have shown that this impairment can be rescued in a young environment *in vivo*. We have also

identified two novel mediators of Treg-driven OPC differentiation, MCAM and ITGA2. This study has

identified novel mechanisms involved in CNS regeneration which need to be further investigated to

determine if they hold therapeutic potential for patients with demyelinating diseases.

Methods

Animals

All mice were on a C57BL/6J background and were either bred in-house or purchased from Charles

River Laboratories, UK. $Foxp3^{tm3(DTR/GFP)Ayr}$/J (Foxp3-DTR) mice were a kindly provided by Prof.

Alexander Rudensky (Memorial Sloan Kettering Institute, New York) and C57BL/6-$Foxp3^{tm1Flv}$/J

Foxp3-IRES-mRFP (Foxp3-RFP) mice were a kind gift from Dr. Rebecca Ingram. Neonatal C57BL6/J P3-

P7 pups were used for OPC and brain slice preparations, which were combined with natural Treg

were isolated exclusively from young (3-4m) and aged (15-18m) C57BL6/J male mice for use with

neonatal mixed sex pups. For *in vivo* Treg adoptive transfer studies, natural Treg were isolated from

young (3-4m) and aged (15m) C57BL6/J female mice or Foxp3-RFP mice, and were injected into both,

female and male recipient Foxp3-DTR mice (2-4m). To deplete Foxp3$^+$ cells in Foxp3-DTR mice,

diphtheria toxin was administered as described below. All animal maintenance and experiments

were done in accordance with the UK Home Office regulation (Project Licences 2789 and 2894) and

were approved by the Queen's University Belfast's Animal Welfare and Ethical Review Committee.

Flow cytometric analysis of healthy murine CNS

Male C57BL/6 mice (8-week-old and 75 and 95-week-old) were injected i.v. with biotinylated CD45

antibody (30-F11, eBioscience) 3 minutes prior to euthanasia via a schedule 1 method. Brain, spinal

cord and spleen tissues were dissected out and placed individually into 5mL cold FACS buffer (2%

FCS in PBS). Single cell suspensions were prepared from brain and spinal cord tissues as previously

described[57]. Briefly, tissues were chopped into small pieces and digested for 30 min at 37°C with 0.4

mg/mL collagenase D (Sigma-Aldrich), 300 µg/mL hyaluronidase (Sigma-Aldrich) and 40 µg/mL

DNase 1 (Sigma-Aldrich) in IMDM supplemented with 2 mM MgCl2, 2 mM CaCl2, 2 mM l-Glutamine

(Sigma-Aldrich), 1 mM Sodium Pyruvate (Gibco) 10 mM HEPES (Gibco), 20% FBS and 1X Gentamycin

(Abcam), followed by filtration (through 100 µm mesh) and enrichment for leukocytes by gradient

centrifugation (40% Percoll GE healthcare, 600 x g, 15 min). To prepare single cell suspensions from

spleen tissues, the spleen was mechanically disrupted between two glass slides, filtered through 100

μm mesh and red blood cells were lysed. Splenocytes were counted using a Countess automated cell

counter. Following cell preparation, non-specific binding was blocked using 2.4G2 supernatant.

Initial staining included eBioscience Fixable Viability Dye e780 (1:4000, eBioscience) and a

fluorescence-tagged streptavidin to identify leukocytes which were from contaminating blood using

the injected CD45-biotin (1:1000, eBioscience, Clone 30-F11). Next, a 1h surface stain was performed

including antibodies for CD45 (1:500; eBioscience, Clone 30-F11), CD4 (1:500; eBioscience, Clone

GK1.5), CD8 (1:1000 Biolegend, Clone 53-6.7), CD62L (1:1000; eBioscience, Clone Mel-14), CD11b

(1:500; eBioscience, Clone M1/70), CD44 (1:500; Biolegend, Clone IM7), CD69 (1:100, eBioscience,

Clone H1.2F3), CD25 (1:200; eBioscience, Clone PC61) and CD19 (1:250, eBioscience, Clone ID3)

Subsequently, cells were fixed and permeabilised with the eBioscience Foxp3 staining kit

(eBioscience) and a 1h intracellular stain for Foxp3 was added (1:100, eBioscience, Clone FJK-16s;

Miltenyi, Clone REA788). For brain and spinal cord, the entire sample was acquired. Flow cytometry

data were acquired on a Cytek Aurora and data collection performed using SpectroFlo (Cytek).

Lysolecithin-induced spinal cord demyelination *in vivo*

Spinal cord demyelination was induced as described previously[27]. In brief, demyelination was

induced in the ventral white matter funiculus of the thoracic spinal cord (between vertebrae T11-12

or T12-13) by the injection of 1.2 μL of 1% (w/v) L-α-Lysophosphatidylcholine (Lysolecithin; Sigma-

Aldrich) under general anaesthesia. At 7-, 10- or 14-days post lesion (dpl), mice were terminally

anaesthetised with intraperitoneal (i.p.) pentobarbital injections and transcardially perfused with

ice-cold phosphate buffered saline (PBS) followed by 4% paraformaldehyde (PFA) (Sigma-Aldrich).

Spinal cords were dissected and immersed overnight in 4% PFA at 4 °C. Next, spinal cords were

cryoprotected with 30% sucrose (Sigma-Aldrich) in PBS for 72h and snap-frozen in OCT (Tissue-Tek).

Frozen spinal cords were cryosectioned at 12 μm thickness and immunostained as described below.

EAE induction

Young (3-4m) and aged (15m) old C57BL/6J female mice were acquired from Charles River Laboratories. Anesthetised mice were immunised with subcutaneous injections of 150 µg of MOG_{35-55} (GenScript) emulsified in 200 µl of complete Freud's adjuvant (CFA) (Incomplete Freund's Adjuvant containing 2.5 mg/mL Mycobacterium tuberculosis H37Ra, both BD Biosciences). At 0- and 2-days post-immunisation, mice were injected with 200 ng of pertussis toxin intraperitoneally (PHZ1174, Thermofisher). EAE clinical signs and weight were monitored daily from 7 days post-immunisation. Mice were euthanised at 36 days post-immunisation by transcardial perfusion with PBS followed by 4% PFA. Spinal cords were dissected and immersed overnight in 4% PFA at 4 °C. Next, spinal cords were cryoprotected with 30% sucrose (Sigma-Aldrich) in PBS for 72h and snap-frozen in OCT (Tissue-Tek). Frozen spinal cords were cryosectioned at 12 µm thickness and immunostained as described below.

In vivo Treg expansion

Young (2-3m) and aged mice (15m) were intraperitoneally injected with IL-2/anti-IL-2 antibody complexes to expand endogenous Treg[58]. In brief, murine IL-2 (1 µg; Peprotech) was mixed with anti-mouse IL-2 (5 µg; Bioxcell, Clone JES6-1A12) in PBS at 37 °C for 30 min. Each mouse was injected daily for 3 consecutive days with IL-2 (1 µg) and anti-IL-2 (5 µg) in 200µl of PBS. Adequate Treg expansion was assessed by flow cytometry as described below. Spinal cord demyelination was induced on the day after the third injection of IL-2/anti-IL-2, by injecting lysolecithin into the ventrolateral white matter as described above. Treg expansion was examined by flow cytometry as described below on day 10dpl in aged mice, while OPC differentiation was examined at 7 dpl in young mice and 10 dpl in aged mice.

In vivo Treg depletion and adoptive transfer of wild type natural Treg

Young (2-4m) and aged (19-23m) male and female Foxp3-DTR mice were injected daily i.p. with

diphtheria toxin (DT; 0.04 µg/g of body weight; Sigma, Cat. No. D0564) for 3 days prior to

demyelination. To maintain endogenous Treg depletion during remyelination, DT (0.04 µg/g of body

weight) was injected i.p. every fourth day. Control animals received 200 µl of saline i.p. Depletion

was confirmed at the endpoint by flow cytometric analysis of endogenous GFP$^+$ natural Treg in

blood, spleen, and lymph nodes (see below). To adoptively transfer young (2-4m) and aged (15m)

natural Treg to Treg-depleted mice, we immunomagnetically isolated natural Treg from wild type

female mice or Foxp3-RFP male and female mice using a CD4$^+$ and CD25$^+$ Treg isolation kit

(STEMCELL Technologies) as described below. For remyelination experiments, mice were injected

i.p. with 10^6 wild type natural Treg (which are resistant to DT) in the 24h prior to lysolecithin-induced

demyelination, while in the case of adoptive transfer of RFP$^+$ Treg to evaluate MCAM and ITGA2

expression, 4 x 10^5 cells per mouse were injected to non-lesioned mice. Reconstitution was

evaluated by flow cytometry in lymph nodes at the endpoint (see below).

OPC isolation and culture

OPCs were isolated from P3-P7 mice based on A2B5 expression. Briefly, pups were culled by an

overdose of pentobarbital (200 mg/mL, 20µl per mouse) and whole brain was dissected and placed

in ice-cold Hibernate-A (ThermoFisher Scientific). Dissected brain was minced using scalpels,

transferred to a 15mL tube, and centrifuged for 1 min at 100 g and 4 °C. Hibernate-A medium was

aspirated, and the pellet was resuspended in 5 mL dissociation media containing 165U of Papain

(Worthington) and DNAse type I (40 µg/mL; Worthington) in hibernate A for 30 min at 37 °C. Upon

digestion papain was washed off with Hanks Buffered Salt Solution (HBSS) (ThermoFisher Scientific)

and cells were centrifuged for 5 min at 300 g and 4 °C. The pellet was resuspended in 4 mL

trituration buffer (Hibernate-A with 2% B27; ThermoFisher Scientific) and 2 mM sodium pyruvate

(ThermoFisher Scientific) and gently triturated ten times using a 5 mL pipette. Tissue was left to

settle for 2 min at room temperature (RT), and supernatant was transferred to a clean 50 mL tube

through a 70 µm strainer (Corning). Remaining tissue was exposed to another 2 mL of trituration

buffer and gently triturated using a glass polished pipette. After trituration, tissue was left to settle

for 2 min at RT and supernatant was transferred to the tube through the 70 µm strainer. This step

was repeated another time with a glass fire-polished pipette of decreased diameter and then with a

1 mL pipette. To further remove debris from the cell suspension, 11.5 mL of 90% Percoll (GE

Healthcare) diluted in 10X PBS (ThermoFisher Scientific) was added and topped up to a final volume

of 45 mL with Dulbecco's Modified Eagle's medium (DMEM) (ThermoFisher Scientific). The cell

suspension was then centrifuged at 800 g for 20 min at 4 °C. Upon centrifugation, the pellet was

washed with HBSS and then resuspended in 10 mL magnetic-activated cell sorting (MACS) buffer

(Hibernate-A with 2 mM sodium pyruvate, 2% B27, 0.5% BSA; ThermoFisher Scientific), 2 mM EDTA

(Sigma-Aldrich) and 10 µg/mL insulin (Sigma-Aldrich), and incubated for 30 min at 37 °C in a 10 cm

petri dish coated with BSC1 Griffonia Simplicifolia Lectin (BSL1, 5 µg/mL)(Vektor Labs) to remove

microglia. Then, the supernatant was collected, and cells were counted and centrifuged at 300 g for

5 min at 4 °C. The pellet was resuspended and incubated with 2 µg of A2B5 antibody (Millipore,

Clone A2B5-105) in 500 µL of MACS buffer per 10 million cells for 25 min on ice, with gentle

resuspensions every 10 min. Cells were washed with HBSS and spun at 300 g for 5min at 4 °C

(Miltenyi Biotec). The pellet was resuspended and incubated with 80 µL of MACS buffer and 20 µL of

anti-IgM microbeads (Miltenyi Biotec) for 15 min. Then, cell suspensions were centrifuged at 300 g

for 5 min at 4 °C, resuspended in 0.5 ml MACS buffer and placed in a MACS mini column (Miltenyi

Biotec) on a MiniMACS Separator (Miltenyi Biotec) and washed with 1.5mL MACS buffer. Once the

liquid passed completely through the column, the column was removed from the stand and placed

in a new 15 mL tube, where 1 mL of OPC media was added to the column and plunged through to

elute the OPCs. OPC media contained DMEM, 2% B27, sodium pyruvate (2 mM), insulin (5 µg/mL),

Trace Elements B (0.01%; Corning), Forskolin (5 µM; Sigma-Aldrich), Biotin (10 ng/mL; Sigma-

Aldrich), Penicillin-streptomycin-glutamine (1%; ThermoFisher Scientific), N-acetyl cysteine (60

µg/mL; Sigma-Aldrich) and 1% SATO stock solution. SATO stock solution contained bovine serum

albumin (BSA) fraction V (0.1 mg/mL; ThermoFisher Scientific), sodium selenite (4 µg/mL; Sigma-

Aldrich), putrescine (1.61 mg/mL; Sigma-Aldrich), apo-transferrin (0.1 mg/mL; Sigma-Aldrich) and

progesterone (4 µg/mL; Sigma-Aldrich). The purified population was counted and diluted in OPC

media for plating. OPCs were plated at a density of 3,000 cells per well in 96-well plates (Falcon)

previously coated with poly-L-lysine (10 µg/mL; Sigma-Aldrich) and laminin (10 µg/mL; Sigma-

Aldrich) diluted in DMEM. OPCs were plated with PDGFaa (20 ng/mL; Peprotech) and NT-3 (10

ng/mL; Peprotech). The following day two thirds of the media was removed (100 µl) and 100 µl of

fresh OPC media with PDGFaa (20 ng/mL) and NT3 (10 ng/mL) were added per well. Cells were

incubated at 37°C and 5% CO_2 for two days. On the third day, media was completely removed and

20,000 Treg in fresh media were added as indicated below.

For transwell assays, OPCs were selected with anti-PDGFRα panning. After microglia depletion in

BSL1-coated petri dishes, the supernatant was transferred to PDGFRα panning plates and incubated

at RT for 1 h. PDGFRα panning plates were coated with goat-anti-rat IgG (H+L) (7.5 µg/mL;

ThermoFisher Scientific) in PBS with 0.2% BSA and rat anti-PDGFRα (1.5 µg/mL in PBS; BD

Biosciences) with 0.2% BSA for 3 h. Supernatants were removed, cells were washed 5-8 times with

PBS and the attached OPCs were scraped, counted, and replated in 24-well plates (Sarstedt) at

10,000 cells per well. On the third day *in vitro* 50,000 Treg were added either directly into the well

with OPCs or in a transwell (Millipore) above the OPCs.

For recombinant ITGA2 and MCAM treatment, OPC were treated with increasing concentrations of

the recombinant proteins in OPC media, or vehicle control, on the day of growth factor removal.

Cells were cultured for 3 days in the case of ITGA2 and 6 days in the case of MCAM without any

further media changes.

At the endpoint (3 or 6 days after treatment) OPC were fixed for 15 min with 4% PFA and washed

twice with PBS prior to immunostaining. Cells were blocked with 5% donkey serum (Sigma-Aldrich)

with 0.1% Triton-X-100 (Sigma-Aldrich) in PBS for 1h at RT. OPCs were then stained overnight at 4 °C

with primary antibodies diluted in blocking solution against Olig2 (1:500, Bio-techne), CNPase (1:500,

Sigma-Aldrich, clone 11-5B), MBP (1:500, Millipore, clone 12), Ki67 (1:300, Abcam, clone SP6), NG2

(1:200, Millipore), and CD3 (1:500, eBioscience). Cells were then washed three times with PBS and

incubated for 1 h at room temperature with secondary antibodies diluted in blocking buffer (Alexa

fluor (AF) 488 donkey anti- rabbit (1:500; ThermoFisher Scientific), AF568 donkey-anti- rat (1:500;

Abcam), AF647 donkey anti-mouse (1:500; Abcam), AF647 donkey anti-rabbit (1:500; ThermoFisher

Scientific) and AF755 donkey anti-goat (1:500; ThermoFisher Scientific) and Höechst (1:10,000;

Sigma-Aldrich). Cells were then washed with PBS three times and stored in 150 µL PBS/well for cell

imaging.

Cell imaging was performed using the CellInsight CX5 high content imaging system (ThermoFisher

Scientific) for 96-plates and EVOS for 24-well plates. Twenty-five separate fields of view of each well

in 96-well plates and the cell populations of interest were quantified using CellInsight CX5 analysis

software and the SpotCounts programme. For 24-well plates we imaged randomly 5 fields of view

per well at 20X magnification and images were analysed manually with Fiji.

Natural Treg isolation

Young (2-4 m) and aged (15-18 m) male mice were culled by CO_2 overdose. Spleens and lymph nodes

were removed and mashed through a 70 µm strainer with a 2 ml syringe plunger. $CD4^+$ T cells were

immunomagnetically purified by negative selection according to manufacturer's instructions

(STEMCELL Technologies). Isolated $CD4^+$ T cells were then subjected to a $CD25^+$ cell isolation

(STEMCELl Technologies) using releasable magnetic spheres (STEMCELL Technologies) according to

the manufacturer's protocol. Purity was confirmed by flow cytometry for CD4, CD25 and Foxp3 (**Sup.**

Fig. 2A). For i.p. injections in adoptive transfer experiments, cells were resuspended at 10^6 cells per

200 µL in saline. For OPC-natural Treg co-cultures 20,000 natural Treg were resuspended in 150 µl of

OPC Brainphys-based media and added to OPCs in 96-well plates. For 24-well transwell plate

experiments 50,000 natural Treg were resuspended in 250 µl OPC-BrainPhys media and added to

transwells above cultured OPCs. OPC BrainPhys-based media contains BrainPhys (STEMCELL

Technologies), B27 (2%; ThermoFisher Scientific), Glutamax (0.5 mM; ThermoFisher Scientific), N-

acetyl-cysteine (60 µg/mL), SATO (1:100, as described above) and insulin (5 µg/mL). In the case of

brain slices, 50,000 Treg were diluted in 5 µL of brain slice media and dropped directly onto brain

slices.

For natural Treg mechanistic studies, OPCs and Treg were isolated as described above. Prior to OPC-

natural Treg co-culture, 120,000 Treg were transferred to a 1.5 mL microcentrifuge tube and

centrifuged at 400 g for 5 min at 4 °C. Treg were resuspended in 100 µL of PBS with 2% FCS and a

blocking antibody (20 µg/mL) or corresponding isotype were added per tube. Natural Treg were

incubated with the blocking antibody for 45 min on ice and then cells were washed with PBS and

centrifuged for 10 min at 400 g and 4 °C. Cells were resuspended in 900 µl of OPC BrainPhys-based

media (see above) and 20,000 Treg in 150 µL of media were added per well in 96-well plates. As

previously, Treg were co-cultured with OPCs for 7 days *in* vitro (DIV) and then fixed and stained as

described above. Blocking antibodies used were anti-NK1.1 PE (eBioscience, clone PK136), anti-

CD62L-APC (eBioscience, clone Mel14), anti-Ly6c1-APC (eBioscience, clone RB6-BC5), rat-isotype-

APC (eBioscience), anti-Itga2 (Abcam, clone EPR5788), anti-MCAM (Abcam, clone EPR3208) and

rabbit-isotype (Vektor labs).

Organotypic brain slice cultures

Brainstem slices from male and female P3 C57BL/6J mice were prepared at 300 µm thickness using a

McIlwain Tissue Chopper as described previously[27]. Slices were separated and placed on Millicell

inserts in individual wells of a 24-well plate (Millipore) with 250 µl brain slice medium containing

46.6% minimum essential medium (ThermoFisher Scientific), 25% Earls balanced salt solution

(Sigma-Aldrich), 25% heat-inactivated horse serum (Thermofisher Scientific), 1% penicillin-

streptomycin (Thermofisher Scientific), 1% glutamax (Thermofisher Scientific) and 1.4% D-glucose

(Sigma). Brain slices were incubated at 37 °C and 5% CO_2 overnight and then media was fully

replaced. On day 3, media was changed again and 50000 young or aged Treg were added directly on

top of each slice in a 5 µl droplet. Brain slices were cultured for a further 7 DIV, with media changes

every other day. On day 10 (7 days post-treatment), brain slices were fixed with 4% PFA for 45 min

and immunostained. In brief, brain slices were blocked for 2 h in 10% normal donkey serum (Sigma-

Aldrich), 1 mM Hepes (Thermofisher Scientific), 1% BSA (ThermoFisher Scientific) and 0.5% Triton X-

100 (Sigma-Aldrich) in PBS. Slices were incubated with primary antibodies for rat anti-MBP (1:500;

Millipore, clone 12) and chicken anti-NFH (1:500; EncorBiotech, polyclonal) in blocking buffer at 4 °C

for two overnights. Brain slices were then washed with PBS-0.01% triton three times for 30 min.

Slices were incubated with secondary antibodies (AF488 donkey anti-chicken IgY (1:500; Abcam,

polyclonal) and AF568 donkey anti-rat IgG (1:500; Abcam, polyclonal) in blocking buffer overnight at

4 °C. Brain slices were washed twice with PBS-0.01% triton and then incubated with Höechst stain

(1:20000; Sigma-Aldrich) for 10 min. Slices were washed with PBS and mounted using fluoromount G

(Thermofisher Scientific). Imaging was performed using a Leica SP8 confocal microscope, using a 63X

oil objective and the Leica Navigator at 0.5 µm intervals over 10 µm. Four fields of view per brain

slice were selected for imaging based on NFH and Höechst staining but blinded to MBP. To quantify

myelination index, the area stained by NFH, MBP and the colocalising area of NFH^+MBP^+ were

measured per stack in Fiji[59] using a Fiji plug-in developed and kindly provided by the Williams

laboratory at the University of Edinburgh[60]. Then, the ratio between total NFH^+ and NFH^+MBP^+ areas

(myelination index) was calculated per z-stack and the average of the myelination index per slice was

calculated.

Flow cytometry

Spleens and lymph nodes were mashed through a 70 µm strainer. For splenocytes, cells were

exposed to red blood cell lysis buffer (STEMCELL Technologies) for 2 min at room temperature. Both,

lymph nodes and splenocytes were then washed with PBS and centrifuged at 300 g for 5 min at 4°C.

Cells were resuspended in 200 µL PBS and stained with a cell viability dye with eFluor 506-UV

viability dye (1:2000; ThermoFisher Scientific) and cell surface stained with antibodies for CD4

(1:500; eBioscience, clone RM4.5), CD25 (1:500; eBioscience, clone PC61.5), ITGA2 (1:50;

eBioscience, Clone DX5) and MCAM (1:100; Biolegend, Clone ME9F1) for 15 min at RT. Cells were

washed with flow cytometry staining buffer (FCSB) (2% FCS in PBS) and centrifuged at 300 g for 5

min at 4 °C. Cells were then fixed with Fix & Perm A (ThermoFisher Scientific) for 10 min at RT.

Fixative was washed off with FCSB and centrifuged at 300 g for 5 min at 4 °C. The pellet was

resuspended in PBS and data were acquired on a FACSCanto II or on a FACS Symphony in the case of

RFP+ adoptive transfer. In the natural Treg depletion experiment, presence of endogenous natural

Treg was determined by the expression of GFP. To determine the purity of natural Treg isolations or

the extent of natural Treg reconstitution by adoptive transfer, cells were washed and centrifuged for

5 min at 300 g and 4 °C. Cells were then resuspended in 100 µL Fix & Perm B (ThermoFisher

Scientific) with an anti-Foxp3 antibody (1:100; eBioscience, clone FJK-16S) overnight at 4 °C. Cells

were then washed with FCSB and centrifuged at 300 g and 4 °C for 5 min. Cells were then

resuspended, data were acquired on a FACSCanto II and analysed using FlowJo software version 9.0

and 10.0 (BD). To calculate cell numbers, singlets were identified by FSC-H versus FSC-A and viable

cells gated for CD3 and CD4, and subsequently CD25 and Foxp3$^+$GFP$^+$ and Foxp3$^+$GFP$^-$ cells. To

evaluate young and aged Treg numbers in blood, 30 µL were incubated with 100 µL of the viability

and cell surface antibody mix for 30 min at room temperature. Cells were then washed with FCSB

and centrifuged at 300 g for 5 min at 4 °C. After resuspension, cells were fixed and lysed in 100 µL of

OptiLyse B (Beckman Coulter) for 10 min at RT, washed with distilled water and then after 15 min,

acquired on a FACSCanto II. After initial data acquisition, cells were centrifuged at 300 g for 5 min at

4 °C and incubated overnight with anti-Foxp3 antibody in Fix & Perm B, as described above. Cells

were washed and data were acquired on a FACSCanto II. To evaluate changes in MCAM and ITGA2

expression upon reconstitution, splenocytes were stained as described above but without Fix &

Perm A or B, to avoid quenching endogenous RFP and GFP signal and data were acquired from live

cells on a BD FACS Symphony. To calculate cell proportions and mean fluorescence intensity, singlets

were identified by FSC-H versus FSC-A and viable cells gated for CD4, and subsequently RFP$^+$ GFP$^-$

Treg. ITGA2 and MCAM expression based on median fluorescent intensity was then evaluated in

RFP$^+$ and ITGA or MCAM$^+$ RFP$^+$ Treg respectively.

Immunofluorescence staining of CNS tissue

Spinal cord sections were dried for 30 min at RT and washed for 10 min in PBS. For Olig2, Ki67, CC1

and ASPA staining, spinal cord sections underwent antigen retrieval at 85 °C for 10 min with 1X

citrate buffer pH 6.0 (Sigma-Aldrich). Tissue sections were washed with PBS and permeabilised with

1% Triton-X-100 in PBS for 30 min at RT. Sections were washed with PBS and incubated with blocking

solution (5% donkey serum; Sigma-Aldrich) diluted in TBS with 0.25% tween (Sigma-Aldrich) for 1h at

RT. If spinal cord sections were incubated with a primary antibody raised in mouse (e.g. anti-CC1)

sections underwent an additional 1 h blocking step at RT using Mouse on Mouse (M.O.M) blocking

reagent (Vector labs) in 5% donkey serum. Then spinal cord sections were incubated with primary

antibodies against Olig2 (1:500; Bio-Techne, polyclonal), anti-APC (1:400, Abcam, clone CC1), anti-

ASPA (1:300, Millipore, polyclonal), anti-NFH (1:500; Abcam, polyclonal), anti-MBP (1:500; Millipore,

clone 12) or anti-Ki67 (1:300; Abcam, clone SP6) overnight at 4 °C. Sections were washed with Tris-

Buffered-Saline (TBS) (ThermoFisher Scientific) with 0.25% Tween (Sigma-Aldrich). Secondary

antibodies including donkey anti-goat AF488, donkey anti-mouse AF568, donkey anti-rabbit AF647

(all 1:500; ThermoFisher Scientific), donkey anti-rat AF568 (1:500; Abcam), donkey anti-rabbit AF647

(1:500, Abcam) and Höechst (1:10000; Sigma-Aldrich) were added for 1 h at room temperature.

Spinal cord sections were then washed with TBS-0.25% Tween twice and mounted with fluoromount

G.

For detection of CD4 and Foxp3, spinal cord sections underwent antigen retrieval at 80 °C for 5 min.

After blocking with 10% NDS, sections were incubated with rat-anti-mouse CD4 (1:500; Biolegend,

Clone RM4.5) and rabbit-anti-mouse Foxp3 (1:500; Biotechne, polyclonal) for 1h at room

temperature, washed and incubated with secondary antibodies as described above.

Image acquisition was performed using the Leica TIRF and Leica DM5500 widefield fluorescent

microscopes. Further image processing was performed in Fiji[59] software and analysis was undertaken

by blinded manual counting. For NFH and MBP wrapping analysis, images were quantified using Cell

Profiler and Cell Profiler Analyst softwares[61,62].

RNA sequencing and analysis

Young (2-4m) and aged (16m-20m) natural Treg were isolated from spleen and lymph nodes by

magnetic-activated cell sorting and purity was checked by flow cytometry as described above (**Sup.**

Fig. 3A). Natural Treg were lysed in 0.5 mL of Trizol (Sigma-Aldrich) by vortexing and pipetting and

frozen at -80 °C. RNA was extracted combining Trizol and Chloform isolations with RNAeasy micro kit

columns (Qiagen). RNA concentration was determined using a Nanodrop, and RNA quality was

assessed by Qubit measurement and in an RNA nanochip Bioanalyzer. Sequencing libraries were

prepared using the Kapa Hyper Preparation kit with riboerase (Roche) following manufacturer's

instructions. Sequencing was performed on the NovaSeq 6000 in a pair-end 75 base pair format and

80 million reads.

The raw sequencing data was checked for lower quality bases and adaptor sequences with FastQC

and quality trimming was performed using CutAdapt[63]. The high-quality trimmed data were mapped

on mm10 mouse reference genome using STAR aligner (v 2.7.0a)[64]. The quantification of each gene

transcripts was performed with ensemble transcript annotation gtf file (GRCm38) using

featureCounts tool (v 2.0.0)[65]. The resultant read counts matrix was analyse using DESeq2 (v1.38.1)[66]

to identify differentially expressed genes between the young and aged natural Treg samples using

Wald tests with the Bonferroni correction for multiple testing. The differentially expressed genes

(DEG) with a multiple testing adjusted $p<0.05$ and fold change >2 were considered for further

downstream analysis. These genes were subjected to gene ontology enrichment analysis using

enrichGO function in clusterProfiler (v4.2) R package and DAVID (https://david.ncifcrf.gov). Over and

underrepresented ontology terms were identified (p value <= 0.05 and Bonferroni correction p

adjustment method). Next, enriched activated and suppressed functional pathways were identified

using gseKEGG function in clusterProfiler with p value<0.05.

Raw data files from RNAseq are available at the GEO database (accession number GSE218804).

Bioinformatic analysis to identify putative Treg mechanisms

To determine the molecular mechanisms associated with Treg-driven OPC differentiation that are

impaired with ageing, we combined our young and aged natural Treg RNA sequencing with data

obtained from Zhang *et al.* 2014 (GSE: 52564)[45]. We first extracted genes that were downregulated

in aged natural Treg and associated with the GO cellular component term "plasma membrane". We

then extracted the FPKM matrix and selected genes that were upregulated in OPC when compared

with newly myelinating oligodendrocytes and oligodendrocytes in the Zhang *et al.* database. We

performed GO cellular component analysis on those OPC-enriched genes and selected genes

associated with the GO term Apical Plasma Membrane. Then, both gene lists were subjected to a

protein-protein interaction analysis using the String open-source tool (www.string-db.org)[46]. Protein-

protein interactions within genes that were associated only with the natural Treg or only the OPC

databases were eliminated, and the remaining downregulated natural Treg membrane proteins

were prioritised considering their significance, fold change and the number of potential interactors

enriched in OPCs. Candidate genes were then selected based on their p-adjusted value, fold

enrichment and the number of interactions with OPC receptors.

Statistical analysis

All statistical analyses were performed in GraphPad Prism (GraphPad Software, Inc. version 9) or R.

First normality of datasets was assessed using Shapiro-Wilk and Kolmogorov-Smirnov tests. For

comparisons between 2 groups, such as young and aged Treg and OPC differentiation (Sup. Fig. 1),

unpaired two-tailed Student's t tests were used if data were normally distributed, and Mann-

Whitney U tests for non-parametric datasets.   When more than two groups were analysed and

datasets were normally distributed, a one-way analysis of variance (ANOVA) was performed

assuming equal variances, followed by Sidak's multiple comparison test. When two factors were

controlling the outcome, such as replicates and treatments, 2-way ANOVA followed by Dunnet's

comparison was used. When datasets were was not normally distributed, a Kruskal-Wallis followed

by Dunn's multiple comparison test was used. For percentage data, arcsin conversion was

performed to analyse the data using parametric tests. For all statistical tests, differences were

considered significant at p<0.05.

Data and code availability

Young and aged natural Treg RNA sequencing data generated in this study are available at GEO

(accession number: GSE218804).

Figure legends

Figure 1: Treg frequency is increased in the aged CNS. A Flow cytometric plot and **B** quantification

showing the proportions of natural Treg (endogenous GFP reporting Foxp3 expression in Foxp3-DTR

mice) in aged blood compared to young circulation (n=30-32 mice, unpaired two-tailed Student's t

test after *arcsin* conversion). **C** Flow cytometric plot and **D** quantification of Treg proportion,

identified by CD4 and Foxp3 expression in young and aged mouse spleens (n=15-32 mice, unpaired

two-tailed Student's t test after *arcsin* conversion). **E** Quantification of tissue resident Treg

proportion, identified by CD69 expression (n=8-13 mice, unpaired two-tailed Student's t test after

arcsin conversion). **F-H** Violin plot-based quantification showing the proportion and count of Treg

and tissue resident Treg in the spinal cord of healthy young and aged mice (n=8-13 mice, unpaired

two-tailed Student's t test after *arcsin* conversion). **I-K** Violin plot-based quantification showing the

proportion and count to Treg and tissue resident Treg in the brain of healthy young and aged mice

(n=8-13 mice, unpaired two-tailed Student's t test after *arcsin* conversion). **L** Immunostaining of

DAPI (blue), CD4 (green) and Foxp3 (magenta) Tregs (highlighted with yellow arrows) in a

demyelinated lesion from mice with EAE (scale bar = 50 μm). **M** Quantification of the density of CD4$^+$

Foxp3$^+$ Treg in lysolecithin-induced demyelinated lesions at 3 and 14 dpl (n=3-7 mice, 2-way

ANOVA). **N** Quantification of the density of CD4$^+$ Foxp3$^+$ Treg in EAE-induced demyelinated lesions at

36 days post-immunisation (n=3-4 mice, unpaired two-tailed Student's t-test).

Figure 2: Expansion of Treg does not rescue impaired OPC differentiation in aged mice. A Flow

cytometric plot showing CD25 and Foxp3 expression of control aged mice and aged mice treated

with intraperitoneal injection of IL-2/anti-IL-2 complexes **B, C** Quantification of CD25$^+$Foxp3$^+$ natural

Treg proportion in a CD4$^+$ T cell population from the spleen and blood of control aged mice and aged

mice treated with intraperitoneal injection of IL-2/anti-IL-2 complexes (n=3, unpaired two-tailed

Student's t test after *arcsin* conversion). **D** Representative images of OLIG2 (cyan) and CC1 (magenta)

immunostaining in young control and Treg expanded demyelinated lesions at 7dpl (scale bar = 50

μm, lesion area demarcated with the dotted bar). **E, F** Quantification of OLIG2$^+$ cell density and

OLIG2$^+$CC1$^+$ cell density in the demyelinated area of young control and Treg expanded mice (n=4-5

mice, unpaired two-tailed Student's t-test). **G** Representative images of OLIG2 (cyan) and CC1

(magenta) immunostaining in aged control and Treg expanded demyelinated lesions at 10dpl (scale

bar = 50 μm, lesion area demarcated with the dotted bar). **H, I** Quantification of OLIG2$^+$ cell density

and OLIG2$^+$CC1$^+$ cell density in the demyelinated area of age control and Treg expanded mice (n=6-7

mice, unpaired two-tailed Student's t-test). **J** Representative images of NFH (cyan) and MBP

(magenta) immunostaining in aged control and Treg expanded demyelinated lesions at 10dpl (scale

bar = 50 μm, lesion area demarcated with the dotted bar). **K, L** Quantification of NFH$^+$ axon density

and NFH$^+$MBP$^+$ axon density in the demyelinated area of aged control and Treg expanded mice (n=6-

7 mice, unpaired two-tailed Student's t-test).

Figure 3: Aged natural Treg demonstrate impaired capacity to drive OPC differentiation and

myelination *in vitro*. A Representative images of OPCs co-cultured with young and aged natural Treg

and immunostained for OLIG2 (cyan) as a pan oligodendrocyte lineage marker, the proliferation

marker Ki67 (magenta). **B** Quantification of OPC proliferation when co-cultured with young and aged

natural Treg (n=6, 1-way ANOVA after *arcsin* conversion). **C** Representative images of OPCs co-

cultured with young and aged natural Treg and immunostained for OLIG2 (cyan) and differentiation

markers CNP (grey) and MBP (magenta) (**C**) (scale bar = 100 μm). **D** Quantification of the proportion

of OPCs reaching early stage-differentiation when exposed to young and aged natural Treg, as

indicated by CNPase staining (n=9, 1-way ANOVA after *arcsin* conversion). **E** Quantification of the

proportion of OPCs expressing late-stage differentiation marker MBP in control and OPCs treated

with young and aged natural Treg (n=9, 1-way ANOVA, Sidak's multiple comparisons test). **F**

Immunohistochemistry of control and young and aged Treg treated cerebellar slices (MBP, magenta

and NFH, green, scale bar = 100 μm). **G** Quantification of myelination index (ratio between MBP and

NFH colocalization area and NFH area) and myelination index fold change in neonatal cerebellar

slices (n=9 mice, 1 brain slice per mouse, 3 independent experiments, 1-way ANOVA after *arcsin*

conversion, Sidak's multiple comparison tests).

Figure 4: Myelin-regenerative capacity of aged Treg is restored in a young environment. A Diagram

explaining the experimental design of *in vivo* Treg depletion, Treg adoptive transfer and spinal cord

demyelination. **B, C** Representative images of immunostaining identifying oligodendrocytes by the

co-localisation of the pan-oligodendrocyte lineage marker OLIG2 (green) with CC1 (magenta, **B**) or

ASPA (magenta, **C**) at 14 dpl (scale bar = 100 μm, demyelination area is highlighted by the white

line). **D-F** Bar graphs show the quantification of total number of oligodendrocyte lineage cells, as

well as CC1-expressing or ASPA-expressing oligodendrocytes in the demyelinated lesions of PBS

control mice, natural Treg-depleted mice and mice depleted of endogenous Treg that received

young or aged natural Treg by adoptive transfer (n=7-9, 1-way ANOVA, Sidak's multiple comparisons

test). **G** Representative images of immunostaining for neurofilament-H (NFH, green) and MBP

(magenta) to quantify myelin wrapping as an early marker of remyelination at 14 dpl (scale bar = 100

μm, demyelination area is highlighted by the white line). **H-J** Quantification shows the total number

of axons, the density of MBP wrapped axons and the percentage of MBP-wrapped axons from the

total number of axons (n=6-9, 1-way ANOVA, Sidak's multiple comparisons test).

Figure 5: Young Treg only partially rescue oligodendrocyte differentiation in aged Treg-depleted

mice. A Representative images of immunostaining identifying oligodendrocytes by the co-

localisation of the pan-oligodendrocyte lineage marker OLIG2 (green) with CC1 (magenta) at 14 dpl

(scale bar = 100 μm, demyelination area is highlighted by the white dotted line). **B-D** Bar graphs

show the quantification of total number of oligodendrocyte lineage cells, as well as Ki67$^+$

proliferating OPCs and CC1$^+$ oligodendrocytes in the demyelinated lesions of PBS-treated control

aged mice, natural Treg-depleted aged mice and aged mice depleted of endogenous Treg that

received young or aged natural Treg by adoptive transfer (n=4-6 mice, 1-way ANOVA, Sidak's

multiple comparisons test).

Figure 6: Ageing significantly alters natural Treg transcriptome. A Principal component analysis

demonstrating the clustering differences between young and aged natural Treg. **B** Volcano plot

demonstrating 1456 genes upregulated and 302 genes downregulated genes in aged natural Treg

relative to young Treg. **C** Heatmap demonstrating hierarchical clustering of the top 50 differentially

expressed genes between young and aged Treg. **D** Graph showing the pathways enriched amongst

differentially expressed genes. **E** Bar graph highlighting the GO biological processes associated with

genes that are upregulated or downregulated in aged Treg. **F** Heatmap showing the normalised

count values for the Treg tissue repair programme identified by Delacher *et al.*[44] **G** Graph showing

the Treg tissue repair programme signature genes score in aged and young natural Treg.

Figure 7: Natural Treg-driven OPC differentiation *in vitro* is contact-dependent. A Representative

images of immunostaining showing cell-to-cell contact between OPCs and Treg in OPC-Treg co-

cultures *in vitro*. OPCs are identified by the co-staining of OLIG2 (cyan) and NG2 (grey), while Treg

are identified by CD3 (red) (scale bar = 50 μm). **B** Representative images showing Treg (CD4$^+$ (green),

Foxp3$^+$ (magenta)) in close proximity or direct contact with OLIG2$^+$ oligodendrocyte lineage cells in

demyelinated lesions in mice with EAE, highlighted by the yellow arrows (scale bar=25μm). **C**

Representative images of immunostaining of OPCs directly co-cultured with young Treg and OPCs

cultured with young Treg in a transwell (OLIG2 (cyan) and MBP (magenta), scale bar=100μm). **D**

Quantification of MBP-expressing oligodendrocytes in control OPCs, OPCs directly co-cultured with

young Treg and OPCs cultured with young Treg in a transwell (n=6, 1-way ANOVA after arcsin

conversion, Sidak's multiple comparison's test). **E** Diagram summarising bioinformatic approaches to

identify protein-protein interactions between OPCs and Treg. **F** Graph showing 21 protein candidates

expressed in the Treg plasma membrane, that are downregulated in aged Treg and have potential

bindings partners enriched in OPCs vs oligodendrocytes. Log$_2$ Change, -Log$_{10}$ (Padj) and the number

of OPC binding partners are indicated (see legend). **G** Bar graphs showing RNAseq normalised count

values for the top 6 candidates.

Figure 8: *Mcam* and *Itga2* contribute to Treg-driven OPC differentiation and are downregulated in

aged Treg. A Representative images of immunostaining showing OPC differentiation in co-culture

with young Treg in the presence or absence of neutralising antibodies against candidate cell surface

mediators (scale bar = 100 μm). **B, C** Bar graphs showing the quantification of OPC differentiation

measured by the fold change in percentage of MBP$^+$ cells and MBP$^+$ area per well (n=7, 2-way

ANOVA, Dunnett's multiple comparison tests). **D** Representative images of immunostaining showing

OPC differentiation in neonatal OPCs treated with recombinant ITGA2 (OLIG2 (cyan), CNPase (grey)

and MBP (magenta), scale bar = 50 μm). **E, F** Bar graphs showing quantification of the proportion of

MBP-expressing oligodendrocytes and total MBP area per well in rITGA2-treated neonatal OPCS at 3

days *in vitro* (n=7 mice, 2 independent experiments, 1-way ANOVA, Dunnett's multiple comparison

test). **G** Sholl analysis showing the number of intersections as a function of the distance from the cell

body to determine changes in the morphology of MBP-expressing oligodendrocytes (n=7 mice, 3

wells per mice and 6-9 cells per well were analysed, 2-way ANOVA).

Supplementary information

Supplementary Figure 1: CD4$^+$ and CD8$^+$ T cells are increased in the aged CNS. A-D Violin plot-based

quantification showing the proportion and count of CD4$^+$ and CD8$^+$ T cells in the spinal cord of

healthy young and aged mice. **E-H** Violin plot-based quantification showing the proportion and count

of CD4$^+$ and CD8$^+$ T cells in the brain of healthy young and aged mice (n=8-13 mice, unpaired two-

tailed Student's t test after *arcsin* conversion).

Supplementary Figure 2: OPC differentiation is impaired in the aged CNS. A Representative images

of OLIG2 (cyan) and CC1 (magenta) in the healthy white matter of young and aged mice (scale bar =

50 μm). **B-D** Bar graph quantification showing OLIG2$^+$ oligodendrocyte lineage cell density,

OLIG2$^+$CC1$^+$ oligodendrocyte density and the proportion of differentiated oligodendrocytes in young

and aged healthy white matter (n=6-8 mice, unpaired two-tailed Student's t-test). **E** Representative

image of OLIG2 (cyan) and CC1 (magenta) immunostaining in lysolecithin-induced demyelinated

white matter of young and aged control mice (scale bar = 50 μm, lesion is demarcated by the white

dotted line). **F** Bar graph quantification showing OLIG2$^+$CC1$^+$ oligodendrocyte density in the

demyelinated lesion area in young and aged control mice (n=4 mice, unpaired two-tailed Student t-

test). **G, H** Bar graph quantification showing the density of CD4$^+$ T cells and CD4$^+$Foxp3$^+$ Treg in the

demyelinated area of young control and Treg expanded mice (n=3 mice, unpaired two-tailed

Student's t-test). **I, J** Bar graph quantification showing the density of CD4$^+$ T cells and CD4$^+$Foxp3$^+$

Treg in the demyelinated area of aged control and Treg expanded mice (n=6-7, unpaired two-tailed

Student's t-test).

Supplementary Figure 3: Endogenous Treg depletion and reconstitution during lysolecithin-

induced demyelination *in vivo*. A Gating strategy followed to identify natural Treg. **B** Bar graph

showing natural Treg isolation purity for *in vitro* experiments and *in vivo* adoptive transfer studies.

Natural Treg are identified by CD4 and CD25 expression (average 91% of the isolated cells for both

young and aged are CD4$^+$CD25$^+$) and by the expression of CD4 and Foxp3 (average 87% of the

isolated cells are positive for CD4 and Foxp3 in both young and aged mouse isolations) (n=14

independent isolations for different adoptive transfer rounds, each pool comprises 2-8 different

mice, Student's t test, after *arcsin* conversion). **C** Flow cytometric plot showing CD4 and GFP

expression to identify endogenous natural Treg in lymph nodes. **D** Bar graph showing quantification

for the proportion of endogenous GFP$^+$ Treg within the CD4$^+$ T cell population in the lymph nodes of

young Foxp3-DTR mice at 14dpl (n=7-9, 1-way ANOVA, after *arcsin* conversion, Sidak's multiple

comparisons test). **E** Flow cytometric plots showing CD25 and Foxp3 expression to detect Treg

reconstitution in lymph nodes after adoptive transfer. **F** Bar graph quantification shows the

proportion of CD25$^+$Foxp3$^+$ cells within the CD4$^+$ T cell population present in the lymph nodes of

young Foxp3-DTR mice at 14dpl (n=7-12, Kruskal Wallis and Dunn's multiple comparisons test). **G**

Representative images of immunostaining showing proliferating OPCs at 14 dpl (OLIG2 (green) and

Ki67 (magenta)) (scale bar = 100 µm, demyelinated area is indicated by the white line). **H-J** Bar graph

showing the areas of demyelination, the density of proliferating OPCs in the lesion and the

proportion of oligodendrocyte lineage cells proliferating in response to damage (n=7-8 mice, 1 way

ANOVA).

Supplementary Figure. 4: A Representative images of immunostaining for neurofilament-H (NFH,

green) and MBP (magenta) to quantify myelin wrapping as an early indicator of remyelination at 14

dpl (scale bar = 50 μm, demyelination area is highlighted by the white dotted line). **B-D**

Quantification shows the total number of axons, the density of MBP wrapped axons and the

percentage of MBP-wrapped axons from the total number of axons (n=4-6, 1-way ANOVA, Sidak's

multiple comparisons test). **E, F** Bar graph quantification showing reconstitution of FOXP3$^+$ Treg

within the CD4$^+$ population and the proportion of CD25$^+$FOXP3$^+$ Treg in CD4$^+$ T cells in the lymph

nodes of aged Foxp3-DTR at the point of sacrifice (n=4-7, Kruskal-Wallis, Dunn's multiple comparison

test).

Supplementary Figure 5: Downstream analysis of differentially expressed genes between young

and aged natural Treg. A Bar graphs showing the purity of young and aged natural Treg subjected to

RNA sequencing (on average 89% of CD4$^+$ cells were Foxp3$^+$, while 94% of CD4$^+$ cells were CD25$^+$)

(n=4-6 mice, Mann-Whitney U test). **B** Bar graph showing normalised counts of natural Treg marker

transcripts *Foxp3*, *Cd4* and *IL2ra*. **C** Heatmap of normalised count values of the Treg tissue repair

programme in male and female Treg. **D** Quantification of normalised count values of the Treg tissue

repair programme in male and female Treg. **E-G** Bar graphs showing the GO cellular component,

molecular function and KEGG pathways enriched within the differentially expressed genes

upregulated (magenta) and downregulated (purple) in aged Treg.

Supplementary Figure 6: MCAM does not directly promote OPC differentiation. A, B Bar graphs

showing the quantification of OPC differentiation in OPC and young Treg co-cultures in the presence

or absence of neutralising antibodies against candidate cell surface mediators, measured by the fold

change in the percentage of MBP$^+$ cells and the fold change in MBP$^+$ area per well (n=5-7, 2-way

ANOVA, Dunnet's multiple comparisons test). **C** Bar graph showing quantification of the proportion

of proliferating OPC (Ki67$^+$ cells) in rMCAM-treated neonatal OPCS at 6 days *in vitro* (n=7 mice, 2

independent experiments, 1-way ANOVA). **D** Bar graph showing quantification of the proportion of

proliferating OPC (Ki67$^+$ cells) in rITGA2-treated neonatal OPCS at 3 days *in vitro* (n=7 mice, 2

independent experiments, 1-way ANOVA, Dunnett's multiple comparison test). **E** Representative

images of immunostaining showing OPC differentiation in neonatal OPCs treated with recombinant

MCAM (OLIG2 (cyan), CNPase (grey) and MBP (magenta), scale bar = 50 µm). **F, G** Bar graphs

showing quantification of the proportion of MBP-expressing oligdoendrocytes and total MBP area

per well in rMCAM-treated neonatal OPCS at 6 days *in vitro* (n=7 mice, 2 independent experiments,

1-way ANOVA).

Supplementary Figure 7: MCAM and ITGA2 levels are restored in aged Treg upon adoptive transfer

into young mice. A Diagram explaining the experimental design of *in vivo* Treg depletion and RFP[+]

Treg adoptive transfer. **B** Gating strategy for flow cytometric analyses**. C** Flow cytometric plots

showing MCAM expression in splenic RFP[+] Treg in control young and aged Foxp3-RFP mice and

young Foxp3-DTR mice that received young or aged RFP[+] cells (data shows 670 cells from the

concatenation of 2-3 mice in each). **D** Bar graph quantification of the proportion of MCAM[+] cells

within RFP population (n=2-3 mice, no statistical analysis performed). **E** Bar graph showing the

median fluorescent intensity of MCAM in MCAM[+]RFP[+] cells (n=2-3 mice, no statistical analysis

performed). **F** Flow cytometric plots showing ITGA2 expression in splenic RFP[+] Treg in control young

and aged Foxp3-RFP mice and young Foxp3-DTR mice that received young or aged RFP[+] cells (data

shows 670 cells from the concatenation of 2-3 mice in each). **G** Bar graph quantification of the

proportion of ITGA2[+] cells within RFP population (n=2-3 mice, no statistical analysis performed). **H**

Bar graph showing the median fluorescent intensity of ITGA2 in MCAM[+]RFP[+] cells (n=2-3 mice, no

statistical analysis performed). statistical analysis performed).

Supplementary Table 1: Statistical analysis of the differentially expressed genes between young and

aged Treg. Related to Fig. 6 and Sup. Fig. 5.

Acknowledgements

We acknowledge extensive technical support from Carmel McVicar, Maria P Athanasios, Michelle

Naughton and the staff of the animal facility. We thank A. Rudensky (Memorial Sloan Kettering

Cancer Centre) for providing Foxp3-DTR mice. We thank the laboratory of Anna Williams (University

of Edinburgh) for the Fiji plug-in to analyse myelination index in brain slice cultures.

This work was supported by the Wellcome Trust (110138/Z/15/Z to DCF), Biotechnology and

Biological Sciences Research Council (BB/J01026X/1 and BB/N003721/1, to DCF), ECTRIMS

postdoctoral fellowship (to AGF), Wellcome ISSF fellowship (to AGF), Miguel Servet Fellowship from

the Spanish Institute of Health Carlos III (CP21/00032 to AGF), Spanish State Research Agency Plan

Generación de Conocimiento 2021 grant (PID2021-124465OA-I00 to AGF), The Leverhulme Trust

(ECF-2014-390, to YD), a postdoctoral fellowship contract from the Generalitat Valenciana

(APOSTD21, to FJR and postgraduate studentship support from the Dept. for the Economy (Northern

Ireland).

Author contributions

958 Experiments were designed by A.G.F., D.C.F., A.L., R.J.M, J.D Y.D. and V.T. Experiments were

959 performed and/or analysed by A.G.F., M.D., E.H., N.V.G., J.W., A.Y., K.M., J.F., S.C.F., T.M., A.D.,

960 F.J.R.B., K.S.R., C.E.M., M.I., J.D., R.P., Y.D. and D.C.F. A.L, R.J.I, R.J.M and C.E.M. provided advice on

961 experimental design and interpretation. Manuscript was written by A.G.F. and D.C.F. with

962 contributions from all authors. D.C.F and A.G.F. oversaw the study.

References

1. Nave, K.-A. & Werner, H. B. Ensheathment and Myelination of Axons: Evolution of Glial Functions.
Annu Rev Neurosci **44**, 197-219 (2021).

2. Griffiths, I. *et al.* Axonal swellings and degeneration in mice lacking the major proteolipid of
myelin. *Science* **280**, 1610–1613 (1998).

3. Lappe-Siefke, C. *et al.* Disruption of Cnp1 uncouples oligodendroglial functions in axonal support
and myelination. *Nat Genet* **33**, 366–374 (2003).

4. Love, S. Demyelinating diseases. *J Clin Pathol* **59**, 1151-9 (2006).

5. Bodini, B. *et al.* Dynamic Imaging of Individual Remyelination Profiles in Multiple Sclerosis. *Ann*
Neurol **79**, 726–738 (2016).

6. Franklin, R. J. M. & ffrench-Constant, C. Regenerating CNS myelin — from mechanisms to
experimental medicines. *Nat Rev Neurosci* **18**, 753–769 (2017).

7. MD, P. C. L., MD, B. Z., MD, P. A. W., MD, P. C. S. & MD, P. B. S. Remyelination in multiple sclerosis:
from basic science to clinical translation. *Lancet* **19**, 678–688 (2020).

8. Ricigliano, V. A. G. *et al.* Spontaneous remyelination in lesions protects the integrity of
surrounding tissues over time in multiple sclerosis. *Eur. J. Neurol.* **29**, 1719–1729 (2022).

9. Shields, S., Gilson, J., Blakemore, W. & Franklin, R. Remyelination occurs as extensively but more
slowly in old rats compared to young rats following fliotoxin-induced CNS demyelination. *Glia* **29**,
1098-1136 (2000).

10. Sim, F. J., Zhao, C., Penderis, J. & Franklin, R. J. M. The age-related decrease in CNS remyelination
efficiency is attributable to an impairment of both oligodendrocyte progenitor recruitment and
differentiation. *Journal of Neuroscience* **22**, 2451–2459 (2002).

11. McMurran, C. E. *et al.* Remyelination in humans due to a retinoid-X receptor agonist is
age-dependent. *Ann Clin Transl Neur* **9**, 1090-1094 (2022)

12. Confavreux, C. & Vukusic, S. Age at disability milestones in multiple sclerosis. *Brain* **129**, 595–605
(2006).

13. Neumann, B. *et al.* Metformin Restores CNS Remyelination Capacity by Rejuvenating Aged Stem
Cells. *Cell Stem Cell* **25**, 473-485.e8 (2019).

14. Fuente, A. G. de la *et al.* Changes in the Oligodendrocyte Progenitor Cell Proteome with Ageing.
Mol. Cell. Proteom. **19**, 1281–1302 (2020).

15. Spitzer, S. O. *et al.* Oligodendrocyte Progenitor Cells Become Regionally Diverse and
Heterogeneous with Age. *Neuron* **101**, 459-471 (2019).

16. Baror, R. *et al.* Transforming growth factor-beta renders ageing microglia inhibitory to
oligodendrocyte generation by CNS progenitors. *Glia* **37**, 2409–11 (2019).

17. Segel, M. *et al.* Niche stiffness underlies the ageing of central nervous system progenitor cells.
Nature **350**, 1199–134 (2019).

18. Ruckh, J. M. *et al.* Rejuvenation of regeneration in the aging central nervous system. *Cell stem*
cell **10**, 96–103 (2012).

19. Natrajan, M. S. *et al.* Retinoid X receptor activation reverses age-related deficiencies in myelin
debris phagocytosis and remyelination. *Brain* **138**, 3581–3597 (2015).

20. Cantuti-Castelvetri, L. *et al.* Defective cholesterol clearance limits remyelination in the aged
central nervous system. *Science* **359**, 684-688 (2018).

21. McNamara, N. B. *et al.* Microglia regulate central nervous system myelin growth and integrity.
Nature 613, 120-129 (2022).

22. Rawji, K. S., Neumann, B. & Franklin, R. J. M. Glial aging and its impact on central nervous system
myelin regeneration. *Ann Ny Acad Sci* **1519**, 34-45 (2022).

23. Rawji, K. S., Martinez, G. A. G., Sharma, A. & Franklin, R. J. M. The Role of Astrocytes in
Remyelination. *Trends in neurosciences* **43**, 596–607 (2020).

24. Franklin, R. J. M., ffrench-Constant, C., Edgar, J. M. & Smith, K. J. Neuroprotection and repair in
multiple sclerosis. *Nature Reviews Neurology* **8**, 624–634 (2012).

25. Burzyn, D. *et al.* A special population of regulatory T cells potentiates muscle repair. *Cell* **155**,
1282–1295 (2013).

26. Ali, N. *et al.* Regulatory T Cells in Skin Facilitate Epithelial Stem Cell Differentiation. *Cell* **169**,
1119-1123.e11 (2017).

27. Dombrowski, Y. *et al.* Regulatory T cells promote myelin regeneration in the central nervous
system. *Nat Neurosci* **20**, 674–680 (2017).

28. McIntyre, L. L. *et al.* Regulatory T cells promote remyelination in the murine experimental
autoimmune encephalomyelitis model of multiple sclerosis following human neural stem cell
transplant. *Neurobiology of Disease* **140**, 104868 (2020).

29. Shi, L. *et al.* Treg cell-derived osteopontin promotes microglia-mediated white matter repair
after ischemic stroke. *Immunity* **54**, 1527-1542.e8 (2021).

30. Garg, S. K. *et al.* Aging is associated with increased regulatory T-cell function. *Aging Cell* **13**, 441–
448 (2014).

31. Elyahu, Y. *et al.* Aging promotes reorganization of the CD4 T cell landscape toward extreme
regulatory and effector phenotypes. *Sci adv* **5**, eaaw8330 (2019).

32. Dulken, B. W. *et al.* Single-cell analysis reveals T cell infiltration in old neurogenic niches. *Nature*
571, 205–210 (2019).

33. Kaya, T. *et al.* CD8+ T cells induce interferon-responsive oligodendrocytes and microglia in white
matter aging. *Nat Neurosci* **25**, 1446–1457 (2022).

34. Groh, J. *et al.* Accumulation of cytotoxic T cells in the aged CNS leads to axon degeneration and
contributes to cognitive and motor decline. *Nat Aging* **1**, 357-367 (2021).

35. Pasciuto, E. *et al.* Microglia Require CD4 T Cells to Complete the Fetal-to-Adult Transition. *Cell*
182, 625-640 (2020).

36. Kuswanto, W. *et al.* Poor Repair of Skeletal Muscle in Aging Mice Reflects a Defect in Local,
Interleukin-33-Dependent Accumulation of Regulatory T Cells. *Immunity* **44**, 355–367 (2016).

37. Ximerakis, M. *et al.* Single-cell transcriptomic profiling of the aging mouse brain. *Nature Neurosci*
22, 1696-1708 (2019).

38. Allen, W. E., Blosser, T. R., Sullivan, Z. A., Dulac, C. & Zhuang, X. Molecular and spatial signatures
of mouse brain aging at single-cell resolution. *Cell* **186**, 194-208(2022).

39. McMurran, C. E., Zhao, C. & Franklin, R. J. M. Toxin-Based Models to Investigate Demyelination
and Remyelination. *Methods Mol. Biol. (Clifton, NJ)* **1936**, 377–396 (2019).

40. Dittmer, M. *et al.* Characterization of a murine mixed neuron-glia model and cellular responses
to regulatory T cell-derived factors. *Molec brain* **11**, 25 (2018).

41. Zhang, H., Jarjour, A. A., Boyd, A. & Williams, A. Central nervous system remyelination in culture
— A tool for multiple sclerosis research. *Expl Neurol* **230**, 138–148 (2011).

42. Kim, J. M., Rasmussen, J. P. & Rudensky, A. Y. Regulatory T cells prevent catastrophic
autoimmunity throughout the lifespan of mice. *Nat Immunol* **8**, 191–197 (2007).

43. Guo, Z. *et al.* DCAF1 regulates Treg senescence via the ROS axis during immunological ageing. *J
Clin Invest* **130**, 5893–5908 (2020).

44. Morales-Nebreda, L. *et al.* Aging imparts cell-autonomous dysfunction to regulatory T cells
during recovery from influenza pneumonia. *JCI Insight* **6**, e141690 (2021).

45. Delacher, M. *et al.* Single-cell chromatin accessibility landscape identifies tissue repair program in
human regulatory T cells. *Immunity* **54**, 702-720 (2021).

46. Zhang, Y. *et al.* An RNA-sequencing transcriptome and splicing database of glia, neurons, and
vascular cells of the cerebral cortex. *J Neurosci* **34**, 11929–11947 (2014).

47. Jensen, L. J. *et al.* STRING 8--a global view on proteins and their functional interactions in 630
organisms. *Nucleic Acids Res* **37**, D412-6 (2009).

48. Linehan, E. *et al.* Aging impairs peritoneal but not bone marrow-derived macrophage
phagocytosis. *Aging cell* **13**, 699–708 (2014).

49. Wang, W., Thomas, R., Oh, J. & Su, D. Accumulation of pTreg cells is detrimental in late-onset
(aged) mouse model of multiple sclerosis. *Aging Cell* **21**, e13630 (2022).

50. Louapre, C. *et al.* A randomized double-blind placebo-controlled trial of low-dose interleukin-2 in
relapsing–remitting multiple sclerosis. *J. Neurol.* **270**, 4403–4414 (2023).

51. Muñoz-Rojas, A. R. & Mathis, D. Tissue regulatory T cells: regulatory chameleons. *Nat Rev
Immunol* **21**, 597-611 (2021).

52. Laursen, L. S., Chan, C. W. & ffrench-Constant, C. An Integrin-Contactin Complex Regulates CNS
Myelination by Differential Fyn Phosphorylation. *J Neurosci* **29**, 9174–9185 (2009).

53. Breuer, J. *et al.* Blockade of MCAM/CD146 impedes CNS infiltration of T cells over the choroid
plexus. *J Neuroinflamm* **15**, 236 (2018).

54. Zondler, L. *et al.* MCAM/CD146 Signaling via PLCγ1 Leads to Activation of β1-Integrins in Memory
T-Cells Resulting in Increased Brain Infiltration. *Front Immunol* **11**, 599936 (2020).

55. Larochelle, C. *et al.* Melanoma cell adhesion molecule–positive CD8 T lymphocytes mediate
central nervous system inflammation. *Ann Neurol.* **78**, 39–53 (2015).

56. Olajuyin, A. M., Olajuyin, A. K., Wang, Z., Zhao, X. & Zhang, X. CD146 T cells in lung cancer: its
function, detection, and clinical implications as a biomarker and therapeutic target. *Cancer Cell Int.*
19, 247 (2019).

57. Şeftalioğlu, A. & Karakoç, L. Expression of CD146 adhesion molecules (MUC18 or MCAM) in the
thymic microenvironment. *Acta Histochem.* **102**, 69–83 (2000).

58. Yshii, L. *et al.* Astrocyte-targeted gene delivery of interleukin 2 specifically increases brain-
resident regulatory T cell numbers and protects against pathological neuroinflammation. *Nat*
Immunol **23**, 878–891 (2022).

59. Webster, K. E. *et al.* In vivo expansion of T reg cells with IL-2–mAb complexes: induction of
resistance to EAE and long-term acceptance of islet allografts without immunosuppression. *J Exp*
Medicine **206**, 751–760 (2009).

60. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**,
676–682 (2012).

61. Stirling, D. R. *et al.* CellProfiler 4: improvements in speed, utility and usability. *BMC*
Bioinformatics **22**, 433 (2021).

62. Dao, D. *et al.* CellProfiler Analyst: interactive data exploration, analysis and classification of large
biological image sets. *Bioinformatics* **32**, 3210–3212 (2016).

63. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
Embnet J **17**, 10–12 (2011).

64. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

65. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2013).

66. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

1102

1103

Fig. 1

Fig. 2

Fig. 4

A

Control   Treg depleted   Treg depleted +Young Treg   Treg depleted +Aged Treg

DAPI / OLIG2 / CC1

Aged Foxp3-DTR

- ☐ Control
- ■ Treg depleted
- ☐ Treg depleted+Young Treg
- ■ Treg depleted+Aged Treg

B

OLIG2$^+$ / mm^2

*
**
*
ns

C

OLIG2$^+$KI67$^+$ / mm^2

ns
ns
ns
ns

D

OLIG2$^+$CC1$^+$ / mm^2

***
ns
**
ns

Fig. 5

Fig. 6

A OPC-nTreg co-culture

OLIG2 / NG2 / CD3

B DAPI / OLIG2 / FOXP3 / CD4

C

Control   +Young Treg No Transwell   +Young Treg Transwell

OLIG2 / MBP

D

% of MBP⁺ cells within OLIG2⁺ population

E

Young vs Aged Treg RNAseq → Downreg Aged Treg → Plasma Membrane

Zhang brain RNAseq J Neuro 2014 → Upreg in OPC vs NFO and OLG → Apical Plasma Membrane

→ **STRING analysis protein-protein interaction**

F

Log2 Change Aged/ Young

Genes: *Acvr1c, Adcy1, Adcy6, Ccr7, Ccr9, Cdh22, Cmklr1, Dapk1, Dnajb4, Itga2, Klra1, Klrb1c, Ly6c1, Mcam, Prex2, Prf1, Ryr2, S1pr1, Sell, Scl16a5, Scl6a19*

-Log₁₀(padj)

OPC Binding partners

G

Ccr7 — Normalised counts — Young / Aged — ***

Itga2 — Normalised counts — Young / Aged — ****

Klrb1c — Normalised counts — Young / Aged — *

Ly6c1 — Normalised counts — Young / Aged — ****

Mcam — Normalised counts — Young / Aged — ****

Sell — Normalised counts — Young / Aged — ****

Fig. 7

Sup. Fig. 1

Sup. Fig. 3

Sup. Fig. 5

Sup. Fig. 6

A

B

C

MCAM-PE-Cy7 vs Treatment group

D

% of MCAM⁺ in RFP⁺ cells vs Treatment group

E

RFP⁺MCAM⁺ population

MFI MCAM-PeCy7

F

ITGA2-APC vs Treatment group

G

% of ITGA2⁺ in RFP⁺ cells

H

RFP⁺ITGA2⁺ population

MFI ITGA2-APC

- 1-Young RFP control
- 2-Aged RFP control
- 3-Young RFP adoptive transferred
- 4-Aged RFP adoptive transferred

Sup. Fig. 6