Item-specific memory reactivation during sleep supports memory consolidation in humans

Jing Liu[1,2], Tao Xia[2,3], Danni Chen[2,3], Ziqing Yao[2,3], Minrui Zhu[2,3],

James W. Antony[4], Tatia M.C. Lee[2,3,5*], Xiaoqing Hu[2,3,6*]

1. Department of Applied Social Sciences, The Hong Kong Polytechnic University, Hong Kong

SAR, China

2. The State Key Laboratory of Brain and Cognitive Sciences, The University of Hong Kong,

Hong Kong SAR, China

3. Department of Psychology, The University of Hong Kong, Hong Kong SAR, China

4. Department of Psychology & Child Development, California Polytechnic State University,

California, USA

5. Laboratory of Neuropsychology and Human Neuroscience, Department of Psychology, The

University of Hong Kong, Hong Kong SAR, China

6. HKU-Shenzhen Institute of Research and Innovation, Shenzhen 518057, China

* Correspondence should be sent to:

T.M.C. L., tmclee@hku.hk; X. H., xiaoqinghu@hku.hk;

Abstract

Memory consolidation stabilizes newly acquired information. Understanding how individual

memories are reactivated during sleep is essential in theorizing memory consolidation. Via

unobtrusively re-playing auditory memory cues to sleeping human participants, we identified the

reactivation of individual memories during slow-wave sleep (SWS). Using representational

similarity analysis (RSA) on cue-elicited electroencephalogram (EEG), we found functionally

segregated item-specific representations: the early post-cue EEG activity (0-2 seconds) contained

comparable representations for memory cues and for non-memory control cues, thus reflecting

sensory processing. Critically, the later EEG activity (2.5-3 s) showed greater item-specific

representations for post-sleep remembered items than for forgotten and control cues,

demonstrating the reactivation and consolidation of individual memories. Moreover, spindles

preferentially supported item-specific memory reactivation for items that were not tested before

sleep. These findings delineated how cue-triggered item-specific memory reactivation, subserved

by spindles during SWS, contributed to memory consolidation. These results will benefit future

research aiming to perturb specific memory episodes during sleep.

Introduction

Newly acquired experiences must undergo consolidation to become long-lasting memories (*1*). Studies using various techniques (e.g., single-unit recordings, fMRI, EEG) suggest repeated, covert reactivation of memories occurs during post-learning non-rapid eye movement (NREM) sleep (*2–8*). Intriguingly, memory reactivation can be manipulated by unobtrusively re-presenting sensory cues associated with wakeful learning during subsequent sleep, a paradigm known as targeted memory reactivation (TMR) (*9–12*). TMR allows researchers to determine when and which memories become reactivated via re-playing stimulus-specific memory cues, which shows promise in modulating the declarative, procedural, and emotional memories, in both clinical and educational settings (*10, 13–16*). However, given the challenges in identifying memory-specific neural ensembles during sleep in humans (*17*), how individual TMR cues reactivate their corresponding memory representations and thereby facilitate memory consolidation remains unclear.

Recent studies using multivariate neural decoding methods have advanced our understanding of TMR-based memory consolidation. First, TMR cue-elicited neural activity contained task-related or category-level memory representations (*18–21*). Second, during NREM sleep, neural representations associated with wakeful retrieval re-emerged rhythmically ~1-Hz following TMR cues (*22*). While these findings suggest that memory reactivation can be detected during NREM sleep, whether TMR cueing triggers the fined-grained neural representations for individual memories remains unknown. Moreover, although the sleeping brain remains its robust neural auditory responses to external stimuli (*23, 24*), how processing auditory memory cues drives the exogenous memory reactivation during sleep is yet to be established.

Converging evidence suggests that the thalamocortical spindles are instrumental for both endogenous and exogenous memory reactivation during NREM sleep (*25, 26*). In TMR, enhanced post-cue spindle-related sigma power not only predicted TMR effects (*27, 28*) but is also correlated with the distinctiveness of category-level memory representations (*19*). Furthermore, TMR effects are abolished when post-cue sigma power is reduced (*29*) and when cueing occurs during the spindle refractory period, i.e., ~3-6 s following a spindle when a spindle

is less likely to occur (*27*). This evidence prompts an intriguing question of whether the spindle

in specific time windows can even support memory reactivation for individual items.

An important factor that may modulate TMR effects and spindle-mediated memory consolidation

is pre-sleep testing. Given that pre-sleep testing could enhance memory via retrieval-induced fast

consolidation processes (*30, 31*), testing may reduce the subsequent TMR effects (*32*) and

weaken the relationship between the neural activity and memory consolidation. Indeed, both

spontaneous sleep and sleep TMR preferentially benefit weaker memories, with sleep spindles

consolidating weak rather than strong memories (*33–36*). Alternatively, pre-sleep testing may

enhance the future relevance of the tested items, which would make these memories

preferentially consolidated due to their motivational salience (*37, 38*). To reconcile these

competing hypotheses, we further examined how pre-sleep testing affects item-specific memory

reactivation and spindles.

We used representational similarity analysis (RSA) on scalp electroencephalogram (EEG) (*19,*

39, 40) to extract item-specific representation during slow-wave sleep (SWS). Employing a

subsequent memory analysis approach to link item-specific representations during sleep with

post-sleep memory performance, we identified two functionally distinct time windows.

Specifically, the early time window (0-2 s post-cue) exhibited item-specific representations for

both the memory cues and the non-memory control cues. However, item-specific representations

in this early time window showed no subsequent memory effect or any difference between

memory items and control items, thus primarily reflected sensory processing of individual

auditory cues. More importantly, a later 2500-2960 ms time window bore relevance with post-

sleep memory performance: remembered items exhibited greater item-specific representations

than both forgotten and control cues. Notably, sleep spindles during this time window supported

item-specific memory reactivation only for pre-sleep untested items, highlighting the nuances of

spindle-mediated mechanisms in memory consolidation.

Results

Behavioral results

Thirty participants (23 females, mean ± SD age, 22.37 ± 2.94 years) were included in the

analyses (see Materials and Methods for participants exclusion criterion). Participants first

studied 96 word-picture (cue-target) pairs, with each pair being repeated three times. After a

distractor (math) task, participants completed a pre-sleep cued recall test, with half (48) of the

learning pairs being tested (i.e., tested items) while the other half (48) were untested (i.e.,

untested items). TMR cues consisted of 24 spoken words from each of the tested and untested

items, respectively, resulting in a total of 48 TMR cues. For tested items, pre-sleep cued recall

performance was matched for subsequently cued and uncued items. For untested items, cues

were pseudo-randomly selected (see Materials and Methods). In addition to these 48 TMR cues,

four spoken words that were not paired with any learning content were included as non-memory

control cues, and they were played the same number of times as TMR cues during sleep. On the

following morning, memory was tested for all 96 learned pairs (see Fig. 1).

We first examined how pre-sleep testing affected TMR effects by conducting a 2 (cued vs.

uncued) by 2 (tested vs. untested) repeated measures ANOVA on post-sleep memory recall

performance. Results revealed neither an interaction effect ($F(1,29) = 0.15$, $p = 0.700$) nor the

TMR main effect (cued vs. uncued, $F(1,29) = 0.11$, $p = 0.739$) (see Fig. S1). The same ANOVA

on the post-sleep memory recognition performance similarly yielded no effects (all ps > 0.373,

see Fig. S1). Despite the absence of significant TMR effects at the group level, substantial

individual differences were observed (see Fig. S1), which prompted further investigations into

individual differences analyses of TMR effects. In the following analyses, we used cued-recall

performance, which is more sensitive than the recognition performance to TMR effects (*10*).

Fig 1. Experimental design. The experiment included four main phases, i.e., learning, a pre-sleep test (cued recall and recognition), targeted memory reactivation (TMR) during slow-wave sleep (SWS), and a post-sleep test. During the learning phase, participants studied word-picture pairs, half of which were tested before sleep (blue rectangle) while half were not (orange rectangle). Auditory cues from half of the tested items and half of the untested items, along with four control cues (grey rectangle), were played during SWS of the overnight sleep. Memory for all pairs was assessed during the post-sleep test (cued recall and recognition) the next morning.

Early neural responses to auditory cues predict TMR effects

Previous studies demonstrated that external auditory stimulation during NREM sleep can

increase EEG power in theta (~4-7 Hz), alpha (~8-12 Hz), and sigma (~11-16 Hz) bands (*19, 41,*

42). In line with these studies and confirming that the auditory cues were indeed processed

during sleep, we found that all memory cues (i.e., including both the tested and untested items)

significantly enhanced EEG power in a broad range of 2-40 Hz during the first 1960 ms as

compare to pre-cue baseline ($p_{\text{corr}} < 0.001$, see Fig. 2A). Notably, two prominent frequency

ranges emerged: a low-frequency range of 2-9 Hz and an extended sigma band of 11-18 Hz.

Following this early cluster, we observed a significant reduction of sigma power in a late cluster

(2320-3380 ms, p_{corr} = 0.005), which potentially reflected the spindle refractoriness (*27*). Further

tests suggested that pre-sleep testing did not modulate cue-elicited EEG power in the above

clusters (all ps > 0.102, see Fig. S2). Together, these results suggest that memory cues were

indeed processed by the sleeping brain during NREM sleep.

To further investigate whether the enhanced EEG power reflects memory-specific processing or

sensory processing to auditory cues, we examined the EEG power locked to the non-memory

control cues. Results showed that control cues also enhanced EEG power in the low-frequency

range and the sigma band within the first 2 s (both ps_{corr} < 0.001), followed by sigma power

reduction (3130-4300 ms, p_{corr} = 0.006) (Fig. 2B). Importantly, direct comparisons between the

EEG power elicited by memory cues and control cues after matching trial numbers revealed no

significant differences (p_{corr} > 0.441, Fig. S2). These findings indicated that the early neural

responses reflected how the sleeping brain reacts to external auditory stimuli, regardless of

memory cues or not.

We next examined whether the cue-elicited EEG power could predict the post-sleep TMR effect

(i.e., cued minus uncued recall performance). Using robust linear regression analyses that

minimized the influence of potential statistical outliers, results revealed that memory cue-elicited

EEG power in the low-frequency range (2-9 Hz) and the sigma band (11-18 Hz) in the early

cluster (within first 2s) positively predicted TMR effects (early low-frequency: adjusted R^2 =

0.28, p = 0.002; early sigma: adjusted R^2 = 0.12, p = 0.032, Fig. 2C). More specifically, theta (4-

7 Hz), and alpha (8-12 Hz) power significantly predicted TMR effects (all ps < 0.026). However,

sigma power reduction during the later time window (i.e., 2320-3380 ms) did not predict TMR

effects (adjusted R^2 = -0.004, p = 0.363). Interestingly, EEG power elicited by non-memory

control cues showed similar prediction effects: early, but not late EEG power positively predicted

the TMR effects (early low-frequency: adjusted R^2 = 0.14, p = 0.025; early sigma: adjusted R^2 =

0.22, p = 0.005; late sigma: adjusted R^2 = 0.064, p = 0.097, Fig. 2C). Directly comparing the

prediction effects of memory and control cues showed no significant differences (all ps > 0.159).

Intriguingly, pre-sleep testing remarkably modulated the relationship between cue-elicited EEG

power and TMR effects. Specifically, cue-elicited EEG power significantly predicted TMR

effects only for untested items (early low-frequency: adjusted $R^2 = 0.32$, $p < 0.001$; early sigma:

adjusted $R^2 = 0.17$, $p = 0.014$; Fig. 2D), but not for tested items (both ps > 0.485). More

importantly, this prediction effect was significantly higher for untested items than for tested

items (both ps < 0.026). Again, EEG power elicited by non-memory control cues showed similar

prediction effects for pre-sleep tested and untested items (see Fig. S2). Taken together, these

results suggested that a stronger early neural response to external auditory stimuli during NREM

sleep predicts TMR efficacy. This association was observed regardless of whether these auditory

stimuli were memory cues or non-memory control cues. Furthermore, this association was

particularly strong for untested items but not for tested items.

Fig 2. TMR cue-elicited EEG spectral power changes in the early time window (i.e., within the first 2 seconds after cue onset) and their association with post-sleep TMR effects. (A) Left panel: memory cues enhanced EEG spectral power across all memory items within an early cluster (circled by the black line), followed by decreased sigma power in a late cluster (circled by the green line), compared to the pre-cue baseline. Right panel: topography plots for the memory cue-elicited low-frequency range and sigma band power in the early cluster. **(B)** Left panel: non-memory control cues enhanced EEG spectral power within the two early clusters (circled by the black line), followed by decreased sigma power in a late cluster (circled by the green line), compared to the pre-cue baseline. Right panel: topography plots for the control cue-elicited low-frequency range and sigma band power in the early clusters. **(C)** Both memory cue and control cue-elicited EEG power in the low-frequency range and the sigma band in the early cluster positively predicted post-sleep TMR effects. The prediction effect did not differ between memory and control cues. **(D)** Memory cue-elicited EEG power in the low-frequency range and

207 the sigma band in the early cluster predicted the post-sleep TMR effects for untested but not for
208 tested items. The p-values indicate the significance of robust linear regression that minimized the
209 potential influence of statistical outliers. *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$.
210

Early EEG activity contains item-specific representations for cue processing

Demonstrating that this early EEG power was elicited by both memory and non-memory control
cues, we hypothesize that this early EEG activity might contain neural representations of
individual auditory cues. To test this hypothesis, we performed the representational similarity
analysis (RSA) between trials of the same cues (within-item [WI] similarity) as well as between
trials of different cues (between-item [BI] similarity) (Fig. 3A). We computed the
representational similarity by correlating the cue-elicited raw EEG pattern (filtered between 0.5-
40 Hz) across all channels within consecutive overlapping time windows of 500 ms, with a 10
ms sliding step, during a 5-second time window after cue onset. The WI and BI similarity were
calculated between trials from two different TMR blocks to control for the temporal proximity
effect. Item-specific representations were indicated by significantly greater WI similarity than BI
similarity. Our analysis revealed significant item-specific representations during 560-1350 ms
after cue onset ($p_{\mathrm{corr}} = 0.008$, Fig. 3B). The same analyses on control items similarly revealed
significant item-specific representations during 1440-1950 ms after cue onset ($p_{\mathrm{corr}} = 0.040$, Fig.
3B). Notably, the relatively delayed onset of item-specific representations for control cues
compared to memory cues may reflect the cost of processing speed for the control cues given
they were less studied pre-sleep (43). Moreover, a repeated measures ANOVA on item type
(memory vs. control) and item-specificity (WI vs. BI) did not reveal any significant clusters
($p_{\mathrm{corr}} > 0.085$).

We next performed the linear regression analysis between control cue-elicited EEG power, which
were sensory-cue driven neural responses, and memory cue-elicited item-specific
representations. The results revealed that both the control cue-elicited low-frequency range and
sigma band EEG power in the early time window significantly and positively predicted the
memory cue-elicited item-specific representations in the 560-1350 ms time window (Low-
frequency: adjusted $R^2 = 0.18$, $p = 0.013$; Sigma: adjusted $R^2 = 0.12$, $p = 0.036$; see Fig. 3C-D).
Additional control analyses also found that memory cue-elicited EEG power in an earlier non-
overlapped time window positively predicted the following item-specific representations for

memory items ($p < 0.001$, see Fig. S3). A similar significant positive prediction effect was also

observed for control items ($p = 0.024$, see Fig. S3).

These results provided converging evidence that the early enhanced EEG power within the first 2

s reflected effective processing of individual auditory cues. Furthermore, the control cue-elicited

EEG power predicted item-specific representations of memory items, highlighting that the level

of the sleeping brain responding to external auditory cues is crucial in shaping item-specific

representations of these cues.

Fig 3. Item-specific representations following TMR cues during slow-wave sleep. (A)
Representational similarity analysis scheme across different TMR blocks. Artifact-free raw EEG
data pattern was correlated between trials with the same auditory cues (within-item similarity,
WI) and between trials with different auditory cues (between-item similarity, BI). Item-specific
representations were examined by contrasting the WI similarity versus BI similarity. **(B)** Item-
specific representations for memory items and control items were significant in the time
windows 560-1350 ms (blue horizontal bar on the top) and 1440-1960 ms (gray horizontal bar on
the top) after cue onset, respectively. **(C-D)** Control cue-elicited low-frequency EEG power and
sigma band EEG power in the early cluster (< 2 s, see Fig. 2B) predicted the item-specific

representations for memory items in the 560-1350 ms time window, respectively. *: $p < 0.05$; **:
$p < 0.01$.
260

Later EEG activity contain item-specific representation for memory reactivation

We next examined our key question: whether TMR cues reactivate item-specific memory
representations during sleep that mediate memory consolidation. Adopting a subsequent memory
approach, we conducted a memory (remember vs. forget) by item-specificity (WI vs. BI)
repeated measures ANOVA. The results indeed revealed a significant interaction effect during
the 2500-2960 ms time window ($p_{corr} = 0.025$, Fig. 4A), suggesting that TMR cues reactivate
item-specific memory representations. Notably, the observed item-specific representations were
above and beyond category-level representations, given that the WI similarity was greater than
the BI similarity even when the BI items were drawn from within-category trial pairs (i.e.,
within-category similarity, WC, see Fig. S4).
271

To further confirm that item-specific representations in the late time window were memory-
related, we compared remembered items with non-memory control cues in an item type
(remember vs. control) by item-specificity (WI vs. BI) two-way repeated measures ANOVA. The
results also showed a significant interaction in a highly overlapped late time window ($p_{corr} =$
0.044, see Fig. S5). Post-hoc analysis [corrected for family-wise error rate (FWER)] on the late
item-specific memory reactivation time window revealed that WI similarity was significantly
greater than BI similarity for remembered items ($t(29) = 3.01$, $p_{FWER} = 0.016$), while there was no
difference between WI and BI similarity for forgotten items ($t(29) = -1.28$, $p_{FWER} = 0.637$) or for
control cues ($t(29) = -0.95$, $p_{FWER} = 1.000$) (Fig. 4B). Furthermore, item-specific representation
(WI - BI similarity) for remembered items was greater than control items ($F(1,29) = 6.31$, $p =$
0.018), while there was no difference between item-specific representations for forgotten and
control items ($F(1,29) = 0.20$, $p = 0.657$). Notably, results remained significant after controlling
for trial pair number differences between WI and BI conditions and between memory and control
items (Fig. S6). In addition, we found that pre-sleep testing did not modulate the item-specific
representations ($p_{corr} > 0.383$, see Fig. S7).
287

Taken together, cue-elicited EEG activity in the early time window contained item-specific
representations for both memory and control items, which may primarily reflect the effective

processing of auditory stimuli during SWS. In contrast, the late time window item-specific

representations emerged specifically for the post-sleep remembered items, which reflected the

memory reactivation processes that support memory consolidation.

Fig 4. Item-specific memory reactivation (A) Item-specificity by post-sleep memory repeated
measures ANOVA revealed a significant interaction effect cluster in 2500-2960 ms time window
(shaded rectangle). **(B)** Post-hoc analyses of the interaction cluster (the shaded rectangle in A)
found WI similarity significantly greater than the BI similarity for remembered items, while
there were no differences between WI and BI similarity for forgotten items or control items.
Item-specific representations for remembered items were greater than both the forgotten items
and control items, while the item-specific representations for forgotten items were not
significantly different from the control items. *: $p < 0.05$; ***: $p < 0.001$; n.s.: not significant.

Spindle activity coordinated item-specific memory reactivation for successfully

remembered untested items

Previous studies suggest that TMR-based memory consolidation is tightly linked with post-cue

spindle activity (*19, 27*). However, it remains unclear how spindle activity is related to memory

reactivation and whether pre-sleep testing could affect the role of spindles in supporting memory

reactivation. To address this question, we first detected discrete spindles on individual trials (Fig.

5A-B, see Materials and Methods) and then calculated spindle probability across all trials for

each time point. We conducted pre-sleep testing (tested vs. untested) by subsequent memory

(remember vs. forget) repeated measures ANOVA across individual time points and identified

two significant clusters showing significant interaction effects (1st cluster: 2344-3166 ms, p_{corr} =

0.002; 2nd cluster: 3526-4118 ms, p_{corr} = 0.026; Fig. 5C). Post-hoc comparisons were performed

by contrasting spindle probabilities across the two clusters. The results showed that, among tested items, the spindle probability for remembered items was not different from that for forgotten items ($t(29) = -2.50$, $p_{FWER} = 0.147$, Fig. 5D). In contrast, among the untested items, spindle probability for remembered items was greater than for forgotten items ($t(29) = 4.50$, $p_{FWER} < 0.001$). Moreover, post-sleep remembered untested items elicited higher spindle probability than both post-sleep remembered tested items ($t(29) = 4.51$, $p_{FWER} < 0.001$) and control items ($t(29) = 5.69$, $p_{FWER} < 0.001$, Fig. 5D), while no significant difference was found between the post-sleep remembered tested items and control items ($t(29) = 1.42$, $p_{FWER} = 1.000$). To ensure that the observed spindle activity difference in the late cluster was not due to the leakage of longer duration or delayed onset of the early spindles, we examined the duration and onset of spindles from the early (0-2 s) time window. The results revealed no significant interaction effect between memory and pre-sleep testing for either the duration or the onset of early spindles (all $ps > 0.370$, see Fig. S8). In addition, spindles that occurred in an extended time window of these two clusters (i.e., 2200-4200 ms) were significantly coupled to the up-state of SOs for all tested and untested items (all $ps < 0.045$) (see Fig. S9).

We next sought to establish the relationship between spindle activity and item-specific memory reactivation. Notably, increased spindles (2344-3166 ms and 3526-4118 ms in Fig. 5C) for post-sleep remembered untested items tended to co-occur in the late item-specific memory reactivation window (2500-2960 ms in Fig. 4A). Importantly, repeating the RSA using the spindle-related sigma power did not find any item-specific representations ($p_{corr} > 0.111$, Fig. S10), ruling out the possibility that late item-specific representations and spindles were different measures of the same phenomena. We then examined how spindles related to item-specific memory reactivation in the overlapping time window (i.e., 2500-2960 ms) for tested and untested items, respectively. Our linear mixed-effect model revealed that there were no significant effects for subsequently remembered or forgotten items among tested items (remember: $\beta = 0.02$, $t(432) = 0.28$, $p = 0.783$; forget: $\beta = -0.01$, $t(258) = -0.16$, $p = 0.875$, Fig. 5E). However, among untested items, cue-elicited spindle probability significantly and positively predicted item-specific memory reactivation for subsequently remembered items ($\beta = 0.26$, $t(239) = 3.678$, $p < 0.001$, Fig. 5F) but not for forgotten items ($\beta = -0.02$, $t(472) = -0.21$, $p = 0.836$). Moreover, among untested items, the effect for remembered items was significantly greater than that for

forgotten items ($\beta = 0.29$, $t(716) = 2.57$, $p = 0.010$). These results suggest that sleep spindles play

a critical role in supporting item-specific memory reactivation and contribute to memory

consolidation, particularly for untested items.

Fig 5. The relationship between spindles and item-specific memory reactivation in the
overlapped late time window (i.e., 2500-2960 ms relative to cue onset). (A) An example of
raw EEG data used for spindle detection. **(B)** EEG signals in (A) filtered between 11-16 Hz, with
a spindle event highlighted in red. **(C)** Significant interaction effects between pre-sleep testing
(tested vs. untested) and subsequent memory effect (remembered vs. forgotten items) on spindle
probability in two clusters (2344-3166 ms and 3526-4118 ms, highlighted in shaded rectangles)
following TMR cues. **(D)** Post-hoc analyses were performed on the averaged data across the two
interaction clusters (shaded rectangle in C). For untested items, post-sleep remembered items
elicited greater spindle probability than forgotten items. Moreover, the spindle probability for
post-sleep remembered untested items was greater than that for both the post-sleep remembered
tested items and control items. **(E)** Among tested items, the spindle probability did not predict
the item-specific memory reactivation for either post-sleep remembered or forgotten items. **(F)**

Among untested items, the spindle probability significantly predicted the item-specific memory
reactivation for post-sleep remembered items but not for forgotten items. The effect for post-
sleep remembered items was greater than that for forgotten items. *: $p < 0.05$; **: $p < 0.01$; ***:
$p < 0.001$.
368

Discussion

We asked fundamental yet unanswered questions in exogenous memory reactivation: how

individual sensory memory cues reactivate the corresponding memory traces during slow-wave

sleep and how such item-specific reactivation led to consolidation. Using targeted memory

reactivation (TMR) and multivariate representational similarity analyses (RSA), we found highly

specific item-level representations, which went beyond prior studies showing neural

reinstatement effects of broader, categorical types (*19–21*). Specifically, both memory cues and

non-memory control cues first elicited item-specific representations, likely reflecting the

successful sensory processing of individual auditory cues. Item-specific memory reactivation

emerged in a late time window that was only present for memory cues, which contributed to

post-sleep memory retention. Critically, increased spindles during this late memory-specific

window preferentially supported item-specific memory reactivation only among pre-sleep

untested items, highlighting that testing can be an important factor that modulates sleep-based

memory consolidation (see Fig. 6).

Fig 6. Schematic depiction of item-specific memory reactivation following TMR cues promotes memory consolidation. Sensory memory cues trigger early item-specific representations during SWS, predominantly reflecting sensory processing. In contrast, item-specific memory reactivation and spindle activity within slow-oscillation up-states in a later time window contribute to memory retention for pre-sleep untested items.

Even during SWS, the brain preserves its capacity to process external sensory information, which is a prerequisite for sensory memory cues to reactivate its associated memories (*23*). Cross-species studies have consistently demonstrated that the primary auditory cortex neuronal responses are well preserved during sleep, with little modulation by different vigilance or consciousness states (*24*, *44*, *45*). Humans research further shows that the sleeping brain can process meaningful auditory stimuli (*46–49*), which are manifested by increased EEG power in both the low-frequency and sigma band immediately following auditory stimuli (*41*, *42*). Consistent with these previous studies, we found that both the memory cues and non-memory control cues enhanced low-frequency activities (1-10 Hz), followed by the extended sigma band activities (11-18 Hz) within approximately the first 2 s post-cue onset. Beyond EEG spectral power changes, our results further revealed that the cue-elicited EEG contained highly specific,

item-specific representations in this early cluster. Moreover, this early item-specific

representation was identified for all memory cues regardless of post-sleep memory performance

and for control cues that were not paired with any prior learning. These early, item-specific

representations may thus reflect the initial processing of individual auditory cues.

While previous studies often found greater EEG power in response to memory cues than control

cues (*19*, *50*), we did not find significant differences. This discrepancy might be caused by the

novelty of the control cues: whereas in these previous studies, control cues were not presented

before the TMR session; here, we presented all memory and control cues at least twice during the

pre-learning familiarization phase. One possibility is that such familiarization would evoke

strong K-complex activities in the sleeping brain (*51*–*53*), thus minimizing the differences

between memory and control cues. Notably, these early EEG responses elicited by the cues

predicted the subsequent behavioral TMR effects, despite almost unitary presentations of

auditory cues across participants. These findings are consistent with previous research that has

linked this early cue-elicited EEG power with behavioral TMR effects (*29*, *50*, *54*, *55*).

Moreover, our results showed that even the increased EEG power elicited by non-memory

control cues could predict the early item-specific representations of the individual memory cues.

Our findings provided new insights that how early EEG responses reflect sensory processing of

memory cues, which paves the pathways for successful memory reactivation in later time

windows.

Indeed, we found compelling evidence that item-specific representations in a late time window

reflected the item-specific memory reactivation. Specifically, post-sleep remembered items

exhibited greater item-specific representations than forgotten items and control cues in this late

cluster. The differences in item-specific representations cannot be exclusively attributed to

different levels of familiarity with sensory cues, given that the cues for post-sleep remembered

and forgotten items were repeated the same number of times during pre-sleep learning. In

addition, the direct comparisons between the item-specific representations of forgotten items and

control items did not yield significant differences. Notably, this late item-specific memory

reactivation window overlaps with the spindle refractory period, highlighting that this period is

instrumental for memory reprocessing during NREM sleep (*25*).

Item-specific memory reactivation following TMR cues could be achieved via a cortical-
hippocampal-cortical loop (*56–58*): effective early processing of auditory cues in the neocortex
reactivate their associated memory traces that are temporarily stored in the hippocampus (*11*, *59–
61*). The reactivated hippocampal memory traces then trigger the reinstatement of the neocortex-
dependent memory representations, which may follow the same principle of pattern completion
as during wakefulness retrieval (*62*, *63*). This cortical-hippocampal-cortical loop during NREM
sleep relies on the timely coupling of thalamocortical spindles to the up-state of slow oscillations
in both humans and rodents, serving as the foundation for systems memory consolidation (*3*, *64–
67*). Critically, our results showed that item-specific memory reactivation, supported by spindles,
during the late window predominantly contributes to memory consolidation. Moreover, the
accompanying spindles were preferentially coupled to the up-state of slow oscillations, which
may have facilitated cross-regions interactions for reactivated memories, particularly the pre-
sleep untested items, to be consolidated (*68–70*). A more precise delineation of the spindle-SO-
coordinated cross-regional interaction awaits future investigations using methods affording both
high spatial and temporal resolutions (e.g., intracranial EEG recordings).

Despite spindles' importance in supporting item-specific memory reactivation, pre-sleep testing
modulated this spindle-mediated consolidation process. Given that pre-sleep testing largely
boosted memory performance relative to untested items in the current study, it is possible that
pre-sleep tested items already underwent retrieval-induced fast consolidation before sleep (*30*).
Thus, spindles may preferentially support item-specific memory reactivation for untested
memories. Corroborating this possibility, previous studies have shown that strong memories did
not further benefit from subsequent spontaneous sleep (*71*) or sleep TMR (*34*). Moreover, sleep
spindles would preferentially consolidate weak over strong memories (*36*), and TMR promoted
consolidation for memories with low (*33*) or moderate (*34*) pre-sleep accuracy. Consistent with
this idea, our results also showed cue-elicited early EEG power was associated with TMR effects
only for pre-sleep untested items.

Our study did not find a significant TMR behavioral effect. A few possibilities might account for
the absence of TMR effects. First, as mentioned above, TMR may be less effective for strong

memories (*33*, *34*). Here, participants may overlearn the word-picture pairs via repeated

encoding and mental rehearsal during the pre-sleep learning, which may contribute to fast

consolidation and thus weaken the TMR effects (*30*). In addition, the pre-sleep recognition task

might serve as the feedback after the pre-sleep test, which further strengthens the pre-sleep

memory for tested items and attenuates the TMR effect. Second, given that TMR effects are

highly sensitive to memory measures (*10*), more precise measurements are desirable to detect the

TMR effects, such as the error distance measures used in previous TMR research (*11*, *12*). Future

word-picture or naturalistic episodic memory paradigms could also include verbal recall to test

how TMR impacts perceptual and conceptual details.

In conclusion, the identification of item-specific memory reactivation and the segregated

processing windows contributed to the mechanistic understanding of how exogenous memory

reactivation unfolds during NREM sleep. Effective processing of individual TMR cues would

drive the reactivation of item-specific memory representations for consolidation, supported by

spindles in a later time window. These findings may ignite new development of sleep-based

memory editing techniques (*72*) in perturbing individual memories: by targeting the underlying

neural activity at the critical memory reactivation time window, techniques can either strengthen

newly acquired knowledge or selectively weaken maladaptive memories.

Materials and Methods

Participants

Thirty healthy, right-handed participants were included in the analyses (23 females, mean ± SD

age, 22.37 ± 2.94 years). To obtain reliable results in the EEG power spectral analysis as well as

the multivariate representational similarity analysis for both the memory items and control items,

we set a minimum requirement of 5 rounds of TMR cueing (i.e., 20 trials). Thus, seven

additional participants failed to meet this criterion due to insufficient or unstable N2 and N3

sleep were excluded from the analysis. All included participants had normal or corrected-to-

normal vision. Participants were pre-screened on sleep by using the questionnaires including the

Pittsburgh Sleep Quality Index (PQSI) and the Insomnia Severity Index (ISI). They all reported

overall good sleep quality and had not taken any medicines to aid sleep in the past month before

the experiment. All participants did not suffer from any neurological or psychiatric disorders.

The study was approved by the Research Ethics Committee of the University of Hong Kong. All

participants gave written informed consent prior to the experiment.

Stimuli

A total of 96 two-character Chinese verbs and 96 target pictures were used in the experiment.

Each verb was randomly paired with a complex visual picture, resulting in 96 cue (word)-target

(picture) pairs. The central element of each picture was from one of four categories (i.e., animals,

electronic devices, plants, and transportation tools), with 24 pictures in each category. For each

picture, a highly similar picture was also selected and served as a lure in the old/new recognition

task. For each unique picture, the target and lure were randomly assigned among participants.

For verbs, visually presented verbs were only used in the familiarization phase, while aurally

presented verbs were used throughout the entire experiment. Four additional two-character

Chinese verbs were presented during the familiarization phase but were never paired with any

pictures. These verbs served as control cues in the TMR. Auditory sounds of the verbs were

generated using the Text-To-Speech of iFLYTEK, with an average duration of 631.70 ms (SD:

55.40 ms).

Procedure

All participants arrived at the sleep lab around 8:30 pm. Participants completed the following

tasks in order: a vigilance task, a stimuli familiarization task, a cue-target associative learning

task, and a pre-sleep memory test. Participants then proceeded to sleep (12 am to 8 am the next

morning), wherein TMR was administered during slow-wave sleep. After ~30 min upon

wakening, participants' vigilance levels were assessed again, followed by the post-sleep memory

test. All behavioral tests were conducted via using the psychopy (version 2020.2.10).

Psychomotor vigilance task

Participants' vigilance levels were assessed using the Psychomotor Vigilance Task (PVT), right

after participants arrived at the sleep lab and the next day morning. During the vigilance task, a

fixation was presented on the center of the screen with a jitter duration in the range of 2-10 s.

Then, the fixation was replaced by a counter counting incrementally from 0 in 10 ms increments.

Participants were instructed to stop the counter by pressing the space bar immediately after they

detected the change of fixation to the number. Response time that appeared on the screen would

serve as feedback on the performance. This task lasted for 5 minutes.

Cue word and picture familiarization task

The familiarization task consists of two sessions, a cue word-familiarization session, and a

picture-familiarization session. In the cue word-familiarization session, each trial started with a

0.3 s fixation, followed by a 0.5 s blank screen. Afterward, a verb was visually presented on the

center of the screen for 2 s, accompanied by its verbalization from the speaker. Participants

judged whether the spoken verbs were clear and recognizable by pressing a button. All 100 verbs

(96 verbs in the word-picture pairs and 4 verbs as control cues) were randomly presented during

this stage, with each verb being presented twice. In the picture familiarization session, each trial

started with 0.3 s fixation, followed by a 0.5 s blank screen. Afterward, a picture and its label

(e.g., for a panda picture, the label would be "panda") were presented on the screen for 2 s.

Participants indicated whether they were familiar with the picture and its name by pressing a

button. Each of the 192 pictures (96 targets + 96 lures) was presented twice. For items that

participants indicate unfamiliar will be presented for another two rounds. At the end of the task,

participants indicated that all stimuli were clearly recognizable and familiar.

Word-picture associative learning task

Participants learned 96 word-picture pairs. Each learning trial consisted of three phases, an

encoding phase, a maintenance phase, and a vividness rating phase. During the encoding phase,

following a 0.3 s fixation and a black screen jittering between 0.9-1.5 s, participants viewed a

picture presented in the center of the screen for 2 s while hearing the spoken verb from the

speaker. Participants were instructed to pay attention to the details of the picture while

memorizing the verb-picture associations during the encoding phase. During the maintenance

phase, the picture disappeared, and participants were asked to maintain the picture in their minds

as vividly as possible for 3s while hearing the spoken verb again. In the vividness rating phase,

participants indicated the vividness of the mental image of the picture on a 1 (not vivid at all) -4

(very vivid) point Likert scale by pressing one of four buttons. The learning task consisted of

three blocks, with each block containing 32 unique verb-picture pairs that were repeated three

times. To reduce the recency effect, participants completed a ~5 min math task after the learning

task.

 Pre-sleep memory test

To understand whether the pre-sleep testing alters TMR effects on memory consolidation,

participants were tested on half of the pairs during the pre-sleep test (i.e., 48 pairs, with 12 pairs

from one of the four picture categories). This test consisted of a cued recall task and a cued

recognition task.

In the cued recall task, each trial started with a 0.3 s fixation, followed by a blank screen (0.9-1.5

s). The spoken verb was played, prompting participants to report whether they could "remember"

or "forget" the corresponding picture. This stage was self-paced so that participants had enough

time to recall. Immediately following this "remember" or "forget" response, participants were

further asked to report the category of the picture by pressing one of four buttons, with each

button indicating one of four categories within 2 s.

To further encourage participants to remember the detailed picture content, instead of the

category information, we further ask participant to perform a cued-recognition task. In this

recognition task, the same half of the pairs were tested. Specifically, each trial started with a

fixation (0.3 s) and was followed by a blank screen (0.9-1.5 s). Participants next saw the picture

presented on the center of the screen while hearing the spoken verb. Participants were asked to

indicate if the picture was the same picture paired with the verb during the previous learning task

by pressing the "Yes" or "No" button. Among 192 recognition trials, 48 trials showed identical

verb-picture associations from the learning task, and 48 trials showed verb-lure picture

associations. The remaining 96 trials consisted of 48 mixed verb-picture associations and 48

mixed verb-lure associations with cue verbs and pictures (or lures) corresponding to different

learning trials.

TMR during NREM sleep

To counterbalance the pre-sleep memory performance between cued and uncued items, we

selected half of the remembered and half of the forgotten pairs (based on the category report

performance in the cued recall task) of the tested items as cued items during TMR. In addition,

we randomly selected half of the untested items to be cued in the TMR. Picture categories were

balanced across cued versus non-cued conditions. Thus, the TMR session contained 48 verbs as

cues, with additional 4 verbs that were not paired with any pictures as control cues. Half of the

cues were from tested trials, with the remaining half from untested trials. The control cues served

as the baseline to examine retrieval-specific neural activity. Therefore, there were 52 unique

sound cues played during sleep.

During the nocturnal sleep, white noise was played in the background throughout the night, with

an intensity of ~ 45 dB, measured by a sound-level meter placed at the same position where

participants laid their heads on the pillow. Experienced experimenters monitored the EEG signals

and visually identified signature EEG events characterizing different sleep stages (e.g., spindles,

K-complex, slow oscillations). Upon detecting stable slow-wave sleep, the experimenter would

begin the TMR, which occurred ~50.04 min (SD: 37.21min) after the start of the sleep phase. On

average, participants were presented with 8.92 (SD: 3.24) rounds of TMR. In each round of

TMR, 52 verb cues were randomly presented with an inter-stimuli interval of 5 ± 0.2 s. After

each round of cueing, the order of the TMR cues was shuffled and replayed again. Each round

was separated by 30 s. The TMR session continued as long as the participants were in slow-wave

sleep in the first 3-4 hours of the nocturnal sleep. Cueing was stopped immediately when

participants showed signs of micro-arousals, awakening or changed to N1 sleep or rapid eye

movement (REM) sleep. Cueing was resumed after the participants returned to stable slow-wave

sleep.

Post-sleep memory test

Approximately 30 minutes after awakening and after the PVT test, participants were tested on all

96 pairs. Similar to pre-sleep memory tests, this test included the cued recall and cued

recognition tasks. In addition, there was a closed-eye mental retrieval task between these two

tasks, in which participants were asked to keep their eyes closed and relax while each of the

verbs was randomly played via the speaker (ISI = 5 ± 0.2 s). The current study focuses on TMR-

based memory consolidation process. Therefore, the post-sleep closed-eye mental retrieval data

will report elsewhere.

EEG recording and preprocessing

We collected the EEG data throughout the experiment except for during the familiarization and

PVT tasks. EEG data were recorded using the amplifier from the eego system (ANT neuro,

Netherlands, https://www.ant-neuro.com) with a sampling rate of 500 Hz from 61 channels

(waveguard EEG caps) that were mounted in the International 10-20 system. Additionally, there

were two electrodes placed on the left and right mastoids, respectively, and one electrode was

placed above the left eye for the EOG measurements. Online EEG recordings were referenced to

the default reference channel (i.e., CPz). For sleep monitoring, another two electrodes were

placed on both sides of the chin to measure the EMG using the bipolar reference. EEG data

collection was started after the impedance of all electrodes was lower than 20 KΩ.

Sleep EEG data preprocessing was performed using EEGLAB (https://sccn.ucsd.edu/eeglab/)

and Fieldtrip toolboxes(*73*), as well as in-house code that was implemented in MATLAB

(MathWorks Inc.). EEG data were first notch filtered at 50 ± 2 Hz, and then bandpass filtered

between 0.5 and 40 Hz. Then continuous sleep EEG data were segmented into 15 s epochs, i.e.,

form -5 s to 10 s relative to the onset of TMR cues. This long epoch was used to eliminate the

edge effect in the later time-frequency analysis. Our main interesting time windows for the sleep

data are from 0 to 5 s relative to the TMR cue onset. Bad epochs are marked based on visual

inspection and rejected from further analysis. Bad channels were marked and interpolated using

spherical interpolation in EEGLAB. Afterward, EEG data were re-referenced to the average of

the artifact-free data.

Time-frequency analysis

We performed time-frequency transformation on the preprocessed EEG data, using the complex

Morlet wavelets (six cycles). Spectral power was extracted from the frequency range of 1-40 Hz,

with a step of 1 Hz and with the time of interest range of [-1 5 s] relative to the TMR cue onset.

The power was down-sampled to 100 Hz after the time-frequency transformation. We

normalized the power data within each frequency bin and each channel by first subtracting the

mean power in the baseline time windows ([-1 -0.5s] relative to the TMR cue onset) and then

dividing the same baseline mean power. Finally, we re-segmented the power data into the 5s

epochs (i.e., [0 5s] relative to the TMR cue onset).

Spindle detection

Individual spindles were detected during the TMR periods following previous studies(*27*).

Specifically, artifact-free EEG data during the TMR periods were first bandpass filtered between

11-16 Hz by using the 4th order two-pass Butterworth filter. Next, the root-mean-square (RMS)

values were calculated for each time point with a moving window of 400 ms. Third, the spindle

amplitude criterion is defined as the mean + 1.5 SD of the RMS signal. Sleep spindles were

detected if the RMS signal consecutively exceeded the amplitude criterion for a duration of 0.5

to 3 s. Previous studies indicated that spindle activity was prominent over anterior-posterior

midlines(*74*). To this end, spindle detection was performed for all seven midline EEG channels

(i.e., "FPz", "Fz", "FCz", "Cz", "Pz", "POz", and "Oz"), separately. We then assign the spindle

value of 1 to the time points where a spindle was detected and 0 otherwise. The spindle

probability was computed as the mean spindle values across trials for each time point in the time

range of [0 5 s] relative to the TMR cue onset and then averaged across all midline channels.

Slow-oscillation detection

The detection of slow oscillations (SOs) was performed on the artifact-free EEG data during the

TMR periods. We first bandpass filtered the EEG data between 0.3 and 1.25 Hz using the two-

pass FIR filter, with the order equalling the three cycles of the low-frequency cut-off. We then

detected the zero-crossings in the filtered signal, and the event durations were calculated as the

temporal distance between two successive positive-to-negative zero-crossings. SO was detected

if the peak-to-peak amplitude was greater than the 75% percentile of the absolute amplitude of

the filtered signal(*75*) and the event duration was between 0.8 s and 2 s. SOs detection was

performed on the Fz, where the amplitude of SOs was prominent according to previous

studies(*76*).

SO-spindle coupling

For TMR trials that showed both the spindle and SOs in the time windows (i.e., 2500-2960 ms),

which showed significant interactions of item-specificity and post-sleep memory effect, we then

performed the SO-spindle coupling. We first filtered the SO-spindle trials in the SO frequency

range (i.e., 0.3-1.25 Hz). We then applied the Hilbert transform to the filtered data to extract the

instantaneous phase of the SOs. To obtain the amplitude of the spindle activities, we filtered the

SO-spindle trials in the spindle frequency range (i.e., 11-16 Hz) and then applied the Hilbert

transform to the filtered data to obtain the instantaneous amplitude of the spindle activities on an

extended time window (i.e., [2.2-4.2 s] relatively to TMR cue onset to ensure at least half of the

SOs cycle was included). We detected the preferred SO phase, which is concurrent with the

maximal spindle amplitude across all trials in each participant, and then tested the distribution of

the preferred SO phases across participants against the uniform distribution using the Rayleigh

test (CircStat toolbox(*77*)).

Representational similarity analysis

To extract item-specific representations, we performed representational similarity analysis (RSA)

between every two clean TMR trials that originated from different TMR rounds. Specifically, the

within-item similarity was calculated as the whole-brain EEG cortical pattern similarity between

the same TMR cues; and the between-item similarity was calculated as the pattern similarity

between different TMR cues. Thus, contrasting within-item similarity versus between-items

similarity enables us to examine whether there are item-specific representations following the

TMR cues.

To characterize the item-specific representations across time, RSA was performed on the

preprocessed raw EEG using sliding time windows of 500 ms, with an incremental step of 10 ms,

resulting in 250*61 (time points * channels) features in each time window. Then, for each time

window, we calculate the similarity between vectorized features of every two trials that were

from different TMR rounds using Spearman's corrections. All the correlation values were Fisher

Z-transformed before further statistical analysis.

Statistics

For the behavioral data, we conducted a two-way ANOVA with TMR effects (cued vs. uncued)

and pre-sleep testing (tested vs. untested) as repeated measures in analyzing the post-sleep

memory performance. We also performed a two-way ANOVA with TMR and time (pre-sleep vs.

post-sleep) as repeated measures in analyzing the memory performance for tested items. The

paired sample *t*-tests were employed to examine the differences between two specific

experimental conditions (e.g., memory performance between pre-sleep tested items and post-

sleep tested items). The relationship between EEG activities during the TMR period and post-

sleep memory performance is assessed by the robust linear regression model, which is less likely

to be affected by potential outliers(*78*).

For the EEG data, multiple comparisons across consecutive time windows were corrected using

the cluster-based nonparametric statistical tests in MATLAB(*79*). Specifically, statistical tests

(e.g., *t*-test) were performed between conditions (e.g., WI vs. BI or tested vs. untested) in

individual time (or time-frequency) windows. Adjacent time (or time-frequency) windows with
statistical values exceeding a threshold ($p < 0.05$) were combined into contiguous clusters.
Cluster-level statistics were computed using the sum of the t values within a cluster. To test the
significance of the time (or time-frequency) cluster, a distribution of cluster-level statistics under
the null hypothesis was constructed by randomly permuting condition labels 1000 times, and the
maximum cluster-level statistic in each permutation was extracted. If no significant cluster was
found for a permutation, a value of 0 was assigned for that permutation. The nonparametric
statistical significance of a cluster was then obtained by calculating the proportion of cluster-
level statistics in the distribution under the null hypothesis that exceeded the empirical cluster-
level statistics. For an identified cluster, we also performed a two-way ANOVA with item-
specificity (WI vs. BI similarity) and pre-sleep testing as repeated measures and a two-way
ANOVA with item-specificity and subsequent memory as repeated measures. The statistical
significance level for all the analyses is set as p-value $< .05$ or corrected p-value (p_{corr}) < 0.05.
712
713

References

1. J. L. McGaugh, Memory--a Century of Consolidation. *Science*. **287**, 248–251 (2000).

2. J. Born, I. Wilhelm, System consolidation of memory during sleep. *Psychological Research*. **76**, 192–203 (2012).

3. J. G. Klinzing, N. Niethard, J. Born, Mechanisms of systems memory consolidation during sleep. *Nat Neurosci*. **22**, 1598–1610 (2019).

4. M. Schönauer, S. Alizadeh, H. Jamalabadi, A. Abraham, A. Pawlizki, S. Gais, Decoding material-specific memory reprocessing during sleep in humans. *Nat Commun*. **8**, 15404 (2017).

5. T. Schreiner, M. Petzka, T. Staudigl, B. P. Staresina, Endogenous memory reactivation during sleep in humans is clocked by slow oscillation-spindle complexes. *Nat Commun*. **12**, 3112 (2021).

6. V. Sterpenich, M. K. M. van Schie, M. Catsiyannis, A. Ramyead, S. Perrig, H.-D. Yang, D. Van De Ville, S. Schwartz, Reward biases spontaneous neural reactivation during sleep. *Nat Commun*. **12**, 4162 (2021).

7. M. A. Wilson, B. L. McNaughton, Reactivation of Hippocampal Ensemble Memories During Sleep. *Science*. **265**, 676–679 (1994).

8. H. Zhang, J. Fell, N. Axmacher, Electrophysiological mechanisms of human memory consolidation. *Nat Commun*. **9**, 4103 (2018).

9. D. Bendor, M. A. Wilson, Biasing the content of hippocampal replay during sleep. *Nat Neurosci*. **15**, 1439–1444 (2012).

10. X. Hu, L. Y. Cheng, M. H. Chiu, K. A. Paller, Promoting memory consolidation during sleep: A meta-analysis of targeted memory reactivation. *Psychological Bulletin*. **146**, 218–244 (2020).

11. B. Rasch, C. Büchel, S. Gais, J. Born, Odor Cues During Slow-Wave Sleep Prompt Declarative Memory Consolidation. *Science*. **315**, 1426–1429 (2007).

12. J. D. Rudoy, J. L. Voss, C. E. Westerberg, K. A. Paller, Strengthening Individual Memories by Reactivating Them During Sleep. *Science*. **326**, 1079–1079 (2009).

13. K. A. Paller, J. D. Creery, E. Schechtman, Memory and Sleep: How Sleep Cognition Can Change the Waking Mind for the Better. *Annual Review of Psychology*. **72**, 123–150 (2021).

14. D. I. Schouten, S. I. R. Pereira, M. Tops, F. M. Louzada, State of the art on targeted memory reactivation: Sleep your way to enhanced cognition. *Sleep Medicine Reviews*. **32**, 123–131 (2017).

15. S. Schwartz, A. Clerget, L. Perogamvros, Enhancing imagery rehearsal therapy for
nightmares with targeted memory reactivation. *Current Biology*. **32**, 4808-4816.e4 (2022).

16. V. Vidal, A. R. Barbuzza, L. M. Tassone, L. I. Brusco, F. M. Ballarini, C. Forcato, Odor
cueing during sleep improves consolidation of a history lesson in a school setting. *Sci Rep*.
12, 10350 (2022).

17. Z. Chen, M. A. Wilson, Deciphering Neural Codes of Memory during Sleep. *Trends
Neurosci*. **40**, 260–275 (2017).

18. S. Belal, J. Cousins, W. El-Deredy, L. Parkes, J. Schneider, H. Tsujimura, A. Zoumpoulaki,
M. Perapoch, L. Santamaria, P. Lewis, Identification of memory reactivation during sleep
by EEG classification. *NeuroImage*. **176**, 203–214 (2018).

19. S. A. Cairney, A. á V. Guttesen, N. El Marj, B. P. Staresina, Memory Consolidation Is
Linked to Spindle-Mediated Information Processing during Sleep. *Current Biology*. **28**,
948-954.e4 (2018).

20. L. K. Shanahan, E. Gjorgieva, K. A. Paller, T. Kahnt, J. A. Gottfried, Odor-evoked category
reactivation in human ventromedial prefrontal cortex during sleep promotes memory
consolidation. *eLife*. **7**, e39681 (2018).

21. B. Wang, J. W. Antony, S. Lurie, P. P. Brooks, K. A. Paller, K. A. Norman, Targeted
Memory Reactivation during Sleep Elicits Neural Signals Related to Learning Content. *J.
Neurosci.* **39**, 6728–6736 (2019).

22. T. Schreiner, C. F. Doeller, O. Jensen, B. Rasch, T. Staudigl, Theta Phase-Coordinated
Memory Reactivation Reoccurs in a Slow-Oscillatory Rhythm during NREM Sleep. *Cell
Reports*. **25**, 296–301 (2018).

23. T. Andrillon, S. Kouider, The vigilant sleeper: neural mechanisms of sensory (de)coupling
during sleep. *Current Opinion in Physiology*. **15**, 47–59 (2020).

24. H. Hayat, A. Marmelshtein, A. J. Krom, Y. Sela, A. Tankus, I. Strauss, F. Fahoum, I. Fried,
Y. Nir, Reduced neural feedback signaling despite robust neuron and gamma auditory
responses during human sleep. *Nat Neurosci*. **25**, 935–943 (2022).

25. J. W. Antony, M. Schönauer, B. P. Staresina, S. A. Cairney, Sleep Spindles and Memory
Reprocessing. *Trends in Neurosciences*. **42**, 1–3 (2019).

26. M. Schönauer, Sleep Spindles: Timed for Memory Consolidation. *Current Biology*. **28**,
R656–R658 (2018).

27. J. W. Antony, L. Piloto, M. Wang, P. Pacheco, K. A. Norman, K. A. Paller, Sleep Spindle
Refractoriness Segregates Periods of Memory Reactivation. *Current Biology*. **28**, 1736-
1743.e4 (2018).

28. L. Fuentemilla, J. Miró, P. Ripollés, A. Vilà-Balló, M. Juncadella, S. Castañer, N. Salord, C.
Monasterio, M. Falip, A. Rodríguez-Fornells, Hippocampus-Dependent Strengthening of
Targeted Memories via Reactivation during Sleep in Humans. *Current Biology*. **23**, 1769–
1775 (2013).

29. T. Schreiner, M. Lehmann, B. Rasch, Auditory feedback blocks memory benefits of cueing
during sleep. *Nat Commun*. **6**, 8729 (2015).

30. J. W. Antony, C. S. Ferreira, K. A. Norman, M. Wimber, Retrieval as a Fast Route to
Memory Consolidation. *Trends in Cognitive Sciences*. **21**, 573–576 (2017).

31. H. L. Roediger, A. C. Butler, The critical role of retrieval practice in long-term retention.
Trends in Cognitive Sciences. **15**, 20–27 (2011).

32. B. H. Joensen, M. O. Harrington, S. C. Berens, S. A. Cairney, M. G. Gaskell, A. J. Horner,
Targeted memory reactivation during sleep can induce forgetting of overlapping memories.
Learn. Mem. **29**, 401–411 (2022).

33. S. A. Cairney, S. Lindsay, J. M. Sobczak, K. A. Paller, M. G. Gaskell, The Benefits of
Targeted Memory Reactivation for Consolidation in Sleep are Contingent on Memory
Accuracy and Direct Cue-Memory Associations. *Sleep*. **39**, 1139–1150 (2016).

34. J. D. Creery, D. Oudiette, J. W. Antony, K. A. Paller, Targeted Memory Reactivation during
Sleep Depends on Prior Learning. *Sleep*. **38**, 755–763 (2015).

35. A. C. Schapiro, E. A. McDevitt, T. T. Rogers, S. C. Mednick, K. A. Norman, Human
hippocampal replay during rest prioritizes weakly learned information and predicts memory
performance. *Nat Commun*. **9**, 3920 (2018).

36. D. Denis, D. Mylonas, C. Poskanzer, V. Bursal, J. D. Payne, R. Stickgold, Sleep Spindles
Preferentially Consolidate Weakly Encoded Memories. *J. Neurosci.* **41**, 4088–4099 (2021).

37. C. Barner, M. Altgassen, J. Born, S. Diekelmann, Effects of sleep on the realization of
complex plans. *Journal of Sleep Research*. **28**, e12655 (2019).

38. I. Wilhelm, S. Diekelmann, I. Molzow, A. Ayoub, M. Mölle, J. Born, Sleep Selectively
Enhances Memory Expected to Be of Future Relevance. *J. Neurosci.* **31**, 1563–1569 (2011).

39. N. Kriegeskorte, M. Mur, P. Bandettini, Representational similarity analysis - connecting
the branches of systems neuroscience. *Frontiers in Systems Neuroscience*. **2** (2008)
(available at https://www.frontiersin.org/articles/10.3389/neuro.06.004.2008).

40. J. Liu, H. Zhang, T. Yu, L. Ren, D. Ni, Q. Yang, B. Lu, L. Zhang, N. Axmacher, G. Xue,
Transformative neural representations support long-term episodic memory. *Science*
advances. **7**, eabg9715 (2021).

41.   M. Wislowska, W. Klimesch, O. Jensen, C. Blume, M. Schabus, Sleep-Specific Processing
of Auditory Stimuli Is Reflected by Alpha and Sigma Oscillations. *J. Neurosci.* **42**, 4711–
4724 (2022).

42.   T. Andrillon, A. T. Poulsen, L. K. Hansen, D. Léger, S. Kouider, Neural Markers of
Responsiveness to the Environment in Human Sleep. *J. Neurosci.* **36**, 6583–6596 (2016).

43.   P. Adank, B. G. Evans, J. Stuart-Smith, S. K. Scott, Comprehension of familiar and
unfamiliar native accents under adverse listening conditions. *Journal of Experimental*
Psychology: Human Perception and Performance. **35**, 520–529 (2009).

44.   E. B. Issa, X. Wang, Sensory Responses during Sleep in Primate Primary and Secondary
Auditory Cortex. *J. Neurosci.* **28**, 14467–14480 (2008).

45.   Y. Nir, V. V. Vyazovskiy, C. Cirelli, M. I. Banks, G. Tononi, Auditory Responses and
Stimulus-Specific Adaptation in Rat Auditory Cortex are Preserved Across NREM and
REM Sleep. *Cerebral Cortex*. **25**, 1362–1378 (2015).

46.   S. Kouider, T. Andrillon, L. S. Barbosa, L. Goupil, T. A. Bekinschtein, Inducing Task-
Relevant Responses to Speech in the Sleeping Brain. *Current Biology*. **24**, 2208–2214
(2014).

47.   G. Legendre, T. Andrillon, M. Koroma, S. Kouider, Sleepers track informative speech in a
multitalker environment. *Nat Hum Behav*. **3**, 274–283 (2019).

48.   F. Perrin, L. García-Larrea, F. Mauguière, H. Bastuji, A differential brain response to the
subject's own name persists during sleep. *Clinical Neurophysiology*. **110**, 2153–2164
(1999).

49.   M. A. Züst, S. Ruch, R. Wiest, K. Henke, Implicit Vocabulary Learning during Sleep Is
Bound to Slow-Wave Peaks. *Current Biology*. **29**, 541-553.e7 (2019).

50.   T. Schreiner, B. Rasch, Boosting Vocabulary Learning by Verbal Cueing During Sleep.
Cerebral Cortex. **25**, 4169–4179 (2015).

51.   C. Blume, R. del Giudice, M. Wislowska, D. P. J. Heib, M. Schabus, Standing sentinel
during human sleep: Continued evaluation of environmental stimuli in the absence of
consciousness. *NeuroImage*. **178**, 638–648 (2018).

52.   I. Holeckova, C. Fischer, M.-H. Giard, C. Delpuech, D. Morlet, Brain responses to a
subject's own name uttered by a familiar voice. *Brain Research*. **1082**, 142–152 (2006).

53.   I. Oswald, A. M. Taylor, M. Treisman, Discriminative responses to stimulation during
human sleep. *Brain*. **83**, 440–453 (1960).

54.   S. Ai, Y. Yin, Y. Chen, C. Wang, Y. Sun, X. Tang, L. Lu, L. Zhu, J. Shi, Promoting
subjective preferences in simple economic choices during nap. *eLife*. **7**, e40583 (2018).

55.   T. Xia, J. W. Antony, K. A. Paller, X. Hu, Targeted memory reactivation during sleep
influences social bias as a function of slow-oscillation phase and delta power.
Psychophysiology. **60**, e14224 (2023).

56.   R. M. W. J. Berkers, M. Ekman, E. V. van Dongen, A. Takashima, M. Barth, K. A. Paller, G.
Fernández, Cued reactivation during slow-wave sleep induces brain connectivity changes
related to memory stabilization. *Sci Rep*. **8**, 16958 (2018).

57.   G. Rothschild, E. Eban, L. M. Frank, A cortical–hippocampal–cortical loop of information
processing during memory consolidation. *Nat Neurosci*. **20**, 251–259 (2017).

58.   P. A. Lewis, D. Bendor, How Targeted Memory Reactivation Promotes the Selective
Strengthening of Memories in Sleep. *Current Biology*. **29**, R906–R912 (2019).

59.   S. Diekelmann, C. Büchel, J. Born, B. Rasch, Labile or stable: opposing consequences for
memory when reactivated during waking and sleep. *Nat Neurosci*. **14**, 381–386 (2011).

60.   J. L. McClelland, B. L. McNaughton, R. C. O'Reilly, Why there are complementary
learning systems in the hippocampus and neocortex: Insights from the successes and
failures of connectionist models of learning and memory. *Psychological Review*. **102**, 419–
457 (1995).

61.   L. R. Squire, Memory and the hippocampus: A synthesis from findings with rats, monkeys,
and humans. *Psychological Review*. **99**, 195–231 (1992).

62.   E. Rolls, The mechanisms for pattern completion and pattern separation in the
hippocampus. *Frontiers in Systems Neuroscience*. **7** (2013) (available at
https://www.frontiersin.org/articles/10.3389/fnsys.2013.00074).

63.   B. P. Staresina, M. Wimber, A Neural Chronometry of Memory Recall. *Trends in Cognitive*
Sciences. **23**, 1071–1085 (2019).

64.   R. F. Helfrich, J. D. Lendner, B. A. Mander, H. Guillen, M. Paff, L. Mnatsakanyan, S.
Vadera, M. P. Walker, J. J. Lin, R. T. Knight, Bidirectional prefrontal-hippocampal
dynamics organize information transfer during sleep in humans. *Nat Commun*. **10**, 3572
(2019).

65.   N. Maingret, G. Girardeau, R. Todorova, M. Goutierre, M. Zugaro, Hippocampo-cortical
coupling mediates memory consolidation during sleep. *Nat Neurosci*. **19**, 959–964 (2016).

66.   A. Mitra, A. Z. Snyder, C. D. Hacker, M. Pahwa, E. Tagliazucchi, H. Laufs, E. C.
Leuthardt, M. E. Raichle, Human cortical–hippocampal dialogue in wake and slow-wave
sleep. *Proceedings of the National Academy of Sciences*. **113**, E6868–E6876 (2016).

67.   H.-V. Ngo, J. Fell, B. Staresina, Sleep spindles mediate hippocampal-neocortical coupling
during long-duration ripples. *eLife*. **9**, e57011 (2020).

68. R. Cox, J. van Driel, M. de Boer, L. M. Talamini, Slow Oscillations during Sleep
Coordinate Interregional Communication in Cortical Networks. *J. Neurosci.* **34**, 16890–
16901 (2014).

69. M. Göldi, E. A. M. van Poppel, B. Rasch, T. Schreiner, Increased neuronal signatures of
targeted memory reactivation during slow-wave up states. *Scientific reports*. **9**, 1–10
(2019).

70. M. Tatsuno, S. Malek, L. Kalvi, A. Ponce-Alvarez, K. Ali, D. R. Euston, S. Grün, B. L.
McNaughton, Memory reactivation in rat medial prefrontal cortex occurs in a subtype of
cortical UP state during slow-wave sleep. *Philosophical Transactions of the Royal Society*
B: Biological Sciences. **375**, 20190227 (2020).

71. L. Himmer, E. Müller, S. Gais, M. Schönauer, Sleep-mediated memory consolidation
depends on the level of integration at encoding. *Neurobiology of Learning and Memory*.
137, 101–106 (2017).

72. E. A. Phelps, S. G. Hofmann, Memory editing from science fiction to clinical practice.
Nature. **572**, 43–50 (2019).

73. R. Oostenveld, P. Fries, E. Maris, J.-M. Schoffelen, FieldTrip: Open Source Software for
Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Computational*
Intelligence and Neuroscience. **2011**, 1–9 (2011).

74. N. Dehghani, S. S. Cash, E. Halgren, Topographical frequency dynamics within EEG and
MEG sleep spindles. *Clinical Neurophysiology*. **122**, 229–235 (2011).

75. R. F. Helfrich, B. A. Mander, W. J. Jagust, R. T. Knight, M. P. Walker, Old Brains Come
Uncoupled in Sleep: Slow Wave-Spindle Synchrony, Brain Atrophy, and Forgetting.
Neuron. **97**, 221-230.e4 (2018).

76. L. Marshall, H. Helgadóttir, M. Mölle, J. Born, Boosting slow oscillations during sleep
potentiates memory. *Nature*. **444**, 610–613 (2006).

77. P. Berens, CircStat: A MATLAB Toolbox for Circular Statistics. *Journal of Statistical*
Software. **31**, 1–21 (2009).

78. C. Yu, W. Yao, Robust linear regression: A review and comparison. *Communications in*
Statistics - Simulation and Computation. **46**, 6261–6282 (2017).

79. E. Maris, R. Oostenveld, Nonparametric statistical testing of EEG- and MEG-data. *Journal*
of Neuroscience Methods. **164**, 177–190 (2007).

Acknowledgments

General: We thank Lingqi Zhang and Zexuan Mu for their help with the data collection.

Funding: The research was supported by the Ministry of Science and Technology of China STI2030-Major Projects (No. 2022ZD0214100), National Natural Science Foundation of China (No. 32171056), General Research Fund (No. 17614922) of Hong Kong Research Grants Council, and the Key Realm R&D Program of Guangzhou (No. 20200703005) to X. H., General Research Fund (No. 17600621) of Hong Kong Research Grants Council to T. L., and Start-up Fund for RAPs under the Strategic Hiring Scheme at The Hong Kong Polytechnic University (Project ID: P0043338) to J. L..

Author contributions: J. L., and X. H. conceived the study and designed the experiment. J. L., and M. Z. conducted the experiment. J. L., T. X., D. C., Z. Y., J. W. A., T. L., and X. H. contributed to data analyses. J.L., X. H. and T.L. drafted the manuscript, with critical feedback from J.W.A. All authors contributed to the revision of the manuscript.

Competing interests: The authors declare that they have no competing interests.

Data and materials availability: All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. All data that support the findings of this study are available in the Open Science Framework (https://osf.io/knb42/?view_only=04d1d0d6deb24c69b34cacce73ed6f03).