AI-assisted selection of mating pairs through simulation-based optimized progeny allocation strategies in plant breeding

Kosuke Hamazaki[1,2] and Hiroyoshi Iwata[1*]

[1]Laboratory of Biometry and Bioinformatics, Department of Agricultural and Environmental

Biology, Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo, Japan.

(current address) [2]Molecular Informatics Team, RIKEN Center for Advanced Intelligence Project

(AIP), RIKEN, Chiba, Japan.

* Correspondence:

Corresponding Author

hiroiwata@g.ecc.u-tokyo.ac.jp

Keywords: Breeding Scheme, Optimization, Progeny Allocation, Forward Simulation, Mating

Pairs, Genetic Diversity, Breeding Values, AI Breeder.

Abstract

Emerging technologies such as genomic selection have been applied to modern plant and animal

breeding to increase the speed and efficiency of variety release. However, breeding requires decisions

regarding parent selection and mating pairs, which significantly impact the ultimate genetic gain of a

breeding scheme. The selection of appropriate parents and mating pairs to increase genetic gain while

maintaining genetic diversity is still an urgent need that breeders are facing. This study aimed to

determine the best progeny allocation strategies by combining future-oriented simulations and

numerical black-box optimization for an improved selection of parents and mating pairs. In this

study, we focused on optimizing the allocation of progenies, and the breeding process was regarded

as a black-box function whose input is a set of parameters related to the progeny allocation strategies

and whose output is the ultimate genetic gain of breeding schemes. The allocation of progenies to

each mating pair was parameterized according to a softmax function of multiple selection criteria.

Different selection criteria are weighted in the function to balance genetic gains and genetic diversity

optimally. The weighting parameters were then optimized by the black-box optimization algorithm

called StoSOO via future-oriented breeding simulations. Simulation studies to evaluate the potential

of our novel method revealed that the breeding strategy based on optimized weights attained almost

10% higher genetic gain than that with an equal allocation of progenies to all mating pairs within just

four generations. Among the optimized strategies, those considering the expected genetic variance of

progenies could maintain the genetic diversity throughout the breeding process, leading to a higher

ultimate genetic gain than those without considering it. These results suggest that our novel method

can significantly improve the speed and efficiency of variety development through optimized

decisions regarding the selection of parents and mating pairs. In addition, by changing simulation

settings, our future-oriented optimization framework for progeny allocation strategies can be easily

implemented into general breeding schemes, contributing to accelerated plant and animal breeding

with high efficiency.

1   Introduction

To meet the increasing demand for agricultural products caused by the population explosion,

developing new cultivars with desired agronomic characteristics, such as high yield, good quality,

and efficient nutrient use, is urgently needed (Capper, 2011). However, in plant breeding, for

instance, conventional breeding methods require several years or longer to produce new cultivars,

even in annual crops, which makes it challenging to meet the increasing food demand in the near

future (Eggen, 2012). Genomic selection (GS) is expected to be a critical methodology for

accelerating the evaluation and selection of superior genotypes that can be implemented in

conventional breeding (Meuwissen et al., 2001; Jannink et al., 2010). In GS, genotypic values of

target traits are predicted using genomic prediction (GP) models based on genome-wide marker data,

and the predicted values are used for selection in breeding schemes (Meuwissen et al., 2001). GS

enables individual-based and crossing-based selection according to the genotypic values predicted by

the models with fewer field evaluations of target traits, which leads to highly efficient and rapid

breeding (Zhong and Jannink, 2007; Jannink et al., 2010; Crossa et al., 2017; Lehermeier et al.,

2017). The superiority of GS over selections based on either phenotypic date or pedigree-derived

prediction (pedigree BLUP) has been reported in many simulations and empirical studies (Goddard,

2009; Hayes et al., 2009; Jannink et al., 2010; Goddard et al., 2011; Crossa et al., 2013; Hickey et al.,

2014; García-Ruiz et al., 2016; Rabier et al., 2016). However, even though GS has contributed to

speeding up breeding during selections, the GS itself does not aim to optimize decisions in breeding.

Thus, the ultimate decisions regarding parent selection and choosing mating pairs as the beginning

step of the breeding scheme still largely rely on breeders' experiences because many data-based

approaches have not been fully utilized by breeders due to target traits' complexity, limited

resources, or implementation difficulty. In other words, the advent of GS has not yet entirely led to

optimized decisions in breeding programs.

To solve the above problems and optimize the decisions regarding (1) selection, (2) mating, and (3)

progeny allocation, there has been much discussion on improving the strategies in breeding schemes.

First, as for the selection strategy, different selection criteria have been used to increase the efficiency

of GS. While various GP models have been developed to improve the accuracy of GS (Meuwissen et

al., 2001; Gianola and van Kaam, 2008; VanRaden, 2008; de los Campos et al., 2009; Jannink et al.,

2010; Habier et al., 2011; de Los Campos et al., 2013; González-Camacho et al., 2016), genomic

estimated breeding value (GEBV), the total sum of estimated additive marker effects, has been

generally utilized for GS as a selection criterion (Meuwissen et al., 2001). Although GEBV is usually

used for a short breeding period with a few cycles of selection and crossing, it does not guarantee an

increased genetic gain in a long-term breeding process (Sonesson et al., 2012; Lin et al., 2017;

Akdemir et al., 2019). Because genetic variance is greatly reduced due to the Bulmer effect (Bulmer,

1971; Van Grevenhof et al., 2012) and genetic diversity is also reduced by inbreeding and random

drift under truncation selection (Falconer and Mackay, 1996; Li et al., 2008), the genetic gain rapidly

reaches a local, rather than global, optimum, i.e., a plateau, in GEBV-based GS cycles (Moeinizade

et al., 2019; Zhang and Wang, 2022). Other selection criteria have been proposed to solve the issue of

GEBV-based GS by maintaining the genetic diversity and further increasing the genetic gain

expected in the long-term breeding program. Weighted genomic estimated breeding value (WGEBV)

is a criterion that utilizes marker allele frequencies to emphasize the marker effects of rare alleles

(Goddard, 2009; Jannink, 2010). Compared to GEBV, WGEBV can maintain genetic diversity in

breeding populations over the long term by suppressing the Bulmer effect (Goddard, 2009; Jannink,

2010). Other selection criteria, such as optimal haploid value (Daetwyler et al., 2015), with an

expected maximum haploid breeding value (Müller et al., 2018), and optimal population value

(Goiffon et al., 2017), have been introduced to maintain genetic diversity by including haplotype

information and have indeed improved the final genetic gains at the end of the breeding scheme in

long-term breeding programs compared to those from GEBV-based GS. Although these selection

criteria are helpful in selecting parental candidates, they do not directly enable the automatic

optimization of decisions regarding the selection of parents. Therefore, breeders must make ultimate

decisions based on these criteria by themselves. Also, an optimal contributions selection (OCS)

method, which aims to optimize decisions regarding parent selection, was proposed to help breeders

decide on more appropriate breeding strategies rather than just use selection criteria (Meuwissen,

1997; Grundy et al., 1998). OCS aims to optimize the expected contribution vector, i.e., the number

of times each individual is used as a parent for the next generation, by maximizing the genetic gains

while constraining the inbreeding in the next generation. Although the efficiency of the OCS-based

GS was evaluated in several simulation studies (Gorjanc et al., 2018; Cowling et al., 2023), OCS still

cannot assist in identifying optimal mating pairs even for the next generation.

Second, as for the problems regarding the selection of appropriate mating pairs, i.e., the mating

design problem, many studies in animal breeding have long been conducted to select optimal pairs

while considering mating constraints (Jansen and Wilton, 1985; Kinghorn, 2011), but they mostly

lacked a long-term perspective. Then, a look-ahead selection (LAS) approach has been proposed as a

method for simultaneously optimizing decisions regarding the selection of appropriate parents and

mating pairs in plant breeding (Moeinizade et al., 2019). LAS can optimize whether each pair of

individuals should be mated to maximize the final genetic gain via look-ahead simulations under two

simplified assumptions: one progeny is obtained from each mating pair, and random mating occurs

after the first selection and mating cycle. Although this approach achieved higher genetic gains than

the previous selection criteria in the final generation by considering optimal mating pairs, the

assumptions were too simple to estimate the final genetic gain in practical breeding programs, and

LAS could not optimize the number of progenies allocated to each mating pair, i.e., progeny

allocation strategy.

Thirdly, as for the progeny allocation strategy, only one study has attempted to optimize the

allocation strategy (Hunter and McClosky, 2016) although it is known to have a significant impact on

the ultimate genetic gain (Wang et al., 2018). In their study, the number of progenies allocated to

each pair was optimized by searching for the Pareto surface in the plane formed by the mean and

variance of progenies in the next generation (Hunter and McClosky, 2016), utilizing the multi-

objective optimal computing budget allocation method (Lee et al., 2010). However, this method lacks

the future-oriented perspectives required for the mid-term/long-term breeding process. Moreover,

there is no study on the allocation optimization of progenies for the final genetic gain under multiple

selection and crossing cycles, although a couple of previous studies tried to optimize breeding

program decisions from other viewpoints (Amini et al., 2021; Diot and Iwata, 2022; Moeinizade et
al., 2022).

As described so far, regardless of the duration of breeding schemes, i.e., short-term or long-term,
how decisions are made significantly impacts the ultimate outcome of the schemes when parent
selection and mating are repeated multiple times. Thus, if the optimization framework regarding the
selection of parents and mating pairs is achieved, it can be widely applied to various types of
breeding programs, i.e., small-scale and large-scale breeding programs. As for small-scale breeding
programs with limited resources, particularly in plant breeding, since many minor crops, including
neglected and underutilized species (NUS), have attracted much attention as breeding targets, there
is increasing demand for realizing more efficient breeding in small-scale programs for minor crops
(Padulosi et al., 2013; Kamenya et al., 2021). Even if NUS has attracted little attention and has been
entirely ignored by plant breeders so far, it is now expected to have the potential to diversify
agriculture and address climate change. Another example of small-scale breeding is breeding
schemes promoted by small and medium enterprises (Zambon et al., 2019). Since there are many
agricultural small and medium enterprises in Asian countries, optimizing decision-making in small-
scale breeding will lead to increased profits in the agricultural market. From these viewpoints, our
study, which helps breeders' decisions on hybridization, is essential to improve breeding efficiency
not only in large-scale but also small-scale breeding programs.

The overall objective of the study was to go beyond the conventional selection criterion-based GS by
optimizing a progeny allocation strategy to maximize the ultimate genetic gains after multiple cycles
of selection of parents and mating pairs. There are two significant issues in this optimization. The
first issue is that it is challenging to describe the breeding process explicitly because the future state
of a breeding population is unknown. The second issue is that breeding is a stochastic rather than
deterministic process since progenies are randomly produced in each generation through meiosis.
Therefore, the specific objectives of this study were (1) to evaluate the final genetic gain via future-
oriented breeding simulations and (2) to achieve an optimal mating pair selection via a numerical
optimization approach conducted by an artificial intelligence (AI) breeder while addressing the two
issues above. First, GS combined with future-oriented breeding simulations enables us to evaluate the
final genetic gain in a breeding scheme with multiple cycles, leading to a solution to the first issue.
Second, the AI breeder in this study referred to a virtual breeder in our computer who can optimize
the decisions regarding the progeny allocation strategy by using both the breeding simulations and

some optimization algorithm. In this study, to design this AI breeder, we applied black-box

optimization by considering the breeding process achieved by the above simulations as a black-box

function whose input is a set of parameters for the allocation strategy and whose output is the

ultimate genetic gain. We employed a black-box optimization algorithm, StoSOO (Valko et al.,

2013), which can also optimize a stochastic black-box function, such as a breeding process,

presenting a solution to the second issue. Multiple selection criteria are used to allocate progenies,

and the optimized progeny allocation can automatically determine the optimal combination of criteria

and weights to be used in the allocation. Further, the optimized progeny allocation can help breeders

select the appropriate mating pairs because the AI breeder can offer information on which cross

combination is prospective by showing the optimized number of progenies in continuous values. In

this study, we mainly focus on optimizing the progeny allocation strategy as a proof of concept. Still,

by changing the simulation settings and the parameters to be optimized, we can easily extend our

framework to assist breeders in making decisions in different types of hybridization, such as parental

selection in bi-parental/multi-parental populations.

165   2   Materials and methods

166   2.1   System design

We assumed that breeders proceed with a breeding scheme according to the progeny allocation

strategy proposed by an "AI breeder", a novel decision-making system in this study (Figure 1A). The

breeders can obtain a set of parameters representing the optimal allocation strategy, $\left\{\hat{\mathbf{h}}^{(\tau)}\right\}_{\tau=0,1,2,3}$, by

passing the information on a parent panel, i.e., marker genotype, genetic marker effects, and

recombination rates between genetic markers, to the AI breeder. Here, we briefly describe the

overview of the framework in which the AI breeder optimizes the strategy for the resource allocation

of progenies.

After receiving the parent panel information, the AI breeder attempts to simulate breeding schemes

by defining the allocation strategy (Figure 1A). At each generation τ in the simulated schemes

($\tau \in \{0,1,2,3\}$ is a current generation number in simulations conducted by the AI breeder, whereas

$t \in \{0,1,2,3\}$ appeared in the subsequent subsections is a current generation number in breeding

schemes proceeded by actual breeders), the breeder first computes the weighted breeding values

(WBV) to select the parent candidates from the current generation. Then, after determining mating

pairs by the diallel crossing between the parent candidates, given the parameter $\mathbf{h}^{(\tau)}$, the breeder

automatically determines the number of progenies allocated to each pair $\mathbf{b}^{(\tau)}$, by utilizing the

information on selection criteria $\mathbf{\Omega}^{(\tau)}$ computed from the marker genotype and effects, such as

breeding values (BV), WBV, and expected genetic variance of progenies (GVP) (Figure 1B). In other

words, we assumed that $\mathbf{b}^{(\tau)}$ is determined by applying the softmax function to the weighted sum of

the selection criteria, as shown in Equation 1.

$$\mathbf{b}^{(\tau)} = \left\lfloor N^{(\text{pop})(\tau+1)} \times \frac{\exp\left(\mathbf{\Omega}^{(\tau)}\, \mathbf{h}^{(\tau)}\right)}{\sum_Q \exp\left(\mathbf{\Omega}^{(\tau)}\, \mathbf{h}^{(\tau)}\right)} \right\rfloor, \#(1)$$

where $N^{(\text{pop})(\tau+1)}$ is the population size in the next generation $\tau + 1$, and $\lfloor x \rfloor$ is the floor function of

x (i.e., the maximum integer that is equal to or lower than x). The softmax function is used to

allocate progenies in proportion to probabilities reflecting the weighted sum of the selection criteria.

Here, $\mathbf{h}^{(\tau)}$ can be interpreted as the weighting parameters that correspond to the importance of each

selection criterion. After determining $\mathbf{b}^{(\tau)}$, the AI breeder can simulate the next generation based on

the principle of meiosis. Then, by repeating the parent selection and mating steps toward the final

generation of the breeding scheme, the AI breeder evaluates the performance of the allocation

strategy with $\mathbf{h}^{(\tau)}$. Thus, the goal of the AI breeder is to estimate $\left\{\mathbf{h}^{(\tau)}\right\}_{\tau=0,1,2,3}$ that maximizes the

final genetic gain via simulations of the breeding process. To optimize the weighting parameters, a

black-box optimization algorithm called StoSOO (Valko et al., 2013) was used in this study (Figure

1A). The StoSOO algorithm is an extension of the SOO algorithm (Munos, 2011; Preux et al., 2014),

which performs global optimization of deterministic black-box functions based on a tree-based

search and can be applied to stochastic black-box functions. Using the StoSOO algorithm with a

finite number of function evaluations can provide a global quasi-optimal solution of an implicit

stochastic function within the number of evaluations. To apply StoSOO to our problem, the AI

breeder performs 50 breeding simulations with one set of $\left\{\mathbf{h}^{(\tau)}\right\}_{\tau=0,1,2,3}$, and the empirical mean of

the 50 final genetic gains is regarded as the objective function for StoSOO. The function evaluations

are repeated 20,000 times in StoSOO, and a set of optimized parameters $\left\{\hat{\mathbf{h}}^{(\tau)}\right\}_{\tau=0,1,2,3}$ is passed from

the AI breeder to the real breeders.

In this study, we evaluated this novel optimization framework by conducting simulation studies as a

proof of concept. We prepared six versions of the optimized allocation strategy according to the

components of $\mathbf{\Omega}^{(\tau)}$ as follows: (a) only BV, (b) WBV, (c) both selection criteria (i.e., BV and

WBV), (d) expected GVP in addition to BV, (e) expected GVP in addition to WBV, and (f) expected

GVP in addition to (c) (Figure 1B). We compared these six optimized strategies to (g) the equal

allocation strategy that equally allocates progenies to each pair, that is, $\mathbf{h}^{(\tau)} = \mathbf{0}$, by conducting

10,000 breeding simulations using the optimized $\left\{\hat{\mathbf{h}}^{(\tau)}\right\}_{\tau=0,1,2,3}$ (Figure 1A). In the following results,

we abbreviate the above strategies as (a) BV, (b) WBV, (c) TBV (Two BVs), (d) BVGVP, (e)

WBVGVP, (f) TBVGVP, and (g) EQ. Here, for the BV and WBV strategies, we only used one

criterion for allocating progenies. For simplicity, we also assumed true marker effects to compute the

selection criteria. In other words, true QTL positions and effects were known throughout the study.

The problem of using the true marker effects will be discussed in the **Discussion** section.

2.2   Overview of the methods

This section provides an overview of the method used in this simulation study (Figure S1). We first

simulated genome-wide markers and QTLs for the parent panel and the corresponding QTL effects.

The selection of parent candidates, determination of mating pairs, and allocation of progenies to each

mating pair were then repeated in sequence to move forward with a breeding scheme until the final

generation. Selection and mating processes were carried out based on two selection criteria: BV and

WBV, and the expected GVP. We compared two breeding strategies regarding the allocation of

progenies: equal allocation and allocation optimized by a black-box optimization algorithm, StoSOO.

These strategies were evaluated by simulating 10,000 breeding schemes based on each strategy.

Each element of the simulation study is described in the following subsections. First, the two

selection criteria, BV and WBV, are described, followed by the simulation of genome-wide markers

and QTLs, the selection and mating processes in breeding schemes, and the evaluation of breeding

schemes. Because optimized allocation requires the simulation and evaluation of breeding schemes,

the method of evaluating breeding schemes is explained before describing the optimized allocation

strategy. The calculation of the GVP required for optimized allocation is presented at the end of this

section.

2.3   Selection criteria

234   We used two selection criteria, BV (Meuwissen et al., 2001) and WBV (Goddard, 2009; Jannink,

235   2010), to select parent candidates and allocate progenies in breeding schemes since these are the two

236   typical breeding values (Figure S1 (1)). Here, we only used WBV for the selection step to maintain

237   the diversity, but we used both criteria for the allocation step.

238   The first criterion, BV, can be expressed as in Equation 2:

$$\mathbf{u}^{(\mathrm{BV})(t)} = \sum_{m=1}^{M} \mathbf{w}_m^{(t)} \alpha_m , \#(2)$$

where M is the number of markers, $N^{(\mathrm{pop})(t)}$ is the number of genotypes for a population with

generation t in a breeding scheme, $\mathbf{u}^{(\mathrm{BV})(t)}$ is an $N^{(\mathrm{pop})(t)} \times 1$ vector of BV for generation t, $\mathbf{w}_m^{(t)}$ is

an $N^{(\mathrm{pop})(t)} \times 1$ vector of marker genotype at m-th marker with scores of -1, 0, or 1 for generation

t, and α_m is a true marker effect at the m-th marker. In this study, we assumed that the true marker

effect α_m was known. BV is a criterion for expressing additive genotypic values directly transmitted

to the next generation and is often used in conventional breeding schemes with GS (Meuwissen et al.,

2001).

The second selection criterion, WBV, can be expressed as in Equation 3:

$$\mathbf{u}^{(\mathrm{WBV})(t)} = \sum_{m=1}^{M} \mathbf{w}_m^{(t)} \alpha_m p_m^{(t)-\frac{1}{2}} , \#(3)$$

where $\mathbf{u}^{(\mathrm{WBV})(t)}$ is an $N^{(\mathrm{pop})(t)} \times 1$ vector of WBV for generation t, $p_m^{(t)}$ is the allele frequency at

the m-th marker for generation t, and the other terms are defined in Equation 2. Emphasizing the

effects of rare alleles, as in Equation 3, leads to the successful maintenance of genetic diversity, thus

WBV can benefit the genetic gain in long-term breeding programs compared to BV (Goddard, 2009;

Jannink, 2010).

2.4   Simulation of the parent panel and genome structure

Genome-wide markers and QTLs for founder haplotypes were generated using the coalescent

simulator GENOME (Liang et al., 2007) (Figure S1 (2)). We assumed a diploid with the number of

chromosomes ten ($2n = 20$), and that there were 500 loci, including both markers and QTLs, on

each chromosome. At first, 4,000 founder haplotypes for each chromosome were independently

generated using GENOME with the following parameters: "-pop 1 4000 -N 4000 -c 1 -pieces 100000

-rec 0.0001 -s 4000 -maf 0.01 -tree 0 -mut 0.00000001". Among the 4,000 loci, 500 loci whose

minor allele frequency (MAF) was equal to or larger than 0.01 were randomly selected.

Subsequently, 2,000 founders were generated by sampling two haplotypes from all founder

haplotypes with replacement. These founders were randomly mated for one generation to simulate

the pre-breeding materials of the 2,000 genotypes. These procedures for simulating the genetic

resources are similar to those in the R package "BreedingSchemeLanguage" by Yabe et al. (Yabe et

al., 2017). From these simulated genetic resources, $N^{(\text{pop})(0)} = 250$ genotypes were selected for the

parent panel of a breeding scheme so that these 250 genotypes represented the simulated genetic

resources in terms of genetic diversity. We applied the k-medoids method to the marker genotypes of

the pre-breeding material using the "pam" function of the R package "cluster" version 2.1.2

(Maechler et al., 2021). In the k-medoids method, after clustering the pre-breeding materials into 250

groups, we selected the medoids as the representative genotypes from each group. This parent panel

was regarded as generation $t = 0$ in a breeding scheme. Here, the parent panel consisted of a

relatively small number of genotypes and was also genetically close to founders compared to usual

breeding populations used in programs for major crops. These simulation settings, including the

assumption for the parent panel, are a little apart from situations in large-scale breeding for major

crops but can be justified when we assume the breeding schemes for the NUS whose breeding has

not been promoted in the past.

2.5   Simulation of QTLs and genotypic values

For each chromosome, 400 loci were randomly sampled from the 500 loci and were regarded as

single nucleotide polymorphism (SNP) markers available for breeders. In simulating QTLs and

phenotypes, we assumed quantitative traits with a simple genetic architecture as target traits, as

described below (Figure S1 (3)). First, among the remaining 100 loci, 2 loci per chromosome were

randomly selected as QTLs (i.e., a total of $M^{(\text{QTL})} = 20$ QTLs). This number of QTLs was

determined assuming the situation where quantitative traits controlled by the relatively small number

of QTLs still remain as target traits in small-scale breeding for the NUS. QTL effects were then

sampled from the normal distribution, as shown in Equation 4.

$$p\left(\boldsymbol{\alpha}^{(\text{QTL})}\right) \sim \text{MVN}\left(\mathbf{0}, \frac{1}{M^{(\text{QTL})}}\mathbf{I}_{M^{(\text{QTL})}}\right), \#(4)$$

where $\boldsymbol{\alpha}^{(\text{QTL})}$ is an $M^{(\text{QTL})} \times 1$ vector of QTL effects and $\mathbf{I}_{M^{(\text{QTL})}}$ is an $M^{(\text{QTL})} \times M^{(\text{QTL})}$ identity

matrix. Here, QTL effects were assumed to be additive for simplicity. The true additive genotypic

values $\mathbf{u}^{(\text{TGV})(0)}$ for the parent panel were then simulated using Equation 2, where each marker was

replaced by each QTL ($M = M^{(\text{QTL})}, \mathbf{w}_m = \mathbf{w}_m^{(\text{QTL})(0)}, \alpha_m = \alpha_m^{(\text{QTL})}; \mathbf{w}_m^{(\text{QTL})(0)}$ is an $N^{(\text{pop})(0)} \times 1$

vector of genotype scores at m-th QTL for the parent panel). Thus, $\mathbf{u}^{(\text{TGV})(0)} = \mathbf{u}^{(\text{BV})(0)}$ when the

true marker effect α_m is known as the setting in this study. Since we used the true marker effects, a

heritability was assumed to be 1 throughout the study.

2.6 AI-assisted breeding scheme

The breeding scheme in this study was carried out according to the following steps (Figure S1 (4)).

Here, we assumed small-scale plant breeding programs with recurrent selection for simplicity and

due to the computational limitations, but changing the settings of the following breeding scheme will

lead to the extension of our framework for various types of breeding programs.

1. Select parent candidates (Figure S1 (4-1))

Based on the order of increasing WBV computed using the true marker (QTL) effects $\boldsymbol{\alpha}^{(\mathrm{QTL})}$ for

the current generation t $(t \in \{0, 1, 2, 3\})$, the top $N^{(\mathrm{sel})(t)}$ genotypes were selected as parent

candidates for the next generation. In this study, we assumed $N^{(\mathrm{sel})(t)}$ was constant over

generations, i.e., $N^{(\mathrm{sel})(t)} = N^{(\mathrm{sel})} = 15$. We used only WBV for the selection step in this study

because selecting parents based on BV often failed to maintain the genetic diversity of the

population when we chose a high selection intensity in the optimized allocation strategy.

2. Determine mating pairs for the next generation (Figure S1 (4-2))

From the selected $N^{(\mathrm{sel})(t)}$ parent candidates, diallel crossing, including selfing, was assumed to

determine the mating pairs for the next generation, that is, $N^{(\mathrm{pair})(t)} = \frac{N^{(\mathrm{sel})(t)}(N^{(\mathrm{sel})(t)}+1)}{2} = 120$

pairs were prepared for crossing.

3. Allocate progenies to each mating pair (Figure S1 (4-3))

The following two strategies were used to allocate progenies to each mating pair determined in

Step 2, and then a crossing table was created. Both strategies generated $N^{(\mathrm{pop})(t+1)}$ progenies for

the next generation. In this study, we assumed $N^{(\mathrm{pop})(t+1)}$ was also constant over generations,

i.e., $N^{(\mathrm{pop})(t+1)} = N^{(\mathrm{pop})} = 250$.

a. The optimized allocation method

One strategy was the optimized allocation method developed in this study, which applied

the softmax function to the weighted sum of the selection criteria computed in Step 1. For

more details, please refer to the **Progenies allocation strategies** subsection.

b. Equal allocation method

We also prepared an equal allocation method that allocated $N^{(\text{pop})(t+1)}$ progenies equally to

the $N^{(\text{pair})(t)}$ mating pairs. Thus, three progenies were allocated to each of the 10 mating

pairs, and two progenies were allocated to each of the remaining 110 mating pairs.

4. Generate progenies for the next generation (Figure S1 (4-4))

According to the crossing table created in Step 3, two gametes were generated for each mating

pair, considering the recombination between markers. Recombination rates between markers

were computed using the Kosambi map function (Kosambi, 1943; Zhao and Speed, 1996) based

on the linkage map generated by GENOME. These two gametes were then combined to create

one new progeny, resulting in a new population (with generation $t + 1$) of $N^{(\text{pop})(t+1)}$ progenies.

5. Repeat the parent selection and mating process

Steps 1-4 were repeated four times until the population reached the final generation, i.e., $t = T -$

1 where T is the final generation number.

Finally, the results for each strategy were evaluated using the true additive genotypic values $\mathbf{u}^{(\text{TGV})(T)}$

for the latest population with generation T. Here, the final generation was set as $T = 4$ in this study,

which was determined assuming the situation where we needed to quickly achieve high genetic gains

in small-scale breeding schemes as one example.

2.7   Evaluation of the outcome of breeding schemes

In one breeding scheme, the true additive genotypic value $\mathbf{u}^{(\text{TGV})(t)}$ was computed for each

generation t (Figure S1 (5)). Then, the top $N^{(\text{top})(t)} = 5$ genotypes were chosen in the order of

increasing $\mathbf{u}^{(\text{TGV})(t)}$, and the empirical mean of the $\mathbf{u}^{(\text{TGV})(t)}$ for these $N^{(\text{top})(t)}$ genotypes, $u^{(\text{top})(t)}$,

was computed to evaluate the population maximum for the breeding strategy. Here, we chose

$N^{(\text{top})(t)} = 5$ genotypes for the evaluation to represent the top individuals for developing a new

variety and to control the stochastic variation between simulations to some extent. For a given

simulation dataset of the parent panel, we conducted 10,000 different breeding schemes for each

strategy and evaluated the empirical mean of the $u^{(\text{top})(t)}$ using these 10,000 simulation results. The

simulation and its evaluation were repeated for 10 replicates of the phenotype simulation with

different QTL positions and effects.

2.8   Progenies allocation strategies

In this study, we developed a novel breeding strategy to quasi-optimize each selection criterion

regarding the allocation of progenies (Figure S1 (6) and Figure 1b). In other words, we determined

the number of progenies allocated to each mating pair at generation τ, $\mathbf{b}^{(\tau)}$, by applying the softmax

function to the weighted sum of the selection criteria instead of the equal allocation strategy

described above, as in Equation 1.

$$\mathbf{b}^{(\tau)} = \left\lfloor N^{(\text{pop})(\tau+1)} \times \frac{\exp\left(\mathbf{\Omega}^{(\tau)} \mathbf{h}^{(\tau)}\right)}{\sum_Q \exp\left(\mathbf{\Omega}^{(\tau)} \mathbf{h}^{(\tau)}\right)} \right\rfloor, \#(1)$$

where $\mathbf{b}^{(\tau)}$ is an $N^{(\text{pair})(\tau)} \times 1$ vector representing the number of progenies allocated to each mating

pair at generation τ, $\mathbf{\Omega}^{(\tau)}$ is an $N^{(\text{pair})(\tau)} \times Q$ matrix of Q different selection criteria for an average

progeny of each mating pair (e.g., BV, WBV, or the genetic variance of progenies in a subsequent

generation (GVP) for each pair), and $\lfloor x \rfloor$ is the floor function (i.e., the maximum integer that is equal

to or lower than x). In the softmax function, we can convert a vector of some weighted goodness for

each mating pair, $\mathbf{\Omega}^{(\tau)} \mathbf{h}^{(\tau)}$, to a vector of probabilities reflecting the relative value of each element

by allocating more progenies to better pairs. Here, $\mathbf{h}^{(\tau)}$ is a $Q \times 1$ vector required for

determining $\mathbf{b}^{(\tau)}$, which corresponds to how much importance is given to each selection criterion.

Since the final genetic gain, $g^{(T)}$, can be represented as the output of the black-box function f whose

input is a set of $\mathbf{h}^{(\tau)}$, i.e., $g^{(T)} = f\left(\{\mathbf{h}^{(\tau)}\}_{\tau=0,1,2,3}\right)$, the main goal of this strategy is to estimate $\mathbf{h}^{(\tau)}$

that maximizes $g^{(T)}$ via breeding simulations and the black-box optimization algorithm. When

$\mathbf{h}^{(\tau)} = \mathbf{0}$, the softmax allocation strategy based on Equation 1 is the same as the equal allocation

strategy. The final genetic gain was defined as the improvement in the true genotypic values of the

top $N^{(\text{top})(t)} = 5$ genotypes of the final population compared with those of the parent panel, that is,

$g^{(T)} = u^{(\text{top})(T)} - u^{(\text{top})(0)}$.

2.9   Optimization of $\mathbf{h}^{(\tau)}$ to maximize the final genetic gain

In this study, we used the StoSOO algorithm (Valko et al., 2013) to optimize the parameter vector

$\mathbf{h}^{(\tau)}$ (Figure S1 (7)). The StoSOO algorithm is an extension of the SOO algorithm (Munos, 2011;

Preux et al., 2014), which performs global optimization of deterministic black-box functions based

on a tree-based search and can be applied to stochastic black-box functions. The StoSOO algorithm

375 with a finite number of function evaluations can provide a global quasi-optimal solution of an

376 implicit stochastic function within the number of evaluations. To apply StoSOO in our study, we

377 performed 50 simulations using one set of $\{\mathbf{h}^{(\tau)}\}_{\tau=0,1,2,3}$, and the empirical mean of the final genetic

378 gains $g^{(T)}$ for these 50 simulations was used as the objective function for StoSOO. Here, we defined

379 the domain of definition as $[0, 2]$ and set 1 as an initial parameter for each element of $\{\mathbf{h}^{(\tau)}\}_{\tau=0,1,2,3}$.

380 The function evaluations were repeated 20,000 times, giving us quasi-optimal solutions,

381 $\{\hat{\mathbf{h}}^{(\tau)}\}_{\tau=0,1,2,3}$, for this optimization problem. We then performed 10,000 simulations based on the

382 estimated $\{\hat{\mathbf{h}}^{(\tau)}\}_{\tau=0,1,2,3}$ to evaluate each optimized strategy developed in this study, as described in

383 the **Evaluation of results** subsection.

385 **2.10 Selection criteria and genetic diversity of progenies as the candidates for $\mathbf{\Omega}^{(\tau)}$**

386 As the candidates for $\mathbf{\Omega}^{(\tau)}$ to compute $\mathbf{b}^{(\tau)}$, we used the two selection criteria, BV and WBV, and the

387 expected GVP for each mating pair (Figure S1 (8)). The BVs and WBVs of the parent candidates

388 were computed using Equations 2 and 3, and the mean of the selection criteria of the two parents for

389 each mating pair was used as one column vector of $\mathbf{\Omega}^{(\tau)}$. This vector, representing BV or WBV for

390 each mating pair, was the same as that for the average progeny of each mating pair in this study

391 because we only assumed additive QTL effects. To compute the GVP, we used Equation 5, based on

392 the idea proposed by Lynch and Walsh (Lynch and Walsh, 1998; Zhong and Jannink, 2007;

393 Lehermeier et al., 2017):

$$v_n^{(\tau)} = \frac{1}{4}\{\boldsymbol{\alpha}^{\mathrm{T}}\mathbf{C}^{(n_{\mathrm{M}})}\boldsymbol{\alpha} + \boldsymbol{\alpha}^{\mathrm{T}}\mathbf{C}^{(n_{\mathrm{P}})}\boldsymbol{\alpha}\}, \#(5)$$

394 where $v_n^{(\tau)}$ is an expected GVP for mating pair n with generation τ, $\boldsymbol{\alpha}$ is a vector of the true marker

395 effects, $\mathbf{C}^{(n_{\mathrm{M}})}$ and $\mathbf{C}^{(n_{\mathrm{P}})}$ are $M \times M$ covariance matrices caused by the segregation of maternal parent

396 n_{M} and paternal parent n_{P} of mating pair n, respectively. A diagonal (m, m) element of $\mathbf{C}^{(n_{\mathrm{M}})}$,

397 $C_{mm}^{(n_{\mathrm{M}})}$, is 0 if an m-th marker of n_{M} is homozygote, and 1 if otherwise (i.e., heterozygote), and a non-

398 diagonal (m_1, m_2) element of $\mathbf{C}^{(n_{\mathrm{M}})}$, $C_{m_1 m_2}^{(n_{\mathrm{M}})}$, is 0 if the m_1-th or m_2-th marker of n_{M} is homozygote,

399 $1 - 2r_{m_1 m_2}$ if both m_1-th and m_2-th markers are heterozygote without recombination, and $2r_{m_1 m_2} -$

400 1 if otherwise (i.e., heterozygote with recombination). Here, $r_{m_1 m_2}$ is the recombination rate between

the m_1-th and m_2-th markers. In this study, after computing the expected GVP for all mating pairs,

$\mathbf{v}^{(\tau)}$ was used as one column vector of $\mathbf{\Omega}^{(\tau)}$.

We prepared six different strategies for the components of $\mathbf{\Omega}^{(\tau)}$ and compared them to the non-

optimized strategy as written in the **System Design** subsection, i.e., (a) BV, (b) WBV, (c) TBV, (d)

BVGVP, (e) WBVGVP, (f) TBVGVP, and (g) EQ. Even when we used only one criterion for the

allocation of progenies in BV and WBV strategies, we optimized the weighting parameter

$\left\{\mathbf{h}^{(\tau)}\right\}_{\tau=0,1,2,3}$ by focusing on the optimization of the gradient of the number of progenies allocated to

each pair against the relative value of that criterion.

3   Results

3.1   Evaluation of convergence conditions of the StoSOO algorithm

For the optimization of the allocation strategy to work, it was necessary to confirm that the solution

obtained from the algorithm used for optimization, the StoSOO algorithm, converged stably. Thus,

we first evaluated the convergence of the solution using the StoSOO algorithm for each strategy. As

described above, the AI breeder conducted 50 simulations of the breeding scheme for one function

evaluation and repeated 20,000 function evaluations to obtain parameter sets that were quasi-

optimized by the StoSOO algorithm. We recorded the quasi-optimized parameter sets for each

evaluation step of the objective function and evaluated the corresponding function values. For each of

the six optimized allocation strategies, the change in the function values was plotted and compared

with the function value based on the equal allocation strategy (Figure S2).

For all strategies on the components of $\mathbf{\Omega}^{(\tau)}$, the function value increased almost monotonically and

reached the quasi-optimized function value after the 20,000 evaluations, which was much larger than

that under the equal allocation strategy (Figure S2). These results indicated that StoSOO successfully

improved the expected final genetic gain evaluated by the simulations and optimized the parameter

set $\left\{\mathbf{h}^{(\tau)}\right\}_{\tau=0,1,2,3}$, resulting in the superiority of the optimized allocation strategy over the equal

allocation strategy.

3.2   Genetic gains over four generations

For all strategies on the components of $\mathbf{\Omega}^{(\tau)}$, we compared the optimized and equal allocation strategies by plotting the change in the genetic gain for each generation (Figure 2A). We then defined genetic gain in each generation as the difference between the empirical means of the true genotypic values of the top $N^{(top)(t)} = 5$ genotypes in the current generation (generation t) and the parent panel (generation 0). Using the optimized allocation strategies obtained after 20,000 function evaluations, we evaluated the expected genetic gains of the seven strategies, including the six optimized and the equal allocation strategies, based on 10,000 breeding simulation results using one replication for the quantitative phenotype simulation.

In the final generation, $t = 4$, all the strategies with optimized allocations showed higher genetic gains than the equal allocation strategy (Figure 2A). Strategies that considered GVP outperformed those without GVP in the final generation, while the strategies that only considered the selection criteria BV and WBV showed little difference in genetic gains.

In the first generation, $t = 1$, the strategies that considered GVP showed lower genetic gains than those without GVP (Figure 2A). After the second generation, $t = 2$, strategies with GVP overtook those without GVP, increasing their advantage over the other strategies through the final generation.

3.3   Genetic gains across different simulation repetitions

To assess the characteristics of each strategy in more detail, we computed the cumulative distribution functions (CDFs) of genetic gain in the final generation, $t = 4$, for each strategy based on 10,000 simulation repetitions. In Figure 2B, the horizontal axis represents the final genetic gain of an individual, whereas the vertical axis represents the percentile of simulation repetitions. Because the 1st and 99th percentiles correspond to the worst and best performances within the 10,000 simulation repetitions, respectively, the more a CDF curve of a strategy moves toward the right and downward in the graph, the better the performance of the strategy is.

Again, the optimized strategies outperformed the equal allocation strategy; among the optimized strategies, those with GVP showed better performance than those without GVP in terms of the CDFs (Figure 2B). The CDF curves for the optimized strategies were parallel to the vertical axis at the location where the genetic gain was approximately 1.15, suggesting a local optimum within the simulation repetitions. In the optimized strategies with GVP, about 50 % of the breeding simulations

459 achieved genetic gains above this local optimum (BVGVP, 55.1 %, WBVGVP, 47.6 %, TBVGVP,

460 49.4 %), which were much higher than those in the equal allocation strategy (EQ, 1.85 %) and

461 strategies without GVP (BV, 16.0 %, WBV, 16.4 %, TBV, 30.1 %).

3.4 Genetic gains in the final generation using ten replications of the phenotype simulation

We also evaluated the genetic gains in the final generation, $t = 4$, using ten replications of the

phenotype simulation to confirm that the optimized strategies showed better results than the equal

allocation strategy for the traits with different quantitative trait loci (QTL) positions and effects

(Figure 2C). When simulating phenotypes for different traits, we changed QTL positions and their

effects, but the number of QTLs and distribution of QTL effects were fixed. Also, the heritabilities

were assumed to be 1 for all ten replications since the true marker effects were utilized for the

optimization.

To reduce the computational time for the optimization in evaluating ten replications of the phenotype

simulation, optimized (or quasi-optimized) allocation strategies were determined after 4,000 function

evaluations. The number of function evaluations was determined to ensure that the function value

reached a certain level, although it did not converge, and to reduce the computation time based on the

results in the subsection that evaluated the convergence conditions of the StoSOO. Here, the StoSOO

algorithm does not necessarily have to be converged since the evaluation for the ten replicates of

phenotype simulation was conducted to prove that the optimization algorithm could improve the

genetic gain to some extent compared to the non-optimized strategy.

Even when the QTL positions and effects of the target trait were changed, the optimized strategies

consistently outperformed the equal allocation strategy, and the strategies with GVP outperformed

those without GVP (Figure 2C). When we evaluated the improvement rate in the final genetic gain of

each optimized strategy compared to the equal allocation strategy using ten replications of the

phenotype simulation, a similar trend was observed between the strategies with and without GVP

(Table 1). In addition, the optimized strategies at least improved the genetic gain compared to the

equal allocation for all ten replications. In particular, the final genetic gains for TBVGVP improved

by 7.22–12.4 % compared to those for the equal allocation strategy within just four generations

(Table 1).

3.5 Genetic diversity over four generations

We compared the optimized and equal allocation strategies in genetic diversity across generations (Figure 2D). The genetic diversity of a population in each generation was calculated as the genetic variance of the true genotypic values in the population.

First, in $t = 1$, all the optimized strategies, except WBVGVP, showed lower genetic variances than those in the parent panel, $t = 0$ (Figure 2D). From $t = 2$ to $t = 4$, genetic diversity was preserved in all strategies compared to $t = 0$. In these generations, strategies with GVP maintained higher genetic variance than those without GVP and the equal allocation strategy. The difference between the strategies with or without GVP was most observed at $t = 3$, followed by a sharp decrease in genetic variance for the former strategies through the final generation, $t = 4$.

In contrast, strategies using different selection criteria (BV or WBV) showed little difference in genetic variance (Figure 2D). The strategy using only WBV as a selection criterion appeared to maintain a higher genetic variance than those using BV or both BV and WBV. However, the difference was much smaller than between strategies with or without GVP.

3.6 Optimized weighting parameters for each strategy

After the 20,000 function evaluations, we obtained a set of optimized weighting parameters $\{\mathbf{h}^{(\tau)}\}_{\tau=0,1,2,3}$ for each allocation strategy (Tables 2 and Tables S1 and S2). A large value of $\mathbf{h}^{(\tau)}$ indicates a large weight of the corresponding criterion.

As for the strategies without GVP, the weighting parameters in generation $\tau = 1$, $\mathbf{h}^{(1)}$, showed the smallest values among all generations, and the weights in the latter generations increased through the final generation (Table 2 and Tables S1 and S2). Using strategies with GVP, the weighting parameters for the selection criteria (BV and WBV) showed higher values in generation $\tau = 1$ than in other generations. The tendency of the selection criteria to increase in later generations was unchanged in the strategies with GVP, and the weights for GVP were similar to those of other selection criteria across generations.

3.7 Optimized progeny allocation in generation $t = 0$

Finally, we present the optimized number of progenies allocated to each mating pair in generation

$t = 0$, $\mathbf{b}^{(0)}$. Here, we compared $\mathbf{b}^{(0)}$ in breeding schemes based on the equal and the six optimized

allocation strategies (Table S3). The trend of which cross is more preferred was almost the same

between the six optimized strategies, but the trend of how progenies are allocated to crosses was

quite different. For example, the optimized strategies without GVP, such as WBV and TBV, tended

to allocate more progenies to the best pair (G0_0669 x G0_0669). In contrast, the strategies with

GVP, such as BVGVP and WBVGVP, tended to allocate progenies to many pairs so that the genetic

diversity was maintained. From the results, we can see which mating pair was more emphasized in

each strategy in not discrete but continuous ways. Thus, even if it is not practical for real breeders to

carry out the exact allocation proposed by the AI breeder, the information on the optimized

continuous progeny allocation will help breeders select appropriate mating pairs.

4   Discussion

In this study, we utilized the StoSOO algorithm to optimize the allocation of progenies and obtained

quasi-optimal solutions for the set of weighting parameters $\mathbf{h}^{(\tau)}$ given a finite number of function

evaluations. Utilizing these quasi-optimized weighting parameters for the allocation strategy can be

justified because the true genetic gain corresponding to each set of parameters almost increased as

StoSOO conducted more function evaluations (Figure S2). Thus, when applying this novel

optimization approach to actual breeding schemes, the appropriate number of function evaluations

will be determined according to the computational resources, and within that number of evaluations,

the quasi-optimized parameters obtained from StoSOO will be used to allocate progenies.

After optimizing the weighting parameters using StoSOO, we compared the optimized allocation

strategies with equal allocation strategies. The simulation results showed that the final genetic gain

was higher in the following order: optimized strategies with GVP (BVGVP, WBVGVP, TBVGVP),

optimized strategies without GVP (BV, WBV, TBV), and non-optimized strategy (EQ) (Figure 2A,

C). These results suggest that the developed optimization framework for the progeny allocation is

efficient in conventional GS and that considering the genetic diversity for later generations in the

optimization framework can further improve the final genetic gains even in mid-term breeding

programs ($T = 4$). In particular, the superiority of the optimized strategies with GVP was significant

because those strategies guaranteed a certain level of the final genetic gains compared to other

strategies, even when random factors, such as recombination between markers and segregation of

alleles, are not realized in a real breeding scheme as they are in the optimization process (Figure 2B).

Thus, because the optimized strategies with GVP are expected to consistently produce better

genotypes than conventional GS, breeders can easily adopt our novel optimization framework in

actual breeding schemes. Also, even when breeders find it difficult to implement the progeny

allocation strategy proposed by the AI breeder as it is in actual breeding schemes, they can use

progeny allocation results to select mating pairs as the results provide the information on which cross

is likely to contribute to the production of better genotypes in the future generation (Table S3).

On the other hand, the impact on the final genetic gain caused by the type of criteria (BV, WBV, or

both) used for allocation was much smaller than that caused by the existence of GVP (Figure 2A, C).

Because WBV was always utilized in the selection step in this study, we succssfully maintained the

genetic diversity throughout the breeding scheme regardless of the allocation strategy (Figure 2D),

which may reduce the superiority of WBV in the allocation step against BV. We do not show the

results here, but when BV was utilized in the selection step, the final genetic gains drastically

decreased compared to when WBV was utilized for selection, regardless of the allocation strategy.

Thus, since the choice of selection criteria in the selection step has a more significant impact on the

final genetic gain than in the allocation step, it is strongly recommended that WBV instead of BV be

used in the selection step when applying the developed method to a breeding scheme.

When focusing on the genetic gains in $t = 1$, the optimized strategies with GVP showed lower

genetic gains than those without GVP (Figure 2A). In contrast, the optimized strategies with GVP

showed larger genetic variances than those without GVP from $t = 1$ (Figure 2D). These results

suggest that the optimized strategies with GVP were able to attain higher genetic gains in the final

generation by their success in maintaining genetic diversity throughout the schemes and converting it

into genetic gain toward the final generation, even in mid-term breeding programs ($T = 4$). When

focusing on the optimized parameters for the strategies with GVP, the optimized weights became

larger after $t = 2$, which meant that the selection intensity automatically became higher in later

generations (Table 2). In addition, when focusing on the optimized weighting parameters for the

strategies without GVP, the optimized weight for BV in $t = 1$ was smaller than that for WBV. This

may be because the larger weight of BV in the earlier generation led to less diversity, and the

optimizer controlled the selection intensity by adjusting the weight to avoid such a situation. Thus,

the developed framework to optimize the allocation strategy of progenies can automatically adjust the

weights in each generation in an interpretable form for breeders and is expected to deal with various

situations, such as breeding programs with varying deadlines T, by estimating proper weights for

each selection criterion. In other words, although we assumed the situation close to small-scale plant

breeding schemes with a relatively small number of breeding cycles and a small population size as

one example of the simulation settings in this study, we can easily apply our novel framework to

larger-scale plant breeding programs, and further animal breeding programs, by considering the

assumed conditions such as mating constraints and changing the simulation settings proceeded by the

AI breeder. Also, although we focused on the allocation strategies to prove that our future-oriented

simulations can optimize the decision-making in breeding, we can further extend our framework to

consider the optimization of other strategies, such as selection intensity or selection criteria used in

the selection step by introducing parameters regarding those strategies to the simulation settings.

The developed future-oriented simulation-based framework achieved much higher final genetic gains

than the GS method without optimized allocation of progenies. In particular, the final genetic gains

for TBVGVP improved by more than 9 % on average compared with those for the non-optimized

equal allocation strategy (Table 1). This improvement was quite significant for the genetic gains in

just four generations because we compared the developed framework to the GS based on WBV

(Goddard, 2009; Jannink, 2010), one of the most promising selection strategies in mid- and long-term

breeding programs to date, not phenotypic selection or conventional GS based on BV (Meuwissen et

al., 2001). Also, since the optimization algorithm was forced to terminate in the middle for the results

of the ten replicates for the phenotype simulation due to the computational limitation, we may still

underestimate our optimized results and can expect higher improvement than that obtained in Table 1.

However, it is important to note that we used true marker effects to compute the selection criteria and

GVP in this study, as the previous studies proposing the selection criteria did (Kemper et al., 2012;

Daetwyler et al., 2015; Goiffon et al., 2017; Müller et al., 2018; Moeinizade et al., 2019). In actual

breeding schemes, we cannot know such true marker effects and must use estimated marker effects

based on GP models. Because the optimization algorithm developed in this study can be largely

influenced by the estimation accuracy of GP models, developing a robust method to optimize the

allocation strategy in such cases is crucial in future studies. The other important factor when

assuming real breeding schemes is the timing of GP model updates. If we can appropriately update

the GP model, the estimation accuracy of the marker effects and optimization performance of the

allocation strategies will be improved, which will directly lead to further improvement in the final

genetic gains. In addition, we can deal with long breeding schemes by conducting appropriate model

updates. Thus, we will further investigate the potential of our optimization framework by assuming

real breeding schemes with the factors discussed above.

Genomic selection is a promising technique that contributes to the acceleration of breeding. In this

study, we introduced and developed a novel framework that can upgrade conventional GS in

breeding schemes by optimizing the allocation strategy of progenies via future-oriented simulations.

To optimize the allocation strategy, we parameterized the strategy via softmax conversion by

utilizing the selection criteria and expected genetic variance of progenies. From the simulation

results, our novel framework with the optimized allocation significantly outperformed the non-

optimized strategies, especially when we added the genetic variance of progenies to the algorithm.

Thus, this future-oriented optimization framework for breeding schemes can contribute to the

acceleration, high efficiency, and optimization of plant breeding at the hybridization stage, which

will lead to the optimization of various kinds of decision-making in plant and animal breeding.

DATA AVAILABILITY STATEMENT

All data were simulated in this study. Because we performed many breeding simulations, we cannot

share all datasets themselves. Instead, we share our scripts, including the simulation of data. The

scripts used in this study are available from the "KosukeHamazaki/SCOBS" repository on GitHub,

https://github.com/KosukeHamazaki/SCOBS. In this study, we utilized the R package

"myBreedSimulatR" to simulate and optimize breeding schemes. "myBreedSimulatR" is available

from the "KosukeHamazaki/myBreedSimulatR" repository on GitHub,

https://github.com/KosukeHamazaki/myBreedSimulatR. This package is a variant of

"breedSimulatR", which is available from the "ut-biomet/breedSimulatR" repository on GitHub,

https://github.com/ut-biomet/breedSimulatR.

AUTHOR CONTRIBUTIONS

KH implemented the simulation of breeding schemes, developed the method for optimizing

allocation strategies in breeding schemes, and drafted the manuscript. KH and HI conceived and

designed the study. HI provided administrative support and supervised the study. All authors have
read and approved the final manuscript.

FUNDING

This study was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI
Grant Number JP 22H02306. The funders had no role in the study design, data collection, analysis,
decision to publish, or preparation of the manuscript.

ACKNOWLEDGMENTS

We are grateful to Mr. Julien Diot, a developer of the "breedSimulatR" package, for fruitful
discussions on the implementation of breeding simulations. We would like to thank Dr. Yohei
Akimoto for fruitful discussions on how to optimize decision-making in a breeding scheme. We are
also grateful to Dr. Bo Zhang for her valuable comments when revising our manuscript. We would
like to thank the Japan Society for the Promotion of Science (JSPS) for funding our study. We would
also like to thank Editage (www.editage.com) for English language editing.

CONFLICT OF INTEREST STATEMENT

Not applicable. The authors declare that they have no competing interests.

SUPPLEMENTAL DATA

The Supplementary Material for this article can be found online at:

REFERENCES

Akdemir, D., Beavis, W., Fritsche-Neto, R., Singh, A. K., and Isidro-Sánchez, J. (2019). Multi-objective optimized genomic breeding strategies for sustainable food improvement. *Heredity* 122, 672–683.

Amini, F., Franco, F. R., Hu, G., and Wang, L. (2021). The look ahead trace back optimizer for genomic selection under transparent and opaque simulators. *Sci. Rep.* 11, 4124.

Bulmer, M. G. (1971). The Effect of Selection on Genetic Variability. *Am. Nat.* 105, 201–211.

Capper, J. L. (2011). Replacing rose-tinted spectacles with a high-powered microscope: The historical versus modern carbon footprint of animal agriculture. *Anim. Front* 1, 26–32.

Cowling, W. A., Castro-Urrea, F. A., Stefanova, K. T., Li, L., Banks, R. G., Saradadevi, R., et al. (2023). Optimal Contribution Selection Improves the Rate of Genetic Gain in Grain Yield and Yield Stability in Spring Canola in Australia and Canada. *Plants* 12. doi: 10.3390/plants12020383.

Crossa, J., Beyene, Y., Kassa, S., Pérez, P., Hickey, J. M., Chen, C., et al. (2013). Genomic prediction in maize breeding populations with genotyping-by-sequencing. *G3* 3, 1903–1926.

Crossa, J., Pérez-Rodríguez, P., Cuevas, J., Montesinos-López, O., Jarquín, D., de Los Campos, G., et al. (2017). Genomic Selection in Plant Breeding: Methods, Models, and Perspectives. *Trends Plant Sci.* 22, 961–975.

Daetwyler, H. D., Hayden, M. J., Spangenberg, G. C., and Hayes, B. J. (2015). Selection on Optimal Haploid Value Increases Genetic Gain and Preserves More Genetic Diversity Relative to Genomic Selection. *Genetics* 200, 1341–1348.

de Los Campos, G., Hickey, J. M., Pong-Wong, R., Daetwyler, H. D., and Calus, M. P. L. (2013). Whole-genome regression and prediction methods applied to plant and animal breeding. *Genetics* 193, 327–345.

de los Campos, G., Naya, H., Gianola, D., Crossa, J., Legarra, A., Manfredi, E., et al. (2009). Predicting quantitative traits with regression models for dense molecular markers and pedigree. *Genetics* 182, 375–385.

Diot, J., and Iwata, H. (2022). Bayesian optimisation for breeding schemes. *Front. Plant Sci.* 13,
1050198.

Eggen, A. (2012). The development and application of genomic selection as a new breeding
paradigm. *Anim. Front.* 2, 10–15.

Falconer, D. S., and Mackay, T. F. C. (1996). *Introduction to Quantitative Genetics*. Harlow, Essex,
UK.: Longmans Green.

García-Ruiz, A., Cole, J. B., VanRaden, P. M., Wiggans, G. R., Ruiz-López, F. J., and Van Tassell,
C. P. (2016). Changes in genetic selection differentials and generation intervals in US
Holstein dairy cattle as a result of genomic selection. *Proc. Natl. Acad. Sci. U. S. A.* 113,
E3995-4004.

Gianola, D., and van Kaam, J. B. C. H. M. (2008). Reproducing kernel hilbert spaces regression
methods for genomic assisted prediction of quantitative traits. *Genetics* 178, 2289–2303.

Goddard, M. (2009). Genomic selection: prediction of accuracy and maximisation of long term
response. *Genetica* 136, 245–257.

Goddard, M. E., Hayes, B. J., and Meuwissen, T. H. E. (2011). Using the genomic relationship
matrix to predict the accuracy of genomic selection. *J. Anim. Breed. Genet.* 128, 409–421.

Goiffon, M., Kusmec, A., Wang, L., Hu, G., and Schnable, P. S. (2017). Improving Response in
Genomic Selection with a Population-Based Selection Strategy: Optimal Population Value
Selection. *Genetics* 206, 1675–1682.

González-Camacho, J. M., Crossa, J., Pérez-Rodríguez, P., Ornella, L., and Gianola, D. (2016).
Genome-enabled prediction using probabilistic neural network classifiers. *BMC Genomics* 17,
208.

Gorjanc, G., Gaynor, R. C., and Hickey, J. M. (2018). Optimal cross selection for long-term genetic
gain in two-part programs with rapid recurrent genomic selection. *Theor. Appl. Genet.* 131,
1953–1966.

Grundy, B., Villanueva, B., and Woolliams, J. A. (1998). Dynamic selection procedures for

constrained inbreeding and their consequences for pedigree development. *Genet. Res.* 72,

159–168.

Habier, D., Fernando, R. L., Kizilkaya, K., and Garrick, D. J. (2011). Extension of the bayesian

alphabet for genomic selection. *BMC Bioinformatics* 12, 186.

Hayes, B. J., Bowman, P. J., Chamberlain, A. J., and Goddard, M. E. (2009). Invited review:

Genomic selection in dairy cattle: progress and challenges. *J. Dairy Sci.* 92, 433–443.

Hickey, J. M., Dreisigacker, S., Crossa, J., Hearne, S., Babu, R., Prasanna, B. M., et al. (2014).

Evaluation of genomic selection training population designs and genotyping strategies in

plant breeding programs using simulation. *Crop Sci.* 54, 1476–1488.

Hunter, S. R., and McClosky, B. (2016). Maximizing quantitative traits in the mating design problem

via simulation-based Pareto estimation. *IIE Trans.* 48, 565–578.

Jannink, J.-L. (2010). Dynamics of long-term genomic selection. *Genet. Sel. Evol.* 42, 35.

Jannink, J.-L., Lorenz, A. J., and Iwata, H. (2010). Genomic selection in plant breeding: from theory

to practice. *Brief. Funct. Genomics* 9, 166–177.

Jansen, G. B., and Wilton, J. W. (1985). Selecting mating pairs with linear programming techniques.

J. Dairy Sci. 68, 1302–1305.

Kamenya, S. N., Mikwa, E. O., Song, B., and Odeny, D. A. (2021). Genetics and breeding for

climate change in Orphan crops. *Theor. Appl. Genet.* 134, 1787–1815.

Kemper, K. E., Bowman, P. J., Pryce, J. E., Hayes, B. J., and Goddard, M. E. (2012). Long-term

selection strategies for complex traits using high-density genetic markers. *J. Dairy Sci.* 95,

4646–4656.

Kinghorn, B. P. (2011). An algorithm for efficient constrained mate selection. *Genet. Sel. Evol.* 43, 4.

Kosambi, D. D. (1943). The estimation of map distances from recombination values. *Ann. Eugen.* 12,

172–175.

Lee, L. H., Chew, E. P., Teng, S., and Goldsman, D. (2010). Finding the non-dominated Pareto set
for multi-objective simulation models. *IIE Trans.* 42, 656–674.

Lehermeier, C., Teyssèdre, S., and Schön, C.-C. (2017). Genetic Gain Increases by Applying the
Usefulness Criterion with Improved Variance Prediction in Selection of Crosses. *Genetics*
207, 1651–1661.

Li, Y., Kadarmideen, H. N., and Dekkers, J. C. M. (2008). Selection on multiple QTL with control of
gene diversity and inbreeding for long-term benefit. *J. Anim. Breed. Genet.* 125, 320–329.

Liang, L., Zöllner, S., and Abecasis, G. R. (2007). GENOME: a rapid coalescent-based whole
genome simulator. *Bioinformatics* 23, 1565–1567.

Lin, Z., Shi, F., Hayes, B. J., and Daetwyler, H. D. (2017). Mitigation of inbreeding while preserving
genetic gain in genomic breeding programs for outbred plants. *Theor. Appl. Genet.* 130, 969–
980.

Lynch, M., and Walsh, B. (1998). *Genetics and Analysis of Quantitative Traits*. Sinauer.

Maechler, M., Rousseeuw, P., Struyf, A., Hubert, M., and Hornik, K. (2021). cluster: Cluster
Analysis Basics and Extensions. Available at: https://CRAN.R-project.org/package=cluster.

Meuwissen, T. H. (1997). Maximizing the response of selection with a predefined rate of inbreeding.
J. Anim. Sci. 75, 934–940.

Meuwissen, T. H., Hayes, B. J., and Goddard, M. E. (2001). Prediction of total genetic value using
genome-wide dense marker maps. *Genetics* 157, 1819–1829.

Moeinizade, S., Hu, G., and Wang, L. (2022). A reinforcement Learning approach to resource
allocation in genomic selection. *Intelligent Systems with Applications* 14, 200076.

Moeinizade, S., Hu, G., Wang, L., and Schnable, P. S. (2019). Optimizing Selection and Mating in
Genomic Selection with a Look-Ahead Approach: An Operations Research Framework. *G3*
9, 2123–2133.

Müller, D., Schopp, P., and Melchinger, A. E. (2018). Selection on Expected Maximum Haploid
Breeding Values Can Increase Genetic Gain in Recurrent Genomic Selection. *G3* 8, 1173–
1181.

Munos, R. (2011). Optimistic Optimization of a Deterministic Function without the Knowledge of its
Smoothness. in *Advances in Neural Information Processing Systems*, eds. J. Shawe-Taylor, R.
Zemel, P. Bartlett, F. Pereira, and K. Q. Weinberger (Curran Associates, Inc.). Available at:
https://proceedings.neurips.cc/paper/2011/file/7e889fb76e0e07c11733550f2a6c7a5a-
Paper.pdf.

Padulosi, S., Bioversity International, Thompson, J., and Rudebjer, P. G. (2013). *Fighting poverty,*
hunger and malnutrition with neglected and underutilized species: needs, challenges and the
way forward. Bioversity International.

Preux, P., Munos, R., and Valko, M. (2014). Bandits attack function optimization. in *2014 IEEE*
Congress on Evolutionary Computation (CEC), 2245–2252.

Rabier, C.-E., Barre, P., Asp, T., Charmet, G., and Mangin, B. (2016). On the Accuracy of Genomic
Selection. *PLoS One* 11, e0156086.

Sonesson, A. K., Woolliams, J. A., and Meuwissen, T. H. E. (2012). Genomic selection requires
genomic control of inbreeding. *Genet. Sel. Evol.* 44, 27.

Valko, M., Carpentier, A., and Munos, R. (2013). Stochastic Simultaneous Optimistic Optimization.
in *Proceedings of the 30th International Conference on Machine Learning* Proceedings of
Machine Learning Research., eds. S. Dasgupta and D. McAllester (Atlanta, Georgia, USA:
PMLR), 19–27.

Van Grevenhof, E. M., Van Arendonk, J. A. M., and Bijma, P. (2012). Response to genomic
selection: the Bulmer effect and the potential of genomic selection when the number of
phenotypic records is limiting. *Genet. Sel. Evol.* 44, 26.

VanRaden, P. M. (2008). Efficient methods to compute genomic predictions. *J. Dairy Sci.* 91, 4414–
4423.

Wang, L., Zhu, G., Johnson, W., and Kher, M. (2018). Three new approaches to genomic selection.
Plant Breed. 137, 673–681.

Yabe, S., Iwata, H., and Jannink, J.-L. (2017). A simple package to script and simulate breeding
schemes: The breeding scheme language. *Crop Sci.* 57, 1347–1354.

Zambon, I., Cecchini, M., Egidi, G., Saporito, M. G., and Colantoni, A. (2019). Revolution 4.0:
Industry vs. Agriculture in a Future Development for SMEs. *Processes* 7, 36.

Zhang, Z., and Wang, L. (2022). A look-ahead approach to maximizing present value of genetic
gains in genomic selection. *G3* 12. doi: 10.1093/g3journal/jkac136.

Zhao, H., and Speed, T. P. (1996). On genetic map functions. *Genetics* 142, 1369–1377.

Zhong, S., and Jannink, J.-L. (2007). Using quantitative trait loci results to discriminate among
crosses on the basis of their progeny mean and variance. *Genetics* 177, 567–576.

FIGURES and TABLES

Figure 1. System design assumed in this study.
(**A**) The framework to optimize decision-making in the breeding scheme. (**B**) The resource allocation
strategy of progenies developed in this study.

Figure 2. Comparison of the seven strategies (six optimized and one non-optimized) in four
ways.

(**A**) Change in the expected genetic gains over four generations. (**B**) Genetic gains across different
simulation repetitions. (**C**) Genetic gains in the final generation using ten replications for the
phenotype simulation. (**D**) Change in the genetic variances over four generations. In all Figures, we
compared the seven strategies: EQ: equal allocation (ABD: black solid, C: grey), BV: optimized
allocation based on BV (ABD: orange dashed, C: yellow), WBV: optimized allocation based on
WBV (ABD: blue dashed, C: light blue), TBV: optimized allocation based on BV and WBV (ABD:
red dashed, C: light pink), BVGVP: optimized allocation based on BV and GVP (ABD: orange solid,

C: yellow), WBVGVP: optimized allocation based on WBV and GVP (ABD: blue solid, C: light

blue), and TBVGVP: optimized allocation based on BV, WBV, and GVP (ABD: red solid, C: light

pink). In Figures A and D, generations 0 and 4 correspond to the populations in the initial and final

generations, respectively. In addition, generations 1–3 correspond to populations in the middle of the

breeding scheme. In Figure B, the horizontal axis shows the final genetic gain of an individual,

whereas the vertical axis represents the percentile of the simulation repetitions. As shown in Figure

C, for each optimized strategy, the weighting parameter was optimized based on 4,000 function

evaluations.

Table 1. Minimum, mean, and maximum improvement rate in the final genetic gain of the optimized

strategies compared to equal allocation strategy among ten different target traits (%)

Method	Min (%)	Mean (%)	Max (%)
BV	0.00	2.97	4.89
WBV	2.07	2.88	3.89
TBV	2.11	3.52	5.86
BVGVP	6.40	8.63	12.4
WBVGVP	6.99	8.28	9.95
TBVGVP	7.22	9.33	12.4

827   **Table 2.** Optimized weighting parameters $\mathbf{h}^{(\tau)}$ in different generations $\tau = 0, \ldots, T-1$ for BV and

828   BVGVP strategies

	BV			BVGVP	
Generation (τ)	BV	GVP		BV	GVP
0	1.00	–		1.00	0.33
1	0.33	–		1.00	1.00
2	1.67	–		1.67	1.67

| | 3 | 1.89 | – | 1.67 | 1.67 |

A

Optimization framework by AI breeder

Forward simulation of breeding programs

$\left(\begin{array}{c}\text{50 simulations to perform one}\\\text{black-box function evaluation}\end{array}\right)$

$g^{(T)} = f(\{h^{(\tau)}\})$

AI breeder

$\{b^{(\tau)}\}$

Breeding population

Selection, Mating & Allocation ($b^{(\tau)}$)

$g^{(T)}$

$\{\Omega^{(\tau)}\}$

Progeny allocation policy (Fig. 1b)

Learning **policy** by **Black-box optimization**

Genetic gain in final generation

$\{h^{(\tau)}\}$

Propose new $\{h^{(\tau)}\}_{\tau=0,...,T-1}$

StoSOO

$g^{(T)}$

Information on initial population

- Marker genotype
- Marker effects
- Recombination rate

20,000 function evaluations

Use optimized $\{\hat{h}^{(\tau)}\}_{\tau=0,1,2,3}$ **for allocation of progenies**

$t = T$

breeder

$\hat{b}^{(0)}$ $\hat{b}^{(1)}$ $\hat{b}^{(2)}$ $\hat{b}^{(3)}$

$t = 0$ $t = 1$ $t = 2$ $t = 3$ $t = 4$

Real breeding programs (Evaluation by $10,000$ simulations)

B

- Breeding values (**BV**)
- Weighted breeding values (**WBV**)
- Genetic variance of progenies (**GVP**)

Comparison of 6 optimized strategies:
Elements consisting of $\Omega^{(\tau)}$ are
1. **BV**   4. BV + GVP (**BVGVP**)
2. **WBV**   5. WBV + GVP (**WBVGVP**)
3. BV + WBV (**TBV**)   6. BV + WBV + GVP (**TBVGVP**)

Q **selection indices**

Pair	WBV	GVP	etc...
P001	1.09	1.01	...
P002	1.91	0.73	...
...	...	$\Omega^{(\tau)}$	...
P119	0.78	1.91	...
P120	1.27	2.35	...

$N^{(pair)(\tau)}$ pairs

Softmax

Pair	parent1	parent2	# of progenies
P001	L001	L001	**?**
P002	L001	L014	**?**
...	...	...	$b^{(\tau)}$
P119	L231	L246	**?**
P120	L246	L246	**?**

$$b^{(\tau)} = \left[N^{(pop)(\tau+1)} \times \frac{\exp(\Omega^{(\tau)} h^{(\tau)})}{\sum_Q \exp(\Omega^{(\tau)} h^{(\tau)})} \right]$$

Optimize $\{h^{(\tau)}\}_{\tau=0,...,T-1}$ **to maximize** $g^{(T)}$ **(Fig. 1a)**

$N^{(pop)(\tau+1)}$: population size τ: current generation in simulation