Deciphering sources of PET signals in the tumor microenvironment

of glioblastoma at cellular resolution

Laura M Bartos[1], Sabrina V Kirchleitner[2], Zeynep Ilgin Kolabas[3,4,5], Stefanie Quach[2], Jens
Blobner[2], Stephan A Mueller[6,7], Selin Ulukaya[3,8], Luciano Hoeher[3], Izabela Horvath[3,9], Karin
Wind-Mark[1], Adrien Holzgreve[1], Viktoria C Ruf[10], Lukas Gold[1], Lea H Kunze[1], Sebastian T
Kunte[1], Philipp Beumers[1], Melissa Antons[1], Artem Zatcepin[1,7], Nils Briel[7,10], Leonie
Hoermann[1], Denise Messerer[11], Peter Bartenstein[1,12], Markus J Riemenschneider[13], Simon
Lindner[1,12], Sibylle Ziegler[1], Jochen Herms[7,10,12], Stefan F Lichtenthaler[6,7,12], Ali Ertürk[3,4,5,12],
Joerg C Tonn[2,14], Louisa von Baumgarten[2,14], Nathalie L Albert[1,14,15*], Matthias Brendel[1,7,12*]

[1]Dept. of Nuclear Medicine, University Hospital of Munich, LMU Munich, Munich, Germany
[2]Dept. of Neurosurgery, University Hospital of Munich, LMU Munich, Munich, Germany
[3]Institute for Tissue Engineering and Regenerative Medicine (iTERM), Helmholtz Center, Neuherberg,
Munich, Germany
[4]Institute for Stroke and Dementia Research (ISD), University Hospital of Munich, Ludwig-Maximilians
University Munich, Munich, Germany
[5]Graduate School of Systemic Neurosciences (GSN), Munich, Germany
[6]Neuroproteomics, School of Medicine, Klinikum rechts der Isar, Technische Universität München,
Munich, Germany
[7]DZNE – German Center for Neurodegenerative Diseases, Munich, Germany
[8]Faculty of Biology, Master of Science Program in Molecular and Cellular Biology, Ludwig-
Maximilians-Universität München, Planegg, Germany
[9]School of Computation, Information and Technology (CIT), TUM, Boltzmannstr. 3, 85748 Garching,
Germany
[10]Center for Neuropathology and Prion Research, LMU Munich, Munich, Germany.
[11]Dept. of Cardiology, University Hospital of Munich, LMU Munich, Munich, Germany
[12]Munich Cluster for Systems Neurology (SyNergy), Munich, Germany
[13]Dept. of Neuropathology, Regensburg University Hospital, Regensburg, Germany
[14]German Cancer Consortium (DKTK), Partner Site Munich, German Cancer Research Center
(DKFZ), Heidelberg, Germany
[15]Bavarian Cancer Research Center (BZKF), Erlangen, Germany.

January 26, 2022

Running title: Cellular origin of PET biomarkers
Word count: 4510
Figures: 6
Tables: 1
Keywords: PET, tumor microenvironment, cell sorting, cellular source, immune biomarker,
glioblastoma

*contributed equally as last authors

First author:
Laura Maria Bartos
Department of Nuclear Medicine

University Hospital of Munich, Marchioninistr.15, 81377 Munich, Germany
Phone: +49 (0) 89 4400 74646
Fax: +49 (0) 89 4400 77646
E-Mail: laura.bartos@med.uni-muenchen.de

52
Corresponding author:
Dr. Matthias Brendel, MHBA
Department of Nuclear Medicine
University Hospital of Munich, Marchioninistr.15, 81377 Munich, Germany
Phone: +49 (0) 89 4400 74650
Fax: +49 (0) 89 4400 77646
E-Mail: matthias.brendel@med.uni-muenchen.de

Abstract

Various cellular sources hamper interpretation of positron-emission-tomography

(PET) biomarkers in the tumor microenvironment (TME). We developed

immunomagnetic cell sorting after *in vivo* radiotracer injection (scRadiotracing) in

combination with 3D-histology via tissue clearing to dissect the cellular allocation of

PET signals in the TME. In SB28 glioblastoma mice, translocator protein (TSPO)

radiotracer uptake per tumor cell was higher compared to tumor-associated

microglia/macrophages (TAMs). Cellular radiotracer uptake was validated by

proteomics and confirmed for *in vitro* samples of patients with glioblastoma. Regional

agreement between PET signals and single cell tracer uptake predicted the individual

cell distribution in 3D-histology. In consideration of cellular tracer uptake and cell type

abundance, tumor cells were the main contributor to TSPO enrichment in

glioblastoma, however proteomics identified potential PET targets highly specific for

TAMs. Combining cellular tracer uptake measures with 3D-histology facilitates precise

allocation of complex PET signal sources and will serve to validate novel TAM-specific

radioligands.

Introduction

The tumor microenvironment (TME) emerged as an acknowledged key component in cancer biology and treatment[1]. In particular, novel immunotherapies against targets of the TME facilitate potentiation of host antitumor immune responses[2]. However, despite the need to monitor novel anti-cancer therapeutics *in vivo*, specific biomarkers for immune cells of the TME are still limited[3]. Recent efforts originated in radiolabeled antibodies for positron-emission-tomography (PET) that specifically target immune cells[4], but the cell-type heterogeneity of the TME hampers cellular allocation and interpretation of such biomarkers. Furthermore, antibody-based PET biomarkers have limited penetrance through the blood-brain-barrier, which limits their use in brain malignancies, such as glioblastoma which represents the most common malignant primary brain tumor with a very poor prognosis[5]. In this disease, 18 kDa translocator protein (TSPO)-PET showed opportunities to assess myeloid cells of the TME as important contributors to immune evasion of glioblastoma[6]. However, the detailed sources of TSPO and other TME biomarkers in glioma have not yet been thoroughly elucidated, which again hampers interpretation and understanding of individual PET images. In glioma, tumor-associated microglia and macrophages (TAMs) show elevated expression of TSPO *in vitro*[7] and *in vivo*[6], acting as a potential signal target of TSPO ligands. However, this finding is a matter of debate since in high-grade tumors, TSPO was found to be highly overexpressed by tumor cells as well[8,9,10].

To overcome the limitations of previous immunohistochemistry to PET comparisons, we aimed to resolve the complex sources of TME biomarkers in glioma at cellular resolution. We therefore established and validated a novel approach of immunomagnetic cell sorting after radiotracer injection (single cell Radiotracing, scRadiotracing)[11] and exploited this technique in a mouse model of experimental glioblastoma. TSPO tracer uptake was measured in single tumor cells and CD11b-positive immune cells of a SB28 glioblastoma mouse model, validated against TSPO protein levels as determined by proteomics as a proof of concept. Furthermore, we investigated associations between single cell TSPO tracer uptake and tumor heterogeneity in TSPO-PET. To bridge the gap between implantation of murine tumors

and spontaneous glioma development in human brain, we transferred TSPO scRadiotracing

to human samples and determined proportions of TSPO tracer uptake *in vitro* in tumor and

immune cells of patients with high-grade and low-grade glioma immediately after surgery. As

a second major innovation, 3D-histology via tissue clearing and light sheet microscopy was

integrated to acknowledge PET signals as a composite of cell type abundance and single cell

tracer uptake. Finally, the full proteome of isolated SB28 tumor cells and TAMs was analyzed

for identification of potential TAM-specific TME radioligand targets with superior binding

capacity when compared to TSPO.

Results

Single cell tracer uptake measures of microglia as calculated by scRadiotracing

correspond to PET signal changes upon microglia depletion

Immune PET biomarkers such as TSPO have the potential to monitor TAMs of the TME, but

the target signal of currently available TSPO radioligands is challenged by several cellular

sources. To validate scRadiotracing for the TSPO tracer [^{18}F]GE-180, we compared the

magnitude of the microglia-specific TSPO-PET signal with the recovered radioactivity in

isolated microglia of untreated mice. First, we investigated the allocation of the TSPO tracer

in the healthy mouse brain. Isolated CD11b(+) microglia (1.2 ± 0.1 x 10^5 cells) of untreated

mice revealed 7.7 ± 0.7 x 10^{-7} percentage radioactivity (normalized to injected dose (ID) and

body weight (BW), %ID*BW) per single cell, whereas enriched ACSA2(+) astrocytes (5.2 ±

0.7 x 10^5 cells) indicated 12.5-fold lower radioactivity per single cell (6.2 ± 0.7 x 10^{-8}

%ID*BW; p = 0.00016; **Fig. 1A**), speaking for strong specificity of the tracer for microglia

over astrocytes. Next, extrapolation by published microglia cell numbers (6% of all brain

cells, 7.63 x 10^6 microglia cells) from dedicated studies[12], whole brain radioactivity (0.19 -

0.30 MBq) and %brain dose per single microglia cell (1.8 x 10^{-6} - 3.6 x 10^{-6}) were used to

calculate the contribution of the microglia population to the radioactivity in the whole brain to

be 17.5% ± 2.2%.

To quantify the contribution of microglia to the overall TSPO-PET signal, we used CSF1R

inhibition via PLX5622 and hereby depleted 96% of all microglia in the mouse brain (**Fig.**

1B). As a result of microglia depletion, the TSPO-PET signal (**Fig. 1C**) in mice at 10.1 ± 2.1

months of age[13,14] showed 18.5% lower standardized uptake values (SUV_H) when compared

to vehicle controls (0.19 ± 0.01 vs. 0.23 ± 0.01; $p < 0.0001$). This decrease corresponded to

the contribution of the microglia cell population extrapolated after scRadiotracing ($p = 0.770$;

Fig. 1D). Thus, TSPO scRadiotracing provided a nearly complete recovery of the microglia-

specific PET signal in healthy mice and suggested capability of the method to be used for

tracer allocation in cells of the TME.

TSPO tracer uptake of tumor cells and TAMs reflects cellular TSPO protein levels in

the SB28 glioblastoma mouse model

In the next step, we applied scRadiotracing to an experimental orthotopic glioblastoma model

(SB28) that closely reflects the human TME[15] to investigate the contribution of tumor cells

and TAMs to the TSPO-PET signal (**Fig. 2A-C**). Using MACS, CD11b(+) immune cells were

isolated from tumor and sham injected brains (TAMs or sham microglia), tumor samples

underwent subsequent tumor cell isolation. Residual cell pellets were analyzed as depleted

fractions. We extracted 1.4 ± 0.4 x 10^5 GFP(+) tumor cells and 6.7 ± 1.7 x 10^5 CD11b(+)

TAMs from tumor mice and 4.2 ± 0.8 x 10^4 CD11b(+) microglia from sham mice (**Fig. 2A**) in

enriched fractions at high purity as determined by flow cytometry (tumor cells: 87% ± 2%;

TAMs: 90% ± 1%; sham microglia: 89% ± 1%; **Fig. 2B-D**). Signal-to-noise-ratios (SNRs) for

radioactivity detection were consistently ≥ 2 (**Fig. 2A**).

Subsequently, single cell tracer uptake of tumor cells and TAMs was quantified to decipher

cellular contributions to the TSPO tracer uptake in glioblastoma. Both, single SB28 tumor

cells (3.53-fold, 1.7 ± 0.2 x 10^{-6} %ID*BW; $p < 0.0001$, 1-way ANOVA; **Fig. 2D**) and TAMs

(2.58-fold, 1.3 ± 0.2 x 10^{-6} %ID*BW; $p = 0.024$, 1-way ANOVA; **Fig. 2D**) indicated higher

TSPO tracer uptake when compared to sham microglia (4.9 ± 0.8 x 10^{-7} %ID*BW). We also

tested if sham injections had an impact on TSPO tracer uptake of microglia and did not find

any remarkable difference in these cells when compared to control microglia of untreated

age-matched mice (p = 0.999, 1-way ANOVA, **Fig. 2D**).

In the direct comparison of tumor cells and TAMs, we noted 1.37-fold higher TSPO tracer

uptake of single SB28 tumor cells when compared to TAMs (p = 0.0010, paired t-test),

indicating a dual cellular allocation of the tracer with even slightly higher single cell uptake of

SB28 tumor cells when compared to the immune cell target population. Importantly, and

highly congruent to scRadiotracing, proteomics revealed increased TSPO protein levels in

SB28 tumor cells (2.98-fold, FDR-corrected p = 0.015) and TAMs (2.04-fold, p = 0.0016)

when compared to control microglia. Although not reaching significance after correction for

multiple comparisons, higher TSPO tracer uptake of SB28 tumor cells over TAMs was also

reflected by higher TSPO protein abundance (1.46-fold, p = 0.357; **Extended Fig. 1**). We

additionally proved the presence of the tracer target in SB28 tumor and immune cells by

revealing that the vast majority of isolated CD11b(+) and GFP(+) cells showed TSPO co-

labeling in flow cytometry (**Fig. 2E,F**), which is in accordance to the previously reported co-

localization of TSPO and GFP/CD11b in immunohistochemistry[16].

To determine the contributions of the distinct cell types with TSPO expression to the

radioactivity in the cell pellets, we calculated a linear regression with normalized radioactivity

per cell count as dependent variable and cellular proportions (GFP(+) tumor cells, CD11b(+)

TAMs and ACSA2(+) astrocytes) of all enriched and depleted fractions as predictors. The

regression model significantly explained the variance in single cell TSPO tracer uptake (F =

6.2, p = 0.0011, R^2 = 0.25, R^2_{adj} = 0.21). Tumor cells showed highest contribution to the

single cell TSPO tracer uptake (β = 0.76 p = 0.00017), followed by TAMs (β = 0.57, p =

0.0032) and astrocytes (β = 0.34, p = 0.0074) (**Fig. 2G**). A global model using measured

radioactivity, total cell count and cell proportions indicated that 76% of the variance in

radioactivity measures could be explained by the cellular component of tumor cells, TAMs

and astrocytes (F = 46.6, p < 0.0001, R^2 = 0.772, R^2_{adj} = 0.756). To question the performed

calculations in depth, we correlated the radioactivity measured in the depleted tumor

fractions, containing 39% ± 2% GFP(+), 7% ± 1% CD11b(+) and 16% ± 3% ACSA2(+) cells,

with estimated summed up radioactivity by single cell tracer uptake and cell count (R = 0.74,

p = 0.0002).

In summary, the established methodology accurately determines radiotracer uptake of single

cells, which closely reflects relative protein abundance in respective cell types. The cellular

allocation of TSPO as a biomarker consist of overexpression in both tumor and immune cells

in SB28 tumors.

Single cell tracer uptake explains inter-individual PET heterogeneity and reveals

dominant association of tumor cells with TSPO-PET signals

To bridge the gap between cellular tracer uptake and tumor imaging, PET was added to the

experimental workflow of scRadiotracing in SB28 glioblastoma mice (**Fig. 3A**). TSPO-PET

indicated heterogeneity of tumor signals 2.5 weeks after surgery (range of SUV_{mean}: 0.937 −

1.825), which were strongly elevated when compared to sham injection (SUV_{mean}: 0.842 ±

0.032; p < 0.0001; **Fig. 3B**). We correlated the individual single cell TSPO tracer uptake of

SB28 tumor cells and TAMs with the TSPO-PET signal in the lesion and observed similar

degrees of strong association for both cell types (**Fig. 3C**). Interestingly, we also found an

inter-correlation between single cell tracer uptake of tumor cells and TAMs (R = 0.758, p =

0.0011), which indicated bidirectional dependence of TSPO enrichment in tumor and immune

cells (**Fig. 3D**). A combined vector of single cell TSPO tracer uptake of both tumor cells and

TAMs strongly correlated with the magnitude of the tumor TSPO-PET signal (R = 0.848, p <

0.0001; **Fig. 3E**). A regression model of TSPO tracer uptake of tumor cells and TAMs

however demonstrated that only tumor cells contributed significantly to the TSPO-PET signal

(tumor cells: β = 0.55, p = 0.038; TAMs: β = 0.35, p = 0.157).

Going into more detail, we interrogated the regional heterogeneity of tumor TSPO-PET

signals and correlated single cell TSPO enrichment with the regional PET signal magnitude.

Therefore, we performed a cluster-based analysis (k-means clustering, 50 clusters defining

intratumoral regions of increasing signal intensity) of TSPO-PET in the tumor lesion (**Fig. 3F**)

and correlated single cell tracer uptake with the PET signal in respective clusters. The

association between single cell TSPO enrichment and TSPO-PET signal intensity in the

cluster followed an inverted U-shape function for tumor cells and TAMs, with higher

agreement between tumor cell TSPO and regional TSPO-PET signal when compared to

TAM TSPO and regional TSPO-PET signal (**Fig. 3F-H**). Strikingly, the association between

TSPO-PET and single cell tracer uptake was increasingly dominated by tumor cell TSPO

enrichment as a linear function of cluster hierarchy (y = 0.004x-0.026; R = 0.947; p < 0.0001;

Fig. 3F-H). This indicates that especially regions with high TSPO-PET signal intensities were

characterized by strong dependence from single tumor cell TSPO. Thus, the combination of

PET and scRadiotracing identified a strong association between PET signals and cellular

tracer uptake, which was characterized by a dominant dependency of TSPO-PET signals

from tumor cells when compared to immune cells.

***In vitro* scRadiotracing in human glioma underlines translational value of the**

methodology and confirms higher TSPO tracer uptake of tumor cells when compared

to TAMs

To test for a potential translational value of single cell tracer uptake measures in human

glioblastoma, we applied *in vitro* scRadiotracing in a cohort of patients with high-grade and

low-grade glioma that underwent biopsy or tumor resection (**Table 1**, **Extended Table 1**).

Tissue samples were investigated immediately after surgery applying *in vitro* [^{18}F]GE-180

incubation of individual single cell suspensions and subsequent cell sorting (**Fig. 4A**). The

rationale was to check for discrepancies of cellular tracer binding between species.

scRadiotracing resulted in enriched fractions with visually well discernible populations of

tumor cells and TAMs (**Extended Fig. 2**). High-grade glioma indicated similar heterogeneity

of tumor-cell-to-TAM ratios in the single cell suspension when compared to low-grade glioma

(**Fig. 4B**). As the purity of tumor and TAM enriched fractions was limited in human samples,

negligibility of TSPO tracer uptake of non-tumor/non-TAM cells was proven by showing that

only the number of isolated TAMs correlated with measured tracer signal in the TAM

enriched cell pellet (β = 0.875, p = 0.0021), whereas non-tumor/non-TAM cells did not

contribute to the overall magnitude of radioactivity in the same sample ($\beta = 0.118$, $p = 0.542$;

Fig. 4C). Single cell TSPO enrichment in TAMs was similar between patients with high-grade

and low-grade glioma ($1.6 \pm 0.9 \times 10^{-7}$ %ID vs. $2.3 \pm 1.0 \times 10^{-7}$ %ID, $p = 0.618$), whereas

TSPO enrichment of tumor cells was higher in patients with high-grade glioma ($4.5 \pm 1.3 \times$

10^{-7} %ID) when compared to patients with low-grade glioma ($1.4 \pm 0.5 \times 10^{-7}$ %ID, $p = 0.047$,

Fig. 4D). Within the group of patients with high-grade glioma, TSPO enrichment of tumor

cells was stronger when compared to TAMs ($p = 0.0014$; **Fig. 4D**). Contrary, TSPO

enrichment of tumor cells showed no significant difference when compared to TAMs within

the group of patients with low-grade glioma. We questioned if previous therapy constitutes a

potential confounder of single cell tracer uptake, but we did not find any significant impact of

a previous therapy (radiation and/or chemotherapy, 11 out of 20) or resection (9 out of 20) on

TSPO enrichment of single tumor cells or TAMs (all $p > 0.05$).

In accordance with the combination of PET and scRadiotracing in the SB28 mouse model,

we correlated single cell TSPO enrichment with TSPO-PET imaging, which was performed in

a subset of patients prior to surgery. TSPO enrichment of tumor cells ($R = 0.934$, $p < 0.0001$)

but not TSPO enrichment of TAMs ($R = 0.295$, $p = 0.328$) was associated with the lesion

signal in TSPO-PET (**Fig. 4E**). Individual patients with varying TSPO-PET signal intensities

but similar FET and (ce)MRI findings showed a strong agreement between lesion signals and

corresponding tumor cell TSPO enrichment (**Fig. 4F**). Taken together, the dominance of

tumor cell radiotracer uptake and the association between scRadiotracing and PET in human

tissue samples closely reflected our findings in SB28 mice. Therefore, *in vitro* scRadiotracing

provides an opportunity to efficiently check for consistency in cellular target binding between

murine and human samples.

The triangle of PET, scRadiotracing and 3D-histology dissects cellular sources of

regional PET signals and pinpoints dominant contribution of tumor cells to the TSPO-

PET signal

Since PET tracer signals are a product of cellular tracer uptake and regional cell type

abundance, we used 3D-histology via tissue clearing and light sheet microscopy (**Fig. 5A**) to

determine absolute and relative cell numbers of GFP(+) tumor cells and CD11b(+) immune

cells throughout entire SB28 tumors in the intact ipsilateral hemispheres. GFP(+) volumes

were 5.5-fold larger when compared to CD11b(+) volumes (18.6 ± 6.8 mm³ vs. 3.4 ± 1.5

mm³, p = 0.037; **Fig. 5B,C**). Confocal microscopy revealed a GFP(+) volume of 4493 µm³ per

SB28 tumor cell and a CD11b(+) volume of 2713 µm³ per TAM (**Fig. 5D**), resulting in higher

cell count of SB28 tumor cells when compared to TAMs in individual SB28 lesions (**Fig. 5E**).

Considering both single cell tracer uptake and cell distributions of tumor cells and TAMs, we

calculated a tumor-cell-to-TAM contribution to the TSPO-PET signals in SB28 lesions of

3.5:1 (**Fig. 5E**).

Next, PET, scRadiotracing and 3D-histology were used as an integrated concept.

Importantly, the regional resolution of 3D-histology could be predicted by combining lesion

PET signals with single cell tracer values (**Fig. 5F,G**, **Extended Fig. 3**). For this purpose, we

used the regional association between single cell tracer uptake of SB28 tumor cells and

TAMs with TSPO-PET (i.e. correlation coefficients per cluster**, Fig. 3F**) as a cluster-based

PET weighting factor. k-means cluster PET signals of all mice that received TSPO-PET

imaging and 3D-histology were multiplied with their respective correlation coefficients and

divided by single cell tracer uptake of both tumor cells and TAMs (**Fig. 2D**) to predict

individual regional abundance (i.e. light sheet signal intensity) for each cell type (**Fig. 5F-H**).

Predicted and standard of truth 3D-histology were strongly correlated (tumor cells: R = 0.624,

p < 0.0001; TAMs: R = 0.304, p < 0.0001; **Fig. 5H; Extended Fig. 3**). Hot spots of GFP-

positivity were co-localized with regions of high TSPO-PET signal intensity (**Fig. 5G**), fitting

to the strong association between tumor cell TSPO and regional TSPO-PET (**Fig. 3F**). The

radioactivity in individual tumors as measured by PET was precisely predicted by aggregated

single cell tracer uptake and cell count of tumor cells and TAMs (**Fig. 5I**). In summary,

exploiting the established methodological combination of PET, scRadiotracing and 3D-

histology enables to pinpoint complex PET signals to their detailed cellular origins.

Proteomics identifies potential TAM-specific radiotracer targets with superior binding capacity when compared to TSPO

Since tumor cells dominate the TSPO-PET signal, the development of novel PET

radiotracers with higher specificity for TAMs could provide added value to glioblastoma

diagnostics. We therefore generated murine proteome data to identify potential TAM-specific

radiotracer targets. Starting from a total of 7869 proteins relatively quantified between TAMs

and microglia of healthy control mice, 1097 showed higher fold-changes than TSPO (**Fig.**

6A,B). 165 of these proteins could not be quantified in SB28 tumor cells and additional 16

proteins showed >10-fold higher levels in TAMs when compared to SB28 tumor cells (**Fig.**

6B), thus providing sufficient specificity for TAM over tumor cell expression.

To translate these findings from mouse to human, we used the Human Protein Atlas[17] to

identify expression of respective human gene homologs in off-target cells in brain as an

index of TAM-specificity (see first set of columns in **Fig. 6C, Extended Fig. 4**). Physiological

expression in resident and invading cells of the tumor microenvironment (see second set of

columns in **Fig. 6C, Extended Fig. 4**) was characterized. In a next step, we ensured low

target expression in other cells in the organism to prevent peripheral radiotracer absorbance,

since this may cause a sink effect and high radiation exposure (see third set of columns in

Fig. 6C, Extended Fig. 4). This selection process resulted in 20 proteins with highly suitable

characteristics as specific immune PET tracer targets (**Fig. 6D**), which were ultimately

categorized into their immune cell expression cluster (**Fig. 6D**). *CYP4F3*, *H2AB1*, *PGR3*,

FCGR3B and *PHF24* were allocated to the neutrophil expression cluster. *ALOX15 and*

RNASE3 were associated with the eosinophil expression cluster and *PGR2* was localized in

the basophil expression cluster, while both expression clusters were shared for *RAB44* and

RNASE2. An association with the monocyte expression cluster was identified for *CLEC6A*,

MPO, *MYO1G* and *SIGLEC1*. Few identified genes belonged to B-cell (*ATP6V1G2* and

SEPTIN1) and T-cell (*STAP1*) expression clusters and group enrichment was found for

LRRC25, *LY9* and *PADI4* (**Fig. 6D**). High protein abundance, which allows sufficient PET

signal-to-noise-ratios, was observed for *ALOX15, PRG2, SIGLEC1, MYO1G* and *MPO*

(Label-free quantification (LFQ) intensity > 18, **Extended Fig. 5**). Radiolabeled ligands with

binding affinity to these proteins may re-enter the workflow of scRadiotracing, 3D-histology

and human translation to test for their cellular specificity *in vivo*.

Discussion

We present the first study that determines *in vivo* radiotracer uptake in the complex TME of

glioblastoma at cellular resolution. After validation of scRadiotracing for the immune target

TSPO, we deciphered the cellular sources of TSPO-PET signals in experimental

glioblastoma and patients with glioma. The established methodology accurately determined

higher radiotracer uptake of single glioblastoma tumor cells when compared to immune cells,

which closely reflected protein levels of respective cell types. The combination of PET and

single cell tracer uptake served to predict 3D-histology. Ultimately, a detailed proteome

analysis facilitated identification of TAM-specific radiotracer targets beyond TSPO.

As a prerequisite of the main analysis, a methodological validation of TSPO radiotracer

measures in single cells had to be addressed. We applied a microglia depletion paradigm[18]

to compare the signal reduction observed by *in vivo* [18F]GE-180 TSPO-PET of mice after

CSF1R inhibition with the total radioactivity attributable to the microglia cell population after

scRadiotracing. Here, we confirmed that scRadiotracing was able to closely recover the

tracer signal reduction observed in PET (**Fig. 1**). By CSF1R inhibition with PLX5622,

microglia cells can be depleted up to 99%, which may be maintained under continuous

treatment for up to 2 months[19], but CSF1R inhibition does not change the TSPO gene

expression of other CNS cells[20]. Thus, we claim that the reduction of TSPO-PET derived

from the nearly complete depletion of microglia after 7-weeks PLX5622 treatment.

Furthermore, we observed 12-fold higher single cell TSPO tracer uptake of microglia when

compared to astrocytes, pinpointing specificity of [18F]GE-180 to microglia in untreated mice.

Additionally, the high congruence between tracer signal in the microglia population and PET

signal reductions after microglia depletion fits to previous investigations that outlined a high

correlation between [^{18}F]GE-180 TSPO-PET and the microglia markers Iba1 and CD68 in

immunohistochemistry[21].

We were able to further validate scRadiotracing in tumor cells and TAMs of the experimental

SB28 glioblastoma model by quantification of relative cellular protein levels of TSPO in

respective cell populations using proteomics. The application of proteomics was particularly

important since transcriptomics does not necessarily reflect the amount of protein that can be

targeted by the radiotracer[22]. We also found that the inter-individual heterogeneity of SB28

and human tumor PET signals was correlated with the magnitude of tracer uptake per each

single cell, which implies that the PET signal is mainly driven by cellular uptake and not by

confounding alterations of the blood-brain-barrier[23] or off target binding[24]. Our observations

of heterogeneous single cell tracer uptake fit to the biological variability of TSPO which has

not only been outlined in glioblastoma[7,25] but also in neurodegenerative diseases[26,27].

Ultimately, the combination of PET and scRadiotracing with 3D-histology confirmed the

methodological capability to quantify cellular sources of *in vivo* PET signals. In this regard,

PET tracer signals are not only a function of cellular uptake but a product of cell type

abundance and single cell tracer uptake, making the novel combination of scRadiotracing

and 3-dimensional assessment of specific cell counts[28] a major achievement of the current

work. Furthermore, scRadiotracing allowed to predict 3D-histology at the individual level,

which may have clinical impact in patients with glioma, where histology can only be acquired

in subsamples of the highly heterogeneous tumor mass[29,30].

In this study, we used TSPO as an established immune PET biomarker in neurological and

neuro-oncological diseases with controversially discussed cellular signal sources. Although

several correlative investigations questioned the cellular sources of TSPO tracer signals in

glioma[6,31], direct measures of radiotracer allocation have not yet been performed and

contributions of tumor and immune cells to the PET signal remained unclear. Our data

indicated higher enrichment of [^{18}F]GE-180 in SB28 tumor cells when compared to TAMs

after *in vivo* tracer administration and similar results were obtained in human high-grade

glioma cells when compared to TAMs of the same tumor after *in vitro* tracer incubation.

These findings were supported by stronger agreement of tumor TSPO-PET signals with

individual tumor cell tracer uptake in mice and humans. 3D-histology in SB28 mice moreover

showed higher abundance of tumor cells when compared to TAMs, pinpointing the

dominance of tumor cells to SB28 tumor TSPO-PET signals. Still, potential discrepancies

between murine and human glioblastoma (e.g. inoculation versus spontaneous tumor

growth) need to be considered. Although the SB28 model is reported to closely mimic the

human TME[15] and our translational data indicate similar results regarding single cell TSPO

enrichment across species, the small sample size of the human analysis in respect of large

heterogeneity in diagnoses and previous therapy strategies remains to be acknowledged.

Further investigations in pre-defined subgroups with larger sample sizes may aid to

disentangle and differentiate therapy-related effects on TSPO expression in cells of the

human TME, i.e. after radiation therapy[32].

A high contribution of tumor cells to the TSPO-PET signal of glioblastoma is in agreement

with preclinical investigations that used TSPO-PET and autoradiography after implantation of

GL261 cells into a full TSPO knock-out [31,33]. However, several preclinical studies have

revealed a distinct co-localization of TSPO and Iba1/CD68 in immunohistochemistry in

different glioblastoma mouse models and concluded that TAMs contribute significantly to the

TSPO-PET signal[7-10]. While some of these studies have already drawn a qualitative

comparison between immunohistochemistry and TSPO-PET quantification by various

ligands[8-10], our study provides first direct evidence for simultaneous allocation of a TSPO-

PET tracer in both cell types. Future research should also compare single cell TSPO

enrichment across the broad spectrum of different glioblastoma models (i.e. GL261) and

primary human glioblastoma cell lines.

Various signal sources with a significant impact on the TSPO-PET signal in the TME call for

detailed interrogation of TSPO as a glioblastoma imaging biomarker. First, lacking specificity

of TSPO-PET signals for TAMs in glioblastoma hampers the biomarker's application as a

mere immune index. However, since TSPO-PET signals allow prognostication of overall

survival[34], TSPO-PET still adds valuable biomarker features to the toolbox of glioblastoma

imaging assessments[35]. These quantitative TSPO-PET data even exceeded the prognostic

value of gene expression levels[36], which could be related to the direct quantification of TSPO

receptor abundance instead of indirect mRNA measures. The pilot dataset of *in vitro*

scRadiotracing in patients with glioma already showed feasibility even with small amounts of

tissue obtained from biopsy. Thus, the current results should encourage to investigate the

separate prognostic value of single cell TSPO enrichment in tumor and immune cells since

such data would potentially result in different treatment strategies. Importantly, distinct

binding of radioligands to different cells of the TME will also impact personalized medicine,

including direct targeting of various tumors with radiotheranostics.

Beyond TSPO, TAM-specific radioligands for assessment of immune cells in the TME could

facilitate monitoring of novel immunotherapies[37]. We used our murine proteomics dataset to

classify TSPO within the spectrum of potential radiotracer targets. Since TSPO was only

moderately elevated in TAMs and surpassed by the TSPO expression in tumor cells, there is

a major need for further improvement, identification and development of TAM-specific

radiotracers. The identified targets need to be further evaluated in terms of ligand

accessibility and stable target expression and screened for available crystal structures or

known binders that aid generation of lead molecules for radiotracer development.

Importantly, several of the identified proteins matched with signature genes of

immunosuppressive (i.e. *ALOX15*[38]) TAM phenotypes or predicted therapy response upon

checkpoint inhibition (i.e. *LY9*[39] or *MPO*[40]), thus indicating their potential as biomarkers of

immunomodulation in cancer. scRadiotracing will provide the opportunity for a precise

determination of the *in vivo* specificity of potential radioligands for these targets in their

developmental process despite the complexity of the TME.

In a nutshell, the established methodology yields robust and reliable quantification of cellular

radiotracer uptake in mice and patients and will serve to disentangle cellular sources of PET

signals of established or newly developed tracers. This is of particular importance since

many potential radiotracer targets are shared between tumor cells, TME cells and resident

cells of the brain[41].

Tables

Table 1

ID	Age (y)	Sex	Dx	Disease duration (m)	WHO grade	rs6971	IDH	Sample acquisition	Previous therapy	[^{18}F]GE-180 TSPO-PET
#1	36.0	m	Astrocytoma	5.6	2	HAB	+	Resection	-	+
#2	59.0	f	Oligodendroglioma	13.6	2	MAB	+	Resection	+	+
#3	50.6	f	Glioblastoma	0.9	4	MAB	-	Resection	+	+
#4	46.9	f	Oligodendroglioma	0.3	2	HAB	+	Resection	-	+
#5	75.7	f	Glioblastoma	0.2	4	MAB	-	Biopsy	-	+
#6	69.1	f	Glioblastoma	1.0	4	LAB	-	Resection	+	-
#7	64.3	f	Oligodendroglioma	17.8	2	-	+	Biopsy	+	-
#8	59.3	m	Astrocytoma	2.6	2	HAB	+	Biopsy	+	+
#9	36.4	f	Oligodendroglioma	1.1	2	HAB	+	Biopsy	-	+
#10	44.0	m	Oligodendroglioma	12.3	3	MAB	+	Biopsy	+	+
#11	72.3	f	Glioblastoma	0.1	4	-	-	Resection	-	-
#12	39.7	f	Astrocytoma	4.5	2	-	+	Biopsy	-	-
#13	27.2	f	Astrocytoma	0.7	2	-	+	Biopsy	-	-
#14	68.7	m	Glioblastoma	0.1	4	LAB	-	Resection	-	+
#15	66.8	f	Oligodendroglioma	25.9	2	-	+	Resection	+	-
#16	58.5	m	Astrocytoma	20.8	3	-	+	Biopsy	+	+
#17	52.8	f	Oligodendroglioma	20.9	3	HAB	+	Biopsy	+	+
#18	49.6	m	Oligodendroglioma	1.2	3	HAB	+	Biopsy	+	+
#19	57.9	m	Astrocytoma	12.9	3	MAB	+	Biopsy	+	-
#20	72.0	m	Glioblastoma	1.4	4	HAB	-	Resection	-	+

452 **Table 1: Characteristics of the human glioma cohort**. rs6971 = TSPO polymorphism rs6971[42], LAB
453 = low-affinity binding status, MAB = medium-affinity binding status, HAB = high-affinity binding status,
454 Dx = Diagnosis, IDH = isocitrate dehydrogenase mutation, m = male, f = female, y = years. See
455 **Extended Table 1** for supplemental data.
456

 Figures and Figure Captions
 Figure 1

Figure 1: TSPO-PET signal reduction in the whole brain after microglia depletion corresponds to the whole brain signal attributable to tracer uptake in microglia.

(**A**) Components of scRadiotracing after *in vivo* tracer injection. Upon *in vivo* tracer injection and magnetic cell separation, cell pellets were analyzed by a high sensitive gamma counter to measure the radioactivity (Bq = Becquerel) in the sample (left panel, n=6, mean±SEM) as well as by flow cytometry to determine the cell count (middle panel, n=6, mean±SEM). After calculation of radioactivity per single cell (scTSPO, normalized to injected dose (ID) and body weight (BW)) in each sample, untreated mice showed significantly higher TSPO tracer uptake in microglia when compared to astrocytes (right panel; each bar represents a single animal; n=6, paired t-test).

(**C**) Representative sagittal Iba1 immunohistochemistry sections of VEH (n=3) and PLX5622 treated (n=3) mice and quantification of stained Iba1 area (%). Unpaired t-test, mean±SEM.

(**B**) Group average TSPO-PET images of healthy mice upon a MRI template after vehicle treatment (VEH, n=14) or CSF1R inhibition (PLX5622, microglia depletion, n=15) indicated a distinct signal reduction in the whole brain of PLX treated animals (-18.5%). SUV_H = heart normalized standardized uptake value (SUV). Unpaired t-test, mean±SEM.

(**D**) Extrapolation of the signal attributable to microglia using published cell numbers[43] and tracer uptake per single microglia cell as calculated by scRadiotracing. The calculated 17.5% contribution of microglia to the whole brain TSPO-PET signal corresponded to the observed TSPO-PET signal reduction after microglia depletion (18.5%). mean±SEM.

Figure 2

Figure 2: scRadiotracing in the SB28 glioblastoma mouse model allows differential assessment of TSPO tracer uptake in tumor cells and TAMs.

(**A**) Acquired components of scRadiotracing after *in vivo* tracer injection in SB28 glioblastoma (n=20) and sham (n=14) mice. Using immunomagnetic cell sorting, tumor cells (GFP(+)) and CD11b(+) TAMs from SB28 bearing mice and CD11b(+) microglia from sham injected mice were enriched, thus leaving the respective residual depleted cell fractions. Absolute cell numbers (left panel) and measured radioactivity (middle panel) resulted in satisfying signal to noise ratios (right panel) for all enriched and depleted fractions investigated. mean±SEM.

(**B**) Immunomagnetic cell sorting of tumor cells and TAMs led to >90% purity in enriched fractions as confirmed by flow cytometry.

(**C**) Distribution of tumor cells (GFP), TAMs (CD11b) and astrocytes (ACSA2) in enriched and depleted fractions of SB28 glioblastoma (n=20) and sham (n=14) mice. High purity was reached for tumor and TAM enriched fractions, while remaining depleted fractions still contained tumor cells but no TAMs. Astrocytes could be detected in both depleted fractions using ACSA2. mean±SEM.

(**D**) Comparison of the single cell TSPO tracer uptake of isolated tumor cells (n=20) and TAMs (n=20) in SB28 glioblastoma mice as well as microglia of sham (n=14) and untreated control (n=8) mice. Tumor cells and TAMs showed significantly higher TSPO tracer uptake per single cell (scTSPO, normalized to injected dose (ID) and body weight (BW)) when compared to isolated microglia of sham and untreated control animals. In SB28 mice, tumor cells showed higher single cell TSPO tracer uptake when compared to TAMs. Single cell tracer uptake of sham microglia showed no difference when compared to control microglia of untreated mice. Paired t-test for tumor cells vs. TAMs, one-way ANOVA for all other comparisons, mean±SEM.

(**E,F**) TSPO co-staining in flow-cytometry shows that nearly all tumor cells (GFP; **E**) and TAMs (CD11b; **F**) were also positive for TSPO. Notably, the minor population of CD11b(-) cells in the TAM enriched fraction did not show positivity for TSPO (**F**, lower left quadrant), confirming the specificity of the TSPO co-staining. Pooled data from n=3 tumors.

(**G**) Regression model including the enriched and depleted fractions of SB28 mice indicated highest contribution of tumor cells (left) to the radioactivity in the sample, followed by TAMs (middle) and astrocytes (right). Linear regression, β = standardized regression coefficient, n=60 samples, error bands represent 95% confidence interval.

Figure 3

Figure 3: Correlation of single cell TSPO tracer uptake of tumor cells and TAMs with TSPO-PET indicates association of TSPO-PET heterogeneity with single cell tracer enrichment.

(**A**) Schematic illustration of the scRadiotracing workflow in the SB28 glioblastoma mouse model including *in vivo* tracer application, PET imaging and cell sorting, resulting in radioactivity per single cell.

(**B**) TSPO-PET imaging indicated significantly higher lesion site signals in SB28 tumor mice (n=15) than in sham (n=14) animals at day 18 after inoculation. Unpaired t-test, mean±SEM.

(**C**) Using the established quantitative endpoint of radioactivity per single tumor cell and TAM, the correlation between scRadiotracing and TSPO-PET showed a strong dependence of PET signals from both tumor and TAM single cell TSPO tracer uptake (scTSPO, normalized to injected dose (ID) and body weight (BW)) in SB28 mice. N=15, R = Pearson's coefficient of correlation. Error bands represent 95% confidence interval.

(**D**) Inter-correlation of scTSPO of tumor cells and TAMs in the SB28 glioblastoma mouse model. N=15, R = Pearson's coefficient of correlation. Error band represents 95% confidence interval.

(**E**) Association between combined scTSPO and heterogeneous TSPO-PET signals. Black bars symbolize the individual TSPO-PET signal for all SB28 mice investigated (n=15). The curves represent scTSPO of tumor cells (orange) and TAMs (blue), both used to calculate a combined vector of cellular tracer uptake (yellow) for each individual animal. Increasing TSPO-PET signals were associated with increasing scTSPO, indicating a dependence of heterogeneous PET signals from single cell tracer uptake. Axial sections of TSPO-PET images upon the individual contrast enhanced computed tomography (ceCT) illustrate inter-individual TSPO-PET signal heterogeneity of SB28 tumors. R = Pearson's coefficient of correlation.

(**F**) Visualization of the regional k-means cluster analysis. A sphere was placed over the entire signal enhancement in TSPO-PET, which was followed by application of k-means clustering. This resulted in 50 volumes of interest (VOIs) defining 50 intratumoral regions of increasing signal intensity (images on the right).
The function between increasing cluster grade and scTSPO to PET correlations of tumor cells and TAMs followed an inverted U-shape (quadratic fit) for both cell types. The correlation coefficients (R) of TSPO-PET signals with scTSPO of tumor cells showed its peak in clusters of higher rank (cluster 27; orange) when compared to the respective cluster peak of TAMs (cluster 21; blue). Increasing cluster hierarchy (i.e. the TSPO-PET hotspot) was associated with predominant dependency of TSPO-PET signal intensity from scTSPO (linear fit of ΔR_{Tumor}-R_{TAM}; grey).

(**G,H**) Strong regional agreement between tumor and TAM cell density (GFP/ CD11b, light sheet microscopy) and regions with high scTSPO-to-PET associations. Coronal and sagittal slices show projections of correlation coefficients onto the k means cluster VOIs of an individual mouse. Notably, both analyses resulted in a characteristic spherical distribution of peak values that encircled the "cold" tumor core. R = Pearson's coefficient of correlation per voxel, arb. = arbitrary units.

Figure 4

Figure 4: Translation of scRadiotracing to human glioma samples confirms higher TSPO tracer uptake in tumor cells and stronger contribution of tumor cells to the TSPO-PET signal when compared to TAMs

(**A**) Schematic illustration of the *in vitro* scRadiotracing workflow in human glioma samples (n=20), leading to calculation of radioactivity per single tumor cell and TAM. Pseudocolor plots derived from flow cytometry show the applied gating strategy. The single cell suspension (left) was separated into TAM-enriched (CD11b(+), second from left) and tumor enriched (third from left) fractions. Tumor cells were defined via GFAP or 5-ALA after confirmation of 5-ALA-positivity during surgery or after confirmation of GFAP-positivity during neuropathological workup of the same samples. 5-ALA positive cells co-localized with GFAP-positive cells in samples where both markers allowed tumor cell identification (right).

(**B**) Relative distribution of tumor cells and TAMs in the single cell suspension of human glioma samples revealed similar heterogeneity in human high-grade glioma (HGG, n=10) compared to low-grade glioma (LGG, n=2). The initial two patients did not receive an analysis of the single cell suspension.

(**C**) Regression model indicated strong contribution of TAMs (CD11b(+) cells) but not of CD11b(-)/non-tumor cells to the measured activity in the investigated TAM-enriched samples of patients that underwent biopsy (n=10). Linear regression, β = standardized regression coefficient, error bands represent 95% confidence interval.

(**D**) Comparison of single cell tracer uptake (scTSPO) of tumor cells and TAMs in samples of human HGG (n=11) and LGG (n=9). Tumor cells of HGG indicated higher scTSPO than tumor cells of LGG, whereas there was no significant difference for TAMs (unpaired t-test). Tumor cells of HGG showed higher scTSPO than TAMs of HGG (paired t-test). Mean±SEM.

(**E**) Correlation of TSPO-PET signals with scTSPO elucidated a strong correlation between PET signals and tumor cell TSPO enrichment and no correlation between PET signals and TAM TSPO enrichment. N=13, R = Pearson's coefficient of correlation. Error bands represent 95% confidence interval.

(**F**) Example of three patients with HGG investigated via *in vitro* scRadiotracing. All three patients showed similar signals in amino acid PET (FET-PET) and only little contrast enhancement in MRI. The patient with high tumoral TSPO-PET signal (upper row, glioblastoma, WHO grade IV, high-affinity binding status (HAB)) showed distinctly more TSPO tracer uptake in both tumor cells and TAMs when compared to the patients with only faint (middle row, astrocytoma, WHO grade III, high-affinity binding status) or low (bottom row, glioblastoma, WHO grade IV, low-affinity binding status (LAB)) tumoral signal in TSPO-PET. The patient with glioblastoma and low-affinity binding status (bottom row) showed nearly absent single cell tracer uptake of TAMs but notable TSPO tracer uptake of tumor cells. T1w = T1-weighted, FLAIR = Fluid-attenuated inversion recovery.

 Figure 5

Figure 5: Integrated analysis of regional PET signals, cellular tracer uptake and 3D-histology via tissue clearing and light sheet imaging

(**A**) A glioblastoma inoculated brain (top left) is cleared with a modified version of 3DISCO[44] and imaged with light sheet florescent microscope (bottom left). The details of the clearing protocol are shown on the right panel.

(**B**) The imaged brains are annotated using a VR-based annotation tool, representative masks of the tumor (middle) and TAMs (right) are demonstrated. This process determined the occupied volume of GFP(+) tumor cells and CD11b(+) TAMs within the whole tumor for each individual mouse using segmentation via syGlass virtual reality tool[45].

(**C**) Quantitative comparison of determined GFP(+) and CD11b(+) volumes within individual SB28 tumors (n=7, paired t-test).

(**D**) Confocal microscopy was used to determine the average GFP(+) volume per single tumor cell and the average CD11b(+) volume per single TAM by automatized count of DAPI nuclear stains within regions of exclusive GFP/CD11b expression. GFP(+) tumor cells accumulated within the tumor core (upper row), whereas the tumor periphery provided areas that were nearly exclusively composed of CD11b(+) TAMs (lower row). The right column illustrates nuclei segmentation using cellpose[46] algorithm, which was applied to determine cell numbers per section.

(**E**) Upper row: Quantitative comparison of the absolute tumor and TAM cell count after extrapolation by individual light sheet volumes (GFP-occupancy and CD11b-occupancy) and averaged GFP/CD11b volumes per single cell (left). Tumor cells revealed higher numerical abundance over TAMs in SB28 tumors (n=7, paired t-test). Lower row: Contribution of tumor cells to the overall TSPO-PET signal exceeded contribution of TAMs (3.5:1). Lines connecting tumor cell and TAM contributions indicate the matrix of variance as determined by means and SEM bands of scTSPO and volume proportions (thick lines: mean of scTSPO x upper/lower SEM Band of tumor volumes; thin lines: upper/lower SEM band of scTSPO x upper/lower SEM Band of tumor volumes).

(**F**) Regional TSPO-PET signals (left) were combined with single cell tracer uptake values of tumor cells and TAMs to predict cell type abundance within individual SB28 tumors. Predicted 3D-histology (middle) shows high regional agreement with standard of truth 3D-histology as obtained by light sheet microscopy (right). The whole set of analyzed mice is presented in **Extended Fig. 3**.

(**G**) Representative magnified areas within the SB28 tumor underline the regional agreement between predicted and standard of truth 3D histology.

(**H**) Quantitative regional correlation of predicted and standard of truth 3D-histology within k means clusters (**Fig. 3**) shows high agreement of predicted tumor cell abundance and moderate agreement of predicted TAM abundance. Light sheet TIFFs were spatially co-registered to PET as 3D RGB images to extract signal intensities of light sheet as a surrogate of cell type abundance in all PET derived clusters. To predict 3D histology (separately for GFP and CD11b), PET signal intensities in each cluster were multiplied with the cluster-based PET correlation coefficient (**Fig. 3F**) and divided by single cell tracer uptake. n=350 single regions from n=7 tumors. R = Pearson's coefficient of correlation. Error bands represent 95% confidence interval. arb. = arbitrary units.

(**I**) PET measured radioactivity of the tumor was predicted by the radioactivity derived from the product of cell type abundance (tumor cells and TAMs) and single cell tracer uptake in all individual mice that underwent PET imaging and light sheet microscopy. R = Pearson's coefficient of correlation. Error bands represent 95% confidence interval. Dashed line represents line of identity (y=x).

Figure 6

Figure 6: Proteome analysis of the SB28 tumor microenvironment identifies TAM-specific radiotracer targets for glioblastoma

(**A**) Volcano plot representation of the different protein levels in isolated TAMs (n = 5) and isolated SB28 tumors cells (n = 3) in comparison with control microglia isolated from age matched mice (wt-microglia; n = 6). TSPO (yellow) showed intermediate elevation of protein levels in TAMs and SB28 tumor cells. 1098 out of 7869 analyzed proteins indicated higher protein levels in TAMs than did TSPO.

(**B**) Selection process of potential radiotracer targets for specific detection of TAMs over SB28 tumor cells. 165 of the identified proteins in TAMs were not detected in SB28 tumor cells and another 16 proteins had >10-fold higher levels in TAMs compared to SB28 tumor cells. A total of 20 proteins showed low cellular off-target sources and were identified as potential TAM radiotracer targets.

(**C**) Heat-map screening for TAM-specificity of the 181 proteins of interest. The Human Protein Atlas[47] was used to determine RNA expression levels of all proteins of interest in i) resident off-target cells of the brain (left cell type columns), ii) resident and infiltrating cells in the presence of glioblastoma (middle cell type columns) and iii) off-target cells of the organism (right cell type columns). Genes are sorted by protein level differences in TAMs compared to control microglia of untreated mice (top to bottom; log2 label-free quantification ratio; first column). 64 proteins with highest elevation (log2 (TAMs/wt-microglia) > 2, left column) are illustrated in **C** and all proteins are provided in **Extended Fig. 4**. Proteins of interest are highlighted in grey.

(**D**) Immune cell expression cluster allocation of proteins of interest. scRNA data of the Human Protein Atlas was used to categorize the final set of identified proteins.

Methods

Study design

The primary goal of the study is to resolve PET tracer signals at cellular resolution. We challenged our novel methodology by targeting TSPO as a biomarker with complex cellular sources in glioblastoma. For this purpose, a combination of radiotracer injection and immunomagnetic cell sorting (MACS) was applied (scRadiotracing[11]). To validate the workflow, the MACS data of microglia in untreated mice were compared to microglia depletion experiments using CSF1R inhibition in combination with [^{18}F]GE-180 TSPO-PET. In mice with orthotopic implanted tumor (SB28, 2.5 weeks post inoculation) or sham injection, the brain was removed 75 minutes after [^{18}F]GE-180 injection and whole-brain activity was measured. The tumor or the injection site was dissected, followed by radioactivity measurement and dissociation of the tumor tissue into a single cell suspension. Tumor cells (negative selection) and TAMs (CD11b) were isolated via magnetic cell sorting using specific antibodies. The radioactivity in the cell-specific pellets was measured by a highly sensitive gamma counter. To determine the absolute cell number in each pellet as well as the proportions (i.e. purity) of the different cell populations, flow cytometry was performed subsequent to cell sorting. TSPO tracer uptake was determined per single cell and compared to TSPO protein expression levels, whereas contributions of the different cell populations to the total activity were calculated by multiple regression. The methodology was translated to human resection and biopsy samples of patients with high-grade and low-grade glioma and *in vitro* scRadiotracing was performed to determine proportions of tumor and immune cell tracer uptake. To acknowledge PET signals as product of cellular uptake and cellular abundance, 3D-histology via light sheet microscopy was performed in PFA perfused mouse brains to obtain absolute cell counts and relative quantitative proportions of glioblastoma and immune cells within the tumor. Then, an integrated model of PET, single cell tracer uptake and 3D-histology was used to disentangle regional PET signals by their distinct cellular components. Finally, TSPO protein levels were characterized within the whole spectrum of

the tumor and immune cell proteome, with the goal to determine TAM-specific radiotracer

targets.

Animals

All animal experiments were performed in compliance with the National Guidelines for Animal

Protection, Germany and with the approval of the regional animal committee (Government of

Upper Bavaria) and overseen by a veterinarian. All animals were housed in a temperature-

and humidity-controlled environment with a 12-h light–dark cycle, with free access to food

(Ssniff, Soest, Germany) and water.

For scRadiotracing experiments, eight-week-old C57BL/6 mice were purchased from Charles

River (Sulzfeld, Germany) and acclimated for at least 1 week. At day 0, the mice were

inoculated with 100,000 SB28-GFP cells suspended in 2 µl of DMEM (Merck, Darmstadt,

Germany) (glioblastoma mice, n=27) or 2 µl of saline (sham mice, n=14). Additional n=8 mice

received no treatment before scRadiotracing, serving as control animals. For inoculation,

mice were anesthetized with intraperitoneal (i.p.) injections of 100 mg/kg ketamine 10% and

10 mg/kg xylazine 2% in 0.9% NaCl. Anesthetized mice were immobilized and mounted onto

a stereotactic head holder (David Kopf Instruments, Tujunga, CA, USA) in the flat-skull

position. After surface disinfection, the skin of the skull was dissected with a scalpel blade.

The scull was carefully drilled with a micromotor high-speed drill (Stoelting Co, Wood Dale,

IL, USA) 2 mm posterior and 1 mm left of the bregma. By stereotactic injection, 1×10^5 cells

were applied with a 10 µl Hamilton syringe (Hamilton, Bonaduz, Switzerland) at a depth of 2

mm below the drill hole. Cells were slowly injected within 1 minute and after a settling period

of another 2 minutes the needle was removed in 1 mm steps per minute. After that, the

wound was closed by suturing. Mice were checked daily for tumor-related symptoms and

sacrificed when tumor burden (i.e. appearance, coordinative deficits, motor symptoms)

reached stop criteria (not reached in any animal). On the last day of the experiment, the mice

were injected with 15 ± 1 MBq [^{18}F]GE-180 into the tail vein before cervical dislocation and

brain extraction at 75 minutes post injection (n=33). N=15 of these glioblastoma and all sham

mice received TSPO-PET imaging directly before brain extraction. Additional n=7

glioblastoma mice received TSPO-PET imaging and perfusion with 4% PFA prior to 3D-

histology by light sheet microscopy. Another n=14 C57BL/6 mice were used for extraction of

naïve control microglia (n=8 scRadiotracing; n=6 proteomics).

For validation of scRadiotracing as a reliable method to determine TSPO-PET signal sources

at cellular resolution, we analyzed combined data of pharmacological depletion of microglia

by CSF1R inhibition[48] which was performed in C57BL/6 mice of two datasets[13,14] by PLX5622

(1200 ppm, n=15) and vehicle controls (n=14), aged 10.1 ± 2.1 months. The TSPO-PET

scan was performed in the last week of seven weeks treatment.

Cell culture

SB28-GFP cells[49] were cultured in DMEM containing MEM non-essential amino acids (1x),

1% Penicillin-Streptomycin solution (Thermo Fisher Scientific, Waltham, MA, USA) and 10%

fetal bovine serum (FBS, Biochrome, Berlin, Germany). Cell cultures were maintained in the

incubator at 37°C in humidified and 5% CO_2-conditioned atmosphere. Cells were passaged

when the cell density in the flask reached 80% confluence.

Radiosynthesis

Automated production of [^{18}F]GE-180 was performed on a FASTlab^{TM} synthesizer with

single-use disposable cassettes. The pre-filled precursor vial was assembled on the cassette

and the cassette was mounted on the synthesizer according to the set-up instructions. The

FASTlab^{TM} control software prompts were followed to run the cassette test and to start the

synthesis. No-carrier-added ^{18}F-fluoride was produced via ^{18}O(p, n)^{18}F reaction by proton

irradiation of ^{18}O-enriched water and delivered to the ^{18}F incoming reservoir. The fully

automated manufacturing process consists of the following steps: trapping of ^{18}F-fluoride on

a QMA cartridge, elution using Kryptofix®222, potassium hydrogen carbonate, water and

acetonitrile, azeotropic drying of ^{18}F-fluoride at 120°C for 9 minutes, labelling of the precursor

in MeCN at 100°C for 6 minutes, dilution of the crude product with water, tC18 cartridge

based purification by use of 20 mL 40% (v/v) Ethanol and 11.5 mL 35% (v/v) Ethanol, elution

of the product with 3.5 mL 55% (v/v) Ethanol and final formulation with phosphate buffer.

RCY 39±7% (n=16) non d. c., synthesis time 43 minutes, RCP ≥98%.

Immunomagnetic cell separation

MACS was performed as described previously[13] with slight modifications for additional tumor

cell isolation. Detailed descriptions of brain dissociation and isolation of different cell types

were as follows.

Mouse brain dissociation

Adult mouse brains were removed after cervical dislocation at 75 minutes post injection (p.i.)

and stored in cold D-PBS. The brains were cut into small pieces and dissociated using

gentleMACS Octo Dissociator with Heaters (Miltenyi Biotec, 130-096-427) in combination

with different Dissociation Kits according to the manufacturer's instructions. Adult Brain

Dissociation Kit, mouse and rat (Miltenyi Biotec, 130-107-677) was used for adult mouse

brain dissociation of WT mice and contralateral hemispheres of glioblastoma mice. Tumor

Dissociation Kit (Miltenyi Biotec, 130-096-730) was used for dissociation of tumor tissue. The

dissociated cell suspension was applied to pre-wet 70-µm Cell Strainer (Miltenyi Biotec, 130-

110-916). The cell pellet was resuspended using cold D-PBS and cold debris removal

solution. Cold D-PBS was gently overlaid on the cell suspension and centrifuged at 4°C and

3000g for 10 minutes with acceleration at 9 and deceleration at 5. The two top phases were

removed entirely. The cell pellets were collected. Non-tumoral cell pellets were additionally

resuspended with 1 mL of cold red blood cell removal solution followed by 10 minutes

incubation. Cell pellets were collected for further applications.

Isolation of tumor cells

Tumor Cell Isolation Kit, mouse (Miltenyi Biotec, 130-110-187) was used according to the

manufacturer's instructions. The prepared cell pellets were resuspended in 80 µl of D-PBS–

0.5% bovine serum albumin (BSA) buffer per 10^7 total cells including red blood cells. 20 µl of

Non-Tumor Cell Depletion Cocktail were added and the suspension was incubated at 4°C in

the dark for 15 minutes. The volume was adjusted to 500 µl per total 10^7 cells with D-PBS–

0.5% BSA buffer before proceeding to magnetic separation. The pre-wet LS columns

(Miltenyi Biotec, 130-042-401) were placed at QuadroMACS Separator (Miltenyi Biotec, 130-

090-976). The cell suspensions were applied onto the column. The columns were washed

with 2 × 1 mL of D-PBS–0.5% BSA buffer. The flow-through containing the unlabeled cells

was collected as the tumor-cell-enriched fractions. The columns were removed from the

magnetic field, and the non-tumor cells were flashed out using 3 mL of D-PBS–0.5% BSA

buffer.

Isolation of TAMs and microglia

TAMs (glioblastoma mice) or microglia (sham injected or untreated control mice) were

isolated from animals using CD11b MicroBeads, human and mouse (Miltenyi Biotec, 130-

049-601) and a MACS separation system (Miltenyi Biotec) as described previously[50,51]. For

murine samples, the prepared cell pellets were resuspended in 90 µl of D-PBS–0.5% BSA

buffer per 10^7 total cells. Ten microliters of CD11b MicroBeads per 10^7 total cells were added

and incubated for 15 minutes in the dark at 4°C. Human samples were resuspended in 80 µl

of D-PBS–0.5% BSA buffer per 10^7 total cells and 20 µl of CD11b MicroBeads per 10^7 total

cells were added and cells were incubated for 15 minutes in the dark at 4°C. Cells were

washed by adding 1 to 2 mL of buffer per 10^7 cells and centrifuged at 300g for 10 minutes.

The cell pellets were resuspended in 500 µl of D-PBS–0.5% BSA. The pre-wet LS columns

(Miltenyi Biotec, 130-042-401) were placed onto a QuadroMACS Separator (Miltenyi Biotec,

130-090-976). The cell suspensions were applied onto the column. The columns were

washed with 3 × 3 mL of D-PBS–0.5% BSA buffer. The flow-through containing the

unlabeled cells was collected as the microglia-depleted fractions. The columns were

removed from the magnetic field, and microglia were flashed out using 5 mL of D-PBS–0.5%

BSA buffer.

Isolation of astrocytes

Adult Brain Dissociation Kit, mouse and rat (Miltenyi Biotec, 130-107-677) was used according to the manufacturer's instructions. The prepared cell pellets were resuspended in 80 µl of AstroMACS separation buffer (Miltenyi Biotec, 130-117-336) per 10^7 total cells. 10 µl of FcR blocking reagent were added and incubated for 10 minutes in the dark at 4°C. 10 µl of Anti-ACSA2 MicroBeads were added and incubated for 15 minutes in the dark at 4°C. Cells were washed by adding 1 mL of AstroMACS separation buffer and centrifuged at 300g for 5 minutes. Cell pellets were resuspended in 500 µl of AstroMACS separation buffer. The pre-wet MS columns (Miltenyi Biotec, 130-042-201) were placed at OctoMACS Separator (Miltenyi Biotec, 130-042-109). The cell suspensions were applied onto the column, followed by washing with 3 × 500 µl of AstroMACS separation buffer. The flow-through was collected containing non-astrocytic cells as an astrocyte-depleted fraction. The columns were removed from the magnetic field, and the astrocytes were flashed out using 3 mL of AstroMACS separation buffer.

Gamma emission measurements

Radioactivity concentrations of cell pellets were measured in a gamma counter (Hidex AMG Automatic Gamma Counter, Mainz Germany), cross-calibrated to the activity in the whole brain, with decay correction to time of tracer injection for final activity calculations.

Flow cytometry

Flow cytometry staining was performed at 4 °C. After gamma emission measurement, the cell suspension was centrifuged at 400g for 5 minutes and the supernatant was aspirated completely. The cell pellet was then resuspended in 100 µl of cold D-PBS containing fluorochrome-conjugated antibodies recognizing mouse CD11b and ACSA2 (Miltenyi Biotec, 130-113-810 and 130-116-247) in a 1:100 dilution and incubated for 10 minutes at 4°C in the dark. Samples were washed with 2 mL of D-PBS and centrifuged for 5 minutes at 400g.

Finally, cell pellets were resuspended in 500 µl of D-PBS and samples were immediately

used for flow cytometry using a MACSQuant® Analyzer. Precision Count Beads (Biolegend,

424902) were added for counting the absolute number of cells for the samples of n=9 mice

measured with a BD LSR Fortessa Cell Analyzer (BD Biosciences, Franklin Lakes, NJ,

USA). Acquired data included absolute cell numbers and purity of GFP(+), CD11b(+) and

ACSA2(+) cells in each sample.

TSPO co-staining was performed for tumor and CD11b enriched cell fractions of n=3 SB28

mice. To this end, cells were permeabilized after initial flow cytometry using Inside Stain Kit

(Miltenyi Biotec, 130-090-477) according to the manufacturer's instructions. Cell pellets were

resuspended in 250 µl of cold D-PBS, 250 µl of Inside Fix Solution was added, cells were

incubated for 20 minutes in the dark at 4°C, then centrifuged at 300g for 5 minutes. Complete

aspiration of supernatant was followed by washing with 1 mL of cold D-PBS and another

centrifugation step at 300g for 5 minutes. Cell pellets were washed with 1 mL of Inside Perm

Solution and centrifuged at 300g for 5 minutes. Supernatant was aspirated, cells were

resuspended in 47 µl of Inside Perm Solution and stained with 3 µl of Anti-PBR antibody

[EPR5384] (abcam, ab199836). After 10 minutes of incubation, cells were washed by adding

1 mL of Inside Perm Solution, centrifuged at 300g for 5 minutes and resuspended in 300 µl

cold D-PBS for flow cytometry analysis.

For subsequent proteome analysis (see below), cell pellets of n=3 tumor enriched and n=5

CD11b enriched fractions of SB28 tumor samples were stored at -80 degree together with

n=6 CD11b enriched fractions (i.e. control microglia) derived from an independent cohort of

age- and sex matched untreated control mice.

Calculation of single cell TSPO tracer signal

Measured radioactivity (Bq) of cell pellets was divided by the specific cell number in the pellet

resulting in calculated radioactivity per cell. Radioactivity per cell was normalized by injected

radioactivity and body weight (i.e. %ID*BW). Published cell numbers of microglia[12] were used

to extrapolate the whole brain radioactivity located in the microglia fraction of untreated mice.

For tumor probes, cell numbers (tumor cells and TAMs) determined by light sheet

microscopy were multiplied with single cell %ID*BW as an estimate of cell fraction specific

contributions to the PET signal.

Human samples and in vitro scRadiotracing

Human tumor tissue samples were acquired during neurosurgical biopsy or open resection

and stored in Tissue Storage Solution (Miltenyi Biotec, 130-100-008) for 2-30 hours until

further processing for scRadiotracing. Details on the patient cohort are provided in the results

section (**Table 1**). Tissue was manually cut into smaller pieces if necessary and dissociated

using Tumor Dissociation Kit, human (Miltenyi Biotec, 130-095-929). Removal of debris and

red blood cells was performed for tumor resections as described above. Single cell

suspensions were incubated *in vitro* at a concentration of 56 ± 8 MBq [^{18}F]GE-180 in 1 mL for

30 minutes, then washed twice with 3 mL cold D-PBS and centrifuged at 400g for 5 minutes.

Cell pellets were resuspended in 100 µl cold D-PBS and stained with fluorochrome-

conjugated antibodies recognizing human CD11b and GFAP (Miltenyi Biotec, 130-113-810

and 130-123-846) as described above. Gamma counter measurement and flow cytometry

analysis of the stained single cell suspension were performed using 10% of the probe after

resuspension in 500 µl. Subsequently, TAM were isolated using CD11b microbeads (Miltenyi

Biotec, 130-093-634), followed by further gamma counter measurement and flow cytometry.

Proportions and cell count of TAM in the resulting cell pellets were determined using CD11b.

For tumor cells, a step-wise validation of the gating strategy was performed. Detection of

GFAP-positive cells was used for n=6 samples where GFAP positivity was validated by

staining during neuropathological workup of the same tumors. N=4 patients received 5-

aminolevulinic acid (5-ALA) guided surgery and differentiation of tumor cell populations by 5-

ALA-positivity in flow cytometry[52]. These n=10 samples were used for cross-validation of

autofluorescence-based gating (excitation wavelength 488 nm, emission wavelengths 655-

730nm)[53] which was applied in another n=10 samples without GFAP- or 5-ALA-positivity.

For samples containing very high proportions of TAMs, in the first step, the radioactivity per

single TAM cell was calculated using the well purified CD11b(+) TAM enriched fraction (see

gating and analysis strategy A in **Extended Fig. 2**). Since the tumor cells in the depleted

fraction were not purified to a satisfactory degree (<70%), the radioactivity attributable to the

fraction of TAM cells was subtracted from the total activity of the depleted fraction and the

remaining radioactivity was divided by the total cell number of tumor cells in the depleted

fraction for calculation of radioactivity per single tumor cell. For samples containing very low

concentrations of TAM, in the first step, the activity per single tumor cell was calculated in the

CD11(-) fraction (see gating and analysis strategy B in **Extended Fig. 2**). Since TAM in the

CD11b(+) fractions were not purified to a satisfactory degree (<70%), the radioactivity

attributable to tumor cells was subtracted from the measured radioactivity in the TAM

enriched fraction and the remaining radioactivity was divided by the total cell number of

CD11b(+) cells in the TAM enriched fraction to obtain radioactivity per TAM. Negligibility of

TSPO tracer uptake by non-TAM and non-tumor cells for the overall radioactivity in the cell

pellets was confirmed by a regression model of cell numbers and gamma emission in TAM

enriched fractions of all biopsy samples.

Small-animal PET/CT

All small animal positron emission tomography (PET) procedures followed an established

standardized protocol for radiochemistry, acquisition, and post-processing[54,55]. For tumor and

sham mice, [^{18}F]GE-180 TSPO small-animal PET (12 ± 1 MBq, n=36) recordings with an

emission window of 0-60 minutes after injection were obtained to measure cerebral TSPO

expression prior to sorting using a Mediso PET/CT system (Mediso, Budapest, Hungary). A

contrast enhanced CT was performed prior to the PET scan. For the mice of the depletion

experiment, a static [^{18}F]GE-180 emission (13 ± 2 MBq) was recorded between 60 and 90

minutes after injection using a harmonized Siemens Inveon DPET (Siemens, Munich,

Germany). All small-animal PET experiments were performed with isoflurane anesthesia

(1.5% at time of tracer injection and during imaging; delivery 3.5 L/min).

All analyses were performed by PMOD (V3.5, PMOD Technologies, Basel, Switzerland)

using CT (tumor and sham mice) and tracer-specific templates (depletion experiment) for

spatial normalization[54]. For tumor mice, a 40-60 minute frame was analyzed and

normalization of activity was performed by SUV. The average TSPO-PET SUV was obtained

from a spherical volume of interest at the tumor site as the primary PET read-out[16]. Within

the tumor volume of interest, we performed a 50-step k means clustering using the PMOD

segmentation tool (10 iterations) to delineate individual clusters of tumor tracer uptake in

each tumor. For correlation analysis with light-sheet microscopy, a 50% threshold was

applied prior to clustering to avoid artificial associations by inclusion of necrotic parts of the

tumor.

For the depletion experiment, normalization of injected activity was performed by the

previously validated myocardium correction method[56] and PET estimates deriving from a

whole brain volume of interest[54] were extracted.

Human PET/CT

TSPO-PET was available for 13 out of 20 (65%) patients that underwent scRadiotracing. All

human TSPO-PET scans were performed on a Biograph 64 PET/CT scanner (Siemens,

Erlangen, Germany). Tracer production and image acquisition were performed as described

previously[57]. Approximately 180 MBq [^{18}F]GE-180 were injected as an intravenous bolus and

summation images 60-80 minutes p.i. were used for image analysis using a Hermes

workstation (Hermes Medical Solutions, Stockholm, Sweden). MRI sequences included

gadolinium-enhanced T1- and T2-weighted images and were used for anatomical mapping of

stereotactic biopsy coordinates and resected tumor mass within the TSPO-PET images

using BRAINLAB ELEMENTS™ (Brainlab AG, Munich, Germany). For correlation analysis

between scRadiotracing and TSPO-PET imaging, the mean SUV was obtained from a 0.2

cm³ (25 voxels) volume-of-interest using the sample coordinates.

Amino acid PET was available for a subset of the investigated patients (80%) and served for

characterization of vital tumor tissue. Approximately 180 MBq [^{18}F]FET were injected and

summation images 20-40 minutes p.i. were analyzed. Tumor uptake was assessed as

maximum standardized uptake value (SUV_{max}). As described previously[58], the mean

background activity was defined as the mean activity of at least 6 crescent-shaped cortical

areas in the healthy contralateral side and SUV_{mean} and SUV_{max} were divided by the mean

background activity to obtain mean and maximal tumor-to-background ratios (TBR_{mean} and

TBR_{max}). The biological tumor volume was semiautomatically delineated using the standard

1.6 x background activity as threshold.

Neuropathological analysis

5 µm thick sections of formalin-fixed and paraffin-embedded (FFPE) tumor tissue were

routinely stained with Hematoxylin and eosin (H&E). In addition, immunohistochemical

stainings using antibodies against GFAP (polyclonal; Agilent Technologies, Santa Clara, CA,

USA), MAP2 (HM-2; Merck, Darmstadt, Germany), IDH 1 (R132H) (QM002; quartett GmbH,

Potsdam, Germany), ATRX (BSB-108; Bio SB, Goleta, CA, USA) and Ki67 (MIB1; Agilent

Technologies, Santa Clara, CA, USA) were routinely performed according to standard

protocols.

1004

Transcardial perfusion, immunohistochemistry and tissue clearing

Mice that were intended for 3D-histology were i.p. injected with 100 mg/kg ketamine 10%

and 10 mg/kg xylazine 2% in 0.9% NaCl. After expiration of the pedal reflex, intracardial

perfusion was performed with 0.1 M PBS (10 U/mL, Ratiopharm) for 6 minutes, followed by

the administration of 4% paraformaldehyde (PFA) in 0.1 M PBS (pH 7.4) (Morphisto,

11762.01000) for another 6 minutes. Afterwards, the brain was removed and post-fixed by

4% PFA for 6-12 hours at 4°C and washed with 0.1 M PBS and stored in 0.1 M PBS.

Samples were subjected to a modified version of vDISCO[44] and SHANEL[59] protocols.

Samples were decolorized with 25% CUBIC solution for 1 day. An extra permeabilization

step was performed using SHANEL reagents: 10% CHAPS with 25% N-

Methyldiethanolamine in dH2O at 37°C overnight in order to access the dense glioblastoma.

Then, samples were further permeabilized with vDISCO permeabilization solution with additional 10% 2- Hydroxypropyl-β-cyclodextrin overnight at 37°C. The next day, antibody labeling and boosting step was performed with the addition of CD11b (1:50) (Miltenyi Biotec, 130-110-611) and GFP-Nanobooster LOT 102 (1:500) (Chromotek, gba647n). The samples were incubated in 37°C for 9 days. After labeling, samples were washed with vDISCO washing solution for 3 hours and 3DISCO clearing[60] with 50-70-90-100-100% tetrahydrofuran (THF) (Sigma, 186562), 1h each, dichloromethane (Sigma, 270997), 30 minutes, was performed. Lastly, samples were placed in BABB (2:1, Benzyl Benzoate, Benzyl alcohol) (Sigma, 24122 and W213802) solution indefinitely for refractive index matching and light sheet fluorescent imaging.

Light sheet fluorescent microscopy

Ultramicroscope Blaze (Miltenyi Biotec) was used to acquire single plane illumination (light sheet) image stacks. Filters used were ex 470/40 nm, em 535/50 nm; ex 545/25 nm, em 605/70 nm; ex 560/30 nm, em 609/54 nm for autofluorescence, CD11b signal and cancer cells respectively. For all scans, 1.66x magnification, 1x2 tiling, 30% overlap, 120 ms exposure time, 100% light sheet width, ~7 μm laser sheet thickness, ~0.31 N/A and 5 μm z-step was used. In order to prevent signal saturation, laser power was adjusted per each sample.

Cryosectioning and immunofluorescence staining

To determine the absolute cell numbers within tumors, after acquiring the light sheet images of brains, tissue was rehydrated for sectioning. After sequential incubation in THF, samples were washed in PBS and placed in 30% sucrose solution for dehydration overnight. The next morning, samples were embedded in OCT and frozen. 15 μm sagittal sections were acquired using a cryostat (CM3050S; 665 Leica, Wetzlar, Germany). Tissue sections blocked 0.2% TritonX-100, 10% DMSO,10% goat serum in 0.1 M PBS for 1 hour, stained, CD11b (1:50) and GFP-Nanobooster LOT 102 (1:500) in 0.2% Tween-20, 5% DMSO, 5% goat serum,

1044   0.001% heparin in 0.1 M PBS and washed with 0.2% Tween-20, 0.001% heparin in 0.1 M

1045   PBS for 3x2 minutes.

1046

Laser scanning confocal microscopy

Leica SP8 was utilized for laser scanning confocal microscopy. Images were acquired with

40x magnification acquired as 8-bits with HC PL APO CS2 40 × /1.30 NA HC PL APO CS2

40×/1.30 NA or a HC PL APO CORR CS2 63×/1.30 NA objective, 1,024 × 1,024 resolution,

200 Hz, at 20–25°C. Tile scans were obtained with z-step size of 1 μm to allow 3D

reconstruction of tissue for depth of 15 μm.

1053

Image processing

Tiles from each brain were stitched as described previously[44]. Stitched images were saved in

TIFF format for further processing. Volumes of segmented CD11b-positivity per TAM and

segmented GFP-positivity per tumor cell were obtained via confocal images to allow

calculation of absolute cell numbers in 3D-histology (see **Equation 1**). Regions of interest

with predominant CD11b (n=5) and cancer cell (n=5) signal were acquired and DAPI nuclear

stains were used to count cells per ROI. To segment single nuclei from cancer cells and

CD11b(+) cells for estimation of absolute cell numbers, cellpose[46] was utilized. The

percentage of fluorescence(+) nuclei/cells was determined via co-localization of nuclei with

adjacent CD11b- and GFP-positivity. Volumes of CD11b- and GFP-positivity were obtained

via assessment of segmented areas per channel (50% max. threshold) for each single layer

after applying a Gaussian filter of 6mm. To account for positive areas not belonging to

captured nuclei (i.e. at the edge of cells) we determined the area of CD11b- and GFP-

positivity remote from nuclei (manual threshold visually matching the cell borders). To

account for caption of variable profiles in single cells, and we assumed a Gaussian

distribution of modelled percentage areas in a fluorescence positive sphere (i.e. cell) with an

internal fluorescence negative sphere (i.e. nucleus), calculating a correction factor of 1.044.

Areas per cell were then extrapolated to volumes per cell using spherical assumption.

Equation 1:

$$cell\ volume\ (\mu m^3) = \frac{4}{3}\pi * \left[\sqrt{\frac{\left(\frac{segmented\ area\ (\mu m^2)}{segmented\ nuclei\ (n)} \times \frac{nuclei\ adjacent\ area\ (\mu m^2)}{total\ area\ (\mu m^2)} \times \frac{all\ nuclei\ (n)}{fluorescence(+)\ nuclei\ (n)} \times 1.044\right)}{\pi}} \right]^3$$

To calculate the volume of tumor cells and TAMs within the tumor in 3D-histology, two

channels were separately segmented using syGlass virtual reality tool[45]. The TIFF-stack was

imported into the syGlass software where the stack is rendered as a 3D image in a virtual

space. The ROI tool was used to annotate the signal of interest with adjustable thresholding

in the 3D space to get the best signal to noise ratio. The annotation mask was exported as

TIFFs for further downstream processing. Based on the VR-aided manual annotations of

tumor and TAM regions, areas were calculated as the sum of the marked mask voxels.

LSFM's resolution of 19.58 mm^3 per voxel was used to compute the total volume in mm^3.

The percentage of TAMs within the tumor was obtained by first filtering the annotated TAM

mask with the tumor area by dot multiplication between the two masks. Then, the number of

obtained voxels was divided by the total amount of tumor voxels. Finally, the PET signal of

the tumor was compared with the aggregated recovered signals of tumor cells and TAMs

(see **Equation 2**).

Equation 2:

$$PET\ signal\ (Bq) = \left(\frac{tumor\ GFP\ volume\ (\mu m^3)}{\frac{GFP\ (\mu m^3)}{tumor\ cell}} \times \frac{Bq}{tumor\ cell} \right) + \left(\frac{tumor\ CD11b\ volume\ (\mu m^3)}{\frac{CD11b\ (\mu m^3)}{TAM}} \times \frac{Bq}{TAM} \right)$$

1087

Proteomics

Isolated TAMs (n=5) and tumor cells (n=3), as well as control microglia isolated from

untreated control mice (n=6) were prepared for mass spectrometry based label free protein

quantification. The cell pellets were lysed in 50 µl of RIPA lysis buffer (50 mM Tris, 150 mM

NaCl, 5 mM EDTA, 1% (v/v) Triton, 0.5% (w/v) sodium deoxycholate, 0.1% (w/v) SDS, pH

8.0) at 4°C with intermediate vortexing. DNA was disrupted by ultrasonication (M220

Focused-ultrasonicator, Covaris, Woburn, MA, USA). The samples were centrifuged for 5

minutes at 16000g and 4°C to remove cell debris and undissolved material. The

supernatants were transferred to protein LoBind tubes (Eppendorf, Hamburg, Germany).

Proteins were reduced at 37°C for 30 minutes with 15 mM dithiothreitol (DTT) followed by

cysteine alkylation with 60 mM iodoacetamide (IAA) for 30 minutes at 20 °C. Excess of IAA

was removed by adding DTT. Detergent removal and subsequent digestion with 0.2 µg LysC

and 0.2 µg trypsin (Promega, Walldorf, Germany) was performed using the single-pot, solid-

phase-enhanced sample preparation as previously described[61]. After vacuum centrifugation,

peptides were dissolved in 20 µl 0.1% formic acid. The peptide amount of tumor cell samples

was estimated using a fluorescence based protein assay (Qubit Protein assay, Thermo

Fisher Scientific, Waltham, MA, USA).

For mass spectrometry, the injected volume was adjusted on the basis of the counted cell

numbers to analyze equal cell numbers with a maximum injection amount of 400 ng of

peptides. The peptides were separated on a nanoElute nanoHPLC system (Bruker, Bremen,

Germany) using a 5 mm trapping column (Thermofisher Scientific, Waltham, MA, USA) and

an in-house packed C18 analytical column (15 cm × 75 µm ID, ReproSil-Pur 120 C18-AQ,

1.9 µm, Dr. Maisch GmbH). Peptides were separated with a binary gradient of water and

acetonitrile (B) containing 0.1% formic acid at flow rate of 300 nL/min (0 min, 2% B; 2 min,

5% B; 70 min, 24% B; 85 min, 35% B; 90 min, 60% B) and a column temperature of 50°C.

The nanoHPLC was online coupled to a TimsTOF pro mass spectrometer (Bruker, Bremen,

Germany) with a CaptiveSpray ion source (Bruker, Bremen, Germany). A Data Independent

Acquisition Parallel Accumulation–Serial Fragmentation (DIA-PASEF) method for spectrum

acquisition. Briefly, ion accumulation and separation using Trapped Ion Mobility

Spectrometry (TIMS) was set to a ramp time of 100 ms. One scan cycle included one TIMS

full MS scan and with 26 windows with a width of 27 m/z covering a m/z range of 350-1002

m/z. Two windows were recorded per PASEF scan. This resulted in a cycle time of 1.4 s.

The data analysis and protein label-free quantification (LFQ) was performed with the

software DIA-NN[62] version 1.8. The data was analyzed using a spectral library including

13912 protein groups and 208218 precursors in 174909 elution groups generated with the

same samples and additional murine microglia samples using DIA-NN searching against a

one protein per gene database from Mus musculus (download date: 2022-01-25, 21994

entries). Trypsin was defined as protease and 2 missed cleavages were allowed. Oxidation

of methionines and acetylation of protein N-termini were defined as variable modifications,

whereas carbamidomethylation of cysteines was defined as fixed modification. The precursor

and fragment ion m/z ranges were limited from 350 to 1002 and 200 to 1700, respectively.

Precursor charge states of 2-4 were considered. The mass accuracy for peptides and

peptide fragments was set to 15 and 20 ppm, respectively. A FDR threshold of 1% was

applied for peptide and protein identifications. Data normalization was disabled to obtain

quantification values, which reflect the protein amounts per cell.

Ratios of TSPO protein expression between SB28 tumor cells, TAMs and control microglia

were compared against corresponding ratios obtained by scRadiotracing. Furthermore, all

proteins were analyzed regarding their SB28 tumor cell and TAM expression levels relative

to control microglia. Potential TAM-specific radiotracer targets were identified in a multi-step

process. First, all proteins with higher levels in TAMs (TAM-to-control-microglia ratio) when

compared to TSPO were selected. Second, among these, we identified proteins that were

not present in SB28 tumor cells or showed high specificity in TAMs over SB28 tumor cells

(TAM-to-tumor cell ratios ≥10). Third, to further interrogate specificity, the Human Protein

Atlas[47] was used to analyze single cell RNA expression levels of brain and non-brain cells for

all identified target proteins that could serve for subsequent development of TAM-specific

radioligands. Fourth, the predominant cell type and the functional immune cell cluster of the

identified proteins for radiotracer development was determined by the Human Protein Atlas.

1145

Statistics

Statistical analyses were performed with Graph Pad Prism (V9, San Diego, CA, USA) and

SPSS (V26, IBM, Armonk, NY, USA). A significance threshold of $p < 0.05$ was considered as

significant for all experiments.

Validation of TSPO scRadiotracing: Radioactivity per cell was compared between microglia

and astrocytes by a paired Student's t-test. In the depletion experiment, TSPO-PET

measures of whole brain were compared between treatment and vehicle by an unpaired

Student's t-test. Iba1 immunoreactivity was compared between treatment and vehicle groups

by an unpaired Student's t-test. The decrease of the TSPO-PET signal in the treatment

group was considered as microglia bound radioactivity in brain and compared to extrapolated

radioactivity of the microglia cell population as assessed by scRadiotracing using a Student's

t-test.

scRadiotracing in SB28 glioblastoma mice: Radioactivity per cell (%ID*BW normalized) was

compared between different cell populations (tumor cells, TAMs, sham microglia, control

microglia) by a one-way analysis of variance with Tukey post hoc test or by a paired

Student's t-test for populations that were present in the same animals. Multiple regression

was used to determine contributions of tumor cells [GFP(+)], microglia [CD11b(+)] and

astrocytes [ACSA2(+)] to the radioactivity in the cell pellet. As a validation, measured

radioactivity in the depleted fraction of the tumor was correlated with predicted radioactivity

by cell count and single cell tracer uptake (Pearson's coefficient of correlation).

Association between scRadiotracing and PET imaging: Lesion site SUV was compared

between SB28 and sham mice by an unpaired Student's t-test. Single cell tracer uptake

(tumor cells, TAMs and a summed vector) was correlated with the tumor PET signal and

between cell types using Pearson's coefficient of correlation and we applied a regression

model with both cell types as predictors and the PET signal as dependent variable. PET

cluster values of the tumor segmentation were correlated with single cell tracer uptake and

plotted as a function of cluster grade, after determining the best curve fit (linear, quadratic,

exponential) by Akaike information criteria. The difference between the cluster agreement of

tumor cells and TAMs was calculated to determine dominant contribution of cell types and

likewise plotted as a function of cluster grade.

Human in vitro scRadiotracing: A regression model with cell count of TAMs and non-

TAM/non-tumor cells as predictors and pellet radioactivity as dependent variable was applied

to test for significant contribution of non-TAM/non-tumor cells to the total radioactivity. Single

cell tracer uptake was compared between cell types and between patients with high- and

low-grade glioma by an unpaired Student's t-test. Single cell tracer uptake (tumor cells and

TAMs) was correlated with the tumor PET signal using Pearson's coefficient of correlation.

Integrated model of PET, scRadiotracing and 3D-histology: Comparisons of GFP(+) and

CD11b(+) volumes and cell counts in SB28 tumors were performed using a paired Student's

t-test. The following analysis was performed using the 50 cluster regions of interest per

individual mouse. TSPO-PET SUVs were multiplied with the regional correlation coefficient of

either tumor or TAM single cell tracer uptake correlations with PET. The individual single cell

tracer uptake was estimated according to TSPO-PET SUV of the tumor (**Fig. 3B**) and applied

to predict light sheet signal intensity per cluster as a surrogate for cell density. This was

performed separately for tumor cells and TAMs. A linear regression was performed to test for

the agreement between predicted and standard of truth cell density per region. Finally, a

combined tumor cell and TAM model was used to predict absolute radioactivity in SB28

tumors as measured by means of TSPO-PET. To this end, the cell count of both tumor cells

and TAMs in seven SB28 tumors was multiplied with the individual single cell tracer uptake

(as estimated according to TSPO-PET SUV of the tumor) and aggregated before correlation

with equally normalized (%ID*BW) PET radioactivity.

Proteomics: The statistical data analysis of the DIA-NN output was performed with the

software Perseus Version 1.6.14.0[63]. Only LFQ intensities from protein groups with at least

two peptides were considered. The protein LFQ intensities were log2 transformed. A 1-way

ANOVA test was applied to determine statistically significant differences between the means

of the three groups. Afterwards, individual Student's t-tests were applied to evaluate proteins

with a significantly different abundance between the TAMs, tumor cells and control microglia.

Additionally, a permutation based false discovery rate estimation was used with a FDR of 5%

at $s_0 = 0.1$ as threshold[64].

All numeric values are reported as average group values ± standard error of the mean

(s.e.m.) unless otherwise indicated.

1206

Data availability

All source data (i.e. dynamic PET imaging data) is available per reasonable request to the corresponding author. Proteomics source data is available as extended data.

Conflict of Interest

NLA and MB are members of the Neuroimaging Committee of the EANM. JCT received research grants from Novocure and Munich Surgical Imaging and a speaker honorarium from Seagen. NLA received funding from Novocure. MB received speaker honoraria from Roche, GE healthcare and Life Molecular Imaging and is an advisor of Life Molecular Imaging. VCR received speaker honoraria from Novocure. All other authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions

LMB: performed tumor inoculation, performed PET acquisition, image analysis and interpretation of PET scans, established and performed scRadiotracing in mice and human tissue samples, performed scRadiotracing analyses, integrated data of 3D-histology and proteomics, wrote the first draft of the manuscript with input of all co-authors. SVK, SQ and JB: contributed and harvested SB28 cells, supported establishment of scRadiotracing, interpreted PET data in the context of human glioblastoma imaging, supported human scRadiotracing by suggesting patients and performing neurosurgical tissue acquisition. ZIK, SU, LH, IH: performed immunohistological analyses and 3D-histology, interpreted histological data and contributed representative images. SAM: performed and interpreted proteomics. KW, AH, LG, LHK: participated in preclinical PET acquisition, image analysis and interpretation. VCR, JH: neuropathological examination and characterization of the tumor samples. STK, PB, MA, LH, DM: supported scRadiotracing data acquisition and interpretation. AZ: performed 3-dimensional PET image reconstruction. NB: performed immunohistological analyses and interpretation of CSF1R inhibition experiments. SL:

1235 performed radiochemistry and TAM target identification. PB, MJR, SZ, JH, SFL, AE, JCT,

1236 LvB, NLA and MB: conception and design, contributed to interpreting data, enhancing

1237 intellectual content of manuscript. All authors contributed with intellectual content and revised

1238 the manuscript.

1239

Funding

This project was partly funded by the Deutsche Forschungsgemeinschaft (DFG, German

Research Foundation) (FOR 2858 project numbers 421887978 and 422188432 and

Research Training Group GRK 2274). NLA is supported by a research grant of the Else

Kröner-Fresenius-Stiftung. MB and SFL were funded by the Deutsche

Forschungsgemeinschaft (DFG) under Germany's Excellence Strategy within the framework

of the Munich Cluster for Systems Neurology (EXC 2145 SyNergy – ID 390857198). SVK

was supported by the Verein zur Förderung von Wissenschaft und Forschung an der

Medizinischen Fakultät der LMU München (WiFoMed) and the Friedrich-Baur-Stiftung.

1249

1250 **Acknowledgments**

1251 We thank Rosel Oos and Giovanna Palumbo for excellent technical support during PET

1252 imaging.

1253

1254

References

1   Jin, M.-Z. & Jin, W.-L. The updated landscape of tumor microenvironment and drug
repurposing. *Signal Transduction and Targeted Therapy* **5**, 166 (2020).
https://doi.org:10.1038/s41392-020-00280-x
2   Pitt, J. M. *et al.* Targeting the tumor microenvironment: removing obstruction to
anticancer immune responses and immunotherapy. *Annals of Oncology* **27**, 1482-1492
(2016). https://doi.org:https://doi.org/10.1093/annonc/mdw168
3   Fernandes, B. *et al.* Potential PET tracers for imaging of tumor-associated
macrophages. *EJNMMI Radiopharmacy and Chemistry* **7**, 11 (2022).
https://doi.org:10.1186/s41181-022-00163-2
4   Kist de Ruijter, L. *et al.* Whole-body CD8+ T cell visualization before and during
cancer immunotherapy: a phase 1/2 trial. *Nature Medicine* **28**, 2601-2610 (2022).
https://doi.org:10.1038/s41591-022-02084-8

1268	5	Low, J. T. *et al.* Primary brain and other central nervous system tumors in the United
1269		States (2014-2018): A summary of the CBTRUS statistical report for clinicians.
1270		*Neurooncol Pract* **9**, 165-182 (2022). https://doi.org:10.1093/nop/npac015
1271	6	Zinnhardt, B. *et al.* TSPO imaging-guided characterization of the immunosuppressive
1272		myeloid tumor microenvironment in patients with malignant glioma. *Neuro-oncology*
1273		**22**, 1030-1043 (2020).
1274	7	Cai, L. *et al.* Glioblastoma Exhibits Inter-Individual Heterogeneity of TSPO and
1275		LAT1 Expression in Neoplastic and Parenchymal Cells. *Int J Mol Sci* **21** (2020).
1276		https://doi.org:10.3390/ijms21020612
1277	8	Pigeon, H. *et al.* TSPO-PET and diffusion-weighted MRI for imaging a mouse model
1278		of infiltrative human glioma. *Neuro-oncology* **21**, 755-764 (2019).
1279	9	Zinnhardt, B. *et al.* Combined PET Imaging of the Inflammatory Tumor
1280		Microenvironment Identifies Margins of Unique Radiotracer Uptake. *Cancer Res* **77**,
1281		1831-1841 (2017). https://doi.org:10.1158/0008-5472.CAN-16-2628
1282	10	Buck, J. R. *et al.* Preclinical TSPO Ligand PET to Visualize Human Glioma
1283		Xenotransplants: A Preliminary Study. *PloS one* **10**, e0141659 (2015).
1284		https://doi.org:10.1371/journal.pone.0141659
1285	11	Bartos, L. M. *et al.* Single cell radiotracer allocation via immunomagnetic sorting
1286		(scRadiotracing) to disentangle PET signals at cellular resolution. *J Nucl Med* (2022).
1287		https://doi.org:10.2967/jnumed.122.264171
1288	12	Dos Santos, S. E. *et al.* Similar Microglial Cell Densities across Brain Structures and
1289		Mammalian Species: Implications for Brain Tissue Function. *J Neurosci* **40**, 4622-
1290		4643 (2020). https://doi.org:10.1523/JNEUROSCI.2339-19.2020
1291	13	Xiang, X. *et al.* Microglial activation states drive glucose uptake and FDG-PET
1292		alterations in neurodegenerative diseases. *Sci Transl Med* **13**, eabe5640 (2021).
1293		https://doi.org:10.1126/scitranslmed.abe5640
1294	14	Shi, Y. *et al.* Long-term diazepam treatment enhances microglial spine engulfment
1295		and impairs cognitive performance via the mitochondrial 18 kDa translocator protein
1296		(TSPO). *Nat Neurosci* **25**, 317-329 (2022). https://doi.org:10.1038/s41593-022-01013-
1297		9
1298	15	Genoud, V. *et al.* Responsiveness to anti-PD-1 and anti-CTLA-4 immune checkpoint
1299		blockade in SB28 and GL261 mouse glioma models. *Oncoimmunology* **7**, e1501137
1300		(2018).
1301	16	Bartos, L. M. *et al.* 18 kDa translocator protein positron emission tomography
1302		facilitates early and robust tumor detection in the immunocompetent SB28
1303		glioblastoma mouse model. *Front Med (Lausanne)* **9**, 992993 (2022).
1304		https://doi.org:10.3389/fmed.2022.992993
1305	17	Uhlén, M. *et al.* Tissue-based map of the human proteome. **347**, 1260419 (2015).
1306		https://doi.org:doi:10.1126/science.1260419
1307	18	Dagher, N. N. *et al.* Colony-stimulating factor 1 receptor inhibition prevents
1308		microglial plaque association and improves cognition in 3xTg-AD mice. *Journal of*
1309		*neuroinflammation* **12**, 1-14 (2015).
1310	19	Acharya, M. M. *et al.* Elimination of microglia improves cognitive function following
1311		cranial irradiation. *Scientific reports* **6**, 1-11 (2016).
1312	20	Elmore, M. R. *et al.* Colony-stimulating factor 1 receptor signaling is necessary for
1313		microglia viability, unmasking a microglia progenitor cell in the adult brain. *Neuron*
1314		**82**, 380-397 (2014).
1315	21	Parhizkar, S. *et al.* Loss of TREM2 function increases amyloid seeding but reduces
1316		plaque-associated ApoE. *Nat Neurosci* **22**, 191-204 (2019).
1317		https://doi.org:10.1038/s41593-018-0296-9

22   Greenbaum, D., Colangelo, C., Williams, K. & Gerstein, M. Comparing protein
abundance and mRNA expression levels on a genomic scale. *Genome Biol* **4**, 117
(2003). https://doi.org:10.1186/gb-2003-4-9-117
23   Zanotti-Fregonara, P. *et al.* Head-to-Head Comparison of (11)C-PBR28 and (18)F-
GE180 for Quantification of the Translocator Protein in the Human Brain. *J Nucl Med*
59, 1260-1266 (2018). https://doi.org:10.2967/jnumed.117.203109
24   Wimberley, C. *et al.* Impact of Endothelial 18-kDa Translocator Protein on the
Quantification of (18)F-DPA-714. *J Nucl Med* **59**, 307-314 (2018).
https://doi.org:10.2967/jnumed.117.195396
25   Foray, C. *et al.* Interrogating Glioma-Associated Microglia and Macrophage
Dynamics Under CSF-1R Therapy with Multitracer In Vivo PET/MRI. *J Nucl Med*
63, 1386-1393 (2022). https://doi.org:10.2967/jnumed.121.263318
26   Palleis, C. *et al.* In Vivo Assessment of Neuroinflammation in 4-Repeat Tauopathies.
Mov Disord **36**, 883-894 (2021). https://doi.org:10.1002/mds.28395
27   Pascoal, T. A. *et al.* Microglial activation and tau propagate jointly across Braak
stages. *Nat Med* **27**, 1592-1599 (2021). https://doi.org:10.1038/s41591-021-01456-w
28   Albert-Smet, I. *et al.* Applications of Light-Sheet Microscopy in Microdevices.
Frontiers in Neuroanatomy **13** (2019). https://doi.org:10.3389/fnana.2019.00001
29   Paulus, W. & Peiffer, J. Intratumoral histologic heterogeneity of gliomas. A
quantitative study. *Cancer* **64**, 442-447 (1989). https://doi.org:10.1002/1097-
0142(19890715)64:2<442::aid-cncr2820640217>3.0.co;2-s
30   Jackson, R. J. *et al.* Limitations of stereotactic biopsy in the initial management of
gliomas. *Neuro Oncol* **3**, 193-200 (2001). https://doi.org:10.1093/neuonc/3.3.193
31   Banati, R. B. *et al.* Selective, high-contrast detection of syngeneic glioblastoma in
vivo. *Sci Rep* **10**, 9968 (2020). https://doi.org:10.1038/s41598-020-67036-z
32   Betlazar, C. *et al.* Mitochondrial Translocator Protein (TSPO) Expression in the Brain
After Whole Body Gamma Irradiation. *Front Cell Dev Biol* **9**, 715444 (2021).
https://doi.org:10.3389/fcell.2021.715444
33   Banati, R. B. *et al.* Positron emission tomography and functional characterization of a
complete PBR/TSPO knockout. *Nat Commun* **5**, 5452 (2014).
https://doi.org:10.1038/ncomms6452
34   Quach, S. *et al.* TSPO PET signal using [(18)F]GE180 is associated with survival in
recurrent gliomas. *Eur J Nucl Med Mol Imaging* (2022).
https://doi.org:10.1007/s00259-022-06006-1
35   Law, I. *et al.* Joint EANM/EANO/RANO practice guidelines/SNMMI procedure
standards for imaging of gliomas using PET with radiolabelled amino acids and
[(18)F]FDG: version 1.0. *Eur J Nucl Med Mol Imaging* **46**, 540-557 (2019).
https://doi.org:10.1007/s00259-018-4207-9
36   Wang, J. *et al.* Inhibition of translocator protein 18 kDa suppressed the progression of
glioma via the ELAV-like RNA-binding protein 1/MAPK-activated protein kinase 3
axis. *Bioengineered* **13**, 7457-7470 (2022).
https://doi.org:10.1080/21655979.2022.2048992
37   Solomon, I. *et al.* CD25-Treg-depleting antibodies preserving IL-2 signaling on
effector T cells enhance effector activation and antitumor immunity. *Nature Cancer* **1**,
1153-1166 (2020). https://doi.org:10.1038/s43018-020-00133-0
38   Wang, Y.-C. *et al.* Targeting monoamine oxidase A-regulated tumor-associated
macrophage polarization for cancer immunotherapy. *Nature Communications* **12**,
3530 (2021). https://doi.org:10.1038/s41467-021-23164-2
39   Sade-Feldman, M. *et al.* Defining T Cell States Associated with Response to
Checkpoint Immunotherapy in Melanoma. *Cell* **175**, 998-1013.e1020 (2018).
https://doi.org:https://doi.org/10.1016/j.cell.2018.10.038

1369	40	Martín-Ruiz, A. *et al.* Effects of anti-PD-1 immunotherapy on tumor regression:
1370		insights from a patient-derived xenograft model. *Scientific Reports* **10**, 7078 (2020).
1371		https://doi.org:10.1038/s41598-020-63796-w
1372	41	Couturier, C. P. *et al.* Single-cell RNA-seq reveals that glioblastoma recapitulates a
1373		normal neurodevelopmental hierarchy. *Nature Communications* **11**, 3406 (2020).
1374		https://doi.org:10.1038/s41467-020-17186-5
1375	42	Berroterán-Infante, N., Tadić, M., Hacker, M., Wadsak, W. & Mitterhauser, M.
1376		Binding Affinity of Some Endogenous and Synthetic TSPO Ligands Regarding the
1377		rs6971 Polymorphism. *International journal of molecular sciences* **20** (2019).
1378		https://doi.org:10.3390/ijms20030563
1379	43	Herculano-Houzel, S. The human brain in numbers: a linearly scaled-up primate brain.
1380		*Front Hum Neurosci* **3**, 31 (2009). https://doi.org:10.3389/neuro.09.031.2009
1381	44	Cai, R. *et al.* Panoptic imaging of transparent mice reveals whole-body neuronal
1382		projections and skull-meninges connections. *Nat Neurosci* **22**, 317-327 (2019).
1383		https://doi.org:10.1038/s41593-018-0301-3
1384	45	Pidhorskyi, S., Morehead, M., Jones, Q., Spirou, G. & Doretto, G. J. a. p. a. syGlass:
1385		interactive exploration of multidimensional images using virtual reality Head-mounted
1386		displays. (2018).
1387	46	Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist
1388		algorithm for cellular segmentation. *Nat Methods* **18**, 100-106 (2021).
1389		https://doi.org:10.1038/s41592-020-01018-x
1390	47	Uhlén, M. *et al.* Tissue-based map of the human proteome. *Science* **347**, 1260419
1391		(2015). https://doi.org:doi:10.1126/science.1260419
1392	48	Spangenberg, E. *et al.* Sustained microglial depletion with CSF1R inhibitor impairs
1393		parenchymal plaque development in an Alzheimer's disease model. *Nat Commun* **10**,
1394		3758 (2019). https://doi.org:10.1038/s41467-019-11674-z
1395	49	Haddad, A. F. *et al.* Mouse models of glioblastoma for the evaluation of novel
1396		therapeutic strategies. *Neurooncol Adv* **3**, vdab100 (2021).
1397		https://doi.org:10.1093/noajnl/vdab100
1398	50	Colombo, A. V. *et al.* Microbiota-derived short chain fatty acids modulate microglia
1399		and promote Abeta plaque deposition. *Elife* **10** (2021).
1400		https://doi.org:10.7554/eLife.59826
1401	51	Xiang, X. *et al.* The Trem2 R47H Alzheimer's risk variant impairs splicing and
1402		reduces Trem2 mRNA and protein in mice but not in humans. *Mol Neurodegener* **13**,
1403		49 (2018). https://doi.org:10.1186/s13024-018-0280-6
1404	52	Smith, S. J. *et al.* Metabolism-based isolation of invasive glioblastoma cells with
1405		specific gene signatures and tumorigenic potential. *Neuro-Oncology Advances* **2**
1406		(2020). https://doi.org:10.1093/noajnl/vdaa087
1407	53	Black, D. *et al.* Characterization of autofluorescence and quantitative protoporphyrin
1408		IX biomarkers for optical spectroscopy-guided glioma surgery. *Scientific Reports* **11**,
1409		20009 (2021). https://doi.org:10.1038/s41598-021-99228-6
1410	54	Brendel, M. *et al.* Glial Activation and Glucose Metabolism in a Transgenic Amyloid
1411		Mouse Model: A Triple-Tracer PET Study. *J Nucl Med* **57**, 954-960 (2016).
1412		https://doi.org:10.2967/jnumed.115.167858
1413	55	Overhoff, F. *et al.* Automated spatial brain normalization and hindbrain white matter
1414		reference tissue give improved [18F]-florbetaben PET quantitation in Alzheimer's
1415		model mice. *Frontiers in neuroscience* **10**, 45 (2016).
1416	56	Deussing, M. *et al.* Coupling between physiological TSPO expression in brain and
1417		myocardium allows stabilization of late-phase cerebral [(18)F]GE180 PET
1418		quantification. *Neuroimage* **165**, 83-91 (2017).
1419		https://doi.org:10.1016/j.neuroimage.2017.10.006

57 Unterrainer, M. *et al.* TSPO PET, tumour grading and molecular genetics in
histologically verified glioma: a correlative (18)F-GE-180 PET study. *Eur J Nucl Med*
Mol Imaging **47**, 1368-1380 (2020). https://doi.org:10.1007/s00259-019-04491-5
58 Unterrainer, M. *et al.* Towards standardization of (18)F-FET PET imaging: do we
need a consistent method of background activity assessment? *EJNMMI Res* **7**, 48
(2017). https://doi.org:10.1186/s13550-017-0295-y
59 Zhao, S. *et al.* Cellular and Molecular Probing of Intact Human Organs. *Cell* **180**, 796-
812 e719 (2020). https://doi.org:10.1016/j.cell.2020.01.030
60 Erturk, A. *et al.* Three-dimensional imaging of solvent-cleared organs using 3DISCO.
Nat Protoc **7**, 1983-1995 (2012). https://doi.org:10.1038/nprot.2012.119
61 Hughes, C. S. *et al.* Single-pot, solid-phase-enhanced sample preparation for
proteomics experiments. *Nat Protoc* **14**, 68-85 (2019). https://doi.org:10.1038/s41596-
018-0082-x
62 Demichev, V., Messner, C. B., Vernardis, S. I., Lilley, K. S. & Ralser, M. DIA-NN:
neural networks and interference correction enable deep proteome coverage in high
throughput. *Nature Methods* **17**, 41-44 (2020). https://doi.org:10.1038/s41592-019-
0638-x
63 Tyanova, S. *et al.* The Perseus computational platform for comprehensive analysis of
(prote)omics data. *Nat Methods* **13**, 731-740 (2016).
https://doi.org:10.1038/nmeth.3901
64 Tusher, V. G., Tibshirani, R. & Chu, G. Significance analysis of microarrays applied
to the ionizing radiation response. *Proc Natl Acad Sci U S A* **98**, 5116-5121 (2001).
https://doi.org:10.1073/pnas.091062498
1443

A

B
Representative VR segmented GFP volume
Representative VR segmented CD11b volume

C
$p = 0.037$
Light sheet volume (mm³)
GFP | CD11b

D
DAPI / Tumor cells / TAMs | DAPI / Tumor cells | DAPI / TAMs | Nuclei segmentation

Inside the tumor

Tumor border

Tumor periphery

E
4493 μm³ GFP / cell
2713 μm³ CD11b / cell

$p = 0.042$
Cell count in SB28 tumors (n)
Tumor cells | TAMs

Tumor cell contr. (%) | TAM contr. (%)
3.5 : 1

F
PET | Predicted 3D histology | 3D histology

Tumor (GFP) / TAMs (CD11b)

G
a

b

H
$R = 0.624$
$p < 0.0001$
Light sheet signal intensity (GFP)
Predicted GFP (arb.)

$R = 0.304$
$p < 0.0001$
Light sheet signal intensity (CD11b)
Predicted CD11b (arb.)

I
PET tumor radioactivity (%ID*BW)
Predicted tumor radioactivity (%ID*BW)
$R = 0.942$
$p = 0.0015$